A NEW COUNTEREXAMPLE TO NGUYEN'S CONJECTURE ON SURFACE FIBRATION †

CAI WENYI¹, XIAO HAO¹, QIAN YUANYUAN², ZHUO LINGYU*

ABSTRACT. Suppose $f:S\to\mathbb{P}^1$ is a fibration of genus g with 3 singular fibers, and two of the fibers are semistable. In 1998, Nguyen conjectured in [Ng] that such kind of fibration does not exist for $g\geq 2$. But in 2013, Cheng Gong, Xin Lu, and Sheng-Li Tan found a counterexample to Nguyen's conjecture for g=2 in [GLT]. Note that such kind of fibration shows strong arithmetic properties, and as such the counterexamples are important, but rare in fact. In this paper, a new counterexample to Nguyen's conjecture for g=2 is constructed.

KEYWORDS. Algebraic surface; Fibration; Singular fiber.

[†] This paper received financial support both from National University Students' Innovation and Entrepreneurship Training Programs, Research on Security of Digital Signature (201710285014Z), and National Natural Science Foundation of China, Research on Surface Fibration with Three Singular Fibers (11401413).

¹Wenyi Cai, the initiator of this project, is a female math-majored junior undergraduate enrolled in the class of 2019 at Soochow University by the time when this paper is finished.

¹Hao Xiao, the main contributor to this paper, is a male math-majored junior undergraduate enrolled in the class of 2019 at Soochow University by the time when this paper is finished.

References

[GLT] C. Gong, X. Lu, and S.-L. Tan, Familles de courbes sur \mathbb{P}^1 avec trois fibres singulières, C. R. Acad. Sci. Paris - Sér. I - Math. **351** (2013), 375-380.

[Ng] K. V. Nguyen, On families of curves over \mathbb{P}^1 with small number of singular fibres, C. R. Acad. Sci. Paris - Sér. I - Math. **326** (1998), 459-463.

School of Mathematical Sciences, Soochow University, Suzhou, Jiangsu 215000, China