

Mathematica Compendium

Miguel Antonio Méndez Hernández

May 20, 2026

Contents

1	Guidelines for Writing Formal Mathematical Proofs	25
2	Logic	27
2.1	Propositional Logic	27
2.1.1	Predicates	27
2.1.2	Predicate Substitution	27
2.1.3	Logic Operators and Truth Tables	28
2.1.4	Tautology and Contradiction	28
2.1.5	Logical Equivalences	29
2.1.6	Truth Tables	30
2.1.7	Disjunctive Normal Form (DNF)	31
2.1.8	Conjunctive Normal Form (CNF)	31
2.1.9	Basis	32
2.1.10	Karnaugh Maps	32
2.1.11	Mathematical Quantifiers	33
2.1.12	Complex Statements	34
3	Set Theory	35
3.0.1	Basics	35
3.0.2	Sets and Classes	37
3.0.3	Axioms of Set Theory	37
3.0.4	Consequences of the Axiom of Replacement	39
3.0.5	Subclasses/Subsets	39
3.0.6	The Universal Class	39
3.0.7	The Empty Class	39
3.0.8	The Empty Set	39
3.0.9	Set Complement Proof	39
3.0.10	Proof The Power Set of a Set is not an Element of the Set	39
3.0.11	Proof of the Union of Subsets	40
3.0.12	The Singleton and Doubleton	40
3.0.13	Ordered Pair	41
3.0.14	The Cartesian Product	41
3.0.15	Laws of Set Algebra	43
3.0.16	Proof of De Morgans's Law for sets and logic	44
3.0.17	Indexed Sets	45
3.0.18	Generalized Union, Intersection and Cartesian Product for Indexed Sets	45
3.0.19	Pairwise Disjoint	45
3.0.20	Disjoint Decomposition	45
3.0.21	Cardinality Laws	45
3.0.22	Cardinality	46
3.0.23	Definition of Countable and Non-countable	46
3.0.24	Countable Union	47

3.0.25	Countability of Subsets	47
3.0.26	Countability of the Rational Numbers	47
3.0.27	Cantor's Diagonal Argument: The Real Numbers are Uncountable	47
3.0.28	The Power Set	48
3.0.29	Set Systems	48
3.0.30	Intersection and Union of Set Systems	48
3.0.31	Equivalence Classes	50
3.0.32	Partitions	50
3.0.33	Relation from the Relation of the Quotient Space and Partition from the Quotient Space	51
3.0.34	Equivalence Relations and Partitions	52
3.0.35	Equivalence Relations under Functions	52
3.0.36	Refinement	52
3.0.37	Quotient Space for Refinements	52
3.0.38	Equivalence Relations and Surjective Mappings	53
3.0.39	Homomorphism Theorem for Sets	53
3.0.40	Representatives	54
3.0.41	Family of Subsets Operations	54
3.0.42	Product of Classes	54
3.0.43	Pigeon Principle	54
3.0.44	The Cantor Set	54
4	Relations, Maps and Functions	59
4.0.1	Relations	59
4.0.2	Types of relations	59
4.0.3	Diagonal Graph	59
4.0.4	Order Relation	60
4.0.5	Strict Order Relation	60
4.0.6	Equivalence relation	60
4.0.7	Equivalence Class	60
4.0.8	Lemmi about Equivalence Classes	61
4.0.9	Quotient Space	61
4.0.10	Map Definition	61
4.0.11	The Set of All Maps	61
4.0.12	The Characteristic Map	61
4.0.13	Functions	62
4.0.14	The Graph	62
4.0.15	Domain	63
4.0.16	The identity	63
4.0.17	Image/Range	63
4.0.18	Composition of Maps	64
4.0.19	Types of Functions	65
4.0.20	Propositions on Images and Pre-images under Set Operations	65
4.0.21	Bijection of the Power Set and 2^A	67
4.0.22	Restriction Function	67
4.0.23	Even Function	67
4.0.24	Odd Function	67
4.0.25	Pre-image	68
4.0.26	Inverse of a Function	68
4.0.27	Transformations of a Function	71
4.0.28	Counting Functions	71
4.0.29	Counting Relations	72
5	Mathematical Proofs	73
5.0.1	Proof by Direct Argument	73
5.0.2	Proof by Contradiction	73
5.0.3	Proof by Induction	74
5.0.4	Strong Induction	75

5.0.5	Proof by Exhaustion	75
5.0.6	Proof by Cases	76
5.0.7	Proof by Construction	76
5.0.8	Proof by Counterexample	76
5.0.9	Proof by Contrapositive	77
5.0.10	Proof by Reduction to Absurdity	77
5.0.11	Proof by Analogy	78
6	Graph Theory	79
6.0.1	Connected Graphs	79
6.0.2	Paths	79
6.0.3	The Adjacency Matrix	80
6.0.4	The Relax Operation	80
7	Order	81
7.0.1	Partially Ordered Classes	81
7.0.2	Lexicographical Ordering	81
7.0.3	Antilexicographical Ordering	81
7.0.4	Comparable	81
7.0.5	Fully Ordered Subclass	81
7.0.6	Initial Segments	82
7.0.7	Cuts	82
7.0.8	Order Preserving Functions	83
7.0.9	Isomorphism	83
7.0.10	Strictly Increasing Function	83
7.0.11	Order by Inverses	83
7.0.12	Common Isomorphism on Partially Ordered Classes	84
7.0.13	Isomorphic Definition	84
7.0.14	Convex	84
7.0.15	Maximal Element	85
7.0.16	Minimal Element	85
7.0.17	Upper Bound	85
7.0.18	Lower Bound	86
7.0.19	Maximum	86
7.0.20	Minimum	86
7.0.21	Supremum and Infimum	86
7.0.22	Duality	86
7.0.23	Greatest and Least Elements under Inclusion	86
7.0.24	Upper and Lower Bounds under Inclusion	87
7.0.25	Suprema and Infima under Inclusion	87
7.0.26	Relation between a Set and Bounds of Bounds	87
7.0.27	Infimum via Lambda Function	88
7.0.28	Bounded	89
7.0.29	Conditionally Complete	89
7.0.30	Lattices	89
7.0.31	Definition of Less Than	89
7.0.32	Sublattices	90
7.0.33	Boolean Algebra	90
7.0.34	Fully Ordered Classes	90
7.0.35	Principle of the Well-ordering	90
7.0.36	Order in Fields	90
8	The Natural Numbers	93
8.0.1	Definition as Sets	93
8.0.2	Addition	93
8.0.3	Inductive Sets	93
8.0.4	Axiom of Infinity	93
8.0.5	Definition of the Sets of Natural Numbers	93
8.0.6	The Standard Order	93

8.0.7	The Peano Axioms	94
8.0.8	Propositions and Proofs	94
8.0.9	Definition of Addition in $\mathbb{N}$	95
9	The Archimedean Principle	97
9.0.1	Equivalent Formulation	97
9.0.2	Applications	97
10	Arithmetic	99
10.0.1	Even and Odd numbers	99
10.0.2	Perfect Squares	99
10.0.3	Division Algorithm	99
10.0.4	Prime Numbers	100
10.0.5	Pythagorean Triplets	100
10.0.6	Divisibility	100
10.0.7	Division	100
10.0.8	Greatest Common Denominator	100
10.0.9	The Euclidean Algorithm	100
10.0.10	Euclidean Algorithm as Linear Map	101
10.0.11	Bezout Identity	102
10.0.12	Euclidean Algorithm as Matrix Multiplication	102
10.0.13	Prime Field	103
10.0.14	Implication of Prime Numbers	103
10.0.15	Fundamental Theorem of Arithmetic	103
10.0.16	Lemmas	104
10.0.17	Proof of the Fundamental Theorem of Arithmetic	104
10.0.18	Least Common Multiple	105
10.0.19	Multiplication Property of the Least Common Multiple	105
10.0.20	Second Proof of the Fundamental Theorem of Arithmetic	105
10.0.21	Goldbach's Conjecture	106
10.0.22	Prime Factorization	106
10.0.23	GCD via Prime factorization	106
11	Real Numbers	107
11.0.1	Properties derivated from the Axioms	107
11.0.2	Existence of Solutions	108
11.0.3	The Completeness Axiom	108
11.0.4	Minima in the Real Numbers	108
11.0.5	Approximation of the Supremum	109
11.0.6	Supremum of the Naturals	109
11.0.7	Remainder of Archimedes and Eudoxus	109
11.0.8	Density of the Rationals in the Reals	110
11.0.9	Existence of Square Roots	110
11.0.10	Irrational Numbers	110
11.0.11	Absolute Value for Real Numbers	110
11.0.12	Norms Remainder	111
11.0.13	Distance	111
11.0.14	Nested Interval Property	111
11.0.15	Definition as Cauchy Sequences	111
11.0.16	Definition	112
12	Complex Numbers	115
12.0.1	The Complex Plane	115
12.0.2	Conjugate of a Complex Number	115
12.0.3	Operations in Cartesian Coordinates	116
12.0.4	Polar Coordinates	116
12.0.5	Multiplication and Division in Polar Coordinates	116
12.0.6	Exponentiation and Roots (De Moivre's Theorem)	117
12.0.7	Example: Solve $z^4 = 1 + \sqrt{3}i$	117

12.0.8	Solving Equations with Complex Numbers	118
12.0.9	The Complex Logarithm	119
12.0.10	Complex Exponents	120
12.0.11	Euler's Formula Proof	120
12.0.12	The Fundamental Theorem of Algebra	121
12.0.13	Complex Numbers as Matrices	121
13	Topology	123
13.0.1	Introduction to topological nomenclature	123
14	Inequalities	125
14.0.1	Triangle Inequality	125
14.0.2	Reciprocal Inequality	125
14.0.3	Bernoulli Inequality	125
14.0.4	Inequalities for Real Numbers	125
14.0.5	Elementary Young Inequalities	126
14.0.6	Inequalities with and without Absolute Value	126
15	Means	129
15.0.1	Arithmetic Mean	129
15.0.2	Geometric Mean	129
15.0.3	Harmonic Mean	129
15.0.4	Quadratic Mean	129
15.0.5	Proof of the Arithmetic Mean-Geometric Mean Inequality	129
15.0.6	Proof of the Harmonic Mean Geometric Mean Inequality	131
16	Solving Polynomial Equations	133
16.0.1	The PQ Formula	133
16.0.2	The Quadratic Formula	133
16.0.3	The Cubic Formula	134
17	The Binomial Coefficient	137
17.0.1	The Binomial Theorem	137
17.0.2	Pascal's Triangle	138
18	Proportionality and the Rule of Three	141
18.0.1	Proportionality	141
18.0.2	The Rule of Three (Simple)	141
18.0.3	The Rule of Three (Compound)	142
19	Factorization	143
19.0.1	Common Factor	143
19.0.2	Difference of Squares	143
19.0.3	Sum of Cubes	143
19.0.4	Difference of Cubes	143
19.0.5	Trinomial: Special Case $x^2 + bx + c$	143
19.0.6	Trinomial: General Form $ax^2 + bx + c$	143
19.0.7	Perfect Square Trinomial	144
19.0.8	Substitution	144
19.0.9	Rationalization of Radicals	144
19.0.10	Horner's Method	144
19.0.11	Long Division of Polynomials	145
20	Partial Fractions	147
20.0.1	The Simplest Case	147
20.0.2	Lemma 2	148
21	Recursion	151
21.0.1	From Recursive to Closed Form	151
21.0.2	Linear Recursion	151

21.0.3	Recursive Inequalities	151
21.0.4	Fibonacci Numbers and Closed Form	152
22	Sum and Product Notation	153
22.0.1	Sum Notation $\sum$	153
22.0.2	Geometric Series	154
22.0.3	Product Notation $\prod$	155
22.0.4	Proof of the geometric Series	155
23	Logarithms	157
23.0.1	Properties of Logarithms	157
23.0.2	Fundamental Identity of Logarithms	157
23.0.3	Change of Base Formula	158
23.0.4	The Chain Rule	158
23.0.5	The Derivative of the Natural Logarithm	158
23.0.6	Proofs of the Properties	158
24	Fractions, Roots, and Exponents	161
24.0.1	Fractions	161
24.0.2	Roots and Exponents	161
25	Generating Functions	163
25.0.1	Connection to Subsets	163
25.0.2	Closed Form of the Fibonacci Sequence	163
26	Modular Arithmetic	167
26.0.1	Residual Classes and Residual Class Rings	167
26.0.2	Properties	167
26.0.3	General Division Rules	167
26.0.4	Using Modular Arithmetic to Find the n-th Digit of a Large Number	168
26.0.5	Gaussian Brackets	168
26.0.6	Residual Class Ring modulo n	169
26.0.7	Equivalence Class of $\mathbb{Z}$	169
27	Relational Algebra	171
27.0.1	Relations	171
27.0.2	The Cartesian Product for Relation Algebra	171
27.0.3	Relational Algebra	171
27.0.4	Domain of an Attribute	172
27.0.5	Types of Keys	176
27.0.6	Relation Schema	176
27.0.7	Constraint	176
27.0.8	Functional Dependencies	176
27.0.9	Armstrong Axioms	177
27.0.10	Attribute Closure	177
27.0.11	Canonical Cover	177
27.0.12	How to find a Candidate Key	178
27.0.13	Losslessness and Maintenance of Dependency	178
27.0.14	Normal Forms	179
27.0.15	Synthesis Algorithm	180
27.0.16	Decomposition Algorithm	181
27.0.17	Optimization	182
28	Geometry	185
28.0.1	Formulas for Area and Perimeter of Geometric Figures	185
28.0.2	Formulas for Area and Volume of 3D Geometric Figures	185
28.0.3	Thales' Theorem	186
28.0.4	Conversion Between Radians and Degrees	187
28.0.5	Geometric Figures with Formulas (2D)	187

28.0.6	3D Geometric Figures with Formulas	188
28.0.7	Axioms of Euclidean Geometry	189
28.0.8	Types of Angles and Angle Relationships	189
28.0.9	Triangle Similarity and Congruence	190
28.0.10	Visualizations of Triangle Similarity and Congruence	191
28.0.11	Triangle Medians, Angle Bisectors, and Altitudes	192
28.0.12	The Midsegment Theorem	193
28.0.13	Intercept Theorems	193
28.0.14	Proof of the Pythagorean Theorem	194
29	Trigonometry	195
29.0.1	Trigonometric Functions	195
29.0.2	SOH-CAH-TOA	195
29.0.3	Pythagorean Identities	195
29.0.4	Sum and Difference Formulas	195
29.0.5	Double Angle Formulas	196
29.0.6	Half Angle Formulas	196
29.0.7	Product to Sum Formulas	197
29.0.8	Power Reducing Formulas	197
29.0.9	Even and Odd Functions	197
29.0.10	Graph Identity	197
29.0.11	Trigonometric Values	198
29.0.12	Unit Circle and Trigonometric Functions	198
29.0.13	Definitions on the Unit Circle:	198
29.0.14	Trigonometric Rules	199
29.0.15	Proof of the Cosine Rule	199
30	Linear Systems of Equations	201
30.1	Linear Systems Standard Form	201
30.2	Linear System as Matrices	201
30.2.1	Solution Set of a Linear System	202
30.2.2	Equality of the Rank $\text{rank}(A) = \text{rank}_S(A)$	202
30.2.3	Solvability of the Linear System $Ax = b$	202
30.2.4	Solution Set of the Linear System $Ax = b$	203
30.2.5	Free Variables	203
30.2.6	Gaussian Elimination	203
30.2.7	Invariability of the Solution Set	204
30.2.8	Gauss-Jordan Elimination	204
30.2.9	Homogeneous Linear Equations	205
30.2.10	Particular Solution	206
30.2.11	General = Particular + Homogeneous	206
30.2.12	Linear Combination Lemma	207
30.2.13	Example: Gaussian Elimination with 3 Equations and 4 Unknowns	207
30.2.14	The Determinant, the cross product and the solutions of linear Systems of Equations	208
30.2.15	Chart for the number of solutions	208
30.2.16	Proof of the Gaussian Elimination	209
30.2.17	Alternative Method for finding the Rank of a matrix	210
30.2.18	Cramer's Rule	210
30.2.19	LU Decomposition	211
30.2.20	Cholesky Decomposition	217
31	Over/Underdetermined Systems of Equations	219
31.0.1	Normal Equations	219
31.0.2	Method of the Least Squares	220
31.0.3	Underdetermined Systems of Equations	221
32	Analytical Geometry	223
32.0.1	Vectors and Points	223
32.0.2	Vector Addition and Scalar Multiplication	223

32.0.3	Equation of a Line	224
32.0.4	Equation of a Plane	224
32.0.5	Dot Product	224
32.0.6	Orthogonal Vectors	225
32.0.7	The Orthogonal Projection	225
32.0.8	Vector Projections in Data Science	226
32.0.9	Dot Product as Projection	226
32.0.10	Metrics for Vectors	227
32.0.11	Angle Relations	227
32.0.12	Line Relations	227
32.0.13	Normalization of a vector	228
32.0.14	The Cross Product	228
32.0.15	Orthogonal vectors in $\mathbb{R}^2$ $\mathbb{R}^3$	230
32.0.16	Hessian Normal Form	230
32.0.17	Converting from the parametric form to the Hessian normal form	231
32.0.18	Converting from the Hessian normal form to the parametric form	231
32.0.19	Properties of lines and planes	231
32.0.20	Convert Normal Vector in Two Direction Vectors	232
32.0.21	Intersection between Line and Plane	232
32.0.22	Distances between points, lines and planes	233
32.0.23	Foot of the Perpendicular and Mirror Point	236
33	Algebraic Structures	237
33.0.1	Operations: Internal and External	237
33.0.2	Properties of Operations	237
33.0.3	Homomorphisms and Isomorphisms	237
33.0.4	Lattice	238
33.0.5	Semigroup $(S, \oplus)$	238
33.0.6	Monoid $(M, \oplus)$	238
33.0.7	Group $(G, \oplus)$	238
33.0.8	Abelian Group $(A, \oplus)$	238
33.0.9	Ring $(R, \oplus, \odot)$	238
33.0.10	Commutative Ring $(R, \oplus, \odot)$	239
33.0.11	Field $(\mathbb{F}, \oplus, \odot)$	239
33.0.12	Vector Space $(V, \oplus, \cdot)$	240
34	Polynomial Rings	241
34.0.1	Degree of a Polynomial	242
34.0.2	Properties of the Polynomial Ring	242
34.0.3	Field Algebra	243
34.0.4	Division for Polynomial	243
34.0.5	Greatest Common Divisor for Polynomials	244
34.0.6	More Properties of Polynomials	244
34.0.7	Euclidean Algorithm for Polynomials	244
34.0.8	Reducible Irreducible	244
34.0.9	Normalized	245
34.0.10	Factorization Property	245
35	Vector Spaces	247
35.0.1	Subspaces	248
35.0.2	Properties of the subspaces	248
35.0.3	Sum of Subspaces	248
35.0.4	Linear Combinations	248
35.0.5	Linear Independence	249
35.0.6	Basis	249
35.0.7	Dimension	249
35.0.8	How to find the basis of a set vector	250
35.0.9	Basis Extension Theorem	250
35.0.10	Epimorphism of Finite Dimensional Vector Spaces	251

35.0.11	Exchange Lemma	251
35.0.12	Dimension of a sum of subspaces	252
35.0.13	Linear Independence of polynomials	252
35.0.14	Interpolation Polynomial	252
36	Dot Product, Euclidean and Unitary Space	255
36.0.1	Outer Product	255
36.0.2	Outer Product	255
36.0.3	Inner Product	255
36.0.4	Trigonometric Definition	256
36.0.5	Standard Inner Product for Complex Number	256
36.0.6	Inner Product on $C[a, b]$	256
36.0.7	Euclidean and Unitary Vector Spaces	256
36.0.8	Norms in Vector Spaces	257
36.0.9	Cauchy-Schwarz Inequality in Unitary Vector Spaces	257
36.0.10	Definiteness of Inner Products	258
36.0.11	Definiteness and Quadratic Forms	258
36.1	Positive Definite Matrices	259
36.1.1	Conditions for Positive Definiteness	259
36.1.2	Minors	260
36.1.3	Connection to Symmetric Matrices	260
36.1.4	Sign of the D_k	261
36.1.5	Negative Quadratic Form	261
36.1.6	Theorems for Non-squares matrices	261
36.1.7	Theorem II	261
37	Orthogonality	263
37.0.1	Orthogonality and Projection	263
37.0.2	Induced Metric	263
37.0.3	Orthogonal Projection and Orthogonal Complement	265
37.0.4	Orthogonality of the basis to the Complement	265
37.0.5	How to find the orthogonal complement	265
37.0.6	Orthogonal and Orthonormal Systems	265
37.0.7	Writing a vector in terms of the Orthogonal Basis	266
37.0.8	Linear Independence of Orthogonal Systems	266
37.0.9	Representation with Respect to an Orthogonal Basis	266
37.0.10	Orthogonal Projection and Direct Sum Decomposition	267
37.0.11	The Gram-Schmidt Process	267
37.0.12	Existence of Orthonormal Bases and Orthogonal Projections	268
37.0.13	Best Approximation	269
38	Matrices	271
38.0.1	Definition of a Matrix	271
38.0.2	Matrix Addition and Subtraction	271
38.0.3	Matrix Multiplication	272
38.0.4	Outer Product for Matrix Multiplication	273
38.0.5	The Transpose of a Matrix	274
38.0.6	The Equivalence of Matrices	274
38.0.7	The Inverse of a Matrix	277
38.0.8	The Rank of a Matrix	279
38.0.9	How to isolate Matrices	279
38.0.10	Column Space	280
38.0.11	Row Space	280
38.0.12	Null Space	280
38.0.13	Linear Maps as matrices	281
38.0.14	Matrix Representation of Linear Maps on Basis vectors	281
38.0.15	Matrices with respect to a basis	282
38.0.16	Example of a complete exercise	282
38.0.17	Diagonal and Triangular Matrices	283

39 Metrics	285
39.1 Metrics for Vectors	285
39.2 Metric for Matrices	285
39.2.1 Spectral Metric	286
39.2.2 Row Metric	286
39.2.3 Column Metric	286
39.2.4 Frobenius Metric	286
40 Linear Maps	287
40.0.1 Types of Linear Maps	287
40.0.2 Properties of Linear Maps	287
40.0.3 Composition as Matrix Multiplication	288
40.0.4 Injectivity and Surjectivity	288
40.0.5 Linear Extension Theorem	288
40.0.6 The Kernel of a Linear Map	289
40.0.7 The Image of a Linear Map	289
40.0.8 The Rank of a Linear Map	289
40.0.9 The Nullity of a Linear Map	290
40.0.10 Proof of the injectivity of the Kernel	290
40.0.11 Dimension Formula for Linear Mappings	290
40.0.12 Linear Maps and Matrices	291
40.0.13 How to identify the type of Linear Map Summary	291
41 The Change Of Basis	293
41.0.1 Pseudo-equality Of Translations	293
41.0.2 Coordinate mapping	293
41.0.3 Coordinates with respect to a basis	294
41.0.4 Translation of Coordinates	294
41.0.5 Change of Basis Theorem	294
41.0.6 Change of Basis in Linear Maps	294
41.0.7 Transformations involving multiple bases	295
42 The Determinant of a Matrix	297
42.0.1 Properties of the determinant	297
42.0.2 Proof of the multiplication of determinants	298
42.0.3 The Leibniz Formula	298
42.0.4 Laplace's Method (Co-factor Expansion)	299
42.0.5 Determinant of the reduced echelon form	301
42.0.6 Gaussian Method for finding determinants	301
43 Elementary Matrix	303
43.0.1 Invertibility	304
43.0.2 Determinants	304
44 Orthogonal Linear Maps	305
44.0.1 Isometry	305
44.0.2 Unit-sphere	305
44.0.3 The Rotation Matrix	305
44.0.4 Orthogonal Matrices	306
44.0.5 The Determinants of Orthogonal Matrices	308
44.0.6 The Orthogonal Group	308
44.0.7 Dot Product	308
44.0.8 Orthonormal Matrices	309
44.0.9 Theorem of the Matrix Representation	310
44.0.10 Change of Basis	310
44.0.11 QR-Decomposition	310
45 Eigenvectors and Eigenvalues	313
45.0.1 Characteristic Polynomial	313

45.0.2	Theorems of Eigenvalues and Eigenvectors	314
45.0.3	Linear Independence Of Eigenvectors With Distinct Eigenvalues	314
45.0.4	The Determinant Via Eigenvalues	314
45.0.5	Eigenvalues Under Isometric Maps	315
45.0.6	Eigenvalues As Powers	315
45.0.7	How to find the Eigenvectors and Eigenvalues	315
45.0.8	Complex Eigenvalues	316
45.0.9	Diagonal Matrices	317
45.0.10	Singular Value Decomposition (SVD)	321
45.0.11	Similar Matrices	322
45.0.12	Jordan Normal Form	323
45.0.13	The Hermitian Transpose	324
45.0.14	Unitary Matrices	324
45.0.15	Unitary Decomposition (Schur's Theorem)	325
45.0.16	Normal Matrices	325
45.0.17	Diagonalisability Criteria Based On Normal Matrices	325
45.0.18	Properties of Unitary Diagonalisable Matrices	326
45.0.19	Hermitian Matrices	326
45.0.20	Real Eigenvalues Via Hermitian Matrices	327
45.0.21	Gerschgorin's Theorem	327
45.0.22	Approximation Of The Differences Of Eigenvalues	327
45.0.23	Eigenvalue Of Block Matrices	328
45.0.24	Roth's Removal Rule	328
45.0.25	Eigenvalues/vectors of Matrices Representing Complex Numbers	329
45.0.26	Real Jordan Form	329
45.0.27	Real Schur Form	329
45.0.28	Orthonormal Standard Form	330
45.0.29	Criterion For Orthonormal Diagonalisation	330
45.0.30	Normal Property Of Symmetric Matrices	330
46	Dual Spaces and Dual Vectors	331
46.0.1	The Dual Basis	331
46.0.2	How to find a Dual Basis	332
46.0.3	Change of Basis in the Dual Space	333
46.0.4	Dimension Theorem	333
46.0.5	The Transpose of a Linear Transformation	333
46.0.6	Connection to the Transpose Matrix	333
46.0.7	The Dual of the Dual Space	336
46.0.8	Basis Theorem for the Dual Space	336
46.0.9	Transpose of the Transpose is T	336
46.0.10	The Annihilator	337
46.0.11	The Duality Argument	338
46.0.12	The Rank A transpose is equal to the rank of A	338
47	Tensors	341
48	The Exponential Function and Relatives	343
48.0.1	Exponential Function as a Series	343
48.0.2	Properties of the Exponential	343
48.0.3	General Definition	343
48.0.4	Difference of Exponentials	343
48.0.5	Sine and Cosine Definitions with e	344
48.0.6	Euler's Formula	344
48.0.7	Reason for the Infinite Series of e (Differentiation)	344
48.0.8	Hyperbolic Functions with e	344
48.0.9	Inverse of the Hyperbolic Trigonometric Functions	345
49	Growth	347
49.0.1	Exponential Growth	347

49.0.2	Logistic Growth	348
50	Asymptotes	351
50.0.1	Vertical Asymptotes	351
50.0.2	Horizontal Asymptotes	351
50.0.3	Oblique (Slant) Asymptotes	352
50.0.4	Summary Table	352
51	Sequences and Series	353
51.0.1	Arithmetic Sequence	353
51.0.2	Geometric Sequence	353
51.0.3	Function Sequence	353
51.0.4	Boundness of a Sequence	354
51.0.5	Convergence	354
51.0.6	The limit of a Sequence is Unique	354
51.0.7	Convergence by Power of One Half	355
51.0.8	Boundness of Convergent Sequences	355
51.0.9	Cauchy Sequence	355
51.0.10	Boundness of Cauchy Sequence	356
51.0.11	Proof that every Convergent Sequence is a Cauchy Sequence	356
51.0.12	Cauchy Sequences are Convergent	356
51.0.13	Definition of ε -Neighborhood	357
51.0.14	Supremum, Infimum, Maximum and Minimum	357
51.0.15	Properties of Suprema and Infima	358
51.0.16	Limit and Accumulation Point	359
51.0.17	The Monotone Convergence Theorem	359
51.0.18	Monotony	360
51.0.19	Monotonicity by Difference and Quotient	360
51.0.20	Divergence	360
51.0.21	Subsequences and Their Properties	360
51.0.22	Proof that -1^n is divergent	362
51.0.23	Bolzano-Weierstraß Theorem	362
51.0.24	Series	363
51.0.25	Power Series	363
51.0.26	Geometric Series	363
51.0.27	Point-wise Convergence	363
51.0.28	Uniform Convergence	363
51.0.29	Cauchy Criterion for Series	364
51.0.30	Interval Nesting and the Interval Nesting Theorem	364
51.0.31	Weierstraß Approximation Theorem	364
51.0.32	Zero Sequence	364
51.0.33	Comparison Criteria for Sequences	365
51.0.34	Properties of Zero Sequences	365
51.0.35	The Monotony Principle	365
51.0.36	Operations on Sequences	366
51.0.37	Boundness of a Function	367
51.0.38	Inverse function due to Monotony	368
51.0.39	Domain Restriction	368
52	Limits	369
52.0.1	Definition of the Limit	369
52.0.2	Alternative Definition	369
52.0.3	Limit Calculation Rules	369
52.0.4	Limits at Infinity	370
52.0.5	Indeterminate Forms	371
52.0.6	Limit of the n-th Root of n	371
52.0.7	Limit of a Composite Function	372
52.0.8	Right and Left Limits	372
52.0.9	L'Hôpital's Rule	372

52.0.10	Limit of a fraction	372
52.0.11	Limit due to Sequence	372
52.0.12	Limit of a Recursive Sequence	373
52.0.13	Important Limits	373
52.0.14	Squeeze Theorem (Sandwich Theorem)	373
52.0.15	The Number e as a Limit	374
52.0.16	Limits of $\arctan$	375
52.0.17	Bolzano-Weierstraß Theorem	375
52.0.18	Uniqueness of the Limit	375
52.0.19	Every Convergent Sequence is a Cauchy Sequence	376
52.0.20	Limsup and Liminf	376
52.0.21	Limit by Bounds	376
53	Convergence Criteria	377
53.0.1	Quotient (Ratio) Test	377
53.0.2	Zero Sequence Criterion	377
53.0.3	Majorant (Comparison Test)	378
53.0.4	Minorant	378
53.0.5	Root Test	378
53.0.6	Leibniz Criterion (Alternating Series Test)	378
53.0.7	Compactness Criterion	378
53.0.8	Cauchy Criterion I	378
53.0.9	Cauchy Criterion II	378
53.0.10	The Cauchy Product	379
53.0.11	Geometric Series	379
53.0.12	Harmonic Series	379
54	Convergence Radius	381
54.0.1	Test of the Borders	381
54.0.2	Approximating the Error	381
54.0.3	Complete Example	382
55	Calculating Series Values	383
55.0.1	Telescoping Sums and Partial Fraction Decomposition	383
56	Transforming Functions into Power Series	385
56.0.1	Common Cases	385
57	Continuity	387
57.0.1	Definition of Continuity at a Point	387
57.0.2	Characterization using sequences	388
57.0.3	Left and Right Continuity	389
57.0.4	Rules of Continuity	389
57.0.5	Continuity and Compactness	389
57.0.6	Extreme-Value Theorem	389
57.0.7	Extreme Points	390
57.0.8	Middle Value Theorem	390
57.0.9	Continuity of the Inverse by a decreasing Function	390
57.0.10	Lipschitz Continuity	390
57.0.11	Fixed Point	390
57.0.12	Fixed Point on Interval	390
57.0.13	Banach Fixed Point Theorem	390
57.0.14	Uniform Continuity	392
57.0.15	Boundedness by Uniform Continuity	393
57.0.16	Cauchy Sequence by Uniform Continuity	393
58	Differentiation	395
58.0.1	Definition of the Derivative	395
58.0.2	Derivative Rules	395

58.0.3	Extrema and Critical Points Test	396
58.0.4	Trigonometric Derivatives	396
58.0.5	Derivatives of Inverse Trigonometric Functions	397
58.0.6	The Differential	397
58.0.7	The Secant Equation and Graphical Meaning	397
58.0.8	Optimization via the Derivative	397
58.0.9	Power Rule Derivation	398
58.0.10	Implicit Differentiation	399
59	Integration	401
59.0.1	Definition (Riemann Integral)	401
59.0.2	Upper and Lower Sums	401
59.0.3	Concrete and Non-Concrete Integrals	401
59.0.4	Rules of Integration	401
59.0.5	Substitution Rule (U-Substitution)	402
59.0.6	Integration by Parts	402
59.0.7	Integration by Partial Fractions	403
59.0.8	Trigonometric Substitution	403
59.0.9	Average Value of a Function	405
59.0.10	Mean Value Theorem for Integrals	405
59.0.11	Length, Area, Volume Formulas	405
59.0.12	Differentiation of Integrals with Variable Bounds	406
59.0.13	Parameter-Dependent Integrals	406
59.0.14	Leibniz Rule for Differentiating Under the Integral Sign	406
59.0.15	Improper Integrals with infinite Bounds	406
59.0.16	Improper Integrals with Discontinuity	406
59.0.17	Convergence Criteria for Improper Integrals	407
59.0.18	Integral Test for Series	407
59.0.19	Integrals with Absolute Value	407
60	Important Theorems of Single Variable Calculus	409
60.0.1	Intermediate Value Theorem	409
60.0.2	Extreme Value Theorem	410
60.0.3	Rolle's Theorem	410
60.0.4	Mean Value Theorem	411
61	Coordinate Systems	413
61.0.1	Cartesian Coordinates	413
61.0.2	Polar Coordinates	413
61.0.3	Cylindrical Coordinates	414
61.0.4	Spherical Coordinates	414
62	Parameterized Curves	415
62.0.1	Definition	415
62.0.2	Purpose	415
62.0.3	From parametric to explicit form	415
62.0.4	How to Parameterize a Function	415
62.0.5	Arclength and t Parameter	416
63	Multi-variable Functions	417
63.0.1	Open Sets	417
63.0.2	Closed Sets	417
63.0.3	Boundedness and Order	417
63.0.4	Sequences	417
63.0.5	Different Norms	418
63.0.6	Accumulation Point	419
63.0.7	Bolzano–Weierstraß Theorem	419
63.0.8	Limits	419
63.0.9	Partial Functions	420

63.0.10 Continuity	420
63.0.11 Uniform and Lipschitz Continuity	422
63.0.12 Epsilon–Delta Criterion	423
63.0.13 Fixpoints	423
63.0.14 Banach Fixed Point Theorem	423
63.0.15 Vector fields	423
64 Multi-variable Differentiation	425
64.0.1 Partial Derivatives	425
64.0.2 Differentiability	425
64.0.3 The Gradient	425
64.0.4 The Tangent Plane	426
64.0.5 The Directional Derivative	427
64.0.6 The Total Differential	428
64.0.7 The Chain Rule	428
64.0.8 Absolute and Relative Error	429
64.0.9 Implicit Differentiation	430
64.0.10 Schwarz’s Theorem (Clairaut’s Theorem)	431
64.0.11 The Jacobian	431
64.0.12 Taylor’s Theorem	432
64.0.13 Relative Extrema	432
64.0.14 Finding Candidate Points for Zeros While Seeking Relative Extrema	433
64.0.15 Lagrange Multipliers	434
64.0.16 The Tangent Vector	437
64.0.17 Linear Regression	439
65 Multidimensional Integrals	441
65.0.1 Arclength of a Curve	441
65.0.2 Line Integrals	441
65.0.3 Line Integrals with respect to a specific variable	443
65.0.4 Flow and Circulation	444
65.0.5 Green’s Theorem Circulation-Curl Form	445
65.0.6 Flux	445
65.0.7 Flux Density	446
65.0.8 Green’s Theorem Divergence-Flux Form	446
65.0.9 Conservative Vectors Fields	446
65.0.10 The Potential Function	446
65.0.11 Double Integral over a Rectangular Region	448
65.0.12 Steps for finding the Region of Integration	449
65.0.13 Integration over Curvilinear Domains	449
65.0.14 Changing the order of integration to solve tricky integrals	450
65.0.15 The Jacobian	451
65.0.16 Integration in Polar Coordinates	453
65.0.17 Improper Integrals over Unbounded Regions	454
65.0.18 Triple Integrals	454
65.0.19 Surfaces	456
65.0.20 Implicit Surface Area	459
65.0.21 Explicit Surface Area	459
65.0.22 Surface Integrals	460
65.0.23 The Orientation of a Surface	460
65.0.24 The Flux of a Surface	460
65.0.25 Curl	460
65.0.26 Stokes’s Theorem	461
65.0.27 Divergence	461
65.0.28 The Divergence Theorem	461
65.0.29 Unified View of Vector Calculus	461
66 Differential Equations	463
66.0.1 Ordinary Differential Equations	463

66.0.2	Initial Value Problems and Solutions	463
66.0.3	Volterra Integral	463
66.0.4	The Picard-Lindelöf Iteration Method	464
66.0.5	Integration case	464
66.0.6	Separation of Variables	464
66.0.7	Geometry of ODE	465
66.0.8	Existence and Uniqueness	465
66.0.9	How to deal with initial conditions	466
66.0.10	Solving DE with Substitution	466
66.0.11	Linear Differential Equations	467
66.0.12	Integrating Factor Method	468
66.0.13	Homogeneous Differential Equations	469
66.0.14	Solving Homogeneous DE I	470
66.0.15	Solving Homogeneous DE I	471
66.0.16	Inhomogeneous Equation	471
66.0.17	Solutions of LDE	471
66.0.18	Existence and Uniqueness Differential Equations	473
66.0.19	Theorem of Superposition and the General Solution	473
66.0.20	Undetermined Coefficients	474
66.0.21	Bernoulli Equation	475
66.0.22	Autonomous Equations	476
66.0.23	Linear Inhomogeneous DEs with Non-Constant Coefficients	477
66.0.24	Power Series Approach	478
66.0.25	Exact Differential Equations	479
66.0.26	Constant Coefficients ODE	481
66.0.27	The Wronskian	484
66.0.28	Linearity Property	484
66.0.29	Variation of Parameters	485
66.0.30	Which method to use depending on the type of differential equations	486
66.0.31	Linear Systems of Differential Equations	486
67	Combinatorics	489
67.0.1	Three Principles of Combinatorics	489
67.0.2	The Basic Counting Principle	489
67.0.3	Permutation	489
67.0.4	Permutation with repetition	489
67.0.5	Combination with repetition	490
67.0.6	Variation	490
67.0.7	Variation without repetition	490
67.0.8	Combination without repetition	490
67.0.9	Multinomial Coefficients	491
67.0.10	Example Problems	491
67.0.11	Disarray	491
67.0.12	Bell Numbers	491
67.0.13	Ramanujan Numbers	491
67.0.14	Stirling Numbers I	492
67.0.15	Stirling Numbers II	492
67.0.16	Lah Numbers	492
67.0.17	Euler's Numbers	492
67.0.18	Catalan Numbers	492
68	Probability	493
68.0.1	Basics	493
68.0.2	The Birthday Paradox	493
68.0.3	The Sample Space	493
68.0.4	Laplace Distribution and Experiment	493
68.0.5	Sigma Algebra	494
68.0.6	Probability Measurement	494

68.0.7	Inclusion-Exclusion Formula for Two Non-Disjoint Sets	496
68.0.8	Inclusion-Exclusion Formula for Three Non-Disjoint Sets	496
68.0.9	General Formula for the Cardinality of the union of sets	496
68.0.10	Stochastic Independence	496
68.0.11	The Product Space	498
68.0.12	The Product Space of Discrete Probability Spaces	498
68.0.13	Bernoulli-Experiments	498
68.0.14	Conditional Probability	499
68.0.15	Multistage Random Experiments	500
68.0.16	Bayes Theorem	501
68.0.17	Law of Large Numbers	502
68.0.18	Strong Law of large Numbers	502
68.0.19	Central Limit Theorem	502
68.0.20	Approximating the Binomial Distribution with the Normal Distribution	503
68.0.21	Theorem of Moivre-Laplace	503
69	Single Variable Distributions	505
69.0.1	Random Variables	505
69.0.2	Indicator Function	506
69.0.3	Distributions	506
69.0.4	Distribution Function	506
69.0.5	Cumulative Probability Calculations	507
69.0.6	The Expected Value for Discrete Random Variables	507
69.0.7	Triangle Inequality for Expected Values	507
69.0.8	Variance	508
69.0.9	Moments	508
69.0.10	Probability Density	509
69.0.11	Expected Value for Continuous Random Variables	509
69.0.12	Transformation Theorem for Densities	510
69.0.13	Quantile	510
69.0.14	The Moment Generating Function	510
69.0.15	Discrete Uniform Distribution	511
69.0.16	Continuous Uniform Distribution	511
69.0.17	Bernoulli Distribution	512
69.0.18	Binomial Distribution	512
69.0.19	Normal Distribution	512
69.0.20	Geometric Distribution	514
69.0.21	Poisson Distribution	514
69.0.22	Hyper-geometric Distribution	515
69.0.23	Exponential Distribution	516
69.0.24	Gamma Distribution	517
69.0.25	Chi-Square Distribution	517
69.0.26	Students t-Distribution	518
69.0.27	Activity Examples	518
70	Multivariable Distributions	519
70.0.1	Measurability of Random Variables and Random Vectors	519
70.0.2	Joint Distribution and Random Vectors	519
70.0.3	Joint Probability Function	520
70.0.4	Marginal Distributions of Discrete Random Variables	520
70.0.5	Multivariate Expected Value for Discrete Random Variables	521
70.0.6	Joint Probability Density Function	522
70.0.7	Multivariate Expected Value for Continuous Random Variables	523
70.0.8	Uniform Distribution on a Measurable Set	523
70.0.9	Probability over Measurable Sets	523
70.0.10	Joint Distribution Function for multiple Variables	523
70.0.11	Marginal Densities	523
70.0.12	Expectation with Respect to a Joint Density	523

70.0.13 Stochastic Independence of Random Variables	524
70.0.14 Independence of Densities	524
70.0.15 Variance for Independent Random Variables	524
70.0.16 Conditional Distribution of Discrete Random Variables	525
70.0.17 Conditional Density for Continuous Random Variables	525
70.0.18 Covariance and Correlation of Random Variables	526
70.0.19 Multivariable Normal Distribution	527
70.0.20 Example: Bivariate Normal Distribution	528
71 Statistics	531
71.0.1 Basic Nomenclature	531
71.0.2 Random Sampling	531
71.0.3 The 5 Number Summary	532
71.0.4 Sample Statistics	532
71.0.5 Finite n-Correction	532
71.0.6 Frequency	533
71.0.7 Order Statistic	533
71.0.8 Median	533
71.0.9 Empirical Quantile	534
71.0.10 Modus	534
71.0.11 Empirical Variance	534
71.0.12 Sigma Rules	534
71.0.13 Absolute Deviation and Range	534
71.0.14 Interquartile Range	535
71.0.15 Coefficient of Variation	535
71.0.16 Covariance	535
71.0.17 Pearson Correlation Coefficient	535
71.0.18 Spearman Correlation Coefficient	536
71.0.19 Empirical Distribution Function	536
71.0.20 Absolute Frequency sum	537
71.0.21 Grouped Data	537
71.0.22 Samples via Random Variables	538
71.0.23 Statistical Model	538
71.0.24 Estimators	538
71.0.25 Consistence of an Estimate	539
71.0.26 Distortion	539
71.0.27 Method of Moments Estimators	539
71.0.28 Expected Value, Standard Deviation and Variance of the Sample Mean	539
71.0.29 Sample Variance as Estimator	540
71.0.30 Normal Approximation to Distribution of the Sample Mean	540
71.0.31 Linear Regression	541
71.0.32 Monte Carlo Sampling	542
71.0.33 Likelihood	542
71.0.34 Maximum Likelihood	543
71.0.35 Confidence Interval	545
71.0.36 Confidence Interval Formal Definition	545
71.0.37 t_{n-1}	545
71.0.38 Confidence Intervals for the Expected Value and Standard Deviation of Normal Distributions	546
71.0.39 Degrees of Freedom	546
71.0.40 Hypothesis Testing	546
71.0.41 Test Statistic	547
71.0.42 Quality, Power and Level	547
71.0.43 Critical Value	548
71.0.44 Test-Statistic	548
71.0.45 p-Value	548
71.0.46 Standard Error	549
71.0.47 General Procedure for Hypothesis Testing	549

71.0.48 Hypothesis Testing on Normal Distributed Random Variables	550
71.0.49 Between-Sample t-Tests	551
71.0.50 Chi-Square-Test	552
71.0.51 Bayesian Inference	552
72 Information Theory	555
72.0.1 Surprise	555
72.0.2 Entropy	555
72.0.3 Mutual Information	555
72.0.4 Cross Entropy	555
72.0.5 KL-Divergence	556
72.0.6 Information Gain	556
72.0.7 Gini Impurity	557
73 Machine Learning	559
73.0.1 Basic Terminology	559
73.0.2 Solution Strategy	559
73.0.3 Central Learning Problems in Machine Learning	559
73.0.4 The Perceptron	560
73.0.5 Pocket Algorithm	562
73.0.6 Preparing the Data Set	562
73.0.7 Linear Models Types of Supervised Learning	563
73.0.8 Classes of Classifications	563
73.0.9 Linear Regression	563
73.0.10 Non-Linear Transformations	564
73.0.11 Approximation and Generalization	564
73.0.12 Bias and Variance	564
73.0.13 Learning Curves	565
73.0.14 Overfitting	565
73.0.15 Underfitting	565
73.0.16 The Augmented Error	565
73.0.17 Regularization	565
73.0.18 Validation	566
73.0.19 Hold-Out-Validation	566
73.0.20 Cross Validation	566
73.0.21 Leave One Out Cross Validation	567
73.0.22 Confusion Matrix	567
73.0.23 Sensitivity and Specificity	567
73.0.24 Learning Principles	567
73.0.25 Logistic Regression	567
73.0.26 Trees	569
73.0.27 Regressions Trees	570
73.0.28 Classification Trees	570
73.0.29 Gini Impurity	571
73.0.30 Information Gain in Machine Learning	571
73.0.31 CART Algorithm	573
73.0.32 Variance Reduction	573
73.0.33 Random Forest	573
73.0.34 Quantifying the Differences between Features	573
73.0.35 Nearest Neighbors	574
73.0.36 k-Nearest Neighbors	574
73.0.37 Support Vector Machine	574
73.0.38 Hard Margin SVM	575
73.0.39 Soft Margin Support Vector Machine	576
73.0.40 Non-Linear Support Vector Machines	577
73.0.41 The Radial Kernel	577
73.0.42 Least Squares Support Vector Machine	578
73.0.43 Dimension Reduction Principal Component Analysis	580

73.0.44	Kernel PCA	583
73.0.45	Kernel PCA Algorithm	586
73.0.46	Linear Discriminant Analysis	586
73.0.47	Naive Bayes	588
73.0.48	KMeans	588
73.0.49	Scaling	589
74	Numerical Mathematics	591
74.1	Floating Point Numbers	591
74.1.1	Normalized Representation	591
74.1.2	Machine Precision	592
74.1.3	Interpretation	592
74.2	Errors	592
74.2.1	Well- and Ill-Posed Problems	592
74.2.2	Error Measurement	592
74.2.3	Error Types	593
74.2.4	Error Criteria	594
74.2.5	Error Kinds	594
74.2.6	Input Error	595
74.2.7	Procedural Error	597
74.2.8	Accumulation Error	597
74.3	Error Analysis For Linear Systems	597
74.3.1	Condition of A Matrix	598
74.3.2	Inequality For Error Analysis in Linear Systems	599
74.3.3	Prager Oettli	600
74.3.4	QR-Decomposition For Numerical Mathematics	601
74.3.5	Householder Algorithm	605
74.3.6	Effect of a Householder Transformation	605
74.4	Iterative Methods	606
74.4.1	Single Step Stationary Iteration Method	607
74.4.2	Error of Single Step Stationary Iteration Method	607
74.4.3	Convergence	607
74.4.4	The Spectral Radius	607
74.4.5	Gelfand	608
74.4.6	Splitting Method For Linear Systems	608
74.4.7	The Residuum As Stopping Criterion	609
74.4.8	Convergence For Semi Positive Definite Matrices	609
74.4.9	Strict Diagonal-dominant	609
74.4.10	The Jacobi Method	609
74.4.11	Gauss-Seidel Method	610
74.4.12	After Iteration	610
74.4.13	Gradient Descent	611
74.5	Iterative Methods For Eigenvalue Problems	612
74.5.1	The Vector Iteration	612
74.5.2	The Jacobi Method	612
74.5.3	The QR Method	613
74.6	Zeros Problems	614
74.6.1	The Bisection Algorithm	614
74.6.2	The Regula-Falsi Algorithm	615
74.6.3	The Secant Algorithm	615
74.6.4	The Newton Algorithm	616
74.6.5	Local Convergence	616
74.6.6	Contraction	616
74.6.7	The Fix Point Theorem of Banach	616
74.6.8	Linear Convergence	616
74.6.9	Convergence With Order m	617
74.6.10	Convergene Of The Differentiable Iteration	617

75 Computability	619
75.0.1 Church Turing Thesis	619
75.0.2 Textual Substitution	619
75.0.3 Leibniz Inference Rule	619
75.0.4 Correctness of Algorithms	619
75.0.5 Termination	622
75.1 Random Access Machine	623
75.2 Complexity Theory	624
75.2.1 Definitions	624
75.2.2 Properties of the Complexity Classes	625
75.2.3 Master's Theorem	625
75.2.4 How To Identify Time Complexity of a Function	626
75.2.5 How to Find Time Complexity In Code	627
75.2.6 Common Time Complexities	628
75.3 State Machines and Languages	629
75.3.1 Languages and Alphabets	629
75.3.2 Inductive Definitions	629
75.3.3 Fixes	630
75.3.4 The Canonical Order	630
75.3.5 Languages	630
75.3.6 Pumping Lemma	630
75.3.7 Algorithm	630
75.3.8 The Decision Problem	631
75.3.9 Recursive Languages	631
75.3.10 Enumeration Problem	631
75.3.11 Grammars	631
75.4 State Machines	632
75.4.1 Deterministic Finite Automaton (DFA)	632
75.4.2 Non-Deterministic Finite Automaton (NFA)	632
75.4.3 Epsilon-NFA (ϵ -NFA)	633
75.4.4 How to Construct a State Machine	633
75.4.5 Regular Expressions	634
75.4.6 State Machine Conversions and Minimization	635
75.4.7 Epsilon-NFA to NFA	637
75.4.8 Minimizing State Machines	641
75.5 Regex	644
75.5.1 Basic Characters	644
75.5.2 Meta Characters (Special Characters)	644
75.5.3 Character Classes	644
75.5.4 Quantifiers	644
75.5.5 Grouping and Capturing	645
75.5.6 Assertions and Anchors	645
75.5.7 Advanced Constructs	645
75.5.8 Examples	645
75.6 Graph Theory	646
75.6.1 Connected Graphs	646
75.6.2 Paths	646
75.6.3 The Adjacency Matrix	646
75.6.4 The Relax Operation	647

Chapter 1

Guidelines for Writing Formal Mathematical Proofs

General Principles

- *Be precise:* use exact mathematical language and avoid vague statements.
- *Be concise but complete:* do not skip logical steps that a knowledgeable reader needs.
- *Explain your reasoning:* a proof is a logical explanation, not merely a chain of equations.
- *Know your audience:* assume familiarity with definitions and standard theorems, but not your personal thought process.

Structure of a Proof

- *Begin with definitions and hypotheses:* For a statement “If P , then Q ,” start with “Assume P .”
- *Direct proofs:* assume P , derive consequences, and conclude Q .
- *Contrapositive proofs:* prove $\neg Q \Rightarrow \neg P$ instead of $P \Rightarrow Q$.
- *Proof by contradiction:* assume the negation of the desired statement and derive a contradiction.
- *Universal statements:* let an arbitrary object satisfying the hypothesis and show the statement holds for it.
- *Existential statements:* construct an example or justify its existence, and verify the required properties.
- *Equality of sets:* prove $A \subseteq B$ and $B \subseteq A$.
- *Uniqueness proofs:* first show existence, then show any two such objects must be equal.

Writing Style

- *Write in complete sentences* with mathematical expressions integrated into the prose.
- *Use transition words* such as “therefore,” “hence,” and “thus” to make logical flow explicit.
- *Avoid informal symbols* such as arrows ($\rightarrow$) in prose except in formal logic contexts.
- *Cite known theorems* when they are used in the argument.
- *Keep notation consistent* throughout the proof.

Using Definitions Explicitly.

- *State relevant definitions clearly* when they are used. For example, to prove continuity at a , one begins: “Let $\varepsilon > 0$. We must find $\delta > 0$ such that ...”

Conclusion.

- *End with a clear concluding statement* such as “Therefore, Q .” or “This completes the proof.”

Common Mistakes to Avoid.

- *Skipping logical steps* or jumping to conclusions.
- *Using examples instead of general arguments.*
- *Assuming the statement* you are trying to prove.
- *Including irrelevant details.*
- *Using ambiguous language* like “clearly” or “obviously” unless truly justified.

Template.

Proof.

[State assumptions clearly.]

[Develop argument using definitions and known theorems.]

[Use clear logical reasoning to reach the conclusion.]

Therefore, [claim to be proved].

\qed

Chapter 2

Logic

2.1 Propositional Logic

Propositional logic is also called Boolean logic as it works on 0 and 1. In propositional logic, we use symbolic variables to represent the logic, and we can use any symbol for a representing a proposition, such p, q, r, s, t , etc. Propositions can be either *true* (1, T) or *false* (0, F) but it cannot be both.

2.1.1 Predicates

It is statement which contains variables linked by some property stated in the language of formal logic whose value can be true or false.

2.1.2 Predicate Substitution

Given a predicate P on the variable v we can substitute every free instance of v with the term v to extend our predicate to another variable space.

$$[t/v]P$$

Types of substitution:

- Full replacement: $[y/x]\forall z((z = x) \wedge (y \in x))$, converted to $\forall z((z = y) \wedge (y \in y))$,
- Partial replacement: $[y/x]x = y \wedge \forall x(x \in y)$, the x of $\forall x(x \in y)$ is not a free variable, hence it can not be replaced completely, just $y = y \wedge \forall x(x \in y)$

Example:

Prove $\forall a(a = a)$:

Let $P := x = a$ be a predicate in x and a . We apply:

$$[a/x](x = a) \text{ to } a = a, \text{ then by extension of our predicate } \forall a(a = a).$$

QED

Prove $\forall a \forall b(a = b \implies b = a)$:

Let $P := x = a$ be a predicate in a and x . We apply:

$$[b/x](x = a) \text{ to } b = a, \text{ then by extension of our predicate } \forall a \forall (a = b \implies b = a).$$

QED

Prove $\forall a(a = b \wedge b = c \implies c = a)$:

We know $a = b \wedge b = a$, let this be our predicate with x and apply the substitution $[c/b]$, which gives us $a = x$, $a = b$ and finally $a = c$.

QED

2.1.3 Logic Operators and Truth Tables

NOT ($\neg$, $\sim$)

p	$\neg p$
0	1
1	0

IFF ($\iff$)

p	q	$p \iff q$
0	0	1
0	1	0
1	0	0
1	1	1

AND ($\wedge$)

p	q	$p \wedge q$
0	0	0
0	1	0
1	0	0
1	1	1

XOR ($\oplus$)

p	q	$p \oplus q$
0	0	0
0	1	1
1	0	1
1	1	0

OR ($\vee$)

p	q	$p \vee q$
0	0	0
0	1	1
1	0	1
1	1	1

NOR ($\downarrow$)

p	q	$p \downarrow q$
0	0	1
0	1	0
1	0	0
1	1	0

IMPLIES ($\implies$)

p	q	$p \implies q$
0	0	1
0	1	1
1	0	0
1	1	1

NAND ($\uparrow$)

p	q	$p \uparrow q$
0	0	1
0	1	1
1	0	1
1	1	0

Implications

As a side note for the *implication* we can think about it as *if, then*. For the cases false, true; and false false; we call them *vacuously* true. From a false premise can come everything out.

2.1.4 Tautology and Contradiction

Tautology: A logical formula that is always true.

Common tautologies

- $p \implies q \vee p$

- $q \implies p \vee q$
- $p \wedge q \implies p$
- $q \wedge q \implies q$
- $[(p \implies q) \wedge (q \implies r)] \implies (p \implies r)$

Contradiction: A formula that is always false.

2.1.5 Logical Equivalences

Commutative Laws

$$p \wedge q \iff q \wedge p \quad p \vee q \iff q \vee p$$

Associative Laws

$$(p \wedge q) \wedge r \iff p \wedge (q \wedge r) \quad (p \vee q) \vee r \iff p \vee (q \vee r)$$

Distributive Laws

$$p \wedge (q \vee r) \iff (p \wedge q) \vee (p \wedge r) \quad p \vee (q \wedge r) \iff (p \vee q) \wedge (p \vee r)$$

Identity Laws

$$p \wedge \mathbf{T} \iff p \quad p \vee \mathbf{F} \iff p$$

Negation Laws

$$p \vee \sim p \iff \mathbf{T} \quad p \wedge \sim p \iff \mathbf{F}$$

Double Negation Law

$$\sim(\sim p) \iff p$$

Idempotent Laws

$$p \wedge p \iff p \quad p \vee p \iff p$$

Universal Bound Laws

$$p \vee \mathbf{T} \iff \mathbf{T} \quad p \wedge \mathbf{F} \iff \mathbf{F}$$

De Morgan's Laws

$$\sim(p \wedge q) \iff (\sim p) \vee (\sim q) \quad \sim(p \vee q) \iff (\sim p) \wedge (\sim q)$$

Absorption Laws

$$p \vee (p \wedge q) \iff p \quad p \wedge (p \vee q) \iff p$$

Conditional Laws

$$(p \implies q) \iff (\sim p \vee q) \quad \sim(p \implies q) \iff (p \wedge \sim q)$$

Complement Law

$$p \vee \neg p \iff \mathbf{T} \quad p \wedge \neg p \iff \mathbf{F}$$

Biconditional Law

$$p \iff q \iff (p \implies q) \wedge (q \implies p)$$

Transitivity Law

$$(p \implies q) \wedge (q \implies r) \implies (p \implies r)$$

Indirect Proof (Contrapositive)

$$(p \implies q) \iff (\neg q \implies \neg p)$$

Disjunctive Syllogism (Disjunctive Exclusion)

$$p \nabla q \equiv (p \vee q) \wedge \neg p \implies q$$

$$p \nabla q \equiv (p \wedge q) \vee \neg(p \wedge q)$$

2.1.6 Truth Tables

Truth tables are a fundamental tool in logic that systematically show the truth value (true or false) of a compound statement for every possible combination of the truth values of its individual component statements.

Essentially, they lay out all the scenarios and the resulting truth of the overall logical expression. This helps determine if an argument is valid, if statements are logically equivalent, or the circumstances under which a complex statement is true or false.

Example:

p	q	$\neg p$	$\neg q$	$\neg p \implies q$	$(\neg p \implies q) \wedge \neg p$	$[(\neg p \implies q) \wedge \neg p] \Rightarrow q$
T	T	F	F	T	F	T
T	F	F	T	T	F	T
F	T	T	F	T	T	T
F	F	T	T	F	F	T

Filling a truth table

To fill a truth table for a logical expression with truth values (True or False), you follow a specific order for the input variables. This order ensures that all possible combinations of truth values for the variables are covered.

General Procedure:

1. **List all possible combinations of truth values for the input variables:** If you have n variables, the number of rows in the truth table will be 2^n . Each variable can be either True (T) or False (F).
2. **Order of the input variables:**
 - Start by filling in the truth values for the first variable. It alternates between True and False every 2^{n-1} rows.
 - Then for the second variable, it alternates every 2^{n-2} rows, and so on.
 - In short: the first variable alternates every other row, the second variable every two rows, the third every four rows, and so on.

Example:

For 3 variables, there are $2^3 = 8$ possible combinations of truth values. The truth values are filled in the following order:

p	q	r	Expression Result
T	T	T	—
T	T	F	—
T	F	T	—
T	F	F	—
F	T	T	—
F	T	F	—
F	F	T	—
F	F	F	—

The pattern for filling the input truth values

- The first column p alternates every 4 rows: ‘T, T, F, F, T, T, F, F’.
- The second column q alternates every 2 rows: ‘T, T, F, F, T, T, F, F’.
- The third column r alternates every row: ‘T, F, T, F, T, F, T, F’.

This ensures that all combinations of p , q , and r are covered, and you can then evaluate the logical expression for each combination.

Truth Table for the Expression $(p \wedge q) \vee q$

p	q	r	$(p \wedge q) \vee r$
T	T	T	T
T	T	F	T
T	F	T	T
T	F	F	F
F	T	T	T
F	T	F	F
F	F	T	T
F	F	F	F

2.1.7 Disjunctive Normal Form (DNF)

Disjunctive Normal Form (DNF) is a standard way of writing a logical expression as a *disjunction(OR)* of *conjunctions (ANDs)*. A DNF expression consists of a series of conjunctions of literals, where each conjunction is connected by disjunctions.

To find the DNF in a truth table take the rows of the final result where there are true statements and bind the propositions that generated it with an AND inside parenthesis. Repeat it with each of true rows and connect all parenthesis with OR’s

Example of DNF:

Consider the logical expression:

$$(p \wedge q) \vee (\neg p \wedge r) \vee (q \wedge \neg r)$$

This is in DNF because it is a disjunction (OR) of conjunctions (ANDs) of literals.

2.1.8 Conjunctive Normal Form (CNF)

Conjunctive Normal Form (CNF) is a standard way of writing a logical expression as a *conjunction (AND)* of *disjunctions (ORs)*. A CNF expression consists of a series of disjunctions of literals, where each disjunction is connected by conjunctions.

To find the CNF proceed just as the DNF but with the false rows and instead of ANDs inside the parenthesis use OR and connect the terms with OR. Also add a negation before each parenthesis.

Example of CNF:

Consider the logical expression:

$$\neg(p \vee q) \wedge \neg(\neg p \vee r) \wedge \neg(q \vee \neg r)$$

This is in CNF because it is a conjunction (AND) of disjunctions (ORs) of literals.

2.1.9 Basis

A *Basis* in the field of logic is set of operations with which you can represent all possible logic operations.

Example: The set $\{\neg, \wedge\}$

2.1.10 Karnaugh Maps

Karnaugh Maps (K-Maps) are a graphical method used to simplify Boolean expressions. The main goal of a K-map is to group adjacent cells that contain 1's in order to simplify the expression. A K-map helps identify common terms, allowing the Boolean expression to be reduced to its simplest form.

Karnaugh Map for Two Variables

Consider the Boolean expression $(A \vee (B \wedge \neg A \wedge \neg B))$.

We first construct a K-map for two variables, A and B . The truth table for this expression gives the following values:

A	B	$(A \vee (B \wedge \neg A \wedge \neg B))$
0	0	0
0	1	1
1	0	1
1	1	1

The corresponding K-map is:

AB	00	01	11	10
Value	0	1	1	1

Here, we group the ones together to simplify the Boolean expression. The simplified expression is:

$$A \vee B$$

Karnaugh Map for Three Variables

Now, let's consider the expression $\neg C$. This expression only depends on one variable, but for illustration, we will use a 3-variable K-map with variables A , B , and C .

The truth table for $\neg C$ is as follows:

A	B	C	$\neg C$
0	0	0	1
0	0	1	0
0	1	0	1
0	1	1	0
1	0	0	1
1	0	1	0
1	1	0	1
1	1	1	0

The corresponding K-map for three variables A , B , and C is:

$AB \backslash C$	0	1
00	1	0
01	1	0
11	1	0
10	1	0

We see that the ones are grouped in a column, leading to the simplified Boolean expression:

$$\neg C$$

Solving a Karnaugh Map

To solve a Karnaugh Map and simplify a Boolean expression, follow these steps:

1. **Determine the Number of Variables:** Decide how many variables are in the Boolean function. This determines the size of the K-Map:
 - 2 variables: 2×2
 - 3 variables: 2×4
 - 4 variables: 4×4
 - etc.
2. **Fill in the K-Map:** Place 1's in the cells that correspond to the min-terms (where the function outputs 1). You may also include do not care conditions (usually denoted as X).
3. **Group the 1's:** Form groups (called *implicants*) of 1's. The groups must follow these rules:
 - Each group must contain 1, 2, 4, 8, ... (powers of 2) 1's.
 - Groups must be rectangular (e.g., 1×2 , 2×2).
 - Groups can wrap around the edges of the K-Map.
 - Try to form the largest groups possible to simplify the expression.
 - Each 1 should be included in at least one group.
4. **Write the Simplified Expression:**
For each group:
 - Identify the variables that are constant (either always 0 or always 1) across the group.
 - Write a product term (AND) using only the constant variables.
 - Combine all product terms with OR operations to get the final simplified SOP (Sum of Products) expression.

2.1.11 Mathematical Quantifiers

Universal Quantifier $\forall$

Means “for all” or “for every”.

Negation: $\neg(\forall x) P(x) \equiv (\exists x) \neg P(x)$ “Not all” is the same as “There exists one that does not”.

Example:

$\forall x \in \mathbb{R}, x^2 \geq 0$ For all real numbers, the square is greater than or equal to zero.

Negated Example:

$\exists x \in \mathbb{R}, x^2 < 0$ There exists a real number whose square is less than zero — this is false.

Existential Quantifier $\exists$

Means “there exists at least one”.

Negation: $\neg(\exists x) P(x) \equiv (\forall x) \neg P(x)$ “There does not exist” is the same as “For all, not”.

Example:

$\exists x \in \mathbb{N}, x > 10$ There exists a natural number greater than 10.

Negated Example:

$\forall x \in \mathbb{N}, x \leq 10$ All natural numbers are less than or equal to 10 — this is false.

Unique Existential Quantifier $\exists!$

Means “there exists exactly one”.

Negation: “Not exactly one” means: $\neg(\exists! x) P(x) \equiv (\forall x) \neg P(x) \vee (\exists x_1 \neq x_2) P(x_1) \wedge P(x_2)$ Either no such x exists, or more than one does.

Example:

$\exists! x \in \mathbb{R}, x + 5 = 0$ There exists exactly one real number such that $x + 5 = 0$.

Negated Example:

$\exists x_1 \neq x_2 \in \mathbb{R}, x_1^2 = 4 \wedge x_2^2 = 4$ There are multiple solutions to $x^2 = 4$.

2.1.12 Complex Statements

=

Chapter 3

Set Theory

Set theory is a foundational branch of mathematics that studies *sets*, which are simply collections of objects. At its core, it deals with the fundamental concepts of membership (whether an object belongs to a set), equality (when two sets are the same), and relationships between sets (like subsets, intersections, and unions).

It might seem simple, but set theory provides the basic language and tools to define and reason about almost all mathematical objects, from numbers and functions to more complex structures. It helps us understand the concept of infinity, organize mathematical ideas logically, and resolve paradoxes that arise from dealing with collections.

In essence, set theory provides the building blocks upon which much of modern mathematics is constructed.

Note, that I will start with sets as such instead of going through classes and then exporting some of their properties to sets. Classes will only be handled shortly with the axioms and some proofs but some of their properties will be directed presented as set properties.

3.0.1 Basics

To be concise, a set is a collection of mathematical objects that can also be other sets. Here is a list of common symbols for set theory.

- $A \subseteq B$: Set A is a subset of B . This means every element of A is also in B .
- $A \subseteq B$ or $A = B$: This notation already includes the possibility that A equals B since a set is always a subset of itself.
- $A \cup B$: The union of sets A and B . It includes all elements that are in A , in B , or in both.
- $A \cap B = A \setminus (A \setminus B)$: The intersection of sets A and B . It includes only the elements that are in both sets.
- $A \setminus B$: The difference of sets. Elements in A that are not in B .
- A^c or $\overline{A}$: The complement of set A . All elements not in A , relative to a universal set.
- $|A|$: The cardinality of set A , which is the number of elements in the set.


Visuals

1. $\emptyset$




1. Empty Set

2. $\{1, 2\}$




2. Set with Two Elements

3. $A \subseteq B$




3. A is a subset of B


5. $A \cup B$




5. Union of A and B



6. Intersection of A and B



7. A minus B



8. Complement of A

Short History Lesson

When set theory was invented back then by Cantor, he had no formal axioms but a set of basic 'common sense' rules he assumed when creating his theory. This choice led to a famous paradox known as *Russell's Paradox*. The paradox states that: "Given a set S which is the set containing all sets which do not contain themselves. The S does not contain itself, but if it does not contain itself it must contain itself". In short $S = \{x : x \notin x\}$, this is a so-called *logical* paradox. This is of course a huge contradiction in the hearth of naive set theory. The choice to take a more formal and axiomatic approach was to eliminate such fundamental faults.

As a short teasing we will declare the *Axiom of Selection*: $\{x \in A : S(x)\}$ this means that we can construct a set of elements of an existing set as long as they satisfy a condition. Now the paradox of $x : x \notin x$ can not be state directly in our new system. We can emulate it by declaring

$$S = \{x \in A \mid x \notin x\}$$

By analyzing the expression we get that, if $S \notin S$ then $S \in S \wedge S \in A$ which will be a contradiction, therefore, $S \notin A$.

Continuing this small historic section, we will also need to limit the power of our condition of statements $S(x)$ to get rid of another type of paradoxes considered as *semantic*.

For example, we could define the following set:

$$\{x \in \mathbb{N} : x \text{ can be described in less than 20 words of the English language}\}$$

Note that this set would need to be finite because it there is only a limited amount of combination of words in the English language. But we could for example say: 'the natural number which can not be described in less than 20 words' this number. This is again an example of an element in a set if and only if it is not in the set. To formalize our conditions we

3.0.2 Sets and Classes

One important distinction is the system of Neumann is notion difference between a *set* and *class*. Classes can not be contained inside sets, while sets can. Formally:

If $X \in Y$ for some class Y then X is a set. If $X \notin Y$ for any class Y then X is a class.

3.0.3 Axioms of Set Theory

I. *Axiom of Selection*: Given a class A , a class S can be build in the following manner $S = \{x \in A \mid P(x)\}$

II. *Axiom of Extensionality*: Let A and B be classes then $A = B \iff \forall X : A \in X \implies B \in X \wedge B \in X \implies A \in X$

- III. *Empty-Set Axiom*: $\emptyset$ is a set.
- IV. *Axiom of Class Construction*: Let $P(x)$ be a statement about x which can be expressed in terms of $\in, \wedge, \vee, \neg, \implies, \exists, \forall$ with variables $x, y, z, \dots$ then there exists a class C whose elements consist of all the elements x which satisfy $P(x)$. $\{x : P(x)\}$
- V. *Axiom of Subset*: Every subclass of a set is a set.
- VI. *Axiom of Paring*: If a and b are sets, then $\{a, b\}$ is a set
- VII. *Axiom of Union*: If A is a set of sets, then $\bigcup_{a \in A} a$ is a set. $\forall a \exists b \forall c (c \in a \implies \forall d (d \in c \implies d \in b))$.
- VIII. *Axiom of the Power Set*: If A is a set then the set of all the subsets of the original set, $\mathcal{P}(A)$, is a set. $\forall a \exists b \forall c (\forall d (f \in c \implies d \in a) \implies d \in b)$.
- IX. *Axiom of Foundation / Regularity*: If A is any set, there is an element $a \in A$ such that $a \cap A = \emptyset$. Alternatively, there exists not such set, which is formed from a chain like: $a_1 \in a_2 \in a_3 \in \dots$
 $\forall a (\exists b (b \in a) \implies \exists c (c \in a \wedge \forall d (d \in c \implies d \notin a)))$
- X. *Axiom of Replacement*: If A is a set and $f : A \rightarrow B$ is a surjective function, then B is a set. Let P be a predicate on a, b, c and let d, e, f, g not be yet introduced,

$$\begin{aligned} \forall a \exists b ((\forall c ((c \in a \wedge \exists d P) \implies \exists e (e \in b \wedge [e/d]P)) \\ \wedge \forall f (f \in b \implies \exists g (g \in a \wedge [g/c][f/d]P))) \end{aligned}$$

- XI. *Axiom of Infinity*: An infinity set exists. $\exists a (\exists b (b \in a) \wedge \forall c (c \in a \implies \exists d (d \in a \wedge c \in d)))$.

- XII. *Axiom of Choice*: Given a non-empty set, $A_1, A_2, \dots$, there is a set containing exactly one element from each A_i .

$$\begin{aligned} \forall a (\forall b (b \in a \implies (\exists c (c \in b) \\ \wedge \forall d ((d \in a \wedge \exists e (e \in b \wedge e \in d)) \implies b = d))) \\ \implies \exists f \forall g (g \in a \implies \exists h ((h \in g \wedge h \in f) \\ \wedge \forall i ((i \in g \wedge i \in f) \implies i = h)))) \end{aligned}$$

Proofs Using the Axioms

- $A = A$ and also $A = B \implies B = A$: The statements is trivial by the first axiom
- $A \subseteq B \wedge B \subseteq A \implies A = B$: $(\forall x \in A \implies x \in B) \wedge (\forall x \in B \implies x \in A)$ Let us suppose for the sake of contradiction that $A \neq B$ then $\exists x \in A : x \notin B$ or the other way around. Note that this statement contradicts our initial conditions. Thus, the statement is true.
- $A \subseteq B \wedge B \subseteq C \implies A \subseteq C$: $(\forall x \in A \implies x \in B) \wedge (\forall x \in B \implies x \in C)$ By a similar manner let us assume that A is not a subset of C this means that for some x the condition holds that $x \in A \wedge x \notin B$ because else it would be in C . Note that because $A \subseteq B$ such an x can not exist due to our initial condition and also $B \subseteq A$ for that case. Thus $A \subseteq C$
- $\emptyset \subseteq A$: By definition $x \in \emptyset \implies x \in A$, by contrapositive $x \notin A \implies x \notin \emptyset$. This is true because of the contrapositive being true.
- Given a proper class $A \subseteq B$ prove that B is a proper class and that the union of two proper classes is also a proper class.

Given $A \subseteq B$, if B would be not be a proper class then by the property which states that every subclass of a class is a set we would have a contradiction. Therefore, B is a proper class.

$A \cap C$ where A and C are proper classes, gives us that their union is also a proper class due to $A, C \subseteq A \cup C$

- $\mathcal{P}(A) \not\subseteq A$: Let $x \in \mathcal{P}(a)$, then $\forall x \in \mathcal{P}(A) \implies x \in A$. By definition, $A \in \mathcal{P}(A)$, but $A \notin A$; hence we have contradiction and the power set of a set is not a subset of the original set.

ΩεΩ

3.0.4 Consequences of the Axiom of Replacement

- Every singleton is a set.
- If A is in a one-to-one correspondence with B , then B is a set.
- If $\{A_i\}_{i \in I}$ is an indexed family of sets and I is a set, then $\{A_i \mid i \in I\}$ is a set.

3.0.5 Subclasses/Subsets

A subset/subclass A of the set/class B is a set which obeys the following property:

$$A \subseteq B \iff \forall x \in A, x \in B \implies x \in B$$

3.0.6 The Universal Class

By the axiom of construction we define $\mathcal{U}$ with $P(x)$ is $x = x$ as the *universal class* of all elements.

It also has the following property

$$A \subseteq \mathcal{U},$$

which gives us the following: $A \cup \mathcal{U} = \mathcal{U}$ and $A \cap \mathcal{U} = A$ for any class or set A .

3.0.7 The Empty Class

The *empty class* is class with no elements.

$$\emptyset = \{x \mid x \neq x\}$$

We can not use existence quantifiers $\exists$ with this class. Every statement with $\forall x \in \emptyset$ is true, because there are no counterexamples. This leads to interesting equalities like $\mathcal{U} = \emptyset$.

We can prove that the empty class exist by letting $\emptyset = \{x \mid x \neq x\}$ this is a perfectly valid statement.

Now we need to prove that it is unique, and this is given by the axiom of extensionality.

□◻◻

3.0.8 The Empty Set

The set that contains no elements $\emptyset$. It is the unique set with zero elements. It inherits the properties of its class counterpart.

It also has the property that $\emptyset \subseteq A$ for any class or set A .

3.0.9 Set Complement Proof

Given $A, B \subseteq M$ then $A \subseteq B \implies B^c \subseteq A^c$

For some $x \in B^c$ we know that $x \notin B$. If $x \in A$ then $x \in B$ which is a contradiction. Hence, $x \notin A$ therefore, for every $x \in B^c$ we conclude $B^c \subseteq A^c$.

□◻◻

3.0.10 Proof The Power Set of a Set is not an Element of the Set

Let X be a set and $Y = \{x \in X \mid x \notin x\}$.

This is a subset of X by definition and implies $Y \in \mathcal{P}(X)$.

Assume $\mathcal{P}(X) \subseteq X$ then $Y \subseteq X, Y \in X$ but

$$Y \in Y \implies Y \notin Y \implies Y \in Y \text{ which is a contradiction}$$

Because our assumption was proven false, then $\mathcal{P}(X)$ is not a subset of X .

□◻

3.0.11 Proof of the Union of Subsets

Let A and B be sets then:

- $A \subseteq B$ if and only if $A \cup B = B$
- $A \subseteq B$ if and only if $A \cap B = A$

Proof:

For $A \subseteq B \iff A \cup B = B$ we know that $x \in A \implies x \in B$ Then

$$\begin{aligned} A \cup B &\iff x \in A \vee x \in B \quad \text{as subset then} \\ &\iff x \in B \vee x \in B \\ &\iff x \in B = B \end{aligned}$$

The other direction will be skipped due to it being very similar to the $\implies$ direction.

Similarly, for $A \subseteq B \iff A \cap B = A$ We know that for some $(y \in B \wedge y \notin A) \notin A \cap B$ because of the definition of the subset. Then this tells us that $\forall x \in A \cap B : x \in A$ thus $A \cap B = A$

□◻

Cut of Set and Complement

$$A \cap B^c = \emptyset \iff A \subseteq B$$

$\implies$:

$$\begin{aligned} A \cap B^c = \emptyset &\implies \nexists x \in A \wedge x \notin B \\ &\implies \forall x \in A : x \in B \\ &\iff A \subseteq B \end{aligned}$$

$\Leftarrow$: Let us assume that A is not a subset of B then

$$\begin{aligned} \exists x \in A \wedge x \notin B &\implies x \in A \wedge x \in B^c \\ &\implies x \in A \cap B^c \implies A \cap B^c \neq \emptyset \end{aligned}$$

□◻

3.0.12 The Singleton and Doubleton

We define the *singleton* as

$$\{a\} := \{x \mid x = a\},$$

and the *doubleton* as

$$\{a, b\} := \{x \mid x = a \vee x = b\},$$

Properties

- $\{a\} = \{b\} \iff a = b$
- $\{a, b\} = \{a\} \iff a = b$
- $\{a, b\} = \{x, y\} \iff (a = x \wedge b = y) \vee (a = y \wedge b = x)$

Proofs:

$\{a\} = \{b\} \iff a = b$, by the axiom of the equality of sets if

$$\exists x \in \{a\} : x \notin \{b\} \implies \{a\} \neq \{b\},$$

which is a contradiction to our condition thus $a = b$. Similarly, we can use the same argument for the other direction of the proof.

□◻◻

$\{a, b\} = \{a\} \iff a = b$ can be proven using the definition and the equivalence $P \vee P = P$

$$\{a, b\} := \{x \mid x = a \vee x = b\} \implies \{x \mid x = a\}$$

□◻◻

$\{a, b\} = \{x, y\} \iff (a = x \wedge b = y) \vee (a = y \wedge b = x)$, we need to first the case $a = b$. In that case our problems simplifies to $\{a\} = \{x, y\}$ which by the axiom guarantees the equality for $\{a\} = \{x\} \vee \{a\} = \{y\}$ because otherwise we will arrive at a contradiction.

For the case $a \neq b$ we have the following cases $(x = a \wedge y = b) \vee (x = b \wedge y = a)$ which against by the use of the axiom of the equality we can discard the cases where the equality does not hold.

□◻◻

3.0.13 Ordered Pair

Let A and B be two non-empty sets and $m \in A, n \in B$. We define

$$(m, n) := \{\{m, n\}, m\}$$

and

$$(n, m) := \{\{m, n\}, n\}$$

as ordered pairs which are elements of $A \times B$ and $B \times A$ respectively.

3.0.14 The Cartesian Product

The Cartesian product of two sets A and B , written $A \times B$, is the set of all ordered pairs in which the first element belongs to A and the second belongs to B :

$$A \times B = \{(a, b) : a \in A, b \in B\}$$

Example:

Table 3.1: Cartesian Product of $A = \{1, 2, 3\}$ and $B = \{4, 5, 6\}$

$A \times B$	$b \in B$		
	4	5	6
$a \in A$			
1	(1, 4)	(1, 5)	(1, 6)
2	(2, 4)	(2, 5)	(2, 6)
3	(3, 4)	(3, 5)	(3, 6)

The general Cartesian product of n sets can be written as:

$$X_{i=1}^{n+1} A_i = (X_{i=1}^n A_i) \times A_{n+1} \quad \text{with} \quad X_{i=1}^1 A_i = A_1$$

When $A_i = M$ for all i :

$$M^n := M \times M \times \cdots \times M = X_{i=1}^n M \quad \text{with} \quad M^1 = M$$

Distributive Law for the Cartesian Product

Let $A, A' \subseteq X$ and $B, B' \subseteq Y$. Then

$$(A \times B) \cap (A' \times B') = (A \cap A') \times (B \cap B').$$

We prove the equality by showing both inclusions.

1. Direction: $(A \times B) \cap (A' \times B') \subseteq (A \cap A') \times (B \cap B')$.

$$\begin{aligned} &\text{Let } (x, y) \in (A \times B) \cap (A' \times B'). \\ &\iff (x, y) \in A \times B \text{ and } (x, y) \in A' \times B' \\ &\iff x \in A, y \in B \text{ and } x \in A', y \in B' \\ &\iff x \in A \cap A' \text{ and } y \in B \cap B' \\ &\iff (x, y) \in (A \cap A') \times (B \cap B'). \end{aligned}$$

2. Direction: $(A \cap A') \times (B \cap B') \subseteq (A \times B) \cap (A' \times B')$.

$$\begin{aligned} &\text{Let } (x, y) \in (A \cap A') \times (B \cap B'). \\ &\iff x \in A \cap A' \text{ and } y \in B \cap B' \\ &\iff x \in A, x \in A' \text{ and } y \in B, y \in B' \\ &\iff (x, y) \in A \times B \text{ and } (x, y) \in A' \times B' \\ &\iff (x, y) \in (A \times B) \cap (A' \times B'). \end{aligned}$$

Thus, both inclusions hold and the equality follows.

□◻◻

Proof that the Cartesian Product is a Set

We need to prove that $A \times B \subseteq \mathcal{P}(A \cup B)$. By the definition of the ordered pair we know that for $x \in A \cup B \implies \{x\} \subseteq A \cup B$ the same for $x, y \in A \cup B$ which implies $\{x, y\} \subseteq A \cup B$. By our axiom of union we know that $\{\{x\}, \{x, y\}\}$ is a set and this is a subset of $\mathcal{P}(\mathcal{P}(A \cup B))$ for free x and y . Hence

$$(x, y) \subseteq A \times B \subseteq \mathcal{P}(\mathcal{P}(A \cup B))$$

□◻◻

Proof the empty set of the Cartesian Product

$A \times B = \emptyset \iff A = \emptyset \vee B = \emptyset$. We will prove the negation of the statement which logically equivalent.

$$A \times B \neq \emptyset \iff A \neq \emptyset \wedge B \neq \emptyset$$

$\implies$:

$$\begin{aligned}
A \times B \neq \emptyset &\implies \exists (x, y) \in A \times B : x \in A \wedge y \in B \\
&\implies \exists x \in A \wedge \exists y \in B \\
&\implies A \neq \emptyset \wedge B \neq \emptyset
\end{aligned}$$

$\Leftarrow$:

$$\begin{aligned}
A \neq \emptyset \wedge B \neq \emptyset &\implies \exists x \in A \wedge \exists y \in B \\
&\implies A \times B \neq \emptyset
\end{aligned}$$

□◻◻

3.0.15 Laws of Set Algebra

Let X be the universal set and $A, B, C \subseteq X$.

- $\emptyset \subseteq A$
- $A \subseteq B \iff A \cap B = A \iff A \cup B = B \iff X \setminus B \subseteq X \setminus A \iff B \subseteq A$
- $A \cup B = B \cup A$
- $A \cap B = B \cap A$
- $(A \cup B) \cup C = A \cup (B \cup C)$
- $(A \cap B) \cap C = A \cap (B \cap C)$
- $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- $A \cup A = A$ and $A \cap A = A$
- $A \setminus B = A \cap (X \setminus B) = A \cap \overline{B}$
- $B = \overline{A} \iff (A \cup B = X \wedge A \cap B = \emptyset)$
- $\overline{A} \cap \overline{B} = \overline{A \cup B}$
- $\overline{A} \cup \overline{B} = \overline{A \cap B}$
- $\overline{\overline{A}} = A$
- $(A \setminus B) \cup (B \setminus A) = A \Delta B$
- $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$
- $A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$

Proof of the Distributive Law

$$\begin{aligned}
&\text{Let } x \in A \cap (B \cup C). \\
&\iff x \in A \text{ and } x \in B \cup C \\
&\iff x \in A \text{ and } (x \in B \text{ or } x \in C) \\
&\iff (x \in A \text{ and } x \in B) \text{ or } (x \in A \text{ and } x \in C) \\
&\iff x \in A \cap B \text{ or } x \in A \cap C \\
&\iff x \in (A \cap B) \cup (A \cap C).
\end{aligned}$$

Thus,

$$A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C).$$

Conversely, let $x \in (A \cap B) \cup (A \cap C)$.

$$\iff x \in A \cap B \text{ or } x \in A \cap C$$

$$\iff (x \in A \text{ and } x \in B) \text{ or } (x \in A \text{ and } x \in C)$$

$$\iff x \in A \text{ and } (x \in B \text{ or } x \in C)$$

$$\iff x \in A \text{ and } x \in B \cup C$$

$$\iff x \in A \cap (B \cup C).$$

Thus,

$$(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C).$$

□◻

3.0.16 Proof of De Morgans's Law for sets and logic

The complement of $A \cup B$ is $\overline{(A \cup B)}$, and the law on disjoint decomposition states:

$$B = \overline{A} \iff (A \cup B = X) \wedge (A \cap B = \emptyset)$$

So define $\overline{C} := A \cup B$ and $D := \overline{A} \cap \overline{B}$, and use Law (11) to show the disjoint decomposition:

$$D = C \iff A \cap B = A \cup B$$

To show:

$$D \cup C = X \iff (\overline{A} \cap \overline{B}) \cup (A \cup B) = X$$

$$\begin{aligned} (\overline{A} \cap \overline{B}) \cup (A \cup B) &= (\overline{A} \cup A \cup B) \cap (\overline{B} \cup A \cup B) \quad (\text{Law (8)}) \\ &= (X \cup B) \cap (X \cup A) \\ &= X \cap X \\ &= X \end{aligned}$$

To show:

$$D \cap C = \emptyset \iff (\overline{A} \cap \overline{B}) \cap (A \cup B) = \emptyset$$

$$\begin{aligned} (\overline{A} \cap \overline{B}) \cap (A \cup B) &= (A \cap B \cap A) \cup (A \cap B \cap B) \quad (\text{Law (7)}) \\ &= (\overline{A} \cap A \cap \overline{B}) \cup (\overline{A} \cap \overline{B} \cap B) \\ &= (\emptyset \cap \overline{B}) \cup (\overline{A} \cap \emptyset) \\ &= \emptyset \cup \emptyset \\ &= \emptyset \end{aligned}$$

□◻

3.0.17 Indexed Sets

Let X be a set, and $A_i \subseteq X$ for all $i \in J$, where I is the *index set*.

If $I = \{1, 2, \dots, n\}$ then we define

$$\{A_i\}_{i \in I},$$

as the set of all classes/sets for which an index in I exists.

We also can define the graph of the sets with their respective indexes:

$$G_{A_i} = A_i \times I = \{(i, A_i) \mid i \in I \wedge A_i \in \{A_i\}_{i \in I}\}$$

3.0.18 Generalized Union, Intersection and Cartesian Product for Indexed Sets

For a class of classes or sets of sets with an index set I we define the generalized *union*, *intersection* and *Cartesian product* as:

$$\bigcup_{i=1}^n A_i := A_1 \cup A_2 \cup \dots \cup A_n = \{x \mid \exists i \in I, (x \in A_i)\}$$

$$\bigcap_{i=1}^n A_i := A_1 \cap A_2 \cap \dots \cap A_n = \{x \mid \forall i \in I, (x \in A_i)\}$$

$$X_{i=1}^n A_i = \{(a_1, \dots, a_n) \mid a_i \in A_i\}$$

3.0.19 Pairwise Disjoint

If I is any set, then $(A_i)_{i \in I}$ are *pairwise disjoint* if and only if:

$$\forall i_1, i_2 \in I, i_1 \neq i_2 \implies A_{i_1} \cap A_{i_2} = \emptyset$$

3.0.20 Disjoint Decomposition

If I is any set, then $(A_i)_{i \in I}$ forms a *disjoint decomposition* of X if and only if:

$$(A_i)_{i \in J} \text{ are pairwise disjoint and } \bigcup_{i \in J} A_i = X$$

3.0.21 Cardinality Laws

Let $A_i, B_j \subseteq X$ for $i \in I$ and $j \in J$. Then the following holds:

- *De Morgan's Laws:*

$$\overline{\bigcap_{i \in I} A_i} = \bigcup_{i \in I} \overline{A_i} \quad \text{and} \quad \overline{\bigcup_{i \in I} A_i} = \bigcap_{i \in I} \overline{A_i}$$

- *Generalized Distributive Laws:*

$$\left(\bigcap_{i \in I} A_i \right) \cup \left(\bigcap_{j \in J} B_j \right) = \bigcap_{i,j} (A_i \cup B_j) \quad \text{with} \quad \bigcap_{i,j} = \bigcap_{(i,j) \in I \times J}$$

$$\left(\bigcup_{i \in I} A_i\right) \cap \left(\bigcup_{j \in J} B_j\right) = \bigcup_{i,j} (A_i \cap B_j) \quad \text{with} \quad \bigcup_{i,j} = \bigcup_{(i,j) \in I \times J}$$

Here, $I = \{1, 2, 3, \dots, n\}$ and $J = \{1, 2, 3, \dots, m\}$. Then:

$$\begin{aligned} I \times J &= \{(i, j) \mid 1 \leq i \leq n, 1 \leq j \leq m, i, j \in \mathbb{N}\} \\ &= \{(1, 1), (1, 2), \dots, (1, m), (2, 1), (2, 2), \dots, (2, m), \dots, (n, 1), (n, 2), \dots, (n, m)\} \end{aligned}$$

3.0.22 Cardinality

The cardinality of a set is the number of elements in that set.

Let A and B be finite sets with $|A| = n$, $|B| = m$, and let X be the finite universal set. Then the following holds:

Cardinality of a Set

$$A = (A \cap B) \cup (A \setminus B), \quad |A| = |A \cap B| + |A \setminus B|$$

Cardinality of the complements

$$\begin{aligned} |A| &= |X \setminus A| = |X| - |A| \\ A \setminus B &= A \cap (X \setminus B), \quad |A \setminus B| = |A| - |A \cap B| \end{aligned}$$

Cardinality of the Cartesian Product

$$|A \times B| = |A| \cdot |B| = n \cdot m$$

*Inclusion-Exclusion Formula for Two Disjoint Sets

$$|A \cup B| = |A| + |B| = n + m$$

Inclusion-Exclusion Formula for Two Non-Disjoint Sets

Let $|A \cap B| = k$, then:

$$|A \cup B| = |A| + |B| - |A \cap B| = n + m - k \quad (\text{since we do not count the intersection twice})$$

Inclusion-Exclusion Formula for Three Non-Disjoint Sets

$$\begin{aligned} |A \cup B \cup C| &= |(A \cup B) \cup C| \\ &= |A \cup B| + |C| - |(A \cup B) \cap C| \\ &= |A| + |B| - |A \cap B| + |C| - |(A \cap C) \cup (B \cap C)| \\ &= |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C| \end{aligned}$$

General Formula for the cardinality of the union of sets

$$\left| \bigcup_{i=1}^n M_i \right| = \sum_{I \subseteq \{1, \dots, n\}, I \neq \emptyset} (-1)^{|I|-1} \left| \bigcap_{i \in I} M_i \right|$$

3.0.23 Definition of Countable and Non-countable

A set A is *infinite countable* or just *countable* if there exist a bijection from $\mathbb{N}$ to A . If A has infinite elements and there is no such bijection, we call it *non-countable*.

3.0.24 Countable Union

Given $A_n, n \in \mathbb{N}$ countable sets, then

$$M := \bigcup_{n \in \mathbb{N}} A_n$$

is also countable.

3.0.25 Countability of Subsets

Given $A \subseteq B$, where B is countable. Then A is also countable. The other case is that if A is non-countable then B must be non-countable as well.

3.0.26 Countability of the Rational Numbers

We show that the set of rational numbers $\mathbb{Q}$ is countable.

Every rational number can be expressed as

$$q = \frac{p}{q}, \quad p \in \mathbb{Z}, \quad q \in \mathbb{N}, \quad q \neq 0.$$

Thus,

$$\mathbb{Q} = \left\{ \frac{p}{q} \mid p \in \mathbb{Z}, \quad q \in \mathbb{N} \right\}.$$

The Cartesian product $\mathbb{Z} \times \mathbb{N}$ is countable because both $\mathbb{Z}$ and $\mathbb{N}$ are countable, and a countable product of countable sets is countable.

Hence, there exists an enumeration $(p_i, q_i)_{i \in \mathbb{N}}$ of all pairs (p, q) with $q \neq 0$. Defining

$$f(i) = \frac{p_i}{q_i},$$

we obtain a surjective mapping $f : \mathbb{N} \rightarrow \mathbb{Q}$.

Therefore, the set of rational numbers $\mathbb{Q}$ is countable.

□

3.0.27 Cantor's Diagonal Argument: The Real Numbers are Uncountable

We show that the set of real numbers in the interval $(0, 1) \subseteq \mathbb{R}$ is uncountable.

Assume, for contradiction, that $(0, 1)$ is countable. Then there exists a surjective mapping

$$f : \mathbb{N} \rightarrow (0, 1),$$

and thus all real numbers in $(0, 1)$ can be listed as

$$f(1) = 0.a_{11}a_{12}a_{13}\dots, \quad f(2) = 0.a_{21}a_{22}a_{23}\dots, \quad f(3) = 0.a_{31}a_{32}a_{33}\dots, \quad \text{and so on.}$$

We now construct a new real number $r \in (0, 1)$ by defining its decimal digits as

$$r = 0.b_1b_2b_3\dots$$

where

$$b_i = \begin{cases} 5, & \text{if } a_{ii} \neq 5, \\ 6, & \text{if } a_{ii} = 5. \end{cases}$$

By construction, r differs from each $f(i)$ in at least the i -th decimal place. Hence, $r \neq f(i)$ for all $i \in \mathbb{N}$, contradicting the assumption that f is surjective.

Therefore, the interval $(0, 1)$, and thus the set of real numbers $\mathbb{R}$, is uncountable.

□◻◻

3.0.28 The Power Set

The Power Set of a set is the set of all subsets of a given set

$$\mathcal{P}(x) := \{M : M \subseteq X\}$$

Its cardinality is 2^n with n being the number of elements in the original set X .

3.0.29 Set Systems

Let A be a non-empty set. A subset $\mathcal{F}$ of the power set of A is called a *set system* or *family of sets* of A .

3.0.30 Intersection and Union of Set Systems

We can define

$$\bigcap \mathcal{F} := \{x \mid \forall F, F \in \mathcal{F} \implies x \in F\}$$

and

$$\bigcup \mathcal{F} := \{x \mid \exists F, F \in \mathcal{F}, x \in F\}.$$

as the *intersection* and *union* of the sets.

The Intersection of the Empty Set is the Universal Class

Proof:

The set intersection of the empty-set gives us the set of all elements that are in every set of the empty-set. We have defined our definition

$$\bigcap \mathcal{F} := \{x \mid \forall F, F \in \mathcal{F} \implies x \in F\}$$

For our case

$$\bigcap \emptyset = \{x \mid \forall F, F \in \emptyset \implies x \in F\}.$$

We can see that that due to the implication our left side is always false which leads our right side to be always true which means that the statement is valid for all x . Resulting in

$$\bigcap \emptyset = \mathcal{U}$$

For the union

$$\bigcup \emptyset = \{x \mid \exists F \in \emptyset, x \in F\}$$

The condition is always false; therefore the union has to be the empty set.

□◻◻

Subset-property of Set Systems

Let $A \in \mathcal{B}$ then $A \subseteq \bigcup \mathcal{B} \wedge \bigcap \mathcal{B} \subseteq A$.

Proof:

$A \in \mathcal{B}$ gives us two cases for $A \subseteq \bigcup \mathcal{B}$.

Case $A = \bigcup \mathcal{B}$: this case is given when A is the only set in $\mathcal{B}$ and the equality holds, because otherwise we will have a contradiction.

Case $A \subset \bigcup \mathcal{B}$: Assume that $\exists x \in A : x \notin \bigcup \mathcal{B} \implies A \notin \mathcal{B}$, and therefore we have a contradiction. Hence, A is a proper subset of the union of the members of $\mathcal{B}$.

For our second statement $\bigcap \mathcal{B} \subseteq A$, we have again the two cases of

Case $\bigcap \mathcal{B} = A$ which is true if A is already the intersection of all members and due to $A \cap A = A$ it is proven true.

For the case $\bigcap \mathcal{B} \subset A$, we assume that the opposite is the case which implies $\exists b \in \bigcap \mathcal{B} : b \notin A$, which is a big contradiction to the definition of $\bigcap \mathcal{B}$. Therefore, the intersection must be a proper subset of A .

□◻◻

One application of this property is for example: Given $\{A_i\}_{i \in I}$ proof that $\bigcap A_i$ is a set.

We know that $\bigcap A_i \subseteq A_i$ for some $i \in I$. Hence, it is a subset which means by the axiom of subsets of classes and sets that it has to be a set.

□◻◻

Inclusion Function

Let $A \subseteq B$ the function

$$E_B(x) : A \rightarrow B, \quad x \mapsto x, \quad \forall x \in A$$

as the *inclusion function* of B in A .

Partition Function

If P is a partition of M then each element of M to exactly one $X \in P$

$$f_P : M \rightarrow P : m \mapsto X \in P \text{ with } m \in X,$$

is a map, called *natural map* of P .

Characteristic Function

Given two sets A and B , we define the *characteristic function* of B as:

$$C_B(x) : A \rightarrow \{0, 1\}, x \mapsto \begin{cases} 0 & x \notin B \\ 1 & x \in B \end{cases}$$

3.0.31 Equivalence Classes

Given a partition P of a set M , the elements of P are called *classes*. If X is a class, then its members are called *representatives* of the class.

A subset of M with an element of each of the classes of M is called the *representative set* or *traversal*.

The function $v : P \rightarrow M$ with $v(X) \in X$ for all $X \in P$ is called *representative function*.

3.0.32 Partitions

Let A be a set. A partition of A is a subset non-empty sets of $\mathcal{P}(A)$ with the following properties:

$$\forall i, j \in I, A_i \cap A_j = \emptyset \quad \text{for } i \neq j$$

and

$$\bigcup_{i \in I} A_i = A \quad \text{or equivalently} \quad \bigcup_{i \in I} A_i \subseteq A$$

Example:

Prove that $B_n = \{m \in \mathbb{Z} \mid \exists q, m = n + 5q\}$ is a partition of $\mathbb{Z}$.

First, we prove that disjoint property which means $m \in B_x \wedge m \in B_y \iff x = y$

For the union, we need to prove that every B_n is not empty which is very easy by letting $m = n + 5 \cdot 0 \implies m = n$ and then combining all of them with the union will give us the entire set of integers because every of the sets B_n has at least the number n in it which can be any integer.

□◻◻

The Quotient Space satisfies the Partition Properties

Given A/R , it satisfies the properties of a partition.

Proof:

For $X, Y \in A/R$, we know by the one of the properties of equivalence classes that $[a]_R \cap [b]_R \neq \emptyset \iff [a]_R = [b]_R$. Therefore, the first condition is satisfied.

To prove: $\bigcup_{[a]_R \in A/R} [a]_R = A$.

$x \in \bigcup A/R \implies x \in [a]_R \in A/R$, this implies that $x \in A$ because $\nexists a \in A, a \notin (x, y) \in R$. This was also already proven in the section subsection dedicated to the quotient space.

□◻◻

Equivalence Relation from a Partition

Given the partition S of the set A , then

$$R_S = \{(a, b) \mid (a, b) \in A \times A, \exists X \in S, a, b \in X\}$$

is an equivalence relation.

Proof:

Reflexivity: $a \in A = \bigcup S \implies a \in X \in S \implies (a, a) \in R_S$.

Symmetry: $a, b \in A \bigcup S \implies a, b \in X \in S \implies (a, b), (b, a) \in R_S$.

Reflexivity: $(a, b), (b, c) \in R_S \implies a, b \in X \in S \wedge b, c \in Y \in S \implies b \in X \cap Y$ by definition $X \cap Y \neq \emptyset \iff X = Y$, therefor, $a, c \in X, (a, c) \in R_S$.

□◻◻

3.0.33 Relation from the Relation of the Quotient Space and Partition from the Quotient Space

Given R on A , $R_{A/R} = R$ with

$$R_{A/R} = \{(a, b) \mid (a, b) \in A \times A, \exists X \in A/R, a, b \in X\}.$$

And for S a partition of A , then $A/R_S = S$.

Proof:

For the first statement we need to prove the condition of an equivalence relation:

Reflexivity: $(a, b) \in R_{A/R} \implies a, b \in [x]_R \in A/R \implies a \sim x \wedge b \sim x \wedge a \sim b \wedge a \sim a \implies (a, a) \in R_{A/R}$.

Symmetry: $(a, b), (b, a) \in R_{A/R} \implies a, b \in [x]_R \in A/R \implies a \sim x \wedge b \sim x \wedge a \sim b \wedge b \sim a \implies (a, b), (b, a) \in R_{A/R}$.

Transitivity: $(a, b), (b, c) \in R_{A/R} \implies a, b \in [x]_R \in A/R \wedge b, c \in [y]_R \in A/R \implies a \sim x \wedge b \sim x \wedge b \sim y \wedge a \sim y \implies (a, b), (b, c), (a, c) \in R_{A/R}$.

Thus, $R_{A/R} = R$. For the other direction $(a, b) \in R \implies b \in [a]_R \in A/R \implies aR_{A/R}b$

For the second part: $A/R_S = S$:

Let S be a partition of A , and $X \in A/R_S$. For $X = [a]_{R_S} = \{x \in A \mid aR_Sx\}$. For a and b to be in a relation they need to be part of the same element $Y \in S$. We need to prove that $Y = [a]_{R_S}$.

This is proven directly by the very definition of the relation R_S which implies that all elements belong to the same set Y . For the other direction if $a, b \in Y \implies aR_Sb \implies b \in [a]_{R_S} \in A/R_S$.

□□□

Properties of Partitions

- Every equivalence relation corresponds to a partition:

$$J : \{R : R \text{ is an equivalence relation on } X\} \rightarrow \{F : F \text{ is a partition of } X\}$$

where $J(R) := X/R$ is a bijection.

- If F is a partition of X , then we can define an equivalence relation R_F by:

$$R_F := \{(x, y) \in X \times X : \exists M \in F \text{ such that } x, y \in M\}$$

Then R_F is an equivalence relation on X .

- Two equivalence classes are either disjoint or equal.

Proof: Equivalence Relations Correspond to Partitions

We want to prove that there exists a one-to-one correspondence between equivalence relations and partitions. We first start by start proving that from a family of equivalence classes we get a partition.

Let $\mathcal{C}$ be the set of all the equivalence classes for an equivalence relation on A , and C_x be some of those sets of equivalence classes.

We must prove that each of its equivalence classes are disjoint if not equal. By the definition if $\exists y \in [a]_{\sim} \wedge y \in [b]_{\sim}$, then this would imply that y is related to both a and b which means that they are in the same equivalence class, because otherwise $a \sim y \wedge b \sim y \iff \neg(a \sim b)$, which is a contradiction. Hence, equivalence classes are either equal or disjoint.

Now, for the union. We can state that $\bigcup [a]_{\sim} \subseteq A$ due to all of its elements forming at least a proper subset of A . The equality, can be shown via the definition of the equivalence class which states that each element in A has its own equivalence class.

Analogously, we can prove it the other way around. By having disjoint sets, we know that they have to be in different equivalence classes $A_i \cap A_j = \emptyset \implies \forall x \in A_i, x \neg (\sim)y, \forall y \in A_j$.

□◻◻

3.0.34 Equivalence Relations and Partitions

Let M be a set.

1. If $\sim$ is an equivalence relation on M , then the equivalence classes defined by

$$[m_0] := [m_0]_{\sim} := \{ m \in M \mid m \sim m_0 \}$$

for $m_0 \in M$ form a *partition* of M . This partition is usually denoted by $M/\sim$.

2. If P is a partition of M , then the relation $\sim_P$ defined by

$$m \sim_P n \quad \text{if and only if there exists } X \in P \text{ such that } m, n \in X,$$

is an *equivalence relation*.

3. The following holds:

$$M/\sim_P = P, \quad \text{for all partitions } P \text{ of } M,$$

and

$$\sim_{(M/\sim)} = \sim, \quad \text{for every equivalence relation } \sim \text{ on } M.$$

3.0.35 Equivalence Relations under Functions

Pre-image

Let $f : A \rightarrow B$ and G be an equivalence relation in B . The *pre-image* of G in A is defined as

$$f^{-1}(G) = \{(x, y) \mid (f(x), f(y)) \in G\},$$

and it is also an equivalence relation.

Restriction

Similarly, we can also define the *restriction* for when $B \subseteq A$

$$G_{[B]} = \{(x, y) \mid x \in B \wedge y \in B \wedge (x, y) \in G\}$$

3.0.36 Refinement

Let G and H be equivalence relations in A . We call G a *refinement* of H if $G \subseteq H$; we also say that G is *finer* than H , and that H is *coarser* than G .

3.0.37 Quotient Space for Refinements

Let G and H be equivalence relations on a set A and let G be a refinement of H . The *quotient* of H by G usually denoted as H/G , is a relation in A/G defined as follows:

$$H/G = \{([x]_G, [y]_G) \mid (x, y) \in H\}$$

In simpler words, it is a set of the pairs consisting of the equivalence classes of a pair $(x, y) \in H$.

The Quotient is an Equivalence Relation

The Quotient is an Equivalence Relation.

Proof:

Reflexivity: For $([x]_G, [y]_G) \in H/G$ we know claim that $[x]_G \sim [x]_G$. By the reflexivity of H we know that $x \sim x$, which means that $(x, x) \in H$; and therefore, $[x]_G \sim [x]_G$.

Symmetry: For $([x]_G, [y]_G) \in H/G$ we know claim that $([x]_G \sim [y]_G) \wedge ([y]_G \sim [x]_G)$. Analogously to our last property, if $(x \sim y \wedge (y \sim x))$ were not true then H would not be an equivalence relation which leads to a contradiction of our initial conditions. Hence, the symmetric property is present.

Transitivity: Lastly, the transitivity we again use the same argument as before using our base elements and pairs from H to arrive at a contradiction, which guarantees us that their equivalence class counterpart are also part of an equivalence relation.

□◻◻

Equivalence Classes under Refinements

Let G and H be equivalence relations in A , suppose $G \subseteq H$. Then $z \in [x]_H \implies [z]_G \subseteq [x]_H$.

Proof:

$z \in [x]_H$ means that $(z, x) \in H$, then

$$\begin{aligned} y \in [z]_G &\implies (y, z) \in G \subseteq H \implies (y, x) \in H \\ &\implies y \in [x]_H \end{aligned}$$

Thus, $[z]_G \subseteq [x]_H$.

□◻◻

This can also be extended for all elements $x \in A$, as $[x]_G \subseteq [x]_H$.

3.0.38 Equivalence Relations and Surjective Mappings

- Let M be a set and $\sim$ an equivalence relation on M . Then there exists a set P and a surjective mapping $\Gamma : M \rightarrow P$ such that $\sim = \sim_\Gamma$. Hence, every equivalence relation can be represented as the equality of images under a surjective mapping.
- If $\Gamma : M \rightarrow N$ is a surjective mapping and $P := M/\sim_\Gamma$ is the partition corresponding to equality of images, then the function

$$f : P \rightarrow N, \quad [m]_{\sim_\Gamma} \mapsto \Gamma(m)$$

is a well-defined bijection.

- If $\Gamma_1 : M \rightarrow P_1$ and $\Gamma_2 : M \rightarrow P_2$ are two surjective mappings, then $\sim_{\Gamma_1} = \sim_{\Gamma_2}$ if and only if there exists a bijection $f : P_1 \rightarrow P_2$ with $\Gamma_2 = f \circ \Gamma_1$


3.0.39 Homomorphism Theorem for Sets

Given $f : M \rightarrow N$ with $M/\sim_f = \{f^{-1}(\{n\}) \mid n \in f(M)\}$

By factorizing f as $f = \bar{f} \circ \iota_f$ with ι_f surjective and $\bar{f}$ injective, we obtain the following:

- The mapping $\iota_f : M \rightarrow M/\sim_f$ is the natural map of the partition $M/\sim_f$ induced by f .
- The mapping $\bar{f} : M/\sim_f \rightarrow N : f^{-1}(\{f(m)\}) \mapsto f(m)$ is well-defined.

The homomorphism theorem for Sets can be visualized through the following commutative diagram:



This is saying us to first form collections of elements and then use our second function to associate that connections to their respective images.

3.0.40 Representatives

Given a set $X \subseteq A$, we consider it a set of *representatives* for a partition S if

$$\forall Y \in S \quad X \cap Y = \{a\}, a \in A$$

3.0.41 Family of Subsets Operations

Let $\mathcal{F}$ be a system of sets (a family of subsets) on the set X . We define:

$$\bigcup_{M \in F} M := \bigcup F := \{x \in X : \text{there exists } M \in F \text{ such that } x \in M\},$$

$$\bigcap_{M \in F} M := \bigcap F := \{x \in X : x \in M \text{ for all } M \in F\}.$$

3.0.42 Product of Classes

Given a family of classes $\{A_i\}_{i \in I}$ we define the *product of classes* as

$$\prod A_i = \{f : I \rightarrow A \mid f(i) \in A_i \wedge i \in I\}$$

Is the set of all functions from I to $\bigcup A_i$.

Properties

- $(\prod_{i \in I} A_i) \cap (\prod_{j \in J} B_j) = \prod_{(i,j) \in I \times J} (A_i \cap B_i)$
- $(\prod_{i \in I} A_i) \cup (\prod_{j \in J} B_j) = \prod_{(i,j) \in I \times J} (A_i \cup B_i)$

3.0.43 Pigeon Principle

For $g : A \rightarrow B$ and $|A| = n + 1$, $|B| = n$ then there exists $b \in B$ such that $|g^{-1}(\{b\})| \geq 2$.

In other words if you have n boxes and $n + 1$ objects, then at least one of those boxes will have more than 2 elements.

3.0.44 The Cantor Set

In the following, we want to study the Cantor set $\mathcal{C} \subseteq \mathbb{R}$.

First, we describe its construction informally: We start with the unit interval $A_0 := [0, 1]$ and remove from it the "middle third" (where the boundary points are not removed). The result A_1 thus consists of the sub-intervals $[0, \frac{1}{3}]$ and $[\frac{2}{3}, 1]$. Now we remove from each of these two sub-intervals the middle third and continue in this manner.

For fixed but arbitrary $n \in \mathbb{N}_0$, A_n thus consists of 2^n pairwise disjoint closed sub-intervals of length 3^{-n} , and A_{n+1} arises from A_n by removing the middle third from each of these sub-intervals. What "remains" is the Cantor set.

Now mathematically precise: For $n \in \mathbb{N}_0$, let

$$A_{n+1} := \frac{A_n}{3} \cup \left(\frac{2+A_n}{3}\right), \text{ with } A_0 = [0, 1]$$

Regarding the notation - for $X \subseteq \mathbb{R}$ and $a, t \in \mathbb{R}$, we define

$$aX := \{ax : x \in X\}$$

$$t + X := \{t + x : x \in X\}$$

Then $\mathcal{C} := \bigcap_{n=0}^{\infty} A_n$.

For the intuition, imagine our fraction squishing all the numbers and then adding two to a copy of those numbers. Then repeating the process multiple times.

Properties:

- The Cantor Set is uncountable
- The Cantor Set has a length of 0
- The set can be interpreted as all infinite sequences of zeroes and ones, when handling left and right as 0 and one.

Proof I: We give a proof by contradiction.

Assume the Cantor set is countable. Then there exists a surjective function

$$c : \mathbb{N} \rightarrow \mathcal{C}.$$

After the first step of the Cantor construction we have

$$A_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1].$$

Since

$$c(1) \in \mathcal{C} \subset A_1,$$

the point $c(1)$ lies in exactly one of the two disjoint intervals. We choose the other interval and call it I_1 .

Thus:

- $I_1 \subset A_1$,
- I_1 is closed,
- $c(1) \notin I_1$.

For the inductive step, assume there exists a closed interval

$$I_n \subset A_n \quad \text{with} \quad c(n) \notin I_n.$$

When passing from A_n to A_{n+1} , the middle third of each interval is removed. Therefore, I_n splits into exactly two disjoint closed sub-intervals of A_{n+1} .

Since $c(n+1)$ can lie in only one of these two intervals, we choose the other and call it I_{n+1} .

Thus:

- $I_{n+1} \subsetneq I_n$,
- I_{n+1} is closed and non-empty,
- $c(n+1) \notin I_{n+1}$.

By construction the nested sequence

$$I_1 \supsetneq I_2 \supsetneq I_3 \supsetneq \cdots$$

is obtained, and the lengths satisfy

$$|I_n| = 3^{-n}.$$

Thus, the lengths converge to 0.

Since the intervals are closed, bounded, nested, and their lengths tend to 0, the interval-nesting principle implies:

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset.$$

There exists exactly one point

$$x \in \bigcap_{n=1}^{\infty} I_n.$$

Since each $I_n \subset A_n$, it follows that

$$x \in \mathcal{C}.$$

Contradiction

Because c is surjective by assumption, there exists a $k \in \mathbb{N}$ such that

$$x = c(k).$$

However, by our construction I_k was chosen so that

$$c(k) \notin I_k,$$

while at the same time

$$x \in I_k.$$

This is a contradiction. Thus, the assumption that the Cantor set is countable leads to a contradiction. Therefore, the Cantor set has to be uncountable.

◻

Proof II:

This is not the most rigorous proof, but accomplishes its goal:

The original interval has length 1 and consists of one unit, the next $1 - \frac{1}{3}$ and consists of two units, this pattern with $2^n \frac{1}{3^n}$ continues forever thus,

$$L = 1 - \sum_{i=1}^{\infty} 2^n \frac{1}{3^n}$$

Note that the sum is a geometric series therefore,

$$L = 1 - 1 = 0$$

QED

Chapter 4

Relations, Maps and Functions

4.0.1 Relations

Given two classes A and B a *relation* R is subset of its Cartesian product $R \subseteq A \times B$.

Example:

Given $A = \{1, 2, 3\}$ find the elements of the relation R defined by $R = \{(a, b) \in A \times A \mid a < b\}$.

$$A \times A = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)\}$$

The relation R would then be:

$$R = \{(1, 2), (1, 3), (2, 3)\}$$

Common types of relations include:

- Functions (special relations where each input has exactly one output)
- Equivalence relations (reflexive, symmetric, and transitive)
- Partial orders (reflexive, antisymmetric, and transitive)

4.0.2 Types of relations

Let A be a set and X be a relation on A .

- *Reflexive:* $\forall a \in A : (a, a) \in X$ (or written as $a \sim a$)
- *Irreflexive:* $\forall a \in A : (a, a) \notin X$
- *Symmetric:* $\forall a, b \in A : (a, b) \in X \rightarrow (b, a) \in X$ (or written as $(a \sim b) \Rightarrow (b \sim a)$)
- *Antisymmetric:* $\forall a, b \in A : (a, b) \in X \text{ and } (b, a) \in X \implies a = b$
- *Transitive:* $\forall a, b, c \in A : (a, b) \in X \text{ and } (b, c) \in X \rightarrow (a, c) \in X$ (or written as $(a \sim b) \text{ and } (b \sim c) \Rightarrow (a \sim c)$)
- *Total:* $\forall a, b \in A : a \neq b \rightarrow (a, b) \in X \text{ or } (b, a) \in X$

4.0.3 Diagonal Graph

Given a relation, the graph I_A is defined as: $\{(x, x) \mid x \in A\}$ and denoted as the *diagonal graph* of A .

Lemmi for Proofs

Let the graph G be a relation, on A .

- G is symmetric if and only if $G = G^{-1}$
- G is antisymmetric if and only if $G \cap G^{-1} \subseteq I_A$
- G is transitive if and only if $G \circ G \subseteq G$.

4.0.4 Order Relation

A relation is called an *order relation* $R \subseteq A \times A$ if it is reflexive, antisymmetric and transitive.

Inverse Order Relation

Given an order relation G on A , then G^{-1} is also an order relation on A .

4.0.5 Strict Order Relation

A relation is called a *strict order relation* $R \subseteq A \times A$ if it is irreflexive, antisymmetric and transitive.

Inverse Strict Order Relation

Given a strict order relation G on A , then G^{-1} is also a strict order relation on A .

4.0.6 Equivalence relation

An *equivalence relation* is a relation $R \subseteq A \times A$ that is symmetric, transitive and reflexive.

Example:

$$R := \{(a, b) \in \mathbb{N} \times \mathbb{N} : a = b\}$$

Properties of Equivalence Relations

Given the equivalence relations G and H , the following properties apply:

- $G \circ H$ is an E.R. if and only if $G \circ H = H \circ G$.
- $G \cup H$ is an E.R. if and only if $G \circ H \subseteq G \cup H \wedge H \circ G \subseteq G \cup H$.

The proofs are left as an exercise for the reader. (Hint: use the first property to prove the second).

Inverse Equivalence Relation

Given an equivalence relation G on A , then G^{-1} is also an equivalence relation on A .

Proof:

4.0.7 Equivalence Class

Let $\sim$ be an equivalence relation on a set A and $a \in A$, the *equivalence class* of a , denoted by

$$[a]_{\sim} := \{y \in A \mid (a, y) \in \sim\}$$

is the set of all elements in A that are related to a .

4.0.8 Lemmi about Equivalence Classes

For $a, b \in A$ then

- $aEb \iff [a]_E = [b]_E$
- $a \neg Eb, (a, b) \notin E \iff [a]_E \cap [b]_E = \emptyset$

Proof I:

$aEb, x \in [a]_E$ tells us that also $xEb \implies x \in [b]_E$. For the other direction using the reflexive property we get $[a]_E = [b]_E \implies aEa \implies a \in [a]_E = [b]_E$

□◻◻

Proof II:

$x \in [a]_E \cap [b]_E \implies xEa, xEb, aEx \implies aEb$ a contradiction. Now, for $[a]_E \cap [b]_E = \emptyset$ we can say $aEb \implies a \in [a]_E, a \in [b]_E$, another contradiction.

□◻◻

4.0.9 Quotient Space

Let $\sim$ be an equivalence relation on a set A . The *quotient space* of A by $\sim$, denoted by $A/\sim$ (or sometimes A/R), is the set of all distinct equivalence classes of elements in A . Formally, it is defined as:

$$A/\sim := \{[x]_\sim \mid x \in A\}$$

The quotient space $A/\sim$ is a partition of the original set A into disjoint equivalence classes. Each element of the quotient space is an equivalence class $[x]$, which itself is a subset of A .

The Union of the Quotient Space

Given A/R , then $\bigcup A/R = A$.

Proof: Suppose $a_0 \in A \wedge a_0 \notin [a]_R, \forall [a]_R \in A/R$. If R is an equivalence relation then $a_0 \in A$ with R implies that $(a_0, a_0) \in R$. Therefore, the only way for a_0 to not belong to any equivalence class is if R were not an equivalence relation, which contradicts our initial condition; hence, we know that the union of the equivalence classes given an equivalence relation must be equal to the original set.

□◻◻

4.0.10 Map Definition

A *map* for two sets A and B is defined as a subset of $A \times B$. And it is denoted by

$$f : A \rightarrow B \subseteq \{(a, b) \mid (a, b) \in A \times B\}$$

4.0.11 The Set of All Maps

For the sets M and N , we define the sets off all maps $f : M \rightarrow N$ as

$$N^M := \{f \mid f : M \rightarrow N\}$$

4.0.12 The Characteristic Map

Given $T \subseteq M$ we define the *characteristic map* as

$$\chi_T : M \rightarrow \{0, 1\} : m \rightarrow \begin{cases} 1, & m \in T \\ 0, & m \notin T \end{cases}$$

This map tell us if an element is in the subset. Also, each map $\alpha : M \rightarrow \{0, 1\}$ defines a unique subset T of M , $T := M \mid \alpha(m) = 1$.

4.0.13 Functions

A subset $f \subseteq X \times Y$ is called a *function* if:

$$\forall x \in X : \exists y \in Y : f(x) = y$$

and

$$(x, y) \in f \text{ and } (x, y') \in f \implies y = y'.$$

That is, for each $x \in X$, there exists at most one $y \in Y$ such that $(x, y) \in f$.

4.0.14 The Graph

We can also consider the graph as a set of ordered pairs x, y where $y = f(x)$.

$$\text{graph}(f) := \{(x, f(x)) \mid x \in X, f(x) \in Y\} = G$$

The Inverse Graph

Let G be a graph. We define

$$G^{-1} := \{(x, y) \mid (y, x) \in G\}$$

as *inverse graph* of G .

The Composite Graph

Let G and H be graphs. We define

$$G \circ H := \{(x, y) \mid \exists z \in (x, z) \in H \wedge (z, y) \in G\}$$

as the *composite graph* of G and H .

- $G \circ (H \circ J) = (G \circ H) \circ J$
- $(G^{-1})^{-1} = G$
- $(G \circ H)^{-1} = H^{-1} \circ G^{-1}$

Proof I:

$$(G \circ H) \circ J = G \circ (H \circ J)$$

$$G \circ H = \{(z, y) \mid \exists w \in (z, w) \in H \wedge (w, y) \in G\}$$

Proceeding

$$(G \circ H) \circ J = \{(x, y) \mid \exists z, w \text{ s.t. } (x, z) \in J \wedge (z, w) \in H \wedge (w, y) \in G\}.$$

For $G \circ (H \circ J)$

$$H \circ J = \{(x, w) \mid \exists z \in (x, z) \in J \wedge (z, w) \in H\}.$$

Proceeding

$$G \circ (H \circ J) = \{(x, y) \mid \exists z, w \in (x, z) \in J \wedge (z, w) \in H \wedge (w, y) \in G\}$$

QED

Proof II:

$$(G^{-1})^{-1} = G$$

$$\begin{aligned} (x, y) \in (G^{-1})^{-1} &\iff (y, x) \in G^{-1} \\ &\iff (x, y) \in (G^{-1})^{-1} \end{aligned}$$

QED

Proof III:

$$(G \circ H)^{-1} = H^{-1} \circ G^{-1}.$$

$$\begin{aligned} (G \circ H)^{-1} &= \{(x, y) \mid \exists z \in (y, z) \in H \wedge (z, x) \in G\} \\ &\iff (y, z) \in H \wedge (z, x) \in G \\ &\iff (z, y) \in H^{-1} \wedge (x, z) \in G^{-1} \end{aligned}$$

QED

Functional Graph

A functional graph is a graph that satisfies the second constraint of functions. Therefore, it corresponds to a function. Hence, all of our graphs properties are applicable to functions.

4.0.15 Domain

The *domain* of a relation R from set A to set B is the set of all elements in A that are related to at least one element in B . Formally, if $R \subseteq A \times B$ is a relation, then the domain of R is defined as:

$$\text{dom}(R) = \{a \in A \mid \exists b \in B \text{ such that } (a, b) \in R\}$$

In other words, the domain consists of all the input values that appear in the ordered pairs of the relation. For example, if $R = \{(1, 4), (2, 5), (3, 4), (2, 6)\}$, then $\text{Dom}(R) = \{1, 2, 3\}$.

4.0.16 The identity

$$\text{id}(f) := \text{id}_X := \{(x, x) \mid x \in X\}$$

4.0.17 Image/Range

The *image* or *range* of a relation R from set A to set B is the set of all elements in B that are related to at least one element in A . Formally, if $R \subseteq A \times B$ is a relation, then the image of R is defined as:

$$\text{ran}(R) = \text{img}(R) = \{b \in B \mid \exists a \in A \text{ such that } (a, b) \in R\}$$

or

$$\text{img}(R[A]) = \{y \in \text{ran}(R) \mid \exists a \in A \text{ such that } xRy\}$$

In other words, the image consists of all the output values that appear in the ordered pairs of the relation. For example, if $R = \{(1, 4), (2, 5), (3, 4), (2, 6)\}$, then $\text{img}(R) = \{4, 5, 6\}$.

Proof that the Domain And Range are Sets

Given a graph G which is a set. Prove that the domain and range are also sets.

$$\forall (x, y) \in G : (x, y) = \{x, \{x, y\}\}$$

Thus, the union of all the elements in G has a structure of the form

$$\bigcup G = \{x, y, z, \dots, \{x, y\}, \{x, z\}, \{y, z\}, \dots\}$$

By iterating one more time with get the union of the union

$$\bigcup \bigcup G = \{x, y, z, \dots\}$$

Hence, by the definition of the domain and the range of the graph we can directly see that they are subsets of the union of the union of the graph G , and thus they are sets.

□◻◻

4.0.18 Composition of Maps

The composition of two functions is the operation of applying one function to the result of another function. If we have functions $f : A \rightarrow B$ and $g : B \rightarrow C$, then the composition of g and f , denoted as $g \circ f$ (read as g composed with f), is a function from A to C defined by:

$$(g \circ f)(a) = g(f(a)) \quad \text{for all } a \in A$$

The composition applies f first, then applies g to the result. Note that the co-domain of f must match the domain of g for the composition to be defined.

The formal definition is

$$g \circ f := \{(s, u) \in S \times U \mid \exists t \in T \text{ such that } (s, t) \in f \wedge (t, u) \in g\}$$

Example:

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = x^2$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $g(x) = x + 3$. Then:

$$(g \circ f)(x) = g(f(x)) = g(x^2) = x^2 + 3$$

$$(f \circ g)(x) = f(g(x)) = f(x + 3) = (x + 3)^2 = x^2 + 6x + 9$$

Note that $g \circ f \neq f \circ g$ in general, which shows that function composition is not commutative.

Domain and Range Under Composition Corollary

Let G and H be graphs, then $\text{img}(H) \subseteq \text{dom}(G) \implies \text{dom}(G \circ H) = \text{dom}(H)$

Proof:

Note that for

$$\forall y \in \text{img}(H) : y \in \text{dom}(G).$$

Also, when constructing $G \circ H$ we get a map from $\text{dom}(H)$ to the $\text{img}(G)$, because for all of our $y \in \text{img}(H)$ we have can always find a pair $(y, w) \in \text{img}(G)$ due to the structure of our graph. Hence, the domain of our composite graph is $\text{dom}(H)$.

□◻◻

4.0.19 Types of Functions

Injective Functions

An injective function (also called a one-to-one function) is a function that maps distinct elements from the domain to distinct elements in the co-domain.

Formally, a function $f : A \rightarrow B$ is injective if for all $a_1, a_2 \in A$:

$$a_1 \neq a_2 \rightarrow f(a_1) \neq f(a_2)$$

Equivalently, using the contrapositive:

$$f(a_1) = f(a_2) \rightarrow a_1 = a_2$$

Injection

Given $T \subseteq M$ a subset of M . The *injection* from T to M defined as

$$\iota_T : T \rightarrow M : t \mapsto t$$

is injective.

Surjective Functions

A surjective function (also called an onto function) is a function whose image equals its co-domain, meaning that every element in the co-domain has at least one pre-image in the domain.

Formally, a function $f : A \rightarrow B$ is surjective if:

$$\forall b \in B, \exists a \in A \text{ such that } f(a) = b$$

Bijection Functions

A bijective function (also called a one-to-one correspondence) is a function that is both injective and surjective. In other words, every element in the co-domain is mapped to by exactly one element in the domain.

Formally, a function $f : A \rightarrow B$ is bijective if it is both:

- Injective: $\forall a_1, a_2 \in A, a_1 \neq a_2 \rightarrow f(a_1) \neq f(a_2)$
- Surjective: $\forall b \in B, \exists a \in A \text{ such that } f(a) = b$

Bijective functions establish a perfect pairing between elements of the domain and co-domain, where each element in the domain corresponds to exactly one element in the co-domain, and vice versa. A bijection allows us to define an inverse function $f^{-1} : B \rightarrow A$.

4.0.20 Propositions on Images and Pre-images under Set Operations

Let $f : X \rightarrow Y$ be a function.

Union and Cut Sets

- For subsets $A_1, A_2 \subseteq X$, we have:

$$f(A_1 \cup A_2) = f(A_1) \cup f(A_2) \quad \text{and} \quad f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

- For subsets $B_1, B_2 \subseteq Y$, we have:

$$f^{-1}(B_1 \cup B_2) = f^{-1}(B_1) \cup f^{-1}(B_2) \quad \text{and} \quad f^{-1}(B_1 \cap B_2) = f^{-1}(B_1) \cap f^{-1}(B_2)$$

Proof:

We prove the second part of 1 and the first part of 2.

Let $y \in f(A_1 \cap A_2)$. Then there exists $x \in A_1 \cap A_2$ such that $f(x) = y$. Since $x \in A_1$ and $x \in A_2$, it follows that $y \in f(A_1)$ and $y \in f(A_2)$, hence $y \in f(A_1) \cap f(A_2)$. Therefore, every element of $f(A_1 \cap A_2)$ is also an element of $f(A_1) \cap f(A_2)$, so:

$$f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$$

Let $x \in f^{-1}(B_1 \cup B_2)$. Then $f(x) \in B_1 \cup B_2$, which means $f(x) \in B_1$ or $f(x) \in B_2$. Thus, $x \in f^{-1}(B_1)$ or $x \in f^{-1}(B_2)$, which implies:

$$x \in f^{-1}(B_1) \cup f^{-1}(B_2)$$

Hence, both sets contain the same elements and are therefore, equal:

$$f^{-1}(B_1 \cup B_2) = f^{-1}(B_1) \cup f^{-1}(B_2)$$

□◻

Union and Cut of the whole Domain and Range

Let $f : X \rightarrow Y$ be a function.

- Let $\mathcal{F}$ be a collection of subsets of X . Then:

$$f\left(\bigcup_{A \in \mathcal{F}} A\right) = \bigcup_{A \in \mathcal{F}} f(A) \quad \text{and} \quad f\left(\bigcap_{A \in \mathcal{F}} A\right) \subseteq \bigcap_{A \in \mathcal{F}} f(A)$$

- Let $\mathcal{G}$ be a collection of subsets of Y . Then:

$$f^{-1}\left(\bigcup_{B \in \mathcal{G}} B\right) = \bigcup_{B \in \mathcal{G}} f^{-1}(B) \quad \text{and} \quad f^{-1}\left(\bigcap_{B \in \mathcal{G}} B\right) = \bigcap_{B \in \mathcal{G}} f^{-1}(B)$$

Proof (partial):

We show the first statement of part (ii); the rest follows analogously.

Let $x \in f^{-1}\left(\bigcup_{B \in \mathcal{G}} B\right)$. Then:

$$f(x) \in \bigcup_{B \in \mathcal{G}} B \Leftrightarrow \exists B \in \mathcal{G} \text{ such that } f(x) \in B \Leftrightarrow \exists B \in \mathcal{G} \text{ such that } x \in f^{-1}(B)$$

Hence:

$$x \in \bigcup_{B \in \mathcal{G}} f^{-1}(B)$$

It follows that:

$$f^{-1}\left(\bigcup_{B \in \mathcal{G}} B\right) = \bigcup_{B \in \mathcal{G}} f^{-1}(B)$$

□◻

Example:

We define $A = \{1, 0\}$ and $B = \{2\}$ and $f(1) = 3, f(2) = 3, f(0) = 1$. Then

$$f(A \cap B) = f(\emptyset) = \emptyset$$

And

$$f(A) \cap f(B) = 3$$

4.0.21 Bijection of the Power Set and 2^A

We denote 2^A as the set of all functions from A to a set of two elements. The statement goes as following: there is a bijection between the power set $\mathcal{P}(A)$ and 2^A .

Proof:

Let $\gamma : \mathcal{P}(A) \rightarrow 2^A$. If $B \in \mathcal{P}(A)$ let

$$\gamma(B) = C_B, \forall B \in \mathcal{P}(A).$$

By this definition we get the properties of a function directly. Now we have to prove that the function is a one-to-one correspondence.

For the injectivity, we already know that each for $\gamma(B) = \gamma(C) \implies B = C$ due to else being the case that we have to elements in the power set which are identical.

For the surjectivity, if $f \in 2^A$, and we let $B = f^{-1}(0)$, then $f = C_B = \gamma(B)$. Thus, γ satisfies the surjectivity.

□◻◻

4.0.22 Restriction Function

Let G be a graph and $B \subseteq \text{dom}(G)$ then we denote the *restriction* of G as:

$$G_{[B]} = \{(x, y) \mid (x, y) \in G \wedge x \in B\}.$$

The range of this restricted map can be denoted as;

$$G(B) = \{y \mid x \in \text{dom}(B) \wedge (x, y) \in G\}$$

4.0.23 Even Function

A real function f is *even* if

- $\forall x \in \text{Dom}(f) \exists -x \in \text{Dom}(f)$
- $\forall x \in \text{Dom}(f)$ the following is true $f(x) = f(-x)$

4.0.24 Odd Function

A real function f is *odd* if

- $\forall x \in \text{Dom}(f) \exists -x \in \text{Dom}(f)$
- $\forall x \in \text{Dom}(f)$ the following is true $f(x) = -f(-x)$

4.0.25 Pre-image

Given $f : M \rightarrow N$. For $n \in N$ we define

$$f^{-1}(\{n\}) := \{m \in M \mid f(m) = n\},$$

as the *pre-image* of n . The complete *pre-image* for $T \subseteq N$ is defined as:

$$f^{-1}(T) := \bigcup_{t \in T} f^{-1}(\{t\}) = \{m \in M \mid f(m) \in T\}$$

4.0.26 Inverse of a Function

The *inverse* of a function $f : A \rightarrow B$ is a function $f^{-1} : B \rightarrow A$ that reverses the operation of f . That is, if f maps an element $a \in A$ to an element $b \in B$, then the inverse function f^{-1} maps b back to a .

Formally, a function $f : A \rightarrow B$ has an inverse $f^{-1} : B \rightarrow A$ if and only if f is bijective (both injective and surjective). The inverse function satisfies the following properties:

$$f^{-1}(f(a)) = a \quad \text{for all } a \in A,$$

$$f(f^{-1}(b)) = b \quad \text{for all } b \in B.$$

In other words, composing a function with its inverse yields the identity function. That is:

$$f^{-1} \circ f = id_A \quad \text{and} \quad f \circ f^{-1} = id_B$$

Where id_A and id_B are the identity functions on sets A and B , respectively.

Proof:

$\Rightarrow$: direction is trivial, due to the definition of a bijective function.

$\Leftarrow$: Our function has the following properties:

$$f^{-1}(f(a)) = a \quad \text{for all } a \in A,$$

$$f(f^{-1}(b)) = b \quad \text{for all } b \in B.$$

The first condition, gives us the injectivity due to the constraint the for all $b \in B$ there exist exactly one $a \in A$ such that $f(a) = b$. Else, the condition would no be obeyed.

The second one gives us the surjectivity of f by providing a definition for each of the $b \in B$. Otherwise, our condition would be violated.

The uniqueness is guaranteed by

$$f^{-1}(b) \neq g^{-1}(b)$$

Gives us a contradiction, if they would have the same properties but were different.

◻◻◻

Steps to Find the Inverse of a Function

To find the inverse of a function $f(x)$, follow these steps:

1. Replace $f(x)$ with y : $y = f(x)$
2. Interchange the variables x and y : $x = f(y)$
3. Solve for y in terms of x : $y = f^{-1}(x)$
4. Verify that the resulting function is indeed the inverse by checking that $f^{-1}(f(x)) = x$ and $f(f^{-1}(x)) = x$

Example:

Let's apply the steps above to find the inverse of $f(x) = 2x + 3$.

Step 1: Replace $f(x)$ with y

$$y = 2x + 3$$

Step 2: Interchange the variables x and y

$$x = 2y + 3$$

Step 3: Solve for y in terms of x

$$\begin{aligned}x &= 2y + 3 \\x - 3 &= 2y \\ \frac{x - 3}{2} &= y\end{aligned}$$

So, the inverse function is:

$$f^{-1}(x) = \frac{x - 3}{2}$$

Step 4: Verify that $f^{-1}(f(x)) = x$ and $f(f^{-1}(x)) = x$

Let's verify $f^{-1}(f(x)) = x$:

$$\begin{aligned}f^{-1}(f(x)) &= f^{-1}(2x + 3) \\&= \frac{(2x + 3) - 3}{2} \\&= \frac{2x}{2} \\&= x\end{aligned}$$

And let's verify $f(f^{-1}(x)) = x$:

$$\begin{aligned}f(f^{-1}(x)) &= f\left(\frac{x - 3}{2}\right) \\&= 2\left(\frac{x - 3}{2}\right) + 3 \\&= (x - 3) + 3 \\&= x\end{aligned}$$

Since both compositions yield the identity function, $f^{-1}(x) = \frac{x-3}{2}$ is indeed the inverse of $f(x) = 2x + 3$.

Properties of the Inverse Function

Let $f : X \rightarrow Y$ be a function.

- Assume $x \sim y$ if $f(x) = f(y)$ so is $\sim$ an equivalence relation.
- Consider the quotient set $X_f := X / \sim$, where $\sim$ is the equivalence relation defined by $x \sim x' \iff f(x) = f(x')$. Let $q_f : X \rightarrow X_f$ be the canonical projection defined by $q_f(x) = [x]_\sim$, and let $\iota_f : f(X) \rightarrow Y$ be the inclusion map, $y \mapsto y$. Then the function

$$\hat{f} : X_f \rightarrow f(X), \quad [x]_\sim \mapsto f(x)$$

is a bijection, and the original map f can be written as the composition

$$f = \iota_f \circ \hat{f} \circ q_f.$$

Here, q_f is surjective, $\hat{f}$ is bijective, and ι_f is injective. This yields the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow q_f & & \uparrow \iota_f \\ X_f & \xrightarrow{\hat{f}} & f(X) \end{array}$$

Left/Right Inverse

Given $f : M \rightarrow N$, then

- f is *injective* if there exists a function $g : N \rightarrow M$ with the property $g \circ f = \text{Id}_M$. Each g is surjective and called the *left inverse* of f .
- f is *surjective* if there exists a function $\gamma : N \rightarrow M$ with the property $f \circ \gamma = \text{Id}_N$. Each γ is injective and called the *right inverse* of f .

Example:

Given $f : M \rightarrow N$ having a right inverse $\gamma : N \rightarrow M$ guarantees the existence of the solution of $f(m) = n$. Set $m = \gamma(n)$ then $f(m) = f(\gamma(n)) = f \circ \gamma = \text{Id}_N(n) = n$.

Proof I:

$\rightarrow$: f is injective, and we want to proof that from $g \circ f = \text{Id}_M$ then g is surjective.

Because f is injective we know that it maps all of N . Suppose, that for some $n \in N, n \notin f(M)$. We also know that $g \circ f = \text{Id}_M$ by definition. Now with our assumption and this knowledge

$$\begin{aligned} m &= \text{Id}_M \\ &= g(f(m)) \\ &= g(n), \text{ for some } n \in N \\ &= m \end{aligned}$$

Therefore, such an n can not exist because of our conditions.

$\leftarrow$: Because g is surjective, we know that all $m \in M$ are mapped and because of the composition and its identity property it is clear that f is injective because $m = \text{Id}_M(m) \in M$ else the identity property would not hold.

4.0.27 Transformations of a Function

Transformations modify the appearance of a function's graph without altering its basic shape. Here, we examine how different algebraic changes to a function $f(x)$ affect its graph:

Vertical Translation: $f(x) + a$

- Shifts the graph *upward* if $a > 0$, and *downward* if $a < 0$.
- Each point on the graph moves vertically by a units.

Horizontal Translation: $f(x + a)$

- Shifts the graph *left* if $a > 0$, and *right* if $a < 0$.
- This is opposite of what might be expected: adding to x shifts the graph in the negative direction.

Vertical Scaling (Stretch/Compression): $af(x)$

- If $|a| > 1$: the graph is *stretched* vertically (taller and narrower).
- If $0 < |a| < 1$: the graph is *compressed* vertically (shorter and wider).
- If $a < 0$: includes a reflection across the x -axis.

Horizontal Scaling (Stretch/Compression): $f(ax)$

- If $|a| > 1$: the graph is *compressed* horizontally (narrower).
- If $0 < |a| < 1$: the graph is *stretched* horizontally (wider).
- If $a < 0$: includes a reflection across the y -axis.

Reflection across the x -axis: $-f(x)$

- Flips the graph upside-down over the x -axis.
- Each point (x, y) becomes $(x, -y)$.

Reflection across the y -axis: $f(-x)$

- Flips the graph left-to-right over the y -axis.
- Each point (x, y) becomes $(-x, y)$.

4.0.28 Counting Functions

Given $f : A \rightarrow B$ with $|A| = n$ and $|B| = m$, we can count the number of different types of functions from A to B :

- *Total functions:* There are m^n total functions from A to B . Each of the n elements in A can be mapped to any of the m elements in B .
- *Injective functions:* If $n \leq m$, the number of injective functions is given by $m \times (m - 1) \times (m - 2) \times \dots \times (m - n + 1)$. This is because the first element in A has m choices, the second has $m - 1$ choices, and so on.
- *Surjective functions:* The number of surjective functions can be calculated using the principle of inclusion-exclusion:

$$\text{Number of surjective functions} = \sum_{k=0}^m (-1)^k \binom{m}{k} (m - k)^n$$

This formula counts all functions and subtracts those that miss at least one element in B .

- *Bijective functions:* If $n = m$, the number of bijective functions (one-to-one and onto) is $n!$ (n factorial), which counts all possible permutations of the n elements in A to the n elements in B .

4.0.29 Counting Relations

Given two sets A and B with $|A| = n$ and $|B| = m$, we can count the number of different types of relations from A to B :

- *Total relations:* There are $2^{n \times m}$ total relations from A to B . Each of the $n \times m$ possible ordered pairs (a, b) can either be included in or excluded from the relation. This comes from the cardinality of the power set of the Cartesian product of $A \times B$.
- *Reflexive relations:* If $A = B$, the number of reflexive relations is $2^{n^2 - n}$. This is because the n diagonal pairs (a, a) must be included, while the remaining $n^2 - n$ pairs can be chosen freely.
- *Symmetric relations:* If $A = B$, the number of symmetric relations is $2^{\frac{n(n+1)}{2}}$. This counts the diagonal pairs and half of the off-diagonal pairs, as each off-diagonal pair (a, b) must be paired with (b, a) .
- *Antisymmetric relations:* If $A = B$, the number of antisymmetric relations is $2^{n^2 - \frac{n(n-1)}{2}}$. This counts all pairs except for those that would violate antisymmetry.
- *Transitive relations:* Counting transitive relations is more complex and does not have a simple closed-form formula. It often requires combinatorial methods or recursive counting techniques.

Chapter 5

Mathematical Proofs

In this section I will provide with some examples of different types of proofs.

5.0.1 Proof by Direct Argument

Assume A to prove to prove B .

$$A \implies B$$

Example I:

For any integer n , if n is even, then n^2 is even. We will prove this theorem by direct argument.

Assume n is an even integer. Then we can write $n = 2k$ for some integer k .

Now, we compute n^2 :

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

This shows that n^2 is even, as it can be expressed as $2m$ where $m = 2k^2$ is an integer. Therefore, we conclude that if n is even, then n^2 is even.

□◻◻

5.0.2 Proof by Contradiction

Assume the negation of A to derive a false statement.

$$A \implies B \iff \neg A \implies F$$

Example I:

If n is an integer such that n^2 is even, then n is even.

We will prove this theorem by contradiction. Assume that n is an integer such that n^2 is even, but n is odd. Then we can write $n = 2k + 1$ for some integer k .

Now, we compute n^2 :

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$$

This shows that n^2 is odd, which contradicts our assumption that n^2 is even. Therefore, our assumption that n is odd must be false, and thus, n must be even.

□◻◻

Example II:

Another example is the proof that there are infinite prime numbers.

Let us define the prime numbers $p_1, \dots, p_n$ where we assume p_n is the biggest prime number. If we construct a number

$$q = 1 + p_1 \cdot \dots \cdot p_n$$

This number is bigger than p_n , but it is not divisible by any of the other prime numbers will give us always a residual of 1. Therefore, the assumption that there is the biggest prime number is false.

□◻◻

Example II:

Suppose, for contradiction, that $\sqrt{2}$ is rational. Then there exist integers p and q with $q \neq 0$, $\gcd(p, q) = 1$, and

$$\sqrt{2} = \frac{p}{q}.$$

Squaring both sides gives

$$2 = \frac{p^2}{q^2} \implies p^2 = 2q^2.$$

Thus, p^2 is even, so p is even. Write $p = 2k$ for some integer k . Then

$$(2k)^2 = 2q^2 \implies 4k^2 = 2q^2 \implies q^2 = 2k^2,$$

so q^2 is even and hence q is even. Therefore, both p and q are even, contradicting $\gcd(p, q) = 1$. This contradiction shows $\sqrt{2}$ is irrational.

□◻◻

5.0.3 Proof by Induction

For all $n \in \mathbb{N}$, the sum of the first n positive integers is given by:

$$S(n) = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

We will prove this theorem by induction on n .

Base Case: For $n = 1$:

$$S(1) = 1 = \frac{1(1+1)}{2}$$

The base case holds.

Inductive Step: Assume that the statement holds for some $n = k$, i.e., assume that:

$$S(k) = 1 + 2 + 3 + \dots + k = \frac{k(k+1)}{2}$$

We need to show that the statement holds for $n = k + 1$:

$$S(k+1) = S(k) + (k+1)$$

By the inductive hypothesis, we have:

$$\begin{aligned}
 S(k+1) &= \frac{k(k+1)}{2} + (k+1) \\
 &= \frac{k(k+1)}{2} + \frac{2(k+1)}{2} \\
 &= \frac{k(k+1) + 2(k+1)}{2} \\
 &= \frac{(k+1)(k+2)}{2}
 \end{aligned}$$

Thus, the statement holds for $n = k + 1$. By the principle of mathematical induction, the statement holds for all $n \in \mathbb{N}$.

□◻◻

5.0.4 Strong Induction

Strong induction is a form of mathematical induction where instead of assuming $A(n)$ is true in our inductive hypothesis, we assume $A(j)$ is true for $1 \leq j \leq n$.

The inductive step: Show $[A(1) \wedge A(2) \dots A(n)] \implies A(n+1)$.

Sometimes, we will prove specific values are true for $1 \leq j \leq n$, and sometimes just assume that $A(j)$ is true for $1 \leq j \leq n$.

Example:

Prove that any integer ≥ 1 can be written as the product of primes.

Let $A(n)$ be that if n is an integer greater than 1, then n can be written as the product of primes.

Basis: $A(2)$ is true because 2 is prime

Inductive hypothesis: $A(j)$ for $2 \leq j \leq n$ is true. **Important!**

Consider the cases $k+1$ is prime or composite.

The first case is true. For the case $n+1$ is not prime then $n+1 = ab$ with $2 \leq a, b \leq n < n+1$ and both for a and b we know that they can be written as the product of primes thus $n+1$ can be written as the product of primes.

□◻◻

5.0.5 Proof by Exhaustion

The only integer solutions to the equation $x^2 + y^2 = 1$ are $(0, 1), (1, 0), (0, -1), (-1, 0)$.

We will prove this theorem by exhaustion. We will check all possible integer values of x and y such that $x^2 + y^2 = 1$.

The possible integer values for x and y are $-1, 0, 1$. We will check each case:

If $x = 0$:

Then $y^2 = 1$ gives $y = 1$ or $y = -1$.

Solutions: $(0, 1), (0, -1)$.

If $x = 1$:

Then $y^2 = 0$ gives $y = 0$.

Solution: $(1, 0)$.

If $x = -1$:

Then $y^2 = 0$ gives $y = 0$.

Solution: $(-1, 0)$.

Thus, the only integer solutions to the equation are:

$$(0, 1), (1, 0), (0, -1), (-1, 0)$$

□◻◻

5.0.6 Proof by Cases

For any integer n , n^2 is even if and only if n is even.

We will prove this theorem by cases.

Case 1: Assume n is even. Then we can write $n = 2k$ for some integer k .

Now, we compute n^2 :

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

This shows that n^2 is even.

Case 2: Assume n is odd. Then we can write $n = 2k + 1$ for some integer k .

Now, we compute n^2 :

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1$$

This shows that n^2 is odd.

Since both cases have been considered, we conclude that n^2 is even if and only if n is even.

□◻◻

5.0.7 Proof by Construction

There exists an irrational number x such that x^2 is rational.

We will construct an irrational number x such that x^2 is rational.

Let $x = \sqrt{2}$. We know that $\sqrt{2}$ is irrational. Now, we compute x^2 :

$$x^2 = (\sqrt{2})^2 = 2$$

Since 2 is a rational number, we have constructed an irrational number $x = \sqrt{2}$ such that $x^2 = 2$ is rational. Therefore, the theorem is proved.

□◻◻

5.0.8 Proof by Counterexample

The statement all prime numbers are odd is false.

To prove this theorem, we will provide a counterexample.

The number 2 is a prime number, as its only divisors are 1 and 2. However, 2 is even, which contradicts the statement that all prime numbers are odd.

Therefore, the statement All prime numbers are odd is false.

□◻◻

5.0.9 Proof by Contrapositive

$$A \implies B \iff \neg B \implies \neg A$$

If n is an integer such that n^2 is odd, then n is odd.

We will prove this theorem by contrapositive. The contrapositive of the statement is: If n is an integer such that n is even, then n^2 is even.

Assume n is even. Then we can write $n = 2k$ for some integer k .

Now, we compute n^2 :

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

This shows that n^2 is even.

Since the contrapositive statement is true, the original statement If n^2 is odd, then n is odd is also true.

□◻◻

5.0.10 Proof by Reduction to Absurdity

The square root of 2 is irrational.

We will prove this theorem by reduction to absurdity. Assume that $\sqrt{2}$ is rational. Then we can write:

$$\sqrt{2} = \frac{p}{q}$$

Where p and q are integers with no common factors (i.e., the fraction is in the simplest form).

Squaring both sides gives:

$$2 = \frac{p^2}{q^2}$$

Rearranging gives:

$$p^2 = 2q^2$$

This implies that p^2 is even, and therefore, p must be even (since the square of an odd number is odd). Let $p = 2k$ for some integer k . Substituting this back into the equation gives:

$$\begin{aligned}(2k)^2 &= 2q^2 \\ 4k^2 &= 2q^2 \\ 2k^2 &= q^2\end{aligned}$$

This implies that q^2 is even, and therefore, q must also be even. Since both p and q are even, they have a common factor of 2, which contradicts our assumption that p and q have no common factors. Therefore, our assumption that $\sqrt{2}$ is rational must be false, and thus, $\sqrt{2}$ is irrational.

□◻◻

5.0.11 Proof by Analogy

The set of rational numbers is dense in the set of real numbers.

We will prove this theorem by analogy.

Consider the set of rational numbers $\mathbb{Q}$ and the set of real numbers $\mathbb{R}$. The density of $\mathbb{Q}$ in $\mathbb{R}$ means that between any two real numbers, there exists a rational number. For example, between the real numbers 1 and 2, we can find the rational number $\frac{3}{2} = 1.5$. Similarly, between any two real numbers a and b (where $a < b$), we can find a rational number $r = \frac{a+b}{2}$.

This shows that the set of rational numbers is dense in the set of real numbers. Therefore, the theorem is proved by analogy.

□◻◻

Chapter 6

Graph Theory

A Graph $G = \langle V, E \rangle$ consists of a finite set of vertices V and a relation $E \subseteq V \times V$ over the set. Depending on the traits of E it can be categorized into

- *Undirected*: E is symmetric.
- *Directed*: E is not symmetric.
- *Complete*: $E = V \times V$ for directed and $|E| \approx |V|(|V| - 1)/2$ for undirected graphs.
- *Loop-free*: $\forall (v, v) \in V \times V : (v, v) \notin E$
- *Weighted*: $G = \langle V, E, \phi \rangle$, where $\phi : E \rightarrow R$ and R is an arbitrary set. It mostly refers to adding a cost to each edge.
- *Dense*: $|E| \approx |V|^2$ for directed and $|E| \approx |V|(|V| - 1)/2$ for undirected graphs.
- *Sparse*: $|E| \ll |V|^2$ for directed and $|E| \ll |V|(|V| - 1)/2$ for undirected graphs.

In computer science the density of a graph is one of the main criteria when modeling a graph.

6.0.1 Connected Graphs

A graph $G = (V, E)$ undirected, then it is called *connected* if for all pairs $(v_i, v_j) \subseteq V \times V$ there exists a path from v_i to v_j . We define E_S the symmetric relation.

$$(v_1, v_2) \in E \iff (v_1, v_2) \in E_S \wedge (v_2, v_1) \in E_S$$

If G is directed then it is connected if and only if its undirected version is connected.

6.0.2 Paths

Given a graph G , we call a *path* a set of vertices if

$$\forall i \in \{1, \dots, k-1\} : (w_i, w_{i+1}) \in E$$

The most common types of paths are

- *Circle*: $w_1 = w_k$
- *Eulerian Circuit*: Each edge will be visited one time. $\forall (s, t) \in E : \exists_1 i : s = w_i \wedge t = w_{i+1}$
- *Hamiltonian Circuit*: Each vertex will be visited one time $\forall v \in V : \exists_1 i : v = w_i$
- *Loop-free*: $\forall (v, v) \in V \times V : (v, v) \notin E$

Path Cost

We define the *path cost* for a weighted graph as

$$\phi(p) = \sum_{i=1}^{k-1} \phi(w_i, w_{i+1})$$

6.0.3 The Adjacency Matrix

For a graph G with vertices $V = \{v_1, \dots, v_n\}$, we define *the adjacency matrix* $A = (a_{ij})$ as an $n \times n$ matrix with

$$a_{ij} = \begin{cases} 1 & \text{if } (v_i, v_j) \in E \\ 0 & \text{else} \end{cases}$$

6.0.4 The Relax Operation

In graph theory when referring to the action of updating the cost of visiting a vertex in a weighted graph we call this operation *relaxation*.

$$\text{relax}(u, v, \text{weight}) = \begin{cases} d[u] + \text{weight}(u, v) & \text{if } d[u] + \text{weight}(u, v) < d[v] \\ d[u] & \text{else} \end{cases}$$

Chapter 7

Order

7.0.1 Partially Ordered Classes

A *partially ordered class* is a pair of objects $\langle A, R \rangle$ where A is a class and R is an order relation in A . Remember that an order relation is one which is : reflexive, antisymmetric and transitive.

7.0.2 Lexicographical Ordering

Let A and B be partially ordered classes; by the *lexicographical ordering* of $A \times B$ we mean the following order relation in $A \times B$: If $(a_1, b_1), (a_2, b_2) \in A \times B$ and $(a_1, b_1) \leq (a_2, b_2)$ if and only if

$$a_1 < a_2$$

or

$$a_1 = a_2 \wedge b_1 \leq b_2$$

7.0.3 Antilexicographical Ordering

Let A and B be partially ordered classes; by the *antilexicographical ordering* of $A \times B$ we mean the following order relation in $A \times B$: If $(a_1, b_1), (a_2, b_2) \in A \times B$ and $(a_1, b_1) \leq (a_2, b_2)$ if and only if

$$b_1 < b_2$$

or

$$b_1 = b_2 \wedge a_1 \leq a_2$$

7.0.4 Comparable

Let A be a partially ordered class. Two elements $x, y \in A$ are considered *comparable* if either $x \leq y$ or $y \leq x$; otherwise, they are said to be *incomparable*.

7.0.5 Fully Ordered Subclass

Let A be a partially ordered class, and let $B \subseteq A$. If two elements of B are comparable, then we call B a *fully ordered subclass* of A , or *linearly ordered subclass* of A , or more commonly, a *chain* of A . If every two elements of A are comparable, then A is also considered as such.

Example:

Let C and D be chains of A and B respectively, prove that $A \times B$ lex. sorted $\implies C \times D$ is chain of $A \times B$.

Proof:

By the contrapositive, if $C \times D$ is not a chain of $A \times B$, then there some element (c, d) can not be compared, because (c, d) can not be compared, but $(c, d) \in A \times B$, then $A \times B$ can not lexicographically sorted.

□◻◻

7.0.6 Initial Segments

Let A be a partially ordered class and suppose $a \in A$. The *initial segment of A determined by a* is the class S_a , defined as follows:

$$S_a = \{x \in A \mid x < a\}.$$

Initial segments are transitive, which means that if we have an initial segment of an initial segment, then the last one is also an initial segment of the parent class.

An initial segment of an initial segment of a class is an initial segment of the same class

Proof:

Let $P = \{x \in A \mid x < a\}$ for some $a \in A$ and let $Q = \{p \in P \mid p < c\}$ for some $c \in P$. By definition $Q \subseteq A$. Because $c \in A$ it is clear that Q is an initial segment of A .

□◻◻

7.0.7 Cuts

If A is a partially ordered class, then a *cut* of A is a pair (L, U) of nonempty subclasses of A with the following properties:

- $L \cup U = \emptyset \wedge L \cup U = A$.
- If $x \in L \wedge y \leq x \implies y \in L$.
- If $x \in U \wedge y \geq x \implies y \in U$.

The two second conditions simply tell us that, we can compare its elements.

Example:

Given the partially ordered classes A and B , and let $A \times B$ be ordered antilexicographically. Prove that (L, U) is a cut of $B \implies (A \times L, A \times U)$ is a cut of $A \times B$

Proof:

Let (L, U) be a cut of B and define

$$L' = A \times L, \quad U' = A \times U.$$

We show (L', U') is a cut of $A \times B$ under the antilexicographical order.

Nonempty, disjoint, and covering:

Suppose $L' \cap U' \neq \emptyset$. Then,

$$\exists(x, y) \in L' \cap U' : x \in A \wedge (y \in L \vee y \in U)$$

Hence, $L \cap U \neq \emptyset$, a contradiction. Therefore, they must be disjoint.

For the covering, we use the distributive property of the Cartesian product under set union, $A \times (L \cup U) = A \times L \cup A \times U = A \times B$.

Downward closure of L' :

Let $(a_1, b_1) \in A \times L$ and $(a_2, b_2) \leq (a_1, b_1)$.

By antilexicographical order,

$$b_2 < b_1 \text{ or } (b_2 = b_1 \wedge a_2 \leq a_1).$$

Since $b_1 \in L$ and L is downward closed in B , we have $b_2 \in L$. Hence $(a_2, b_2) \in A \times L$.

Upward closure of U' :

Let $(a_1, b_1) \in A \times U$ and $(a_1, b_1) \leq (a_2, b_2)$. Then

$$b_1 < b_2 \text{ or } (b_1 = b_2 \wedge a_1 \leq a_2).$$

Since $b_1 \in U$ and U is upward closed in B , we have $b_2 \in U$. Thus $(a_2, b_2) \in A \times U$.

Therefore $(A \times L, A \times U)$ is a cut of $A \times B$.

□◻

7.0.8 Order Preserving Functions

Let A and B be partially ordered classes; a function $f : A \rightarrow B$ is said to *increasing* or *decreasing*, if it satisfies the following condition: For every two elements $x, y \in A$,

$$x \leq y \implies f(x) \leq f(y).$$

7.0.9 Isomorphism

If f is bijective and $x \leq y \iff f(x) \leq f(y)$ for every two elements $x \in A, y \in B$, we call it an *isomorphism*.

7.0.10 Strictly Increasing Function

We consider f strictly increasing if it satisfies that:

$$x < y \implies f(x) < f(y).$$

Example:

Prove that if $f : A \rightarrow B$ is an injective, increasing function, then it is strictly increasing.

Proof:

Suppose it is not increasingly increasing. The negation is $x < y \implies f(x) \geq f(y)$.

Now we have that $x \leq y \implies f(x) \geq f(y) \wedge x \leq y \implies f(x) \leq f(y) \iff f(x) = f(y)$ for some $x, y \in A$, hence we have a contradiction to the injectivity.

□◻

7.0.11 Order by Inverses

Let A and B be partially ordered classes and let $f : A \rightarrow B$ be a bijective function. Then f is an isomorphism if and only if $f : A \rightarrow B$ and $f^{-1} : B \rightarrow A$ are increasing functions.

Proof:

$$f(x) \leq f(y) \implies f^{-1}(f(x)) \leq f^{-1}(f(y)) \implies x \leq y$$

By the isomorphic property we get the increasing property for f and also for f^{-1} because else for $f(x) \leq f(y) : y \leq x$ violates our initial condition that $x \leq y$

$$x \leq y \implies f(x) \leq f(y) \implies f^{-1}(f(x)) \leq f^{-1}(f(y))$$

Thus, f and f^{-1} are increasing functions.

□◻◻

7.0.12 Common Isomorphism on Partially Ordered Classes

Let A , B and C be partially ordered classes.

- $I_A : A \rightarrow A$ with $I_A(x) = x$, $x \in A$ is an isomorphism.
- If $f : A \rightarrow B$ is an isomorphism, then $f^{-1} : B \rightarrow A$ is an isomorphism.
- If $f : A \rightarrow B$ and $g : B \rightarrow C$ is an isomorphism, then $g \circ f : A \rightarrow C$ is an isomorphism.

Proof I:

Let $I_A = f$, for $x, y \in A$ we have that $x \leq y$ gives us $x < y \iff f(x) < f(y)$ and $x = y \iff f(x) = f(y)$.

□◻◻

Proof II:

In one of our previous results we got that f^{-1} is increasing if and only if f is an isomorphism. We prove that f^{-1} is also one by assuming it is not, then $\exists y \in B : f(x) \neq y, \forall x \in A$ which is a contradiction.

□◻◻

Proof III:

We keep in mind the bijectivity of f and g and state the following

$$x \leq y \iff f(x) \leq f(y) \iff g(f(x)) \iff (f(y))$$

□◻◻

Example:

Given the isomorphism $f : A \rightarrow B$, if (L, U) is a cut of A , then $(f(L), f(U))$ is a cut of B .

Proof:

Suppose $(f(L), f(U))$ is not a cut, then $y \in f(L) \cap f(U)$, then $y \in f^{-1}(f(L)) \cap f^{-1}(f(U))$ but this is a contradiction. Therefore, $(f(L), f(U))$ must be disjoint.

For the covering, we use a similar argument $f(L) \cup f(U) \neq B$, then f would not be bijective, hence, another contradiction.

For the downward and upward closure, we know that our function targets the whole set B and by its isomorphic properties it respects the order. Therefore, this properties are also satisfied.

□◻◻

7.0.13 Isomorphic Definition

Let A , and B be partially ordered classes. If there exist an order preserving isomorphism between the two classes, we say that A is *isomorphic* with B .

Note, that the isomorphic property allows us to define an equivalence relation between classes.

7.0.14 Convex

Let A be a partially ordered class and $C \subseteq A$. If for $a \in C$ and $b \in C$ the following holds $a \leq x \leq b, \forall x \in C$, then we consider C *convex*.

Properties

Let A , and B be partially ordered classes

- If $C \subseteq B$ is convex and $f : A \rightarrow B$ is an increasing function, then $f^{-1}(C)$ is also convex.
- If H is the functional equivalence relation, and $f : A \rightarrow B$ is an increasing function, then all equivalence classes of A/H are convex subsets of A .

Proof I:

Given $f(a), f(b) \in C$, then using the properties of the increasing functions we get $f^{-1}(f(a)) \leq f^{-1}(f(x)) \leq f^{-1}(f(b))$ which gives us $a \leq x \leq b$ which is the definition of a convex class.

□◻◻

Proof II:

Given the equivalence relation given by the function f we have that for every member of an arbitrary equivalence class $f(a) = f(x) = f(b)$ can be written as $f(a) \leq f(x) \leq f(b)$ hence we the definition of convex classes.

□◻◻

7.0.15 Maximal Element

An element $m \in A$ is considered *maximal element* of A under the partially order relation R if $\forall x \in A$

$$x \leq m$$

and if

$$x \geq m \implies x = m.$$

Or equivalently m under the relation rule does not relate to anything except itself:

$$\forall (m, y) \in R \implies m = y \text{ or equivalently } (x, m) \in R, \forall x \in A.$$

7.0.16 Minimal Element

An element $m \in A$ is considered *minimal element* of A under the partially order relation R if $\forall x \in A$

$$x \geq m$$

and if

$$x \leq m \implies x = m.$$

Or m is under the relation rule does not relate to anything except itself:

$$\forall (x, m) \in R \implies m = x \text{ or equivalently } (m, x) \in R, \forall x \in A.$$

7.0.17 Upper Bound

Let $B \subseteq A$. An *upper bound* of B in A is an element $a \in A$ such that $a \geq x, \forall x \in B$. Equivalently

$$\forall x \in B, (x, a) \in R.$$

Set of Upper Bounds

Let $B \subseteq A$, we define the function v as

$$v(B) := \{a \in A \mid a \geq x, \forall x \in B\}.$$

7.0.18 Lower Bound

Let $B \subseteq A$. A *lower bound* of B in A is an element $a \in A$ such that $a \leq x, \forall x \in B$. Equivalently

$$\forall x \in B, (a, x) \in R$$

Set of Lower Bounds

Let $B \subseteq A$, we define the function λ as

$$\lambda(B) := \{a \in A \mid a \leq x, \forall x \in B\}.$$

7.0.19 Maximum

An element x is a *maximum* of a partially ordered class A if

$$(a, x) \in R, \forall a \in A.$$

7.0.20 Minimum

An element x is a *minimum* of a partially ordered class A if

$$(x, a) \in R, \forall a \in A.$$

7.0.21 Supremum and Infimum

Given a partially ordered class $B \subseteq A$, the least upper bound is called the *supremum* $s \in A$ and defined as:

$$\forall b \in B, (b, s) \in R \text{ and } \forall a \in A, [(\forall b \in B, (b, a) \in R) \implies (s, a) \in R],$$

and the greatest lower bound is its *infimum* $i \in A$ is defined as:

$$\forall b \in B, (i, b) \in R \text{ and } \forall a \in A, [(\forall b \in B, (a, b) \in R) \implies (a, i) \in R].$$

7.0.22 Duality

Let $\mathcal{S}$ be a statement about partially ordered classes with $\geq, \leq$. If we replace the occurrences of $\geq, \leq$ with their contrary, then the resulting statement $\mathcal{S}'$ is called the *dual* of $\mathcal{S}$.

We know that given an order relation G on A , then G^{-1} is also an order relation on A . For the dual this means that for any true statement on G the dual of the statement is also true for G^{-1} .

7.0.23 Greatest and Least Elements under Inclusion

If A has a greatest element a , and B has a greatest element b , and $B \subseteq A$, then

$$b \leq a.$$

Proof:

$$a \geq x, \forall x \in A \implies a \geq b, \forall b \in B \implies a \geq b$$

□◻◻

Dual: If A has a least element a , and B has a least element b , and $B \subseteq A$, then

$$b \geq a.$$

7.0.24 Upper and Lower Bounds under Inclusion

Let B and C be subclasses of A . If $B \subseteq C$, then

$$v(C) \subseteq v(B).$$

Proof:

$x \in v(C) \implies x \geq y, \forall y \in C$ because $y \in B \implies b \in C$, then $x \geq y, \forall y \in B \implies x \in v(B)$.

The opposite is not always true, due to the fact that if $z \in v(B)$, such that $z \in C$, but $z \notin v(C)$, which is possible if z is not an upper bound of C , then the theorem does not hold.

□◻◻

Dual: If $B \subseteq C$, then

$$\lambda(C) \subseteq \lambda(B).$$

7.0.25 Suprema and Infima under Inclusion

Let B and C be subclasses of A , and suppose that B and C each has a sup in A . If $B \subseteq C$, then

$$\sup B \leq \sup C.$$

Proof:

By the dual of the greatest element under inclusion applied to $v(C) \subseteq v(B)$,

$$\sup(B) \leq \sup(C).$$

□◻◻

Dual: If B and C each has an inf in A , and if $B \subseteq C$, then

$$\inf B \geq \inf C.$$

7.0.26 Relation between a Set and Bounds of Bounds

Let B be a subclass of A . Then

$$B \subseteq v(\lambda(B)).$$

Proof: Suppose $x \in B$; for each $y \in \lambda(B)$, $y \leq x$; hence $x \in v(\lambda(B))$.

□◻◻

Dual:

$$B \subseteq \lambda(v(B)).$$

7.0.27 Infimum via Lambda Function

Let $B \subseteq A$, then

$$\sup(\lambda(B)) \in A \implies \inf(B) \in A \wedge \inf(B) = \sup(\lambda(B))$$

Dual: $\inf(v(B)) \in A \implies \sup(B) \in A \wedge \sup(B) = \inf(v(B))$.

Proof:

Let $s = \sup(\lambda(B))$, $\forall b \in B, s \leq b$; hence, s is a lower bound of B . Suppose $\inf(B) \neq s$, then $s > \inf(B) \vee s < \inf(B)$.

Case $s > \inf(B)$: this leads to a contradiction because if $s \in A$ and s is a lower bound of B , but it is greater than its infimum then $\inf(B)$ can not be the greatest lower bound of B , which does not make sense by the definition.

Case $s < \inf(B)$: $\inf(B)$ is a lower bound of B ; therefore, $\inf(B) \in \lambda(B)$. If $s < \inf(B)$ then s can not be the supremum of $\lambda(B)$ since there exists a greater element in it; hence $s = \inf(B)$.

□

Example Exercises

1. Prove that given $B \subseteq A$, $\sup(B) = b \in A \implies \lambda(v(B)) \cap v(B) = \{b\}$

Proof:

By definition $b \in v(B)$. Also, by the definition of $\lambda(v(B))$:

$$\{a \in A \mid x \leq b, \forall b \in v(B)\};$$

we can guarantee that $b \in \lambda(v(B))$. It is leftover to prove that the intersection only consists of b .

Suppose that there also exists another x in the intersection such that $x \neq b$, then $x \in v(B) \wedge x \in \lambda(v(B))$.

If $x \in v(B)$ and $x \neq b$, then $x < b$ or $x > b$ which when handled separately lead to the following conclusion:

If $x < b$, then b is not the least upper bound of B which is a contradiction to the definition of the supremum. Hence, by the $\wedge$ in our original statement, we arrive at the conclusion that such an x can not exist.

□

2. Let $B \subseteq A$, $C \subseteq A$, then $\sup(B) = \sup(C) \implies v(B) = v(C)$.

Proof:

By definition for $\sup(B) \geq b, \forall b \in B$, and also $\sup(B) \in v(B)$, but because it is the least upper bound we can write

$$v(B) = \{a \in A \mid a \geq \sup(B)\}$$

Therefore, if $\sup(B) = \sup(C) = s$, then $v(B) = \{a \in A \mid a \geq s\}$, and because they are shared then $v(C) = \{a \in A \mid a \geq s\}$

□

3. Suppose $B \subseteq A$; prove that $v(B) = v(\lambda(v(B)))$.

For $x \in v(B)$ holds that $x \geq b, \forall b \in B$, then for $y \in \lambda(v(B))$ holds $y \leq z, \forall z \in v(B)$ which implies that $y \leq b, \forall b \in B$. Finally, if $y \leq b, b \in B$, then for $w \in v(\lambda(v(B)))$ it must be true that $w \geq y, \forall y \in \lambda(v(B))$, which by transitivity implies that $w \geq b, \forall b \in B$; hence, $v(B) = v(\lambda(v(B)))$.

□

7.0.28 Bounded

Let A be a partially ordered class. If A has both a lower and an upper bound then it is *bounded*.

7.0.29 Conditionally Complete

Let A be a partially ordered class. If every non-empty subclass of A that is bounded has a sup, then A is said to be *conditionally complete*.

7.0.30 Lattices

Let A be a partially ordered class. If every doubleton $\{x, y\}$ in A has a sup and inf then A is called a *lattice*.

When dealing with lattices, we denote $\sup(\{x, y\}) := x \vee y$ and $\inf(\{x, y\}) := x \wedge y$.

Properties

For $a, b \in A$

1. $a \leq a \vee b$ and $b \leq a \vee b$
2. $a \geq a \wedge b$ and $b \geq a \wedge b$
3. If $a \leq c$ and $b \leq c$, then $a \vee b \leq c$
4. If $a \geq c$ and $b \geq c$, then $a \wedge b \geq c$
5. $a \vee a = a$ and $a \wedge a = a$
6. $a \vee b = b \vee a$ and $a \wedge b = b \wedge a$
7. $(a \vee b) \vee c = a \vee (b \vee c)$ and $(a \wedge b) \wedge c = a \wedge (b \wedge c)$
8. $(a \vee b) \wedge a = a$ and $(a \wedge b) \vee a = a$

Proof 3:

To prove: $a \vee a = a$ and $a \wedge a = a$.

$a \vee a \geq x, \forall x \in \{a\}$. Using the definition of the infimum we can also prove the statement for infimum.

The other statements are easily proved by cases.

□◻

7.0.31 Definition of Less Than

Let A be a *lattice*, we define the symbol $\leq$ as:

$$x \leq y \iff x \vee y = y$$

Example I:

Given a lattice A with $\leq$, prove that it is an order relation.

Reflexivity: $(a, a) \in R$ because $a \leq a$ implies $a \vee a = a$.

Antisymmetry: $a \leq b \implies a \vee b = b$ if by supposition $b \leq a$, then $b \vee a = a$, but our property tells us that $a \vee b = b \vee a$; therefore, $a = b$.

Transitivity: $(a, b), (b, z) \in R \implies (a, z) \in R$, if $a \vee b = b$ and $b \vee z = z$, then $a \leq b$ and $b \leq z$, hence; $a \leq b \vee z = z \implies (a, z) \in R$.

□◻

Example II:

Given $a \leq b$ and $c \leq d$, prove that $a \vee c \leq b \vee d$.

Proceeding by cases note that: $a \geq c$ delivers $a \vee b \vee d$. If $b \geq c$, then $a \leq b$; if $d \geq b$ then $a \leq d$ by transitivity. Similarly, for the case $a \vee c = c$, then if $b \geq d$ gives us $c \leq b$; else if $b \leq d$ then $c \leq b \leq d$.

□◻◻

Distributive Laws for Lattices

- $x \wedge (y \vee z) \geq (x \wedge y) \vee (x \wedge z)$.
- $(x \vee y) \wedge z \leq (x \vee y) \wedge (x \vee z)$.

7.0.32 Sublattices

Let A be a lattice, and let B be a subclass of A . If

$$x \in B \text{ and } y \in B \implies x \vee y \in B \text{ and } x \wedge y \in B,$$

the B is called a *sublattice* of A .

7.0.33 Boolean Algebra

A *Boolean algebra* is defined to be a lattice A with the following additional properties:

1. $\exists 0 \in A$ and $1 \in A$ such that $x \vee 0 = x$ and $x \wedge 1 = x$.
2. $\forall x \in A, \exists \bar{x} \in A : x \wedge \bar{x} = 0$ and $x \vee \bar{x} = 1$.
3. $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ and $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ Let A be a partially ordered class; A is called a *complete lattice* if every subclass of A has a sup. Alternatively, if every subclass of A has an inf.

In other words 1 the biggest and 0 the smallest element in A .

Distributive Lattices

A *distributive lattice* is boolean algebra.

7.0.34 Fully Ordered Classes**7.0.35 Principle of the Well-ordering**

Every non-empty set of natural numbers has a minimum.

7.0.36 Order in Fields

An *ordered field* is a field $\mathbb{F}$ equipped with a total order $<$ that is compatible with the field operations $+$ and $\cdot$. That is, the order satisfies both algebraic and ordering properties.

Order Axioms for Fields

A field F is called an *ordered field* if it satisfies the following properties for all $a, b, c \in F$:

- I. *Trichotomy*: Exactly one of the following holds:

$$a < b, \quad a = b, \quad a > b$$

- II. *Transitivity*: If $a < b$ and $b < c$, then $a < c$.

- III. *Additive Compatibility*: If $a < b$, then $a + c < b + c$.

IV. *Multiplicative Compatibility*: If $0 < a$ and $0 < b$, then $0 < ab$.

These properties ensure that arithmetic operations respect the order structure.

Consequences of the Order Axioms

- $a < b \implies -b < -a$
- $a < b \wedge c < 0 \implies ac > bc$
- $a < 0, b < 0 \implies a + b < 0$ the contrary applies to the multiplication $ab > 0$
- $a > 0, b > 0 \implies a + b > 0$
- Squares are always non-negative: $a^2 \geq 0$
- The order is total: any two elements are comparable

Examples of Ordered Fields

- $\mathbb{Q}$: Rational numbers with the usual order
- $\mathbb{R}$: Real numbers with the usual order
- $\mathbb{C}$: Complex numbers are not an ordered field, since $i^2 = -1 < 0$ would violate positivity of squares

Chapter 8

The Natural Numbers

8.0.1 Definition as Sets

We define the members of $\mathbb{N}$ as:

- $0 := \emptyset$
- $1 := \{0\} = \{\emptyset, \{\emptyset\}\}$
- $2 := \{0, 1\} = \{\emptyset, \{\emptyset, \{\emptyset\}\}\}$
- $n := \{n_1, \dots, 0\}$

8.0.2 Addition

$$S(n) := n \cup n := n + 1$$

8.0.3 Inductive Sets

A set I is inductive if

$$0 \in I, x \in I \implies S(x) \in I.$$

8.0.4 Axiom of Infinity

An inductive set exists.

8.0.5 Definition of the Sets of Natural Numbers

$$\mathbb{N} := \{x \mid x \in I, \text{ for every inductive set } I\}$$

Proof that $\mathbb{N}$ is inductive for every inductive set $I, \mathbb{N} \subseteq I$:

$\forall I, \emptyset \in I$, hence $0 \in \mathbb{N}$, for $n \in \mathbb{N}$, let I be inductive then

$$S(n) \in I \therefore S(n) \in \mathbb{N}$$

8.0.6 The Standard Order

We define $<$ for the natural numbers as:

$$\forall m, n \in \mathbb{N}, m < n \iff m \in n.$$

Lemma

- $0 < n, \forall n \in \mathbb{N}$
- $\forall k, n \in \mathbb{N}, k < S(n) \iff k < n \vee k = n$

Proof:

Suppose for the sake of contradiction that $\exists n \in \mathbb{N}, n \neq 0 : n < 0$. By definition $n < 0 \iff n \in 0 = \emptyset$. Hence, we arrive at our desired contradiction.

For the second part of the lemma $k < S(n) \iff k < n \vee k = n$:

$$\begin{aligned}
 k < S(n) &\implies k \in \{n\} \cup n \iff k \in n \vee k = n \\
 &\iff k < n \vee k = n \\
 &\iff k < n
 \end{aligned}$$

□◻◻

8.0.7 The Peano Axioms

We will define the set of natural numbers, $\mathbb{N}$, via the following 9 axioms. These axioms are known as the *Peano Axioms*. The first 4 axioms define equality on the set $\mathbb{N}$.

- I. *Axiom of Reflexivity*: For every $x \in \mathbb{N}$, we have $x = x$.
- II. *Axiom of Symmetry*: For every $x, y \in \mathbb{N}$, if $x = y$ then $y = x$.
- III. *Axiom of Transitivity*: For every $x, y, z \in \mathbb{N}$, if $x = y$ and $y = z$ then $x = z$.
- IV. *Axiom of the Closure of Equality*: For all x, y , if $x \in \mathbb{N}$ and $x = y$, then $y \in \mathbb{N}$.

The remaining 5 axioms define the structure of $\mathbb{N}$:

- V. $1 \in \mathbb{N}$
- VI. If $x \in \mathbb{N}$, then the successor $S(x) \in \mathbb{N}$.
- VII. There is no $x \in \mathbb{N}$ such that $S(x) = 1$.
- VIII. For all $x, y \in \mathbb{N}$, if $S(x) = S(y)$, then $x = y$.
- IX. Let $P(x)$ be a statement about the natural number x . If:
 - $P(1)$ is true, and
 - for all $n \in \mathbb{N}$, if $P(n)$ is true, then $P(S(n))$ is also true,
 then $P(x)$ is true for all $x \in \mathbb{N}$.

(Mathematical Induction)

As shorthand, we denote:

$$S(1) = 2, \quad S(S(1)) = 3, \quad S(S(S(1))) = 4, \quad \text{and so on.}$$

8.0.8 Propositions and Proofs

Proposition 1: $n \neq m \implies S(n) \neq S(m)$

Proof:

We will prove this by contradiction. Assume $n \neq m$ and $S(n) = S(m)$. By Axiom 8, we have $n = m$, which is a contradiction. Therefore, $S(n) \neq S(m)$.

Proposition 2: For any $n \in \mathbb{N}$, $n \neq S(n)$

Proof:

$$M = \{n \in \mathbb{N} \mid n \neq S(n)\}$$

By $1 \neq S(1)$ for any $n \in \mathbb{N}$, this implies that 1 is part of the set M . This implies that $S(n) \neq S(S(n)) \implies S(n) \in M$ By Axiom 9 $M = \mathbb{N}$

Proposition 3: $n \neq 1 \implies \exists m \in \mathbb{N} \mid n = S(m)$

Proof:

$$M = \{1\} \cup \{n \in \mathbb{N} \mid \text{Proposition 3 is true}\}$$

We know that 1 is in the set M . And by proposition 1 we know that

$$S(n) = S(S(m)) \implies S(n) \in M$$

And by Axiom 9 we know that $M = \mathbb{N}$

8.0.9 Definition of Addition in $\mathbb{N}$

For any pair $n, m \in \mathbb{N}$ there is a unique way to define

$$\text{Add}(n, m) = n + m$$

1. **Base Case:** $n + 1 = S(n)$
2. **Inductive Step:** $n + S(m) = S(n + m) \iff S(n + m)$

Uniqueness:

Suppose: A & B satisfy our conditions. Fix n and then let $M = \{m \in \mathbb{N} \mid A(n, m) = B(n, m)\}$. Then

$$A(n, 1) = S(n) = B(n, 1) \implies 1 \in M$$

$$m \in M \implies A(n, m) = B(n, m) \implies A(n, S(m)) = S(A(n, m)) = S(B(n, m)) = B(n, S(m))$$

$$\implies A(n, S(m)) = B(n, S(m))$$

by Axiom 9 we know that $M = \mathbb{N}$ and $A = B$

Construction: For $n = 1$ Define $A(n, m) = S(m)$

1. $A(1, 1) = S(1) = S(1)$
2. $A(1, S(m)) = S(A(1, m)) = S(S(m))$

Define: $A(S(n), S(m)) = S(A(n, m))$

1. $A(S(n), 1) = S(A(n, 1)) = S(S(n))$
2. $A(S(n), S(m)) = S(A(n, m)) = S(S(m))$

Commutativity of Addition:

The proposition says $n + m = m + n$ for any $n, m \in \mathbb{N}$. We will prove this by induction on m . Fix n and consider $M = \{m \in \mathbb{N} \mid A(n, m) = A(m, n)\}$

Now recall that $A(n, 1) = S(n)$

For $n = 1$:

$$A(n, 1) = S(1) = A(1, n) \implies 1 \in \mathbb{N}$$

$$\text{also } A(n, k) = 1 + k \implies 1 + m = S(m) \implies 1 + m = m + 1 \implies 1 \in \mathbb{N}$$

$$\text{Suppose: } n \in \mathbb{N} \implies n + m = m + n \text{ or } A(n, m) = A(m, n)$$

$$\text{By construction } A(S(n), m) = S(A(n, m)) \text{ and by definition } A(S(n), m) = S(A(n, m)) = A(m, S(n)) = S(n) + m = m + S(n) \implies S(n) \in M \implies \text{by induction } M = \mathbb{N}.$$

Chapter 9

The Archimedean Principle

If a and b are real numbers with $a > 0$, then there exists $n \in \mathbb{N}$ such that $na > b$.

Equivalently, for any real number $x \in \mathbb{R}$, there exists a natural number $n \in \mathbb{N}$ such that $n > x = \frac{b}{a}$.

Proof:

We prove the Archimedean Principle by contradiction.

Assume that there exists some real number $x \in \mathbb{R}$ such that $n \leq x$ for all $n \in \mathbb{N}$. That is, x is an upper bound for the set $\mathbb{N} \subset \mathbb{R}$, baked up by the Completeness Axiom of the real numbers.

Let $s = \sup(\mathbb{N})$ be the least upper bound of $\mathbb{N}$. Then,

$$s - 1 < s \implies \exists m \in \mathbb{N} \text{ s. t. } m > s - 1$$

But, if we add 1 to both sides we get

$$s < m + 1 \in \mathbb{N}$$

This is a contradiction to the assumption that s is the supremum of the naturals. Hence there is not a real number x which is greater than all natural numbers.

□◻◻

9.0.1 Equivalent Formulation

For any $\varepsilon > 0$, there exists $n \in \mathbb{N}$ such that $\frac{1}{n} < \varepsilon$

9.0.2 Applications

1. **Density of Rational Numbers:** The Archimedean Principle helps in proving that between any two real numbers, there exists a rational number.
2. **Limits and Infinitesimals:** It ensures that sequences like $\left\{\frac{1}{n}\right\}$ converge to 0, foundational in real analysis and calculus.
3. **Bounding Functions:** It is used in analysis to show that functions do not grow faster than natural numbers in certain contexts.
4. **Non-Existence of Infinitely Small Numbers:** The principle implies that real numbers do not contain infinitesimals (nonzero numbers smaller than all $\frac{1}{n}$), distinguishing $\mathbb{R}$ from non-standard number systems.

Example:

Prove that $\forall \varepsilon > 0, \exists N \in \mathbb{N} : \frac{1}{n} < \varepsilon, \forall n \geq N$

We know that

$$\begin{aligned} \frac{1}{N} &< \varepsilon \\ \varepsilon &> \frac{1}{N} \geq \frac{1}{n} \text{ by transitivity} \\ \varepsilon &> \frac{1}{n} \end{aligned}$$

We also get $N > \frac{1}{\varepsilon}$.

And by the Archimedean Principle we let $b = 1$, $a = \varepsilon$ and $n = N$. Then by the transitivity we get out proof.

□◻◻

Chapter 10

Arithmetic

10.0.1 Even and Odd numbers

A natural number a is *even* if $a = 2k$ with $k \in \mathbb{N}$.

A natural number b is *odd* if $b = 2q - 1$ with $q \in \mathbb{N}$.

10.0.2 Perfect Squares

A natural number a has a *perfect square* if $\sqrt{a} \in \mathbb{N}$.

10.0.3 Division Algorithm

Given any two integers a and b with $a > 0$, there exists unique integers q and r such that $b = qa + r$ with $0 \leq r < a$.

Proof:

Let $S = \{b - xa \mid b - xa \geq 0, x \in \mathbb{Z}\}$

By the well ordering principle of the natural numbers we know that our set has minimum r .

First we prove $0 \leq r$

If $b \geq 0$ then let $x = 0 \implies b \in S$. For $b < 0$ then let $x = b$. Thus

$$b - ba = b(1 - a) \implies b - ba \in S$$

Now to prove that $r < a$, assume $r \geq a$, but then

$$\begin{aligned} b - qa &\geq a \\ b - qa - a &\geq 0 \\ b - a(q + 1) &\geq 0 \\ b - a(q + 1) &\geq 0 \implies b - a(q + 1) \in S \end{aligned}$$

And $b - a(q + 1) < b - aq = r \implies r < a$ therefore, we have a contradiction.

Now we will prove the uniqueness.

Assume $pq + r = a = pq' + r'$ with $q \neq q'$ and $r \neq r'$

$$r - r' = a(q - q') = 0$$

Using the property of inverses of $\mathbb{Z} \setminus \{0\}$ as a Field we know that $r = r'$ and $q = q'$.

10.0.4 Prime Numbers

A *prime number* $p \in \mathbb{N} \setminus \{1\}$ fulfills the following condition:

- I. p is only divisible by itself and 1.

10.0.5 Pythagorean Triplets

Pythagorean triplets are 3 natural numbers such that

$$a^2 + b^2 = c^2$$

10.0.6 Divisibility

A number $a \in \mathbb{Z}$ is divisible a number $b \in \mathbb{Z}$ if

$$a = kb \quad k \in \mathbb{Z}$$

10.0.7 Division

Given $a, b \in \mathbb{Z}$. There exists unique $q, r \in \mathbb{Z}$ such that

$$a = qb + r \quad 0 \leq r < |b|$$

10.0.8 Greatest Common Denominator

For two integers $a, b \in \mathbb{Z}$ with $(a, b) \neq (0, 0)$, the positive integer t satisfying

- $t \mid a$ and $t \mid b$,
- for all $c \in \mathbb{Z}$ with $c \mid a$ and $c \mid b$, it follows that $c \mid t$,

is called the *greatest common divisor* (gcd) of a and b denoted as $t = \gcd(a, b)$.

If $t = 1$, then a and b are said to be *coprime* or *relatively prime*.

10.0.9 The Euclidean Algorithm

If $a = bq + r$ then $\gcd(a, b) = \gcd(b, r)$.

Input: $a, b \in \mathbb{Z}$, $a \neq 0 \neq b$. **Output:** The greatest common divisor t of a and b .

Algorithm:

1. If $a = 0$, then return $|b|$. If $b = 0$, return $|a|$.
2. Set $r_1 := a$; $r_2 := b$; $k := 2$.
3. As long as $r_k \neq 0$, define $r_{k+1} := r_{k-1} \bmod r_k$.
4. As soon as $r_k = 0$, return r_{k-1} .

In simpler terms, as long the remainder is not zero, we keep dividing the previous remainder by the current remainder and updating the remainders. When we reach a remainder of zero, the previous remainder is the gcd.

Proof:

Notice, that the algorithm terminates because the sequence given r_n is monotone decreasing due to $0 \leq r < b$ and the well ordering principle of the natural numbers.

For the correctness: given is $a = bq + r$ to show that $\gcd(a, b) = \gcd(b, r)$. This is like saying

$$\begin{aligned}
\gcd(a, b) &= \gcd(b, r) \\
\gcd(b, r) &= \gcd(r, r_1) \\
\gcd(r, r_1) &= \gcd(r_1, r_2) \\
&\vdots \\
\gcd(r_{n-1}, r_n) &= \gcd(r_n, 0) = r_n
\end{aligned}$$

Let d be a common divisor of a, b . Then $d \mid a$ and $d \mid b$, so $d \mid ((a - bq) = r) \implies d \mid r$.

Let e be a common divisor of b, r . Then $e \mid b$ and $e \mid r$, so $e \mid ((bq + r) = a) \implies e \mid a$.

Thus, the set of common divisors of a, b is the same as the set of common divisors of b, r , so their greatest common divisors are equal.

□◻◻

Example:

Find $\gcd(252, 198)$:

$$\begin{aligned}
r_{n-1} &= q_{n-1}r_n + r_{n+1} \\
252 &= 1 \cdot 198 + 54 \\
198 &= 3 \cdot 54 + 36 \\
54 &= 1 \cdot 36 + 18 \\
36 &= 2 \cdot 18 + 0
\end{aligned}$$

So, $\gcd(252, 198) = 18$.

10.0.10 Euclidean Algorithm as Linear Map

Given $E := \{(a, b)^T \mid a, b \in \mathbb{Z}, b \neq 0\}$ and

$$\varepsilon : E \rightarrow \mathbb{Z}^{2 \times 1} : \begin{bmatrix} a \\ b \end{bmatrix} \mapsto \begin{bmatrix} b \\ a \bmod b \end{bmatrix}$$

Then the Euclidean algorithm can be described as repeated application of ε until the second component is zero. The first component at that point is the $\gcd$.

$$\varepsilon^{k-1} \begin{bmatrix} a \\ b \end{bmatrix} = \varepsilon \circ \dots \circ \varepsilon \begin{bmatrix} t \\ 0 \end{bmatrix}$$

$$t = \gcd(a, b).$$

Also, we can express ε as a matrix multiplication:

$$\varepsilon \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & -q \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} b \\ r \end{bmatrix}$$

with $a = qb + r$ and $q \in \mathbb{Z}, 0 \leq q < |b|$.

10.0.11 Bezout Identity

This is one of the most important properties of the $\gcd$. Given $a, b \in \mathbb{Z}$, not both zero, there exist integers $\alpha, \beta \in \mathbb{Z}$ such that

$$\gcd(a, b) = a\alpha + b\beta$$

This equation comes directly from *Euclid's Algorithm*. For the intuition, when writing your iterations you can manipulate them to arrive at the solution.

Proof:

Given

$$M := \{a\alpha + b\beta \mid a\alpha + b\beta > 0 \wedge \alpha \in \mathbb{Z} \wedge \beta \in \mathbb{Z}\}$$

M is not empty because of $\alpha, \beta \in M$, and it has a minimum by the well ordering principle of the natural numbers which we will call c .

Then, let $k \in M$. Then there exist $q \in \mathbb{N}_0$ and $r \in \mathbb{N}_0$ with $r < c$ such that

$$k = qc + r$$

because k and c are elements of M we should be able to find r_1, r_2, s_1, s_2 such that

$$k = r_1a + s_1b \quad c = r_2a + s_2b$$

Now let us

$$\begin{aligned} k &= r_1a + s_1b = qc + r \\ r_1a + s_1b &= q(r_2a + s_2b) + r \\ r &= r_1a + s_1b - q(r_2a + s_2b) \\ r &= r_1a + s_1b - qr_2a - qs_2b \\ r &= (r_1 - qr_2)a + (s_1 - qs_2)b \end{aligned}$$

$r \geq 0$ and $r < c$, but if $r \neq 0$ then $r \in M$ would be the minimum which is a contradiction to the assumption that c is the minimum, also $r = 0$, $k = qc$. Thus, *every element of M is multiplicity of c* . Therefore, a, b are also divisible by c .

Now we need to show that c is in fact the greatest common divisor. $c = r_2a + s_2b$

$$c = a\alpha_0 + b\beta_0$$

Because $c \mid a \wedge c \mid b$ we know that $c \mid a\alpha_0 + b\beta_0$. We also know that $c \mid \gcd(a, b)$ and thus, $|c| = |g| = \gcd(a, b)$ but because both are positive $c = g$.

□◻◻

10.0.12 Euclidean Algorithm as Matrix Multiplication

Input: $a, b \in \mathbb{Z}$, $a \neq 0 \neq b$.

Output: $t = \gcd(a, b)$ and $\alpha, \beta \in \mathbb{Z}$ with $\alpha a + \beta b = t$.

Algorithm:

1. If $a = 0$, return $|b|$, $\alpha = 0$, and $\beta = b/|b|$. If $b = 0$, return $|a|$, $\alpha = a/|a|$, and $\beta = 0$.

2. Compute, for $r_1 := a$ and $r_2 := b$, the sequence (r_n) using Algorithm 5.18, i.e. $r_{n-1} = q_{n-1}r_n + r_{n+1}$ with $q_{n-1} \in \mathbb{Z}$ and $r_k = 0$.
3. Define

$$A_n := \begin{bmatrix} 0 & 1 \\ 1 & -q_{n-1} \end{bmatrix}$$

for $1 \leq n \leq k-2$. Then the first row of the matrix $A_{k-2} \cdots A_1$ returns the pair α, β

10.0.13 Prime Field

Given $p \in \mathbb{N}, p > 1$ is prime, then $\mathbb{Z}/p\mathbb{Z}$ is a field with p elements. It can also be noted $\mathbb{F}_p$

Proof:

We need to show that for $a + p\mathbb{Z} \neq p\mathbb{Z}$ there exists some $b + p\mathbb{Z}$ such that $(a + p\mathbb{Z})(b + p\mathbb{Z}) = 1 + p\mathbb{Z}$.

For $a \notin p\mathbb{Z}$ we know that $\gcd(a, p) = 1$ thus, we do not have zero divisors.

10.0.14 Implication of Prime Numbers

For $a, b \in \mathbb{Z}$ and a prime number p

$$p \mid ab \implies (p \mid a) \vee (p \mid b)$$

Proof:

“ $\Rightarrow$ ” Let p be a prime and $a, b \in \mathbb{Z}$ with $p \mid ab$. Assume $p \nmid a$. Then there exist $\alpha, \beta \in \mathbb{Z}$ with $\alpha p + \beta a = 1$. Since $p \mid ab$, it follows that $p \mid \alpha p b + \beta a b = (\alpha p + \beta a)b = b$.

“ $\Leftarrow$ ” Let $c \in \mathbb{N}$ be a divisor of p . Then there exists $d \in \mathbb{N}$ with $p = cd$. In particular $1 \leq c \leq p$ and $1 \leq d \leq p$ and $p \mid cd$. By our assumption it follows that $p \mid c$ or $p \mid d$, i.e. $p \leq c$ or $p \leq d$. Together with $c \leq p$ and $d \leq p$ this forces $c = 1$ or $c = p$. Hence, p has no nontrivial divisors and is prime.

□◻◻

10.0.15 Fundamental Theorem of Arithmetic

Every integer greater than 1 can be written in the form

$$p_1^{n_1} p_2^{n_2} \cdots p_k^{n_k}$$

Where $n_i \geq 0$ and the p_i are distinct primes. The factorization is unique, except possibly for the order of the factors.

Example.

$$4312 = 2 \cdot 2156 = 2 \cdot 2 \cdot 1078 = 2 \cdot 2 \cdot 2 \cdot 539 = 2 \cdot 2 \cdot 2 \cdot 7 \cdot 77 = 2 \cdot 2 \cdot 2 \cdot 7 \cdot 7 \cdot 11$$

That is,

$$4312 = 2^3 \cdot 7^2 \cdot 11$$

Approach to the proof:

- The Fundamental Theorem of Arithmetic says that every integer greater than 1 can be factored uniquely into a product of primes.
- Euclid’s lemma says that if a prime divides a product of two numbers, it must divide at least one of the numbers.

- The least common multiple $[a, b]$ of nonzero integers a and b is the smallest positive integer divisible by both a and b .

10.0.16 Lemmas

Lemma

If $m \mid pq$ and $\gcd(m, p) = 1$, then $m \mid q$.

Proof:

Write $1 = \gcd(m, p) = am + bp$ for some $a, b \in \mathbb{Z}$. Then

$$q = amq + bpq$$

Since $m \mid amq$ and $m \mid bpq$ (because $m \mid pq$), we conclude $m \mid q$.

□◻◻

Lemma

If p is prime and $p \mid a_1 a_2 \cdots a_n$, then $p \mid a_i$ for some i .

Proof (Case $n = 2$):

Suppose $p \mid a_1 a_2$, and $p \nmid a_1$. Then $\gcd(p, a_1) = 1$, and by the previous lemma, $p \mid a_2$.

For general $n > 2$: Assume the result is true for $n-1$. Suppose $p \mid a_1 a_2 \cdots a_n$. Group as $(a_1 a_2 \cdots a_{n-1}) a_n$.

By the $n = 2$ case, either $p \mid a_n$ or $p \mid a_1 a_2 \cdots a_{n-1}$, and by induction, $p \mid a_i$ for some i .

□◻◻

10.0.17 Proof of the Fundamental Theorem of Arithmetic

Existence:

Use induction on $n > 1$. Let $X = (X_1, \dots, X_n)^T : \Omega \rightarrow X$ be a random variable with statistical model $P = \{P_\theta : \theta \in \Theta\}$. Furthermore, let $x_1, \dots, x_n$ be a sample and let $\alpha \in [0, 1]$.

A mapping I that assigns to each sample $x = (x_1, \dots, x_n)^T \in X$ a subset $I(x) \subset \Theta$ is called a $(1 - \alpha)$ -confidence region for θ if

$$\mathbb{P}_\theta(\theta \in I(X_1, \dots, X_n)) \geq 1 - \alpha$$

for all $\theta \in \Theta$.

If $\Theta \subset \mathbb{R}$ and $I(x)$ is an interval for all $x \in X$, then I is called a $(1 - \alpha)$ -confidence interval.

Base case: $n = 2$ is prime.

Inductive step: If n is prime, done. Otherwise, $n = ab$, with $1 < a, b < n$. By induction, both a and b factor into primes, so n does too.

Uniqueness:

Suppose:

$$p_1^{m_1} \cdots p_j^{m_j} = q_1^{n_1} \cdots q_k^{n_k}$$

With all p_i and q_i distinct primes.

Since p_1 divides the LHS, it divides the RHS. So $p_1 \mid q_i^{n_i}$ for some i , hence $p_1 = q_i$. Re-order so $p_1 = q_1$. Then:

If $m_1 > n_1$, divide both sides by $q_1^{n_1}$:

$$p_1^{m_1-n_1} \cdots p_j^{m_j} = q_2^{n_2} \cdots q_k^{n_k}$$

But then p_1 divides LHS but not RHS, contradiction. So $m_1 = n_1$. Cancel and repeat.

Eventually, all p_i match with some q_i , and the exponents are equal. So the factorizations are the same up to order.

□◻◻

10.0.18 Least Common Multiple

The least common multiple of a and b , denoted $[a, b]$, is the smallest positive integer divisible by both.

Example:

$$[6, 4] = 12, \quad [33, 15] = 165$$

10.0.19 Multiplication Property of the Least Common Multiple

$$[a, b] \cdot \gcd(a, b) = ab$$

Proof:

Let:

$$a = p_1 \cdots p_l q_1 \cdots q_m, \quad b = q_1 \cdots q_m r_1 \cdots r_n$$

Then:

$$\begin{aligned} \gcd(a, b) &= q_1 \cdots q_m \\ [a, b] &= p_1 \cdots p_l q_1 \cdots q_m r_1 \cdots r_n \\ ab &= p_1 \cdots p_l q_1^2 \cdots q_m^2 r_1 \cdots r_n \end{aligned}$$

So:

$$[a, b] \cdot \gcd(a, b) = ab$$

Example:

$$\gcd(36, 90) = 18, \quad [36, 90] = 180, \quad 36 \cdot 90 = 3240 = 18 \cdot 180$$

10.0.20 Second Proof of the Fundamental Theorem of Arithmetic

Given $n \in \mathbb{N}$ has 1 as a prime factor. Suppose that every natural number smaller than n can be factorized into primes.

If n is a prime number then it is itself its prime factorization. Else. there are $b, c \in \mathbb{Z}$ with $n = bc$ $1 < b, c < n$. By our induction's supposition, b and c have also a prime factorization.

$$b = p_1 \cdot p_2 \cdots p_r \text{ and } c = q_1 \cdot q_2 \cdots q_s$$

with p_i, p_j $1 \leq i \leq r$ and $1 \leq j \leq s$ prime numbers. The product is the prime factorization of the n .

The uniqueness is given by *the implication of prime numbers*.

□◻◻

10.0.21 Goldbach's Conjecture

Every even integer greater than 4 can be written as the sum two primes.

10.0.22 Prime Factorization

To find the *prime factorization* of a number a the algorithm goes as follows

- Let n be the first prime number.
- Try dividing a by n . If $n \mid a$ note the number and set $a = a \div n$ to repeat this step. If n did not divide a set n to the next prime number and repeat this step.
- When 1 was reached using the previous step then the algorithm terminates and the noted numbers is our prime factorization of a .

10.0.23 GCD via Prime factorization

The gcd of two or more number can also be found with the following algorithm.

- Find the prime factorization of all target numbers.
- Note the number which appear in both prime factorization always taking the one with least exponent.
- Multiply the noted prime factors together.

Chapter 11

Real Numbers

A way of defining the *real numbers* is via the *Cauchy sequences*. Another way is via the axioms of a field with the order axioms and the *completeness axiom*.

- *Associativity*: $(a * b) * c = a * (b * c)$ for all $a, b, c \in S$
- *Commutativity*: $a * b = b * a$ for all $a, b \in S$
- *Identity Element*: There exists $e \in S$ such that $a * e = e * a = a$ for all $a \in S$
- *Inverse Element*: For every $a \in S$, there exists $a^{-1} \in S$ such that $a * a^{-1} = a^{-1} * a = e$
- *Distributivity*: $a \circ (b \bullet c) = (a \circ b) \bullet (a \circ c)$ and/or $(b \bullet c) \circ a = (b \circ a) \bullet (c \circ a)$

These rules apply for the addition and multiplication for real numbers.

We also refresh the order axioms.

I. *Trichotomy*: Exactly one of the following holds:

$$a < b, \quad a = b, \quad a > b$$

II. *Transitivity*: If $a < b$ and $b < c$, then $a < c$.

III. *Additive Compatibility*: If $a < b$, then $a + c < b + c$.

IV. *Multiplicative Compatibility*: If $0 < a$ and $0 < b$, then $0 < ab$.

11.0.1 Properties derivated from the Axioms

Some proofs of these properties can be found in the section about the algebraic structures.

- The neutral and inverse elements of the addition and multiplication are well defined.
- $a + b = c$ and $a + c = 0 \implies b = c$
- $a \neq 0$, $ab = 1$ and $ac = 1 \implies b = c$
- $a \cdot 0 = 0$
- $-a = (-1)a$
- $-(-a) = a$
- $(-1)^2 = 1$
- $(a^{-1})^{-1} = a, \forall \in \mathbb{R}, a \neq 0$

11.0.2 Existence of Solutions

Given $a, b \in \mathbb{R}$, then

$$\exists x \in \mathbb{R}, \text{ such that } a + x = b$$

Also, if $a \neq 0$

$$\exists x \in \mathbb{R}, \text{ such that } ax = b$$

Proof:

For the first proposition, let $x = -a + b$

$$\begin{aligned} a + x &= a - a + b \\ &= b \end{aligned}$$

For the second, let $x = a^{-1}b$

$$\begin{aligned} ax &= aa^{-1}b \\ &= 1 \cdot b \end{aligned}$$

□

11.0.3 The Completeness Axiom

All non-empty upper-bounded subsets of the real numbers have a *supremum* $s \in \mathbb{R}$.

11.0.4 Minima in the Real Numbers

As a consequence of the last axiom each non-empty lower-bounded subset of real numbers has a minimum.

Proof:

The main idea is that the supremum times minus 1 of a set is the minimum of another set.

We define $-A := \{-a \mid a \in A\}$ then $-A \neq \emptyset$ and is upper-bounded. By definition s is less or greater than all the element of the set A .

$$\begin{aligned} s &\leq a, \forall a \in A \\ -s &\geq -a, \forall a \in A \end{aligned}$$

Thus, $-s = \sup -A$.

Now we want to prove that $-\sup -A$ is first, the lower bound of A and second, it is the greatest upper bound.

For the first part

$$\begin{aligned} \sup -A &\geq -a \forall a \in A \\ -\sup -A &\leq a \forall a \in A \end{aligned}$$

Thus, $-\sup -A$ is in fact a lower bound of A . For the second part let us assume that there exists some element s such that $s > -\sup -A$ and $s \leq a, \forall a \in A$, then

$$-s < \sup -A \text{ and } -s \geq -a, \forall a \in A$$

This is a contradiction because we have an upper bound, which is not an upper bound. Therefore, $-\sup -A$ is the infimum of A .

□◻◻

11.0.5 Approximation of the Supremum

Given a non-empty, upper-bounded set $A \subseteq \mathbb{R}$ and $s = \sup A$. Then there is for each $y < s$ an $x \in A$, such that

$$y < x \leq s$$

An analogous theorem can be written for the minimum as well.

Proof:

If $x \in A$ then the proof is completed. For the case $s \notin A$, if there existed some $y^* \in \mathbb{R}$ where $y^* < s$ and $y^* \geq x$ for all $x \in A$ then we still have a contradiction to the assumption that s is the supremum of A .

□◻◻

11.0.6 Supremum of the Naturals

$\mathbb{N}$ is not upper-bounded.

Proof:

Assume $a = \sup \mathbb{N}$. Since $\mathbb{N} \subset \mathbb{R}$ is not empty it has an upper-bound by the completeness axiom.

By the definition of the supremum $a - 1 \in \mathbb{N}$ is not an upper-bound of $\mathbb{N}$, therefore $a - 1 < k, k \in \mathbb{N}$.

Then, the following also applies

$$a < k + 1$$

This is a contradiction to the assumption that a our supremum, thus our assumption has been proven false.

11.0.7 Remainder of Archimedes and Eudoxus

Archimedes

For all pair of positive real numbers x, y exists some $n \in \mathbb{N}$ such that:

$$nx > y$$

Eudoxus

For all positive real numbers there exists an $n \in \mathbb{N}$, such that

$$\frac{1}{n} < x$$

Proof Eudoxus:

$x \in \mathbb{R}_{>0} \implies \frac{1}{x} \in \mathbb{R}_{>0}$. By the Archimedean property, $\exists n \in \mathbb{Z}_{>0}, \frac{1}{x} < n$. Equivalently $\frac{1}{n} < x$.

□◻◻

11.0.8 Density of the Rationals in the Reals

For each pair $a, b \in \mathbb{R}$ there exists some $q \in \mathbb{Q}$ such that

$$a < q < b$$

Proof:

Let $q = \frac{m}{n}, m \in \mathbb{Z}, n \in \mathbb{N}$ to prove $a < \frac{m}{n} < b$ equivalently $an < m < bn$.

For $b - a > 0$ we know $\frac{1}{n} < b - a$ or $a < b - \frac{1}{n}$, by the Archimedean property.

Next take the smallest $m \in \mathbb{Z}$ such that $m - 1 \leq na < m$ or $a < \frac{m}{n}$. This gives us the first half of our theorem.

Finally, for the second half

$$\begin{aligned} m - 1 &\leq na \\ &= na + 1 \\ &< n(b - \frac{1}{n}) + 1 \text{ using our result from the Archimedean property} \\ &= nb \end{aligned}$$

This tells us that $m < nb \implies \frac{m}{n} < b$.

□◻◻

11.0.9 Existence of Square Roots

For each $x \in [0, \infty]$, $\exists r \in [0, \infty)$ s. d. $r^2 = x$. Also, $\sqrt{x} := r$ and we call $x \mapsto \sqrt{x}$ the *root function*.

Proof:

Let $T = \{x \in [0, \infty) \mid x < y\}$ and let us define $a = y^2$ as the supremum of T .

To prove that the equality holds we will prove that the other order cases can not be true.

Assume $a < y$, because $a - y < 0$, we know that $\frac{1}{n} < a - y$ and $a + \frac{1}{n} < y$. Thus, $a + \frac{1}{n} \in T$, but because a is the supremum this contradiction because our supremum is not the smallest upper-bound.

For the second case, assume $a > y$, by applying the same reasoning with $a - \frac{1}{n}$, we will arrive at a similar contradiction

Assume $a > y$, then $a - y > 0 \implies \frac{1}{n} < a - y, \implies a - \frac{1}{n} < y$

But then $a - \frac{1}{n}$ is an upper bound and this contradicts our assumption. Thus, by elimination $a^2 = y$

□◻◻

11.0.10 Irrational Numbers

$$\mathbb{R} \setminus \mathbb{Q} \text{ or } \mathbb{Q}^C := \{x \in \mathbb{R} \mid x \notin \mathbb{Q}\}$$

11.0.11 Absolute Value for Real Numbers

$$|x| := \sqrt{x * x}$$

11.0.12 Norms Remainder

Properties of a norm

- $|x| \geq 0, \forall x \in \mathbb{R}$
- $|x| = 0, \implies x = 0$
- $|\alpha x| = |\alpha||x|, \forall x, \alpha \in \mathbb{R}$
- $|x + y| \leq |x| + |y|, \forall x, y \in \mathbb{R}$

11.0.13 Distance

We define the *distance* between two real numbers as

$$d(x, y) = |x - y|$$

With the following properties

- $d(x, y) \geq 0, \forall x, y \in \mathbb{R}$
- $d(x, y) = 0, \iff x = y$
- $d(x, y) = d(y, x)$
- $d(x, y) \geq d(x, z) + d(z, y), \forall x, y, z \in \mathbb{R}$

11.0.14 Nested Interval Property

For each $n \in \mathbb{N}$, assume we are given a closed interval $I_n = [a_n, b_n] = \{x \in \mathbb{R} \mid a_n \leq x \leq b_n\}$. Also assume each I_n contains I_{n+1} . Then the nested sequence of closed intervals $I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$ has a non-empty intersection.

Proof:

By definition every I_n is non-empty, thus the sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are well-defined. Also, every set $A = \{a_n : n \in \mathbb{N}\}$ is bounded above by the corresponding b_n , and this is guaranteed by the axiom of completeness.

Let $x = \sup A$. Then consider $I_n = [a_n, b_n]$

$$a_n \leq x \leq b_n, \forall n \in \mathbb{N} \implies x \in I_n$$

Thus, $x \in \bigcap_{n=1}^{\infty} I_n$ and the intersection is non-empty.

□◻◻

11.0.15 Definition as Cauchy Sequences

Let K be an ordered field. On the set

$$\text{ch}(K) := \{x : \mathbb{N} \rightarrow K \mid x \text{ is a Cauchy sequence}\}$$

and on the set

$$c(K) := \{x : \mathbb{N} \rightarrow K \mid x \text{ is a convergent sequence}\}$$

we can define an addition and a multiplication as follows:

If $x = (x_n)_{n \in \mathbb{N}}$ and $y = (y_n)_{n \in \mathbb{N}}$ are Cauchy sequences (respectively, convergent sequences), then their sum is defined as

$$x + y := (x_n)_{n \in \mathbb{N}} + (y_n)_{n \in \mathbb{N}} := (x_n + y_n)_{n \in \mathbb{N}}$$

and their product is defined as

$$x \cdot y := (x_n)_{n \in \mathbb{N}} \cdot (y_n)_{n \in \mathbb{N}} := (x_n \cdot y_n)_{n \in \mathbb{N}}$$

The sum and product satisfy all field axioms except for the existence of the multiplicative inverse. The zero element is $0_{\mathbb{N}} = (0, 0, \dots)$, the unit element is $1_{\mathbb{N}} = (1, 1, \dots)$, and the additive inverse of $x = (x_n)_{n \in \mathbb{N}}$ is $-x = (-x_n)_{n \in \mathbb{N}}$. We demonstrate the distributive law as an example:

Let $x = (x_n)_{n \in \mathbb{N}}, y = (y_n)_{n \in \mathbb{N}}, z = (z_n)_{n \in \mathbb{N}}$ be Cauchy sequences (convergent sequences). Then we have:

$$\begin{aligned} x(y + z) &= (x_n)_{n \in \mathbb{N}} \cdot ((y_n)_{n \in \mathbb{N}} + (z_n)_{n \in \mathbb{N}}) \\ &= (x_n)_{n \in \mathbb{N}} \cdot (y_n + z_n)_{n \in \mathbb{N}} \\ &= (x_n(y_n + z_n))_{n \in \mathbb{N}} \\ &= (x_n y_n + x_n z_n)_{n \in \mathbb{N}} \\ &= (x_n y_n)_{n \in \mathbb{N}} + (x_n z_n)_{n \in \mathbb{N}} \\ &= xy + xz. \end{aligned}$$

We now aim to construct the ordered field $\mathbb{R}$ of the real numbers; it will have the following properties:

- There exists an injective mapping $j : \mathbb{Q} \rightarrow \mathbb{R}$ which respects addition, multiplication, and order, such that the following holds:

For all $z, w \in \mathbb{R}$ with $z < w$, there exists an $x \in \mathbb{Q}$ such that

$$z < j(x) < w$$

- Every Cauchy sequence in $\mathbb{R}$ converges.

Via j , we identify $\mathbb{Q}$ with $j(\mathbb{Q})$ and consider $\mathbb{Q}$ as a subset of $\mathbb{R}$. In $\mathbb{R}$, the following will additionally hold:

- For all $y > 0$ and $n \in \mathbb{N}$, the equation $x^n = y$ has a solution.
- Every bounded above subset of $\mathbb{R}$ has a supremum.

We define the following relation on the set $\text{ch}(\mathbb{Q})$ of all Cauchy sequences in $\mathbb{Q}$:

$$x \sim y \quad \text{if and only if} \quad x - y \text{ is a null sequence}$$

That is, $(x_n)_{n \in \mathbb{N}} \sim (y_n)_{n \in \mathbb{N}}$ if and only if

$$x_n - y_n \rightarrow 0 \quad (n \rightarrow \infty)$$

11.0.16 Definition

The set

$$\mathbb{R} := \{[x]_{\sim} : x \in \text{ch}(\mathbb{Q})\}$$

is called the set of real numbers.

Analogous to the construction of the rational numbers, the real numbers consist of equivalence classes. Roughly speaking, an equivalence class consists of those Cauchy sequences in $\mathbb{Q}$ that exhibit the same limit behavior.

Equipped with the addition

$$+ : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, \quad [x], [y] \mapsto [x] + [y] := [x + y]$$

and the multiplication

$$\cdot : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, \quad [x], [y] \mapsto [x] \cdot [y] := [xy]$$

$\mathbb{R}$ is a field. The zero element is $[0_{\mathbb{N}}]$, and the unit element is $[1_{\mathbb{N}}]$.

Chapter 12

Complex Numbers


A complex number is a number of the form:

$$z = a + bi,$$

where $a, b \in \mathbb{R}$, and i is the imaginary unit defined by $i^2 = -1$. The set of all complex numbers is denoted by $\mathbb{C}$.

12.0.1 The Complex Plane

Complex numbers can be represented graphically in the *complex plane*, where the horizontal axis represents the real part and the vertical axis the imaginary part.



The point $1 + i$ is located at (1,1), showing 1 unit on the real axis and 1 unit on the imaginary axis.

12.0.2 Conjugate of a Complex Number

The *conjugate* of a complex number $z = a + bi$ is denoted $\bar{z}$ and is defined as:

$$\bar{z} = a - bi$$

Geometrically, it reflects the point z across the real axis in the complex plane. Conjugates are useful in division and in finding the modulus, since:

$$z \cdot \bar{z} = a^2 + b^2 = |z|^2$$

12.0.3 Operations in Cartesian Coordinates

Let $z_1 = a + bi$ and $z_2 = c + di$ be two complex numbers.

Addition:

$$z_1 + z_2 = (a + c) + (b + d)i$$

Multiplication:

$$z_1 \cdot z_2 = (ac - bd) + (ad + bc)i$$

Quotient:

$$\frac{z_1}{z_2} = \frac{(a + bi)(c - di)}{(c + di)(c - di)} = \frac{(a + bi)(c - di)}{c^2 + d^2} = \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2}$$

12.0.4 Polar Coordinates

A complex number can also be expressed in polar form as:

$$z = r(\cos \theta + i \sin \theta) = re^{i\theta},$$

where:

$$r = |z| = \sqrt{a^2 + b^2} \quad (\text{modulus})$$
$$\theta = \arg(z) = \tan^{-1} \left(\frac{b}{a} \right) \quad (\text{argument})$$

The value of θ depends on the quadrant where the complex number lies:

- Quadrant I: $a > 0, b > 0$ - use $\tan^{-1}(b/a)$
- Quadrant II: $a < 0, b > 0$ — add π to $\tan^{-1}(b/a)$
- Quadrant III: $a < 0, b < 0$ — add π to $\tan^{-1}(b/a)$
- Quadrant IV: $a > 0, b < 0$ — use $\tan^{-1}(b/a)$

12.0.5 Multiplication and Division in Polar Coordinates

Given:

$$z_1 = r_1 e^{i\theta_1}, \quad z_2 = r_2 e^{i\theta_2},$$

Multiplication:

$$z_1 \cdot z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

Division:

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}$$

This polar form is especially useful in simplifying powers and roots of complex numbers using De Moivre's Theorem.

12.0.6 Exponentiation and Roots (De Moivre's Theorem)

Let $z = r(\cos \theta + i \sin \theta) = re^{i\theta}$ be a complex number in polar form.

Exponentiation

To raise z to the power $n \in \mathbb{N}$, we use De Moivre's Theorem:

$$z^n = r^n(\cos(n\theta) + i \sin(n\theta)) = r^n e^{in\theta}$$

Roots of Complex Numbers

To find the n th roots of a complex number $z = re^{i\theta}$, we use the formula:

$$z^{1/n} = r^{1/n} \left(\cos \left(\frac{\theta + 2k\pi}{n} \right) + i \sin \left(\frac{\theta + 2k\pi}{n} \right) \right), \quad k = 0, 1, \dots, n-1$$

This yields n distinct roots, each separated by an angle of $\frac{2\pi}{n}$ in the complex plane.

12.0.7 Example: Solve $z^4 = 1 + \sqrt{3}i$

Step 1: Convert RHS to polar form.

Let $w = 1 + \sqrt{3}i$. Real part: $a = 1$, Imaginary part: $b = \sqrt{3}$

$$\begin{aligned} r = |w| &= \sqrt{1^2 + (\sqrt{3})^2} = \sqrt{1+3} = 2 \\ \theta = \arg(w) &= \tan^{-1} \left(\frac{\sqrt{3}}{1} \right) = \frac{\pi}{3} \end{aligned}$$

So,

$$w = 2 \left(\cos \left(\frac{\pi}{3} \right) + i \sin \left(\frac{\pi}{3} \right) \right)$$

Step 2: Solve $z^4 = w \rightarrow z = w^{1/4}$

Using the root formula:

$$z_k = 2^{1/4} \left(\cos \left(\frac{\pi + 2k\pi}{4} \right) + i \sin \left(\frac{\pi + 2k\pi}{4} \right) \right), \quad k = 0, 1, 2, 3$$

So the four roots are:

$$\begin{aligned} z_0 &= 2^{1/4} \left(\cos \left(\frac{\pi}{4} \right) + i \sin \left(\frac{\pi}{4} \right) \right) \\ z_1 &= 2^{1/4} \left(\cos \left(\frac{5\pi}{4} \right) + i \sin \left(\frac{5\pi}{4} \right) \right) \\ z_2 &= 2^{1/4} \left(\cos \left(\frac{9\pi}{4} \right) + i \sin \left(\frac{9\pi}{4} \right) \right) = 2^{1/4} \left(\cos \left(\frac{3\pi}{4} \right) + i \sin \left(\frac{3\pi}{4} \right) \right) \\ z_3 &= 2^{1/4} \left(\cos \left(\frac{13\pi}{4} \right) + i \sin \left(\frac{13\pi}{4} \right) \right) \end{aligned}$$

These represent the four complex 4th roots of $1 + \sqrt{3}i$, equally spaced around the circle of radius $2^{1/4}$ in the complex plane.

12.0.8 Solving Equations with Complex Numbers

Solving equations in $\mathbb{C}$ can involve various forms. Here are the most common cases:

Linear Equations:

Solve for z in $az + b = 0$, where $a, b \in \mathbb{C}$, $a \neq 0$:

$$z = -\frac{b}{a}$$

Equations Involving the Conjugate:

Solve for z in equations like $z + \bar{z} = 4$. Let $z = x + iy$, then $\bar{z} = x - iy$. So:

$$z + \bar{z} = 2x \quad \rightarrow \quad x = 2 \quad \Rightarrow \quad z = 2 + iy$$

The imaginary part remains free unless further constraints are given.

Modulus Equations:

Solve $|z| = r$. Let $z = x + iy$, then:

$$\sqrt{x^2 + y^2} = r \rightarrow x^2 + y^2 = r^2$$

This is a circle of radius r centered at the origin in the complex plane.

Equations Involving $z \cdot \bar{z}$

Recall $z \cdot \bar{z} = |z|^2$. For example, solve:

$$z \cdot \bar{z} = 9 \rightarrow |z| = 3$$

Again, a circle in the complex plane of radius 3.

Quadratic Equations:

Complex roots occur naturally. For example:

$$z^2 + 1 = 0 \rightarrow z^2 = -1 \Rightarrow z = \pm i$$

General Polynomial Equations:

Use De Moivre's Theorem or polar form. Example:

$$z^n = w \rightarrow z_k = \sqrt[n]{|w|} \cdot e^{i\left(\frac{\arg(w) + 2k\pi}{n}\right)}, \quad k = 0, 1, \dots, n-1$$

Example 1:

Solve:

$$\begin{aligned} \left(\frac{2+3i}{1+i} + \frac{4+5i}{2-2i} \right) \hat{z} &= \frac{i+2}{i} \\ \left(\frac{-3-i}{4} \right) \hat{z} &= \frac{i+2}{i} \\ \hat{z} &= \frac{i+2}{i} : \frac{-3-i}{4} \end{aligned}$$

Example 2:

Solve:

$$z - 3i + (2 - i)\hat{z} + 2 = 0$$

In this case we let $z = x + iy$ and $\hat{z} = x - iy$.

$$\begin{aligned} z &= 3i - (2 - i)\hat{z} - 2 \\ x + iy &= 3i - (2 - i)(x - iy) - 2 \\ x + iy &= 3i - [2x - 2yi - xi + yi^2] - 2 \\ x + yi &= 3i - 2 + 2x + 2yi + xi + y \\ x + yi &= (y - 2 - 2x) + i(3 + 2y + x) \\ x &= y - 2 - 2x \quad y = 3 + 2y + x \\ x &= \frac{y - 2}{3} \quad y = 3 + 2y + \frac{y - 2}{3} = -7 \\ x &= \frac{-7 - 2}{3} = -3 \\ z &= -3 - 7i \quad \hat{z} = -3 + 7i \end{aligned}$$

12.0.9 The Complex Logarithm

The logarithm of a complex number is multivalued due to the periodic nature of the complex exponential.

Let $z = re^{i\theta}$ with $r > 0$, $\theta \in \mathbb{R}$. Then:

$$\log z = \ln r + i(\theta + 2\pi k), \quad k \in \mathbb{Z}$$

Here:

- $\ln r$ is the natural (real) logarithm of the modulus.
- θ is the principal argument $\arg(z) \in (-\pi, \pi]$.
- The term $2\pi k$ accounts for the infinitely many branches of the logarithm in $\mathbb{C}$.

Principal Value:

The principal value of the complex logarithm is often written:

$$\text{Log } z = \ln |z| + i \text{Arg}(z), \quad \text{where } \text{Arg}(z) \in (-\pi, \pi]$$

Example:Let $z = -1$. Then:

$$|z| = 1, \quad \arg(z) = \pi, \quad \rightarrow \log(-1) = i(\pi + 2\pi k), \quad k \in \mathbb{Z}$$

$$\rightarrow \text{Log}(-1) = i\pi$$

The multivalued nature of $\log z$ is crucial in advanced complex analysis, especially in defining analytic continuations and branch cuts.

12.0.10 Complex Exponents

$a^x \approx 1(a + \alpha \frac{x}{N})^N \rightarrow e^z := \lim_{N \rightarrow \infty} (1 + \frac{z}{N})^N$ The process above is called linearization of the exponential function by zooming $\alpha \approx \frac{dy}{dx}$

For $e^{ci} = \cos \theta + i \sin \theta$ every exponentiation of a complex number is a rotation in the complex plane.

$$e^{ic} = \lim_{N \rightarrow \infty} \left(1 + \frac{ic}{N}\right)^N$$

Now imagine that in a sector of a circumference you put triangles one above the other with base of length one and a height of $\frac{C}{N}$ and an angle of δ

$$\tan \delta \approx \delta \text{ for } \delta \ll 1$$

$$1 + \frac{ci}{N} = 1 \angle \frac{c}{N}, N \gg 1$$

$$e^{ic} = \lim_{N \rightarrow \infty} \left(1 + \frac{c}{N}\right)^N \rightarrow e^{ci} = 1 \angle c = \cos c + i \sin c$$

12.0.11 Euler's Formula Proof

We know that

$$e^{i\pi} = -1 \text{ and } e^{i\theta} = |r|(\cos \theta + i \sin \theta)$$

$$e^z = \lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n \implies e^{i\pi} = \lim_{n \rightarrow \infty} \left(1 + \frac{i\pi}{n}\right)^n = -1$$

$$\implies \lim_{n \rightarrow \infty} |r_n| = 1 \quad \lim_{n \rightarrow \infty} \theta = 0 +$$

Now we can demonstrate the formula.

$$|r_n| = \left(1 + \left|\frac{z}{n}\right|\right)^n \implies \left(\sqrt{1 + \frac{\pi^2}{n^2}}\right)^n$$

$$\theta = \sum_{k=1}^n n \arctan \frac{\pi}{2} = n \arctan \frac{\pi}{n}$$

$$\lim_{n \rightarrow \infty} \left(\sqrt{1 + \frac{\pi^2}{n^2}}\right)^n = \lim_{n \rightarrow \infty} \left(1 + \frac{\pi^2}{2n}\right)^{\frac{n}{2}} = \lim_{n \rightarrow \infty} e^{\ln\left(1 + \frac{\pi^2}{2}\right) \frac{n}{2}} = e^0 = 1$$

$$\lim_{n \rightarrow \infty} \theta = \lim_{n \rightarrow \infty} n \arctan \frac{\pi}{n} = \lim_{n \rightarrow \infty} n^{-1} \arctan \frac{\pi}{n} = 0$$

Thus, for $e^{i\pi} = 1$ for $x = \pi \forall x \in \lim r_n(x) = 1$ and $\lim \theta(x) = x$

ΩℰΩ

12.0.12 The Fundamental Theorem of Algebra

Given any positive integer $n \geq 1$ and any choice of complex numbers $a_0, a_1, \dots, a_n$, such that $a_n \neq 0$, the polynomial equation

$$a_n z^n + \dots + a_1 z + a_0 = 0$$

has at least one solution $z \in \mathbb{C}$.

12.0.13 Complex Numbers as Matrices

Given a complex number $z = a + bi$, we can represent it as a real 2x2 matrix:

$$z \leftrightarrow \begin{bmatrix} a & -b \\ b & a \end{bmatrix},$$

with the linear function $f : M_2(\mathbb{R}) \rightarrow \mathbb{C}$ defined by:

$$f\left(\begin{bmatrix} a & -b \\ b & a \end{bmatrix}\right) = a + bi$$

By taking the determinant of this matrix gives us the modulus squared of the complex number:

$$\det \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = a^2 + b^2 = |z|^2$$

Note, that this matrix representation preserves addition and multiplication:

$$\begin{aligned} \begin{bmatrix} a_1 & -b_1 \\ b_1 & a_1 \end{bmatrix} + \begin{bmatrix} a_2 & -b_2 \\ b_2 & a_2 \end{bmatrix} &= \begin{bmatrix} a_1 + a_2 & -(b_1 + b_2) \\ b_1 + b_2 & a_1 + a_2 \end{bmatrix} \\ \begin{bmatrix} a_1 & -b_1 \\ b_1 & a_1 \end{bmatrix} \cdot \begin{bmatrix} a_2 & -b_2 \\ b_2 & a_2 \end{bmatrix} &= \begin{bmatrix} a_1 a_2 - b_1 b_2 & -(a_1 b_2 + b_1 a_2) \\ a_1 b_2 + b_1 a_2 & a_1 a_2 - b_1 b_2 \end{bmatrix} \end{aligned}$$

Hence,

$$M_2(\mathbb{R}) \supseteq \left\{ \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \right\} \cong \mathbb{C}$$

Chapter 13

Topology

In this section, we introduce essential vocabulary used in topology. Each term is accompanied by a brief explanation and its formal mathematical definition.

13.0.1 Introduction to topological nomenclature

Open Ball $B_\varepsilon(x)$

For a metric space (X, d) , the *open ball* centered at $x \in X$ with radius $\varepsilon > 0$ is defined as:

$$B_\varepsilon(x) := \{y \in X \mid d(x, y) < \varepsilon\} = (x - \varepsilon, x + \varepsilon)$$

It represents the set of all points within distance ε from x , excluding the boundary.

Open Set

A subset $U \subseteq X$ of a topological space is called *open* if for every point $x \in U$, there exists an $\varepsilon > 0$ such that the open ball $B_\varepsilon(x) \subseteq U$. Intuitively, an open set contains none of its boundary points and every point has some wiggle room around it.

Example:

We will prove that $(0, 1)$ is an open set.

This means $x_0 > 0$ and $1 - x_0 > 0$. First we define

$$\varepsilon := \frac{1}{2} \min x_0, 1 - x_0$$

Then from $|y - x_0| \leq \varepsilon$ we get

$$\begin{aligned} y \geq x_0 |y - x_0| &\geq x_0 - \varepsilon \geq \frac{1}{2}x_0 > 0, \\ y \leq x_0 |y - x_0| &\leq x_0 + \varepsilon \leq x_0 + \frac{1}{2}(1 - x_0) = \frac{1}{2} + \frac{1}{2}x_0 < 1 \end{aligned}$$

□◻◻

The Interior of a Set

The *interior* denoted as A° of a set A is the set of all inner points of A .

Closed Set

A subset $A \subseteq X$ is called *closed* if its complement $X \setminus A$ is open. Equivalently, A contains all its limit points. That is, A is closed if it includes its boundary.

Example: $[0, 1]$

Boundary Point

A point $x \in X$ is a *boundary point* of a set $A \subseteq X$ if every open ball around x contains both points in A and in $X \setminus A$. The set of all boundary points is called the *boundary* of A , denoted ∂A .

Accumulation Point / Limit Point

A point $x \in X$ is an *accumulation point* of a set $A \subseteq X$ if every open ball $B_\varepsilon(x)$ contains a point of $A \setminus \{x\}$. In other words, points of A cluster arbitrarily close to x , even if $x \notin A$.

Isolated Point

A point $x \in A$ is an *isolated point* if there exists $\varepsilon > 0$ such that $B_\varepsilon(x) \cap A = \{x\}$. That is, x stands alone in A without other points of A nearby.

Example:

-1 is an isolated point of $-1 \cup (0, 1)$

Boundary

The set $A \cup \{ \text{all boundary points} \}$ is called the *boundary* of A . It is denoted as $\overline{A}$.

Compact Set

A set $K \subseteq X$ is *compact* if every open cover of K has a finite sub-cover. In $\mathbb{R}^n$, this is equivalent to K being closed and bounded (by the Heine–Borel theorem).

Dense Set

A subset $D \subseteq X$ is *dense* in X if every point $x \in X$ is either in D or is a limit point of D . Equivalently, the closure of D is X , i.e., $\overline{D} = X$.

Chapter 14

Inequalities

14.0.1 Triangle Inequality

- $|x + y| \leq |x| + |y|$
- $|x - y| \geq |x| - |y|$
- $||x| - |y|| \leq |x - y|$
- $xy \leq |x| \cdot |y|$

14.0.2 Reciprocal Inequality

$$a < b < c \implies \frac{1}{a} > \frac{1}{b} > \frac{1}{c}$$

14.0.3 Bernoulli Inequality

Given $x \in \mathbb{R}, x > -1, x \neq 0$, then for $n \in \mathbb{N}, n \geq 2$, we have

$$(1 + x)^n > 1 + nx$$

Proof:

For $n = 2$

$$(1 + x)^2 > 1 + 2x.$$

Then, for $n + 1$ with $n > 2$:

$$\begin{aligned}(1 + x)^{n+1} &> 1 + (n + 1)x \\ (1 + x)^n &> 1 + (n + 1)x\end{aligned}$$

□

14.0.4 Inequalities for Real Numbers

For $x, y \in \mathbb{R}$ the following holds:

$$x < y + \varepsilon, \forall \varepsilon > 0 \iff x \leq y$$

$$|x| < \varepsilon, \forall \varepsilon > 0 \iff x = 0$$

Proof:

For the first, if $x > y$, then I can choose some epsilon only at some point where $y + \varepsilon > x$; therefore, not all epsilons satisfy the inequality.

For the second one, if $x \neq 0$, then $|x| \neq 0 \wedge |x| > 0$; therefore, the only valid option is for x to be zero.

□◻◻

14.0.5 Elementary Young Inequalities

For all $x, y \in \mathbb{R}$, the following holds:

$$xy \leq \frac{1}{2}x^2 + \frac{1}{2}y^2.$$

And this implies:

$$\sqrt{xy} \leq \frac{1}{2}x + \frac{1}{2}y, \quad \forall x, y \geq 0$$

and

$$xy \leq \frac{\varepsilon}{2}x^2 + \frac{1}{2\varepsilon}y^2, \quad \forall x, y \in \mathbb{R}, \varepsilon > 0.$$

14.0.6 Inequalities with and without Absolute Value

Inequalities are like normal *equalities*, but with ranges of solutions instead of one particular solution. The inequalities without absolute value are straightforward to solve, just like equations, with the particularity that when dividing both sides by a negative number, the relation $<, >, \geq, \leq$ has to be converted to its opposite.

The case with absolute value is similar, but here we have to account for different cases depending on the expression inside the $|\circ|$.

Example:

Solve $|x + 1| \leq 10 + |2x - 6|$.

Main cases:

$$F_1 : x + 1 \geq 0 \implies x \geq -1$$

$$F_2 : x + 1 < 0 \implies x < -1$$

Subcases:

$$F_{1,1} : 2x - 6 \geq 0 \implies x \geq 3$$

$$\begin{aligned} x + 1 &\geq 10 + 2x - 6 \\ x &\leq 5 \\ \mathcal{L} &= [3; \infty) \end{aligned}$$

$$F_{1,2} : 2x - 6 < 0 \implies x < 3$$

$$\begin{aligned}
x + 1 &\geq 10 - (2x - 6) \\
x &\geq 5 \\
\mathcal{L} &= (-\infty; 3]
\end{aligned}$$

$$F_{2,1} : 2x - 6 \geq 0 \implies x \geq 3$$

$$\begin{aligned}
-(x + 1) &\geq 10 + 2x - 6 \\
x &\leq \frac{-5}{3} \\
\mathcal{L} &= \emptyset
\end{aligned}$$

$$F_{2,2} : 2x - 6 < 0 \implies x < 3$$

$$\begin{aligned}
-(x + 1) &\geq 10 - (2x - 6) \\
x &\leq 17 \\
\mathcal{L} &= (-\infty, 3)
\end{aligned}$$

Union of the solutions:

$$\mathcal{L} = (-\infty; 3] \cup [3; \infty) \cup \emptyset \cup (-\infty, 3) = (-\infty, \infty)$$

Chapter 15

Means

15.0.1 Arithmetic Mean

The *arithmetic mean* of n numbers $x_1, x_2, \dots, x_n$ is given by:

$$A = \frac{x_1 + x_2 + \cdots + x_n}{n}$$

15.0.2 Geometric Mean

The *geometric mean* of n numbers $x_1, x_2, \dots, x_n$ is given by:

$$G = \sqrt[n]{x_1 \cdot x_2 \cdots x_n}$$

15.0.3 Harmonic Mean

The *harmonic mean* of n numbers $x_1, x_2, \dots, x_n$ is given by:

$$H = \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \cdots + \frac{1}{x_n}}$$

15.0.4 Quadratic Mean

The *quadratic mean* of n numbers $x_1, x_2, \dots, x_n$ is given by:

$$Q = \sqrt{\frac{x_1^2 + x_2^2 + \cdots + x_n^2}{n}}$$

15.0.5 Proof of the Arithmetic Mean-Geometric Mean Inequality

Let $a_1, a_2, \dots, a_n > 0$. We will prove by induction that:

$$\frac{a_1 + a_2 + \cdots + a_n}{n} \geq \sqrt[n]{a_1 a_2 \cdots a_n},$$

with equality if and only if $a_1 = a_2 = \cdots = a_n$.

Base Case: $n = 2$

We want to prove:

$$\frac{a_1 + a_2}{2} \geq \sqrt{a_1 a_2}$$

Let $a_1, a_2 > 0$. Then, by the identity

$$\left(\frac{a_1 - a_2}{2}\right)^2 \geq 0,$$

we get

$$\frac{a_1^2 - 2a_1a_2 + a_2^2}{4} \geq 0 \Rightarrow a_1^2 + a_2^2 \geq 2a_1a_2.$$

So,

$$(a_1 + a_2)^2 \geq 4a_1a_2 \Rightarrow \left(\frac{a_1 + a_2}{2}\right)^2 \geq a_1a_2,$$

and taking square roots gives the desired result:

$$\frac{a_1 + a_2}{2} \geq \sqrt{a_1a_2}.$$

For $n \geq 2$:

$$\begin{aligned} A_{n+1} &:= \left(\sum_{i=1}^{n+1} a_i\right) / (n+1) = \frac{a_1 + a_2 + \cdots + a_n + a_{n+1}}{n+1} \\ G_{n+1} &:= \sqrt[n+1]{a_1a_2 \cdots a_na_{n+1}} = \sqrt[n+1]{(a_1a_2 \cdots a_n)a_{n+1}} \\ A_{n+1}^{n+1} &:= \left(\frac{a_1 + a_2 + \cdots + a_{n+1}}{n+1}\right)^{n+1} = A_{n+1}^{n+1} \\ G_{n+1}^{n+1} &:= \left(\sqrt[n+1]{a_1a_2 \cdots a_na_{n+1}}\right)^{n+1} = (a_1a_2 \cdots a_n)a_{n+1} = \sqrt[n]{(a_1a_2 \cdots a_n)^n} a_{n+1}^{n+1} = G_n^n a_{n+1}^{n+1}. \end{aligned}$$

Then

$$G_{n+1}^{n+1} A_{n+1}^{n-1} = G_n^n a_{n+1}^{n+1} A_{n+1}^{n-1} \leq A_n^n A_{n+1}^{n-1}.$$

This comes from:

$$\begin{aligned} G_n &\leq A_n \\ G_n^n &\leq A_n^n \\ G_n^n a_{n+1} &\leq A_n^n a_{n+1} \\ G_n^n a_{n+1}^{n+1} A_{n+1}^{n-1} &\leq A_n^n A_{n+1}^{n-1} = \left(A_n^n (a_{n+1} A_{n+1})^{n-1}\right)^{\frac{n}{n}} \\ &\leq A_n^n \left(\frac{a_{n+1} + A_{n+1} + \cdots + A_{n+1}}{n}\right)^n \\ &\quad A_n^n \left(\frac{a_{n+1} + (n-1)A_{n+1}}{n}\right)^n \\ &\quad \left(A_n \frac{a_{n+1} + (n-1)A_{n+1}}{n}\right)^n. \end{aligned}$$

Note that:

$$\begin{aligned} \left(A_n \frac{a_{n+1} + (n-1)A_{n+1}}{n} \right) &\rightarrow \left(\sqrt{A_n \frac{a_{n+1} + (n-1)A_{n+1}}{n}} \right)^2 \\ &\leq \left(\frac{A_n + \frac{a_{n+1} + (n-1)A_{n+1}}{n}}{2} \right)^{2n}. \end{aligned}$$

Now, with power of n , we have

$$\begin{aligned} \left(A_n + \frac{a_{n+1} + (n-1)A_{n+1}}{n} \right)^n &\leq \left(\frac{A_n + \frac{a_{n+1} + (n-1)A_{n+1}}{n}}{2} \right)^{2n} \\ &= \left(\frac{A_n}{2} + \frac{a_{n+1} + (n-1)A_{n+1}}{2n} \right)^{2n} \\ &= \left(\frac{A_n n}{2n} + \frac{a_{n+1} + (n-1)A_{n+1}}{2n} \right)^{2n} \\ &= \left(\frac{A_n n + a_{n+1} + (n-1)A_{n+1}}{2n} \right)^{2n} \\ &= \left(\frac{A_{n+1}(n+1) + (n-1)A_{n+1}}{2n} \right)^{2n} \\ &= \left(\frac{2nA_{n+1}}{2n} \right)^{2n} = (A_{n+1})^{2n}. \end{aligned}$$

Now, we have proven that $G_{n+1}^{n+1} A_{n+1}^{n-1} \leq (A_{n+1})^{2n}$. By dividing both sides by A_{n+1}^{n-1} , we get

$$G_{n+1}^{n+1} \leq A_{n+1}^{n+1}.$$

□□□

15.0.6 Proof of the Harmonic Mean Geometric Mean Inequality

Let $a_1, a_2, \dots, a_n > 0$. We will prove the inequality.

We know that $G_n \leq A_n$, which implies

$$\begin{aligned} G_n &\leq A_n \\ \sqrt[n]{x_1 \dots x_2 \dots x_n} &\leq \frac{x_1 + x_2 + \dots + x_n}{n} \\ \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}} &\leq \sqrt[n]{x_1 \cdot x_2 \dots x_n}. \end{aligned}$$

This concludes the proof of the harmonic mean-geometric mean inequality.

□□□

Chapter 16

Solving Polynomial Equations

In this section, we discuss the solution formulas for polynomial equations of degrees 2 and 3: the PQ formula, the ABC formula, and the Cubic formula. We also derive each of them step by step.

16.0.1 The PQ Formula

The PQ formula solves quadratic equations of the form:

$$x^2 + px + q = 0.$$

Derivation

To derive the PQ formula, we complete the square

$$\begin{aligned}x^2 + px + q &= 0 \\x^2 + px &= -q \\x^2 + px + \left(\frac{p}{2}\right)^2 &= -q + \left(\frac{p}{2}\right)^2 \\ \left(x + \frac{p}{2}\right)^2 &= \left(\frac{p}{2}\right)^2 - q \\x + \frac{p}{2} &= \pm \sqrt{\left(\frac{p}{2}\right)^2 - q} \\x &= -\frac{p}{2} \pm \sqrt{\left(\frac{p}{2}\right)^2 - q}.\end{aligned}$$

PQ formula:

$$x = -\frac{p}{2} \pm \sqrt{\left(\frac{p}{2}\right)^2 - q}$$

16.0.2 The Quadratic Formula

The general quadratic equation is:

$$ax^2 + bx + c = 0 \quad \text{with } a \neq 0$$

Derivation of the PQ-Formula

We normalize the equation by dividing through by a and complete the square:

$$\begin{aligned}
ax^2 + bx + c &= 0 \\
x^2 + \frac{b}{a}x + \frac{c}{a} &= 0 \\
x^2 + \frac{b}{a}x &= -\frac{c}{a} \\
x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 &= -\frac{c}{a} + \left(\frac{b}{2a}\right)^2 \\
\left(x + \frac{b}{2a}\right)^2 &= \frac{b^2 - 4ac}{4a^2} \\
x + \frac{b}{2a} &= \pm \frac{\sqrt{b^2 - 4ac}}{2a} \\
x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}
\end{aligned}$$

ABC formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

16.0.3 The Cubic Formula

To solve a general cubic equation:

$$ax^3 + bx^2 + cx + d = 0,$$

we first reduce it to a depressed cubic using a substitution.

Step 1: Depress the cubic

Let $x = t - \frac{b}{3a}$, then the equation becomes:

$$t^3 + pt + q = 0$$

with:

$$p = \frac{3ac - b^2}{3a^2}, \quad q = \frac{2b^3 - 9abc + 27a^2d}{27a^3}$$

Step 2: Solve the depressed cubic using Cardano's method

Assume a solution of the form

$$t = u + v.$$

Then substitute and simplify

$$(u + v)^3 + p(u + v) + q = 0.$$

Expanding and setting

$$u^3 + v^3 + (3uv + p)(u + v) + q = 0.$$

To eliminate the $(u + v)$ term, set:

$$3uv + p = 0 \quad \Rightarrow \quad uv = -\frac{p}{3}.$$

Now

$$u^3 + v^3 = -q.$$

Let

$$u^3 = A, \quad v^3 = B \quad \Rightarrow \quad A + B = -q, \quad AB = -\frac{p^3}{27}.$$

These are the roots of the quadratic

$$z^2 + qz - \frac{p^3}{27} = 0;$$

solve for A and B , then take cube roots to get u and v . The final solution is:

$$x = u + v$$

Cardano's Formula (for depressed cubic)

$$x = \sqrt[3]{-\frac{q}{2} + \sqrt{\left(\frac{q}{2}\right)^2 + \left(\frac{p}{3}\right)^3}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\left(\frac{q}{2}\right)^2 + \left(\frac{p}{3}\right)^3}}$$

This formula gives one real root. The other roots (if real) can be found using trigonometric or complex methods depending on the discriminant.

Chapter 17

The Binomial Coefficient

In this short section, we will cover the definition and properties of the *binomial coefficient*.

$$\binom{n}{k} := \frac{n!}{k!(n-k)!},$$

or as a product:

$$\prod_{j=1}^n \frac{n-k+j}{j},$$

with n and k as natural numbers (for the moment).

Properties

- $\binom{n}{k} = \binom{n}{n-k}$
- $\binom{n}{k-1} = \binom{n}{k} \frac{n-k+1}{k}$
- $\sum_{k=0}^n \binom{n}{k} = 2^n$
- $\sum_{k=0}^n \binom{n}{k} = \binom{n+1}{n+1}$
- $\binom{n}{0} = 1 = \binom{n}{n}$
- $\binom{n}{k} + \binom{n}{k+1} = \binom{n+1}{k+1}$
- $\binom{n}{k} = (-1)^k \binom{k-n-1}{k}$

17.0.1 The Binomial Theorem

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

Proof:

We will prove that the equality holds for any n .

For n , we have $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$.

For $n+1$, we get:

$$(a+b)^{n+1} = \sum_{k=0}^{n+1} \binom{n+1}{k} a^{n+1-k} b^k.$$

Assume that the statement is true for some integer $n \geq 0$.

Now, for some greater n ,

Using the induction hypothesis:

We distribute:

In the second sum, change the index of summation by setting $j = k + 1$, then rename j back to k :

Now, combine both sums:

where we use the convention $\binom{n}{k} = 0$ for $k < 0$ or $k > n$.

By *Pascal's Identity*:

Therefore,

၇၉၇

17.0.2 Pascal's Triangle

This is a triangle built by the formula $\binom{n+1}{k+1} = \binom{n}{k} + \binom{n}{k+1}$.

138

Here is the triangle, but with the binomial coefficients:

$$\begin{array}{ccccccc}
 n = 0 & & & & & & \\
 n = 1 & & & & & & \\
 n = 2 & & & & & & \\
 n = 3 & & & & & &
 \end{array}
 \begin{array}{ccccccc}
 & & & & \binom{0}{0} & & \\
 & & & & \binom{1}{0} & \binom{1}{1} & \\
 & & & \binom{2}{0} & \binom{2}{1} & \binom{2}{2} & \\
 & \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \binom{3}{3} & &
 \end{array}$$

It can be used with the Binomial Theorem to get the coefficients of a polynomial of degree n .

Example:

$$(a + b)^2 = 1a^2b^0 + 2a^1b^1 + 1a^0b^2,$$

or

$$(a - b)^2 = 1a^2b^0 - 2a^1b^1 + 1a^0b^2.$$

Chapter 18

Proportionality and the Rule of Three

18.0.1 Proportionality

Two quantities are said to be *proportional* if their ratio remains constant.

Direct Proportionality

Two quantities a and b are in *direct proportion* if:

$$\frac{a}{b} = k \quad \Rightarrow \quad a = k \cdot b,$$

where k is the constant of proportionality.

Example:

If 2 pencils cost 1, then 4 pencils cost 2. The ratio is constant: $\frac{2}{1} = \frac{4}{2}$.

Inverse Proportionality

Two quantities a and b are in *inverse proportion* if their product is constant:

$$a \cdot b = k$$

Example:

If 4 workers finish a job in 6 hours, then 2 workers would need 12 hours:

$$4 \cdot 6 = 2 \cdot 12 = 24$$

18.0.2 The Rule of Three (Simple)

The *Rule of Three* is a method used to find a fourth value when three values are known and a proportional relationship is assumed.

Direct Rule of Three (Simple)

Given: $a : b = c : x$, solve for x :

$$x = \frac{b \cdot c}{a}$$

Example:

If 3 apples cost 6, how much do 5 apples cost?

$$x = \frac{6 \cdot 5}{3} = 10$$

Inverse Rule of Three (Simple)

If the relationship is inverse:

$$a : b = x : c \quad \Rightarrow \quad x = \frac{a \cdot c}{b}$$

Example:

If 5 people finish a task in 8 hours, how long will 10 people need?

$$x = \frac{5 \cdot 8}{10} = 4$$

18.0.3 The Rule of Three (Compound)

The *Compound Rule of Three* (or *composed rule of three*) involves more than two variables.

Example:

If 4 machines produce 120 items in 5 hours, how many items will 6 machines produce in 8 hours?

Step 1: Set up the proportional relationships:

$$\text{Items} \propto \text{Machines} \quad (\text{direct}) \quad \text{Items} \propto \text{Time} \quad (\text{direct})$$

Step 2: Adjust the quantities:

Initial: 4 machines, 5 hrs $\rightarrow$ 120 items

New: 6 machines, 8 hrs $\rightarrow x$ items

Step 3: Apply proportionality:

$$x = 120 \cdot \frac{6}{4} \cdot \frac{8}{5} = 120 \cdot 1.5 \cdot 1.6 = 288$$

Answer: 288 items.

Summary Table

Type	Formula
Direct Proportion	$x = \frac{b \cdot c}{a}$
Inverse Proportion	$x = \frac{a \cdot c}{b}$
Compound Rule of Three	Multiply by all direct ratios and divide by all inverse ratios

Chapter 19

Factorization

Factorization is the process of writing a mathematical expression as a product of its factors. This is a fundamental technique in algebra, used to simplify expressions, solve equations, and analyze functions.

19.0.1 Common Factor

Factor out the greatest common divisor (GCD) of all terms.

Example:

$$6x^2 + 9x = 3x(2x + 3)$$

19.0.2 Difference of Squares

$$a^2 - b^2 = (a - b)(a + b)$$

19.0.3 Sum of Cubes

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$$

19.0.4 Difference of Cubes

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

Example:

$$x^3 + 8 = (x + 2)(x^2 - 2x + 4)$$

19.0.5 Trinomial: Special Case $x^2 + bx + c$

This is the case where $a = 1$. Find two numbers whose product is c and whose sum is b .

Example:

$$x^2 + 5x + 6 = (x + 2)(x + 3)$$

19.0.6 Trinomial: General Form $ax^2 + bx + c$

As in the previous case, we are going to use a similar procedure, but we start by multiplying and dividing by $\frac{a}{a}$, and then proceed with the rest. Look at the example:

$$\begin{aligned}
3x^2 - 5x - 2 &= \frac{3(3x^2 - 5x - 2)}{3} \\
&= \frac{(3x)^2 - 5(3x) - 6}{3} \\
&= \frac{(3x - 6)(3x + 1)}{3} \\
&= (x - 2)(3x + 1)
\end{aligned}$$

19.0.7 Perfect Square Trinomial

These follow the identities:

$$(a + b)^2 = a^2 + 2ab + b^2$$

$$(a - b)^2 = a^2 - 2ab + b^2$$

Example:

$$x^2 + 6x + 9 = (x + 3)^2$$

19.0.8 Substitution

Substitute a more complex expression with a single variable, factor, then back-substitute.

Example:

Given $x^4 + 2x^2 + 1$. Let $y = x^2$, then

$$y^2 + 2y + 1 = (y + 1)^2 = (x^2 + 1)^2$$

19.0.9 Rationalization of Radicals

Rationalizing removes radicals from the denominator.

Example (Single Radical):

$$\frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$$

Example (Binomial):

$$\frac{1}{\sqrt{3} + 1} = \frac{\sqrt{3} - 1}{(\sqrt{3} + 1)(\sqrt{3} - 1)} = \frac{\sqrt{3} - 1}{2}$$

19.0.10 Horner's Method

This method is used to divide a polynomial by a binomial of the form $(x - r)$.

We can apply this method by following these steps:

1. Write the coefficients of the polynomial $P_1(x)$.
2. Find the factors of the term with x^0 .
3. Bring down the first coefficient.
4. Multiply it by r , and add to the next coefficient. Repeat until you reach the term with x^0 .

5. If the final addition gives zero, then you have found a zero of the polynomial. Now, it can be written in the following form: $P_2(x)(x-r)$, with $P(x)$ being the new polynomial whose coefficients were computed during the process to get the zero.

Example:

Given $P_1(x) = x^3 - 6x^2 + 11x - 6$.

The factors of -6 are $\{\pm 1, \pm 2, \pm 3, \pm 6\}$. We use $r = 1$:

$$\begin{array}{r|rrrr} 1 & 1 & -6 & 11 & -6 \\ & & 1 & -5 & 6 \\ \hline & 1 & -5 & 6 & 0 \end{array} \implies P_2(x) = (x^2 - 5x + 6)(x - 1)$$

Now, by repeating the process with $r = 2$:

$$\begin{array}{r|rrr} 2 & 1 & -5 & 6 \\ & & 2 & -6 \\ \hline & 1 & -3 & 0 \end{array} \implies P_3(x) = (x - 3)(x - 2)(x - 1)$$

The final result is $(x - 3)(x - 2)(x - 1)$.

19.0.11 Long Division of Polynomials

Another way of factorization is the so-called *long division for polynomials*. This works exactly as the division algorithm for real numbers, but we use powers of x and the scalars of $\mathbb{F}$ for the multiplication step.

Example:

$$\begin{array}{r} x^2 + 2x + 2 \\ x - 1 \overline{) \begin{array}{r} x^3 + x^2 - 1 \\ - x^3 + x^2 \\ \hline 2x^2 \\ - 2x^2 + 2x \\ \hline 2x - 1 \\ - 2x + 2 \\ \hline 1 \end{array}} \end{array}$$

with the result: $x^2 + 2x + 2 + \frac{1}{x-1}$.

Chapter 20

Partial Fractions

20.0.1 The Simplest Case

In the most common partial fraction decomposition, we split:

$$\frac{N(x)}{(x - a_1)(x - a_2) \cdots (x - a_d)}$$

into a sum of the form:

$$\frac{A_1}{x - a_1} + \cdots + \frac{A_d}{x - a_d}.$$

We now show that this decomposition can always be achieved, under the assumption that the a_i are all different and $N(x)$ is a polynomial of degree at most $d - 1$.

Lemma 1

Let $N(x)$ and $D(x)$ be polynomials of degree n and d , respectively, with $n \leq d$. Suppose that a is not a root of $D(x)$. Then there exists a polynomial $P(x)$ of degree $< d$ and a number A such that:

$$\frac{N(x)}{D(x)(x - a)} = \frac{P(x)}{D(x)} + \frac{A}{x - a}.$$

Proof:

Let $z = x - a$. Define:

$$\tilde{N}(z) = N(z + a), \quad \tilde{D}(z) = D(z + a).$$

Then

$$\frac{\tilde{N}(z)}{\tilde{D}(z)z} = \frac{\tilde{P}(z)}{\tilde{D}(z)} + \frac{A}{z} \Rightarrow \frac{\tilde{P}(z)z + A\tilde{D}(z)}{\tilde{D}(z)z}$$

We equate

$$\tilde{P}(z)z + A\tilde{D}(z) = \tilde{N}(z)$$

Choosing $A = \frac{\tilde{N}(0)}{\tilde{D}(0)}$, the constant terms match. The remainder has no constant term and is divisible by z , so:

$$\tilde{P}(z)z = \tilde{N}(z) - A\tilde{D}(z)$$

Thus, $\tilde{P}(z)$ is a polynomial of degree $< d$.

Recursive Decomposition

Now, consider

$$\frac{N(x)}{(x-a_1)(x-a_2)\cdots(x-a_d)}.$$

Apply Lemma 1 recursively

$$\frac{N(x)}{(x-a_1)\cdots(x-a_d)} = \frac{A_1}{x-a_1} + \frac{P(x)}{(x-a_2)\cdots(x-a_d)}.$$

Then

$$\frac{P(x)}{(x-a_2)\cdots(x-a_d)} = \frac{A_2}{x-a_2} + \frac{Q(x)}{(x-a_3)\cdots(x-a_d)}.$$

Continue until

$$\frac{N(x)}{(x-a_1)\cdots(x-a_d)} = \frac{A_1}{x-a_1} + \cdots + \frac{A_d}{x-a_d}.$$

20.0.2 Lemma 2

Let $N(x)$ and $D(x)$ be polynomials of degree n and d respectively, with $n < d + m$. Suppose that a is NOT a zero of $D(x)$. Then there is a polynomial $P(x)$ of degree $p < d$ and numbers $A_1, \dots, A_m$ such that

$$\frac{N(x)}{D(x)(x-a)^m} = \frac{P(x)}{D(x)} + \frac{A_1}{x-a} + \frac{A_2}{(x-a)^2} + \cdots + \frac{A_m}{(x-a)^m}.$$

Proof:

To save writing, let $z = x - a$. Then $\tilde{N}(z) = N(z+a)$ and $\tilde{D}(z) = D(z+a)$ are polynomials of degree n and d respectively, $\tilde{D}(0) = D(a) \neq 0$ and we have to find a polynomial $\tilde{P}(z)$ of degree $p < d$ and numbers $A_1, \dots, A_m$ such that

$$\begin{aligned} \frac{\tilde{N}(z)}{\tilde{D}(z)z^m} &= \frac{\tilde{P}(z)}{\tilde{D}(z)} + \frac{A_1}{z} + \frac{A_2}{z^2} + \cdots + \frac{A_m}{z^m} \\ &= \frac{\tilde{P}(z)z^m + A_1z^{m-1}\tilde{D}(z) + A_2z^{m-2}\tilde{D}(z) + \cdots + A_m\tilde{D}(z)}{\tilde{D}(z)z^m} \end{aligned}$$

or equivalently, such that

$$\tilde{P}(z)z^m + A_1z^{m-1}\tilde{D}(z) + A_2z^{m-2}\tilde{D}(z) + \cdots + A_{m-1}z\tilde{D}(z) + A_m\tilde{D}(z) = \tilde{N}(z)$$

Now look at the polynomial on the left-hand side. Every single term on the left-hand side, except for the very last one, $A_m\tilde{D}(z)$, has at least one power of z . So the constant term on the left-hand side is exactly the constant term in $A_m\tilde{D}(z)$, which is $A_m\tilde{D}(0)$. The constant term on the right-hand side is $\tilde{N}(0)$. So

the constant terms on the left and right-hand sides are the same if we choose $A_m = \frac{\tilde{N}(0)}{\tilde{D}(0)}$. Recall that $\tilde{D}(0) \neq 0$. Now move $A_m \tilde{D}(z)$ to the right-hand side.

$$\tilde{P}(z)z^m + A_1 z^{m-1} \tilde{D}(z) + A_2 z^{m-2} \tilde{D}(z) + \cdots + A_{m-1} z \tilde{D}(z) = \tilde{N}(z) - A_m \tilde{D}(z).$$

The constant terms in $\tilde{N}(z)$ and $A_m \tilde{D}(z)$ are the same, so the right-hand side contains no constant term and the right-hand side is of the form $\tilde{N}_1(z)z$ with $\tilde{N}_1$ a polynomial of degree at most $d+m-2$. (Recall that $\tilde{N}$ is of degree at most $d+m-1$ and $\tilde{D}$ is of degree at most d .) Divide the whole equation by z .

$$\tilde{P}(z)z^{m-1} + A_1 z^{m-2} \tilde{D}(z) + A_2 z^{m-3} \tilde{D}(z) + \cdots + A_{m-1} \tilde{D}(z) = \tilde{N}_1(z).$$

Now, we can repeat the previous argument. The constant term on the left-hand side, which is exactly $A_{m-1} \tilde{D}(0)$ matches the constant term on the right-hand side, which is $\tilde{N}_1(0)$ if we choose $A_{m-1} = \frac{\tilde{N}_1(0)}{\tilde{D}(0)}$. With this choice of A_{m-1} .

$$\tilde{P}(z)z^{m-1} + A_1 z^{m-2} \tilde{D}(z) + A_2 z^{m-3} \tilde{D}(z) + \cdots + A_{m-2} z \tilde{D}(z) = \tilde{N}_1(z) - A_{m-1} \tilde{D}(z) = \tilde{N}_2(z)z.$$

With $\tilde{N}_2$ a polynomial of degree at most $d+m-3$. Divide by z and continue. After m steps like this, we end up with

$$\tilde{P}(z)z = \tilde{N}_{m-1}(z) - A_1 \tilde{D}(z).$$

After having chosen $A_1 = \frac{\tilde{N}_{m-1}(0)}{\tilde{D}(0)}$. There is no constant term on the right side so that $\tilde{N}_{m-1}(z) - A_1 \tilde{D}(z)$ is of the form $\tilde{N}_m(z)z$ with $\tilde{N}_m$ a polynomial of degree $d-1$. Choosing $\tilde{P}(z) = \tilde{N}_m(z)$ completes the proof.

Now back to

$$\frac{N(x)}{(x-a_1)^{n_1} \times \cdots \times (x-a_d)^{n_d}}.$$

Apply Lemma 2, with $D(x) = (x-a_2)^{n_2} \times \cdots \times (x-a_d)^{n_d}$, $m = n_1$ and $a = a_1$. It says

$$\frac{N(x)}{(x-a_1)^{n_1} \times \cdots \times (x-a_d)^{n_d}} = \frac{P(x)}{(x-a_2)^{n_2} \times \cdots \times (x-a_d)^{n_d}} + \frac{A_{1,1}}{x-a_1} + \frac{A_{1,2}}{(x-a_1)^2} + \cdots + \frac{A_{1,n_1}}{(x-a_1)^{n_1}}.$$

Apply Lemma 2 a second time, with $D(x) = (x-a_3)^{n_3} \times \cdots \times (x-a_d)^{n_d}$, $N(x) = P(x)$, $m = n_2$ and $a = a_2$. And so on. Eventually, we end up with

$$\frac{N(x)}{(x-a_1)^{n_1} \times \cdots \times (x-a_d)^{n_d}} = \left[\frac{A_{1,1}}{x-a_1} + \cdots + \frac{A_{1,n_1}}{(x-a_1)^{n_1}} \right] + \cdots + \left[\frac{A_{d,1}}{x-a_d} + \cdots + \frac{A_{d,n_d}}{(x-a_d)^{n_d}} \right].$$

Chapter 21

Recursion

Recursion is a term used to refer to something defined in terms of itself. An example is the factorial: $n! = (n-1)!$, with $0! = 1$ being the base case where the recursion stops.

21.0.1 From Recursive to Closed Form

Often, a sequence defined recursively can be rewritten in an explicit or closed form.

Recursive form defines a_n based on previous terms (e.g., $a_n = a_{n-1} + 2$).

Closed form expresses a_n directly in terms of n , with no reference to previous terms.

21.0.2 Linear Recursion

Linear recursion has the form:

$$\begin{aligned}a_n &= \lambda a_{n-1} + c \\a_1 &= \lambda a_0 + c \implies a_2 = \lambda(\lambda a_0 + c) + c \\ \implies a_n &= \lambda^n a_0 + (1 + \lambda + \lambda^2 + \cdots + \lambda^{n-1})c.\end{aligned}$$

Therefore,

$$a_n = \lambda^n a_0 + \frac{\lambda^n - 1}{\lambda - 1} c.$$

General Method for Linear Recursion

1. Calculate some terms and observe the difference or quotient between them. Maybe the pattern is clear, and you can build a geometric or arithmetic sequence directly.
2. Identify the type (e.g., linear, homogeneous, constant coefficients).
3. Solve the characteristic equation (if linear).
4. Use initial conditions to determine constants.

21.0.3 Recursive Inequalities

Recursive inequalities bound a sequence rather than define it exactly.

Example:

Suppose

$$a_{n+1} \leq \frac{1}{2}a_n + 3.$$

To analyze

- Guess a bound (e.g., show $a_n \leq M$ for all n).
- Use induction or iteration to justify convergence or boundedness.
- Compare with a simpler sequence (e.g., geometric series).

These techniques are common in analysis, approximation, and numerical algorithms.

21.0.4 Fibonacci Numbers and Closed Form

The Fibonacci sequence is defined recursively:

$$F_0 = 0, \quad F_1 = 1, \quad F_n = F_{n-1} + F_{n-2}.$$

The characteristic equation is:

$$x^2 - x - 1 = 0 \Rightarrow x = \varphi = \frac{1 + \sqrt{5}}{2}, \quad \psi = \frac{1 - \sqrt{5}}{2}.$$

The closed form (Binet's Formula) is

$$F_n = \frac{1}{\sqrt{5}} (\varphi^n - \psi^n).$$

- $\varphi \approx 1.618$ is the golden ratio
- This formula demonstrates exponential growth of F_n .

Chapter 22

Sum and Product Notation

multiplication, respectively. These notations are essential for working with sequences, series, and algebraic expressions. In mathematics, the sum and product notations are compact ways to represent repeated addition and multiplication, respectively. These notations are essential for working with sequences, series, and algebraic expressions.

22.0.1 Sum Notation $\sum$

The summation symbol $\sum$ represents the addition of a sequence of terms:

$$\sum_{i=m}^n a_i = a_m + a_{m+1} + \cdots + a_n$$

Here, i is the index of summation, m is the lower bound, and n is the upper bound.

Linearity:

$$\sum_{i=m}^n (a_i + b_i) = \sum_{i=m}^n a_i + \sum_{i=m}^n b_i$$

$$\sum_{i=m}^n c \cdot a_i = c \cdot \sum_{i=m}^n a_i$$

Splitting:

$$\sum_{i=m}^n a_i = \sum_{i=m}^k a_i + \sum_{i=k+1}^n a_i \quad (m \leq k < n)$$

Change of Index

Let $j = i + k$, then:

$$\sum_{i=m}^n a_i = \sum_{j=m+k}^{n+k} a_{j-k}$$

Example:

$$\sum_{i=1}^4 a_i = \sum_{j=2}^5 a_{j-1}$$

Power Sums and Their Formulas

$$\begin{aligned}\sum_{i=1}^n i &= \frac{n(n+1)}{2} \\ \sum_{i=1}^n i^2 &= \frac{n(n+1)(2n+1)}{6} \\ \sum_{i=1}^n i^3 &= \left[\frac{n(n+1)}{2} \right]^2 \\ \sum_{i=1}^n i^k &= (\text{Higher-order polynomial in } n)\end{aligned}$$

Derivation of $\sum_{i=1}^n i^2$

We use the method of finite differences or induction. Assume a quadratic form:

$$\sum_{i=1}^n i^2 = An^3 + Bn^2 + Cn$$

Plug in small values of n (e.g., 1, 2, 3), solve the system of equations to find:

$$A = \frac{1}{3}, \quad B = \frac{1}{2}, \quad C = \frac{1}{6} \Rightarrow \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

Telescoping Sum

A telescoping sum is a sum in which intermediate terms cancel out, leaving only the first and last terms.

$$\sum_{k=m}^n (a_k - a_{k+1}) = a_m - a_{n+1}$$

Example:

$$\sum_{i=1}^n \left(\frac{1}{i} - \frac{1}{i+1} \right) = 1 - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \cdots + \frac{1}{n} - \frac{1}{n+1} = 1 - \frac{1}{n+1}$$

22.0.2 Geometric Series

Finite Geometric Series:

For a geometric sequence $a, ar, ar^2, \dots, ar^{n-1}$:

$$\sum_{i=0}^{n-1} ar^i = a \cdot \frac{1-r^n}{1-r}, \quad r \neq 1$$

Infinite Geometric Series:

If $|r| < 1$, then:

$$\sum_{i=0}^{\infty} ar^i = \frac{a}{1-r}$$

22.0.3 Product Notation $\prod$

The product notation $\prod$ represents repeated multiplication:

$$\prod_{i=m}^n a_i = a_m \cdot a_{m+1} \cdot \cdots \cdot a_n$$

Multiplication:

$$\prod_{i=m}^n (a_i \cdot b_i) = \left(\prod_{i=m}^n a_i \right) \cdot \left(\prod_{i=m}^n b_i \right)$$

Power Rule:

$$\prod_{i=m}^n a^k = a^{k(n-m+1)}$$

Change of Index

Let $j = i + k$, then:

$$\prod_{i=m}^n a_i = \prod_{j=m+k}^{n+k} a_{j-k}$$

Example:

$$\prod_{i=1}^3 a_i = \prod_{j=2}^4 a_{j-1}$$

Telescoping Product

A telescoping product occurs when consecutive terms simplify or cancel each other.

$$\prod_{k=m}^n \frac{a_k}{a_{k+1}} = \frac{a_m}{a_{n+1}}$$

Example:

$$\prod_{i=1}^n \frac{i}{i+1} = \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4} \cdots \frac{n}{n+1} = \frac{1}{n+1}$$

22.0.4 Proof of the geometric Series

Let $S_n = \sum_{k=0}^n aq^k$ with $|q| < 1$.

Then $S_n q = a(\sum_{k=0}^{n+1} q^k) = a(\sum_{k=0}^n q^k + q^{n+1})$.

$$a + S_n q = \sum_{k=0}^n aq^k + aq^{n+1} = S_n + aq^{n+1}$$

Finally, we subtract S_n and a , and then divide by $q - 1$:

$$S_n = \frac{a - aq^{n+1}}{q - 1}$$

Chapter 23

Logarithms

The *logarithm* of a number x with base b is the exponent to which the base must be raised to produce that number. It is denoted as:

$$\log_b(x) = y \iff b^y = x,$$

where $b > 0$, $b \neq 1$, and $x > 0$.

$$\log_2(8) = 3 \quad \text{because} \quad 2^3 = 8.$$

We can also approximate a logarithm by dividing our target x by the base b repeatedly until we get 1 or less.

23.0.1 Properties of Logarithms

- $\log_b(xy) = \log_b(x) + \log_b(y)$
- $\log_b\left(\frac{x}{y}\right) = \log_b(x) - \log_b(y)$
- $\log_b(x^k) = k \cdot \log_b(x)$
- $\log_b(b) = 1$
- $\log_b(1) = 0$
- $\log_{b^k}(x^w) = \frac{w}{k} \cdot \log_b(x)$
- $\log_b\left(\frac{1}{x}\right) = -\log_b(x)$
- $\log_b(b^x) = x$
- $\log_b(x) = \frac{\log_k(x)}{\log_k(b)}$ for any positive $k \neq 1$
- $e^{\ln(x)} = x$
- If $0 < a < 1$, then $\ln(a)$ is a negative number.
- $\log(x) < \log(y)$ for $x < y$
- $\log(x+1) \leq x$ for $x > -1$
- For all $c > 0$, there is some M such that $\log(x) < cx$ for all $x \in (0, \infty)$, $x > M$

23.0.2 Fundamental Identity of Logarithms

$$a^{\log_a(x)} = x$$

23.0.3 Change of Base Formula

$$\log_b(x) = \frac{\log_k(x)}{\log_k(b)},$$

where k is any positive number different from 1.

Proof:

23.0.4 The Chain Rule

$$\log_y(a) \log_a(b) = \log_y(b)$$

Proof:

23.0.5 The Derivative of the Natural Logarithm

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Let $f(x) = \ln(x)$, then

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{\ln(x+h) - \ln(x)}{h} \\ &= f'(x) = \lim_{h \rightarrow 0} \frac{\ln\left(\frac{x+h}{x}\right)}{h} \\ &= f'(x) = \lim_{h \rightarrow 0} \frac{\ln\left(1 + \frac{h}{x}\right)}{h} \\ &= f'(x) = \lim_{h \rightarrow 0} \ln\left(1 + \frac{h}{x}\right)^{\frac{1}{h}} \end{aligned}$$

Now, let $n = \frac{h}{x}$.

$$\begin{aligned} f'(x) &= \lim_{n \rightarrow 0} \ln(1+n)^{\frac{x}{h} \frac{1}{x}} \\ f'(x) &= \lim_{n \rightarrow 0} \frac{1}{x} \ln(1+n)^{\frac{1}{n}} \\ f'(x) &= \frac{1}{x} \ln\left(\lim_{n \rightarrow 0} (1+n)^{\frac{1}{n}}\right) \\ f'(x) &= \frac{1}{x} \end{aligned}$$

Therefore, the derivative of the natural logarithm is:

$$\frac{d}{dx} \ln(x) = \frac{1}{x}$$

QED

23.0.6 Proofs of the Properties

Product

Let $a^n a^m = a^{n+m}$, and $a^n = x$, and $a^m = y$, then

$$\log_a x = n \text{ and } \log_a y = m \rightarrow \log_a(xy) = n + m$$

$$\log_a(xy) = \log_a x + \log_a y$$

□◻◻

Quotient

Let $\frac{a^n}{a^m} = a^{n-m}$, and $a^n = x$, and $a^m = y$, then as in the previous proof,

$$\log_a \left(\frac{x}{y} \right) = \log_a x - \log_a y$$

□◻◻

Power

Let $(a^n)^m = a^{nm}$, and $a^n = x$, and $a^{nm} = y$.

$$\log_x y = m \text{ and } \log_a x = n \text{ thus, } \log_a y = (\log_a x)m$$

□◻◻

Power Rule: General Case

Let $(a^n)^m = b^x$, then $\log_{a^n} b^x = m$. Now, let us manipulate the original expression:

$$\begin{aligned} \log_a (a^n)^m &= \log_a b^x \\ nm \log_a a &= x \log_a b \\ m &= \frac{x}{n} \log_a b = \log_{a^n} b^x \end{aligned}$$

□◻◻

Chapter 24

Fractions, Roots, and Exponents

This is a brief chapter to review the properties of fractions, roots, and exponents. It is not a complex chapter, but it is useful to include it in the compendium.

The following statements apply for $a, b \in \mathbb{R}$.

24.0.1 Fractions

- $\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}$
- $\frac{a}{b} \pm \frac{c}{d} = \frac{ad \pm bc}{bd}$
- $\frac{a}{b} \div \frac{c}{d} = \frac{ad}{bc}$
- $\frac{a}{b} = \frac{c}{d} \iff ad = bc$
- $\frac{ac}{bc} = \frac{a}{b}$

24.0.2 Roots and Exponents

- $\sqrt{a} \cdot \sqrt{b} = \sqrt{ab}$
- $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$
- $\sqrt[n]{\sqrt[m]{a}} = \sqrt[nm]{a}$
- $a \sqrt[n]{b} = \sqrt[n]{a^n b}$
- $a^m \cdot a^n = a^{m+n}$
- $a^m \div a^n = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^{-n} = \frac{1}{a^n}$
- $a^{\frac{m}{n}} = \sqrt[n]{a^m}$
- $a^{\frac{1}{2}} = \sqrt{a}$
- $a^{\frac{1}{3}} = \sqrt[3]{a}$
- $a^{\frac{1}{n}} = \sqrt[n]{a}$
- $a^0 = 1$
- $a^1 = a$

Chapter 25

Generating Functions

Given a sequence $\{a_n\}_{n \geq 0}$, its *generating function* is defined as:

$$F(x) = \sum_{n=0}^{\infty} a_n x^n.$$

Here, x is an indeterminate, and the series is usually treated formally, meaning we do not worry about convergence unless the context requires it.

25.0.1 Connection to Subsets

There is a deep similarity between generating functions and the process of forming subsets. Consider the generating function for the set $\{1, x\}$:

$$(1 + x)^n = \sum_{k=0}^n \binom{n}{k} x^k.$$

This expansion encodes all subsets of an n -element set, where each term x^k corresponds to selecting a subset of size k . The coefficient $\binom{n}{k}$ gives the number of such subsets.

In this way, generating functions can be viewed as a powerful algebraic analogue of subset selection and counting, with the exponents representing sizes or weights, and the coefficients representing counts.

Generating functions serve as a bridge between algebra and combinatorics, allowing abstract operations to yield concrete combinatorial results.

25.0.2 Closed Form of the Fibonacci Sequence

Let us use generating functions to find a closed form for the Fibonacci numbers.

We know that $F_0 = 1$ and $F_1 = 1$, and for $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$.

Now, let us use the following function:

$$F(x) = \sum_{n=0}^{\infty} F_n x^n,$$

which is going to have the Fibonacci numbers as the coefficients of the power series.

$$1 + x + 2x^2 + 3x^3 + 5x^4 + \dots,$$

But before that, we have to look at when our recursive formula is defined.

In this case, $n = 2$. Thus, we have to write our expression in the following way:

$$\begin{aligned} F(x) &= 1 + x + \sum_{n=2}^{\infty} F_n x^n \\ &= 1 + x + \sum_{n=2}^{\infty} F_{n-1} x^n + \sum_{n=2}^{\infty} F_{n-2} x^n. \end{aligned}$$

Here, we are going to adjust our expression by making our n be equal.

$$\begin{aligned} F(x) &= 1 + x + \sum_{n=2}^{\infty} F_{n-1} x^n + \sum_{n=2}^{\infty} F_{n-2} x^n \\ F(x) &= 1 + x + x \left(\sum_{n=2}^{\infty} F_{n-1} x^{n-1} \right) + x^2 \left(\sum_{n=2}^{\infty} F_{n-2} x^{n-2} \right). \end{aligned}$$

Now, we are going to reset our sum to start from zero by adding a smart zero to our term with $n - 1$. This allows us to shift the sum to start from 0, because when $n = 1$, $F_{n-1} x^{n-1}$ turns into $F_0 x^0$.

$$\begin{aligned} F(x) &= 1 + x + x \left(-F_0 x^0 + F_0 x^0 + \sum_{n=2}^{\infty} F_{n-1} x^{n-1} \right) + x^2 \left(\sum_{n=2}^{\infty} F_{n-2} x^{n-2} \right) \\ F(x) &= 1 + x + x \left(-F_0 x^0 + \sum_{n=1}^{\infty} F_{n-1} x^{n-1} \right) + x^2 \left(\sum_{n=2}^{\infty} F_{n-2} x^{n-2} \right) \\ F(x) &= 1 + x + x \left(\sum_{n=0}^{\infty} F_n x^n \right) + x^2 \left(\sum_{n=2}^{\infty} F_{n-2} x^{n-2} \right) \\ F(x) &= 1 + x + xF(x) - x + x^2 F(x), \end{aligned}$$

which, at the end, simplifies to:

$$F(x) = \frac{1}{1 - x - x^2}.$$

If you start with $F_0 = 0$ then the result is

$$F(x) = \frac{x}{1 - x - x^2}.$$

denominator. Now, we can use partial fraction decomposition to make this expression even more explicit by using the following expression, where α and β are zeros of the polynomial in the denominator.

$$\frac{1}{(1 - \alpha x)(1 - \beta x)}.$$

The zeros of the polynomial are:

$$x_1 = \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad x_2 = \frac{1 - \sqrt{5}}{2}.$$

Now, we can write our expression as:

$$F(x) = \frac{1}{1-x-x^2} = \frac{1}{\left(1-x \cdot \frac{1+\sqrt{5}}{2}\right) \left(1-x \cdot \frac{1-\sqrt{5}}{2}\right)},$$

which, after performing the partial fraction decomposition, leaves us with:

$$F(x) = \left(\left(\frac{1}{\sqrt{5}}(1 + \sqrt{5}) \right)^n - \left(\frac{1}{\sqrt{5}}(1 - \sqrt{5}) \right)^n \right).$$

□◻◻

Chapter 26

Modular Arithmetic

We define an equivalence relation

$$a \cong b \pmod n \iff n|(a - b) \wedge a, b \in \mathbb{Z}.$$

This means that both have the same remainder when divided by n .

26.0.1 Residual Classes and Residual Class Rings

Because we have an equivalence relation, we can classify the partitions of our set $\mathbb{Z}$ into classes, which we will call *residual classes*.

Example:

$$[13]_7 + \{m \in \mathbb{Z} \mid m \cong 13 \pmod 7\}$$

The set of all residual classes can be written as $\mathbb{Z}/n\mathbb{Z}$, where n is the modulus.

26.0.2 Properties

Addition

$$(a + b) \pmod n = (a \pmod n + b \pmod n) \pmod n$$

Multiplication

$$(a * b) \pmod n = (a \pmod n * b \pmod n) \pmod n$$

Exponentiation

$$(a^m) \pmod n = ((a \pmod n)^n) \pmod n$$

Division

$$\left(\frac{a}{b}\right) \pmod n \neq \frac{a \pmod n}{b \pmod n}$$

26.0.3 General Division Rules

Sum Rules

If the number a can be expressed as a sum and b is a common divisor of all the terms, then b is also a divisor of a .

Example:

$742 = 700 + 42$. Then 7, as a common divisor of 700 and 42, is a divisor of 742.

Product Rule

If the number a can be written as a product of two factors and b is a divisor of one of these factors, then b is also a divisor of a .

Example:

$1500 = 15 \cdot 100$. Then 3 and 5 are divisors of 15, and 2, 4, 5, 10, 25, and 50 are divisors of 100—thus, all are divisors of 1500.

Divisor Pair Rules

If the number a has b as a divisor, then $\frac{a}{b}$ is also a divisor of a .

Example:

3 is a divisor of 315. Then $105 = \frac{315}{3}$ is also a divisor of 315.

Divisor Rule

If the number a has b as a divisor, then every divisor of b is also a divisor of a .

Example:

45 is a divisor of 450. Then 3, 5, 9, and 15 (divisors of 45) are also divisors of 450.

Difference Rule

If the number a can be expressed as a difference and b is a common divisor of the minuend and subtrahend, then b is also a divisor of a .

Example:

$686 = 700 - 14$. Then 7, as a common divisor of 700 and 14, is a divisor of 686.

Relative Prime Rule

If the number a has two relatively prime numbers b and c as divisors, then their product $b \cdot c$ is also a divisor of a .

Example:

1540 has the relatively prime numbers 7 and 11 as divisors. Then $77 = 7 \cdot 11$ is also a divisor of 1540.

26.0.4 Using Modular Arithmetic to Find the n-th Digit of a Large Number

What is the last digit of the number 13^{2023} ?

Note that $13 \bmod 10 = 3$ and $13^2 \bmod 10 = ((13 \bmod 10)(13 \bmod 10)) \bmod 10 = 9$.

We can continue this process, and we will notice a pattern. After some iterations, we will know that

$$(13^{2020}13) \bmod 10 = 7.$$

26.0.5 Gaussian Brackets

in specific ways: The Gaussian brackets refer to two closely related functions used to round real numbers to integers in specific ways:

Floor Function

The floor function, denoted by $\lfloor x \rfloor$, gives the greatest integer less than or equal to a real number x .

Example:

$$\lfloor 3.7 \rfloor = 3, \lfloor -1.2 \rfloor = -2$$

Ceiling Function

The ceiling function, denoted by $\lceil x \rceil$, gives the smallest integer greater than or equal to a real number x .

Example:

$$\lceil 3.7 \rceil = 4, \lceil -1.2 \rceil = -1$$

and rounding operations, where precise integer bounds of real numbers are required. These functions are particularly useful in number theory, computer science, and rounding operations, where precise integer bounds of real numbers are required.

26.0.6 Residual Class Ring modulo n

Given $n \in \mathbb{N}$, we define the relation $\sim_n$ for $\mathbb{Z}$.

$$a \sim b \iff n \mid (a - b).$$

Also, a/n and b/n have the same remainder.

26.0.7 Equivalence Class of $\mathbb{Z}$

For $i, n \in \mathbb{Z}$

$$\bar{i} := i + n\mathbb{Z} = \{i + na \mid a \in \mathbb{Z}\}$$

and

$$\mathbb{Z}_N = \mathbb{Z}/n\mathbb{Z} = \{\bar{0} := n\mathbb{Z}, \bar{1} := 1 + n\mathbb{Z}, \dots, \overline{n-1} := n-1 + n\mathbb{Z}\},$$

is the set of all equivalence classes for $\mathbb{Z}$ with respect to the relation $k \sim l \iff n \mid (k - l)$. We define the addition and multiplication for $\mathbb{Z}/n\mathbb{Z}$ using the representatives:

$$(a + n\mathbb{Z}) + (b + n\mathbb{Z}) := (a + b) + n\mathbb{Z}$$

and

$$(a + n\mathbb{Z}) \cdot (b + n\mathbb{Z}) := ab + n\mathbb{Z}$$

For all $a, b \in \mathbb{Z}$. Therefore,

$$\bar{a} + \bar{b} = \overline{a+b}, \quad \bar{a} \cdot \bar{b} = \overline{ab}.$$

Chapter 27

Relational Algebra

27.0.1 Relations

In mathematics, a *relation* is a set formed from the Cartesian product of two sets. It contains ordered pairs (a, b) .

27.0.2 The Cartesian Product for Relation Algebra

The Cartesian Product for our entities is defined as:

$$S \times T := \bigcup_{(s_1, \dots, s_n) \in S} \left[\bigcup_{(t_1, \dots, t_k) \in T} \{(s_1, \dots, s_n, t_1, \dots, t_k)\} \right]$$

Simplified: each element of S is combined with each element of T . The size of this special product is $|S \times T| = |S| \cdot |T|$.

In relational algebra, relations represent tables and their rows are the tuples.

27.0.3 Relational Algebra

Relational Algebra is a *formal, procedural query language* for the relational model. It provides a set of operators that take relations (tables) as input and produce relations as output.

Axioms of Relational Algebra

Relational Algebra is based on *set theory* and satisfies the following axioms:

- *Closure*: Every operation produces a relation.
- *Set Semantics*: Relations do not contain duplicate tuples.
- *Order Irrelevance*: The order of tuples and attributes does not matter.
- *Composability*: Operations can be nested arbitrarily.

Example:

If $\sigma_{id>1}(A)$ returns a relation, it can be used as input to another operator such as projection.

Relations

In database theory, a *relation* is a finite set of tuples over a fixed set of attributes.

Formally:

$$R \subseteq D_1 \times D_2 \times \dots \times D_n$$

Example:

Relation $A(id, value)$ is a set of tuples:

$$A = \{(1, X), (2, Y)\}$$

Identifier and Schema

We define:

$$\text{ident}(R) = \{a_i\}_i \quad \text{and} \quad \text{schema}(R) = \{[a_1 : T_1, \dots, a_n : T_n]\}.$$

ident returns the names of the attributes of the relation, while schema returns the attributes and their corresponding types.

27.0.4 Domain of an Attribute

The *domain* of an attribute a in a relation R is the set of all possible values that a can take in R .

Cartesian Product for Relations

The *Cartesian product* combines every tuple of one relation with every tuple of another.

$$S \times T = \bigcup_{(s_1, \dots, s_n) \in S} \left[\bigcup_{(t_1, \dots, t_n) \in T} \{(s_1, \dots, s_n, t_1, \dots, t_k)\} \right]$$

With size $|S \times T| = |S| \times |T|$.

```
SELECT * FROM T, S;
-- or equivalently:
SELECT * FROM T CROSS JOIN S;
```

Union ($\cup$)

Combines tuples from two relations with the *same schema*.

$$A \cup B$$

For the union of relations they have to be *union compatible* which means:

- They have the same numbers of elements in the tuple.
- The relations need to have the same columns.

```
SELECT * FROM T
UNION
SELECT * FROM S;
```

Set Difference ($\setminus$)

Returns tuples in one relation but not in another.

$$A \setminus B$$

```
SELECT * FROM T
EXCEPT
SELECT * FROM S;
-- or in some SQL dialects:
SELECT * FROM T
```

```
MINUS
SELECT * FROM S;
```

Intersection $\cap$

Returns tuples common to both relations.

$$A \cap B = A \setminus (A \setminus B)$$

```
SELECT * FROM T
INTERSECT
SELECT * FROM S;
```

Selection σ

Filters tuples based on a predicate.

$$\sigma_{\theta}(S) = \{s \in S \mid s \text{ satisfies } \Theta\}$$

```
SELECT * FROM T WHERE a > 10;
```

Projection π

Selects specific attributes.

$$\prod_{a_{i1}, \dots, a_{ik}} (S) \{(a_{i1}, \dots, a_{ik}) \mid a \in S\}$$

This gives us the same tuples of the original relational but only with the columns we selected.

It can also be noted as π .

```
SELECT a, b FROM T;
```

Symmetric Difference $\triangle$

Tuples in either relation but not in both.

$$A \triangle B = (A \setminus B) \cup (B \setminus A)$$

```
(SELECT * FROM T EXCEPT SELECT * FROM S)
UNION
(SELECT * FROM S EXCEPT SELECT * FROM T);
```

Rename ρ

Renames relations or attributes.

$$\rho_{A'(key, data)}(A)$$

```
SELECT a AS x, b AS y, c AS z FROM T;
-- or rename the table:
SELECT * FROM T AS T_new;
```

with

with allows us to store products of specific operation with a special name which can be used later in other operations.

Join $\bowtie$

A *theta join* is the Cartesian product plus a selection.

$$S \bowtie_{\theta} T = \sigma_{\theta}(S \times T)$$

Inner Join $\bowtie$

It is a *theta join* combines tuples that satisfy a condition.

$$S \bowtie_{\theta} T = \sigma_{\theta}(S \times T)$$

$$S \bowtie_{\theta} T = \sigma_{\theta}(S \times T)$$

```

SELECT * FROM T INNER JOIN S ON T.a = S.d;
-- or with theta condition:
SELECT * FROM T INNER JOIN S ON T.a < S.d;

```

Equi-Join

An *equi-join* is a join where the join condition consists only of equality comparisons between attributes of the two relations.

$$R \bowtie_{R.a_1=S.b_1 \wedge \dots \wedge R.a_k=S.b_k} S.$$

The name and domain of the attributes must be the same.

$$R \bowtie_{R.a_1=S.b_1 \wedge \dots \wedge R.a_k=S.b_k} S = \sigma_{R.a_1=S.b_1 \wedge \dots \wedge R.a_k=S.b_k}(R \times S)$$

```

SELECT * FROM T JOIN S ON T.a = S.d AND T.b = S.e;

```

Natural Join

It is a special case of the equi-join which can only be performed only if there is a common attribute.

$$R \bowtie S$$

Let $C = \text{ident}(R) \cap \text{ident}(S)$ be the common attributes.

$$R \bowtie S = \pi_{\text{ident}(R) \cup \text{ident}(S)} \left(\sigma_{\bigwedge_{c \in C} R.c=S.c} (R \times S) \right)$$

```

SELECT * FROM T NATURAL JOIN S;
-- assumes T and S have common column names

```

Semi-Join

It is a case where we only care about the attribute names but not its values.

$$R \ltimes S$$

$$R \ltimes S = \pi_{\text{ident}(R)}(R \bowtie S)$$

Or alternatively:

$$R \ltimes S = \pi_{\text{ident}(R)} \left(\sigma_{\bigwedge_{c \in C} R.c=S.c} (R \times S) \right)$$

where $C = \text{ident}(R) \cap \text{ident}(S)$.

```

SELECT T.* FROM T WHERE EXISTS
(SELECT 1 FROM S WHERE T.a = S.d);
-- or using IN:
SELECT * FROM T WHERE a IN (SELECT d FROM S);

```

Anti Semi-Join

In this case, it works very similar to the semi join, but we take the attributes which do not participate in the natural join.

$$R \triangleright S$$

$$R \triangleright S = R \setminus (R \bowtie S) = R \setminus \pi_{\text{ident}(R)}(R \bowtie S)$$

```

SELECT T.* FROM T WHERE NOT EXISTS
(SELECT 1 FROM S WHERE T.a = S.d);
-- or using NOT IN:
SELECT * FROM T WHERE a NOT IN (SELECT d FROM S);

```

Outer Join

Preserves non-matching tuples.

Left Outer Join: It returns all tuples from the left relation and matching tuples from the right relation.

Right Outer Join: It returns all tuples from the right relation and matching tuples from the left relation.

Full Outer Join: It returns all tuples from both relations, filling in NULLs for non-matching tuples.

```

-- Left Outer Join:
SELECT * FROM T LEFT OUTER JOIN S ON T.a = S.d;

-- Right Outer Join:
SELECT * FROM T RIGHT OUTER JOIN S ON T.a = S.d;

-- Full Outer Join:
SELECT * FROM T FULL OUTER JOIN S ON T.a = S.d;

```

Division $\div$

Used to express "for all" queries. It can only be performed when the relation on the right is a proper subset of the left relation.

Let

- $R(X, Y)$
- $S(Y)$

Then:

$$R \div S = \{x \mid \forall y \in S : (x, y) \in R\}$$

$$R \div S = \pi_X(R) \setminus \pi_X((\pi_X(R) \times S) \setminus R),$$

where $X = \text{ident}(R) \setminus \text{ident}(S)$.

Example Concept

If $A(\text{id}, \text{value})$ contains all values in $B(\text{value})$, division returns the corresponding ID's.

```

-- Find tuples in T that match all tuples in S
SELECT DISTINCT T.a FROM T
WHERE NOT EXISTS (
  SELECT S.d FROM S
  WHERE NOT EXISTS (
    SELECT * FROM T AS T2
    WHERE T2.a = T.a AND T2.b = S.d
  )
);

```

Assignment (=)

This is just using to store the result of a relational algebra operation.

```

-- Create a view or temporary table
CREATE VIEW Result AS SELECT * FROM T WHERE a > 5;
-- or
CREATE TEMPORARY TABLE Result AS SELECT * FROM T WHERE a > 5;

```

27.0.5 Types of Keys

A *key* is an attribute or combination of attributes (composite key) which uniquely identifies an entity.

- *Candidate Key*: The term refers to the attributes which are suited for being the key.
- *Irreducible*: A key has to be as atomic as possible.
- *Super key*: It consists of a candidates key's attributes + potentially some extra attributes.
- *Alternate Key*: This is an alternative primary key. This means that the UNIQUE constraints will be imposed on the attribute/s.
- *Surrogate Key*: This key has only meaning inside the database. Example: ID $\rightarrow$ Person's Name.
- *Foreign Key*: This key is an attribute of one table which references another table.

27.0.6 Relation Schema

The $(\mathcal{R}, \tau)$ represents a *relation schema*. We can constrain our relations to obey certain rules. Thus, limiting our space. Here, $\mathcal{R}$ is the set of all relations of n elements and τ is the set of n -tuple (attributes).

The rules we use to limit the space of allowed relations are called *restrictions*.

This is the analog of using CREATE TABLE in SQL. We are creating rules for the tuples inside the relation.

Also, this is just the definition of the name of the relations and the types of its attributes.

27.0.7 Constraint

We denote that $d[x]$ is a function from $x \rightarrow R^+$, where R^+ is the relation R without the attribute x . This is called the *constraint*.

27.0.8 Functional Dependencies

A *functional relation* is dependence between one or more attributes commonly noted $A \rightarrow B$.

Formally, given $\alpha \subseteq \text{ident}(R)$ and $\beta \subseteq \text{ident}(R)$ and $D \subseteq \text{dom}(R)$ $\alpha \rightarrow \beta$, β is dependent on α if

$$\forall d_1, d_2 \in D : d_1[\alpha] = d_2[\alpha] \implies d_1[\beta] = d_2[\beta].$$

In simple terms the notation $X \rightarrow Y$ means "we know Y if we know X ".

A complete dependence is when $A \rightarrow B$ is when we can not further reduce the tuple A so that we still get B .

27.0.9 Armstrong Axioms

For a relation $R = \{a_1, \dots, a_n\}$ with the schema $\alpha, \beta, \gamma \subseteq \text{ident}(R)$.

The Armstrong axioms are given:

- I. *Reflexivity*: for all $\beta \subseteq \alpha$ holds $\alpha \rightarrow \beta$
- II. *Transitivity*: $\alpha \rightarrow \beta \wedge \beta \rightarrow \gamma \implies \alpha \rightarrow \gamma$
- III. *Connectivity*: $\alpha \rightarrow \beta \implies \alpha\gamma \rightarrow \beta\gamma$
- IV. *Union*: $\alpha \rightarrow \beta \wedge \alpha \rightarrow \gamma \implies \alpha \rightarrow \beta\gamma$
- V. *Decomposition*: $\alpha \rightarrow \beta\gamma \implies \alpha \rightarrow \beta \wedge \alpha \rightarrow \gamma$
- VI. *Pseudotransitivity*: $\alpha \rightarrow \beta \wedge \beta\gamma \rightarrow \delta \implies \alpha\gamma \rightarrow \delta$

These are not strictly axioms, but they are called that way.

27.0.10 Attribute Closure

The *attribute closure* $AC(FD, x)$ is the set of all attributes reachable via functional dependency including the starting point.

Example:

$R = \{A, B, C, D\}$ with $FD = \{A \rightarrow B, B \rightarrow C, AB \rightarrow CD\}$.

- $AC(FD, A) = \{A, AB, ABC, ABCD\} = ABCD$
- $AC(FD, B) = \{B, BC\} = BC$

27.0.11 Canonical Cover

We consider two sets of functional dependencies if and only if their attribute closures FD^+ are identical.

For a schema R with a set of functional dependencies FD we denote FC^c as *the canonical cover* of FD if the following criteria are satisfied:

- $FD^c \cong FD, FD^{c+} = FD^+$
- For all $\alpha \rightarrow \beta \in FD^c$ are minimal in the sense that
 - $\forall a \in \alpha : FD^c \setminus (\alpha \rightarrow \beta) \cup (\alpha \setminus a \rightarrow \beta) \neq FD^c$
 - $\forall b \in \beta : FD^c \setminus (\alpha \rightarrow \beta) \cup (\alpha \rightarrow \beta \setminus b) \neq FD^c$
- All left side attributes appear only one time

The following steps are always performed for the whole set of functional dependencies one after the other.

Left-Reduction

For all $\alpha \rightarrow \beta \in FD$ check if there is a redundant $a \in \alpha$. $\beta \subseteq AC(FD, \alpha \setminus a)$. If that is the case then replace $\alpha \rightarrow \beta$ with $\alpha \setminus a \rightarrow \beta$.

After that

Right-Reduction

For all $\alpha \rightarrow \beta \in FD$ check if there is a redundant $b \in \beta$. $\beta \subseteq AC(FD \setminus \{\alpha \rightarrow \beta\} \cup \{\alpha \rightarrow \beta \setminus b, \alpha\})$. If that is the case then replace $\alpha \rightarrow \beta$ with $\alpha \rightarrow \beta \setminus b$.

Then

Elimination

Remove $\alpha \rightarrow \emptyset$.

And finally

Union

If $\alpha \rightarrow \beta_1$ and $\alpha \rightarrow \beta_2$ then combine them into $\alpha \rightarrow \beta_1, \beta_2$.

Example:

Given are $R = \{A, B, C, D, E, F\}$, $FD = \{ABC \rightarrow DEF, A \rightarrow BC, D \rightarrow EF\}$.

Left-Reduction:

First iteration, $ABC \rightarrow DEF$

- Is A redundant? No: $AC(F, BC) = BC$
- Is B redundant? Yes: $AC(F, AC) = AC \implies ABC \implies ABCDEF$. Thus, we have $AC \rightarrow DEF, A \rightarrow BC, D \rightarrow EF$.
- Is C redundant? Yes: $AC(F, A) = A \implies ABC \implies ABCDEF$. Thus, we have $A \rightarrow DEF, A \rightarrow BC, D \rightarrow EF$.

Right-Reduction

Using $\{A \rightarrow DEF, A \rightarrow BC, D \rightarrow EF\}$ we proceed with the first dependency $A \rightarrow DEF$.

- Is D redundant? No, without it, we can not reach EF .
- Is E redundant? Yes: $AC(A \rightarrow DF, A \rightarrow BC, A \rightarrow EF, A) = A \implies ADF \implies ADEFBC$. Thus, we have $F = \{A \rightarrow DF, A \rightarrow BC, D \rightarrow EF\}$.
- Is F redundant? Yes: $AC(A \rightarrow D, A \rightarrow BC, D \rightarrow EF, A) = A \implies ABC \implies ABCDEF$. Thus, we have $F = \{A \rightarrow D, A \rightarrow BC, D \rightarrow EF\}$.

Elimination and Union

$$FC^c = \{A \rightarrow BCD, D \rightarrow EF\}$$

Important Notes

- By the left-reduction we only care about fd's which have at least two attributes on the right side.
- By the right-reduction we only care about the fd's with one attribute at the right.

27.0.12 How to find a Candidate Key

Given a set of attributes $\{A, B, C, D\}$ with the following functional dependencies:

$$A \rightarrow B, C \rightarrow B, A \rightarrow D$$

The set is already left-reduced.

- $AC(F, A) = ABD$
- $AC(F, C) = CB$

We get $A \rightarrow BD, C \rightarrow B$.

For the right reduction:

- Is B redundant? Yes

Finally, we get our candidate key $AC \rightarrow ABCD$.

27.0.13 Losslessness and Maintenance of Dependency

Given a relation which needs to be split into more relations:

- *Losslessness:* After the split, the information in the original relation needs to be accessible via joins.
- *Maintenance of Dependency:* The original functional dependencies do not change after the split.

27.0.14 Normal Forms

A *normalized* table are protected from data inconsistencies and make the tables more readable.

I Normal Form

- No row order for information.
- No Mixing data types in the same column.
- No Tables without a primary key.
- No Repeating groups.

This normal is mostly given in relational algebra problems.

II Normal Form:

Each non-key attribute must depend on the entire primary key.

The relation is in the second normal form if it is already in the first normal form and each **non-key** attribute is functionally dependent on the **key-candidate**.

Example I:

$R = \{A, B, C, D\}$, $FD = \{A \rightarrow BC, B \rightarrow A\}$ not in 2NF due to D not being present.

Example II:

$R = \{A, B, C, D\}$, $FD = \{A \rightarrow B, B \rightarrow A, AD \rightarrow C\}$

Candidates AD, BD , not prime attribute C . Thus, it obeys the 2NF.

III Normal Form

A schema R is in the 3NF if for all $\alpha \rightarrow \beta \in FD$ at least one of the following statements is true:

- $\alpha \rightarrow \beta$ is trivial $\beta \subseteq \alpha$
- β has only prime attributes
- α is a superkey

The relation is in the third normal form if it is already in the second normal form and each *non-key* attribute is not dependent on other *non-key* attribute.

Boyce-Codd III Normal Form

A schema R is in the BCNF if for all $\alpha \rightarrow \beta \in FD$ at least one of the following statements is true:

- $\alpha \rightarrow \beta$ is trivial $\beta \subseteq \alpha$
- α is a superkey

Every attribute in a table should depend on the key, the whole key, and nothing but the key.

Example I

$R = \{A, B, C, D\}$, $FD = \{A \rightarrow BC, C \rightarrow D\}$, not in 3NF due to D being dependent on C .

IV Normal Form

Multivalued dependencies in a table must be multivalued dependencies on the key.

V Normal Form

Our table in the forth normal form cannot be describable as the logical result of joining some other tables.

27.0.15 Synthesis Algorithm

The *synthesis algorithm* transforms a relation schema into a set of schemas that are in Third Normal Form (3NF), while ensuring *lossless join* and *dependency preservation*. The algorithm proceeds as follows:

- Compute the *canonical cover* F_c of the given set of functional dependencies F :
 - Decompose all dependencies so that each has a single attribute on the right-hand side.
 - Remove extraneous attributes from the left-hand sides.
 - Eliminate redundant dependencies.
- For each functional dependency $\alpha \rightarrow \beta \in F_c$, construct a relation schema

$$R_k = \alpha \cup \beta.$$

All dependencies whose attributes are contained in R_k are associated with this schema.

- If none of the generated schemas contains a candidate key of the original relation, create an additional schema whose attributes form a candidate key.
- If a schema R_k is completely contained in another schema R_m , remove R_k , since it is redundant.

Example:

Given

$$R = \{A, B, C, D, E\}$$

with

$$F = \{ADE \rightarrow BE, B \rightarrow ACD, D \rightarrow AC, ACE \rightarrow BD, C \rightarrow B\}.$$

Step 1: Right-hand side decomposition

$$\begin{aligned} &ADE \rightarrow B, \quad ADE \rightarrow E, \quad B \rightarrow A, \quad B \rightarrow C, \quad B \rightarrow D, \\ &D \rightarrow A, \quad D \rightarrow C, \quad ACE \rightarrow B, \quad ACE \rightarrow D, \quad C \rightarrow B \end{aligned}$$

Step 2: Left reduction

- $ADE \rightarrow B$: A and E are extraneous since $D^+ = \{D, B, A, C, E\}$, hence $D \rightarrow B$.
- $ADE \rightarrow E$ is trivial and removed.
- $ACE \rightarrow B$: A and E are extraneous, yielding $C \rightarrow B$.
- $ACE \rightarrow D$: A and E are extraneous, yielding $C \rightarrow D$.
- All other dependencies already have minimal left-hand sides.

Current set:

$$\{D \rightarrow B, B \rightarrow A, B \rightarrow C, B \rightarrow D, D \rightarrow A, D \rightarrow C, C \rightarrow B, C \rightarrow D\}$$

Step 3: Remove redundant dependencies

- $D \rightarrow A$ is redundant since $D \rightarrow C$ and $C \rightarrow B$ and $B \rightarrow A$.
- $B \rightarrow C$ is redundant since $B \rightarrow D$ and $D \rightarrow C$.
- $C \rightarrow D$ is redundant since $C \rightarrow B$ and $B \rightarrow D$.

Thus, the canonical cover is:

$$F_c = \{B \rightarrow A, B \rightarrow D, D \rightarrow C, C \rightarrow B\}$$

Step 4: Construct relation schemas

$$\begin{aligned} R_1 &= \{B, A\}, \\ R_2 &= \{B, D\}, \\ R_3 &= \{D, C\}, \\ R_4 &= \{C, B\} \end{aligned}$$

Step 5: Remove redundant schemas

None of these schemas is contained in another, so all are kept.

Step 6: Test for candidate keys

For the original relation $R = \{A, B, C, D, E\}$, attribute E does not appear on any right-hand side and must be part of every key.

$$(BE)^+ = \{A, B, C, D, E\}$$

so $\{B, E\}$ is a candidate key.

Since no schema contains $\{B, E\}$, we add:

$$R_5 = \{B, E\}$$

Final 3NF Decomposition:

$$\{\{B, A\}, \{B, D\}, \{D, C\}, \{C, B\}, \{B, E\}\}$$

27.0.16 Decomposition Algorithm

The *decomposition algorithm* transforms a relation schema into Boyce-Codd Normal Form (BCNF) by repeatedly decomposing relations that violate the BCNF condition. A relation schema R is in BCNF if for every non-trivial functional dependency $\alpha \rightarrow \beta$ on R , α is a superkey of R .

- If R is already in BCNF, stop
- Otherwise, choose a functional dependency $\alpha \rightarrow \beta$ that violates BCNF, meaning α is not a superkey of R
- Decompose R into two schemas

$$R_1 = \alpha \cup \beta \quad \text{and} \quad R_2 = R \setminus (\beta \setminus \alpha)$$

- Apply the algorithm recursively to R_1 and R_2 until all resulting schemas are in BCNF

Example:

Given

$$R = \{A, B, C, D\},$$

with

$$F = \{A \rightarrow B, B \rightarrow C, C \rightarrow D\}.$$

Step 1: Check BCNF

$A \rightarrow B$ violates BCNF since A is not a superkey of R

Step 2: First decomposition

Using $A \rightarrow B$, decompose into

$$R_1 = \{A, B\} \quad \text{and} \quad R_2 = \{A, C, D\}$$

Step 3: Check BCNF of R_1

In $R_1(A, B)$, $A \rightarrow B$ holds and A is a key, so R_1 is in BCNF

Step 4: Check BCNF of R_2

From $B \rightarrow C$ and $C \rightarrow D$, we obtain $A \rightarrow C$ and $A \rightarrow D$, but A is not a key of R_2 , so BCNF is violated by $A \rightarrow C$

Step 5: Second decomposition

Using $A \rightarrow C$, decompose R_2 into

$$R_3 = \{A, C\} \quad \text{and} \quad R_4 = \{A, D\}$$

Step 6: Check BCNF of remaining schemas

In $R_3(A, C)$, A is a key, so R_3 is in BCNF

In $R_4(A, D)$, $A \rightarrow D$ holds and A is a key, so R_4 is in BCNF

Final BCNF Decomposition:

$$\{\{A, B\}, \{A, C\}, \{A, D\}\}$$

27.0.17 Optimization

The following equivalences hold for relational algebra expressions:

- $\sigma_{C_1 \wedge C_2 \wedge \dots \wedge C_n} \cong \sigma_{C_1}(\sigma_{C_2}(\dots(\sigma_{C_n}(R))\dots))$
- $\sigma_{C_1}(\sigma_{C_2}(R)) \cong \sigma_{C_1}(\sigma_{C_2}(R))$
- If $L_1 \subseteq L_2 \subseteq \dots \subseteq L_n$ then

$$\pi_{L_1}(\pi_{L_2}(\dots(\pi_{L_n}(R))\dots)) \cong \pi_{L_1}(R)$$

- $\pi_{A_1, \dots, A_n}(\sigma_C(R)) \cong \sigma_C(\pi_{A_1, \dots, A_n}(R))$
- $\times, \cup, \cap$ and $\bowtie$ are commutative
- Pushing Selections, select as early as possible $\sigma_C(R \bowtie_{C_J} S) \cong \sigma_C(R) \bowtie_{C_J} S$

Special Case:

$$\sigma_{C_1 \wedge C_2}(R \bowtie_{C_J} S) \cong \sigma_{C_1}(R) \bowtie_{C_J} \sigma_{C_2}(S)$$

- $\pi_L(R \bowtie_{C_J} S) \cong \pi_{A_1, \dots, A_n}(R) \bowtie_{C_J} \pi_{b_1, \dots, b_n}(S)$
- $\times, \cup, \cap$ and $\bowtie$ are associative

- σ is distributive with $\cup, \cap, \setminus$

$$\sigma_C(R\Theta S) = \sigma_C(R)\Theta\sigma_C(S)$$

- π is distributive with $\cup$

$$\pi_C(R \cup S) = \pi_C(R) \cup \pi_C(S)$$

- De Morgam for Joins with selections

$$\neg(A \wedge B) = \neg A \vee \neg B \quad \neg(A \vee B) = \neg A \wedge \neg B$$

- $\sigma_C(R \times S) \cong R \bowtie_C S$

Chapter 28

Geometry

28.0.1 Formulas for Area and Perimeter of Geometric Figures

- **Square**

- Area: $A = a^2$
- Perimeter: $P = 4a$

- **Rectangle**

- Area: $A = l \cdot w$
- Perimeter: $P = 2(l + w)$

- **Circle**

- Area: $A = \pi r^2$
- Circumference: $C = 2\pi r$

- **Triangle (General)**

- Area: $A = \frac{1}{2}b \cdot h$
- Perimeter: $P = a + b + c$

- **Equilateral Triangle**

- Area: $A = \frac{\sqrt{3}}{4}a^2$
- Perimeter: $P = 3a$

- **Isosceles Triangle**

- Area: $A = \frac{b}{4}\sqrt{4a^2 - b^2}$

- **Trapezoid**

- Area: $A = \frac{1}{2}(a + b)h$
- Perimeter: $P = a + b + c + d$

- **Parallelogram**

- Area: $A = b \cdot h$
- Perimeter: $P = 2(a + b)$

28.0.2 Formulas for Area and Volume of 3D Geometric Figures

- **Cube**


- Surface Area: $A = 6a^2$

- Volume: $V = a^3$
- **Cylinder**
 - Surface Area: $A = 2\pi r(h + r)$
 - Volume: $V = \pi r^2 h$
- **Cone**
 - Surface Area: $A = \pi r(r + l)$ with l = slant height
 - Volume: $V = \frac{1}{3}\pi r^2 h$
- **Sphere**
 - Surface Area: $A = 4\pi r^2$
 - Volume: $V = \frac{4}{3}\pi r^3$
- **Square Pyramid**
 - Surface Area: $A = a^2 + 2a \cdot l$
 - Volume: $V = \frac{1}{3}a^2 h$
- **Triangular Pyramid (Tetrahedron)**
 - Volume: $V = \frac{1}{3}A_b \cdot h$
- **Prism (General)**
 - Surface Area: $A = 2A_b + P_b \cdot h$
 - Volume: $V = A_b \cdot h$

28.0.3 Thales' Theorem

Thales' Theorem states: If A , B , and C are points on a circle where the line AB is the diameter, then the angle $\angle\alpha + \beta$ is a right angle.

$$\angle\alpha + \beta = 90^\circ \quad \text{if } AB \text{ is a diameter of the circle}$$



Proof:

Note that we have two Isosceles triangles where the sides AO , AC and OC have all the same length, being this the radius of the circle.

Because they are Isosceles then two of their angles have to be equal and the sum of the angles of this two triangle has to add up to be 180° . Therefore,

$$\begin{aligned}\alpha + \alpha + \beta + \beta &= 180 \\ 2\alpha + 2\beta &= 180 \\ \alpha + \beta &= 90\end{aligned}$$

□◻◻

28.0.4 Conversion Between Radians and Degrees


To convert between radians and degrees, use the following formulas:

$$\begin{aligned}\text{Degrees} &= \text{Radians} \times \left(\frac{180^\circ}{\pi} \right) \\ \text{Radians} &= \text{Degrees} \times \left(\frac{\pi}{180^\circ} \right)\end{aligned}$$


28.0.5 Geometric Figures with Formulas (2D)

Square and Rectangle


$$A = a^2 \quad P = 4a$$



$$A = l \cdot w \quad P = 2(l + w)$$




Circle and Triangle Types




Circle

$$A = \pi r^2 \quad C = 2\pi r$$



Equilateral


$$A = \frac{\sqrt{3}}{4} a^2$$



Isosceles


$$A = \frac{b}{4} \sqrt{4a^2 - b^2}$$

Scalene Triangle, Trapezoid, Parallelogram




Scalene Triangle

$$A = \frac{1}{2}b \cdot h$$



Trapezoid

$$A = \frac{1}{2}(a + b)h$$




Parallelogram

$$A = b \cdot h \quad P = 2(a + b)$$


28.0.6 3D Geometric Figures with Formulas

Cube and Cylinder



Cube


$$A = 6a^2 \quad V = a^3$$



Cylinder


$$A = 2\pi r(h + r) \\ V = \pi r^2 h$$

Cone and Sphere



Cone


$$A = \pi r(r + l) \\ V = \frac{1}{3}\pi r^2 h$$



Sphere

$$A = 4\pi r^2 \quad V = \frac{4}{3}\pi r^3$$


Pyramid and Prism



Square Pyramid

$$A = a^2 + 2al$$

$$V = \frac{1}{3}a^2h$$



Triangular Prism

$$A = 2A_b + P_b h$$

$$V = A_b \cdot h$$

28.0.7 Axioms of Euclidean Geometry

Euclidean geometry is founded on a set of basic assumptions known as axioms or postulates, originally formulated by the Greek mathematician Euclid in his work *Elements*. These axioms form the logical basis for plane geometry and include the following:

- I. *Line Postulate*: A straight line segment can be drawn joining any two points.
- II. *Extension Postulate*: Any straight line segment can be extended indefinitely in a straight line.
- III. *Circle Postulate*: Given any straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
- IV. *Right Angle Postulate*: All right angles are congruent.
- V. *Parallel Postulate*: If a line segment intersects two straight lines and makes the interior angles on the same side less than two right angles, then the two lines, if extended indefinitely, meet on that side.

These five postulates define the framework of classical geometry in a flat (Euclidean) space. Notably, the fifth postulate—the parallel postulate—has been the subject of much scrutiny and led to the development of non-Euclidean geometries when it was altered or replaced.

28.0.8 Types of Angles and Angle Relationships

An angle is the distance between two rays or intercepting lines.

- If angle is less than 90° it is an *acute* angle
- If it is exactly 90° it is a *right* angle
- It is more than 90° but less than 180° it is called an *obtuse* angle

Now let us look at other types of angles

Vertical Angles

When two lines intersect, they form two pairs of opposite (or vertical) angles. Vertical angles are always congruent. That is, if two angles are vertical, their measures are equal:

$$\angle A = \angle B$$

For example, if two lines intersect and form angles of 60° and 120° , the angles opposite each other are both 60° and both 120° .

Adjacent Angles

Adjacent angles are two angles that share a common side and a common vertex, and do not overlap. When two lines intersect, each pair of adjacent angles forms a straight angle (or a linear pair), summing to 180° :

$$\angle 1 + \angle 2 = 180^\circ$$

This is known as the Linear Pair Postulate.

Complementary Angles

Complementary angles are two angles whose measures add up to 90° :

$$\angle X + \angle Y = 90^\circ$$

Complementary angles do not have to be adjacent, but if two intersecting lines form a right angle, the two acute angles adjacent to the right angle may be complementary.

Angles Generated by Traversals

When a transversal crosses two or more lines, it forms several pairs of angles with specific relationships. Some key types include:

- *Corresponding Angles*: Angles in the same relative position at each intersection. If the lines are parallel, corresponding angles are congruent:

$$\angle 1 \cong \angle 2$$

- *Alternate Interior Angles*: Angles that lie between the two lines but on opposite sides of the transversal. They are congruent if the lines are parallel:

$$\angle 3 \cong \angle 4$$

- *Alternate Exterior Angles*: Angles outside the two lines and on opposite sides of the transversal. Also, congruent when the lines are parallel:

$$\angle 5 \cong \angle 6$$

- *Consecutive (Same-Side) Interior Angles*: Angles on the same side of the transversal and inside the parallel lines. These are supplementary:

$$\angle 7 + \angle 8 = 180^\circ$$

28.0.9 Triangle Similarity and Congruence

Triangles can be compared using criteria of similarity or congruence, depending on whether their shapes are the same or both shape and size are identical.

Triangle Congruence

Two triangles are *congruent* if all corresponding sides and angles are equal. This means the triangles are identical in shape and size. Common criteria for triangle congruence include:

- *SSS (Side-Side-Side)*: All three pairs of corresponding sides are equal.
- *SAS (Side-Angle-Side)*: Two pairs of sides and the angle between them are equal.
- *ASA (Angle-Side-Angle)*: Two pairs of angles and the included side are equal.

- *AAS (Angle-Angle-Side)*: Two angles and a non-included side are equal.
- *HL (Hypotenuse-Leg)*: For right triangles, if the hypotenuse and one leg are equal.

Triangle Similarity


Two triangles are *similar* if their corresponding angles are equal and their corresponding sides are in proportion. This means they have the same shape but not necessarily the same size. Criteria for similarity include:

- *AA (Angle-Angle)*: Two pairs of corresponding angles are equal.
- *SAS (Side-Angle-Side)*: One pair of corresponding angles is equal, and the sides including those angles are in proportion.
- *SSS (Side-Side-Side)*: All three pairs of corresponding sides are in proportion.

28.0.10 Visualizations of Triangle Similarity and Congruence


SSS (Side-Side-Side)

A triangle is uniquely determined if all three sides are known.




SAS (Side-Angle-Side)

A triangle is uniquely determined if two sides and the included angle are known.




ASA (Angle-Side-Angle)

A triangle is uniquely determined if two angles and the included side are known.



AAS (Angle-Angle-Side)


A triangle is uniquely determined if two angles and a non-included side are known.



SSA (Side-Side-Angle)

Two sides and a non-included angle do *not always* determine a unique triangle. This case is ambiguous and can result in:

- One triangle (when the opposite side is long enough),
- Two triangles (ambiguous case),
- No triangle (when the side is too short).



28.0.11 Triangle Medians, Angle Bisectors, and Altitudes

In any triangle, there are special line segments that connect vertices to specific points on the opposite sides. These segments play key roles in triangle geometry and have unique properties.

Median

A *median* of a triangle is a line segment that connects a vertex to the midpoint of the opposite side. Every triangle has three medians, and they intersect at a single point called the *centroid*, which is the triangle's center of mass. The centroid divides each median in a ratio of 2 : 1, with the longer part closer to the vertex.

Angle Bisector

An *angle bisector* is a line segment that divides an angle of the triangle into two equal parts and extends to the opposite side. All three angle bisectors intersect at a point called the *incenter*, which is the center of the triangle's incircle (the largest circle that fits inside the triangle and touches all three sides).

Altitude

An *altitude* of a triangle is a perpendicular segment from a vertex to the line containing the opposite side (not necessarily within the triangle). Altitudes intersect at a point called the *orthocenter*. Unlike medians and bisectors, altitudes can fall outside the triangle in obtuse triangles.

Centroid $\rightarrow$ Intersection of medians

Incenter $\rightarrow$ Intersection of angle bisectors

Orthocenter $\rightarrow$ Intersection of altitudes

28.0.12 The Midsegment Theorem

The *Midsegment Theorem* states that the segment connecting the midpoints of two sides of a triangle is:

- *Parallel* to the third side of the triangle
- *Half the length* of the third side

Let $\triangle ABC$ be a triangle, and let D and E be the midpoints of sides AB and AC , respectively. Then the segment $\overline{DE}$ is the *midsegment*, and we have:

$$\overline{DE} \parallel \overline{BC} \quad \text{and} \quad DE = \frac{1}{2}BC$$

This theorem is useful in coordinate geometry, triangle proofs, and for simplifying complex geometric relationships within triangles.

28.0.13 Intercept Theorems


The intercept theorems describe relationships between segment lengths when two rays from a point intersect two parallel lines. They are based on similar triangles and allow us to calculate unknown lengths using proportions.

First Intercept Theorem

If two rays start from a common point and are intersected by two parallel lines, then:

$$\frac{a}{a'} = \frac{b}{b'},$$


where a and b are segments on one ray, and a' , b' are the corresponding segments on the other ray.



Second Intercept Theorem (General Form)

If a ray intersects two parallel lines, the segments from the origin point to the lines are in the same ratio as the segments along the parallels:

$$\frac{SA}{SA'} = \frac{SB}{SB'} \quad \text{and} \quad \frac{AB}{A'B'} = \frac{SA}{SA'}$$



28.0.14 Proof of the Pythagorean Theorem

We want to prove that $c^2 = a^2 + b^2$ is a true statement.

Let us build a Square with area c^2 and take our original triangle and put it on the sides on the square.

This will create a square of sides $(a + b)^2$ which is bigger than the inner square c^2 .

Now let us find the area c^2 using what we know about this shape.

$$\begin{aligned} (a + b)^2 &= 4\left(\frac{ab}{2}\right) + c^2 \\ (a + b)^2 &= 2ab + c^2 \\ a^2 + 2ab + b^2 &= 2ab + c^2 \\ a^2 + b^2 &= c^2 \end{aligned}$$

QED

Chapter 29

Trigonometry

In this section, we will cover a variety of trigonometric functions and their properties.

29.0.1 Trigonometric Functions

- $\sin(x) = \frac{1}{\csc(x)}$
- $\cos(x) = \frac{1}{\sec(x)}$
- $\tan(x) = \frac{\sin(x)}{\cos(x)} = \frac{1}{\cot(x)}$
- $\csc(x) = \frac{1}{\sin(x)}$
- $\sec(x) = \frac{1}{\cos(x)}$
- $\cot(x) = \frac{\cos(x)}{\sin(x)} = \frac{1}{\tan(x)}$

29.0.2 SOH-CAH-TOA


- $\sin(x) = \frac{\text{opposite}}{\text{hypotenuse}}$
- $\cos(x) = \frac{\text{adjacent}}{\text{hypotenuse}}$
- $\tan(x) = \frac{\text{opposite}}{\text{adjacent}}$

29.0.3 Pythagorean Identities

- $\sin^2(x) + \cos^2(x) = 1$
- $1 + \tan^2(x) = \sec^2(x)$
- $1 + \cot^2(x) = \csc^2(x)$

29.0.4 Sum and Difference Formulas

- $\sin(a + b) = \sin(a) \cos(b) + \cos(a) \sin(b)$
- $\sin(a - b) = \sin(a) \cos(b) - \cos(a) \sin(b)$
- $\cos(a + b) = \cos(a) \cos(b) - \sin(a) \sin(b)$
- $\cos(a - b) = \cos(a) \cos(b) + \sin(a) \sin(b)$
- $\tan(a + b) = \frac{\tan(a) + \tan(b)}{1 - \tan(a) \tan(b)}$
- $\tan(a - b) = \frac{\tan(a) - \tan(b)}{1 + \tan(a) \tan(b)}$



$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

29.0.5 Double Angle Formulas

- $\sin(2x) = 2 \sin(x) \cos(x)$
- $\cos(2x) = \cos^2(x) - \sin^2(x) = 2 \cos^2(x) - 1 = 1 - 2 \sin^2(x)$
- $\tan(2x) = \frac{2 \tan(x)}{1 - \tan^2(x)}$
- $\csc(2x) = \frac{2 \csc(x) \sec(x)}{1 - \tan^2(x)}$
- $\sec(2x) = \frac{1 + \tan^2(x)}{2 \tan(x)}$
- $\cot(2x) = \frac{1 - \tan^2(x)}{2 \tan(x)}$

29.0.6 Half Angle Formulas

- $\sin\left(\frac{x}{2}\right) = \pm \sqrt{\frac{1 - \cos(x)}{2}}$
- $\cos\left(\frac{x}{2}\right) = \pm \sqrt{\frac{1 + \cos(x)}{2}}$
- $\tan\left(\frac{x}{2}\right) = \frac{\sin(x)}{1 + \cos(x)} = \frac{1 - \cos(x)}{\sin(x)} = \frac{\tan(x)}{1 + \tan^2(x)}$
- $\csc\left(\frac{x}{2}\right) = \frac{1}{\sin\left(\frac{x}{2}\right)} = \pm \sqrt{\frac{2}{1 - \cos(x)}}$
- $\sec\left(\frac{x}{2}\right) = \frac{1}{\cos\left(\frac{x}{2}\right)} = \pm \sqrt{\frac{2}{1 + \cos(x)}}$
- $\cot\left(\frac{x}{2}\right) = \frac{1}{\tan\left(\frac{x}{2}\right)} = \frac{1 + \cos(x)}{\sin(x)} = \frac{1 - \tan^2(x)}{2 \tan(x)}$

29.0.7 Product to Sum Formulas

- $\sin(a) \sin(b) = \frac{1}{2}[\cos(a - b) - \cos(a + b)]$
- $\cos(a) \cos(b) = \frac{1}{2}[\cos(a - b) + \cos(a + b)]$
- $\sin(a) \cos(b) = \frac{1}{2}[\sin(a + b) + \sin(a - b)]$
- $\cos(a) \sin(b) = \frac{1}{2}[\sin(a + b) - \sin(a - b)]$

29.0.8 Power Reducing Formulas

- $\sin^2(x) = \frac{1 - \cos(2x)}{2}$
- $\cos^2(x) = \frac{1 + \cos(2x)}{2}$
- $\tan^2(x) = \frac{1 - \cos(2x)}{1 + \cos(2x)}$
- $\csc^2(x) = \frac{1}{\sin^2(x)} = \frac{2}{1 - \cos(2x)}$
- $\sec^2(x) = \frac{1}{\cos^2(x)} = \frac{2}{1 + \cos(2x)}$
- $\cot^2(x) = \frac{1}{\tan^2(x)} = \frac{1 + \cos(2x)}{1 - \cos(2x)}$

29.0.9 Even and Odd Functions

- $\sin(-x) = -\sin(x)$
- $\cos(-x) = \cos(x)$
- $\tan(-x) = -\tan(x)$
- $\csc(-x) = -\csc(x)$
- $\sec(-x) = \sec(x)$
- $\cot(-x) = -\cot(x)$

29.0.10 Graph Identity


$$\cos(x) = \sin\left(x + \frac{\pi}{2}\right)$$

29.0.11 Trigonometric Values

θ (radians)	$\sin(\theta)$	$\cos(\theta)$	$\tan(\theta)$
0	0	1	0
$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$\frac{\pi}{2}$	1	0	undefined
$\frac{2\pi}{3}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$-\sqrt{3}$
$\frac{3\pi}{4}$	$\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	-1
$\frac{5\pi}{6}$	$\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{\sqrt{3}}$
π	0	-1	0
$\frac{7\pi}{6}$	$-\frac{1}{2}$	$-\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$\frac{5\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2}$	1
$\frac{4\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$	$\sqrt{3}$
$\frac{3\pi}{2}$	-1	0	undefined
$\frac{5\pi}{3}$	$-\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$-\sqrt{3}$
$\frac{7\pi}{4}$	$-\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	-1
$\frac{11\pi}{6}$	$-\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$-\frac{1}{\sqrt{3}}$
2π	0	1	0

29.0.12 Unit Circle and Trigonometric Functions

The unit circle is a circle with a radius of 1 centered at the origin. Trigonometric functions like $\sin(\theta)$, $\cos(\theta)$, and $\tan(\theta)$ can be defined based on the coordinates of a point on the circle corresponding to angle θ .



29.0.13 Definitions on the Unit Circle:

- $\cos(\theta)$ is the *x-coordinate* of the point on the circle.
- $\sin(\theta)$ is the *y-coordinate*.

- $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$ is the ratio of the opposite side to the adjacent side in the triangle.

29.0.14 Trigonometric Rules

- $\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$
- $c^2 = a^2 + b^2 - 2ab \cos \gamma$
- $\frac{a-b}{a+b} = \frac{\tan \frac{1}{2}(\alpha-\beta)}{\tan \frac{1}{2}(\alpha+\beta)}$

29.0.15 Proof of the Cosine Rule

The *Cosine Rule* relates the lengths of the sides of a triangle to the cosine of one of its angles. For any triangle ABC with sides $a = BC$, $b = AC$, and $c = AB$, and angle $\gamma = \angle ACB$, the cosine rule states:

$$c^2 = a^2 + b^2 - 2ab \cos(\gamma)$$

Proof:

Consider triangle ABC , and place it on the coordinate plane such that point c is at the origin $(0, 0)$, point B is at $(a, 0)$, and point A lies somewhere in the plane. Let angle $\gamma = \angle ACB$.

Let the coordinates of point A be $(b \cos \gamma, b \sin \gamma)$, since $b = AC$ and we're using polar coordinates to define its location relative to point c .

Using the distance formula to find side $c = AB$, we get:

$$\begin{aligned} c^2 &= (a - b \cos \gamma)^2 + (0 - b \sin \gamma)^2 \\ &= a^2 - 2ab \cos \gamma + b^2 \cos^2 \gamma + b^2 \sin^2 \gamma \\ &= a^2 + b^2 (\cos^2 \gamma + \sin^2 \gamma) - 2ab \cos \gamma \\ &= a^2 + b^2 - 2ab \cos \gamma \end{aligned}$$

Since $\cos^2 \gamma + \sin^2 \gamma = 1$.

Thus, we have proven that:

$$c^2 = a^2 + b^2 - 2ab \cos \gamma$$

□◻◻

Chapter 30

Linear Systems of Equations

30.1 Linear Systems Standard Form

A linear system of equations is a set of equations that can be expressed in the form:

$$\begin{aligned}a_{1,1}x_1 + a_{1,2}x_2 + \cdots + a_{1,n}x_n &= b_1 \\a_{2,1}x_1 + a_{2,2}x_2 + \cdots + a_{2,n}x_n &= b_2 \\&\vdots \\a_{m,1}x_1 + a_{m,2}x_2 + \cdots + a_{m,n}x_n &= b_m,\end{aligned}$$

where $a_{i,j}$ are the coefficients of the variables x_j , and b_i are the constants on the right-hand side of the equations. The goal is to find the values of $x_1, x_2, \dots, x_n$ that satisfy all equations simultaneously.

More formally, given M, N with $f : M \rightarrow N$. For each $n \in N$ is the corresponding *linear system of equation* given by

$$f(m) = n$$

With the solution set

$$f^{-1}(\{n\}) = \{m \in M \mid f(m) = n\}$$

With the following types of solutions depending on the nature of f

- *At least one solution* if f is surjective.
- *One solution and no more* if f is injective/surjective.

30.2 Linear System as Matrices

A linear system can be represented in matrix form as:

$$Ax = b$$

Where A is the coefficient matrix, x is the vector of variables, and b is the vector of constants. The coefficient matrix A is an $m \times n$ matrix, where m is the number of equations and n is the number of variables. The vector x is an $n \times 1$ column vector, and the vector b is an $m \times 1$ column vector.

30.2.1 Solution Set of a Linear System

In the context of Definition 6.1, the solution set $L(A, b)$ of the system of equations associated with (A, b) is defined as

$$L(A, b) := \{x \in \mathbb{F}^n \mid Ax = b\}.$$

Reduced Echelon Form

For a matrix $A \in \mathbb{F}^{m \times n}$ with rows $Z_1, \dots, Z_n$.

We define $i \in \underline{m}$ the i -th *echelon index* $Ei_i(A)$ as

$$Ei := \min\{j \in \underline{n} \mid A_{i,j} \neq 0\}$$

We define the *echelon form* as the sequence

$$Ei(A) := (Ei_1(A), Ei_2(A), \dots, Ei_s(A))$$

If the index of all $Ei_i(A)$ coincides with the column number, then we call it the *strict echelon form*.

30.2.2 Equality of the Rank $\text{rank}(A) = \text{rank}_S(A)$

We have $\text{rank}(A) = \text{rank}_S(A)$.

Proof:

For $x = (x_1, \dots, x_n)^T \in \mathbb{F}^n$, it holds that

$$Ax = \sum_{i=1}^n x_i a_i.$$

It follows that

$$\begin{aligned} \text{rank}(A) &= \text{rank}(L_A) = \dim(\text{img}(L_A)) = \dim(\{L_A(x) \mid x \in \mathbb{F}^n\}) = \dim(\{Ax \mid x \in \mathbb{F}^n\}) \\ &= \dim\left\{\sum_{i=1}^n x_i a_i \mid x_i \in K\right\} = \dim(L(a_1, \dots, a_n)) = \text{rank}_S(A). \end{aligned}$$

30.2.3 Solvability of the Linear System $Ax = b$

The linear system $Ax = b$ is solvable if and only if

$$\text{rank}(a_1, \dots, a_n) = \text{rank}(a_1, \dots, a_n, b).$$

More concisely, we write

$$\text{rank}(A) = \text{rank}(A, b).$$

Proof:

We have

$$Ax = (a_1, \dots, a_n)x = x_1 a_1 + \dots + x_n a_n = b.$$

Therefore, a solution vector x exists if and only if

$$b \in L(a_1, \dots, a_n) \iff \text{rank}(A) = \text{rank}(A, b),$$

where $L(a_1, \dots, a_n)$ denotes the linear span of $a_1, \dots, a_n$.

30.2.4 Solution Set of the Linear System $Ax = b$

Let $x_s \in \mathbb{F}^n$ be a solution of $Ax = b$. Then the solution set is given by

$$L(A, b) = x_s + \ker(A) = \{x_s + x \mid x \in \ker(A)\}.$$

Proof:

$$x \in \ker(A) \iff Ax = 0 \iff A(x_s + x) = Ax_s + Ax = Ax_s = b.$$

30.2.5 Free Variables

In a system of linear equations, a *free variable* is a variable whose column in the coefficient matrix does not correspond to a pivot column after row reduction. Free variables are not directly solved by any equation in the system; instead, they can take any real value, and the values of the leading (or dependent) variables are expressed in terms of them.

Free variables arise naturally in systems with infinitely many solutions and play a key role in describing the solution set. Given a system of with p equations and q unknowns then if $q > p$ then there are $q - p$ free variables.

Example:

Consider the system

$$\begin{cases} x_1 + 2x_2 - x_3 = 4 \\ 3x_1 + 6x_2 - 3x_3 = 12 \end{cases}$$

which row reduces to

$$\begin{bmatrix} 1 & 2 & -1 \\ 0 & 0 & 0 \end{bmatrix}.$$

Here, the pivot is in the first column corresponding to x_1 , while x_2 and x_3 are free variables. Assigning parameters to x_2 and x_3 , the general solution can be expressed in terms of these free variables.

30.2.6 Gaussian Elimination

Gaussian elimination is a method for solving linear systems by transforming the system into an upper triangular form. The steps involved in Gaussian elimination are:

1. *Forward elimination*: Transform the system into an upper triangular form by eliminating the variables from the equations.
2. *Back substitution*: Solve for the variables starting from the last equation and substituting back into the previous equations.

The forward elimination process involves performing row operations on the augmented matrix $[A|b]$ to create zeros below the diagonal. The row operations include:

- Swapping two rows
- Multiplying a row by a non-zero scalar
- Adding or subtracting a multiple of one row from another row

These steps are defined as linear maps applied to a system like $f(m) = n$ to $(g \circ f)(m) = g(n)$. We need to be able to find the respective inverses to get to the result $(g \circ f)^{-1}(g(n)) = n$. This is accomplished via the *elementary matrices*.

Once the matrix is in upper triangular form, back substitution is used to find the values of the variables. The last equation gives the value of the last variable, which can then be substituted into the previous equations to find the other variables.

Procedure

Starting with $Ax = b$

$$\left[\begin{array}{cccc|c} a_{1,1} & a_{1,2} & \cdots & a_{1,n} & b_1 \\ a_{2,1} & a_{2,2} & \cdots & a_{2,n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m,1} & a_{m,2} & \cdots & a_{m,n} & b_m \end{array} \right]$$

For eliminating the elements underneath a pivot element we apply

$$\text{Row}_n - \frac{a_{n,i}}{a_{n-1,i}} \text{Row}_{n-1},$$

where n the index of the target row and i the index of the column.

We denote the coefficient $\frac{a_{n,i}}{a_{n-1,i}} = l_{n,i}$

This results into

$$\left[\begin{array}{cccc|c} a_{1,1} & a_{1,2} & \cdots & a_{1,n} & b_1 \\ 0 & a_{2,2} - a_{1,2} \frac{a_{2,1}}{a_{1,1}} & \cdots & a_{2,n} - a_{1,n} \frac{a_{2,1}}{a_{1,1}} & b_2 - b_1 \frac{a_{2,1}}{a_{1,1}} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & \lambda & \gamma \end{array} \right]$$

With λ and γ being a complex sums of results of the other manipulations.

30.2.7 Invariability of the Solution Set

Let $f : M \rightarrow N$ and $n \in N$. For every bijective map $g : N \rightarrow N$, the following holds for the fiber of f over n :

$$f^{-1}(\{n\}) = (g \circ f)^{-1}(\{g(n)\}).$$

Proof:

The first inclusion does not use the bijectivity of g :

Let $m \in f^{-1}(\{n\})$, i.e. $f(m) = n$. Then $g(f(m)) = g(n)$, i.e. $m \in (g \circ f)^{-1}(\{g(n)\})$. For the reverse implication one needs a left inverse of g , which of course exists because g is bijective.

□◻◻

30.2.8 Gauss-Jordan Elimination

Gauss-Jordan elimination is an extension of Gaussian elimination that transforms the matrix into reduced row echelon form (RREF). In RREF, each leading entry in a row is 1, and all entries above and below the leading entry are zeros. The steps involved in Gauss-Jordan elimination are:

1. Forward elimination: Transform the system into an upper triangular form.
2. Back substitution: Transform the upper triangular matrix into RREF by eliminating the entries above the leading 1s.

3. Solve for the variables directly from the RREF matrix.

The Gauss-Jordan elimination method is particularly useful for finding the inverse of a matrix, as it can be applied to the augmented matrix $[A|I]$, where I is the identity matrix. If the left side of the augmented matrix becomes I , then the right side will be the inverse of A .

Example:

Solve the following system of equations using Gaussian elimination:

$$\begin{aligned}2x + 3y + z &= 1 \\4x + y - z &= 2 \\-2x + y + 3z &= 3\end{aligned}$$

Solution: The augmented matrix for the system is:

$$\left[\begin{array}{ccc|c} 2 & 3 & 1 & 1 \\ 4 & 1 & -1 & 2 \\ -2 & 1 & 3 & 3 \end{array} \right]$$

Performing row operations to eliminate the variables, we can transform the matrix into upper triangular form:

$$\left[\begin{array}{ccc|c} 1 & \frac{3}{2} & \frac{1}{2} & \frac{1}{2} \\ 0 & -5 & -3 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

Now, we can perform back substitution to find the values of x , y , and z :

$$\begin{aligned}z &= 1 \\-5y - 3z &= 0 \implies y = -\frac{3}{5} \\2x + 3y + z &= 1 \implies x = \frac{1}{5}\end{aligned}$$

Thus, the solution to the system is:

$$\begin{aligned}x &= \frac{1}{5} \\y &= -\frac{3}{5} \\z &= 1\end{aligned}$$

30.2.9 Homogeneous Linear Equations

A linear equation is said to be *homogeneous* if its constant term is zero. That is, it can be written in the form:

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = 0$$

Such equations always have at least the trivial solution $x_1 = x_2 = \cdots = x_n = 0$.

30.2.10 Particular Solution

A *particular solution* to a linear system of equations is a specific solution that satisfies the system. It can be found using various methods, including substitution, elimination, or matrix methods. A particular solution is not unique; there may be multiple particular solutions depending on the system. A particular solution can be found by substituting specific values for the variables and solving for the remaining variables. For example, in the system

$$\begin{aligned}2x + 3y &= 5 \\4x - y &= 1\end{aligned}$$

we can substitute $x = 1$ into the first equation to find y :

$$\begin{aligned}2(1) + 3y &= 5 \\3y &= 3 \\y &= 1\end{aligned}$$

Thus, $(x, y) = (1, 1)$ is a particular solution to the system. However, this is not the only solution; other values of x may yield different values of y .

30.2.11 General = Particular + Homogeneous

The *general solution* of a linear system of equations is the complete set of solutions that satisfy the system. It can be expressed as the sum of a particular solution and the general solution of the associated homogeneous system.

The general solution can be written as:

$$x = x_p + x_h$$

Where x_p is a particular solution to the non-homogeneous system, and x_h is the general solution to the homogeneous system. The homogeneous system is obtained by setting the right-hand side of the equations to zero:

$$Ax = 0$$

The theorem says:

Any linear system's solution set has the form:

$$\left\{ \vec{p} + c_1\vec{\beta}_1 + \cdots + c_k\vec{\beta}_k \mid c_1, \dots, c_k \in \mathbb{R} \right\},$$

where $\vec{p}$ is a particular solution to the system, and the vectors $\vec{\beta}_1, \dots, \vec{\beta}_k$ form a basis of the solution space to the corresponding homogeneous system. The number k equals the number of *free variables* the system has after applying Gaussian elimination.

Proof:

We want to prove that every linear system of equations $Ax = b$ which has at least one solution fulfills the following properties

1. $L = \{Ax = 0\} \neq \emptyset$
2. The general solution is linear combination of the particular solutions plus the homogeneous solution.

For the first property the solution of $Ax = 0$ is the $\ker(A)$ more precisely $\ker(\phi_A)$ and thus, it is not empty because $0 \in \ker(A)$

The second proposition follows for the second one because the addition of the zero vector to a linear combination of other vectors just gives you the original combination by the axioms of the vector spaces.

□◻

30.2.12 Linear Combination Lemma

Any linear combination of linear combinations is a linear combination.

30.2.13 Example: Gaussian Elimination with 3 Equations and 4 Unknowns

Consider the following system of linear equations:

$$\begin{aligned}x_1 + 2x_2 + x_3 + x_4 &= 4 \\2x_1 + 5x_2 + x_3 + 3x_4 &= 10 \\x_1 + 3x_2 + 2x_3 + 2x_4 &= 7\end{aligned}$$

Step 1: Augmented Matrix

$$\begin{bmatrix}1 & 2 & 1 & 1 & 4 \\2 & 5 & 1 & 3 & 10 \\1 & 3 & 2 & 2 & 7\end{bmatrix}$$

Step 2: Eliminate below pivot in column 1

- Row 2 = Row 2 – 2 × Row 1
- Row 3 = Row 3 – Row 1

$$\begin{bmatrix}1 & 2 & 1 & 1 & 4 \\0 & 1 & -1 & 1 & 2 \\0 & 1 & 1 & 1 & 3\end{bmatrix}$$

Step 3: Eliminate below pivot in column 2

- Row 3 = Row 3 – Row 2

$$\begin{bmatrix}1 & 2 & 1 & 1 & 4 \\0 & 1 & -1 & 1 & 2 \\0 & 0 & 2 & 0 & 1\end{bmatrix}$$

Step 4: Back Substitution

From Row 3:

$$2x_3 = 1 \implies x_3 = \frac{1}{2}$$

From Row 2:

$$x_2 - x_3 + x_4 = 2 \implies x_2 = 2 + x_3 - x_4 = 2 + \frac{1}{2} - x_4 = \frac{5}{2} - x_4$$

From Row 1:

$$x_1 + 2x_2 + x_3 + x_4 = 4 \implies x_1 = 4 - 2x_2 - x_3 - x_4$$

Substitute:

$$x_1 = 4 - 2\left(\frac{5}{2} - x_4\right) - \frac{1}{2} - x_4 = 4 - 5 + 2x_4 - \frac{1}{2} - x_4 = -1 - \frac{1}{2} + x_4 = -\frac{3}{2} + x_4$$

General Solution

Let $x_4 = t$ (free variable), then:

$$\begin{aligned} x_1 &= -\frac{3}{2} + t \\ x_2 &= \frac{5}{2} - t \\ x_3 &= \frac{1}{2} \\ x_4 &= t \end{aligned} \quad \text{with } t \in \mathbb{R}$$

Solution Set:

$$\left\{ \begin{bmatrix} -\frac{3}{2} \\ \frac{5}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix} + t \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \\ 1 \end{bmatrix} \mid t \in \mathbb{R} \right\}$$

30.2.14 The Determinant, the cross product and the solutions of linear Systems of Equations

A linear system of three equation has the following properties:

- There is a unique solution if the determinant of the coefficient matrix is non-zero.

$$\langle (a \times b), c \rangle = \det(a, b, c) \neq 0$$


- There are infinitely many solutions if the determinant of the coefficient matrix is zero.


$$\langle (a \times b), c \rangle = \det(a, b, c) = 0$$

- There is no solution if the determinant of the coefficient matrix is zero and the system is inconsistent.

$$\langle (a \times b), c \rangle = 0$$

30.2.15 Chart for the number of solutions





30.2.16 Proof of the Gaussian Elimination

The row operations do not change the solution set.

Proof:

A linear system of equations of the form $Ax = b$, $A \in \mathbb{F}^{m \times n}$ with some solution x can be multiplied by some elementary matrix C_1, C_2, C_3 which correspond to a row operation $CAx = Cb$.

The Equality is kept and therefore, $L(A, b) \subset L(CA, Cb)$. If $x \in L(CA, Cb)$ then

$$CAx = Cb \implies C^{-1}CAx = C^{-1}Cb \implies Ax = b$$

Because of the invertibility of elementary matrices we can be sure that $L(CA, Cb) \subset L(A, b)$.

Now for the case that $L(A, b) = \emptyset$. If we assume that $L(CA, CB) \neq \emptyset$, then there would be some $x \in L(CA, CB)$ with $CAx = Cb$. By multiplying both side by C^{-1} we get $Ax = b$ and therefore, $L(A, b) \neq \emptyset$, which is a contradiction to our original assumption $L(A, b) = \emptyset$.

□□□

Now we only have to prove that each regular matrix can be transformed in the reduced echelon form or even row reduced echelon form. This is the equivalent of saying that the row operations do not change the rank of the matrix.

Proof:

If C is an elementary matrix and $A \in \mathbb{F}^{m \times n}$ some matrix. The statement $rg(CA) = rg(A)$ is true because of the invertibility of the elementary matrices and the isomorphism in vector spaces.

Given $A = (a_1, a_2, \dots, a_n)$. The exchange of two columns preserves $L := L(a_1, \dots, a_n)$, and also the $rg_s(A)$. Concerning the scaling of a column by some λ $L := L(a_1, \dots, \lambda a_k, \dots, a_n)$ also preserves $dim(L) = rg_s(A)$. And finally, the addition $L = L(a_1, \dots, a_i + \lambda a_k, \dots, a_n)$ also preserves $rg_s(A) = rg(A)$. Thus, $rg(CA) = rg(A)$.

□□□

The only part left is to prove that if A has an inverse, then with use of the row operations we can transform it in to a matrix with an upper triangular with also no zero element on the main diagonal.

Proof:

First we eliminate all the elements under the main diagonal for all columns until the column $k - 1$. With this, we have created a new matrix $A^{(\ell)} = C_\ell \cdots C_1 A$ which looks like:

$$A^{(\ell)} = \begin{bmatrix} a_{1,1}^{(\ell)} & \cdots & a_{1,k-1}^{(\ell)} & a_{1,k}^{(\ell)} & a_{1,k+1}^{(\ell)} & \cdots & a_{1,n}^{(\ell)} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & a_{k-1,k-1}^{(\ell)} & a_{k-1,k}^{(\ell)} & a_{k-1,k+1}^{(\ell)} & \cdots & a_{k-1,n}^{(\ell)} \\ 0 & \cdots & 0 & a_{k,k}^{(\ell)} & a_{k,k+1}^{(\ell)} & \cdots & a_{k,n}^{(\ell)} \\ 0 & \cdots & 0 & \vdots & a_{k+1,k+1}^{(\ell)} & \cdots & a_{k+1,n}^{(\ell)} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & a_{n,k}^{(\ell)} & \cdots & \cdots & a_{n,n}^{(\ell)} \end{bmatrix}$$

Because of $rg(A) = rg(A^{(\ell)})$, $a_{i,i}^{(\ell)} \neq 0$ for $1 \leq i < k$. All of these elements will not change for the rest of the Gaussian Elimination. Now we want to build another matrix with row operations $A^{(\mu)}$ where $a_{k,k}^{(\mu)} \neq 0$ and $a_{i,k}^{(\mu)} = 0$ for $i > k$. If $a_{k,k}^{(\ell)} \neq 0$ then (i, k) -th element will become 0 by subtracting $\frac{a_{i,k}^{(\ell)}}{a_{k,k}^{(\ell)}}$ time the k -th row of the i -th row. So can the k -th column be transformed with a series of row operations to the desired form. But if $a_{k,k}^{(\ell)} = 0$, then you have to swap it with another row under it. This is the critical point of the algorithm: What if there is no row, also $a_{i,k}^{(\ell)} = 0$ for $i \geq k$? Then $A^{(\ell)}$ would be:

$$A^{(\ell)} = \begin{bmatrix} a_{1,1}^{(\ell)} & \cdots & a_{1,k-1}^{(\ell)} & a_{1,k}^{(\ell)} & a_{1,k+1}^{(\ell)} & \cdots & a_{1,n}^{(\ell)} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & a_{k-1,k-1}^{(\ell)} & a_{k-1,k}^{(\ell)} & a_{k-1,k+1}^{(\ell)} & \cdots & a_{k-1,n}^{(\ell)} \\ 0 & \cdots & 0 & 0 & a_{k,k+1}^{(\ell)} & \cdots & a_{k,n}^{(\ell)} \\ 0 & \cdots & 0 & \vdots & a_{k+1,k+1}^{(\ell)} & \cdots & a_{k+1,n}^{(\ell)} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & \cdots & a_{n,n}^{(\ell)} \end{bmatrix}$$

But, that would imply that the k -th column is a linear combination of the other $k - 1$ columns, and then $rg(A^{(\ell)}) \leq n - 1$ which is a contradiction to the assumption $rg(A^{(\ell)}) = n$

□◻◻

30.2.17 Alternative Method for finding the Rank of a matrix

For a matrix $A \in \mathbb{F}^{(m \times n)}$. The biggest submatrix of the size $r \times r$ with a non-zero determinant gives us the *rank* of the matrix. $rg(A) = r$.

30.2.18 Cramer's Rule

For a square matrix $A = (a_1, \dots, a_n)$ and $x, b \in \mathbb{F}^n$ with $Ax = b$ with $\det A \neq 0$ then

$$A_i = (a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n), i = 1, \dots, n.$$

The solution i is then

$$x_i = \frac{\det(A_i)}{\det(A)}$$

Proof:

Given a matrix $X_i \in \mathbb{F}^{n \times n}$ by

$$\begin{bmatrix} 1 & \cdots & x_1 & \cdots \\ \vdots & \ddots & \vdots & 0 \\ & x_{i-1} & & \\ 0 & x_i & & \\ & x_{i+1} & & \\ & \vdots & \ddots & \\ & x_n & & 1 \end{bmatrix}$$

$X_i = (e_1, \dots, e_{i-1}, e_i, e_{i+1}, \dots, e_n)$. By using an n -times the Laplace formula in the first row shows $|X_i| = x_i$.

$$\begin{aligned} AX_i &= A(e_1, \dots, e_{i-1}, e_i, e_{i+1}, \dots, e_n) \\ AX_i &= A(a_1, \dots, a_{i-1}, Ax, a_{i+1}, \dots, a_n) \\ AX_i &= A(a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n) = A_i \end{aligned}$$

This implies $|A_i| = |AX_i| = |A||X_i|$ therefore, $x_i = \frac{|A_i|}{|A|}$.

QED

Another way of thinking about this in a more geometric way is to first think at the determinant as the area/volume/etc. that is spanned by the basis vector and the stretching factor of the transformation $\det(A)$. Now visualize your matrix A as a linear transformation with a non-zero determinant and think about $\det(A_i)$ as the area/volume/etc. which is spanned by the basis vector but with one of the vector substituted by b .

Now in two dimensions, think of the area spanned by the basis vectors $A = xy$. The signed area of the original parallelogram gets stretched by $\det(A)$ this gives us $Area = \det(A)y \implies y = \frac{Area}{\det(A)}$. This also works for x .

30.2.19 LU Decomposition

Given a regular system $Ax = b$ with $a_{k,k}^{(k-1)} \neq 0$, then the Gaussian elimination delivers a so called *LU-Decomposition*

$$A = LU$$

of A , where L is a under triangular matrix with main diagonal consisting of only of ones and U being a upper triangular matrix. $a_{k,k}^{(k-1)} \neq 0$ depicts the elements of the main diagonal for the $k - 1$ matrix generated after applying a Gauss operation.

Derivation

Given our matrix A , we want to eliminate the coefficients underneath the main diagonal. This is accomplished by applying an elementary matrix of the form:

$$F = \begin{bmatrix} 1 & & & \\ & \ddots & & \\ & & 1 & \\ & & \lambda & 1 \\ & & & & \ddots \end{bmatrix}$$

Hence, to get to the row-echelon-form we apply a series of such F matrices, where for the elimination of the k -th row we use the *Frobenius* matrix

$$F_k = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & -l_{k+1,k} & 1 & \\ & & -l_{n,k} & & \ddots \end{bmatrix}$$

where $-l_{i,k} = \frac{a_{i,k}^{(k-1)}}{a_{k,k}^{(k-1)}}$.

For example:

$$A = \begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{bmatrix}, a_{1,1} \neq 0$$

To eliminate $a_{2,1}$ we need to subtract $\frac{a_{2,1}}{a_{1,1}}$ times the first row.

$$F = \begin{bmatrix} 1 & 0 \\ -\frac{a_{2,1}}{a_{1,1}} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -l_{2,1} & 1 \end{bmatrix}$$

The multiplication results into

$$FA = \begin{bmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} - \frac{a_{1,1}a_{2,1}}{a_{1,1}} & a_{2,2} - \frac{a_{2,1}}{a_{1,1}} \end{bmatrix} = \begin{bmatrix} a_{1,1} & a_{1,2} \\ 0 & a_{2,2} - \frac{a_{2,1}}{a_{1,1}} \end{bmatrix}$$

Note that the main diagonal does not have to be normalized.

By repeating this process we get

$$U = A^{(n-1)} = F_{n-1} \cdots F_1 A$$

and

$$y = b^{(n-1)} = F_{n-1} \cdots F_1 b$$

Due to all F_k being regular, we get that $Ax = b$ and $Ux = y$ are equivalent. Hence, by applying the inverses

$$A = F_1^{n-1} \cdots F_{n-1} U$$

This chain of the inverses is called

$$L = \begin{bmatrix} 1 & & & & \\ l_{2,1} & \ddots & & & \\ \vdots & & 1 & & \\ & & & 1 & \\ l_{n,1} & \cdots & l_{n,n-1} & 1 \end{bmatrix}$$

The individual members are

$$F_k^{-1} = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & l_{k+1,k} & 1 & \\ & & l_{n,k} & & 1 \end{bmatrix}$$

Therefore $A = LU$ and $Ly = b$.

Algorithm

Overview:

1. Get L and U via Gaussian Elimination.
2. Solve $Ly = b$.
3. Solve $Ux = y$.

Concrete algorithm:

Start with the your $n \times n$ matrix perform the following steps across all of the entries in the main diagonal and its corresponding sub-square-matrix.

1. **Division:** For the next $k + n$ rows divide the elements in the current column by $a_{k,k}$. Only to apply to the $k \times k$ matrix.
2. **Column Elimination:** Subtract from the current row a_{k+n} -times the k row, also your current first row. Apply only for the inner sub-matrix $(k - 1) \times (k - 1)$
3. **Scoping:** Now proceed with the next diagonal element which is going to give you an inner $(n - 1) \times (n - 1)$ matrix. To repeat the process. Note that when performing the row addition, this will only affect the inner matrix; hence, our old results remain intact.
4. **Getting L and U :** After the previous process, the resulting matrix will be divided into two matrices. L consisting on the elements underneath the main diagonal with it being all ones and above all zeros; and U begin the rest upper triangular matrix.

Example:

Given

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & 4 & 0 \\ 2 & 7 & 1 \end{bmatrix}, b = \begin{bmatrix} 7 \\ 12 \\ 19 \end{bmatrix} \text{ and } b' = \begin{bmatrix} 3 \\ 4 \\ 9 \end{bmatrix}$$

solve using the LU -decomposition.

First iteration:

$$\begin{bmatrix} 2 & 1 & 1 \\ 4 & 4 & 0 \\ 2 & 7 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 2 & 4 & 0 \\ 1 & 7 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 2 & 2 & -2 \\ 1 & 6 & 0 \end{bmatrix}$$

Second iteration;

$$\begin{bmatrix} 2 & 1 & 1 \\ 2 & 2 & -2 \\ 1 & 6 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 2 & 2 & -2 \\ 1 & 3 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 1 \\ 2 & 2 & -2 \\ 1 & 3 & 6 \end{bmatrix}$$

Hence

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 3 & 1 \end{bmatrix} \quad U = \begin{bmatrix} 2 & 1 & 1 \\ 0 & 2 & -2 \\ 0 & 0 & 6 \end{bmatrix}$$

We solve $Ly = b$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 3 & 1 \end{bmatrix} y = \begin{bmatrix} 7 \\ 12 \\ 19 \end{bmatrix} \iff y = \begin{bmatrix} 7 \\ -2 \\ 18 \end{bmatrix}$$

Then we solve $Ux = y$

$$\begin{bmatrix} 2 & 1 & 1 \\ 0 & 2 & -2 \\ 0 & 0 & 6 \end{bmatrix} x = \begin{bmatrix} 7 \\ -2 \\ 18 \end{bmatrix} \iff x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

Extended Algorithm for Any Regular Matrix

We want to expand the algorithm to be compatible with the swap rows operations. Given a regular matrix $A^{(k-1)} = F_{k-1} \cdots F_1 A$, where after a Gaussian operations we get $a_{k,k}^{(k-1)} = 0$, then there exists in the k -th. row underneath at a least a $a_{i,k}^{(k-1)} \neq 0, i > k$, so that we can swap the rows.

Proof:

Let us suppose, that there does not exists such an element; then the k -th column must be a linear combination of the other columns because there is a free variable, which contradicts our initial assumption. Note that due to F_i also being regular, the multiplication of $A^{(k-1)}$ must also be regular. Therefore, it is always possible to swap rows.

□◻◻

Derivation of the Algorithm

We start with the same setup as before, but now we also have

$$P = \begin{bmatrix} 1 & & & & & & & & \\ & \ddots & & & & & & & \\ & & 0 & & & & 1 & & \\ & & & 1 & & & & & \\ & \vdots & & & \ddots & & \vdots & & \\ & & 1 & & & 0 & & & \\ & & & & & & 1 & & \\ & & & & & & & \ddots & \\ & & & & & & & & 1 \end{bmatrix}$$

which the following properties $P = P^T$ and $PP = I$ which means $P = P^{-1}$

Our U and L becomes

$$U = F_{n-1}P_{n-1} \cdots F_1P_1A$$

$$\hat{L} = (F_{n-1}P_{n-1} \cdots F_1P_{-1})^{-1}$$

And $U = \hat{L}^{-1}A$, the question is how to get $\hat{L}$ to always be a lower triangular matrix.

For that we use the inverse properties of permutation matrices to multiply by the identity matrix

$$I = P_1P_2 \cdots P_{n-1}P_{n-1} \cdots P_2P_1$$

by associativity.

We get

$$\begin{aligned}
U &= F_{n-1}P_{n-1} \cdots F_1P_1A \\
&= F_{n-1}P_{n-1} \cdots F_1P_1 \cdot (P_1P_2 \cdots P_{n-1}P_{n-1} \cdots P_2P_1)A \\
&= (F_{n-1}P_{n-1} \cdots F_1P_1 \cdot P_1P_2 \cdots P_{n-1})P_{n-1} \cdots P_2P_1A \\
&= (F_{n-1}P_{n-1} \cdots F_1P_1 \cdot P_1P_2 \cdots P_{n-1})PA \\
&= L^{-1}PA
\end{aligned}$$

Where

$$\begin{aligned}
L &= (F_{n-1}P_{n-1} \cdots F_1P_1 \cdot P_1P_2 \cdots P_{n-1})^{-1} \\
&= P_{n-2} \cdots P_2F_1^{-1}P_2F_2^{-1} \cdots P_{n-1}F_{n-1}^{-1}
\end{aligned}$$

Now L is lower triangular matrix, because of the following inductive proof:

Let $L_1 = F_1^{-1}$, $L_k = P_kL_{k-1}P_kF_k^{-1}$ for $k = 2, \dots, n-1$. This is $L_{n-1} = L$

We show that all L_k are lower triangular matrices.

Base Case: For $k = 1$

$$L_1 = \begin{bmatrix} 1 & & & \\ l_{2,1} & \ddots & & \\ \vdots & & 1 & \\ & & & 1 \\ l_{l-n,1} & & & & \ddots \end{bmatrix}$$

Inductive Step:

For $L_2 = P_2L_1P_2F_2^{-1}$:

In P_2L_1 are the second and j row changed

$$P_2L_1 = \begin{bmatrix} 1 & & & \\ l_{21} & \ddots & & 1 \\ l_{j,1} & & 1 & \\ & \vdots & & 1 \\ l_{l-n,1} & & & & \ddots \end{bmatrix}$$

Then in $P_2L_1P_2$ the change get undone; and finally, when multiplied by

$$F_2^{-1} = \begin{bmatrix} 1 & & & \\ & \ddots & & \\ l_{3,2} & 1 & & \\ \vdots & & 1 & \\ l_{n,2} & & & \ddots \end{bmatrix}$$

we get

$$L_2 = P_2 L_1 P_2 F_2^{-1} = \begin{bmatrix} 1 & & & \\ l_{2,1} & \ddots & & \\ \vdots & l_{3,2} & 1 & \\ & \vdots & & 1 \\ l_{n,1} & l_{n,2} & & \ddots \end{bmatrix}$$

Hence, by induction L_k is always a lower triangular matrix.

□◻◻

Formally we state that given a regular matrix $A \in \mathbb{R}^{n \times n}$, there are also permutations $P_k, k = 1, \dots, n-1$ such that $P = P_{n-1} \cdots P_1$ for which

$$PA = LU$$

where L and U are lower and upper triangular matrices respectively.

Extended LU-Decomposition Algorithm

The only important change for the algorithm is that for each row swaps to our matrix $A^{(i)}$ we perform the same on the matrix $P^{(i)}$. The process for the Gaussian elimination remains the same.

Overview:

1. Decompose A into L, U and P .
2. Solve $Ly = Pb$. By first permuting b just like A and then $Ly = Pv$
3. Solve $Ux = y$.

Example:

Find the L, U and P for

$$A = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 2 & 1 & 2 & 0 \\ 4 & 4 & 0 & 0 \\ 2 & 3 & 1 & 0 \end{bmatrix}$$

We now use a vector p of size $n \times 1$ where the j -th entry p_j is read as: switch the rows j with p_j

We swap rows 1 and 2, giving us $p_1 = 2$:

$$\begin{bmatrix} 2 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 4 & 4 & 0 & 0 \\ 2 & 3 & 1 & 0 \end{bmatrix}$$

Proceeding with the algorithm as for the simple LU-decomposition

$$\begin{bmatrix} 2 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 4 & 4 & 0 & 0 \\ 2 & 3 & 1 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 2 & 2 & -4 & 0 \\ 1 & 2 & -1 & 0 \end{bmatrix} (p_2 = 3) \rightarrow \begin{bmatrix} 2 & 1 & 2 & 0 \\ 2 & 2 & -4 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 2 & -1 & 0 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} 2 & 1 & 2 & 0 \\ 2 & 2 & -4 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & -1 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 2 & 1 & 2 & 0 \\ 2 & 2 & -4 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 3 & 0 \end{bmatrix} (p_3 = 4) \rightarrow \begin{bmatrix} 2 & 1 & 2 & 0 \\ 2 & 2 & -4 & 0 \\ 1 & 1 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Hence

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad U = \begin{bmatrix} 2 & 1 & 2 & 0 \\ 0 & 2 & -4 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad v = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$$

Now the solving part is equal for both procedures:

First $Ly = Pb$, where Pb is the permuted vector using p .

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} y = \begin{bmatrix} 5 \\ 4 \\ 5 \\ 3 \end{bmatrix} \iff y = \begin{bmatrix} 5 \\ -6 \\ 6 \\ 3 \end{bmatrix}$$

Finally, $Ux = y$:

$$\begin{bmatrix} 2 & 1 & 2 & 0 \\ 0 & 2 & -4 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} 5 \\ -6 \\ 6 \\ 3 \end{bmatrix} \iff x = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \end{bmatrix}$$

Greatest Pivot Strategy

As a side note, to mitigate the rounding errors in scientific computing, the swap row strategy is modified to always take the row with biggest element in the current column.

30.2.20 Cholesky Decomposition

This is a special a special case of the LU -Decomposition. Given $A \in \mathbb{R}^{n \times n}$ a symmetric, positive definite matrix it can be decomposed as

$$A = \tilde{L}D\tilde{L}^T,$$

where $\tilde{L}$ is a lower triangular matrix with 1's in the main diagonal and D a diagonal matrix with positive diagonal elements. (*Rational Cholesky Decomposition*).

Or $A = GG^T$, G is lower triangular matrix. (*Cholesky Decomposition*).

Algorithm for the Cholesky Decomposition

1. Find the LU -decomposition of A .
2. D is given by the entries in the diagonal of U .
3. Determine V by taking L^T , replacing its main diagonal with the diagonal of U and then dividing each row with the pivot elements of D . For symmetric matrices $V = L^T$.
4. Find the root of D by splitting it into two diagonal matrices which consist on the square roots of the elements in the main diagonal. Take one of them and call it S .
5. Find $G = LS$.
6. Finally $A = GG^T$.

Example:

Find the Cholesky decomposition of

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & -2 \\ 1 & -2 & 14 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & -2 \\ 1 & -2 & 14 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -3 \\ 1 & -3 & 13 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -3 \\ 1 & -3 & 13 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -3 \\ 1 & -3 & 4 \end{bmatrix}$$

We get

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & -3 & 1 \end{bmatrix} \quad U = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 0 & 0 & 4 \end{bmatrix} \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

Then, we get

$$S = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} \quad V = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 0 & 0 & 2 \end{bmatrix}$$

And finally,

$$Q = LS = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & -3 & 2 \end{bmatrix}$$

We got $A = QQ^T$.

Chapter 31

Over/Underdetermined Systems of Equations

Let us start with a little motivation for this chapter. Imagine that you have collected some data points, and you want to relate them via a linear function $Ax = (\alpha, \beta, \dots)^T$. This would probably not work because it will be impossible to connect all the points, but you could make an approximation which is the line $\alpha x_1 + \beta x_2 + \dots$ that is the closest to all the points in your data set. Thus, we will find an approximation b that is close to the image of A ($Im(A)$).

For this purpose will use a theorem that says that the best approximation is the orthogonal projection of $p_A(b)$. This projection can be found if there exist an orthonormal basis of the subspace we want to project our vector onto.

31.0.1 Normal Equations

Given a projection $p_A(b)$ onto the subspace U generated by $A = (a_1, \dots, a_n)$ then there exists an $x \in \mathbb{R}^n$ such that

$$p_A(b) = \sum_{k=1}^n x_k a_k = Ax$$

Also, because of the orthogonal property then

$$\begin{aligned} b - p_A(b) \perp U &\implies b - Ax \in U^\perp \\ &\implies b - Ax \perp a_k \quad \forall k \\ &\implies \langle a_k, b - Ax \rangle = 0 \quad \forall k \\ &\iff A^T(b - Ax) = \vec{0} \\ &\implies A^T b = Ax \end{aligned}$$

This equation $A^T b = Ax$ is called the *normal equation*.

A *normal equation* of a matrix $A \in \mathbb{R}^{(m \times n)}$ has a solution. If $rg(A) = n$ then

$$x = (A^T A)^{-1} A^T b$$

is a unique solution to the system. The term $(A^T A)^{-1} A^T$ is called the *Generalized Inverse* of A .

Proof:

The statement is true because of the existence of the orthogonal projection which give us a concrete basis. Also, the fact that $rg(A) = n$ also implies that. And also because of the invertibility of $(A^T A)^{-1}$.

31.0.2 Method of the Least Squares

For a linear system of equations $Ax = b$ with $A \in \mathbb{R}^{(m \times n)}$, $b \in \mathbb{B}^m$ and $m \geq n$. For the case $rg(A) = n$ we have

$$x_s = (A^T A)^{-1} A^T b$$

and

$$\|b - Ax_s\| = \min_{z \in \mathbb{R}^n} \|b - Az\|$$

Here the vector z is called *approximation solution via the method of the least squares*.

Proof:

We know that because of $rg(A) = n$ our solution is unique. And that $p_A(b) = Ax$ is our orthogonal projection. We can use Pythagoras theorem to

$$\begin{aligned} \|b - Az\|^2 &= \|(b - p_A(b)) + (p - p_A(b))\|^2 \\ &= \|(b - p_A(b))\|^2 + \|(p - p_A(b))\|^2 \\ &\geq \|(b - p_A(b))\|^2 \\ &= \|b - Ax\|^2 \end{aligned}$$

Thus, $p_A(b)$ gives us the smallest distance to the $Img(A)$

Example I:

$$x = 1, x = -1$$

$$\|Ax - b\|^2 = (x + 1)^2 + (x - 1)^2 = 2x^2 + 2$$

Then $\|Ax - b\|^2$ gives us the smallest distance for $x = 0$.

Example II:

We can also take the last example and solve it with an alternative method.

$$A = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, x = (x), b = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Let us use the normal equations

$$A^T Ax = A^T b$$

$$(1, 1) \begin{bmatrix} 1 \\ 1 \end{bmatrix} x = (1, 1) \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\implies 2x = 0$$

Example 3:

Given are

$$x_1 = 1, x_2 = 2, x_1 + x_2 = -1$$

with the corresponding matrix

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}$$

and solution vector

$$b = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

Now let us build the normal equation

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix} \vec{x} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Now with Gaussian elimination we get $x = (-\frac{1}{3}, \frac{2}{3})^T$.

Finally, we can use the $\vec{x}$ to solve $Ax - b$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} -\frac{1}{3} \\ \frac{2}{3} \end{bmatrix} - \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix} = \frac{4}{3} \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix}$$

And therefore, $\|Ax - b\|^2 = \frac{48}{9}$.

31.0.3 Underdetermined Systems of Equations

For a linear system $Ax = b$ such that $A \in \mathbb{R}^{m \times n}$, $x \in \mathbb{R}^n$, $b \in \mathbb{R}^m$ and $m \leq n$. For $rg(A) = m$ we get the solution vector

$$x_s = A^T(AA^T)^{-1}b$$

Also, x_s is the solution with the smallest length.

$$\|x_s\| = \min_{Ax=b} \|x\|,$$

and x_s is orthogonal to the homogenous system $Az = 0$ which means

$$x_s \perp \ker(A)$$

Finally, we call for $rg(A) = m$, $A^T(AA^T)^{-1}$ the *Generalized Inverse* of A .

Proof:

First, we show the invertibility of AA^T because this implies that x_s is defined.

Given $x \in \ker(AA^T)$. This implies that

$$AA^T x = 0$$

which implies

$$x^T AA^T x = 0$$

because of $x^T A = A^T x$ and this gives us

$$\|A^T x\| = 0$$

therefore, $x \in \text{kern}(A^T)$.

Now because of $\text{rg}(A^T) = \text{rg}(A) = m$ we get $\dim(\text{kern}(A^T)) = 0$, thus $x = \vec{0}$. Because of that $\text{kern}(AA^T) = 0$ and $\text{rg}(AA^T) = m$ thus, the invertibility is proven.

x_s is a solution of $Ax = b$ because of

$$Ax_s = A(A^T(AA^T)^{-1}b) = AA^T(AA^T)^{-1}b = b$$

$x_s \perp \text{kern}(A)$ because for $z \in \text{kern}(A)$

$$\langle z, x_s \rangle = z^T A^T (AA^T)^{-1} b = \langle Az, (AA^T)^{-1} b \rangle = 0$$

due to $Az = 0$.

Finally, thanks to the general solution theorem we know that $x = x_s + z$. Using Pythagoras and by taking into consideration $x_s \perp \text{kern}(A)$

$$\|x\|^2 = \|x_s\|^2 + \|z\|^2 \geq \|x_s\|^2$$

Example:

We have the equation $2x_1 + 2x_2 - x_3 = 6$.

In this case the solution build a plane in $\mathbb{R}^3$, thus the solution is the normal vector that cuts the plane. We build the hessian normal form.

$$\frac{1}{3}(2x_1 + 2x_2 - x_3) = 2$$

whose solution is $x = \frac{2}{3}(2, 2, -1)^T$.

Now with our method: $A = (2, 2, -1)$, $x = (x_1, x_2, x_3)$, $b = 6$

$$x = A^T(AA^T)^{-1}b = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix} \left((2, 2, -1) \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix} \right)^{-1} 6$$

$$x = \frac{2}{3}(2, 2, -1)^T$$

Chapter 32

Analytical Geometry

In this section, we will cover the topics for the geometry of $\mathbb{R}^2$ and $\mathbb{R}^3$. And maybe also in higher dimensions.

32.0.1 Vectors and Points

In analytical geometry, points and vectors are the basic elements.

A point in $\mathbb{R}^3$ is represented as $\vec{P} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$. Or as $(x_1, x_2, \dots, x_n)$.

A vector is an object with direction and magnitude (in this case), also represented as $\vec{v} = \begin{bmatrix} v_x \\ v_y \\ v_z \end{bmatrix}$.

Or as $(x_1, x_2, \dots, x_n)^T$.

32.0.2 Vector Addition and Scalar Multiplication

Given two vectors $\vec{a}$ and $\vec{b}$:

$$\vec{a} \pm \vec{b} = \begin{bmatrix} a_x \\ a_y \end{bmatrix} \pm \begin{bmatrix} b_x \\ b_y \end{bmatrix} = \begin{bmatrix} a_x \pm b_x \\ a_y \pm b_y \end{bmatrix}$$

For a scalar λ and vector $\vec{a}$:

$$\lambda \vec{a} = \lambda \begin{bmatrix} a_x \\ a_y \end{bmatrix} = \begin{bmatrix} \lambda a_x \\ \lambda a_y \end{bmatrix}$$




Figure 32.1: Vector addition: $\vec{a} + \vec{b}$

32.0.3 Equation of a Line

A line is defined by a point $\vec{P}$ and a direction vector $\vec{v}$:

$$\vec{r}(t) = \vec{P} + t\vec{v}, \quad t \in \mathbb{R}$$

32.0.4 Equation of a Plane

A plane is defined by a point $\vec{P}$ and a normal vector $\vec{n}$:

$$P : \vec{x} = \vec{s} + \lambda_1 \vec{v}_1 + \lambda_2 \vec{v}_2$$

32.0.5 Dot Product

The inner product of vectors $\vec{a}$ and $\vec{b}$ is a function $\langle x, y \rangle := V \times V \rightarrow \mathbb{K}$ with the following properties:

$\forall \vec{a}, \vec{b} \in V$:


- $\langle \vec{a}, \vec{b} \rangle = \langle \vec{b}, \vec{a} \rangle$
- $\langle \vec{a}, \vec{b} + \vec{c} \rangle = \langle \vec{a}, \vec{b} \rangle + \langle \vec{a}, \vec{c} \rangle$
- $\langle \vec{a}, \lambda \vec{b} \rangle = \lambda \langle \vec{a}, \vec{b} \rangle$
- $\langle \vec{a}, \vec{b} \rangle = 0 \Leftrightarrow \vec{a} = \vec{0} \vee \vec{b} = \vec{0}$

Formula:

$$\langle \vec{a}, \vec{b} \rangle = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n = \sum_{i=1}^n a_i b_i$$

Proof:

Let us look at the following Triangle:



We can use the Cosine law: $\|v - u\|^2 = \|v\|^2 + \|u\|^2 - 2\|u\|\|v\|\cos\theta$.

Now we expand:

$$\|u\|^2 = u_1^2 + u_2^2 \quad \text{and} \quad \|v\|^2 = v_1^2 + v_2^2$$

Then: $\|v - u\|^2 = (v_1 - u_1)^2 + (v_2 - u_2)^2$ and by re-grouping after expansion we get

$$\langle v, u \rangle = v_1 u_1 + v_2 u_2 = \|v\|\|u\|\cos\theta$$

ΩεΩ

And if we manipulate this expression we can also prove the angle formulas. And with that information we can prove that

$$\cos \frac{\pi}{2} = \frac{\langle v, u \rangle}{\|v\| \|u\|} = 0$$

Only if the Dot Product is equal to zero so, like that we have also proved that only orthogonal vector have a dot product of zero.

32.0.6 Orthogonal Vectors

Two vectors $\vec{a}$ and $\vec{b}$ are orthogonal if:

$$\langle \vec{a}, \vec{b} \rangle = 0$$

32.0.7 The Orthogonal Projection

The *orthogonal projection of vector $\vec{a}$ onto vector $\vec{b}$* is given by:

$$\text{Proj}_{\vec{b}}(\vec{a}) = \frac{\langle \vec{a}, \vec{b} \rangle}{\|\vec{b}\|^2} \cdot \vec{b}$$

Proof:

We know that

$$p \parallel q \implies p = \alpha q \quad \forall \alpha \in \mathbb{F} \text{ and } \forall p, q \in V$$

Then let us define the orthogonal projection of vector $\vec{a}$ on vector $\vec{b}$ as p . We want to find α such that $p = \alpha q$ where $q = \vec{b}$. We also know that the vector $a - p$ is orthogonal to q :

$$\begin{aligned} \langle a - p, q \rangle = 0 &\implies \langle a - \alpha q, q \rangle = 0 \\ &\implies \langle a, q \rangle - \alpha \langle q, q \rangle = 0 \quad \text{by the properties of the inner product} \\ \alpha &= \frac{\langle a, q \rangle}{\langle q, q \rangle} \end{aligned}$$

Therefore, the orthogonal projection of vector $\vec{a}$ on $\vec{b}$ is given by multiplying $\vec{b}$ by the scalar α .

□◻◻




Figure 32.2: Orthogonal projection of $\vec{a}$ onto $\vec{b}$

32.0.8 Vector Projections in Data Science

In data science, vector projections are used in various applications such as dimensionality reduction, feature extraction, and data visualization. For example, in Principal Component Analysis (PCA), data points are projected onto the directions of maximum variance to reduce the dimensionality of the dataset while preserving as much information as possible.

We find a unit vector in the direction of interest (e.g., a principal component) $\vec{u}_i$ onto which we want to project our data points $\vec{x}_j$. Note we can find multiple such vectors $\vec{u}_1, \vec{u}_2, \dots, \vec{u}_k$ to project onto a lower-dimensional subspace.

$$\text{Proj}_{\vec{u}}(\vec{v}) = \langle \vec{v}, \vec{u} \rangle \vec{u} = k\vec{u}$$

Proof:

Let us start with the derivation. We have a vector $\vec{v}$, and we want to project it onto a unit vector $\vec{u}$. The

$$\vec{u} = \frac{\vec{v}}{\|\vec{v}\|}$$

and

$$p = k\vec{u} \quad \text{and} \quad d = \vec{v} - p$$

also $\langle d, u \rangle = 0$

Analog to the derivation of the formula of the orthogonal projection we get

$$d = \vec{v} - k\vec{u} \implies \langle \vec{v} - k\vec{u}, \vec{u} \rangle = 0 \implies \langle \vec{v}, \vec{u} \rangle - k\langle \vec{u}, \vec{u} \rangle = 0$$

Notice that $\langle \vec{u}, \vec{u} \rangle = 1$ because $\vec{u}$ is a unit vector. Therefore,

$$k = \langle \vec{v}, \vec{u} \rangle$$

Thus, the projection of $\vec{v}$ onto the unit vector $\vec{u}$ is given by:

$$\text{Proj}_{\vec{u}}(\vec{v}) = \langle \vec{v}, \vec{u} \rangle \vec{u} = k\vec{u}$$

QED

,

32.0.9 Dot Product as Projection

The *dot product* of two vectors can be interpreted as projecting one vector onto the other and then multiplying the length of the projecting vector to the magnitude of the other. For real vectors.

This comes from the fact that taking a dot product of real vectors is the same as multiplying one of them with the linear transformation formed by the transpose version of the other vector.

$$\langle \vec{a}, \vec{b} \rangle = \vec{a}^T \vec{b}$$

The transformation will take a vector from our space and take it to the number line, and it is also linear. In our case,

$$\text{Proj} = \begin{bmatrix} u_{x_1} & u_{x_2} & \cdots & u_{x_n} \end{bmatrix}$$

is defined this way because when we project our basis vector $\vec{b}_i$ onto the line our vector $\vec{u}$ because of the symmetry we will get the same value as if we were projecting $\vec{u}$ onto the line $\vec{b}_i$ lives in.

32.0.10 Metrics for Vectors

The metric of a vector $\vec{a}$ is described by different norms:

It also has the following properties:

- $\|\vec{a}\| \geq 0$ and $\|\vec{a}\| = 0 \Leftrightarrow \vec{a} = \vec{0}$
- $\|\lambda\vec{a}\| = |\lambda|\|\vec{a}\|$
- $\|\vec{a} + \vec{b}\| \leq \|\vec{a}\| + \|\vec{b}\|$ (Triangle inequality)
- $\|\vec{a} + \vec{b}\|^2 = \|\vec{a}\|^2 + \|\vec{b}\|^2 + 2\langle\vec{a}, \vec{b}\rangle$ (Pythagorean theorem)

These are the most common metric, although we will be using primarily the euclidean metric:

- *Euclidean metric:*

$$\|\vec{a}\| = \sqrt{\langle\vec{a}, \vec{a}\rangle} = \sqrt{a_1^2 + a_2^2 + \cdots + a_n^2}$$

- *Manhattan metric:*

$$\|\vec{a}\|_1 = |a_1| + |a_2| + \cdots + |a_n|$$

- *Maximum metric:*

$$\|\vec{a}\|_\infty = \max(|a_1|, |a_2|, \dots, |a_n|)$$

32.0.11 Angle Relations

The angle θ between two vectors is:

$$\cos \theta = \frac{\langle\vec{a}, \vec{b}\rangle}{\|\vec{a}\|\|\vec{b}\|}, \quad \theta = \arccos\left(\frac{\langle\vec{a}, \vec{b}\rangle}{\|\vec{a}\|\|\vec{b}\|}\right)$$

Line-Line Angle

Use direction vectors $\vec{v}_1$ and $\vec{v}_2$:

$$\theta = \arccos\left(\frac{\langle\vec{v}_1, \vec{v}_2\rangle}{\|\vec{v}_1\|\|\vec{v}_2\|}\right)$$

Line-Plane Angle

Let $\vec{v}$ be the line's direction and $\vec{n}$ the plane's normal:

$$\theta = \arcsin\left(\frac{\langle\vec{v}, \vec{n}\rangle}{\|\vec{v}\|\|\vec{n}\|}\right)$$

Plane-Plane Angle

Angle between planes is angle between their normals:

$$\theta = \arccos\left(\frac{\langle\vec{n}_1, \vec{n}_2\rangle}{\|\vec{n}_1\|\|\vec{n}_2\|}\right)$$

32.0.12 Line Relations

Two lines can be:

- *Identical*: same direction vector and point
- *Parallel*: direction vectors are proportional
- *Intersecting*: one solution for t_1, t_2 such that $\vec{r}_1(t_1) = \vec{r}_2(t_2)$

- *Skew*: not parallel, do not intersect

To find the relation:

1. Check if direction vectors are scalar multiples $\Rightarrow$ parallel
2. Solve $\vec{P}_1 + t\vec{v}_1 = \vec{P}_2 + s\vec{v}_2$ for t and s :
 - Solution exists $\Rightarrow$ intersect
 - No solution $\Rightarrow$ skew
3. If same point and direction vector $\Rightarrow$ identical

32.0.13 Normalization of a vector

To normalize a vector $\vec{a}$, we divide it by its length:

$$\hat{\vec{a}} = \frac{\vec{a}}{\|\vec{a}\|} = \frac{\begin{bmatrix} a_x \\ a_y \\ a_z \end{bmatrix}}{\sqrt{a_x^2 + a_y^2 + a_z^2}} = \begin{bmatrix} \frac{a_x}{\|\vec{a}\|} \\ \frac{a_y}{\|\vec{a}\|} \\ \frac{a_z}{\|\vec{a}\|} \end{bmatrix}$$

32.0.14 The Cross Product

The cross product of two vectors $\vec{a}$ and $\vec{b}$ in $\mathbb{R}^3$ is defined as:

$$\vec{a} \times \vec{b} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix}$$

or

$$\det \left(\begin{bmatrix} \hat{i} & v_1 & w_1 \\ \hat{j} & v_2 & w_2 \\ \hat{k} & v_3 & w_3 \end{bmatrix} \right) = \hat{i}(v_2w_3 - v_3w_2) - \hat{j}(v_1w_3 - v_3w_1) + \hat{k}(v_1w_2 - v_2w_1)$$

The cross product has the following properties:

- $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$
- $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$
- $(\lambda_1\vec{a}) \times (\lambda_2\vec{b}) = \lambda_1\lambda_2(\vec{a} \times \vec{b})$
- $\|\vec{a} \times \vec{b}\| = \|\vec{a}\|\|\vec{b}\|\sin(\theta)$
- $\langle \vec{a}, (\vec{b} \times \vec{c}) \rangle = 0$ (scalar triple product)
- $\vec{a} \times \vec{b} = \vec{0} \Leftrightarrow \vec{a} = \lambda\vec{b}$ for some $\lambda \in \mathbb{R}$ (parallel vectors)
- $\vec{a} \times \vec{b} = \vec{0} \Leftrightarrow \vec{a} = \vec{0} \vee \vec{b} = \vec{0}$ (zero vector)

The cross product is not defined in $\mathbb{R}^2$. The cross product is not commutative, but it is associative:

$$(\vec{a} \times \vec{b}) \times \vec{c} = \vec{a} \times (\vec{b} \times \vec{c})$$

The cross product is distributive over vector addition:

$$\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$$

The cross product is anti-commutative:

$$\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$$

The cross product is not associative:

$$(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$$

The cross product is not distributive over scalar multiplication:

$$\lambda(\vec{a} \times \vec{b}) \neq (\lambda\vec{a}) \times \vec{b}$$

The length of the cross product in $\mathbb{R}^3$ is the area of the parallelogram spanned by the two vectors.

Determinant Trick

For two vectors $\vec{a}, \vec{b}$ in $\mathbb{C}^3$ we can compute the cross product by:

$$\vec{n} = \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix}$$

Connection to the Volume of the Parallelepiped

We want to define a linear transformation f which relates the volume of the parallelepiped formed by three vectors, the determinant; and the dot product of a input vector of the transformation with some vector $\vec{p}$.

$$f\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) = \det\left(\begin{bmatrix} x & v_1 & w_1 \\ y & v_2 & w_2 \\ z & v_3 & w_3 \end{bmatrix}\right) = \left\langle \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix}, \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right\rangle$$

This gives us another version of the determinant trick view previously and we know that this gives us a vector perpendicular to $\vec{v}$ and $\vec{w}$. The motivation behind this weird formula is that if you compute both sides of the equation you will get the coordinates $\vec{p}$ from the determinant trick.

This whole equality can be related to the geometric meaning of the cross product. Remember that the dot product of two vectors can be interpreted as projecting one onto another and then multiplying the length of the projecting to the magnitude of the other. This is exactly the same as projecting our input vector to a line which is perpendicular to the parallelogram and then multiplying it by the area which will gives us the volume.

Derivation of the formula

We define $\vec{n}$ as a vector perpendicular to $\vec{v}$ and $\vec{w}$. Therefore,

$$\vec{v} \cdot \vec{n} = 0 \quad \text{and} \quad \vec{w} \cdot \vec{n} = 0$$

Writing this out in components, let $\vec{v} = (v_1, v_2, v_3)$, $\vec{w} = (w_1, w_2, w_3)$, and $\vec{n} = (n_1, n_2, n_3)$. Then:

$$v_1 n_1 + v_2 n_2 + v_3 n_3 = 0$$

$$w_1 n_1 + w_2 n_2 + w_3 n_3 = 0$$

Now we multiply the first equation by $-w_1$, the second by v_1 and add them

$$n_2(w_2 v_1 - v_2 w_1) + n_3(w_3 v_1 - v_3 w_1) = 0$$

We know that the equation $pc_1 + qc_2 = 0$ has the results $c_1 = -q$ and $c_2 = p$. Thus,

$$n_2 = v_3w_1 + w_3v_1$$

$$n_3 = v_1w_2 - w_1v_2$$

Finally, by substituting into the original equation and solving we get

$$n_1 = v_2w_3 - v_3w_2$$

Thus, we have gotten the formula for the cross product.

□◻◻

32.0.15 Orthogonal vectors in $\mathbb{R}^2$ $\mathbb{R}^3$

Orthogonal vectors in $\mathbb{R}^2$

1. Interchange the components
2. Change the sign of one component

Orthogonal vectors in $\mathbb{R}^3$

1. Interchange two components
2. the one that was not changed, set to zero
3. Change the sign of first component

Example:

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \vec{b} = \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$$

32.0.16 Hessian Normal Form

The Hessian normal form is a way of expressing the equation of a plane in three-dimensional space using a normalized normal vector. It is particularly useful in computational geometry and physics, where signed distances from points to planes are important.

Geometric Interpretation

The Hessian normal form represents a plane by specifying:

- A unit normal vector $\vec{n} = (a, b, c)$ to the plane,
- and the shortest distance d from the origin to the plane.

This form is derived by normalizing the general plane equation. A plane in 3D can be written as:

$$ax + by + cz + d = 0,$$

where (a, b, c) is a normal vector to the plane and d is the dot product of the normal vector with a point p . If we divide all terms by $\sqrt{a^2 + b^2 + c^2}$, we normalize the normal vector:

$$\frac{a}{\sqrt{a^2 + b^2 + c^2}}x + \frac{b}{\sqrt{a^2 + b^2 + c^2}}y + \frac{c}{\sqrt{a^2 + b^2 + c^2}}z + \frac{d}{\sqrt{a^2 + b^2 + c^2}} = 0.$$

Let:

$$\vec{n} = \left(\frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}} \right), \quad d' = \frac{d}{\sqrt{a^2 + b^2 + c^2}},$$

then the equation becomes:

$$\vec{n} \cdot \vec{r} + d' = 0,$$

which is the Hessian normal form. Here, $\vec{r} = (x, y, z)$ is any point on the plane, and $\vec{n}$ is the unit normal.

Signed Distance to the Plane

This form allows easy calculation of the signed distance from any point $\vec{p}$ to the plane:

$$\text{distance} = \vec{n} \cdot \vec{p} + d',$$

which is positive if $\vec{p}$ lies on the same side of the plane as the normal vector.

Derivation Illustration

To visualize the derivation, imagine a plane with unit normal vector $\vec{n}$, and a point P in space. The shortest distance from P to the plane is the projection of the vector $\vec{p} - \vec{q}$ onto $\vec{n}$, where $\vec{q}$ is any point on the plane. This leads to:

$$\text{distance} = (\vec{p} - \vec{q}) \cdot \vec{n}.$$

This gives the signed distance formula and thus, motivates the Hessian form.

32.0.17 Converting from the parametric form to the Hessian normal form

Steps:

1. Find the normal vector $\vec{n}$ of the plane
2. Normalize the normal vector
3. Find the distance d from the origin to the plane
4. Write the Hessian normal form

32.0.18 Converting from the Hessian normal form to the parametric form

Steps:

1. Find a point on the plane
2. Find two direction vectors in the plane
3. Write the parametric form

32.0.19 Properties of lines and planes

- Two planes are parallel if their normal vectors are scalar multiples of each other.
- A line and a plane are parallel if the direction vector of the line is orthogonal to the normal vector of the plane.
- A line intersects a plane if there exists a point on the line that satisfies the equation of the plane.
- Two planes intersect in a line if their normal vectors are not parallel.
- Three planes can intersect in a point, a line, or not at all.

- If we have a line G and a point on the line, for every vector $\vec{n}$ that is orthogonal to the direction vector of the line: $x \in G \iff \langle x, \vec{n} \rangle = 0$
- If p and q are two points in the line G with a normal vector then $\langle p, \vec{n} \rangle = \langle q, \vec{n} \rangle$
- Let E be a plane with the origin p and the direction vector $\vec{v}$ and $\vec{w}$, then there exist a normal vector and $x \in E \iff \langle x, \vec{n} \rangle = \langle p, \vec{n} \rangle$

32.0.20 Convert Normal Vector in Two Direction Vectors

Steps:

1. Given the normal vector $\vec{n} = (a, b, c)$, interchange a and b and multiply b by -1
2. Set the other component to 0. This gives you the first direction vector $\vec{v} = (-b, a, 0)$
3. Take the original normal vector $\vec{n}$ and interchange a and c and multiply c by -1
4. Set the other component to 0. This gives you the second direction vector $\vec{w} = (-c, 0, a)$

32.0.21 Intersection between Line and Plane

To find the intersection between a line and a plane, we can use the following steps:

1. Write the parametric form of the line: $\vec{r}(t) = \vec{P} + t\vec{v}$, where $\vec{P}$ is a point on the line and $\vec{v}$ is the direction vector.
2. Write the equation of the plane in Hessian normal form: $\langle \vec{n}, \vec{x} - \vec{P} \rangle = 0$, where $\vec{n}$ is the normal vector and $\vec{P}$ is a point on the plane.
3. Substitute the parametric form of the line into the equation of the plane.
4. Solve for t to find the intersection point.
5. Substitute t back into the parametric form of the line to find the intersection point.

Example:

Given the line:

$$g(t) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

and the plane:

$$E(u, m) = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} + u \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + m \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix},$$

we want to find the intersection point.

Step 1: Determine the normal vector of the plane using the cross product of the two direction vectors:

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

$$\vec{n} = \vec{v}_1 \times \vec{v}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix}$$

Step 2: Use the normal vector and a point on the plane to write the plane equation:

$$\langle \vec{n}, \vec{x} - \vec{Q} \rangle = 0, \quad \vec{Q} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$$

Step 3: Plug the line into the plane equation:

$$\vec{x}(t) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \Rightarrow \vec{x}(t) - \vec{Q} = \begin{bmatrix} t \\ t-1 \\ t-1 \end{bmatrix}$$

Now compute the dot product:

$$\langle \vec{n}, \vec{x}(t) - \vec{Q} \rangle = \begin{bmatrix} -1 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} t \\ t-1 \\ t-1 \end{bmatrix} = -t + (t-1) + (t-1) = t-2$$

Step 4: Solve for t :

$$t-2=0 \Rightarrow t=2$$

Step 5: Substitute $t=2$ into the line:

$$\vec{x}(2) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$$

Result: The line intersects the plane at the point

$$\begin{bmatrix} 2 \\ 2 \\ 3 \end{bmatrix}$$

Proof:

$$E : \langle x, n \rangle = \langle p, n \rangle$$

$$G : x = p + t \cdot v$$

$$\langle x, n \rangle = c$$

$$\langle p + t \cdot v, n \rangle = c$$

$$\langle p, n \rangle + t \cdot \langle v, n \rangle = c$$

$$t = \frac{c - \langle p, n \rangle}{\langle v, n \rangle}$$

□◻◻

32.0.22 Distances between points, lines and planes

Distance between two points

The distance between two points $\vec{P}_1$ and $\vec{P}_2$ in $\mathbb{R}^n$ is given by:

$$d(\vec{P}_1, \vec{P}_2) = \|\vec{P}_1 - \vec{P}_2\| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + \cdots + (z_1 - z_2)^2}$$

where $\vec{P}_1 = (x_1, y_1, \dots, z_1)$ and $\vec{P}_2 = (x_2, y_2, \dots, z_2)$.

Distance between a point and a hyperplane

The distance between a point $\vec{P}$ and a hyperplane defined by the equation $\langle \vec{n}, \vec{x} - \vec{P}_0 \rangle = 0$ is given by:

$$d(\vec{P}, \text{hyperplane}) = \frac{|\langle \vec{n}, \vec{P} - \vec{P}_0 \rangle|}{\|\vec{n}\|}$$

where $\vec{P}_0$ is a point on the hyperplane and $\vec{n}$ is the normal vector of the hyperplane.

Distance between two lines

The distance between two lines in $\mathbb{R}^3$ can be calculated using the formula:

$$d = \frac{|\langle \vec{v}_1 \times \vec{v}_2, \vec{P}_2 - \vec{P}_1 \rangle|}{\|\vec{v}_1 \times \vec{v}_2\|}$$

where $\vec{P}_1$ and $\vec{P}_2$ are points on the two lines, and $\vec{v}_1$ and $\vec{v}_2$ are the direction vectors of the lines.

Distance between a point and a line

The distance between a point $\vec{P}$ and a line defined by the parametric equation $\vec{r}(t) = \vec{P}_0 + t\vec{v}$ is given by:

$$d(\vec{P}, \text{line}) = \frac{\|\vec{v} \times (\vec{P} - \vec{P}_0)\|}{\|\vec{v}\|}$$

where $\vec{P}_0$ is a point on the line and $\vec{v}$ is the direction vector of the line.

Distance between two planes

The distance between two parallel planes defined by the equations $\langle \vec{n}, \vec{x} - \vec{P}_1 \rangle = 0$ and $\langle \vec{n}, \vec{x} - \vec{P}_2 \rangle = 0$ is given by:

$$d = \frac{|\langle \vec{n}, \vec{P}_2 - \vec{P}_1 \rangle|}{\|\vec{n}\|}$$

where $\vec{P}_1$ and $\vec{P}_2$ are points on the two planes, and $\vec{n}$ is the normal vector of the planes.

Distance between a point and a plane

The distance between a point $\vec{P}$ and a plane defined by the equation $\langle \vec{n}, \vec{x} - \vec{P}_0 \rangle = 0$ is given by:

$$d(\vec{P}, \text{plane}) = \frac{|\langle \vec{n}, \vec{P} - \vec{P}_0 \rangle|}{\|\vec{n}\|}$$

where $\vec{P}_0$ is a point on the plane and $\vec{n}$ is the normal vector of the plane.

Example:

Distance Between Two Skew Lines

To find the shortest distance between two skew lines, we use the formula:

$$\text{distance} = \frac{|\langle (\vec{P}_2 - \vec{P}_1), (\vec{v}_1 \times \vec{v}_2) \rangle|}{\|\vec{v}_1 \times \vec{v}_2\|}$$

Where:

- $\vec{P}_1$ and $\vec{P}_2$ are points on each line,
- $\vec{v}_1$ and $\vec{v}_2$ are the direction vectors,
- $\vec{v}_1 \times \vec{v}_2$ is the cross product of the direction vectors.

Given:

$$g_1 : \vec{r}_1(a) = \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix} + a \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \quad g_2 : \vec{r}_2(b) = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + b \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

Step 1: Set

$$\vec{P}_1 = \begin{bmatrix} 2 \\ 2 \\ 2 \end{bmatrix}, \quad \vec{v}_1 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

$$\vec{P}_2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

Step 2: Compute the vector between base points:

$$\vec{P}_2 - \vec{P}_1 = \begin{bmatrix} 1-2 \\ 2-2 \\ 3-2 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Step 3: Compute the cross product:

$$\vec{v}_1 \times \vec{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \times \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} (1)(1) - (1)(2) \\ (1)(3) - (0)(1) \\ (0)(2) - (1)(3) \end{bmatrix} = \begin{bmatrix} -1 \\ 3 \\ -3 \end{bmatrix}$$

Step 4: Compute scalar triple product:

$$\langle (\vec{P}_2 - \vec{P}_1), (\vec{v}_1 \times \vec{v}_2) \rangle = \left\langle \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 3 \\ -3 \end{bmatrix} \right\rangle = (-1)(-1) + (0)(3) + (1)(-3) = 1 + 0 - 3 = -2 \Rightarrow |\dots| = 2$$

Step 5: Magnitude of the cross product:

$$\|\vec{v}_1 \times \vec{v}_2\| = \sqrt{(-1)^2 + 3^2 + (-3)^2} = \sqrt{1 + 9 + 9} = \sqrt{19}$$

Final Answer:

$$\text{distance} = \frac{2}{\sqrt{19}} \approx 0.458$$

$$\text{Shortest distance between the lines is } \frac{2}{\sqrt{19}}$$

32.0.23 Foot of the Perpendicular and Mirror Point

Foot of the Perpendicular

The *foot of the perpendicular* from a point $\vec{P}$ to a line (or plane) is the point on the line (or plane) where the perpendicular from $\vec{P}$ meets it.

Line case:

Given a line in parametric form:

$$g : \vec{r}(t) = \vec{A} + t\vec{v}$$

and a point $\vec{P}$ not on the line, the foot of the perpendicular $\vec{F}$ satisfies:

$$(\vec{P} - \vec{F}) \perp \vec{v} \Rightarrow (\vec{P} - (\vec{A} + t\vec{v})) \cdot \vec{v} = 0$$

Solve this inner product for t , then compute:

$$\vec{F} = \vec{A} + t\vec{v}$$

Plane case:

Given a plane in normal form:

$$\langle \vec{n}, \vec{x} - \vec{Q} \rangle = 0$$

then the foot of the perpendicular from point $\vec{P}$ to the plane is:

$$\vec{F} = \vec{P} - ((\vec{P} - \vec{Q}) \cdot \vec{n}) \cdot \vec{n}$$

Mirror Point

The *mirror point* (or reflected point) of $\vec{P}$ across a line or plane is the point $\vec{P}'$ such that the midpoint between $\vec{P}$ and $\vec{P}'$ is the foot of the perpendicular.

Formula:

$$\vec{P}' = 2\vec{F} - \vec{P}$$

Where $\vec{F}$ is the foot of the perpendicular from $\vec{P}$ to the line or plane.

Chapter 33

Algebraic Structures

Algebraic structures are mathematical systems consisting of a set equipped with one or more operations that satisfy certain axioms. They provide a unified language to study various objects in mathematics, from numbers and matrices to functions and vector spaces. Understanding these structures is fundamental in abstract algebra and has applications in computer science, cryptography, coding theory, and physics.

33.0.1 Operations: Internal and External

An *internal composition law* is a binary operation that takes two elements from a set and returns another element in the same set. Formally, for a set S and operation $\circ$, we have:

$$\oplus : S \times S \rightarrow S$$

An *external composition law* involves a second set acting on the structure, such as scalar multiplication in vector spaces:

$$\oplus : \mathbb{F} \times V \rightarrow V$$

where $\mathbb{F}$ is a field and V is a vector space.

33.0.2 Properties of Operations

Let $\odot$ and $\otimes$ be binary operations on a set S . The most important properties include:

- *Associativity*: $(a \odot b) \odot c = a \odot (b \odot c)$ for all $a, b, c \in S$
- *Commutativity*: $a \odot b = b \odot a$ for all $a, b \in S$
- *Identity Element*: There exists $e \in S$ such that $a \odot e = e \odot a = a$ for all $a \in S$
- *Inverse Element*: For every $a \in S$, there exists $a^{-1} \in S$ such that $a \odot a^{-1} = a^{-1} \odot a = e$
- *Distributivity*: $a \otimes (b \odot c) = (a \otimes b) \odot (a \otimes c)$ and/or $(b \odot c) \otimes a = (b \otimes a) \odot (c \otimes a)$

33.0.3 Homomorphisms and Isomorphisms

Let $(G, \oplus)$ and $(H, \oplus')$ be two algebraic structures.

A *homomorphism* is a function $\varphi : G \rightarrow H$ such that:

$$\varphi(a \oplus b) = \varphi(a) \oplus \varphi(b), \quad \forall a, b \in G$$

An *isomorphism* is a bijective homomorphism. If such a map exists, we say the structures are *isomorphic*, written as $G \cong H$.

33.0.4 Lattice

A *lattice* $(L, \wedge, \vee)$ is an algebraic structure with two binary, associative and commutative operations, which satisfies the following properties:

$$a \vee (a \wedge b) = a$$

and

$$a \wedge (a \vee b) = a$$

for all $a, b \in L$.

33.0.5 Semigroup $(S, \oplus)$

A *semigroup* is a set S equipped with a binary operation $\oplus$ that is *associative*. This means:

$$(a \oplus b) \oplus c = a \oplus (b \oplus c) \quad \text{for all } a, b, c \in S.$$

There is no requirement for an identity element or inverses. Semigroups capture the essence of combining elements consistently (e.g., string concatenation).

33.0.6 Monoid $(M, \oplus)$

A *monoid* builds on a semigroup by adding an *identity element* e such that:

$$a \oplus e = e \oplus a = a \quad \text{for all } a \in M.$$

This structure is useful when an operation must have a “do nothing” element, like 0 for addition or 1 for multiplication.

33.0.7 Group $(G, \oplus)$

A *group* is a monoid where every element has an *inverse*:

$$\text{For each } a \in G, \text{ there exists } a^{-1} \in G \text{ such that } a \oplus a^{-1} = a^{-1} \oplus a = e.$$

Groups model reversible processes and are foundational in symmetry and abstract algebra.

33.0.8 Abelian Group $(A, \oplus)$

An *Abelian group* (or commutative group) is a group where the operation is *commutative*:

$$a \oplus b = b \oplus a \quad \text{for all } a, b \in A.$$

This property makes Abelian groups especially important theory and linear algebra.

33.0.9 Ring $(R, \oplus, \odot)$

A *ring* is a set with two operations:

- $(R, \oplus)$ is an Abelian group.
- $(R, \odot)$ is a semigroup (associative multiplication).
- Multiplication distributes over addition:

$$a \odot (b \oplus c) = a \odot b \oplus a \odot c.$$

Rings generalize arithmetic of integers, where addition and multiplication interact in a structured way.

33.0.10 Commutative Ring $(R, \oplus, \odot)$

A *commutative ring* is a ring where multiplication is also *commutative*:

$$a \odot b = b \odot a \quad \text{for all } a, b \in R.$$

This is the kind of ring most often encountered in basic algebra, such as the integers $\mathbb{Z}$.

33.0.11 Field $(\mathbb{F}, \oplus, \odot)$

A *field* is a commutative ring where every nonzero element has a *multiplicative inverse*:

$$(\mathbb{F} \setminus \{0\}, \odot) \text{ is an Abelian group.}$$

Fields like $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$ allow division (except by zero), enabling the full range of arithmetic operations.

Field Proofs

Inverses are well-defined:

We know $a + -a = 0$. Now let us say $-a' \neq -a$ and $a - a' = 0$.

$$\begin{aligned} a + -a &= a - a' & | -a \\ -a &= -a' \end{aligned}$$

This contradicts our initial assumption, therefore, they are the same. Analog for the multiplication

$$\begin{aligned} a \cdot a^{-1} &= a \cdot a'^{-1} & | *a^{-1} \\ a^{-1} &= a'^{-1} \end{aligned}$$

□◻◻

Neutral Element for the Addition is well-defined:

We know $a + 0 = a$. Now let us say $0' \neq 0$ and $a + 0' = a$.

$$\begin{aligned} a + 0 &= a + (a - a) = a + 0' \\ a - a + a &= a + 0' \\ a &= a + 0' & | -a \\ 0 &= 0' \end{aligned}$$

There is a contradiction to the assumption that there are more than one neutral element for the addition. Thus, it is well-defined.

Neutral Element for the Multiplication is well-defined: This proof is very similar to the other, thus it will be skipped.

□◻◻

Minus times Minus is Plus:

We will prove that $ab = (-a)(-b)$ for some elements of a field.

$$\begin{aligned}
ab &= (-a)(-b) \\
ab + (-a)b &= (-a)(-b) + (-a)b \\
b(a + (-a)) &= (-a)((-b) + b) = 0
\end{aligned}$$

From that point, for the right side we can do the following

$$\begin{aligned}
(-a)((-b) + b) &= 0 \\
(-a)(-b) + (-a)b &= 0 \\
(-a)(-b) &= -(-a)b
\end{aligned}$$

Then, by looking on the left side of the original equality

$$\begin{aligned}
b(a + (-a)) &= 0 \\
ab + (-a)b &= 0 \\
ab &= -(-a)b
\end{aligned}$$

Thus, we have got our equality $ab = (-a)(-b)$.

□◻◻

33.0.12 Vector Space $(V, \oplus, \cdot)$

A *vector space* over a field $\mathbb{F}$ consists of:

- An Abelian group $(V, \oplus)$ for vector addition.
- A scalar multiplication $\cdot : \mathbb{F} \times V \rightarrow V$ satisfying:
 - Distributivity over vector addition: $a \cdot (v_1 + v_2) = a \cdot v_1 + a \cdot v_2$
 - Distributivity over field addition: $(a + b) \cdot v = a \cdot v + b \cdot v$
 - Associativity: $a \cdot (b \cdot v) = (ab) \cdot v$
 - Identity: $1 \cdot v = v$

Vector spaces provide the foundation for linear algebra, where scalars from a field combine with vectors from a set.

Chapter 34

Polynomial Rings

Given a field $\mathbb{F}$ with $\mathbb{F}^{\mathbb{Z}_{\geq 0}}$ we define the multiplication

$$(a_0, a_1, \dots) \cdot (b_0, b_1, \dots) := (c_0, c_1, \dots)$$

Where

$$c_0 := a_0 b_0, \quad c_1 := a_0 b_1 + a_1 b_0 \quad c_2 := a_0 b_2 + a_1 b_1 + a_2 b_0$$

In general for $n \geq 0$

$$c_n := \sum_{i=0}^n a_i b_{n-i} = a_0 b_n + a_1 b_{n-1} + \dots + a_n b_0$$

And the addition as

$$(a_i)_{i \in \mathbb{N}} + (b_i)_{i \in \mathbb{N}} = (a_i + b_i)_{i \in \mathbb{N}}$$

This $(\mathbb{F}^{\mathbb{Z}_{\geq 0}}, +, \cdot)$ is also noted as $\mathbb{F}[[\mathbb{F}]]$ and called the *power-ring* over $\mathbb{F}$. Also, instead of $(a_0, a_1, \dots)$ we write

$$f = \sum_{i=0}^{\infty} a_i X^i$$

For $X := (0, 1, 0, 0, \dots)$. We call this ring also the *formal power series* over $\mathbb{F}$.

The members $f = (a_0, a_1, \dots) \in \mathbb{F}[[X]]$ are called *polynomials*

For the finite case we write

$$f = \sum_{i=0}^n a_i X^i$$

The set of all polynomials reachable via the multiplication vector-space properties of $\mathbb{F}[[X]]$ is called the *polynomial ring* $\mathbb{F}[X]$. The field before the $[X]$ tells us which elements are allowed as coefficients.

Example:

For $f := (1, 2, 3, 4, \dots) = 1 + 2X + 3X^2 + 4X^3$

$$f \cdot 4 = (4, 1, 5, 2, 0, 0, \dots) = 4 + 1X + 5X^2 + 2X^3$$

$$f \cdot 3X = (0, 3, 6, 2, 5, 0, 0, \dots) = 3X + 6X^2 + 2X^3 + 5X^4$$

$$f \cdot 2X^2 = (0, 0, 2, 4, 6, 1, 0, 0, \dots) = 2X^2 + 4X^3 + 6X^4 + X^5$$

$$f \cdot X^3 = (0, 0, 0, 1, 2, 3, 4, 0, 0, \dots) = X^3 + 2X^4 + 3X^5 + 4X^6$$

34.0.1 Degree of a Polynomial

The *degree* of a polynomial is given by the n in $\sum_{i=0}^n a_i X^i$ with $f \neq \vec{0}$. Also, note that we define $\deg(0) := -\infty$

Degree Addition

For $f, g \in \mathbb{F}[X] \setminus \{0\}$ the following holds

$$\deg(f + g) \leq \max(\deg(f), \deg(g))$$

Degree Multiplication

For $f, g \in \mathbb{F}[X] \setminus \{0\}$ the following holds

$$\deg(fg) \leq \max(\deg(f) + \deg(g))$$

34.0.2 Properties of the Polynomial Ring

Neutral Element

$$1 := (1, 0, 0, \dots) \in \mathbb{F}[X]$$

is the neutral element of the multiplication in $\mathbb{F}[[X]]$ and $\mathbb{F}[X]$.

X_a

$$X_a = (0, a_0, a_1, \dots) \forall a \in \mathbb{F}[[X]]$$

Multiplication

$$X^i X^j = X^{i+j}, \text{ with } X^0 := 1$$

Identity

Given $f \in \mathbb{F}[X]$ with degree n then

$$f = a_0 + a_1 X + \dots + a_n X^n$$

gives a clear identification of $\mathbb{F}$ with

$$\{0\} \cup \{f \in \mathbb{F}[X] \mid \deg(f) = 0\}$$

Vector space

$$\mathbb{F}[X]_{\deg < n} := \{0\} \cup \{f \in \mathbb{F}[X] \mid \deg(f) < n\}$$

is a vector space.

34.0.3 Field Algebra

Given a ring R with the one of the vector space with the field $\mathbb{F}$. We call R an *associative algebra* with the one if only if

$$a(rs) = (ar)s = r(as)$$

for all $r, s \in R$ and $a \in \mathbb{F}$.

34.0.4 Division for Polynomial

Given the field $\mathbb{F}$ then

- $\mathbb{F}[X]$ is a commutative ring with one and exactly one commutative $\mathbb{F}$ -Algebra with 1.
- For $f, g \in \mathbb{F}[X]$ with $g \neq 0$ there are well-defined $q, r \in \mathbb{F}[X]$ with

$$f = qg + r \text{ and } \deg(r) < \deg(g) \vee r = 0$$

Proof:

For the first lemma, to show that the multiplication is commutative we just use the commutative property of commutativity of our field $\mathbb{F}$.

The distributive property for $f, g, h \in \mathbb{F}[X]$ and the scalar multiplication are also easy to show using our definitions. We will show the associativity.

We will use induction: If $\deg(h) = 0$, then $h \in \mathbb{F}$ and the associativity holds because of the associativity for scalars.

Let us assume that it holds for $\deg(h) \leq n$. If $\deg(h) > 0$ then

$$h = \hat{h} + a_{n+1}X^{n+1} \text{ with } \hat{h} \in \mathbb{F}[X]$$

is a polynomial of degree at most n or 0. Due to that fact, $(fg)\hat{h} = f(g\hat{h})$

Induction step:

$$\begin{aligned} (fg)h &= (fg)(\hat{h} + a_{n+1}X^{n+1}) \\ &= fg\hat{h} + fga_{n+1}X^{n+1} \\ &= fg\hat{h} + f(ga_{n+1}X^{n+1}) \\ &= f(g\hat{h} + ga_{n+1}X^{n+1}) \\ &= f(g(\hat{h} + a_{n+1}X^{n+1})) \\ &= f(gh) \end{aligned}$$

□

As for the second lemma, given $\deg(f) = m, \deg(g) = n, f = \sum_{i=0}^m a_i X^i, g = \sum_{i=0}^n b_i X^i$. If $f = 0$ then $q = 0$ and $r = 0$. If $m < n$ we are done, because $q = 0, r = f$.

For the case $m \geq n$ we will show the existence of q, r per induction using m . For $m = 0$ are $n = 0$, $f = a_0, g = b_0$. Set $q = \frac{a_0}{b_0}$ and $r = 0$, and we get $f = qg + r$.

If $m > 0$ and we assume that for each polynomial h with $\deg(h) \leq m - 1$ there exists polynomials $\hat{q}, \hat{r}$ such that $\deg(\hat{r}) < \deg(g)$ and $h = \hat{q}g + \hat{r}$.

Define $\hat{f} := f - \frac{a_m}{b_n} X^{m-n}g$. Then $\deg(\hat{f}) \leq m - 1$. There also exist $\hat{q}, \hat{r}$ with $\deg(\hat{r}) < \deg(g)$ and $\hat{f} = \hat{q}g + \hat{r}$. Then

$$f = \hat{f} + \frac{a_m}{b_n} X^{m-n} g = \hat{q}g + \hat{r} + \frac{a_m}{b_n} X^{m-n} g = \left(\hat{q} + \frac{a_m}{b_n} X^{m-n} \right) g + \hat{r}$$

Hence, with $q = \hat{q} + \frac{a_m}{b_n} X^{m-n}$ and $r = \hat{r}$, existence is shown.

For uniqueness: Let $f = qg + r$ and $f = \hat{q}g + \hat{r}$ with $\deg(r) < \deg(g)$ and $\deg(\hat{r}) < \deg(g)$. Then it follows that

$$r - \hat{r} = (q - \hat{q})g.$$

If $q - \hat{q} \neq 0$, then $\deg(r - \hat{r}) \geq \deg(g)$, which is a contradiction. Thus, $q = \hat{q}$ and $r = \hat{r}$.

□◻

34.0.5 Greatest Common Divisor for Polynomials

For two polynomials $f, g \in \mathbb{F}[X]$ with $(f, g) \neq (0, 0)$. The polynomial $t \in \mathbb{F}[X]$ with

- $t \mid f$ and $t \mid g$
- For all $c \in \mathbb{F}[X]$ with $c \mid f$ and $c \mid g$ holds $c \mid t$

is considered the *greatest common divisor* of f and g . If $t = 1$ then f, g are *relative prime*.

34.0.6 More Properties of Polynomials

- In $\mathbb{Z}$ one has $|ab| = |a||b|$; in $\mathbb{F}[X]$ one has $\deg(fg) = \deg(f) + \deg(g)$.
- The gcd of two polynomials $f, g \in \mathbb{F}[X]$ with $(f, g) \neq (0, 0)$ is unique only up to factors in $\mathbb{F}[X]^*$, that is, up to factors in $\mathbb{F}[X]$ of degree 0. Notation: $t = \gcd(f, g)$ actually also denotes every multiple at with $a \in \mathbb{F} \setminus \{0\}$ as a gcd of f and g .
- In both rings there is a division with remainder, and therefore $\mathbb{F}[X]$ also has a Euclidean algorithm for computing the gcd and an extended Euclidean algorithm for determining Bézout coefficients. Here the degree of a polynomial plays the role of the absolute value in $\mathbb{Z}$. We give the first part of the algorithm again.
- Furthermore, one also has the analogue of prime numbers in $\mathbb{F}[X]$, namely irreducible polynomials, and correspondingly a unique prime factorization for polynomials.

34.0.7 Euclidean Algorithm for Polynomials

Input: $f, g \in \mathbb{F}[X]$, $(f, g) \neq (0, 0)$.

Output: The greatest common divisor t of f and g .

Algorithm:

1. If $f = 0$, then return g . If $g = 0$, return f .
2. Set $r_1 := f$, $r_2 := g$, $k := 2$.
3. As long as $r_k \neq 0$, define $r_{k+1} := r_k$.
4. As soon as $\deg(r_k) = 0$, return r_{k-1} .

Example:

34.0.8 Reducible Irreducible

A polynomial $p \in \mathbb{F}[X] \setminus \{0\}$ is called *irreducible* if p is not a unit (that is, $p \notin \mathbb{F}[X]^*$) and if for $f, g \in \mathbb{F}[X]$ the following holds: $p = f \cdot g \Rightarrow f \in \mathbb{F}[X]^* \vee g \in \mathbb{F}[X]^*$. An element $h \in \mathbb{F}[X] \setminus \{0\}$ for which $h = f \cdot g$ holds with $f, g \in \mathbb{F}[X] \setminus \mathbb{F}[X]^*$ is called *reducible*.

34.0.9 Normalized

A polynomial $f \in \mathbb{F}[X] \setminus \{0\}$ with $f = \sum_{i=0}^n a_i X^i$ is considered *normalized* if $a_n = 1$.

34.0.10 Factorization Property

Each polynomial $f \in \mathbb{F}[X] \setminus \{0\}$ has from to one to the order of the factors unique factorizations in a normalized irreducible form and a unit.

Chapter 35

Vector Spaces

Given a field $\mathbb{F}$. A *vector space* is a set $(V, \mathbb{F}, \oplus, \odot)$ with two operations, vector addition and scalar multiplication, such that:

- I. $(a \oplus b)\vec{v} = a\vec{v} \oplus b\vec{v}, \forall a, b \in \mathbb{F} \text{ and } \vec{v} \in V$
- II. $(\vec{v} \oplus \vec{u})a = a\vec{v} \oplus a\vec{u}, \forall a \in \mathbb{F} \text{ and } \vec{v}, \vec{u} \in V$
- III. $(ab) \odot \vec{v} = a(b \odot \vec{v}), \forall a, b \in \mathbb{F} \text{ and } \vec{v} \in V$
- IV. $1 \odot \vec{v} = \vec{v}, \forall \vec{v} \in V$

The scalar multiplication is defined as

$$\mathbb{F} \times V \rightarrow V : (a, v) \mapsto a \odot v$$

Also using these axioms, we get

- $\vec{v} \odot 0 = \vec{0}$
- $(-1) \odot \vec{v} = -1 \odot \vec{v}$

Proof:

For the first property, $0\vec{v} = (0 + 0)\vec{v} = 0\vec{v} + 0\vec{v}$ then

$$0\vec{v} = 0\vec{v} + 0\vec{v} \implies \vec{0} = 0\vec{v}$$

For the second property,

$$\begin{aligned} (-1)\vec{v} &= (1 - 2)\vec{v} \\ (-1)\vec{v} &= \vec{v} - 2\vec{v} \\ (-1)\vec{v} &= -\vec{v} \end{aligned}$$

□

Examples:

1. The set of all n -tuples of real numbers $\mathbb{R}^n$ is a vector space over the field of real numbers $\mathbb{R}$.
2. The set of all polynomials of degree less than or equal to n is a vector space over the field of real numbers $\mathbb{R}$.
3. The set of all continuous functions from $\mathbb{R}$ to $\mathbb{R}$ is a vector space over the field of real numbers $\mathbb{R}$.
4. The set of all $m \times n$ matrices with real entries is a vector space over the field of real numbers $\mathbb{R}$.
5. The set of all linear maps is also a vector space.

35.0.1 Subspaces

A subset W of a vector space V is a *subspace* of V if:

- I. The zero vector $\vec{0} \in W$.
- II. For all $\vec{u}, \vec{v} \in W$, $\vec{u} + \vec{v} \in W$.
- III. For all $a \in F$ and $\vec{v} \in W$, $a\vec{v} \in W$.

If W is a subspace of V , we write $W \subseteq V$.

A subspace subset W of V is only a subspace if and only if $W \neq \emptyset$ and for all $v, w \in W$ and all $a \in \mathbb{F}$ then $v + aw \in W$

Proof:

$\Rightarrow$: $\vec{0} \in W$ thus $W \neq \emptyset$. Also, by the axioms $v + aw \in W$.

$\Leftarrow$: $W \neq \text{emptyset} \implies \exists w \in W$ because $\mathbb{F}$ is a field then $-1 \in \mathbb{F}$ and therefore, $w - w = 0 \in W$. For $v + aw \in W$ we choose $a = 1$ or $v = 0$, and we see that the addition and multiplication is also defined for our vector space.

35.0.2 Properties of the subspaces

- The intersection of two or more subspaces $\bigcap_{W \in V} W$ is a subspace.
- The union of two subspaces is not necessarily a subspace.
- The sum of two subspaces U and W is defined as:

$$U + W = \{\vec{u} + \vec{w} : \vec{u} \in U, \vec{w} \in W\}$$

The sum of two subspaces is a subspace.

35.0.3 Sum of Subspaces

- The *direct sum* of two subspaces U and W is defined as:

$$U \oplus W = \{\vec{u} + \vec{w} : \vec{u} \in U, \vec{w} \in W\}$$

The direct sum of two subspaces is a subspace.

- The direct sum of two subspaces is the smallest subspace containing both subspaces, such that $U \cap W = \{\vec{0}\}$.
- The direct sum of two subspaces is denoted by $U \oplus W$.

The sum of two subspaces is the smallest subspace containing both subspaces.

Vector Space Product

The *product* $\langle M \rangle$ of $M \subseteq V$ is the intersection of all subspaces of V which contain M

$$\langle M \rangle := \bigcap_{W \subseteq V, M \subseteq W} W$$

35.0.4 Linear Combinations

A *linear combination* of vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ in a vector space V is an expression of the form:

$$a_1\vec{v}_1 + a_2\vec{v}_2 + \dots + a_n\vec{v}_n$$

Where $a_1, a_2, \dots, a_n$ are scalars from the field F . The set of all linear combinations of a set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is called the **span** of those vectors, denoted by $\text{span}(\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n)$. The span of a set of vectors is a subspace of the vector space V .

The span of a set of vectors is the smallest subspace containing those vectors.

The set of all linear combinations of a set of vectors $B = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is given by $\mathcal{L}(B)$

35.0.5 Linear Independence

A set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ in a vector space V is said to be *linearly independent* if the only solution to the equation:

$$\lambda_1 \vec{v}_1 + \lambda_2 \vec{v}_2 + \dots + \lambda_n \vec{v}_n = 0$$

or

$$\sum_{i=1}^n \lambda_i \vec{v}_i = 0$$

is $\lambda_1 = \lambda_2 = \dots = \lambda_n = 0$. If there exists a non-trivial solution to this equation, then the set of vectors is said to be linearly dependent. A set of vectors is linearly independent if and only if the only linear combination of those vectors that equals the zero vector is the trivial combination where all coefficients are zero.

Properties of the linear independence

- A set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is linearly independent if and only if the only linear combination of those vectors that equals the zero vector is the trivial combination where all coefficients are zero.
- If a set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is linearly independent, then any subset of that set is also linearly independent.
- If a set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is linearly dependent, then at least one vector in that set can be expressed as a linear combination of the others.

35.0.6 Basis

- A *basis* of a vector space V is a set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ that is linearly independent and spans the vector space V .
- The number of vectors in a basis of a vector space is called the *dimension* of the vector space.
- The dimension of a vector space V is denoted by $\dim(V)$.
- If V has a finite basis, then it is said to be *finite-dimensional*.
- If V does not have a finite basis, then it is said to be *infinite-dimensional*.

Given $B = (v_1, \dots, v_n)$. B is a basis of V if and only if

$$\lambda = \lambda_B : \mathbb{F}^n \rightarrow V : (a_1, \dots, a_n) \mapsto \sum_{i=1}^n a_i v_i$$

is an isomorphism. This means that each vector in V can be uniquely represented as a linear combination of the basis vectors.

35.0.7 Dimension

The *dimension* of a vector space V is the number of vectors in a basis of V .

- The dimension of a vector space is denoted by $\dim(V)$.
- The dimension of a vector space can be finite or infinite.
- If the dimension of a vector space is finite, then it is said to be *finite-dimensional*.
- If the dimension of a vector space is infinite, then it is said to be *infinite-dimensional*.

35.0.8 How to find the basis of a set vector

To find the basis of a set of vectors, we can use the following steps:

1. Write the vectors as columns of a matrix.
2. Row-reduce the matrix to echelon form.
3. The non-zero rows of the echelon form matrix correspond to the basis of the vector space spanned by the original set of vectors.

The number of non-zero rows in the echelon form matrix is equal to the dimension of the vector space spanned by the original set of vectors.

The basis of a vector space is not unique. Different bases can span the same vector space.

35.0.9 Basis Extension Theorem

Let V be a vector space over a field K , and let

$$v_1, \dots, v_r, \quad w_1, \dots, w_s \in V.$$

Suppose that $(v_1, \dots, v_r)$ is a linearly independent tuple and that

$$\text{span}(v_1, \dots, v_r, w_1, \dots, w_s) = V.$$

Then it is possible to extend $(v_1, \dots, v_r)$ to a basis of V by possibly adding suitable vectors from the set $\{w_1, \dots, w_s\}$.

Proof:

If $\text{span}(v_1, \dots, v_r) = V$, the statement is obvious. So assume

$$\text{span}(v_1, \dots, v_r) \neq V.$$

Then there exists at least one w_i such that $w_i \notin \text{span}(v_1, \dots, v_r)$; otherwise, if all $w_i \in \text{span}(v_1, \dots, v_r)$, then

$$\text{span}(v_1, \dots, v_r, w_1, \dots, w_s) = \text{span}(v_1, \dots, v_r) = V,$$

which contradicts our assumption that $\text{span}(v_1, \dots, v_r) \neq V$.

The tuple $(w_i, v_1, \dots, v_r)$ is linearly independent, because of

$$\sum_{j=1}^r \lambda_j v_j + \lambda w_i = 0$$

It follows that $\lambda = 0$ (since $w_i \notin \text{span}(v_1, \dots, v_r)$), and then also $\lambda_j = 0$ for all j because the v_j are linearly independent.

Possibly, $(w_i, v_1, \dots, v_r)$ is still not a basis of V . Then we repeat the previous step and keep adding further w_i until the tuple extends $(v_1, \dots, v_r)$ to a basis of V . This process terminates after finitely many steps, since

$$\text{span}(v_1, \dots, v_r, w_1, \dots, w_s) = V.$$

□◻◻

Every finitely generated vector space V has a basis.

35.0.10 Epimorphism of Finite Dimensional Vector Spaces

A finite dimensional vector space is an *epimorphic* image of Field^n where n is the number of dimensions.

Proof:

Given $\{v_1, \dots, v_n\} \subseteq V$ a generating system of V and $X = (v_1, \dots, v_n) \in V^n$

$$\lambda : K^n \rightarrow V : a \mapsto a_1 X_1 + \dots + a_n X_n$$

35.0.11 Exchange Lemma

Let $(v_1, \dots, v_n)$ be a base of V and $(w_1, \dots, w_m)$ be generating system of V . Then, for every v_i , there exists a w_j such that if we replace v_i by w_j in the tuple $(v_1, \dots, v_n)$, it still forms a basis of V . We can exchange $m - n$ vectors in this way.

Proof I:

Let $(v_1, \dots, v_n)$ and $(w_1, \dots, w_m)$ be two bases of V . Suppose we remove v_i from the first basis. The truncated tuple $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n)$ satisfies

$$\text{span}(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n) \neq V,$$

because if $\text{span}(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n) = V$, then v_i would lie in the span of the remaining vectors and could be written as a linear combination of them. This would contradict the assumption that $(v_1, \dots, v_n)$ is linearly independent and a basis of V .

By the Basis Extension Theorem, we can extend the truncated tuple $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n)$ to a basis of V by adding vectors from $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n, w_1, \dots, w_m)$. Therefore, by the Basis Extension Theorem, there exists a w_j such that

$$w_j \notin \text{span}(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n),$$

and the tuple $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n, w_j)$ is linearly independent.

If this tuple does not form a basis, we can again apply the Basis Extension Theorem and add one of the vectors $v_1, \dots, v_n$ to complete the basis. Clearly, the only possibility is to add v_i , but this would imply that the tuple $(v_1, \dots, v_n, w_j)$ is not a basis, as w_j would then be linearly dependent on the other vectors. Therefore, $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_n, w_j)$ must form a basis of V .

□◻◻

Proof II:

Let $A = (v_1, \dots, v_n)$ and $B = (w_1, \dots, w_m)$ be two bases of V . We prove the statement by induction on m .

For $m = 1$ $A \cup B = (v_1, \dots, v_n, w_1)$ is linearly dependent because A is a basis of V . Thus, there exist $\lambda_1, \dots, \lambda_n = w_1$.

Choose $\lambda_j \neq 0$ and define

$$v_j = -\frac{1}{\lambda_j}(\lambda_1 v_1 + \dots + \lambda_{j-1} v_{j-1} + \lambda_{j+1} v_{j+1} + \dots + \lambda_n v_n - w_1)$$

If we remove v_j from A and add w_1 we get a basis of V . This is true because let

$$\tilde{\lambda}_1 v_1 + \dots + \lambda_{j-1} v_{j-1} + \tilde{\lambda}_j w_1 + \lambda_{j+1} v_{j+1} + \dots + \tilde{\lambda}_n v_n = 0$$

Assume $\tilde{\lambda}_j \neq 0$ then we can express w_1 as a linear combination of the other vectors. This is a contradiction because this would mean that we can write w_1 as another linear combination of the vectors in A which is not possible because B is a basis of V .

For we repeat this process $n - m$ times we get the desired result.

□◻◻

35.0.12 Dimension of a sum of subspaces

Let U and W be two subspaces of a vector space V . Then the dimension of the sum of the two subspaces is given by:

$$\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$$

35.0.13 Linear Independence of polynomials

Let P_n be the vector space of polynomials of degree at most n . The set of polynomials $\{1, x, x^2, \dots, x^n\}$ is a basis for P_n . The dimension of P_n is $n + 1$.

- The set of polynomials $\{1, x, x^2, \dots, x^n\}$ is linearly independent.
- The set of polynomials $\{1, x, x^2, \dots, x^n\}$ spans the vector space P_n .
- The dimension of P_n is $n + 1$.

To prove that the set of polynomials $\{1, x, x^2, \dots, x^n\}$ is linearly independent, we can use the following steps:

1. Assume that there exists a linear combination of the polynomials that equals zero:

$$a_0 + a_1x + a_2x^2 + \dots + a_nx^n = 0,$$

where $a_0, a_1, \dots, a_n$ are scalars.

2. Since the left-hand side is a polynomial of degree at most n , it can only be equal to zero if all coefficients are zero.
3. Therefore, we have $a_0 = a_1 = \dots = a_n = 0$, which proves that the set of polynomials $\{1, x, x^2, \dots, x^n\}$ is linearly independent.

So you only have to prove that the set of coefficients vector is linearly independent.

35.0.14 Interpolation Polynomial

Given the $n + 1$ points (x_k, y_k) , with $0 \leq k \leq n$ and all x_k distinct, there exists exactly one polynomial $p_n \in P_n$ such that $y_k = p_n(x_k)$ for all $0 \leq k \leq n$. This polynomial is called the interpolation polynomial.

Proof:

The uniqueness follows immediately. We prove the existence by induction on n . For $n = 0$, choose $p_0(x) = y_0$.

Now assume the statement is true for $n - 1$. Let the polynomial p_{n-1} interpolate the points $(x_0, y_0), \dots, (x_{n-1}, y_{n-1})$.

Define

$$p_n(x) = p_{n-1}(x) + q(x),$$

where

$$q(x) = \frac{(x - x_0)(x - x_1) \cdots (x - x_{n-1})}{(x_n - x_0)(x_n - x_1) \cdots (x_n - x_{n-1})} (y_n - p_{n-1}(x_n)).$$

We have $q \in P_n$, and it follows that $p_n \in P_n$. Furthermore, $q(x_k) = 0$ for $k \leq n - 1$ because a linear factor in the numerator always vanishes at x_k . Therefore, $p_n(x_k) = y_k$ for $k \leq n - 1$. Additionally, we have

$$q(x_n) = y_n - p_{n-1}(x_n),$$

so that $p_n(x_n) = y_n$.

□♦♦

Example:

Consider the three points $(-2, 1)$, $(-1, -1)$, and $(1, 1)$. These points uniquely define an interpolating parabola p_2 . This parabola can be determined using the definition of p_n . For hand calculations and a few points to interpolate, the following approach is also useful. The general form of the polynomial is

$$p_2(x) = ax^2 + bx + c.$$

Substituting the three points into this form gives the system of equations:

$$\begin{aligned} 1 &= a + b + c && \text{(from the point (1, 1))} \\ -1 &= a - b + c && \text{(from the point (-1, -1))} \\ 1 &= 4a - 2b + c && \text{(from the point (-2, 1))} \end{aligned}$$

This leads to the system of equations:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 4 & -2 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

Solving this system gives $a = 1$, $b = 1$, and $c = -1$, so the interpolation polynomial is

$$p_2(x) = x^2 + x - 1.$$

Chapter 36

Dot Product, Euclidean and Unitary Space

36.0.1 Outer Product

The *outer product* of two vectors $a \in \mathbb{F}^m$ and $b \in \mathbb{F}^n$ is defined as the matrix

$$a \otimes b := ab^T = \begin{bmatrix} a_1 b_1 & a_1 b_2 & \dots & a_1 b_n \\ a_2 b_1 & a_2 b_2 & \dots & a_2 b_n \\ \vdots & \vdots & \ddots & \vdots \\ a_m b_1 & a_m b_2 & \dots & a_m b_n \end{bmatrix} \in \mathbb{F}^{m \times n}.$$

36.0.2 Outer Product

The *outer product* of two vectors $a \in \mathbb{F}^m$ and $b \in \mathbb{F}^n$ is defined as the matrix

$$a \otimes b := ab^T = \begin{bmatrix} a_1 b_1 & a_1 b_2 & \dots & a_1 b_n \\ a_2 b_1 & a_2 b_2 & \dots & a_2 b_n \\ \vdots & \vdots & \ddots & \vdots \\ a_m b_1 & a_m b_2 & \dots & a_m b_n \end{bmatrix} \in \mathbb{F}^{m \times n}.$$

36.0.3 Inner Product

Let V be a vector space over a field $\mathbb{F}$. A mapping $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{F}$ is called an inner product (or inner product) if the following conditions are satisfied:

Symmetry

For all $a, b \in V$:

$$\langle a, b \rangle = \begin{cases} \langle b, a \rangle & \text{if } K = \mathbb{R}, \\ \overline{\langle b, a \rangle} & \text{if } K = \mathbb{C}. \end{cases}$$

Linearity in the First Argument

For all $a, b, c \in V$:

$$\langle a, b + c \rangle = \langle a, b \rangle + \langle a, c \rangle$$

and

$$\langle a + b, c \rangle = \langle a, c \rangle + \langle b, c \rangle.$$

Homogeneity in the First Argument

For all $\alpha \in \mathbb{F}$, we have:

$$\langle \alpha a, b \rangle = \alpha \langle a, b \rangle = \begin{cases} \langle a, \alpha b \rangle & \text{if } \mathbb{F} = \mathbb{R}, \\ \langle a, \alpha b \rangle & \text{if } \mathbb{F} = \mathbb{C}. \end{cases}$$

Positive Definiteness

For all $a \in V \setminus \{0\}$:

$$\langle a, a \rangle > 0,$$

and

$$\langle 0, 0 \rangle = 0.$$

It also can be interpreted as a linear transformation that takes a vector and maps it to a real number via matrix vector multiplication. Also, a more practical way of thinking about the *dot product* is the question: How much are two vectors pointing in the same direction?

36.0.4 Trigonometric Definition

Think about the orthogonal projection as a triangle. Now, remember that to find the adjacent side x you need to take $x = \cos(\theta)h$, which is the value α in our projection formula $p_a(b) = \frac{\langle a, b \rangle}{\langle a, a \rangle} \alpha a$. Here we add $\|a\|\|b\|$ to make this function linear.

$$\langle a, b \rangle = \|a\|\|b\|\cos(\theta)$$

36.0.5 Standard Inner Product for Complex Number

Let $a = (a_i)_{i=1}^n$ and $b = (b_i)_{i=1}^n$ be vectors in $\mathbb{C}^n$. The standard inner product is defined by

$$\langle a, b \rangle := \sum_{i=1}^n a_i \overline{b_i} = a^T \overline{b}$$

and by the *hermitian inner product*

$$\langle a, b \rangle := \overline{a^T b}$$

Here the *symmetry* is given by $\langle a, b \rangle = \overline{\langle b, a \rangle}$.

36.0.6 Inner Product on $C[a, b]$

Let $f, g \in C[a, b]$. The inner product on $C[a, b]$ is defined by

$$\langle f, g \rangle := \int_a^b f(x) \cdot g(x) dx.$$

36.0.7 Euclidean and Unitary Vector Spaces

A real vector space equipped with an inner product is called a *Euclidean vector space*, while a complex vector space with an inner product is called a *unitary vector space*.

36.0.8 Norms in Vector Spaces

Let V be a K -vector space and $a, b \in V$. A function $\|\cdot\| : V \rightarrow \mathbb{R}$ is called a norm if and only if the following conditions hold:

- $\|a\| \in \mathbb{R}$,
- $\|a\| \geq 0$,
- $\|a\| = 0 \iff a = 0$,
- $\forall \lambda \in K, \|\lambda a\| = |\lambda| \|a\|$,
- $\|a + b\| \leq \|a\| + \|b\|$.

Induced Norm by a Inner Product

As in the special case $V = \mathbb{R}^n$, an inner product induces a norm. In a unitary (or Euclidean) space, the inner product induces a (standard) norm defined by

$$\|\cdot\| = \sqrt{\langle \cdot, \cdot \rangle}.$$

36.0.9 Cauchy-Schwarz Inequality in Unitary Vector Spaces

In all unitary vector spaces V , the Cauchy-Schwarz inequality holds:

$$|\langle a, b \rangle| \leq \|a\| \|b\| \quad \forall a, b \in V.$$

Proof of the Triangle Inequality

Both sides of the triangle inequality are real and, in particular, non-negative. Therefore, it is sufficient to prove that the squares of both sides satisfy the desired inequality, i.e., we need to show:

$$\langle a + b, a + b \rangle \leq (\|a\| + \|b\|)^2.$$

First, we expand the left-hand side:

$$\langle a + b, a + b \rangle = \langle a, a \rangle + \langle a, b \rangle + \langle b, a \rangle + \langle b, b \rangle.$$

Since $\langle b, a \rangle = \overline{\langle a, b \rangle}$, we have:

$$\langle a, b \rangle + \langle b, a \rangle = 2 \operatorname{Re} \langle a, b \rangle.$$

Now, we know that the absolute value of a complex number is always greater than or equal to its real part, so:

$$2 \operatorname{Re} \langle a, b \rangle \leq 2 |\langle a, b \rangle|.$$

Using the Cauchy-Schwarz inequality, we can further bound this by:

$$2 \operatorname{Re} \langle a, b \rangle \leq 2 \|a\| \|b\|.$$

Thus, we have:

$$\langle a + b, a + b \rangle \leq \langle a, a \rangle + 2 \|a\| \|b\| + \langle b, b \rangle.$$

Using the definition of the norm, $\|a\|^2 = \langle a, a \rangle$ and $\|b\|^2 = \langle b, b \rangle$, we obtain:

$$\langle a + b, a + b \rangle \leq \|a\|^2 + 2\|a\|\|b\| + \|b\|^2.$$

This is exactly the expansion of $(\|a\| + \|b\|)^2$, which completes the proof.

□◻◻

36.0.10 Definiteness of Inner Products

The concept of the *inner product* plays a major role in linear algebra, and there are various manifestations of this element. We are going to see how inner products are constructed based on the properties we are interested in. Those being linearity for the columns, symmetry and positive definiteness.

Let us start with a matrix $A \in \mathbb{R}^{n \times n}$ and a map $(\cdot, \cdot)_A : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ more specific $(x, y)_A := \langle x, Ay \rangle$ using the standard Euclidean inner product.

For our first property, the linearity; it is granted for every matrix $A = (a_1, a_2, \dots, a_n)$ because of the rules of matrix-vector multiplication. For example for the basis vectors e_i, e_j we get

$$(e_i, e_j)_A = \langle e_i, Ae_j \rangle = e_i^T Ae_j = e_i^T a_j = a_{ij}$$

Therefore, $(e_i, e_j)_A = (e_j, e_i)_A \iff a_{ij} = a_{ji}$. This means that A has to be symmetric. As a side note, the choice for the Euclidean inner product for the structure of our map is optional. Any kind of inner product would also do the job. This implies that each inner product can be described via a matrix.

36.0.11 Definiteness and Quadratic Forms

Given a symmetric $A \in \mathbb{R}^{n \times n}$ then:

1. The map $x \rightarrow \langle x, Ax \rangle$ is called the *Quadratic Form*
2. A is *positive definite* if $\langle x, Ax \rangle > 0 \forall x \in \mathbb{R}^n \setminus \{0\}$
3. A is *negative definite* if $\langle x, Ax \rangle < 0 \forall x \in \mathbb{R}^n \setminus \{0\}$
4. A is *positive semidefinite* if $\langle x, Ax \rangle \geq 0 \forall x \in \mathbb{R}^n \setminus \{0\}$
5. A is *negative semidefinite* if $\langle x, Ax \rangle \leq 0 \forall x \in \mathbb{R}^n \setminus \{0\}$
6. A is *indefinite* if $\exists x, y \in \mathbb{R}^n \setminus \{0\} : \langle x, Ax \rangle < 0 \wedge \langle y, Ay \rangle > 0$

$(\cdot, \cdot)_A$ is only an inner product if A is symmetric and positive definite.

Example:

Given is the term $x_1^2 - x_3^2 + x_1x_4$, show that it is a quadratic form.

First write the formula for the quadratic form $\langle x, Ax \rangle$ where A is real symmetric matrix

$$\left\langle \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}, \begin{bmatrix} a & b & c & d \\ e & f & g & h \\ i & j & k & l \\ m & n & o & q \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \right\rangle$$

$$\left\langle \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}, \begin{bmatrix} ax_1 + bx_2 + cx_3 + dx_4 \\ ex_1 + fx_2 + gx_3 + hx_4 \\ ix_1 + jx_2 + kx_3 + lx_4 \\ mx_1 + nx_2 + ox_3 + px_4 \end{bmatrix} \right\rangle$$

$$x_1(ax_1 + bx_2 + cx_3 + dx_4) + x_2(ex_1 + fx_2 + gx_3 + hx_4) + x_3(ix_1 + jx_2 + kx_3 + lx_4) + x_4(mx_1 + nx_2 + ox_3 + px_4)$$

Now we have to compare the coefficients with the original expression $x_1^2 + (0)x_2^2 - x_3^2 + x_1x_4$.

$$\begin{aligned}
& ax_1^2 + bx_1x_2 + cx_1x_3 + dx_4x_4 + \\
& ex_2x_1 + fx_2^2 + gx_2x_3 + hx_2x_4 + \\
& ix_3x_1 + jx_3x_2 + kx_3^2 + lx_3x_4 + \\
& mx_4x_1 + nx_4x_2 + ox_4x_3 + px_4^2
\end{aligned}$$

Note that the only quadratic terms present in the expression are x_1^2 and $-x_3^2$ while the others are zero. This means that they have to disappear in the main diagonal. Also, the only term that x_1 is also multiplying is x_4 therefore all the others have to be zero like all other non-quadratic terms which are not present. Thus

$$\begin{bmatrix} 1 & 0 & 0 & d \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ m & 0 & 0 & 0 \end{bmatrix}$$

Now the question is what are d and m because A has to be symmetric, but they can not be 1 because then we would have $2x_1x_4$. Because of that, we will take the only available option $\frac{1}{2}$.

$$\begin{bmatrix} 1 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ \frac{1}{2} & 0 & 0 & 0 \end{bmatrix}$$

And now we are done.

36.1 Positive Definite Matrices

A matrix $A \in \mathbb{R}^{n \times n}$ is called *positive definite* if it is hermitian and if the quadratic form

$$\langle x, Ax \rangle > 0 = x^T Ax > 0, \quad \forall x \in \mathbb{R}^n, \quad x \neq 0.$$

Properties of Positive Definite Matrices

- All eigenvalues of A are positive.
- If A is symmetric, then A is positive definite if and only if all eigenvalues of A are positive.
- A is invertible and A^{-1} is also positive definite.
- The determinant of A is positive.
- The main diagonal entries of A are all positive.
- The quadratic form $\langle x, Ax \rangle$ is strictly convex.

36.1.1 Conditions for Positive Definiteness

A is positive definite if and only if one of the following conditions is satisfied:

$$A_s = \frac{1}{2} (A + A^T)$$

is also positive definite.

Proof:

A is positive definite then:

$$\langle x, Ax \rangle = \langle A_x, x \rangle = (Ax)^T x = x^T A^T x$$

Therefore,

$$x^T Ax = x^T (A + A^T) x = 2x^T A_s x$$

□◻

36.1.2 Minors

For a matrix $A \in \mathbb{R}^{n \times n}$ the starting from the left $k \times k$ -submatrices $A_k = (a_{ij})_{i,j=1}^k$ of A . The $D_k = \det(A_k)$ is called a *main minor* or *minor* of A .

Minors and Definiteness

Let A be real, square symmetric matrix, then

- A is positive definite if all main minors are positive.
- A is negative definite if all main minors are negative.
- A is positive semidefinite if all main minors are greater or equal to zero but not all zero.
- A is negative semidefinite if all main minors are less or equal to zero but not all zero.
- A is indefinite if all main minors are equal to zero.

36.1.3 Connection to Symmetric Matrices

For a real symmetric matrix A the following statements are equivalent.

- A is positive definite.
- All eigenvalues of A are positive.
- All main minors of A are positive.

Continuing, A is positive semidefinite, if all eigenvalues of A are not negative and the main minors are all not positive.

Example: Symmetry and Positive Definiteness of BAB^T and $B^T AB$

Let $A \in \mathbb{R}^{n \times n}$ be a symmetric matrix and $B \in \mathbb{R}^{m \times n}$ be a real matrix. We study the symmetry and positive definiteness of the matrices $C = BAB^T \in \mathbb{R}^{m \times m}$ and $D = B^T AB \in \mathbb{R}^{n \times n}$.

(i) Symmetry of BAB^T :

$$(BAB^T)^T = (B^T)^T A^T B^T = BAB^T,$$

since $(B^T)^T = B$ and $A^T = A$. Thus, BAB^T is symmetric.

(ii) Symmetry of $B^T AB$:

$$(B^T AB)^T = B^T A^T B = B^T AB,$$

since A is symmetric. Hence, $B^T AB$ is symmetric as well.

To determine when BAB^T and $B^T AB$ are positive definite, we use the quadratic form criterion.

(i) Positive definiteness of $C = BAB^T$:

Let $x \in \mathbb{R}^m$, $x \neq 0$. Consider:

$$x^T C x = x^T B A B^T x.$$

Let $y = B^T x \in \mathbb{R}^n$. Then:

$$x^T BAB^T x = y^T Ay.$$

Since A is positive definite, $y^T Ay > 0$ for all $y \neq 0$. Therefore, $x^T Cx > 0$ if and only if $B^T x \neq 0$ for all $x \neq 0$, which holds if and only if $\ker(B^T) = \{0\}$, or equivalently, B has full column rank.

(ii) Positive definiteness of $D = B^T AB$:

Let $z \in \mathbb{R}^n$, $z \neq 0$. Then:

$$z^T Dz = z^T B^T ABz = (Bz)^T A(Bz).$$

Let $w = Bz$. Since A is positive definite, $w^T Aw > 0$ if $w \neq 0$. Thus, D is positive definite if and only if $Bz \neq 0$ for all $z \neq 0$, i.e., if B has full rank (i.e., $\ker(B) = \{0\}$).

In summary:

- BAB^T is positive definite if B has full column rank.
- $B^T AB$ is positive definite if B has full rank.

□◻◻

36.1.4 Sign of the D_k

For the main minors D_k of a negative definite matrix it applies that

$$(-1)^k D_k = (-1)^k \det(A_k) = \det(-A_k) > 0,$$

because $-A$ is positive definite.

36.1.5 Negative Quadratic Form

If A is negative definite then

$$-\langle x, Ax \rangle = \langle x - Ax, x \rangle > 0$$

36.1.6 Theorems for Non-squares matrices

Theorem I

Given $m \geq n$ and $A \in \mathbb{R}^{m \times n}$ with $rg(A) = n$. Then $A^T A$ is positive definite. For $rg(A) < n$ is $A^T A$ positive semidefinite.

Proof:

Because of $(A^T A)^T = A^T A$ is $A^T A$ symmetric. Therefore, it applies that

$$\langle x, A^T Ax \rangle = x^T A^T Ax = (Ax)^T Ax = \langle Ax, Ax \rangle = \|Ax\|^2 \geq 0$$

Also, $A^T A$ is positive semidefinite, and it is also invertible for $rg(A) = n$. If there were an $x \neq 0$ such that $\|Ax\| = 0$ this would imply that $Ax = 0$, also $A^T(Ax) = 0$ therefore $x \in \ker(A^T A)$. And $A^T A$ would not be invertible. Finally, $\langle x, A^T Ax \rangle > 0 \forall x \neq 0$, and $A^T A$ is symmetric and positive definite.

36.1.7 Theorem II

Given $m \leq n$ and $A \in \mathbb{R}^{m \times n}$. Then $A^T A$ is positive semidefinite and even positive definite if $rg(A) = m$.

Chapter 37

Orthogonality

37.0.1 Orthogonality and Projection

Let $a, b \in V$. The vectors a and b are orthogonal to each other if

$$\langle a, b \rangle = 0.$$

This is written as $a \perp b$.

Also, the Pythagorean Theorem holds

$$\|a + b\|^2 = \|a\|^2 + \|b\|^2,$$

for $a \perp b$ in all unitary vector spaces.

For the orthogonal projection $p_b(a)$ of a vector a onto b , with $b \neq 0$, the formula in every unitary vector space is:

$$p_b(a) = \frac{\langle a, b \rangle}{\langle b, b \rangle} b.$$

37.0.2 Induced Metric

Given $(V, \|\cdot\|)$ a metric space there is an inner product $\langle \cdot, \cdot \rangle$ on $\mathbb{F}$ such that $\|x\|^2 = \langle x, x \rangle$ if and only if the *parallelepiped equation* holds.

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2), \quad \forall x, y \in V$$

In such a case, the inner product is given by

$$\langle x, y \rangle = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2)$$

Proof:

$\Rightarrow$: If $\|x\|^2 = \langle x, x \rangle$ then

$$\begin{aligned} \|x + y\|^2 + \|x - y\|^2 &= \langle x + y, x + y \rangle + \langle x - y, x - y \rangle \\ &= 2(\langle x, x \rangle + \langle y, y \rangle) \\ \langle x, x + y \rangle + \langle y, x + y \rangle + \langle x, x - y \rangle - \langle y, x - y \rangle &= 2(\langle x, x \rangle + \langle y, y \rangle) \\ \langle x, x \rangle + \langle x, y \rangle + \langle y, y \rangle + \langle x, y \rangle + \langle x, x \rangle - \langle x, y \rangle - \langle y, x \rangle + \langle y, y \rangle &= 2(\langle x, x \rangle + \langle y, y \rangle) \\ 2\langle x, x \rangle + 2\langle y, y \rangle &= 2(\langle x, x \rangle + \langle y, y \rangle) \end{aligned}$$

$\Leftarrow$: The parallelepiped equation holds, then want to show that the equation

$$f(x, y) = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2)$$

Satisfies the properties of a inner product:

- $f(x, y) \geq 0$ trivial.
- $f(x, y) = 0 \iff x = y = 0$ is also trivial and
- $f(x, y) = f(y, x)$ is also trivial by our definition.
- $f(\alpha x + y, z) = \alpha f(x, y) + f(y, z)$ Must be shown in a two step process.

First we show that $f(x + y, z) = f(x, z) + f(y, z)$:

$$\begin{aligned} 4(f(x, y) + f(z, y)) &= \|x + y\|^2 - \|x - y\|^2 + \|z + y\|^2 - \|z - y\|^2 \\ &= \|x + y\|^2 + \|z\|^2 \\ &\quad - \|x - y\|^2 - \|z\|^2 \\ &\quad + \|z + y\|^2 + \|x\|^2 \\ &\quad - \|z - y\|^2 - \|x\|^2 \end{aligned}$$

Hence, by adding this clever zero and using the parallelepiped equation we get

$$\begin{aligned} 2(f(x, y) + f(z, y)) &= \|x + y + z\|^2 + \|x + y - z\|^2 \\ &\quad - \|x - y + z\|^2 - \|x - y - z\|^2 \\ &\quad + \|z + y + x\|^2 + \|z + y - x\|^2 \\ &\quad - \|z - y + x\|^2 - \|z - y - x\|^2 \\ &= 2(\|x + z + y\|^2 - \|x + z - y\|^2) \\ &= 2f(x + z, y) \end{aligned}$$

Therefore, $f(x + z, y) = f(x, y) + f(z, y)$.

Now we need to prove that $f(\alpha x, y) = \alpha f(x, y)$:

For the Natural starting with $n = 2$: $f(2x, y) = f(x + x, y) = f(x, y) + f(x, y) = 2f(x, y)$, hence using induction for $n \in \mathbb{N}$ with base case 2 and summation, the prove is concluded.

For the Rationals, $r \in \mathbb{Q}$ with $r = \frac{m}{n}$ we have that

$$f(rx, y) = f\left(\frac{mx}{n}, y\right) = mf\left(\frac{1}{n}x, y\right) = \frac{m}{n}nf\left(\frac{1}{n}x, y\right) = \frac{m}{n}f(x, y)$$

Finally, for the Reals; each Real number can be approximated as a limit of an infinite sequence of Rationals.

For $\alpha \in \mathbb{R}$ there is $r_K \in \mathbb{Q}$ such that $\lim_{k \rightarrow \infty} r_k = \alpha$.

Given that f is continuous we can then use the following

$$f\left(\lim_{k \rightarrow \infty} r_K x, y\right) = \lim_{k \rightarrow \infty} f(r_K x, y) = \lim_{k \rightarrow \infty} r_k f(x, y) = \alpha f(x, y)$$

QED

37.0.3 Orthogonal Projection and Orthogonal Complement

Let U be a finitely generated subspace of V and $a \in V$. A vector $p_U(a) \in U$ is called the orthogonal projection of a onto U if

$$a - p_U(a) \perp u \quad \forall u \in U$$

The question arises about well-definedness, i.e., whether such a vector $p_U(a)$ always exists and if it is unique. The following concept helps in the discussion of uniqueness.

For $M \subseteq V$, the *orthogonal complement* of M is defined as

$$M^\perp = \{v \in V \mid v \perp u \forall u \in M\}.$$

- $M^\perp$ is a subspace of V .
- Let U be a subspace of V . Then, we have $U \cap U^\perp = \{0\}$.

Proof:

We need to check the closure property. For $u \in M$, $x, y \in M^\perp$, and $\lambda \in \mathbb{R}$, we have:

$$\begin{aligned} \langle x + y, u \rangle &= \langle x, u \rangle + \langle y, u \rangle = 0 \\ \langle \lambda x, u \rangle &= \lambda \langle x, u \rangle = 0. \end{aligned}$$

Let $a \in U \cap U^\perp$. Then, we have $\langle a, u \rangle = 0$ for all $u \in U$, because $a \in U^\perp$, and in particular, $\langle a, a \rangle = 0$ since $a \in U$, which implies $a = 0$.

37.0.4 Orthogonality of the basis to the Complement

Let U be as before, and let $(u_1, \dots, u_m)$ be a basis of U . For $v \in V$, we have:

$$v \in U^\perp \quad \text{if and only if} \quad \langle v, u_i \rangle = 0 \quad \forall 1 \leq i \leq m.$$

37.0.5 How to find the orthogonal complement

Let U be a finitely generated subspace of V with basis $(u_1, \dots, u_m)$. To find the orthogonal complement $U^\perp$, we can solve the system of equations:

$$\langle v, u_i \rangle = 0 \quad \forall 1 \leq i \leq m.$$

This system can be expressed in matrix form as $A \cdot v = 0$, where A is the matrix whose rows are the vectors u_i and v is the vector we want to find in $U^\perp$.

37.0.6 Orthogonal and Orthonormal Systems

Let $B = (v_1, \dots, v_m)$ be an m -tuple of vectors in $V \setminus \{0\}$.

- B is called an orthogonal system in V if all the vectors v_i are pairwise orthogonal.
- An orthogonal system is called an orthonormal system if, in addition, $\|v_i\| = 1$ for all $i = 1, \dots, m$.
- An orthogonal system that forms a basis of V is called an orthogonal basis of V .
- An orthonormal system that forms a basis of V is called an orthonormal basis of V .

Notation:

$$\text{OG-System} \quad \text{OG-Basis} \quad \text{ON-System} \quad \text{ON-Basis}$$

Using the Kronecker delta symbol $\delta_{i,j}$:

$$\delta_{i,j} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j, \end{cases}$$

we have $\langle v_i, v_j \rangle = \delta_{i,j}$ for any orthonormal system.

37.0.7 Writing a vector in terms of the Orthogonal Basis

Example:

The vectors

$$a_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \quad a_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

are orthogonal and normalized, but do not form a basis. Thus, they constitute an *orthonormal system*.

Example:

The vectors

$$a_1 = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}, \quad a_2 = \begin{bmatrix} -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}, \quad a_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

form a basis, are orthogonal and normalized. Therefore, they form an *orthonormal basis*.

37.0.8 Linear Independence of Orthogonal Systems

An orthogonal system $(v_1, \dots, v_n)$ is linearly independent.

Proof:

Let $\sum_{i=1}^n \lambda_i v_i = 0$. Then, for any v_j , it follows that

$$\left\langle \sum_{i=1}^n \lambda_i v_i, v_j \right\rangle = \sum_{i=1}^n \lambda_i \langle v_i, v_j \rangle = 0.$$

Due to the orthogonality of the system, all terms vanish except $\lambda_j \langle v_j, v_j \rangle$, so

$$\lambda_j \langle v_j, v_j \rangle = 0.$$

Since $v_j \neq 0$ in any orthogonal system, we have $\langle v_j, v_j \rangle = \|v_j\|^2 > 0$, implying $\lambda_j = 0$ for all $1 \leq j \leq n$.

□

37.0.9 Representation with Respect to an Orthogonal Basis

Let $B = (v_1, \dots, v_n)$ be an orthogonal basis of a vector space V . Then for every $v \in V$, we have:

$$v = \sum_{k=1}^n \frac{\langle v, v_k \rangle}{\langle v_k, v_k \rangle} v_k,$$

that is, the coordinates of v with respect to the basis B are given by

$$\left(\frac{\langle v, v_k \rangle}{\|v_k\|^2} \right)_{1 \leq k \leq n}^T.$$

37.0.10 Orthogonal Projection and Direct Sum Decomposition

Let $B = (v_1, \dots, v_m)$ be an orthogonal system in V , and let $U = L(B)$ be the subspace of V spanned by B .

- For every $v \in V$, the orthogonal projection of v onto U is given by:

$$p_U(v) = \sum_{i=1}^m \frac{\langle v, v_i \rangle}{\langle v_i, v_i \rangle} v_i.$$

- Every vector $v \in V$ can be uniquely written as the sum $v = p_U(v) + w$, with $w \in U^\perp$. In this case, we have:

$$w = v - p_U(v).$$

- $V = U \oplus U^\perp$, which means that every vector in V can be uniquely decomposed into a sum of a vector from U and a vector from $U^\perp$.
- If $\dim(V) = n$, then:

$$\dim(U) + \dim(U^\perp) = n \quad \text{for every subspace } U.$$

37.0.11 The Gram-Schmidt Process

The Gram-Schmidt process is a method for orthonormalizing a set of vectors in an inner product space. Given a finite set of linearly independent vectors $(v_1, \dots, v_n)$, the process generates an orthonormal basis $(u_1, \dots, u_n)$ as follows:

1. Set $u_1 = \frac{v_1}{\|v_1\|}$.
2. For $k = 2, \dots, n$:
 - (a) Set $w_k = v_k - \sum_{j=1}^{k-1} \langle v_k, u_j \rangle u_j$.
 - (b) Set $u_k = \frac{w_k}{\|w_k\|}$.
3. The resulting set $(u_1, \dots, u_n)$ is an orthonormal basis of the subspace spanned by $(v_1, \dots, v_n)$.

The Gram-Schmidt process can be applied to any finite set of linearly independent vectors in an inner product space, and it is particularly useful for constructing orthonormal bases in Euclidean spaces.

Proof:

Let W be a non-zero finite-dimensional subspace of an inner product space and let $(u_1, \dots, u_n)$ be a basis for W . We will show that an *orthonormal basis* exists.

Let $v_1 = u_1$, then compute the vector u_2 orthogonal to the space W_1 spanned by v_1

$$v_2 = u_2 - \text{proj}_{w_1} u_2 = u_2 - \frac{\langle u_2, v_1 \rangle}{\|v_1\|} v_1$$

Then normalize the vector.

We can continue this approach to of projecting the vector of u_i orthogonal to the space W_i spanned by all basis previously computed v_i basis vectors and then normalize it until we get our orthonormal basis. Note, that because our previous vectors formed a basis we can be sure that during this process no zero vector will be generated because then we will have some u_i that is a linear combination of the other vectors. Thus, this would be a contradiction.

□◻◻

Example:

Let $v_1 = (1, 0, 0)$, $v_2 = (1, 1, 0)$, and $v_3 = (1, 1, 1)$. The Gram-Schmidt process yields:

$$u_1 = \frac{v_1}{\|v_1\|} = (1, 0, 0), \quad u_2 = \frac{v_2 - \langle v_2, u_1 \rangle u_1}{\|v_2 - \langle v_2, u_1 \rangle u_1\|} = (0, 1, 0),$$

$$u_3 = \frac{v_3 - \langle v_3, u_1 \rangle u_1 - \langle v_3, u_2 \rangle u_2}{\|v_3 - \langle v_3, u_1 \rangle u_1 - \langle v_3, u_2 \rangle u_2\|} = (0, 0, 1).$$


Thus, the orthonormal basis is $(1, 0, 0), (0, 1, 0), (0, 0, 1)$.

37.0.12 Existence of Orthonormal Bases and Orthogonal Projections

- Every finitely generated unitary vector space has an orthonormal basis.
- This is in general false for vector spaces that are not finitely generated.
- We now provide the existence proof of the orthogonal projection.
- Let V be a unitary vector space and U a finitely generated subspace. Then for every $v \in V$, the orthogonal projection $p_U(v)$ of v onto U exists.
- Let V be a finitely generated unitary vector space and U any subspace. Then we have $V = U \oplus U^\perp$, and

$$\dim(V) = \dim(U) + \dim(U^\perp).$$

- Every hyperplane in $\mathbb{R}^n$ admits a normal form; the normal vector is unique up to scalar multiplication.
- Let V be as above and let $v_1, \dots, v_m \in V$. If it is possible to construct orthonormal vectors $w_1, \dots, w_m$ from them using the Gram-Schmidt process, then $(v_1, \dots, v_m)$ are linearly independent.

Visualization of the Gram-Schmidt process in 2 dimension

37.0.13 Best Approximation

Let V be a unitary vector space and U a finitely generated subspace of V . A vector $v* \in U$ is called Best Approximation in U on v , if:

$$\|v* - v\| = \inf_{x \in U} \|x - v\|$$

also

$$\|v* - P_u(v)\| = \inf_{u \in U} \|u - v\|$$

Proof:

Let $x \in V$, v be some point and $v* = \text{proj}_V v$. Then

$$v - x = v - v* + v* - x$$

where $v - v* \in V^\perp$ and $v* - x \in V$. By the Pythagorean Theorem

$$\|x - v\|^2 = \|v* - v\|^2 + \|v* - x\|^2$$

Therefore

$$\|v* - v\|^2 \leq \|v* - x\|^2$$

□

Chapter 38

Matrices

In this section, we explore the fundamental concepts of matrices, their operations, and important properties that form the foundation of linear algebra.

38.0.1 Definition of a Matrix

A matrix is a rectangular array of numbers, symbols, or expressions arranged in rows and columns. Formally, an $m \times n$ matrix A consists of mn elements a_{ij} where $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$:

Formally

$$A : m \times n \rightarrow \mathbb{F} : (i, j) \mapsto a_{i,j}$$

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

The set of all $m \times n$ matrices with real entries is denoted by $\mathbb{R}^{m \times n}$. Special cases include:

- Square matrix: matrix with the same number of rows and columns ($m = n$)
- Column vector: $m \times 1$ matrix
- Row vector: $1 \times n$ matrix
- Identity matrix I_n : An $n \times n$ matrix with ones on the main diagonal and zeros elsewhere
- Zero matrix: A matrix where all entries are zero

38.0.2 Matrix Addition and Subtraction

Matrix addition and subtraction are defined for matrices of the same dimensions.

$$c_{ij} = a_{ij} \pm b_{ij} \quad \text{for all } i = 1, 2, \dots, m \text{ and } j = 1, 2, \dots, n$$

Matrix addition/subtraction satisfies the following properties:

$$\begin{aligned} A \pm B &= B \pm A && \text{(Commutativity)} \\ (A \pm B) + C &= A \pm (B \pm C) && \text{(Associativity)} \\ A \pm O &= A && \text{(Identity element)} \\ A + (-A) &= O && \text{(Inverse element)} \end{aligned}$$

Where O is the zero matrix.

38.0.3 Matrix Multiplication

Matrix multiplication is defined between matrices where the number of columns in the first matrix equals the number of rows in the second matrix.

For $A \in \mathbb{C}^{m \times p}$ and $B \in \mathbb{C}^{p \times n}$, their product $C = AB \in \mathbb{C}^{m \times n}$ is defined as:

$$c_{ij} = \sum_{k=1}^p a_{ik}b_{kj} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{ip}b_{pj}$$

Matrix multiplication is defined this way to think as each operation as dot product of two vectors.

An example of this is thinking about the dot product as the total price of a purchase in store. The total price is the sum of the amounts which are given by the number of items per price. Now to extend this idea for matrices we think about the number of items as a vector of size i and the prices of each item as another vector p_1 with dot product $\langle i, p_1 \rangle$. If we want to by other items with different prices, but in the same amounts given by the vector i we get a matrix $A = (p_1, \dots, p_n)$. Now it makes sense that, if we take three different arrays of products we get a vector of three components which represent each of the total amounts to pay.

Thus, each operation in matrix multiplication is taking the dot product of the *transpose* of the current row with the column vector of the other matrix.

$$A \times \vec{v} = \begin{bmatrix} \langle a_1^T, v \rangle \\ \langle a_2^T, v \rangle \\ \vdots \\ \langle a_n^T, v \rangle \end{bmatrix}$$

As for the case with multiple columns

$$A \times B = \begin{bmatrix} \langle a_1^T, b_j \rangle & \dots & \langle a_1^T, b_n \rangle \\ \vdots & \dots & \vdots \\ \langle a_n^T, b_j \rangle & \dots & \langle a_n^T, b_n \rangle \end{bmatrix}$$

Matrix multiplication satisfies the following properties:

$$\begin{aligned} A(BC) &= (AB)C && \text{(Associativity)} \\ A(B+C) &= AB+AC && \text{(Left distributivity)} \\ (A+B)C &= AC+BC && \text{(Right distributivity)} \\ AI_n &= A \quad \text{and} \quad I_m A = A && \text{(Identity)} \end{aligned}$$

Note that matrix multiplication is generally not commutative, i.e., $AB \neq BA$ in most cases.

Example:

Consider the matrices A and B given by:

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & 0 & -2 \end{bmatrix} \in \mathbb{R}^{2 \times 3} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 \\ -1 & 3 \\ 4 & 0 \end{bmatrix} \in \mathbb{R}^{3 \times 2}$$

To compute the product $C = AB \in \mathbb{R}^{2 \times 2}$, we calculate each entry c_{ij} using the formula:

$$c_{ij} = \sum_{k=1}^3 a_{ik} b_{kj}$$

Let's calculate each entry of C :

$$\begin{aligned} c_{11} &= a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31} \\ &= 2 \cdot 1 + 3 \cdot (-1) + 1 \cdot 4 \\ &= 2 - 3 + 4 = 3 \end{aligned}$$

$$\begin{aligned} c_{12} &= a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} \\ &= 2 \cdot 2 + 3 \cdot 3 + 1 \cdot 0 \\ &= 4 + 9 + 0 = 13 \end{aligned}$$

$$\begin{aligned} c_{21} &= a_{21}b_{11} + a_{22}b_{21} + a_{23}b_{31} \\ &= 1 \cdot 1 + 0 \cdot (-1) + (-2) \cdot 4 \\ &= 1 + 0 - 8 = -7 \end{aligned}$$

$$\begin{aligned} c_{22} &= a_{21}b_{12} + a_{22}b_{22} + a_{23}b_{32} \\ &= 1 \cdot 2 + 0 \cdot 3 + (-2) \cdot 0 \\ &= 2 + 0 + 0 = 2 \end{aligned}$$

Therefore, the product $C = AB$ is:

$$C = AB = \begin{bmatrix} 3 & 13 \\ -7 & 2 \end{bmatrix}$$

Let's verify that matrix multiplication is not generally commutative by attempting to compute BA :

Since B is a 3×2 matrix and A is a 2×3 matrix, the product BA would be a 3×3 matrix. However, this calculation cannot be performed since the number of columns in B (which is 2) does not equal the number of rows in A (which is 2). Thus, BA is undefined, demonstrating that matrix multiplication is not always commutative.

38.0.4 Outer Product for Matrix Multiplication

The *outer product* of two vectors $a \in \mathbb{F}^m$ and $b \in \mathbb{F}^n$ is defined as the matrix

$$a \otimes b := ab^T = \begin{bmatrix} a_1b_1 & a_1b_2 & \dots & a_1b_n \\ a_2b_1 & a_2b_2 & \dots & a_2b_n \\ \vdots & \vdots & \ddots & \vdots \\ a_mb_1 & a_mb_2 & \dots & a_mb_n \end{bmatrix} \in \mathbb{F}^{m \times n}.$$

For the case where $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$

$$A \times B = \sum_{k=1}^n a_k \otimes b_k = \sum_{k=1}^n a_k b_k^T$$

where a_k is the k -th column of A and b_k is the k -th row of B .

38.0.5 The Transpose of a Matrix

The transpose of a matrix $A \in \mathbb{R}^{m \times n}$, denoted $A^T \in \mathbb{R}^{n \times m}$, is obtained by interchanging rows and columns:

$$(A^T)_{ij} = a_{ji} \quad \text{for all } i = 1, 2, \dots, n \text{ and } j = 1, 2, \dots, m$$

Properties of the Transpose

$$\begin{aligned} (A^T)^T &= A \\ (A + B)^T &= A^T + B^T \\ (AB)^T &= B^T A^T \\ (\alpha A)^T &= \alpha A^T \quad \text{for any scalar } \alpha \\ (A^{-1})^T &= (A^T)^{-1} \\ \text{rank}(A) &= \text{rank}(A^T) \\ \det(A) &= \det(A^T) \\ \text{eig}(A) &= \text{eig}(A^T) \text{ where } \text{eig}() \text{ are the eigenvalues} \end{aligned}$$

The transpose can also help us interpret the dot product of two vectors being 0 as a matrix multiplication. Here are given $A = (a_1, \dots, a_n)$ with Ax as some vector and b some other vector and a_k is some row of A .

$$\langle a_k, b - Ax \rangle = 0 \quad \forall k \iff A^T(b - Ax) = \vec{0}$$

Proof that the inverse of the transpose is the transpose of the inverse

To prove: $(A^{-1})^T = (A^T)^{-1}$

$$\begin{aligned} (A^{-1})^T A^T &= (A^T)^{-1} A^T \\ (AA^{-1})^T &= (A^T)^{-1} A^T \\ I^T &= I \end{aligned}$$

□◻◻

38.0.6 The Equivalence of Matrices

Two matrices A and B are said to be equivalent if one can be transformed into the other through a finite sequence of elementary row operations. We write $A \sim B$ to denote this equivalence.

The elementary row operations are:

- Interchanging two rows: $R_i \leftrightarrow R_j$
- Multiplying a row by a non-zero scalar: $R_i \mapsto \alpha R_i$ where $\alpha \neq 0$
- Adding a multiple of one row to another: $R_i \mapsto R_i + \alpha R_j$ where $i \neq j$

Matrix equivalence is an equivalence relation, satisfying reflexivity, symmetry, and transitivity. Equivalent matrices represent the same linear system in different bases.

Row Echelon Form (REF)

A matrix is in row echelon form if:

- All rows consisting entirely of zeros are at the bottom of the matrix.

- The leading entry (first non-zero element) of each non-zero row is to the right of the leading entry of the row above it.
- All entries in a column below a leading entry are zeros.

Reduced Row Echelon Form (RREF)

A matrix is in reduced row echelon form if:

- It is in row echelon form.
- Each leading entry is 1.
- Each leading entry is the only non-zero entry in its column.

The Gauss-Jordan elimination algorithm proceeds as follows:

- Start with the leftmost non-zero column.
- Find the pivot (non-zero element) in this column. If necessary, swap rows to move a non-zero element to the pivot position.
- Divide the pivot row by the pivot value to make the pivot equal to 1.
- Eliminate all other entries in the pivot column by subtracting appropriate multiples of the pivot row.
- Cover the pivot row and column, and repeat steps 1-4 on the submatrix until all rows are processed.
- For RREF, eliminate all entries above each pivot as well.

Example:

We'll use Gauss-Jordan elimination to solve the linear system:

$$\begin{aligned} 2x + y - z &= 8 \\ -3x - y + 2z &= -11 \\ x + y + z &= 3 \end{aligned}$$

First, we set up the augmented matrix:

$$\left[\begin{array}{ccc|c} 2 & 1 & -1 & 8 \\ -3 & -1 & 2 & -11 \\ 1 & 1 & 1 & 3 \end{array} \right]$$

Now we apply Gauss-Jordan elimination to transform this into reduced row echelon form:

Step 1: We'll choose the first element in the first row as our pivot. Let's first swap row 1 and row 3 to get a simpler pivot:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ -3 & -1 & 2 & -11 \\ 2 & 1 & -1 & 8 \end{array} \right]$$

Step 2: Eliminate the first elements in rows 2 and 3:

Row 2 + 3 × Row 1:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 2 & 5 & -2 \\ 2 & 1 & -1 & 8 \end{array} \right]$$

Row 3 - 2 × Row 1:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 2 & 5 & -2 \\ 0 & -1 & -3 & 2 \end{array} \right]$$

Step 3: Make the pivot in row 2 equal to 1 by dividing the entire row by 2:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 1 & \frac{5}{2} & -1 \\ 0 & -1 & -3 & 2 \end{array} \right]$$

Step 4: Eliminate the second element in rows 1 and 3:

Row 1 - Row 2:

$$\left[\begin{array}{ccc|c} 1 & 0 & -\frac{3}{2} & 4 \\ 0 & 1 & \frac{5}{2} & -1 \\ 0 & -1 & -3 & 2 \end{array} \right]$$

Row 3 + Row 2:

$$\left[\begin{array}{ccc|c} 1 & 0 & -\frac{3}{2} & 4 \\ 0 & 1 & \frac{5}{2} & -1 \\ 0 & 0 & -\frac{1}{2} & 1 \end{array} \right]$$

Step 5: Make the pivot in row 3 equal to 1 by multiplying the entire row by -2:

$$\left[\begin{array}{ccc|c} 1 & 0 & -\frac{3}{2} & 4 \\ 0 & 1 & \frac{5}{2} & -1 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

Step 6: Eliminate the third element in rows 1 and 2:

Row 1 + $\frac{3}{2} \times$ Row 3:

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & \frac{5}{2} & -1 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

Row 2 - $\frac{5}{2} \times$ Row 3:

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 + 5 = 4 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

The matrix is now in reduced row echelon form:

$$\left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 4 \\ 0 & 0 & 1 & -2 \end{array} \right]$$

This corresponds to the system:

$$\begin{aligned} x &= 1 \\ y &= 4 \\ z &= -2 \end{aligned}$$

Therefore, the solution to the original system is $x = 1$, $y = 4$, and $z = -2$.

38.0.7 The Inverse of a Matrix

For a square matrix $A \in \mathbb{R}^{n \times n}$, the inverse matrix A^{-1} (if it exists) satisfies:

$$AA^{-1} = A^{-1}A = I_n$$

Properties of the Inverse

$$(A^{-1})^{-1} = A$$

$$(AB)^{-1} = B^{-1}A^{-1}$$

$$(A^T)^{-1} = (A^{-1})^T$$

$$\det(A^{-1}) = \frac{1}{\det(A)}$$

$$(cA)^{-1} = \frac{1}{c}A^{-1}$$

$$\det(A^{-1}) = \det(A)^{-1}$$

$$\text{rank}(A^{-1}) = \text{rank}(A)$$

$$\text{eig}(A^{-1}) = \text{eig}(A)^{-1}$$

Finding the Inverse

There are several methods to find the inverse of a matrix:

Gauss-Jordan Method

Form the augmented matrix $[A|I_n]$ and apply Gauss-Jordan elimination to transform it into $[I_n|A^{-1}]$:

1. Create the augmented matrix $[A|I_n]$
2. Apply row operations to transform the left side into I_n
3. The right side will be A^{-1}

Adjoint Method

For an $n \times n$ matrix A :

$$A^{-1} = \frac{1}{\det(A)} \text{adj}(A)$$

Where $\text{adj}(A)$ is the adjoint (or adjugate) of A , defined as the transpose of the co-factor matrix.

A matrix is invertible if and only if its determinant is non-zero. Such matrices are called non-singular or regular matrices.

Example:

Let's find the inverse of the matrix:

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$$

We form the augmented matrix $[A|I_3]$:

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Now we apply Gauss-Jordan elimination:

Step 1: Make the first pivot equal to 1 by dividing the first row by 2:

$$\left[\begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Step 2: Eliminate the first element in rows 2 and 3:

Row 2 - 3 × Row 1:

$$\left[\begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{3}{2} & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Row 3 - 2 × Row 1:

$$\left[\begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{1}{2} & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

Step 3: Make the second pivot equal to 1 by multiplying the second row by 2:

$$\left[\begin{array}{ccc|ccc} 1 & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ 0 & 1 & -1 & -3 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

Step 4: Eliminate the second element in row 1 and the third element in row 2:

Row 1 - $\frac{1}{2}$ × Row 2:

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & -1 & -3 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

Row 2 + Row 3:

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & 0 & -4 & 2 & 1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

Step 5: Eliminate the third element in row 1:

Row 1 - Row 3:

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & -1 & -1 \\ 0 & 1 & 0 & -4 & 2 & 1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right]$$

The right side of the augmented matrix now gives us A^{-1} :

$$A^{-1} = \begin{bmatrix} 3 & -1 & -1 \\ -4 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$$

To verify, we can check that $AA^{-1} = I_3$:

$$\begin{aligned}
AA^{-1} &= \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix} \begin{bmatrix} 3 & -1 & -1 \\ -4 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix} \\
&= \begin{bmatrix} 2 \cdot 3 + 1 \cdot (-4) + 1 \cdot (-1) & 2 \cdot (-1) + 1 \cdot 2 + 1 \cdot 0 & 2 \cdot (-1) + 1 \cdot 1 + 1 \cdot 1 \\ 3 \cdot 3 + 2 \cdot (-4) + 1 \cdot (-1) & 3 \cdot (-1) + 2 \cdot 2 + 1 \cdot 0 & 3 \cdot (-1) + 2 \cdot 1 + 1 \cdot 1 \\ 2 \cdot 3 + 1 \cdot (-4) + 2 \cdot (-1) & 2 \cdot (-1) + 1 \cdot 2 + 2 \cdot 0 & 2 \cdot (-1) + 1 \cdot 1 + 2 \cdot 1 \end{bmatrix} \\
&= \begin{bmatrix} 6 - 4 - 1 & -2 + 2 + 0 & -2 + 1 + 1 \\ 9 - 8 - 1 & -3 + 4 + 0 & -3 + 2 + 1 \\ 6 - 4 - 2 & -2 + 2 + 0 & -2 + 1 + 2 \end{bmatrix} \\
&= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I_3
\end{aligned}$$

This confirms that we have correctly found the inverse of matrix A .

38.0.8 The Rank of a Matrix

The rank of a matrix A , denoted $\text{rank}(A)$ or $\text{rg}(A)$, is the dimension of the column space (or equivalently, the row space) of A .

Equivalent definitions of rank include:

- The maximum number of linearly independent columns of A
- The maximum number of linearly independent rows of A
- The order of the largest non-zero minor of A
- The number of non-zero rows in any row echelon form of A

38.0.9 How to isolate Matrices

Like in other types of equations we can isolate a matrix by doing the proper operations.

- $A \pm B = C \implies A = C \mp B$
- $AB = C \implies ABB^{-1} = CB^{-1} \implies A = CB^{-1}$
- $AB = C \implies A^{-1}AB = A^{-1}C \implies B = A^{-1}C$

Finding the Rank

To find the rank of a matrix:

1. Transform the matrix into row echelon form using Gauss-Jordan elimination
2. Count the number of non-zero rows in the resulting matrix

Properties of Rank

$$\begin{aligned}
\text{rank}(A) &\leq \min(m, n) \text{ for } A \in \mathbb{R}^{m \times n} \\
\text{rank}(A^T) &= \text{rank}(A) \\
\text{rank}(AB) &\leq \min(\text{rank}(A), \text{rank}(B)) \\
\text{rank}(A + B) &\leq \text{rank}(A) + \text{rank}(B)
\end{aligned}$$

For a square matrix $A \in \mathbb{R}^{n \times n}$, the following are equivalent:

- A is invertible
- $\text{rank}(A) = n$

- $\det(A) \neq 0$
- The columns of A are linearly independent
- The rows of A are linearly independent
- $Ax = 0$ has only the trivial solution $x = 0$

38.0.10 Column Space

The column space of A , denoted $\text{Col}(A)$, is the span of the columns of A :

$$\text{Col}(A) = \{\vec{y} \in \mathbb{R}^m : \vec{y} = A\vec{x} \text{ for some } \vec{x} \in \mathbb{R}^n\}$$

This is also called the range or image of the linear transformation represented by A . The dimension of the column space equals the rank of A .

38.0.11 Row Space

The *row space* of A , denoted $\text{Row}(A)$, is the span of the rows of A : It is also perpendicular to the *Null space*.

$$\text{Row}(A) = \text{Col}(A^T)$$

The dimension of the *row space* also equals the rank of A .

38.0.12 Null Space

The null space (or kernel) of A , denoted $\text{Null}(A)$ or $\text{Ker}(A)$, is the set of all vectors that A maps to zero:

$$\text{Null}(A) = \{\vec{x} \in \mathbb{R}^n : A\vec{x} = \vec{0}\}$$

The dimension of the null space is called the nullity of A , denoted $\text{nullity}(A)$.

Left Null Space

The left null space of A is the null space of A^T :

$$\text{Null}(A^T) = \{\vec{y} \in \mathbb{R}^m : A^T \vec{y} = \vec{0}\} = \{\vec{y} \in \mathbb{R}^m : \vec{y}^T A = \vec{0}^T\}$$

The Rank-Nullity Theorem connects these spaces:

$$\text{rank}(A) + \text{nullity}(A) = n$$

To find a basis for these spaces:

- *Column space*: Take the linearly independent columns of A .
- *Row space*: Take the non-zero rows from any row echelon form of A .
- *Null space*: Solve the homogeneous system $A\vec{x} = \vec{0}$ and express the general solution in terms of free variables.

38.0.13 Linear Maps as matrices

Linear maps between vector spaces V and W can be represented by matrices. In the following way

$$A\vec{x} = \sum_{i=1}^n x_i \vec{a}_i$$

where $\vec{a}_i$ are images of the basis vector of V as columns of A and $\vec{x} = (x_1, \dots, x_n)^T$.

38.0.14 Matrix Representation of Linear Maps on Basis vectors

Let V and W be two $\mathbb{F}$ -vector spaces, $\beta = \{v_1, \dots, v_n\}$ a basis of V and $\gamma = \{w_1, \dots, w_m\}$ a basis of W , and let $f : V \rightarrow W$ be linear. Then there exists a unique matrix $A = (a_{ij}) \in \mathbb{F}^{m \times n}$ with

$$f(v_j) = \sum_{i=1}^m a_{ij} w_i \quad \forall j = 1, \dots, n$$

Example:

Let $V = \mathbb{R}^2$, $W = \mathbb{R}^3$, and define a linear map $f : V \rightarrow W$ by:

$$f\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x + 2y \\ 3x + y \\ 4y \end{bmatrix}$$

We aim to find the matrix representation $[M]_{\beta}^{\gamma}(f) \in \mathbb{R}^{3 \times 2}$ of f , with respect to the standard bases:

$$\beta = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \subseteq V$$
$$\gamma = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\} \subseteq W$$

Step 1: Apply f to the basis vectors of V

$$f\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 + 2 \cdot 0 \\ 3 \cdot 1 + 0 \\ 4 \cdot 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 0 \end{bmatrix}$$
$$f\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 + 2 \\ 0 + 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}$$

Step 2: Form the matrix $[M]_{\beta}^{\gamma}(f)$

The columns of the matrix are the coordinate vectors of $f(v_1)$ and $f(v_2)$ in basis γ :

$$[M]_{\beta}^{\gamma}(f) = \begin{bmatrix} 1 & 2 \\ 3 & 1 \\ 0 & 4 \end{bmatrix}$$

Step 3: Verify the matrix representation

We check that:

$$f(v_j) = \sum_{i=1}^3 a_{ij}w_i \quad \text{for } j = 1, 2$$

$$f(v_1) = 1w_1 + 3w_2 + 0w_3$$

$$f(v_2) = 2w_1 + 1w_2 + 4w_3$$

This confirms the matrix representation is correct.

38.0.15 Matrices with respect to a basis

Expressing a linear map in terms of a different basis means that given two vector-spaces V and W with the basis $\beta = \{v_1, \dots, v_n\}$ and $\gamma = \{w_1, \dots, w_n\}$ respectively. Then the map $f : V \rightarrow W$ with a matrix representation A whose columns are vectors in the basis β can be expressed in terms of the basis γ using the following steps:

1. Determine the images of the basis vectors of V under f which will give us the matrix A_β . If not already given.
2. For every basis vector w_i of W apply $A_\beta w_i$. This is going to give you a vector k_i
3. Express each k_i as a linear combination of the basis vectors of W . Which means finding coefficients λ_k such that $\sum_{k=1}^n \lambda_k w_k = k_i$. This can be found using Gaussian elimination.
4. Form the matrix A_γ using each collection of coefficients λ_k as columns, where each column corresponds to the image of a basis vector v_j of V .

Another much faster, method to express a linear map in terms of another basis is to use the change of basis formula. If φ_γ is the change of basis matrix from γ to the canonical basis formed by the basis vector of W , and φ_γ^{-1} is the change of basis matrix from the canonical basis to γ , then:

$$A_\gamma = \varphi_\gamma^{-1} A_\beta \varphi_\gamma$$

The intuition behind is that in the previous method we are solving the system of equations of $A_\gamma \vec{x}_i = A_\beta w_i$ for each basis vector w_i in γ . Which is equivalent of $\vec{x}_i = \varphi_\gamma^{-1} A_\beta w_i$ when using Gaussian elimination. If you then put all $\vec{x}_i$ in order in a matrix then you get A_γ .

38.0.16 Example of a complete exercise

1. Determination of the kernel
2. Determination of the dimension of the kernel
3. Determination of the rank (dimension formula)
4. Determination of the image

Example:

Given

$$f \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 + x_2 \\ x_1 - x_2 + x_3 \\ 4x_1 - x_2 + 2x_3 \end{bmatrix}.$$

It should be shown that f is linear, and $\ker(f)$, $\text{img}(f)$ and their dimensions should be determined.

A direct proof of linearity is easily possible. Instead, we give the transformation matrix A . The images of the (canonical) basis vectors are

$$f \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}, \quad f \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}, \quad f \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}.$$

We obtain

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 1 & -1 & 1 \\ 4 & -1 & 2 \end{bmatrix}.$$

But now it must be shown that indeed $f(x) = Ax \quad \forall x \in \mathbb{R}^3$ holds, by, for example, calculating both Ax and $f(x)$ for a general x and showing equality: Here, with $x = (x_1, x_2, x_3)^T$

$$A \cdot x = \begin{bmatrix} 2 & 1 & 0 \\ 1 & -1 & 1 \\ 4 & -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 + x_2 \\ x_1 - x_2 + x_3 \\ 4x_1 - x_2 + 2x_3 \end{bmatrix}.$$

This obviously corresponds to $f(x)$, so that by Theorem 4.36, the map f is linear. To determine the kernel, one has to solve the system of linear equations

$$\begin{array}{cccc} 2 & 1 & 0 & 0 \\ 1 & -1 & 1 & 0 \\ 4 & -1 & 2 & 0 \end{array}$$

Which corresponds to the equation $Ax = f(x) = 0$. Gaussian elimination yields $x_3 = \lambda'$; $x_2 = \frac{2}{3} \cdot \lambda'$; $x_1 = -\frac{1}{3} \cdot \lambda'$, thus, with $\lambda = \frac{1}{3}\lambda'$:

$$\ker(f) = \left\{ x = \lambda \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix} \mid \lambda \in \mathbb{R} \right\}$$

It follows that $\dim(\ker(f)) = 1$ and because of $\dim(V) = 3$ from the dimension formula $\dim(\text{img}(f)) = 2$.

$\text{img}(f)$ corresponds to the linear span of the columns of the matrix. One chooses consequently $\dim(\text{img}(f))$ column vectors, e.g., the first ones, and tests if they are linearly independent. In the concrete case, this is obvious, because the second column is not a multiple of the first. It follows therefore,

$$\text{img}(f) = \left\{ x \mid x = \lambda \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix} + \mu \begin{bmatrix} 1 \\ -1 \\ -1 \end{bmatrix}, \quad \lambda, \mu \in \mathbb{R} \right\}$$

38.0.17 Diagonal and Triangular Matrices

A matrix is called *Diagonal* if the only non-zero elements are the elements on the main diagonal. A matrix is *Over Triangular* if all elements under or above the main diagonal are all zeroes.

Properties of Diagonal and Triangular Matrices

$$\det(A) = \prod_{i=1}^n a_{ii}$$

$$\text{eig}(A) = \{a_{1,1}, a_{2,2}, \dots, a_{n,n}\}$$

Chapter 39

Metrics

39.1 Metrics for Vectors

The metric of a vector $\vec{a}$ is described by different norms:

It also has the following properties:

- $\|\vec{a}\| \geq 0$ and $\|\vec{a}\| = 0 \Leftrightarrow \vec{a} = \vec{0}$
- $\|\lambda\vec{a}\| = |\lambda|\|\vec{a}\|$
- $\|\vec{a} + \vec{b}\| \leq \|\vec{a}\| + \|\vec{b}\|$ (Triangle inequality)
- $\|\vec{a} + \vec{b}\|^2 = \|\vec{a}\|^2 + \|\vec{b}\|^2 + 2\langle\vec{a}, \vec{b}\rangle$ (Pythagorean theorem)

These are the most common metric, although we will be using primarily the euclidean metric:

- *Euclidean metric:*

$$\|\vec{a}\| = \sqrt{\langle\vec{a}, \vec{a}\rangle} = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$$

- *Manhattan metric:*

$$\|\vec{a}\|_1 = |a_1| + |a_2| + \dots + |a_n|$$

- *Maximum metric:*

$$\|\vec{a}\|_\infty = \max(|a_1|, |a_2|, \dots, |a_n|)$$

39.2 Metric for Matrices

Given a metric $\|\cdot\|$ over $\mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$, we define

$$\|A\| := \sup_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \max_{\|x\|=1} \|Ax\|,$$

as the *matrix metric* for $\mathbb{R}^{n \times n}$.

- This norm is dependent on the chosen vector metric, hence is it called an *induced matrix metric*.
- The vector metric and the induced metric norm are compatible, which means:

$$\|Ax\|_{\text{vector}} \leq \|A\|_{\text{matrix}} \cdot \|x\|_{\text{vector}}.$$

- For matrix metrics the following holds:

$$\|AB\| \leq \|A\| \cdot \|B\|.$$

- For $\|\mathbb{R}^{n \times n}\|$ are all matrix metrics equivalent.

39.2.1 Spectral Metric

For $\|\cdot\|_2$ there is the so called *spectral metric*.

$$\|A\|_2 = \sqrt{\rho(A^T A)},$$

where the *spectral radius* $\rho(B)$ is the by absolute-value-sorted greatest eigenvalue of B .

39.2.2 Row Metric

For $\|\cdot\|_\infty$ there is the *row-sum metric*

$$\|A\|_\infty = \max_{i=1,\dots,n} \sum_{j=1}^n |a_{ij}|.$$

39.2.3 Column Metric

For $\|\cdot\|_1$ we get the *column-sum metric*

$$\|A\|_1 = \max_{j=1,\dots,n} \sum_{i=1}^n |a_{ij}|.$$

39.2.4 Frobenius Metric

The *frobenius metric* given by

$$\|A\|_F = \sqrt{\text{dig}(A^T, A)}$$

is not induced, but for it holds

$$\|AB\|_F \leq \|A\|_F \cdot \|B\|_F,$$

and is due to

$$\|Ax\|_2 \leq \|A\|_F \cdot \|x\|_2, \forall x$$

it is compatible with the vector metric $\|\cdot\|$. More precisely, $\|A\|_F$ is an upper bound for $\|A\|_2$.

Chapter 40

Linear Maps

A linear map $f : V \rightarrow W$ is a function that satisfies the following properties:

- $f(v_1 + v_2) = f(v_1) + f(v_2)$ for all $v_1, v_2 \in V$.
- $f(\lambda v) = \lambda f(v)$ for all $v \in V$ and $\lambda \in \mathbb{R}$.
- $f(0) = 0$.

This kind of function is also called *Homomorphism*. If $V = W$ is it called *Endomorphism*.

40.0.1 Types of Linear Maps

- *Injective or Monomorphism* (One-to-One): A linear map $f : V \rightarrow W$ is injective if $f(v_1) = f(v_2)$ implies $v_1 = v_2$.
- *Surjective or Epimorphism* (Onto): A linear map $f : V \rightarrow W$ is surjective if for every $w \in W$, there exists a $v \in V$ such that $f(v) = w$.
- *Bijjective or Isomorphism*: A linear map $f : V \rightarrow W$ is bijective if it is both injective and surjective.

Isomorphic if there exists a bijective linear map between them. In this case, we can say that the two vector spaces are *isomorphic*, and we write $V \cong W$.

40.0.2 Properties of Linear Maps

- The composition of two linear maps is a linear map.
- The inverse of a bijective linear map is also a linear map.
- The zero map $f : V \rightarrow W$ defined by $f(v) = 0$ for all $v \in V$ is a linear map.
- The identity map $\text{id}_V : V \rightarrow V$ defined by $\text{id}_V(v) = v$ for all $v \in V$ is a linear map.
- The sum of two linear maps $f : V \rightarrow W$ and $g : V \rightarrow W$ is a linear map defined by $(f + g)(v) = f(v) + g(v)$.
- The scalar multiplication of a linear map $f : V \rightarrow W$ by a scalar c is a linear map defined by $(cf)(v) = c(f(v))$.
- The composition of linear maps is associative, i.e., $(f \circ g) \circ h = f \circ (g \circ h)$.
- The composition of linear maps is distributive over addition, i.e., $f \circ (g + h) = f \circ g + f \circ h$.
- The composition of linear maps is compatible with scalar multiplication, i.e., $(cf) \circ g = c(f \circ g)$.
- Linear Maps compose a vector space over the field of scalars.

$$(\text{Hom}(V, W), +, \cdot)$$

40.0.3 Composition as Matrix Multiplication

Given $A \in \mathbb{F}^{m \times n}$, $B \in \mathbb{F}^{n \times l}$ and $\phi_A : \mathbb{F}^{n \times 1} \rightarrow \mathbb{F}^{m \times 1}$ and $\phi_B : \mathbb{F}^{l \times 1} \rightarrow \mathbb{F}^{n \times 1}$ induced linear maps then

•

$\phi_A \circ \phi_B$ is a linear map

•

$$\phi_A \circ \phi_B = \phi_{AB}$$

Proof I:

$$\begin{aligned} (\varphi_A \circ \varphi_B)(au + bv) &= \varphi_A(\varphi_B(au + bv)) \\ &= \varphi_A(a\varphi_B(u) + b\varphi_B(v)) \\ &= a\varphi_A(\varphi_B(u)) + b\varphi_A(\varphi_B(v)) \\ &= a(\varphi_A \circ \varphi_B)(u) + b(\varphi_A \circ \varphi_B)(v) \end{aligned}$$

□

Proof II:

$$\begin{aligned} (\varphi_A \circ \varphi_B)(x) &= \varphi_A(\varphi_B(x)) \\ &= \varphi_A(Bx) \\ &= A(Bx) \\ &= (AB)x \\ &= \varphi_{AB}(x) \end{aligned}$$

□

40.0.4 Injectivity and Surjectivity

Let $A \in \mathbb{F}^{m \times n}$. Then:

φ_A is injective if and only if the fiber of φ_A over the zero column $0_m \in \mathbb{F}^{m \times 1}$ consists only of the zero column $0_n \in \mathbb{F}^{n \times 1}$.

φ_A is surjective if and only if every column in $\mathbb{F}^{m \times 1}$ can be written as a linear combination of the columns of A .

40.0.5 Linear Extension Theorem

Let $\beta = \{v_1, \dots, v_n\}$ be a basis of the finite dimensional vector space V and $w_1, \dots, w_n$ be arbitrary vector in W . Then $\exists L$ with $L : V \rightarrow W$ such that $T(v_i) = w_i$.

Proof:

Define T as follows: Whenever $a = \lambda_1 v_1 + \dots + \lambda_n v_n$ then $T(a) = \lambda_1 w_1 + \dots + \lambda_n w_n$.

We know it is unique because for $\lambda_1, \dots, \lambda_n \neq \gamma_1, \dots, \gamma_n$ if we assume $a - b = \vec{0}$ then:

$$a - b = (\lambda_1 - \gamma_1)v_1 + \dots + (\lambda_n - \gamma_n)v_n = \vec{0}$$

If and only if $\lambda_1, \dots, \lambda_n = \gamma_1, \dots, \gamma_n$. Thus, it works uniquely for each vector.

For the linearity, using the definition of linearity and T this part becomes trivial.

Finally, for the uniqueness of T itself, let us assume there is a map U which $U(v_i) = T(v_i)$. Because we can write a vector a linear combination we will arrive at the fact that

$$U(x) = \lambda_1 U(v_1) + \cdots + \lambda_n U(v_n) = \lambda_1 T(v_1) + \cdots + \lambda_n T(v_n)$$

Therefore, $U = T$.

□◻◻

40.0.6 The Kernel of a Linear Map

The *kernel* of a linear map $f : V \rightarrow W$ is the set of all vectors in V that are mapped to the zero vector in W :

$$\ker(f) = \{v \in V \mid f(v) = 0\}$$

and also $\ker(f) \subseteq V$.

To compute the *kernel* of the linear map, we just solve the homogenous system of equations $Ax = \vec{0}$ where A is the matrix representation of f . The number of free variables is the $\dim(\ker(f))$.

Proof:

$\ker(f) \neq \emptyset$ because $0 \in \ker(f)$ (at least zero should be in the kernel by the definition of the vector space and linear maps). Also, for $v, w \in \ker(A)$ we know that $f(av + bw) = af(v) + bf(w) = a0 + b0 = 0$.

□◻◻

The Nullity

The *nullity* of a linear map $f : V \rightarrow W$ is the dimension of its kernel:

$$\text{nullity}(f) = \dim(\ker(f))$$

40.0.7 The Image of a Linear Map

The image of a linear map $f : V \rightarrow W$ is the set of all vectors in W that can be expressed as $f(v)$ for some $v \in V$:

$$\text{img}(f) = \{w \in W \mid w = f(v) \text{ for some } v \in V\}$$

Also, $\text{img}(f) \subseteq W$

To find $\dim(\text{im}(f))$, compute the rank of the matrix representing f . This is the number of pivot columns in the row-reduced form of A .

Proof:

$f(0) \in \text{img}(f) \implies \text{img}(f) \neq \emptyset$. Therefore, for $v_1, v_2 \in V, \exists w_1, w_2 \in \text{img}(f)$ with $w_1 = f(v_1)$ and $w_2 = f(v_2)$ due to

$$aw_1 + bw_2 = af(v_1) + bf(v_2) = f(av_1 + bv_2) \in \text{img}(f)$$

□◻◻

40.0.8 The Rank of a Linear Map

The rank of a linear map $f : V \rightarrow W$ is the dimension of its image:

$$\text{rank}(f) = \dim(\text{img}(f))$$

40.0.9 The Nullity of a Linear Map

The nullity of a linear map $f : V \rightarrow W$ is the dimension of its kernel:

$$\text{nullity}(f) = \dim(\ker(f))$$

40.0.10 Proof of the injectivity of the Kernel

Suppose we have a linear map f and two vectors v_1 and v_2 which are not equal. We are going to assume that they both map to the $\vec{0}$ therefore:

$$f(v_1) = f(v_2) = \vec{0}$$

Because of the linearity of the map we can write:

$$f(v_1 - v_2) = f(v_1) - f(v_2) = \vec{0} - \vec{0} = \vec{0}$$

And that contradicts the assumption that v_1 and v_2 are not equal. Therefore, the kernel of a linear map is injective.

40.0.11 Dimension Formula for Linear Mappings

Let $f : V \rightarrow W$ be linear and $\dim(V) = n$. Then

$$\dim(\ker(f)) + \text{rank}(f) = n.$$

Remember that $\ker(f)$ forms a subspace of V and therefore, $\dim(\ker(f)) := r \leq n$. We extend an arbitrary basis $(v_1, \dots, v_r)$ of $\ker(f)$ to a basis $(v_1, \dots, v_r, v_{r+1}, \dots, v_n)$ of V . Setting $w_{r+i} = f(v_{r+i})$ for $i = 1, \dots, n - r$, we have $\forall v \in V$:

$$\begin{aligned} f(v) &= f(\lambda_1 v_1 + \dots + \lambda_r v_r + \lambda_{r+1} v_{r+1} + \dots + \lambda_n v_n) \\ &= \lambda_1 \underbrace{f(v_1)}_{=0} + \dots + \lambda_r \underbrace{f(v_r)}_{=0} + \lambda_{r+1} f(v_{r+1}) + \dots + \lambda_n f(v_n) \\ &= \lambda_{r+1} f(v_{r+1}) + \dots + \lambda_n f(v_n) \\ &= \lambda_{r+1} w_{r+1} + \dots + \lambda_n w_n \end{aligned}$$

Thus, $\text{img}(f) = \text{span}(w_{r+1}, \dots, w_n)$. We now show that $w_{r+1}, \dots, w_n$ are linearly independent. Let

$$\lambda_{r+1} w_{r+1} + \dots + \lambda_n w_n = 0$$

From

$$0 = \lambda_{r+1} w_{r+1} + \dots + \lambda_n w_n = f(\lambda_{r+1} v_{r+1} + \dots + \lambda_n v_n)$$

it follows that

$$\lambda_{r+1} v_{r+1} + \dots + \lambda_n v_n \in \ker(f)$$

Therefore,

$$\lambda_{r+1} v_{r+1} + \dots + \lambda_n v_n = \lambda_1 v_1 + \dots + \lambda_r v_r$$

For some $\lambda_1, \dots, \lambda_r$. Since $v_1, \dots, v_n$ are linearly independent, we have $\lambda_1 = \dots = \lambda_n = 0$. Thus, the vectors $w_{r+1}, \dots, w_n$ are linearly independent.

It follows that $\dim(\text{img}(f)) = \text{rank}(f) = n - r$ and therefore,

$$\dim(\ker(f)) + \dim(\text{img}(f)) = \dim(\ker(f)) + \text{rank}(f) = r + n - r = n$$

□◻

40.0.12 Linear Maps and Matrices

Let V and W be finite-dimensional vector spaces over the same field $\mathbb{F}$. If $\dim V = n$ and $\dim W = m$, then a linear map $f : V \rightarrow W$ can be represented by an $m \times n$ matrix. The action of the linear map on a vector can be expressed as:

$$f(x) = A \cdot x,$$

where A is the matrix representation of f and v is the vector represented in a column format. The columns of the matrix A are the images of the basis vectors of V under the linear map f .

This fact comes from f being linear which means that

$$f(x) = x_1 f(e_1) + \dots + x_n f(e_n)$$

with e_i being a basis vector. And from that expression we can derive $(f(e_1), \dots, f(e_n))x$.

We can state that matrices *induce* linear maps which we are denoted as φ_A .

40.0.13 How to identify the type of Linear Map Summary

We want to summarize how to identify the type of linear map given $f : V \rightarrow W$,

- *Injective*: A linear map f is injective if and only if

$$\dim(\ker(f)) = 0.$$

- *Surjective*: The map f is surjective if every element of W is the image of some element in V . This occurs when

$$\dim(\text{img}(f)) = \dim(W).$$

That is, the image of f are the linear independent vector of f .

- *Bijjective*: A linear map is bijective if it is both injective and surjective. Therefore, it must satisfy both conditions:

$$\dim(\ker(f)) = 0 \quad \text{and} \quad \dim(\text{im}(f)) = \dim(W).$$

- *Neither*: If neither of the above conditions is satisfied, the map is neither injective nor surjective.

Example:

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by the matrix

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 2 & 3 & 2 \end{bmatrix}.$$

To identify the type of this linear map:

- Compute $\dim(\ker(f))$: By reducing the matrix A to row echelon form, we find that $\dim(\ker(f)) = 1$. This means f is not injective.
- Compute $\dim(\operatorname{im}(f))$: The image corresponds to the column space of A . Since there are 2 linearly independent rows in the row echelon form, we have $\dim(\operatorname{im}(f)) = 2$, which equals $\dim(W) = 2$. Therefore, f is surjective.

Thus, f is surjective but not injective.

Chapter 41

The Change Of Basis

Given a vector $\vec{x} = \lambda_1 \vec{v}_1 + \cdots + \lambda_n \vec{v}_n$ with $\vec{v}_1, \dots, \vec{v}_n \in B_v$ being the basis vectors.

Now imagine having another basis, coordinates, $\vec{w}_1, \dots, \vec{w}_n \in B_w$. It is important to understand that the basis vectors $\vec{w}_1, \dots, \vec{w}_n$ are written in terms of **our basis**, because for the perspective of the user of the basis, the basis vectors are not different from the canonical basis.

The translation between the two basis is given in the following way:

$$B_w \vec{\mu}_w = \vec{\lambda}_v, \quad \vec{\mu}_w = B_w^{-1} \vec{\lambda}_v$$

This works because from the perspective of the user of the basis, the basis vectors are not different from the canonical basis, therefore we can write the basis vectors in terms of the our basis and the apply a transformation to get the coordinates in the our basis.

If we then want to get the vector in the basis B_w we need to apply the inverse of B_w to the vector in the basis B_v .

41.0.1 Pseudo-equality Of Translations

Note we can write the following pseudo-equality for our translations

$$B_v \vec{\lambda}_v = B_w \vec{\mu}_w$$

This generalizes the notion of the change of basis, by not limiting ourselves to only the canonical basis.

Note, that we are cheating a bit here with the equal notation. Now by taking the corresponding inverse we can write the change of basis as desired.

41.0.2 Coordinate mapping

Let V be a $\mathbb{F}$ -vector space with a basis $\beta = \{v_1, \dots, v_n\}$. Then there exists exactly one isomorphism

$$\varphi_\beta : \mathbb{F}^n \rightarrow V$$

such that

$$\varphi_\beta(e_i) = v_i, \quad \text{for } 1 \leq i \leq n.$$

which is called the *coordinate mapping* of V with respect to the basis β .

41.0.3 Coordinates with respect to a basis

The isomorphism φ_β from the previous theorem is called the *coordinate mapping*, and for $v \in V$, we define

$$(\vec{v})_\beta := \varphi_\beta^{-1}(v) \in \mathbb{F}^n$$

as the *coordinates of v with respect to β* .

41.0.4 Translation of Coordinates

We define the *Translation* of a vector to another basis as:

$$T_\beta^\gamma = \varphi_\beta^{-1} \circ \varphi_\gamma$$

where φ_β and φ_γ are the matrices made of the basis vector of the bases β and A with respect to the standard basis, respectively.

$$\begin{array}{ccc} & V & \\ \varphi_\gamma \nearrow & & \nwarrow \varphi_\beta \\ \mathbb{F}^n & \xrightarrow{T_\beta^\gamma = \varphi_\beta^{-1} \circ \varphi_\gamma} & \mathbb{F}^n \end{array}$$

41.0.5 Change of Basis Theorem

Let $v \in V$ be arbitrary, with $(\vec{v})_\gamma = (x_1, \dots, x_n)^T$ and $(\vec{v})_\beta = (y_1, \dots, y_n)^T$. Then the following holds:

$$(\vec{v})_\beta = \varphi_\beta^{-1} \circ \varphi_\gamma(\vec{v})_\gamma$$

In vector notation with the translation:

$$\begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} = T_\beta^A \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$$

41.0.6 Change of Basis in Linear Maps

Let V and W be finitely generated $\mathbb{F}$ -vector spaces with bases γ and β , respectively, and let $f \in \text{Hom}(V, W)$. Then we have

$$M_\beta^\gamma(f) = \varphi_\beta^{-1} \circ f \circ \varphi_\gamma$$

Thus, $M_\beta^\gamma(f)$ tells you to take vector in the basis γ , translate it to the canonical basis, apply the linear map f and then convert the result to the basis β .

$$\begin{array}{ccc} \mathbb{F}^n & \xrightarrow{M_\beta^\gamma} & \mathbb{F}^m \\ \searrow \varphi_\gamma & & \swarrow \varphi_\beta \\ & V \xrightarrow{f} W & \end{array}$$

41.0.7 Transformations involving multiple bases

Let V and W be finitely generated vector spaces with bases γ and γ' for V , and β and β' for W , respectively. Furthermore, let $f : V \rightarrow W$ be a linear map. Then the following holds:

$$M_{\beta'}^{\gamma'}(f) = T_{\beta'}^{\beta} \cdot M_{\beta}^{\gamma}(f) \cdot (T_{\gamma'}^{\gamma})^{-1}$$

More explicitly, we can write this as:

$$\begin{aligned} M_{\beta'}^{\gamma'}(f) &= T_{\beta'}^{\beta} \cdot M_{\beta}^{\gamma}(f) \cdot (T_{\gamma'}^{\gamma})^{-1} \\ &= (\varphi_{\beta'}^{-1} \circ \varphi_{\beta}) \circ (\varphi_{\beta}^{-1} \circ f \circ \varphi_{\gamma}) \circ (\varphi_{\gamma'}^{-1} \circ \varphi_{\gamma})^{-1}. \\ &= (\varphi_{\beta'}^{-1} \circ \varphi_{\beta}) \circ (\varphi_{\beta}^{-1} \circ f \circ \varphi_{\gamma}) \circ (\varphi_{\gamma}^{-1} \circ \varphi_{\gamma'}). \end{aligned}$$

We can visualize this as a commutative diagram where when take an inverse we reverse the direction of the arrows:

$$\begin{array}{ccccc} \mathbb{R}^n & & \xrightarrow{M_{\beta}^{\gamma}} & & \mathbb{R}^m \\ & \searrow \varphi_{\gamma} & & \swarrow \varphi_{\beta} & \\ & & V & \xrightarrow{f} & W \\ & \nearrow \varphi_{\gamma'} & & \nwarrow \varphi_{\beta'} & \\ \mathbb{R}^n & & \xrightarrow{M_{\beta'}^{\gamma'}} & & \mathbb{R}^m \end{array}$$

$\downarrow T_{\gamma'}^{\gamma}$ $\downarrow T_{\beta'}^{\beta}$

Chapter 42

The Determinant of a Matrix

The determinant of a square matrix $A \in \mathbb{R}^{n \times n}$, denoted $\det(A)$ or $|A|$, is a scalar value that provides important information about the matrix, including whether it is invertible and the volume scaling factor of the linear transformation represented by A . The determinant can be computed using various methods, including the Laplace expansion, row reduction, or the Leibniz formula.

The determinant of a 2×2 matrix is given by:

$$\det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

For a 3×3 matrix, the determinant can be computed using the rule of Sarrus or the co-factor expansion:

$$\det(A) = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

The determinant of larger matrices can be computed using co-factor expansion along any row or column:

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det(A_{ij}),$$

where A_{ij} is the $(n-1) \times (n-1)$ sub-matrix obtained by deleting the i -th row and j -th column of A .

42.0.1 Properties of the determinant

- $\det(A) = 0$ if and only if A is singular (not invertible).
- $\det(AB) = \det(A) \cdot \det(B)$ for any square matrices A and B of the same size.
- $\det(A^T) = \det(A)$.
- If a row (or column) of A is multiplied by a scalar α , then $\det(A)$ is multiplied by α .
- If two rows (or columns) of A are swapped, then $\det(A)$ changes sign.

$$\det(a, b, c) = -\det(b, a, c)$$

- If a row (or column) of A is added to another row (or column), then $\det(A)$ remains unchanged.
- If one of the columns is a linear combination of the others, then $\det(A) = 0$.
- The determinant of the identity matrix I_n is 1.

- The determinant of can split into the sum of more determinants:

$$\det(a, b, c + d) = \det(a, b, c) + \det(a, b, d)$$

- The determinant of a diagonal matrix is the product of its diagonal entries.
- For elementary matrices $\det C1 = 1$, $\det C2 = -1$, $\det C3 = \lambda$.
- For $A \in \mathbb{R}^{n \times n}$ $\det(\lambda A) = \lambda^n \det(A)$

42.0.2 Proof of the multiplication of determinants

For $A, B \in \mathbb{F}^{n \times n}$, it holds that

$$\det(AB) = \det(A) \det(B).$$

Proof:

If $\det(A) = 0$ or $\det(B) = 0$, then $\text{rank}(L_A) < n$ or $\text{rank}(L_B) < n$.

Thus, $\text{rank}(L_{AB}) < n$, and the statement is trivial.

So let us assume that A and B are invertible, i.e., $\text{rank}(A) = \text{rank}(B) = n$.

We now assume that in such a case, a matrix can always be transformed into reduced row echelon form using the row operations from Gaussian elimination.

Then, $A = C_k \cdots C_1$ for certain elementary matrices C_i , and therefore:

$$\begin{aligned} \det(AB) &= \det(C_k \cdots C_1 B) \\ &= \det(C_k) \det(C_{k-1} \cdots C_1 B) \\ &\vdots \\ &= \det(C_k) \cdots \det(C_1) \det(B) \\ &= \det(C_k \cdots C_1) \det(B) \\ &= \det(A) \det(B). \end{aligned}$$

Therefore, also if A is invertible, then

$$\det(A^{-1}) = (\det(A))^{-1}.$$

Proof:

It holds that

$$\det(A) \det(A^{-1}) = \det(AA^{-1}) = \det(E) = 1.$$

42.0.3 The Leibniz Formula

Let $A = (a_{ij}) \in \mathbb{F}^{n \times n}$ be a square matrix over a field K . The determinant of A is defined via the **Leibniz formula** as:

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \cdot a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)},$$

where:

- S_n is the set of all permutations of $\{1, 2, \dots, n\}$,
- $\text{sgn}(\sigma)$ is the sign of the permutation σ , equal to $+1$ for even permutations and -1 for odd ones.

Derivation

Step 1: Expansion using multilinearity

Each column vector a_j of the matrix A can be written as a linear combination of the standard basis vectors $e_1, \dots, e_n$:

$$a_j = \sum_{i=1}^n a_{ij} e_i.$$

Using the multilinearity of the determinant function (i.e., linearity in each column), we expand:

$$\det(a_1, a_2, \dots, a_n) = \det\left(\sum_{i_1} a_{i_1 1} e_{i_1}, \sum_{i_2} a_{i_2 2} e_{i_2}, \dots, \sum_{i_n} a_{i_n n} e_{i_n}\right).$$

By multilinearity, this becomes a sum over all n -tuples $(i_1, i_2, \dots, i_n)$:

$$= \sum_{i_1, i_2, \dots, i_n=1}^n a_{i_1 1} a_{i_2 2} \cdots a_{i_n n} \cdot \det(e_{i_1}, e_{i_2}, \dots, e_{i_n}).$$

Step 2: Eliminate repeated indices

If two indices $i_j = i_k$ for $j \neq k$, then the corresponding determinant is zero (because the determinant is alternating and thus, zero for repeated columns). Therefore, only terms with pairwise distinct indices remain. These correspond to permutations of $\{1, 2, \dots, n\}$.

Let $\sigma \in S_n$ be such a permutation. Then we can write:

$$\det(A) = \sum_{\sigma \in S_n} a_{\sigma(1)1} a_{\sigma(2)2} \cdots a_{\sigma(n)n} \cdot \det(e_{\sigma(1)}, e_{\sigma(2)}, \dots, e_{\sigma(n)}).$$

Step 3: Sign of the permutation

Since the determinant of $(e_{\sigma(1)}, \dots, e_{\sigma(n)})$ is $\text{sgn}(\sigma)$, we obtain:

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \cdot a_{\sigma(1)1} a_{\sigma(2)2} \cdots a_{\sigma(n)n}.$$

Equivalently, by re-indexing:

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \cdot \prod_{i=1}^n a_{i\sigma(i)}.$$

This formula expresses the determinant as a sum over all permutations of the indices $\{1, \dots, n\}$, where each term is a product of matrix entries taken from different rows and columns, weighted by the sign of the corresponding permutation.

42.0.4 Laplace's Method (Co-factor Expansion)

The determinant of an $n \times n$ matrix $A = [a_{ij}]$ can be computed using the *Laplace expansion*, also known as co-factor expansion. This method expands the determinant along a specific row or column.

Expansion Along Row i :

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det(M_{ij})$$

Expansion Along Column j :

$$\det(A) = \sum_{i=1}^n (-1)^{i+j} a_{ij} \det(M_{ij})$$

Where:

- a_{ij} is the element in the i -th row and j -th column of matrix A .
- M_{ij} is the *minor* of a_{ij} , i.e., the determinant of the submatrix formed by removing the i -th row and j -th column from A .
- $(-1)^{i+j}$ is the *sign factor*, determining the sign of each co-factor term.

Steps:

1. **Choose a Row or Column:** Select any row or column of the matrix. It's often easiest to choose one with many zeros.

2. **For Each Element:** For each element, a_{ij} , in the chosen row or column:

Find the Minor, M_{ij} : The minor M_{ij} is the determinant of the sub-matrix formed by deleting the i -th row and the j -th column of the original matrix.

Find the Co-factor, C_{ij} : The co-factor C_{ij} is the minor multiplied by a sign factor:

$$C_{ij} = (-1)^{i+j} M_{ij}$$

The term $(-1)^{i+j}$ gives a checkerboard pattern of signs:

$$\begin{bmatrix} + & - & + & - & \cdots \\ - & + & - & + & \cdots \\ + & - & + & - & \cdots \\ - & + & - & + & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

3. **Calculate the Determinant:** The determinant of the matrix, A , is the sum of the products of the elements in the chosen row or column and their corresponding co-factors.

Expansion along the i -th row:

$$\det(A) = \sum_{j=1}^n a_{ij} C_{ij} = a_{i1} C_{i1} + a_{i2} C_{i2} + \cdots + a_{in} C_{in}$$

Expansion along the j -th column:

$$\det(A) = \sum_{i=1}^n a_{ij} C_{ij} = a_{1j} C_{1j} + a_{2j} C_{2j} + \cdots + a_{nj} C_{nj}$$

Both expansions give the same result.

Example (3×3 Matrix):

Let

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Expanding along the first row:

$$1. a_{11}: M_{11} = \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix}, C_{11} = +M_{11}$$

$$2. a_{12}: M_{12} = \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix}, C_{12} = -M_{12}$$

$$3. a_{13}: M_{13} = \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}, C_{13} = +M_{13}$$

Therefore,

$$\det(A) = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$

Example of Determinant Calculation

Let's calculate the determinant of the matrix:

$$A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 2 \\ 0 & 1 & 1 \end{bmatrix}$$

Using the rule of Sarrus for 3×3 matrices:

$$\begin{aligned} \det(A) &= 2 \cdot 0 \cdot 1 + 1 \cdot 2 \cdot 3 + 3 \cdot 1 \cdot 1 - (3 \cdot 0 \cdot 0 + 1 \cdot 2 \cdot 2 + 2 \cdot 1 \cdot 1) \\ &= 0 + 6 + 3 - (0 + 4 + 2) \\ &= 9 - 6 = 3 \end{aligned}$$

Thus, the determinant of matrix A is $\det(A) = 3$.

Now consider the following 4×4 matrix:

$$A = \begin{bmatrix} 2 & 1 & 3 & 2 \\ 4 & 0 & -1 & 3 \\ -2 & 3 & 1 & 5 \\ 1 & -1 & 0 & 2 \end{bmatrix}$$

For a 4×4 matrix, we can use Laplace Method along the first row:

$$\begin{aligned} \det(A) &= a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} + a_{14}C_{14} \\ &= 2 \cdot \det \begin{bmatrix} 0 & -1 & 3 \\ 3 & 1 & 5 \\ -1 & 0 & 2 \end{bmatrix} - 1 \cdot \det \begin{bmatrix} 4 & -1 & 3 \\ -2 & 1 & 5 \\ 1 & 0 & 2 \end{bmatrix} \\ &\quad + 3 \cdot \det \begin{bmatrix} 4 & 0 & 3 \\ -2 & 3 & 5 \\ 1 & -1 & 2 \end{bmatrix} - 2 \cdot \det \begin{bmatrix} 4 & 0 & -1 \\ -2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \end{aligned}$$

$$\det(A) = 145$$

42.0.5 Determinant of the reduced echelon form

The determinant has another trait that states that if your matrix A is REF then its determinant is given by the product of the main diagonal.

42.0.6 Gaussian Method for finding determinants

Another method for finding determinants is using the row operations we already knew plus also column operations but with the catch that for the column case we have to take into consideration how operating with columns affect the determinant, in contrast to the row operations, except for scaling a row by some λ in this case we would need to undo the change by multiplying the determinant by $\frac{1}{\lambda}$.

Example:

We are given the matrix:

$$A = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}$$

We apply Gaussian elimination:

Swap $R_2 \leftrightarrow R_3$:

$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

Now the matrix is upper triangular. The determinant is the product of diagonal elements times -1 for one row swap:

$$\det(A) = (-1)^1 \cdot (1) \cdot (-1) \cdot (-1) \cdot (3) = -3$$

$$\det(A) = -3$$

Chapter 43

Elementary Matrix

We are going to define 3 matrices that can be obtained from the Elementary Row Operations on the *Identity Matrix* and call them *Elementary Matrices*.

C1: Row Addition

$$C_1 := \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & \lambda & 1 & \\ & & & & \ddots \end{bmatrix} \in \mathbb{F}^{n \times n} \quad \text{or} \quad C_1 := \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & \lambda & \\ & & & 1 & \\ & & & & \ddots \end{bmatrix} \in \mathbb{F}^{n \times n}$$

C2: Row swap

$$C_2 := \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 0 & 1 & \\ & & 1 & 0 & \\ & & & & \ddots \\ & & & & & 1 \end{bmatrix} \in \mathbb{F}^{n \times n}$$

C3: Row Scaling

$$C_3 := \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & \lambda & & \\ & & & \ddots & \\ & & & & 1 \end{bmatrix} \in \mathbb{F}^{n \times n}$$

Each of them also encodes the row Operations from where they came from. So now Gaussian Elimination can be encoded as matrix multiplication.

Formally,

$$C_1 := \text{Add}_m(i, j, \lambda) := I_m + a e_i f_j : m \times m \rightarrow \mathbb{F} : (s, t) \mapsto \begin{cases} 1, & s = t \\ \lambda, & (s, t) = (i, j) \\ 0, & \text{else} \end{cases}$$

$$C_2 := \text{Swap}_m(i, j) := (e_1, \dots, e_{i-1}, e_j, e_{j+1}, \dots, e_{j-1}, e_i, e_{j+1}, \dots, e_m) : m \times m \rightarrow \mathbb{F} : (s, t) \mapsto \begin{cases} 1, & s = t \notin \{i, j\} \\ 1, & (s, t) \in \{(i, j), (j, i)\} \\ 0, & \text{else} \end{cases}$$

$$C_3 := \text{Mul}_m(i, \lambda) := I_m + (a - 1)e_i f_j : m \times m \rightarrow \mathbb{F} : (s, t) \mapsto \begin{cases} 1, & s = t \neq i \\ \lambda, & s = t = i \\ 0, & \text{else} \end{cases}$$

43.0.1 Invertibility

Because every row operation is invertible now we can also claim that Elementary Matrices are invertible and they also represented that inverted row operation.

Formally,

- $\text{Add}_m(i, j, a)$ has the inverse $\text{Add}_m(i, j, -a)$.
- $\text{Mul}_m(i, \lambda)$ has the inverse $\text{Mul}_m(i, \lambda^{-1})$.
- $\text{Swap}_m(i, j)$ has itself as the inverse.

43.0.2 Determinants

The determinants corresponding to each matrix are:

$$\begin{aligned} \det C1 &= 1 \\ \det C2 &= -1 \\ \det C3 &= \lambda \end{aligned}$$

Chapter 44

Orthogonal Linear Maps

In this section we are going to take a look at some special type of linear maps that are called *orthogonal* and at the term *isometry*. It will be useful to remember some concepts about the dot product such as $\langle x, x \rangle = x^T \cdot x = \|x\|^2$.

44.0.1 Isometry

Given two vector spaces $(V, \|\cdot\|_V)$ and $(W, \|\cdot\|_W)$. Then a map $f : V \rightarrow W$ is called *isometric* if and only if $\|x - y\|_V = \|f(x) - f(y)\|_W$.

It also has the following traits:

1. *Length-Preserving*: $\|x\|_V = \|y\|_W \quad \forall x \in V$.
2. *Angle-Preserving*: $\angle(x, y) = \angle(f(x), f(y)) \quad \forall x, y \in V$
3. *Linear*: If $f(x + \lambda y) = f(x) + \lambda f(y)$

44.0.2 Unit-sphere

For $(V, \mathbb{R})$ with metric $\|\cdot\|$ we denote

$$S = \{x \mid x \in V : \|x\| = 1\}$$

as the *unit-sphere*.

44.0.3 The Rotation Matrix

This matrix is a common example of an isometric linear map, because of the way it transforms the space.

For the 2D we have

$$\begin{bmatrix} \cos(\phi) & -\sin(\phi) \\ \sin(\phi) & \cos(\phi) \end{bmatrix}$$

The derivation comes from the addition theorems for $\sin(\phi)$ and $\cos(\phi)$.

Given two vectors $\vec{v}$ and $\vec{g}$ whose angles are α and $\theta = \alpha + \beta$. We can rotate the vector $\vec{v}$ by some β to transform into $\vec{g}$.

$$\begin{aligned} x &= r \cos(\alpha) \\ y &= r \sin(\alpha) \end{aligned}$$

Then

$$\begin{aligned}x' &= r' \cos(\alpha + \beta) \\y' &= r' \sin(\alpha + \beta)\end{aligned}$$

And therefore

$$\begin{aligned}x' &= r' \cos(\alpha) \cos(\beta) - r' \sin(\alpha) \sin(\beta) \\y' &= r' \cos(\alpha) \sin(\beta) + r' \sin(\alpha) \cos(\beta)\end{aligned}$$

Because, a rotation does not change the magnitude we can say that $r = r'$. Thus

$$\begin{aligned}x' &= r \cos(\alpha) \cos(\beta) - r \sin(\alpha) \sin(\beta) \\y' &= r \cos(\alpha) \sin(\beta) + r \sin(\alpha) \cos(\beta)\end{aligned}$$

Now notice that we have our x and y inside the expression

$$\begin{aligned}x' &= x \cos(\beta) - y \sin(\beta) \\y' &= x \sin(\beta) + y \cos(\beta)\end{aligned}$$

which gives us

$$\begin{bmatrix} \cos(\phi) & -\sin(\phi) \\ \sin(\phi) & \cos(\phi) \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

□◻◻

Rotation Matrices For $\mathbb{R}^3$

$$R_x(\phi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\phi) & -\sin(\phi) \\ 0 & \sin(\phi) & \cos(\phi) \end{bmatrix}$$

$$R_y(\phi) = \begin{bmatrix} 1 & 0 & \sin(\phi) \\ 0 & 1 & 0 \\ -\sin(\phi) & 0 & \cos(\phi) \end{bmatrix}$$

$$R_z(\phi) = \begin{bmatrix} \cos(\phi) & -\sin(\phi) & 0 \\ \sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

44.0.4 Orthogonal Matrices

A matrix $A \in \mathbb{R}^{n \times n}$ is *orthogonal* if its columns vectors form an orthonormal basis.

The set of all *orthogonal* matrices is $O(n)$.

Properties of orthogonal matrices

1. $A \in O(n)$.
2. If A has an inverse then $A^{-1} = A^T$.
3. $A^T \in O(n)$.
4. $\|Q\|_2 = \sup_{x \neq 0} \frac{\|Qx\|}{\|x\|_2} = 1$.

Proof 1-2:

$$\begin{aligned}
 A^T = A^{-1} &\iff A^T A = E \\
 &\iff (A^T A)_{ij} = \langle a_i, a_j \rangle = \delta_{ij} \quad 1 \leq i, j \leq n \\
 &\iff a_i \perp a_j, i \neq j, \text{ and } \|a_i\| = 1, 1 \leq i \leq n \\
 &\iff A \in O(n)
 \end{aligned}$$

Proof 1-3:

$$\begin{aligned}
 A \in O(n) &\iff \\
 &\iff A^{-1} = A^T \\
 &\iff (A^T)^{-1} = (A^T)^T \\
 &\iff (A^T)^T = A \\
 &\iff A^T \in O(n)
 \end{aligned}$$

Example of a proof

Prove that H_n is *orthogonal* for all $\vec{u} \in \mathbb{R}^n \setminus \{0\}$.

$$H_n = I_n - 2 \frac{uu^T}{u^T u}$$

I_n is the identity matrix.

Proof:

$$\begin{aligned}
 (H_n)^2 &= I \\
 &= \left(I_n - 2 \frac{uu^T}{u^T u} \right)^2 \\
 &= I_n^2 - 4 \frac{uu^T}{u^T u} + 4 \left(\frac{uu^T}{u^T u} \right)^2 \\
 &= I_n - 4 \frac{uu^T}{u^T u} + 4 \frac{uu^T uu^T}{(u^T u)^2} \\
 &= I_n - 4 \frac{uu^T}{u^T u} + 4 \frac{u(u^T u)u^T}{(u^T u)^2} \\
 &= I_n - 4 \frac{uu^T}{u^T u} + 4 \frac{uu^T}{u^T u} \\
 &= I_n
 \end{aligned}$$

QED

44.0.5 The Determinants of Orthogonal Matrices

The *determinant* of this kind of matrices can only be either positioning 1 or negative 1.

Proof:

We know that $\det(I) = 1$, therefore

$$\begin{aligned}\det(I) &= \det(AA^{-1}) \\ &= \det(AA^T) \\ &= \det(A) \det(A^T) \\ &= (\det(A))^2 \\ &= \|\det(A)\| = 1\end{aligned}$$

□◻

44.0.6 The Orthogonal Group

$O(n)$ with the operation matrix multiplication forms the *orthogonal group*.

44.0.7 Dot Product

Given $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ then this two statements are equivalent.

$$\langle f(x), f(y) \rangle = \langle x, y \rangle \quad \forall x, y \in \mathbb{R}^n$$

and

f preserves the angles and length.

f is also called *orthogonal*.

Proof $\Rightarrow$:

$$\|x\|^2 = \langle x, x \rangle = \langle f(x), f(x) \rangle = \|f(x)\|^2$$

This implies the length preservation. The isometry gives us:

$$\begin{aligned}\|f(x) - f(y)\|^2 &= \langle f(x) - f(y), f(x) - f(y) \rangle \\ &= \langle f(x), f(x) \rangle - 2 \langle f(x), f(y) \rangle + \langle f(y), f(y) \rangle \\ &= \langle x, x \rangle - 2 \langle x, y \rangle + \langle y, y \rangle \\ &= \langle x - y, x - y \rangle = \|x - y\|^2\end{aligned}$$

Then

$$\begin{aligned}\angle(f(x), f(y)) &= \arccos \left(\frac{\langle f(x), f(y) \rangle}{\|f(x)\| \cdot \|f(y)\|} \right) \\ &= \arccos \left(\frac{\langle x, y \rangle}{\|x\| \cdot \|y\|} \right) \\ &= \angle(x, y)\end{aligned}$$

□◻

Proof $\Leftarrow$:

With $\alpha = \angle(x, y) = \angle(f(x), f(y))$ gilt:

$$\langle f(x), f(y) \rangle = \cos(\alpha) \cdot \|f(x)\| \cdot \|f(y)\| = \cos(\alpha) \cdot \|x\| \cdot \|y\| = \langle x, y \rangle$$

□◻◻

44.0.8 Orthonormal Matrices

A matrix $Q \in \mathbb{R}^{n \times n}$ is *orthonormal* if for each of its column vectors q_j , they are pairwise orthonormal.

$$(q_i, q_j) = \delta_{i,j} = \begin{cases} 1 & i = j \\ 0 & \text{else} \end{cases}$$

Properties of Orthonormal Matrices

If $Q, \tilde{Q} \in \mathbb{R}^{n \times n}$ are orthonormal matrices, the following applies:

1. $QQ^T = Q^TQ = I$, i.e. $Q^{-1} = Q^T$.
2. $\kappa_2(Q) = 1$.
3. $Q\tilde{Q}$ is an orthonormal matrix.

Proof 1:

When applying the matrix product, we note that our repeated inner product is applied to each of the flipped columns. Since by our definition

$$(q_i, q_j) = \delta_{i,j} = \begin{cases} 1 & i = j \\ 0 & \text{else} \end{cases}$$

We get a matrix with 1's at the entries $QQ^T_{i,j}$ if $i = j$ and else 0. in other words the identity.

□◻◻

Proof 2:

For all $x \in \mathbb{R}^n$ holds that

$$\|Qx\|_2^2 = (Qx)^T(Qx) = x^T Q^T Q x = x^T x = \|x\|_2^2$$

Since orthonormal matrices do not preserve the length. Therefore $\|Q\|_2 = 1$ and analogously, the same for Q^{-1} hence

$$\kappa_2(Q) = \|Q\|_2 \|Q^{-1}\| = 1$$

□◻◻

Proof 3:

Applying our first property, we need to show that $(Q\tilde{Q})^T = (Q\tilde{Q})^{-1}$.

$$(Q\tilde{Q})^T(\tilde{Q}Q) = Q^T \tilde{Q}^T (\tilde{Q}Q) = Q^T I Q = Q^T Q = I$$

Therefore the columns are pairwise orthonormal.

□◻◻

44.0.9 Theorem of the Matrix Representation

The mapping $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is orthogonal if and only if the associated matrix representation Q (with respect to the standard basis) is an orthogonal matrix, i.e., $Q \in O(n)$.

Proof $\Rightarrow$:

f is linear, and thus has a matrix representation Q . It follows that

$$\langle Qx, Qy \rangle = \langle Q^\top Qx, y \rangle = \langle x, y \rangle \quad \forall x, y \in \mathbb{R}^n,$$

thus,

$$\langle Q^\top Qx - x, y \rangle = 0 \quad \forall x, y.$$

Choosing $y = Q^\top Qx - x$, we get

$$\langle Q^\top Qx - x, Q^\top Qx - x \rangle = 0 \quad \forall x,$$

and hence $(Q^\top Q - I)x = 0 \quad \forall x$. It follows that $Q^\top Q = I$, and by Theorem 7.3, $Q \in O(n)$.

□◻◻

Proof $\Leftarrow$:

Follows by the same calculation as above.

□◻◻

44.0.10 Change of Basis

Let f be orthogonal. Then the matrix representation of f with respect to any orthonormal basis is orthogonal.

Proof:

Let E be the standard basis and B an orthonormal basis. Let $Q = M_E^E(f)$ and $S = T_E^B$ be the change-of-basis matrix from B to E . Since the columns of S are the basis vectors of B , we have $S \in O(n)$.

$$M_B^B(f) = S^{-1}QS \in O(n),$$

due to the group property of $O(n)$.

□◻◻

44.0.11 QR-Decomposition

A use case of orthogonal matrices is the *QR-Decomposition*. The goal is that given a system $Ax = b$ with a matrix $A \in \mathbb{R}^{m \times n}$, $m \geq n$ and $rg(A) = n$ we can find two matrices Q and R such that $A = QR$ with $R \in \mathbb{R}^{n \times n}$ and $Q \in \mathbb{R}^{m \times n}$.

R is an upper triangular matrix and Q is orthogonal for the columns.

We know that the best approximation by the method of the least squares is given by

$$x_s = (A^T A)^{-1} A^T b$$

Because of the orthogonality of the columns $rg(Q) = rg(A)$, therefore, it can be inverted. Hence, we get:

$$\begin{aligned}
x_s &= (A^T A)^{-1} A^T b \\
&= ((QR)^T (QR))^{-1} (QR)^T b \\
&= (R^T Q^T Q R)^{-1} (QR)^T b \\
&= (R^T R)^{-1} R^T Q^T b \\
&= R^{-1} (R^T)^{-1} R^T Q^T b \\
&= R^{-1} Q^T b \\
Rx_s &= Q^T b
\end{aligned}$$

Thus, we can solve the linear system by finding Q which is matrix with an orthonormal basis of the vector space which can be found via the Gram-Schmidt algorithm. The matrix R consists of the matrix of the inner products our A matrix and Q

$$R = \begin{bmatrix} \langle q_1, a_1 \rangle & \langle q_1, a_2 \rangle & \cdots & \langle q_1, a_n \rangle \\ 0 & \langle q_2, a_2 \rangle & \cdots & \langle q_2, a_n \rangle \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \langle q_n, a_n \rangle \end{bmatrix}$$

Or even simpler $R = Q^T A$.

Example:

Find the *QR-Decomposition* of the matrix A given by:

$$\begin{bmatrix} 1 & 3 \\ 1 & -1 \end{bmatrix}$$

First, we find the orthonormal basis of the columns of our matrix A

$$q_1 = \frac{a_1}{\|a_1\|} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$r_2 = a_2 - \langle a_2, q_1 \rangle q_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\rangle \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

$$q_2 = \frac{1}{\sqrt{8}} \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

$$Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{2}{\sqrt{8}} \\ \frac{1}{\sqrt{2}} & -\frac{2}{\sqrt{8}} \end{bmatrix}$$

$$R = \begin{bmatrix} \sqrt{2} & \sqrt{2} \\ 0 & \sqrt{8} \end{bmatrix}$$

Chapter 45

Eigenvectors and Eigenvalues

Eigenvector:

An eigenvector of a square matrix A is a non-zero vector $\vec{v}$ that, when multiplied by A , results in a vector that is a scalar multiple of itself. In other words, the direction of the vector $\vec{v}$ remains unchanged when the linear transformation represented by A is applied to it.

Eigenvalue:

The scalar multiple, denoted by λ , is called the eigenvalue associated with the eigenvector v . It represents the factor by which the eigenvector is scaled when transformed by the matrix A .

Mathematically, the relationship between a square matrix A , an eigenvector $\vec{v}$, and its corresponding eigenvalue λ is expressed by the following equation:

$$A\vec{v} = \lambda\vec{v}$$

Where $A = f : V \rightarrow V$ is an *endomorphism*, $\vec{v} \neq \vec{0}$ and $\lambda \in \mathbb{C}$.

The way we find a method to find them is by manipulating the equation above

$$\begin{aligned} A\vec{v} &= \lambda\vec{v} \\ A\vec{v} - \lambda\vec{v} &= \vec{0} \\ A\vec{v} - \lambda I\vec{v} &= \vec{0} \\ (A - \lambda I)\vec{v} &= \vec{0} \end{aligned}$$

Here note that if $(A - \lambda I)$ were invertible then we could multiply both sides by its inverse and then get $\vec{v} = \vec{0}$. But we do not want that therefore, $(A\vec{v} - \lambda I)$ must be linear transformation whose determinant is 0 which means that it can not be invertible.

$$\begin{aligned} (A - \lambda I)\vec{v} &= \vec{0} \\ \det(A - \lambda I) &= 0 \end{aligned}$$

45.0.1 Characteristic Polynomial

The polynomial which we get from the determinant $\det(A\vec{v} - \lambda I) = 0$ is called the characteristic polynomial of A .

$$\mathcal{X}_A = \det(A - \lambda I) = 0$$

Eigenspace

The *Eigenspace* of an eigenvalue are the solution of the homogeneous system $A - \lambda I$.

$$\text{Eig}(f; \lambda) = \ker(A - \lambda I)$$

45.0.2 Theorems of Eigenvalues and Eigenvectors

Theorem I

Given an eigenvalue λ of f and $v_1, \dots, v_n$ being the eigenvectors of f with λ then $v \in L(v_1, \dots, v_k) \setminus \{0\}$ is an eigenvector of f with λ .

Theorem II

For $\lambda \in \mathbb{C}$ is $\text{Eig}(f; \lambda) := \{c \in V | f(v) = \lambda v\}$ the *Eigenspace* of f with λ and, it is a subspace of V .

Theorem III

For $\lambda \neq \gamma$ $\text{Eig}(f; \lambda) \cap \text{Eig}(f; \gamma) = \{0\}$.

45.0.3 Linear Independence Of Eigenvectors With Distinct Eigenvalues

Eigenvectors with different eigenvalues are linearly independent.

Proof:

If $\alpha v + \beta w = 0$, where v and w are eigenvector of $A \in \mathbb{C}^{n \times n}$; then it must be true that

$$\begin{aligned}\alpha \lambda v + \beta \lambda w &= 0 \\ \alpha \mu v + \beta \mu w &= 0\end{aligned}$$

With $\lambda \neq \mu$ being eigenvalues of A .

Also we have that

$$0 = A(\alpha v + \beta w) = \alpha \lambda v + \beta \mu w$$

By subtracting this equation to the previous two equations we get

$$\begin{aligned}0 &= \alpha(\lambda - \mu)v \\ 0 &= \beta(\mu - \lambda)w\end{aligned}$$

But since $\mu \neq \lambda$ and $v, w \neq \vec{0}$ it must be that $\alpha = \beta = 0$. Thus, they are linearly independent.

□◻◻

45.0.4 The Determinant Via Eigenvalues

Given the following eigenvalues $\lambda_1, \dots, \lambda_n$, then the determinant of the matrix A is given by

$$\det(A) = \prod_{i=1}^n \lambda_i$$

Proof:

Given the eigenvalues $\lambda_1, \dots, \lambda_n$ are the roots of the characteristic polynomial

$$\begin{aligned}
\det(A - \lambda I) &= p(\lambda) = (-1)^n (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots \\
&= (-1)(\lambda - \lambda_1)(-1)(\lambda - \lambda_2) \cdots \\
&= (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots
\end{aligned}$$

Now, by setting $\lambda = 0$ we get

$$\det(A) = \prod_{i=1}^n \lambda_i$$

Or by using the definition of the diagonal matrix:

$$\begin{aligned}
A &= XDX^{-1} \\
\det(A) &= \det(XDX^{-1}) \\
\det(A) &= \det(X) \det(D) \det(X^{-1}) \\
\det(A) &= \det(D) = \prod_{i=1}^n \lambda_i
\end{aligned}$$

QED

45.0.5 Eigenvalues Under Isometric Maps

For all linear isometric map f with some eigenvalue λ it applies that

$$\|x\| = \|f(x)\| = \|\lambda x\| = \|\lambda\| \|x\|$$

Also, $\lambda = 1$.

An example for this are rotation matrices which map to the same space and preserve the length and whose eigenvalue are complex with a modulus of length of one.

45.0.6 Eigenvalues As Powers

For some eigenvector x with an eigenvalue λ of A . It applies that

$$A^k x = \lambda^k x$$

45.0.7 How to find the Eigenvectors and Eigenvalues

To find the eigenvalues and eigenvectors of a square matrix A , we solve the eigenvalue equation:

1. Form the characteristic equation:

Rewrite the equation $A\vec{v} = \lambda\vec{v}$ as $(A - \lambda I)\vec{v} = \vec{0}$, where λI is the identity matrix times λ because this matrix encodes the multiplication by some scalar.

To have a non-trivial solution $\vec{0}$ for $\vec{v}$, the matrix $(A - \lambda I)$ must be singular like said before, which means its determinant must be zero because otherwise $\vec{v}$ would be $\vec{0}$. Thus, we have the characteristic equation:

$$\det(A - \lambda I) = 0$$

2. Solve for the eigenvalues:

Solve the characteristic equation for λ . The solutions $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of the matrix A .

3. Find the eigenvectors:

For each eigenvalue λ_i , substitute it back into the equation $(A - \lambda_i I)\vec{v} = \vec{0}$ and solve for the vector $\vec{v}$. The non-zero solutions for $\vec{v}$ are the eigenvectors corresponding to the eigenvalue λ_i . Note that we are going to get infinite solutions thus, we have to write our vector with dependence on the free parameters.

Example:

Find the eigenvalues and vectors of

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

Let us build the characteristic equation and solve it

$$\det \begin{bmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{bmatrix} = 0$$

$$(1 - \lambda)^2 - 4 = 0$$

$$1 - 2\lambda + \lambda^2 - 4 = 0$$

$$\lambda^2 - 2\lambda - 3 = 0$$

$$(\lambda - 3)(\lambda + 1) = 0$$

We have got the eigenvalues $\lambda_1 = 3$ and $\lambda_2 = -1$. Now let us continue by substituting our eigenvalues in the original equation.

$$\begin{bmatrix} 1 - \lambda_2 & 1 \\ 4 & 1 - \lambda_2 \end{bmatrix} = \vec{0}$$

$$\begin{bmatrix} 1 - 2 & 1 \\ 4 & 1 - 2 \end{bmatrix} = \vec{0}$$

$$\begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} = \vec{0}$$

Using Gaussian elimination we get:

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix}$$

Thus, let $v_2 = -2$ and $v_1 = \frac{2}{2} = 1$. This give us the first eigenvector and all of it scalar versions.

$$e_1 = \left\{ c \begin{bmatrix} 1 \\ -2 \end{bmatrix} \mid c \in \mathbb{R} \right\}$$

The process for the other eigenvalue is exactly the same.

45.0.8 Complex Eigenvalues

This is a special case of *eigenvalues* which are not as easy to visualize.

Example:

Find the eigenvalues and vectors of

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$$

$$\det \left(\begin{bmatrix} -\lambda & 1 \\ -1 & -\lambda \end{bmatrix} \right) = 0$$

$$\lambda^2 + 1 = 0$$

$$\lambda = \pm i$$

Now the corresponding eigenvectors.

For $\lambda = i$

$$\begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \rightarrow \begin{bmatrix} 1 & i \\ 0 & 0 \end{bmatrix}$$

Therefore, our first eigenvector is

$$c \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

For $\lambda = -i$

$$\begin{bmatrix} i & 1 \\ -1 & i \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -i \\ 0 & 0 \end{bmatrix}$$

Therefore, the second eigenvector is

$$c \begin{bmatrix} i \\ 1 \end{bmatrix}$$

45.0.9 Diagonal Matrices

A *diagonal matrix* is matrix of dimensions $n \times n$ where only the entries of the main diagonal are not zero. This kind of matrices make a lot of the computations in linear algebra trivial. Therefore, if our matrix is not diagonal we are interested in finding a way of transforming the original matrix into a diagonal matrix.

A matrix is diagonalisable if and only if the eigenvectors of it are linear independent.

Complex Diagonalization

Given a matrix $A \in \mathbb{C}^{n \times n}$ with n distinct eigenvalues then is A diagonalisable. But this does not guaranty the diagonalisability for $\mathbb{R}$.

Geometric and Algebraic Multiplicity

Given a matrix $A \in \mathbb{C}^{n \times n}$ with λ_i as an eigenvalue:

The multiple occurrences of the zero λ_i in $\mathcal{X}_A$ is called *algebraic multiplicity* $\text{alg}(\lambda)_i$.

Continuing, $\text{geo}(\lambda_i) := \dim(\text{Eig}(A; \lambda_i))$ is the *geometrical multiplicity* of λ_i i.e. the number of linearly independent eigenvectors associated with λ_i .

Also, A is diagonalisable if and only if the *geometric multiplicity* of every eigenvalue is equal to its *algebraic multiplicity*.

Example:

This example shows a matrix which is not diagonalisable:

$$\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

The corresponding multiplicities are:

$$\begin{aligned}\mathcal{X} &= \det(A - \lambda I) = 0 \\ (1 - \lambda)^2 &= 0\end{aligned}$$

With eigenvalue $\lambda = 1$.

This results into $\text{alg}(1) = 2$ and the $\text{geo}(1) = 1$ because of

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

More precisely $\dim(\text{Eig}(A; 1)) = 1$.

Example With Complex Eigenvalues

Given is the rotation matrix

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

With the characteristic polynomial

$$\begin{aligned}\mathcal{X} &= (\cos(\theta) - \lambda)^2 + \sin^2(\theta) = 0 \\ &= \cos^2(\theta) - 2\cos(\theta)\lambda + \lambda^2 + \sin^2(\theta) = 0 \\ &= \lambda^2 - 2\cos(\theta)\lambda + 1 = 0\end{aligned}$$

Using the quadratic formula gives us

$$\begin{aligned}\lambda_{1,2} &= \cos(\theta) \pm \sqrt{\cos^2(\theta) - 1} \\ &= \cos(\theta) \pm \sqrt{-\sin^2(\theta)} \\ &= \cos(\theta) \pm i\sin(\theta) = e^{\pm i\theta}\end{aligned}$$

Therefore, our eigenvalues are complex. Now, is the matrix diagonalisable? The answer is yes.

Because of $(\cos(\theta) - \lambda)^2$ gives us $\text{alg}(\lambda) = 2$.

$$\text{alg}(e^{i\theta}) = 2, \quad \text{alg}(e^{-i\theta}) = 2$$

And

$$\text{geo}(e^{i\theta}) = 2, \quad \text{geo}(e^{-i\theta}) = 2$$

For $\lambda = e^{i\theta}$ we get two linearly independent vectors

$$\begin{bmatrix} \cos(\theta) - \cos(\theta) - i\sin(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \cos(\theta) - i\sin(\theta) \end{bmatrix}$$

$$\begin{bmatrix} -i \sin(\theta) & -\sin(\theta) \\ \sin(\theta) & -i \sin(\theta) \end{bmatrix}$$

For $\lambda = e^{-i\theta}$ we also get two linearly independent vectors

$$\begin{bmatrix} \cos(\theta) - \cos(\theta) + i \sin(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \cos(\theta) + i \sin(\theta) \end{bmatrix}$$

$$\begin{bmatrix} i \sin(\theta) & -\sin(\theta) \\ \sin(\theta) & i \sin(\theta) \end{bmatrix}$$

Eigenvalues of Real Symmetric Matrices

Eigenvalues of real symmetric matrices are real and also they respective eigenvectors.

Proof:

By definition, we have that $Ax = \lambda x$. Let us multiply both sides by $\bar{x}^T$. Note that because x is a real vector then the conjugate does not change anything.

$$\begin{aligned} Ax &= \lambda x \\ \bar{x}^T Ax &= \bar{x}^T \lambda x \\ \bar{x}^T Ax &= \lambda \bar{x}^T x \\ \bar{x}^T Ax &= \lambda \langle \bar{x} x \rangle \\ \bar{x}^T Ax &= \lambda \|x\|^2 \\ \frac{\bar{x}^T Ax}{\|x\|^2} &= \lambda \end{aligned}$$

With this we know that the denominator is a real number thus, if we prove that the numerator is also real we are done. For this we will use the properties of the conjugate.

$$\begin{aligned} \bar{x}^T Ax &= \overline{\bar{x}^T Ax} \\ &= \overline{\bar{x}^T} \overline{Ax} \\ &= x^T \overline{Ax} \\ &= (\overline{Ax})^T x \\ &= (\bar{A} \bar{x})^T x \\ &= \bar{x}^T \bar{A}^T x \end{aligned}$$

Notice that we ended just where we started. Therefore, the conjugate had no effect on whole term, thus it is real-valued.

□◻◻

Orthogonality of Eigenvectors

Given a symmetric matrix $A \in \mathbb{R}^{n \times n}$ with $\lambda \neq \mu$ as two eigenvalues of A with eigenvectors v and w . Then v and w are *orthogonal* to each other.

Proof:

$$\langle w, Av \rangle = w^T Av = (A^T w)^T v = (Aw)^T v = \langle Aw, v \rangle$$

Therefore, $\lambda \langle v, w \rangle = \langle Av, w \rangle = \langle v, Aw \rangle = \mu \langle v, w \rangle$. Because $\mu \neq \lambda$ and $\lambda - \mu \langle v, w \rangle = 0$ the vectors must be orthogonal.

Eigenvalue Decomposition of Symmetric Matrices

Given a symmetric matrix $A \in \mathbb{R}^{n \times n}$ with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. Then because of the orthogonality of its eigenvectors and the fact that they can be normalized write the matrix Q as an orthogonal matrix, and therefore

$$A = QDQ^T$$

Matrix Diagonalization

Diagonalizing a matrix involves finding a diagonal matrix that is similar to the given matrix. A square matrix A is diagonalisable if there exists an invertible matrix X made of eigenvectors of A such that $X^{-1}AX = D$, where D is a diagonal matrix made of eigenvalues.

The order of the eigenvalue and eigenvectors matters if you choose $\lambda_1, \lambda_2, \dots, \lambda_n$ then the order of the eigenvectors $v_1, v_2, \dots, v_n$ must also correspond. This process only works if A has unique eigenvalues or if they are duplicates they have to be linearly independent.

The process of diagonalisation is as follows:

1. Find the eigenvalues and eigenvectors of A .

2. Form the matrix X :

Create a matrix X whose columns are the linearly independent eigenvectors of A .

3. Form the diagonal matrix D :

Create a diagonal matrix D whose diagonal entries are the eigenvalues of A , corresponding to the order of the eigenvectors in P . That is, if the i -th column of P is the eigenvector corresponding to the eigenvalue λ_i , then the i -th diagonal entry of D is λ_i .

4. Verify the diagonalisation:

Check that $X^{-1}AX = D$.

A matrix A is diagonalisable if and only if it has n linearly independent eigenvectors, where n is the size of the matrix.

Example:

Consider the matrix

$$A = \begin{bmatrix} -3 & -4 \\ 5 & 6 \end{bmatrix}$$

The characteristic equation is

$$\det(A - \lambda I) = \det \begin{bmatrix} -3 - \lambda & -4 \\ 5 & 6 - \lambda \end{bmatrix} = (-3 - \lambda)(6 - \lambda) + 20 = \lambda^2 - 3\lambda + 2 = 0$$

$$(\lambda - 1)(\lambda - 2)$$

Solving for λ , we get $\lambda_1 = 1$ and $\lambda_2 = 2$.

For $\lambda_1 = 1$:

$$(A - I)\vec{v} = \begin{bmatrix} -4 & -4 \\ 5 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} -4 & -4 & 0 \\ 5 & 5 & 0 \end{array} \right] \Rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 5 & 5 & 0 \end{array} \right] \Rightarrow \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Our free variable is y therefore

$$\vec{v} = \left\{ y \begin{bmatrix} -1 \\ 1 \end{bmatrix} \mid y \in \mathbb{R} \right\}$$

Now choose $y = 1$ which gives us $x = -1$. Now we get the vector

$$\vec{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

For $\lambda_2 = 2$ the process is similar, so we are going to skip it. We get the vector

$$\vec{y} = \begin{bmatrix} -\frac{4}{5} \\ 1 \end{bmatrix}$$

Now we will complete the process by writing D , X , and X^{-1} .

$$D = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} X = \begin{bmatrix} -1 & -\frac{4}{5} \\ 1 & 1 \end{bmatrix} X^{-1} = \begin{bmatrix} -1 & \frac{4}{5} \\ -1 & -1 \end{bmatrix}$$

Normal Matrices

A matrix $A \in \mathbb{C}^{n \times n}$ is considered *normal* if and only if

$$A\overline{A^T} = \overline{A^T}A$$

For real matrices, this condition simplifies to $AA^T = A^T A$.

Also, then there exist with the euclidean inner product in $\mathbb{C}^n$ an orthonormal basis of eigenvector and therefore, A is diagonalisable.

This implies that

1. All real symmetric matrices are diagonalisable, and their eigenvalues are real.
2. All real antisymmetric matrices $A^T = -A$ are diagonalisable, and their eigenvalues are purely imaginary or 0.
3. All orthogonal matrices are diagonalisable.

45.0.10 Singular Value Decomposition (SVD)

The *Singular Value Decomposition (SVD)* is a fundamental matrix factorization technique in linear algebra. It expresses any real matrix $A \in \mathbb{R}^{n \times n}$ as a product of three specific matrices:

$$A = USV^T$$

where:

- $U \in \mathbb{R}^{n \times n}$ is an orthogonal matrix whose columns are called the *left singular vectors* of A . A rotation.
- $V \in \mathbb{R}^{n \times n}$ is an orthogonal matrix whose columns are the *right singular vectors* of A . A stretching with dimension reduction.
- $S \in \mathbb{R}^{n \times n}$ is a diagonal matrix with non-negative entries $\sigma_i \geq 0$ on the diagonal, called the *singular values* of A . Another rotation.

Example:

Let $A \in \mathbb{R}^{n \times n}$ be a real square matrix with SVD given by:

$$A = USV^T$$

We show that:

1. The columns of U are eigenvectors of AA^T ,
2. The columns of V are eigenvectors of $A^T A$,
3. The eigenvalues of both AA^T and $A^T A$ are the squares of the singular values σ_i^2 .

Proof:

For AA^T , we compute:

$$AA^T = (USV^T)(VS^T U^T) = USV^T VS^T U^T = USS^T U^T = US^2 U^T$$

Since SS^T is diagonal with entries σ_i^2 , this is an eigenvalue decomposition of AA^T . Thus, the columns of U are eigenvectors of AA^T , and the corresponding eigenvalues are σ_i^2 .

For $A^T A$, we have:

$$A^T A = (VS^T U^T)(USV^T) = VS^T U^T USV^T = VS^T SV^T = VS^2 V^T$$

Again, $S^T S$ is diagonal with entries σ_i^2 , so the matrix $A^T A$ is diagonalized by V , and the columns of V are eigenvectors of $A^T A$, with eigenvalues σ_i^2 .

□◻◻

45.0.11 Similar Matrices

Given $A \in \mathbb{C}^{n \times n}$ and $X \in \mathbb{C}^{n \times n}$ then if

$$A = XDX^{-1} \iff X^{-1}AX = D$$

we denote XDX^{-1} as *similar* to A .

Case I

A is diagonalisable with eigenvalues and eigenvectors.

$$S^{-1}AS = D = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \ddots & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & & \lambda_n \end{bmatrix}$$

Case II

Any complex eigenvalue must come in complex conjugate pairs if A is real valued.

$$\lambda_1 = a + ib, \lambda_2 = a - bi$$

$$S^{-1}AS = D = \begin{bmatrix} a & b \\ -b & a \end{bmatrix} \rightarrow e^{At} = \begin{bmatrix} \cos(bt) & \sin(bt) \\ -\sin(bt) & \cos(bt) \end{bmatrix} e^{at}$$

Case III

Repeated eigenvalues and not enough vector in the null-space.

$$S^{-1}AS = D = \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}$$

Equality Of Eigenvalues

Given $A \in \mathbb{C}^{n \times n}$ and $X \in \mathbb{C}^{n \times n}$, both regular, then XAX^{-1} has the same eigenvalues as A .

Proof:

$$\begin{aligned} \det(A - \lambda I) &= \det(XAX^{-1} - \lambda I) \\ &= \det(XAX^{-1} - \lambda XIX^{-1}) \\ &= \det(X[A - \lambda I]X^{-1}) \\ &= \det(X) \det(A - \lambda I) \det(X^{-1}) \\ &= \det(A - \lambda I) \\ &= \mathcal{X}_A \end{aligned}$$

Thus, similar matrices have the same **eigenvalues with their correspondent multiplicity**.

QED

45.0.12 Jordan Normal Form

Let $A \in \mathbb{C}^{n \times n}$, it is diagonalisable if

$$X^{-1}AX = D \iff A = XDX^{-1}$$

where D is a diagonal matrix.

Generalizing, we can also find a matrix $J \in \mathbb{C}^{n \times n}$ in the *Jordan normal form* such that $A = SJS^{-1}$. The matrix J is then considered the most not diagonalisable matrix similar to our matrix A . This comes in handy when we have repeated eigenvalues with enough eigenvectors.

$$A = XJX^{-1}, J = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_n \end{bmatrix}$$

Whereas J is a *block-diagonal matrix*

$$J_i = \begin{bmatrix} \lambda_i & 1 & & \\ & \ddots & \ddots & \\ & & \lambda_i & 1 \\ & & & \lambda_i \end{bmatrix}$$

For each of these blocks the one corresponding eigenvalue is repeated across the diagonal

If A has exactly n linear independent eigenvectors then J is just D .

Algorithm For Jordan Blocks

- Determine the eigenvalues of A and their algebraic multiplicities.
- For each eigenvalue, determine the size of the largest Jordan block by finding the smallest k such that $\text{nullity}((A - \lambda I)^k) = \text{nullity}((A - \lambda I)^{k+1})$.
- Construct the Jordan normal form J by placing the Jordan blocks corresponding to each eigenvalue along the diagonal, with the size of each block determined in the previous step.

Example:

Given the matrix

$$A = \begin{bmatrix} 5 & 4 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 3 \end{bmatrix}$$

The characteristic polynomial is

$$\mathcal{X}_A = (5 - \lambda)(1 - \lambda)(3 - \lambda)$$

Delivering the eigenvalues $\lambda_1 = 5$, $\lambda_2 = 1$, and $\lambda_3 = 3$ with algebraic multiplicity of 1 for each.

Hence the Jordan normal form of A is

$$J = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{bmatrix}$$

If we had a matrix with repeated eigenvalues and not enough eigenvectors, for example

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$$

The characteristic polynomial is

$$\mathcal{X}_A = (2 - \lambda)^2$$

Delivering the eigenvalue $\lambda = 2$ with algebraic multiplicity of 2. However, there is only one linearly independent eigenvector, so the Jordan normal form of A is

$$J = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$$

Note that the Jordan block has a size equal to the algebraic multiplicity of the eigenvalue, and the superdiagonal entry is 1, indicating that there is a single Jordan block for the eigenvalue 2.

45.0.13 The Hermitian Transpose

For a matrix $A \in \mathbb{C}^{n \times m}$, the *hermitian transpose* A^H is the transpose of the conjugate matrix of a matrix A .

$$A^H = \bar{A}^T$$

45.0.14 Unitary Matrices

Unitary Matrices are matrices whose columns form orthonormal vectors and

$$U^H = U^{-1}$$

45.0.15 Unitary Decomposition (Schur's Theorem)

For each matrix $A \in \mathbb{C}^{n \times n}$, there exist a unitary matrix $U \in \mathbb{C}^{n \times n}$ such that

$$A = UTU^H, \quad T = \begin{bmatrix} \lambda_1 & t_{1,2} & \cdots & t_{1,n} \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & t_{n-1,n} \\ 0 & \cdots & 0 & \lambda_{n,n} \end{bmatrix}$$

As a reminder $U^H = \overline{U^T}$.

Unitary Diagonalisation

For a matrix $A \in \mathbb{C}^{n \times n}$, it is considered *unitary diagonalisable* if there exist a matrix $U \in \mathbb{C}^{n \times n}$ such that

$$A = U\Lambda U^H, \quad \Lambda = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \lambda_{n,n} \end{bmatrix}$$

45.0.16 Normal Matrices

A matrix $A \in \mathbb{C}^{n \times n}$ is considered *normal* if $AA^H = A^H A$

45.0.17 Diagonalisability Criteria Based On Normal Matrices

A matrix $A \in \mathbb{C}^{n \times n}$ is unitary diagonalisable if and only if it is normal.

Proof:

A is unitary diagonalisable $\Rightarrow A$ is normal:

Then there exists a unitary matrix U such that $A = U\Lambda U^H$. Therefore,

$$A^H A = (U\Lambda U^H)^H (U\Lambda U^H) = U\Lambda^H U^H U\Lambda U^H = U\Lambda^H \Lambda U^H$$

and

$$AA^H = (U\Lambda U^H)(U\Lambda U^H)^H = U\Lambda U^H U\Lambda^H U^H = U\Lambda\Lambda^H U^H$$

Hence, $AA^H = A^H A$, so A is normal.

A is normal $\Rightarrow A$ is unitary diagonalisable:

By Schur's theorem, there exists a unitary matrix U such that $A = UTU^H$, where T is upper triangular.

Since A is normal, we have:

$$AA^H = UTU^H (UT^H U^H) = UTT^H U^H$$

and

$$A^H A = (UTU^H)^H (UTU^H) = UT^H U^H UTU^H = UT^H T U^H$$

Since $AA^H = A^H A$, it follows that $TT^H = T^H T$. This implies that T is a normal upper triangular matrix, which must be diagonal. Therefore, $T = \Lambda$ is diagonal, and we have $A = U\Lambda U^H$, showing that A is unitary diagonalisable.

Example

The matrix

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 2 & 0 & 1 \end{bmatrix}$$

is normal and unitary diagonalisable. We can verify that $AA^H = A^H A$ and then find a unitary matrix U such that $A = U\Lambda U^H$, where Λ is a diagonal matrix containing the eigenvalues of A .

$$\Lambda = \begin{bmatrix} 3 & 0 & 0 \\ 0 & -\sqrt{3}i & 0 \\ 0 & 0 & \sqrt{3}i \end{bmatrix} \quad U = \begin{bmatrix} \frac{\sqrt{3}}{3} & -\frac{\sqrt{3}}{6} - \frac{i}{2} & -\frac{\sqrt{3}}{6} + \frac{i}{2} \\ \frac{\sqrt{3}}{3} & -\frac{\sqrt{3}}{6} + \frac{i}{2} & -\frac{\sqrt{3}}{6} - \frac{i}{2} \\ \frac{\sqrt{3}}{3} & \frac{\sqrt{3}}{3} & \frac{-1+i\sqrt{3}}{2\sqrt{3}} \end{bmatrix}$$

45.0.18 Properties of Unitary Diagonalisable Matrices

For matrices $A \in \mathbb{C}^{n \times n}$ that are unitary diagonalisable $A = U\Lambda U^H$, the following properties hold:

- For all eigenvalues λ_i its geometric and algebraic multiplicity are the same.
- The column vectors of U form an orthonormal basis of $\mathbb{C}^n$. This means that for a vector x

$$x = \sum_{i=1}^n x_i u_i, \quad x_i = \langle u_i, x \rangle_2 = u_i^H x$$

and

$$Ax = \sum_{i=1}^n x_i A u_i = \sum_{i=1}^n x_i \lambda_i u_i$$

- For any vector $x, y \in \mathbb{C}^n$ we have that:

$$\langle x, y \rangle_2 = x^H y = \sum_{i=1}^n \bar{x}_i y_i, \quad \|x\|_2^2 = \sum_{i=1}^n |x_i|^2$$

45.0.19 Hermitian Matrices

A matrix $M \in \mathbb{C}^{n \times m}$ with the following property:

$$M^H = M$$

is considered *hermitian*.

Properties of Hermitian Matrices

- All eigenvalues of a hermitian matrix are real.
- Hermitian matrices are normal and therefore unitary diagonalisable.
- Symmetric matrices are a special case of hermitian matrices where all entries are real.

45.0.20 Real Eigenvalues Via Hermitian Matrices

Given a normal matrix $A \in \mathbb{C}^{n \times n}$, the all eigenvalues of A are real if and only if A is hermitian.

Proof:

$\Rightarrow$: A is normal which means that it is unitary diagonalisable. Therefore, there exists a unitary matrix U such that $A = U\Lambda U^H$, where Λ is a diagonal matrix containing the eigenvalues of A .

Since all eigenvalues are real, Λ is a real diagonal matrix. Thus, $A^H = (U\Lambda U^H)^H = U\Lambda^H U^H = U\Lambda U^H = A$, showing that A is hermitian.

$\Leftarrow$: If A is hermitian, we know that $A^H = A$. Since A is also normal, it is unitary diagonalisable. Therefore, there exists a unitary matrix U such that $A = U\Lambda U^H$, where Λ is a diagonal matrix containing the eigenvalues of A .

$$A^H = (U\Lambda U^H)^H = U\Lambda^H U^H = U\Lambda U^H = A$$

Since, $\Lambda^H = \Lambda$, follows that Λ is a real diagonal matrix, and thus all eigenvalues of A are real.

□□□

45.0.21 Gerschgorin's Theorem

Given $A \in \mathbb{C}^{n \times n}$ and

$$K = \bigcup_{i=1}^n K_i, \quad K_i = \{z \in \mathbb{C} : |z - a_{i,i}| \leq R_i\}, \quad R_i = \sum_{j \neq i} |a_{i,j}|$$

Then, all eigenvalues of A are contained in K , $\lambda_i \in K \forall i$.

Example:

45.0.22 Approximation Of The Differences Of Eigenvalues

Given a symmetric matrix $A \in \mathbb{R}^{n \times n}$ with distinct eigenvalues and $c \in \mathbb{R}$ and $x \in \mathbb{R}^n \setminus \{\vec{0}\}$, then

$$\min_{i=1, \dots, n} |\lambda_i - c| \leq \frac{\|Ax - cx\|}{\|x\|} = \frac{\|d\|}{\|x\|}$$

Proof:

Because of the distinct eigenvalues of A we know that there exists an eigenbasis such that we can write x as

$$\sum_{i=1}^n \alpha_i v_i$$

We can write

$$Ax = \sum_{i=1}^n \alpha_i A v_i$$

Because of the eigenbasis we get

$$Ax = \sum_{i=1}^n \alpha_i \lambda_i v_i$$

Resulting into

$$Ax - cx = \sum_{i=1}^n \alpha_i \lambda_i v_i - \sum_{i=1}^n \alpha_i c v_i$$

Which results into

$$\sum_{i=1}^n \alpha_i v_i (\lambda_i - c) = d$$

Using the inner product

$$\|d\|_2^2 = \sum_{i=1}^n \alpha_i^2 (\lambda_i - c)^2$$

Turning our focus to the minimal of the distance $\min_{i=1,\dots,n} |\lambda_i - c|$, let us define m as the number associated with the minimum.

This gives us the following:

$$\begin{aligned} \sum_{i=1}^n \alpha_i^2 (\lambda_i - c)^2 &\geq m^2 \sum_{i=1}^n \alpha_i^2 \\ \frac{\|d\|^2}{\|x\|^2} &\geq m^2 \\ \frac{\|d\|}{\|x\|} &\geq \min_{i=1,\dots,n} |\lambda_i - c| \end{aligned}$$

□◻◻

45.0.23 Eigenvalue Of Block Matrices

Given a block matrix

$$A = \begin{bmatrix} B & C \\ 0 & D \end{bmatrix} \in C^{n \times n}, \quad B \in C^{p \times p}, \quad C \in C^{p \times (n-p)}, \quad D \in C^{(n-p) \times (n-p)}$$

Its eigenvalues are the eigenvalues of B and D .

Proof:

This fact comes from the determinant of a block matrix

$$\det(A - \lambda I) = \det(B - \lambda I) \cdot \det(D - \lambda I)$$

□◻◻

45.0.24 Roth's Removal Rule

The matrices

$$A = \begin{bmatrix} B & C \\ 0 & D \end{bmatrix} \quad \tilde{A} = \begin{bmatrix} B & 0 \\ 0 & D \end{bmatrix}$$

are similar

$$A = V \tilde{A} V^{-1}$$

if and only if the *Sylvester equation* has a solution.

$$BX - XD = C$$

In such a case, the matrix V is given by

$$V = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix}, \quad V^{-1} = \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix}$$

45.0.25 Eigenvalues/vectors of Matrices Representing Complex Numbers

Given a complex number $z = a + ib$ we can represent it as a matrix

$$A = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

Its eigenvalues are z and $\bar{z}$ and the corresponding eigenvectors are

$$v_z = \begin{bmatrix} \frac{a}{b} - \frac{a-ib}{b} \\ 1 \end{bmatrix}, \quad v_{\bar{z}} = \begin{bmatrix} \frac{a}{b} - \frac{a+ib}{b} \\ 1 \end{bmatrix}$$

45.0.26 Real Jordan Form

For every matrix $A \in \mathbb{R}^{n \times n}$ there exists a regular matrix $V \in \mathbb{R}^{n \times n}$ such that

$$A = VJV^{-1}, \quad J = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_k \end{bmatrix} \quad J_i = \begin{bmatrix} \Lambda_i & I & & \\ & \ddots & \ddots & \\ & & \Lambda_i & I \\ & & & \Lambda_i \end{bmatrix}$$

For $\lambda_i \in \mathbb{R}$ is $\Lambda_i = \lambda_i$ else

$$\Lambda_i = \begin{bmatrix} a_i & b_i \\ -b_i & a_i \end{bmatrix}$$

where $\lambda_i = a_i \pm jb_i \in \mathbb{C}$ are the complex conjugate eigenvalues of A .

45.0.27 Real Schur Form

For every matrix $A \in \mathbb{R}^{n \times n}$ there exists an orthogonal matrix $Q \in \mathbb{R}^{n \times n}$ such that

$$A = QTQ^T, \quad T = \begin{bmatrix} T_{1,1} & T_{1,2} & \cdots & T_{1,m} \\ 0 & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & T_{m-1,m} \\ 0 & 0 & 0 & T_{m,m} \end{bmatrix}$$

where $T_{i,i} \in \mathbb{R}$ or $T_{i,i} \in \mathbb{R}^{2 \times 2}$.

45.0.28 Orthonormal Standard Form

A matrix $A \in \mathbb{R}^{2 \times 2}$ with complex eigenvalues $\lambda = a + bi$, $\bar{\lambda} = a - bi$ is then transformable to the *orthonormal standard form*

$$T = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

If A is normal.

45.0.29 Criterion For Orthonormal Diagonalisation

A matrix $A \in \mathbb{R}^{n \times n}$ is orthonormal diagonalisable i.e

$$A = QDQ^T, \quad D = \begin{bmatrix} \Lambda_1 & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & 0 \\ 0 & \cdots & 0 & \Lambda_n \end{bmatrix}, \quad \Lambda_i \in \mathbb{R} \text{ or } \Lambda_i = \begin{bmatrix} a_i & b_i \\ -b_i & a_i \end{bmatrix}$$

if and only if A is normal, i.e. $AA^T = A^T A$.

Example:

The matrix

$$A = \begin{bmatrix} 41 & 12 & 60 \\ 12 & 34 & 45 \\ -60 & 45 & 50 \end{bmatrix}$$

is normal; hence it is unitary diagonalisable $A = U\Lambda U^H$.

Its eigenvalues are $\lambda = 25$, $\lambda = 50 - 75i$ and $\lambda = 50 + 75i$.

$$\Lambda = \begin{bmatrix} 25 & 0 & 0 \\ 0 & 50 - 75i & 0 \\ 0 & 0 & 50 + 75i \end{bmatrix}$$

The eigenvectors will be skipped for the sake of brevity, but they are 3.

We can diagonalize A as follows stated in the criterion for orthonormal diagonalisation.

$$\Lambda = \begin{bmatrix} 25 & 0 & 0 \\ 0 & 50 & -75 \\ 0 & 75 & 50 \end{bmatrix}$$

Note that the complex eigenvalues are represented as 2×2 blocks in the diagonal of Λ .

Then Q is given by the matrix of the corresponding eigenvectors of A and we can verify that $A = Q\Lambda Q^T$.

$$Q = \begin{bmatrix} \frac{3}{5} & 0 & \frac{4}{5} \\ -\frac{4}{5} & 0 & -\frac{3}{5} \\ 0 & 1 & 0 \end{bmatrix}$$

45.0.30 Normal Property Of Symmetric Matrices

All real symmetric matrices are normal, and therefore they are orthonormal diagonalisable. And also all of their eigenvalues are real by the properties of hermitian matrices.

Chapter 46

Dual Spaces and Dual Vectors

Given a $\mathbb{F}$ -vector space V , we define V^* *Dual Space* of V , as the set of linear transformations from $f : V \rightarrow K$. f is called the *Linear Functional*.

The vector inside the space V^* are called *dual vectors* or *covariant vectors* while vectors in the space V are called *contravariant vectors*.

Example:

$V = \mathbb{R}^3$, with $f : V \rightarrow \mathbb{R}$ and $f(x, y, z) = x + y + z$

46.0.1 The Dual Basis

Given a finite dimensional $\mathbb{F}$ -vector space V . Let $\beta = \{v_1, \dots, v_n\}$ be a basis of V let $f \in V^*$. The set of functionals $\beta^* = \{f_1, \dots, f_n\}$ of V^* . Defined by:

$$f_i(v_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases} = \delta_{ij}$$

Is called a *Dual Basis*. The definition above is saying that only one of the basis vector of V is going to be associated with a function of the basis β .

Proof:

To prove a set of vectors is basis we have to prove that they are linearly independent and that they span the entire vector space.

Linear Independence:

We want that

$$\sum_{i=1}^n \lambda_i f_i(v) = 0 \text{ only if all } \lambda_i = 0 \text{ for all } v \in V$$

Now just choose n -times $v = v_j, 1 \leq j \leq n$ and you are going to realize that because $f_i(v_j) = 1$ only if $j = i$ you are going to get $\lambda_j f_j(v_j) + 0 + \dots + 0 = 0$. Therefore, by repeating this process you will arrive at the conclusion that all lambdas have to zero. Thus, the linear Independence is shown.

Span:

Suppose $f \in V^*$ is arbitrary and $f = \sum_{i=1}^n \lambda_i f_i(v)$ for all $v \in V$. Using the previously collected knowledge, which tells us that for $v = v_j$ the linear combination of $\sum_{i=1}^n \lambda_i f_i(v_j)$ is

$$\sum_{i=1}^n \lambda_i f_i(v_j) = \lambda_j f_j(v_j) = \lambda_j$$

Now let us guess $f = f(v_1)f_1 + \cdots f(v_n)f_n = g$. If $f(v_i) = g(v_i)$ for all basis vector of V .

Note for

$$g(v_i) = f(v_1)f_1(v_i) + \cdots + f(v_i)f_i(v_i) + \cdots f(v_n)f_n(v_i)$$

All terms are zero except for $f(v_i)f_i(v_i)$ which equals $f(v_i) = g(v_i)$.

QED

46.0.2 How to find a Dual Basis

We know that to find the coefficients of a linear combination $f = a_1f_1 + a_2f_2 + \cdots + a_nf_n$ in the space V^* with the basis $\beta = \{f_1, \dots, f_n\}$ we just need to evaluate the function f at each of the basis vectors of the space V which are $\{v_1, \dots, v_n\}$.

The process for solving this kind of problems can be summarized into the following steps using multiple systems of equations.

1. For each f_i repeat the following steps except for the last one.
2. Use the linearity property of vector to write non-canonical basis vectors as a linear combination $f_i(\vec{k}) = \sum \lambda_k f_i(\vec{v}_k)$, do this for every vector in the basis of V with the current f_i . This will set up the left-hand side of the system of equations.
3. Use the definition of the dual basis for setting up the right-hand side of linear system
4. Solve the system of equations
5. If a f was given then use the found basis to write $f = \sum_{i=1}^n f(v_i)f_i$

Example:

Given the space $V = \mathbb{R}^n$ with the basis $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}$ find $\beta^* = \{f_1, f_2\}$.

Because of linearity

$$\begin{aligned} f_1(2, 1) &= 2f_1(\vec{e}_1) + 1f_1(\vec{e}_2) = 1 \\ f_1(3, 1) &= 3f_1(\vec{e}_1) + 1f_1(\vec{e}_2) = 0 \end{aligned}$$

We get a linear system of equations. By solving the system we get $f_1(1, 0) = -1$ and $f_2(0, 1) = 3$. Thus, $f_1(x, y) = x(-1) + y(3)$.

And for f_2 it is the same process

$$\begin{aligned} f_2(2, 1) &= 2f_2(\vec{e}_1) + 1f_2(\vec{e}_2) = 0 \\ f_2(3, 1) &= 3f_2(\vec{e}_1) + 1f_2(\vec{e}_2) = 1 \end{aligned}$$

This time we get $f_2 = 2x - 5y$. Thus, our basis is $\beta^* = \{-x + 3y, 2x - 5y\}$, and if we let f be $f = 2x - 5$ we get that it can be decomposed into:

$$\begin{aligned} f &= f(2, 1)f_1 + f(3, 1)f_2 \\ &= (-1)f_1 + 1f_2 \end{aligned}$$

46.0.3 Change of Basis in the Dual Space

Given some finite $\mathbb{F}$ -vector space V with basis β and β' . Then matrix for the change of basis $\beta^* \rightarrow \beta'^*$ is given by

$$(T^{-1})^T = (T^T)^{-1}$$

Where $T = \beta^{-1}\beta'$.

Example:

Given are $\beta = \{(1, 1); (-1, 1)\}$ and $\beta' = \{(2, 3), (1, 1)\}$.

$$T = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 2 & 1 \\ 3 & 1 \end{bmatrix}$$

Then

$$(T^T)^{-1} = \begin{pmatrix} 0 & 1 \\ 2 & -5 \end{pmatrix}$$

46.0.4 Dimension Theorem

If the dimension of V is finite then V and V^* are *isomorphic*.

46.0.5 The Transpose of a Linear Transformation

Let V and W be finite dimensional vector spaces with the field K . Let $T : V \rightarrow W$ be a linear transformation. Now define

$$T^T : W^* \rightarrow V^*$$

with an input $g \in W^*$ where $g : W \rightarrow K$; and an output $T^T(g) \in V^*$ with $T^T(g) : V \rightarrow K$. Therefore, $T^T(g)(\vec{x}) = g(T(\vec{x}))$ for all $\vec{x} \in V$.

46.0.6 Connection to the Transpose Matrix

Let $T : V \rightarrow W$ be a linear transformation, $\beta = \{v_1, \dots, v_n\}$ be a basis of V , $\gamma = \{w_1, \dots, w_n\}$ be a basis of W . The matrix of the change of the transformation is given by $A = [T]_{\beta}^{\gamma}$ which takes vector from β to γ .

Then let $\beta^* = \{f_1, \dots, f_n\}$ be the dual basis of V^* and $\gamma^* = \{g_1, \dots, g_n\}$ the dual basis of W^* . Then

$$[T^T]_{\gamma^*}^{\beta^*} = A^T$$

Which takes functions from γ^* to β^* .

Proof:

Because $T(v_i) \in W$ we can write

$$T(v_i) = \sum_{k=1}^n A_{ki} w_k$$

as a vector in terms of the basis γ .

For $T^T : W^* \rightarrow V^*$ we can write $T^T(g)$ as a linear combination of the basis vectors β^*

$$T^T(g_j) = \sum_{i=1}^n T^T(g_j)(v_i) f_i = \sum_{i=1}^n g(T(v_i)) f_i = \sum_{i=1}^n A_{ij} f_i$$

where $A = (T^T(g_1), \dots, T^T(g_n))$.

We have assumed that $T^T(g)(\vec{x}) = g(T(\vec{x}))$. Now let us apply this to any $v_j \in V$ and $g_i \in W^*$

$$\begin{aligned} T^T(g_j)(v_i) &= g_j(T(v_i)) \\ &= g_j\left(\sum_{k=1}^n A_{ki} w_k\right) = \sum_{k=1}^n A_{ki} g_j(w_k) \\ &= A_{ji} \end{aligned}$$

This sum is equals to A_{ji} for when $g_j(w_j)$. Therefore, we have flipped the columns and rows. Thus, we have shown that

$$[T^T]_{\gamma^*}^{\beta^*} = A^T$$

Because we have shown it for the basis vectors it gets extended to all of it linear combinations.

□◻◻

Example:

Let $P_1\mathbb{R}$ be the space of polynomials of degree at most 1 and $W = \mathbb{R}^2$. Also, let $T : V \rightarrow W$ be defined by $T(p) = \begin{bmatrix} p(0) - 2p(1) \\ p(0) + p'(0) \end{bmatrix}$

$$\text{For example } T(1 + 2x) = \begin{bmatrix} 1 - 2(1 + 2) \\ 1 + 2 \end{bmatrix} = \begin{bmatrix} -5 \\ 3 \end{bmatrix}$$

If $f(a, b) = a - 2b \in W^*$ find $T^T(f)$.

The output of $T^T(f)$ is in V^* and because we are using the transpose we can write it as $T^T(f)(p) = f(T(p))$ and

$$\begin{aligned} f(a, b) &= a - 2b \text{ then} \\ f(T(p)) &= f((p(0) - 2p(1), p(0) + p'(0))) \\ &= (p(0) - 2p(1)) - 2(p(0) + p'(0)) \\ &= p(0) - 2p(1) - 2p(0) - 2p'(0) \\ T^T(f)(p) &= -p(0) - 2p(1) - 2p'(0) \end{aligned}$$

Now calculate the matrix of T and T^T . The basis of V is $\{1, x\}$ and the basis of W is $\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$.

First we know that $\beta^* = \{f_1, f_2\}$ and $\gamma^* = \{g_1, g_2\}$, and we remember that

$$f_i(v_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases} = \delta_{ij}$$

To find the matrix of T^T which is a 2×2

$$T^T = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

We need to evaluate $T^T(g_1)$ and $T^T(g_2)$ and then write them in terms of β^* .

$$T^T(g_1) = af_1 + cf_2$$

Now we evaluate at the basis vectors of V

$$T^T(g_1)(1) = af_1(1) + cf_2(1) = a$$

Now we use the definition of T

$$\begin{aligned} g_1(T(1)) &= g_1\left(\begin{bmatrix} 1-2 \\ 1+0 \end{bmatrix}\right) = g_1\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) \\ &= g_1\left(-1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = -1 \end{aligned}$$

Thus, we have $a = -1$.

Now lets repeat this for x

$$T^T(g_1)(x) = af_1(x) + cf_2(x) = c$$

$$\begin{aligned} g_1(T(x)) &= g_1\left(\begin{bmatrix} 0-2 \\ 0+1 \end{bmatrix}\right) = g_1\left(\begin{bmatrix} -2 \\ 1 \end{bmatrix}\right) \\ &= g_1\left(-2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = -2 \end{aligned}$$

Thus, we have $c = -2$.

Now if you repeat this for g_2 you will get $b = 1$ and $d = 0$.

Thus, the matrix of T^T is

$$T^T = \begin{bmatrix} -1 & 1 \\ -2 & 0 \end{bmatrix}$$

And the matrix of T is calculated by applying $T(1)$ and $T(x)$ and then writing them in terms of the basis γ .

This gives us

$$T(1) = \begin{bmatrix} 1-2 \\ 1+0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\text{Which is } -1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

And

$$T(x) = \begin{bmatrix} 0-2 \\ 0+0 \end{bmatrix} = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$$

$$\text{Which is } -2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0 \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Thus, the matrix of T is

$$T = \begin{bmatrix} -1 & -2 \\ 1 & 0 \end{bmatrix}$$

46.0.7 The Dual of the Dual Space

Given a dual space $V^* = L(V, \mathbb{F})$ then

$$(V^*)^* = L(V^*, \mathbb{F})$$

We also define the linear functional $\varphi : V \rightarrow V^{**}$ where $\varphi(x) = \hat{x}$ and $\hat{x}(f) = f(x)$ for all $f \in V$.

Isomorphism of the Dual Space

If V is finite dimensional then V^{**} and V are isomorphic.

Isomorphism from V to V^{**}

If V is finite dimensional then the map $\phi : V \rightarrow V^{**}$ is isomorphic.

46.0.8 Basis Theorem for the Dual Space

Every basis β^* of V^* is the dual basis of some basis β of V .

Proof:

Let $\beta^* = \{f_1, \dots, f_n\}$ be a basis of V^* . We can define a basis β of V with help of the following:

Let us find a basis $\beta^{**} = \{\mathcal{F}_1, \dots, \mathcal{F}_n\}$ of V^{**} . The functional $\varphi : V \rightarrow V^{**}$ is an isomorphism and $\mathcal{F}_i = \hat{v}_i$ for some $v_i \in V$.

Let us choose a basis $\beta = \{v_1, \dots, v_n\}$ of V and use β^* . By definition

$$f_i(v_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases} = \delta_{ij}$$

Using the isomorphism φ we get

$$f_j(v_i) = \hat{v}_i(f_j) = \mathcal{F}_i(f_j) = \begin{cases} 1 & \text{if } j = i \\ 0 & \text{if } j \neq i \end{cases} = \delta_{ij}$$

□◻◻

46.0.9 Transpose of the Transpose is T

We define $T^{TT}V^{**} \rightarrow W^{**}$ by

$$T^{TT}(\hat{x}) = \hat{x}(T^T)$$

and claim $T^{TT} = T$.

Proof:

In this proof we know that $T^T : W^* \rightarrow V^*$. Now let us declare our basis $\beta = \{v_1, \dots, v_n\}$ for V , $\gamma = \{w_1, \dots, w_n\}$ for W , $\beta^* = \{f_1, \dots, f_n\}$ for V^* , $\gamma^* = \{g_1, \dots, g_n\}$ for W^* , $\beta^{**} = \{\mathcal{F}_1, \dots, \mathcal{F}_n\}$ for V^{**} and finally $\gamma^{**} = \{\mathcal{G}_1, \dots, \mathcal{G}_n\}$ for W^{**} .

When we evaluate at $g : W \rightarrow \mathbb{F}$ we get

$$\begin{aligned}
T^{TT}(\hat{x})(g) &= \hat{x}(T^T)(g) \\
&= \hat{x}(T^T(g)) \text{ by associativity} \\
&= T^T(g)(x) \text{ using the definition of } \hat{x} \\
&= g(T)(x) \text{ by associativity} \\
&= g(T(x)) \\
&= T(\hat{x})(g)
\end{aligned}$$

Here $T(\hat{x}) \in W^{**}$. Finally

$$\begin{aligned}
T^{TT}(\hat{x})(g) &= T(\hat{x})(g) \\
T^{TT}(\hat{x}) &= T(\hat{x}) \\
T^{TT} &= \hat{T} \\
T^{TT} &= T \text{ by ignoring the hats}
\end{aligned}$$

They are not strictly the same, but because of the isomorphism we can set them equal.

□

46.0.10 The Annihilator

We define the *Annihilator* of a subspace of V as

$$W^\circ = \{f \in V^* \mid f(w) = 0, \forall w \in W\}$$

We also get the following formula

$$\dim(W^\circ) + \dim(W) = \dim(V)$$

whose proof is similar to the one of the rank nullity theorem.

Proof:

Let $\{w_1, \dots, w_{k-1}, w_k, w_{k+1}, \dots, w_n\}$ be a basis of W with dimension k . Let us extend the basis to a basis of V by $\beta = \{w_1, \dots, w_{k-1}, w_k, w_{k+1}, \dots, w_n\}$ with dimension n

Now let us claim that the vector $f_{k+1}, \dots, f_n$ is a basis of W° . This implies that for every linear combination of vector from v_1 to v_k which equals w .

$$f_{k+1}(w) = \sum_{i=1}^k f_{k+1}(\lambda_i v_i) = 0$$

Therefore, $f_{k+1}, \dots, f_n$ are in W° . Now we have to show that they are linearly independent.

For $f \in W^\circ$

$$f(x) = \sum_{i=0}^n f(v_i) f_i(x) = f(v_{k+1}) f_{k+1}(x) + \dots f(v_n) f_n(x)$$

Therefore we get that for some f in the annihilator we get a linear combination of the our chosen basis.

□

46.0.11 The Duality Argument

$$T \text{ is surjective} \iff T^T \text{ is injective}$$

Proof:

Let us assume the opposite

$$\neg T \text{ is surjective} \iff \neg T^T \text{ is injective}$$

If T is not surjective we can assume that the image of it is a strict subset of W which we will call W' . We can also define a function $g \in W^*$ which is not 0 everywhere except for the vector in W' .

Then if we use the definition of T^T which tells us that when evaluating at g which is not the zero functional gives us

$$T^T(g)(x) = g(T(x)) = 0 \text{ because } T(x) \in W'$$

This concludes the proof in one way because we have found some some linear combination for the zero vector different from the trivial solution.

For the other way of the if and only if, we know that T^T is not injective. This means that multiple vector get transformed into the zero vector under T^T . Therefore, if suppose T is surjective and

$$\begin{aligned} T^T(g)(x) &= 0 \text{ for some } g \neq 0 \\ g(T(x)) &= 0, \quad \forall x \in V \end{aligned}$$

Then for $y = T(x)$ is always 0 but because we have assume that $g \neq 0$ we have a contradiction. T must not surjective and by getting rid of the negation on both sides we have proven the statement.

□

46.0.12 The Rank A transpose is equal to the rank of A

For this proof we will claim:

$$(T^T) = (\text{img}(T))^\circ$$

The right part tell us that for some $g \in (T^T)$, $T^T(g) = f_0$. But because we can write

$$T^T(g)(x) = g(T(x)) = 0 \text{ where } T(x) \in \text{img}(T)$$

We know that g is in the annihilator of the image of T .

With this knowledge we can know proof $\text{rank}(A) = \text{rank}(A^T)$ using the rank nullity theorem

$$\begin{aligned} \dim((T^T)) + \dim(\text{img}(T^T)) &= \dim(W^*) = \dim(W) \\ \dim((\text{img}(T)^\circ)) + \dim(\text{img}(T^T)) &= \dim(W) \end{aligned}$$

For $\dim(W)$ we get

$$\begin{aligned} \dim((\text{img}(T)^\circ)) + \dim(\text{img}(T)) &= \dim(W) \\ \dim((\text{img}(T)^\circ)) &= \dim(W) - \dim(\text{img}(T)) \end{aligned}$$

By substituting back into the original equation

$$\begin{aligned}\dim((\text{img}(T)^\circ)) + \dim(\text{img}(T^T)) &= \dim(W) \\ \dim(W) - \dim(\text{img}(T)) + \dim(\text{img}(T^T)) &= \dim(W) \\ \dim(\text{img}(T^T)) &= \dim(\text{img}(T))\end{aligned}$$

□◻◻

Chapter 47

Tensors

A *tensor* T of rank (k, l) is a multilinear map from k dual vectors and l vectors to the $\mathbb{F}$.

$$T : V^* \times \cdots \times V^* \times V \times \cdots \times V \rightarrow \mathbb{F}$$

Example:

Suppose $k = 1$, $l = 1$ and our field is $\mathbb{R}$. Then a tensor $T(\varphi, v) \rightarrow \mathbb{R}$ where $\varphi = \alpha f + \beta g$, $f, g \in V^*$ and $v = \gamma v_1 + \lambda v_2$, $v_1, v_2 \in V$ and $\alpha, \beta, \gamma, \lambda \in \mathbb{R}$. We get

$$T(\varphi, v) = T(\alpha f + \beta g, \gamma v_1 + \lambda v_2) = \alpha \gamma T(f, v_1) + \alpha \lambda T(f, v_2) + \beta \gamma T(g, v_1) + \beta \lambda T(g, v_2)$$

where each term is a real number.

Another examples: a rank $(0, 1)$ tensor is dual vector and a rank $(1, 0)$ this is a vector and finally a scalar is rank $(0, 0)$

Chapter 48

The Exponential Function and Relatives

The exponential function and its related functions such as sine, cosine, and the hyperbolic functions are fundamental in both pure and applied mathematics. This section explores these functions through their series expansions and interrelationships.

$$\exp(x) := e^x := \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n$$

48.0.1 Exponential Function as a Series

The exponential function e^x is defined as an infinite series:

$$e^x := \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots$$

This series converges for all real and complex values of x .

48.0.2 Properties of the Exponential

- $\exp(x) < \exp(y), \forall x, y \in \mathbb{R} : x < y$
- $\exp(x) \geq 1 + x, \forall 1 + x$
- $|\exp(x) - 1| \leq \frac{|x|}{1-|x|}$ for all $x \in \mathbb{R}$ such that $|x| < 1$
- For a large enough x : $\exp(x) > x^n, \forall n \in \mathbb{N}$ and $x > 4n^2$

48.0.3 General Definition

For $a > 0$ and $x \in \mathbb{R}$ we define

$$a^x := \exp(x \ln(a)) = e^{x \ln(a)}$$

48.0.4 Difference of Exponentials

For $a, b \in \mathbb{R}$ and $n \in \mathbb{N}$

$$a^n - b^n = (a - b) \sum_{k=1}^n a^{n-k} b^{k-1}$$

48.0.5 Sine and Cosine Definitions with e

Using Euler's identities and Taylor series, sine and cosine can be defined in terms of exponential functions:

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i}, \quad \cos x = \frac{e^{ix} + e^{-ix}}{2}$$

These definitions are consistent with their Taylor expansions:

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}, \quad \cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

48.0.6 Euler's Formula

One of the most elegant formulas in mathematics is Euler's formula:

$$e^{ix} = \cos x + i \sin x$$

This identity provides a deep connection between exponential and trigonometric functions and is valid for all real (and complex) x .

Special Case:

$$e^{i\pi} + 1 = 0$$

This is known as Euler's Identity and is often celebrated as one of the most beautiful equations in mathematics.

48.0.7 Reason for the Infinite Series of e (Differentiation)

The exponential series:

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

has the remarkable property that its derivative is the same function:

$$\frac{d}{dx} e^x = \sum_{n=1}^{\infty} \frac{nx^{n-1}}{n!} = \sum_{n=1}^{\infty} \frac{x^{n-1}}{(n-1)!}$$

Now replace the index $m = n - 1$:

$$= \sum_{m=0}^{\infty} \frac{x^m}{m!} = e^x$$

This shows that e^x is the unique function (up to constant multiples) that is equal to its own derivative.

48.0.8 Hyperbolic Functions with e

The hyperbolic functions are analogs of the trigonometric functions but defined using the exponential function:

$$\begin{aligned}\sinh x &= \frac{e^x - e^{-x}}{2} \\ \cosh x &= \frac{e^x + e^{-x}}{2} \\ \tanh x &= \frac{\sinh x}{\cosh x} = \frac{e^x - e^{-x}}{e^x + e^{-x}}\end{aligned}$$

These functions satisfy identities similar to those of trigonometric functions, for example:

$$\cosh^2 x - \sinh^2 x = 1$$

They naturally arise in the study of differential equations, special relativity, and the geometry of hyperbolas.

48.0.9 Inverse of the Hyperbolic Trigonometric Functions

- $\sinh^{-1}(x) = \ln(x + \sqrt{x^2 + 1})$
- $\cosh^{-1}(x) = \ln(x + \sqrt{x^2 - 1})$
- $\tanh^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$
- $\coth^{-1}(x) = \frac{1}{2 \ln \left(\frac{x+1}{x-1} \right)}$
- $\operatorname{sech}^{-1} = \ln \left(\frac{1 + \sqrt{1-x^2}}{x} \right)$
- $\operatorname{csch}^{-1} = \ln \left(\frac{1}{x} + \frac{\sqrt{1+x^2}}{|x|} \right)$

Chapter 49

Growth

Growth models are fundamental in understanding population dynamics, resource consumption, and many natural and artificial processes. Two primary models of growth are the **exponential** and the **logistic** growth models. Each of these has discrete and continuous forms.

49.0.1 Exponential Growth

Exponential growth describes a process that increases at a rate proportional to its current value.

Discrete Exponential Growth

$$P_n = P_0 \cdot (1 + r)^n$$

Where:

- P_n is the population at time step n
- P_0 is the initial population
- r is the growth rate per time step
- n is the number of time steps

Example:

If $P_0 = 100$ and $r = 0.1$, after 5 steps:

$$P_5 = 100 \cdot (1 + 0.1)^5 = 100 \cdot 1.61051 \approx 161.05$$

Continuous Exponential Growth

$$P(t) = P_0 \cdot e^{rt}$$


Where:

- $P(t)$ is the population at time t
- e is Euler's number (≈ 2.718)
- r is the continuous growth rate

Example:

If $P_0 = 100$, $r = 0.1$, and $t = 5$:

$$P(5) = 100 \cdot e^{0.5} \approx 100 \cdot 1.64872 \approx 164.87$$



49.0.2 Logistic Growth

Logistic growth occurs when a population grows rapidly at first and then slows as it approaches a maximum sustainable size (carrying capacity).

Discrete Logistic Growth

$$P_{n+1} = P_n + r \cdot P_n \cdot \left(1 - \frac{P_n}{K}\right)$$


Where:

- K is the carrying capacity
- r is the intrinsic growth rate

Example:

With $P_0 = 10$, $r = 0.5$, $K = 100$:

$$P_1 = 10 + 0.5 \cdot 10 \cdot \left(1 - \frac{10}{100}\right) = 10 + 0.5 \cdot 10 \cdot 0.9 = 10 + 4.5 = 14.5$$



Continuous Logistic Growth


$$P(t) = \frac{K}{1 + \left(\frac{K-P_0}{P_0}\right) e^{-rt}}$$

Example:

With $P_0 = 10$, $K = 100$, and $r = 0.5$, at $t = 5$:

$$P(5) = \frac{100}{1 + \left(\frac{90}{10}\right) e^{-0.5 \cdot 5}} = \frac{100}{1 + 9e^{-2.5}} \approx \frac{100}{1.7387} \approx 57.52$$

Continuous Logistic Growth



Chapter 50

Asymptotes

An *asymptote* of a curve is a line that the curve approaches but never touches as the input or output grows infinitely large in magnitude. There are three main types of asymptotes: vertical, horizontal, and oblique (slant). Each one describes a different long-term behavior of a function.

50.0.1 Vertical Asymptotes

A *vertical asymptote* is a vertical line $x = a$ where the function grows without bound as x approaches a from the left or right.

$$\lim_{x \rightarrow a^-} f(x) = \pm\infty \quad \text{or} \quad \lim_{x \rightarrow a^+} f(x) = \pm\infty$$

How to Find Vertical Asymptotes

1. Find values of x that make the denominator of a rational function zero.
2. Eliminate removable discontinuities (common factors).
3. For each remaining zero in the denominator, check if the limit tends to infinity.

Example:

$$f(x) = \frac{1}{x-2} \Rightarrow x = 2 \text{ is a vertical asymptote}$$

50.0.2 Horizontal Asymptotes

A *horizontal asymptote* is a horizontal line $y = L$ that the function approaches as x tends to infinity or negative infinity:

$$\lim_{x \rightarrow \pm\infty} f(x) = L$$

How to Find Horizontal Asymptotes (Rational Functions)

1. Compare the degrees of the numerator (n) and denominator (m):
 - If $n < m$: Horizontal asymptote at $y = 0$
 - If $n = m$: Horizontal asymptote at $y = \frac{\text{leading coefficient of numerator}}{\text{leading coefficient of denominator}}$
 - If $n > m$: No horizontal asymptote (check for slant)

Example:

$$f(x) = \frac{2x^2 + 3}{x^2 - 1} \Rightarrow y = \frac{2}{1} = 2 \text{ is the horizontal asymptote}$$

50.0.3 Oblique (Slant) Asymptotes

An *oblique or slant asymptote* occurs when the degree of the numerator is exactly one more than the degree of the denominator.

$$\lim_{x \rightarrow \infty} [f(x) - (mx + b)] = 0$$

How to Find Oblique Asymptotes

1. Perform polynomial long division (or synthetic division) on the rational function.
2. The quotient (ignoring the remainder) is the equation of the slant asymptote.

Example:

$$f(x) = \frac{x^2 + 1}{x - 1}$$

Long division gives:

$$\frac{x^2 + 1}{x - 1} = x + 1 + \frac{2}{x - 1} \Rightarrow \text{Slant asymptote: } y = x + 1$$

50.0.4 Summary Table

Type	Equation	Occurs When
Vertical	$x = a$	Denominator $\rightarrow 0$ (non-removable)
Horizontal	$y = L$	$\lim_{x \rightarrow \pm\infty} f(x) = L$
Oblique	$y = mx + b$	Degree numerator = degree denominator + 1

Chapter 51

Sequences and Series

A sequence is an ordered list of numbers, usually defined by a formula or a recurrence relation. It is one of the fundamental objects of study in mathematical analysis.

51.0.1 Arithmetic Sequence

An *arithmetic sequence* is a sequence where the difference between consecutive terms is constant.

$$a_n = a_1 + (n - 1)d$$

- a_1 is the first term
- d is the common difference $d = a_{n+1} - a_n$

Example:

$$a_n = 3 + (n - 1) \cdot 2 = 2n + 1 \quad (\text{Odd numbers})$$

51.0.2 Geometric Sequence

A *geometric sequence* is a sequence where each term is obtained by multiplying the previous term by a fixed non-zero constant.

$$a_n = a_1 \cdot r^{n-1}$$

- a_1 is the first term
- r is the common ratio $r = \frac{a_{n+1}}{a_n}$

Example:

$$a_n = 2 \cdot 3^{n-1} = 2, 6, 18, 54, \dots$$

51.0.3 Function Sequence

A *function sequence* is sequence composed of functions and its noted like:

$$(f)_n \text{ with } x^t := \{f : D \rightarrow R \mid f \text{ is a function}\}$$

51.0.4 Boundness of a Sequence

To show the boundness of a sequence we can use the definition of the *supremum* or *infimum* and approximations.

One useful definition is that a sequence is bounded below and above if

$$|a_n| \leq M, \forall n \geq 1$$

Example:

Given is $a_n = \frac{1}{2n+3}$

$$\left| \frac{1}{2n+3} \right| \leq \frac{1}{2n+3} < \frac{1}{2n} < \frac{1}{n} \leq 1$$

Because, we have found a concrete bound we can state that the sequence is bounded.

51.0.5 Convergence

A sequence (a_n) *converges* to a limit L if for every $\varepsilon > 0$, there exists an integer n_0 such that for all $n \geq n_0$:

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : \forall n \geq N |a_n - L| < \varepsilon$$

In this case, we write:

$$\lim_{n \rightarrow \infty} a_n = L$$

This also means that $\{a_n - L\}_{n=1}^{\infty}$ is a zero sequence.

Properties of convergent Sequences

Given the convergent sequences $\{a_n\}_{n=1}^{\infty}, \{b_n\}_{n=1}^{\infty}$ with a and b as the respective limits. For a bounded sequence $\{c_n\}_{n=1}^{\infty}$ then the following applies:

- $\{a_n\}_{n=1}^{\infty}$ is bounded
- $\{\alpha a_n + \beta b_n\}_{n=1}^{\infty}, \alpha, \beta \in \mathbb{R}$ converges towards $\alpha a + \beta b$
- $\{|a_n|\}_{n=1}^{\infty}$ converges towards $|a|$ and $\{\sqrt{a_n}\}_{n=1}^{\infty}$ towards $\sqrt{|a|}$
- $\{a_n \cdot b_n\}_{n=1}^{\infty}$ converges towards $a \cdot b$
- $\{a_n \cdot c_n\}_{n=1}^{\infty}$ is bounded
- If $a_n \neq 0, \forall n \in \mathbb{N}$ and $a \neq 0$ then $\lim_{n \rightarrow \infty} \frac{1}{a_n} = \frac{1}{a}$

51.0.6 The limit of a Sequence is Unique

Given a convergent sequence a_n then its limit is unique.

Proof:

Suppose that a_n has two different limits a and b . Let $\varepsilon = \frac{|a-b|}{2} > 0$

Because we have distinct limit we write

$$\forall \varepsilon > 0 \exists i \in \mathbb{N} : \forall i \geq N_1 |a_i - a| < \frac{|a-b|}{2}$$

and

$$\forall \varepsilon > 0 \exists j \in \mathbb{N} : \forall j \geq N_2 |a_j - b| < \frac{|a - b|}{2}$$

Let $M = \max N_1, N_2$. Then

$$|a - b| = |a - a_n + a_n - b| \leq |a_n - a| + |a_n - b| < 2\varepsilon = \varepsilon > 0$$

Thus, $|a - b| < |a - b|$ and this is a contradiction. The limit must be unique.

□◻◻

51.0.7 Convergence by Power of One Half

If for a sequence a_n the following holds then it converges

$$|a_n - a_{n+1}| \leq \frac{1}{2^n}$$

51.0.8 Boundness of Convergent Sequences

Let a_n be a convergent sequence then a_n is bounded

Proof:

Because a_n converges, it has a limit a . Now, let $\varepsilon = 1$, and let us use the definition of convergence.

$$\begin{aligned} -\varepsilon &< |a_n - a| < \varepsilon \\ -1 &< a_n - a < 1 \\ a - 1, a_n &< 1 + a \text{ for all } n > n_0 \end{aligned}$$

Until, this point we have proof the boundness for all $n > n_0$. To guarantee that the sequence is also bounded for $1 \leq n \leq n_0$ we define

$$U := \max(x \mid x \in a_n, n \leq n_0, a + 1)$$

and

$$L := \min(x \mid x \in a_n, n \leq n_0, a - 1)$$

Thus, our sequence is bounded

$$L \leq a_n \leq U$$

□◻◻

51.0.9 Cauchy Sequence

A sequence (a_n) is a *Cauchy sequence* if for every $\varepsilon > 0$, there exists an integer n_0 such that for all $n, m \geq n_0$:

$$\forall \varepsilon > 0 \exists n, m, N \in \mathbb{N} : |a_n - a_m| < \varepsilon \text{ with } n, m \geq N$$

Every convergent sequence is a Cauchy sequence. In complete metric spaces (like $\mathbb{R}$), the converse also holds.

Example: Convergence Exercise

Let $a_i = \frac{1}{i}$ and $\varepsilon = 0.01$. We want to find i_0 such that:

$$\left| \frac{1}{n} - 0 \right| < \varepsilon \implies \frac{1}{i} \leq \frac{1}{n} < \varepsilon \rightarrow \frac{1}{\varepsilon} < n$$

$$\Rightarrow \frac{1}{i} < 0.01 \Rightarrow i > 100 \Rightarrow i_0 = 101$$

51.0.10 Boundness of Cauchy Sequence

Every Cauchy sequence is *bounded*.

51.0.11 Proof that every Convergent Sequence is a Cauchy Sequence

Given a sequence a_n then a_n is convergent if and only if it is a Cauchy sequence.

$\rightarrow$: a_n is convergent therefore, the following holds

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : \forall n \geq N |a_n - L| < \varepsilon$$

Thus, we can use the triangle inequality for our definition of the Cauchy sequence and the following definition

$$|a_n - a| < \frac{\varepsilon}{2}$$

$$|a_m - a| < \frac{\varepsilon}{2}$$

This is allowed we know that a_n converges. We choose the max of each of the indices for which the inequality holds. Then

$$|a_n - a_m + a - a| \leq |a_n - a| + |a_m - a| < 2 \frac{\varepsilon}{2} = \varepsilon$$

$\leftarrow$: a_n is a Cauchy sequence, thus

$$\forall \varepsilon > 0 \exists n, m, N \in \mathbb{N} : |a_n - a_m| < \varepsilon \text{ with } n, m \geq N$$

a_n is bounded because else $|a_n - a_m| > \varepsilon$ for $n, m \geq N_0$. Therefore, by Bolzano-Weierstraß Theorem it must converge

□

51.0.12 Cauchy Sequences are Convergent

Given a Cauchy sequence a_n then it converges

Proof:

We know that a_n is bounded and by the Bolzano Weierstraß Theorem it has convergent subsequence.

$$\forall \varepsilon > 0 \exists n, m, N_1 \in \mathbb{N} : |a_n - a_m| < \frac{\varepsilon}{2} \text{ with } n, m \geq N_1$$

And

$$\forall \varepsilon > 0 \exists n, m, N_2 \in \mathbb{N} : |a_{n_k} - a| < \frac{\varepsilon}{2} \text{ with } n \geq N_2$$

Then let $N = \max N_1, N_2$ therefore

$$|a_n - a_m| = |a_n + a_{n_{N+1}} - a_{n_{N+1}} - a_m| \leq |a_n - a_{n_{N+1}}| + |a_m - a_{n_{N+1}}| = 2\frac{\varepsilon}{2} = \varepsilon$$

Thus, it converges.

□◻◻

51.0.13 Definition of ε -Neighborhood

The ε -neighborhood of a point $a \in \mathbb{R}$ is the set:

$$B_\varepsilon(a) = \{x \in \mathbb{R} \mid |x - a| < \varepsilon\}$$

This is the open interval $(a - \varepsilon, a + \varepsilon)$.

51.0.14 Supremum, Infimum, Maximum and Minimum

The *supremum* sup of a set $A \subseteq \mathbb{R}$ is the least upper bound: the smallest real number s such that $a \leq s$ for all $a \in A$.

$$x \leq s, \forall x \in A$$

$$x \leq s', \forall x \in A \implies s' \geq s$$

The *maximum* is the largest element in the set, if it exists.

The *minimum* is the smallest element in the set, if it exists.

Finally we define $\sup \emptyset := -\infty$ and $\inf \emptyset := +\infty$

Example I:

For $A = (0, 1)$:

- $\sup A = 1$ (not in A)
- $\inf A = 0$ (not in A)
- $\max A$ and $\min A$ do not exist

Example II:

For $(0, 5)$ and $(0, 5]$, the supremum is 5.

Example III:

For the real numbers we can define $\sqrt{2}$ as the supremum of the set $\{x \in \mathbb{R} \mid x^2 < 2\}$.

The *infimum* inf of a set A is the greatest lower bound: the largest number t such that $a \geq t$ for all $a \in A$.

$$x \geq s, \forall x \in A$$

$$x \geq s', \forall x \in A \implies s' \leq s$$

Example IV:

Prove that $\sup\{\frac{n}{n+1} \mid n \in \mathbb{N}\} = 1$.

To prove we need to show that 1 is an upper bound and then that $\forall \varepsilon > 0$ $a - \varepsilon$ is not an upper bound.

$$1 \geq \frac{n}{n+1}$$

$$n+1 \geq n$$

Thus, 1 is in fact an upper bound.

For the second part, let $\varepsilon > 0$, and we want to prove that $\exists n \in \mathbb{N} : \frac{n}{n+1} > 1 - \varepsilon$

$$\frac{n}{n+1} > 1 - \varepsilon$$

$$\varepsilon > 1 - \frac{n}{n+1}$$

$$\varepsilon > \frac{n+1-n}{n+1}$$

$$\varepsilon > \frac{1}{n+1}$$

$$n+1 > \frac{1}{\varepsilon}$$

$$n > \frac{1}{\varepsilon} - 1$$

By the Archimedian principle we now that we can always find a bigger natural number such that $n > \frac{1}{\varepsilon} - 1$. Now we can reverse the steps in the inequality, and we will get our original inequality.

□◻◻

Alternative Characterization of the Supremum

Given a non-empty set $A \subseteq \mathbb{R}$. The $s \in \mathbb{R}$ is only a supremum of A if

$$(\nexists s \in A, s < x) \wedge (s' \in \mathbb{R}, s' < s \implies \exists x \in A, x > s')$$

51.0.15 Properties of Suprema and Infima

For non-empty sets $A, B \subseteq \mathbb{R}^+$

- $\sup AB = \sup A \cdot \sup B$
- $\inf AB = \inf A \cdot \inf B$
- $\sup(A + B) = \sup A + \sup B$
- $\inf(A + B) = \inf A + \inf B$
- $\sup(A/B) = \sup(A)/\inf(B)$
- $\sup(-A) = -\inf A$
- $\inf(-A) = -\sup A$

For the sake of completeness we will prove one of these properties:

Proof:

By definition $\forall ab \in AB$ the following holds:

$$ab \leq \sup A \cdot \sup B \implies \sup AB \leq \sup A \cdot \sup B$$

Because $\sup A$ and $\sup B$ not necessarily belong to A or B . With this, we already have the half of the proof. Now we need to prove that $\sup AB \geq \sup A \cdot \sup B$.

Start with

$$\begin{aligned} ab &\leq \sup AB \\ a &\leq \frac{\sup AB}{b} \end{aligned}$$

Therefore, $\frac{\sup AB}{b}$ is an upper bound of A . This means that:

$$\sup A \leq \frac{\sup AB}{b} \implies \sup A \cdot b \leq \sup AB$$

From this we can also derive that

$$\sup A \cdot b \leq \sup AB \implies b \leq \frac{\sup AB}{\sup A}$$

Thus, because $\frac{\sup AB}{\sup A}$ is an upper bound of B we have that:

$$\sup B \leq \frac{\sup AB}{\sup A} \implies \sup A \cdot \sup B \leq \sup AB$$

Combining both parts of the proof we have that:

$$\sup AB = \sup A \cdot \sup B$$

□◻◻

The other proofs can be proven using similar arguments.

51.0.16 Limit and Accumulation Point

- A *limit* of a sequence (a_n) is the value the terms get arbitrarily close to as $n \rightarrow \infty$.
- An *accumulation point* (or limit point) of a set A is a point x such that every neighborhood of x contains infinitely many points of A .

Example:

The sequence $a_n = (-1)^n + \frac{1}{n}$ has two accumulation points: 1 and -1 .

51.0.17 The Monotone Convergence Theorem

Let a_n be a monotone sequence. Then a_n converges if and only if it is bounded.

Proof:

a_n converges; thus there exists a limit and therefore, the sequence is bounded.

The sequence a_n is bounded and monotone. Therefore, there exist both infimum and supremum. We will prove the case for the supremum

$$a \geq a_i \forall a_i \in a_n \implies a - \varepsilon < a_i$$

Suppose a is the limit then for all $i > n_0, n_0 \in \mathbb{N}$

$$\begin{aligned} |a_i - a| &< \varepsilon \\ |a - a_i| &< \varepsilon \\ |a - a_i| = a - a_i &< \varepsilon \\ a - \varepsilon &< a_i \end{aligned}$$

Therefore, the supremum is the limit.

□◻◻

51.0.18 Monotony

We define a real valued function f as *monotone increasing/decreasing*

$$x < y \implies f(x) \leq f(y)$$

$$x < y \implies f(x) \geq f(y)$$

We call it *strictly monotone increasing or decreasing* if $f(x) < f(y)$ or $f(x) > f(y)$.

51.0.19 Monotonicity by Difference and Quotient

A is considered *monotone* if either for all $n \in \mathbb{N}$ the following holds

$$a_{n+1} \geq a_n \text{ or } a_{n+1} \leq a_n$$

Monotonicity by Difference:

- If $a_{n+1} - a_n \geq 0$ for all n , then the sequence is non-decreasing.
- If $a_{n+1} - a_n < 0$, it is non-increasing.

Monotonicity by Quotient: (Typically for positive sequences)

- If $\frac{a_{n+1}}{a_n} \geq 1$, then the sequence is non-decreasing.
- If $\frac{a_{n+1}}{a_n} < 1$, it is non-increasing.

51.0.20 Divergence

A sequence *diverges* if it does not converge. That is, there is no real number L such that:

If it diverges towards some infinity then

$$\forall M \in \mathbb{R} \exists n, N \in \mathbb{N} : |a_n| \geq |M| \forall n \geq N$$

- A sequence can diverge to ∞ or $-\infty$
- A sequence can also oscillate without approaching any limit

Example:

The sequence $a_n = (-1)^n$ diverges since it oscillates between 1 and -1.

51.0.21 Subsequences and Their Properties

A *subsequence* of a sequence (a_n) is a sequence (a_{n_k}) where (n_k) is a strictly increasing sequence of natural numbers.

Properties:

- Every subsequence of a convergent sequence converges to the same limit.
- A sequence converges if and only if all of its subsequences converge to the same limit.
- A bounded sequence always has at least one convergent subsequence (Bolzano-Weierstraß theorem).
- $n_k \geq k$

Existence of the Accumulation Point as Limit

A sequence $\{x_n\}_{n=1}^{\infty}$ has x_0 as accumulation point if and only if there exists a subsequence which converges towards x_0 .

Proof:

Suppose that x_0 is an accumulation point then $(x_0 - 1, x_0 + 1)$ contains a sequence member $x_{n_0}, n_0 \in \mathbb{N}$. Let

$$n_1 := n_0$$

For each $k \in \mathbb{N}$ exists some $N(k) \in \mathbb{N}$ such that $N(k) > N(k-1)$ and

$$x_{N(k)} \in (x_0 - \frac{1}{k}, x_0 + \frac{1}{k})$$

Now we set

$$n_k = N(k)$$

For each subsequence $\{x_{n_k}\}_{k=1}^{\infty}$ the following holds

$$|x_{n_k} - x_0| \leq \frac{1}{k}$$

Therefore, the subsequence converges also towards x_0 .

Other way of the proof, if the sequence $\{x_{n_k}\}_{k=1}^{\infty}$ converges towards x_0 then there exists a $\delta > 0$ and $K \in \mathbb{N}$ such that

$$x_{n_k} \in (x_0 - \delta, x_0 + \delta), \forall k \geq K$$

Therefore, each neighborhood has infinite sequence members.

□◻◻

Monotone Subsequence

Every sequence a_n has monotone subsequence.

Proof:

Let a_n be a sequence and a_m be a peak if $\forall a_i \in a_n, i > m, m, i \in \mathbb{N}$ the following holds $a_i \leq a_m$.

Now consider the case that a_n has infinitely many peaks. Let a_{n_k} be the sequence of the k -th peak. This gives us that each peak is greater or equal than the other, therefore we have monotone growing subsequence. If we define the peaks with $a_i \geq a_m$ we also would get monotone sequence, but a decreasing instead.

For the case that there are only a finite number of peaks, let a_n be the last peak $a_i \geq a_m$. If we can always find another a_{i+1} such that $a_{i+1} \geq a_n$, therefore we can find monotone increasing subsequence. Similarly, the case for the decreasing case can be shown.

□◻◻

Convergence due to Subsequences

A sequence $\{a_n\}_{n=1}^{\infty}$ converges to a if and only if each of its subsequences also converges to a .

Proof:

Let a_{n_k} be a subsequence of a_n . Let $\varepsilon > 0$.

$$\exists n_0 \in \mathbb{N}, |a_n - a| < \varepsilon \forall n > n_0$$

We want to show

$$\exists n_1 \in \mathbb{N}, |a_{n_k} - a| < \varepsilon \forall k > n_1$$

with $n_k \geq k$.

Let $n_1 = n_0$ then if $k > n_1$, $k > n_0$ and $n_k \geq k > n_0$. Thus, for all $k > n_1$, $|a_{n_k} - a| < \varepsilon$.

For the other way of the proof: assume every subsequence of a_n converges to a . a_n is itself a subsequence of thus, it converges to a .

□◻◻

51.0.22 Proof that -1^n is divergent

Let us assume that a limit exists then

$$|-1^n - L| < \varepsilon \forall n \geq N$$

Let us choose $\varepsilon = \frac{1}{2}$.

Now our sequence bounces between -1 and 1 thus we can say that

$$|-1 - L| < \frac{1}{2}$$

$$|1 - L| < \frac{1}{2}$$

Now let us use the triangle inequality a

$$2 = |-1 - L + 1 - L| \leq |-1 + L| + |1 - L| < 2 \frac{1}{2} = 2$$

This is a contradiction. Thus, such a limit can not exist

□◻◻

51.0.23 Bolzano-Weierstraß Theorem

Every bounded sequence in $\mathbb{R}$ has at least one convergent subsequence.

Equivalently, every bounded sequence of real numbers has at least one accumulation point.

Proof:

Let a_n be a bounded sequence, and let a_{n_k} be a monotone subsequence given by the monotone subsequence theorem. a_{n_k} is bounded because original sequence is bounded.

Due to the monotone convergent theorem, we also know that the subsequence converges. Therefore, our theorem is proven.

□◻◻

51.0.24 Series

A *series* is the sum of the terms of a sequence:

$$\sum_{n=1}^{\infty} a_n$$

We define the partial sums $S_n = \sum_{k=1}^n a_k$. If the sequence (S_n) converges to a limit S , then the series is said to converge:

$$\sum_{n=1}^{\infty} a_n = S$$

51.0.25 Power Series

A *power series* is a series of the form:

$$\sum_{n=0}^{\infty} a_n (x - x_0)^n$$

Where x_0 is the center of the expansion. Power series converge within a radius R :

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}}$$

51.0.26 Geometric Series

There two types of *geometric series*.

1. *Finite geometric series*:

$$\sum_{n=0}^{r-1} aq^n = a \cdot \frac{1 - q^r}{1 - q} \quad \text{for } q \neq 1$$

2. *Infinite geometric series*:

$$\sum_{n=0}^{\infty} aq^n = \frac{a}{1 - q} \quad \text{for } |q| < 1$$

51.0.27 Point-wise Convergence

A sequence of functions (f_n) converges *point-wise* to a function f on a set A if:

$$\forall x \in A, \quad \lim_{n \rightarrow \infty} f_n(x) = f(x)$$

Point-wise convergence does not preserve properties like continuity or differentiability.

51.0.28 Uniform Convergence

A sequence of functions (f_n) converges *uniformly* to f on A if:

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n \geq n_0, \forall x \in A, \quad |f_n(x) - f(x)| < \varepsilon$$

Key Property: Uniform convergence preserves continuity, integration, and differentiation under certain conditions.

51.0.29 Cauchy Criterion for Series

A series $\sum a_n$ converges if and only if for every $\varepsilon > 0$ there exists n_0 such that:

$$\left| \sum_{k=m}^n a_k \right| < \varepsilon \quad \forall n > m \geq n_0$$

51.0.30 Interval Nesting and the Interval Nesting Theorem

Let (I_n) be a sequence of closed intervals:

$$I_n = [a_n, b_n] \quad \text{with } I_{n+1} \subseteq I_n, \text{ and } \lim_{n \rightarrow \infty} (b_n - a_n) = 0$$

Then, the intersection contains exactly one point:

$$\bigcap_{n=1}^{\infty} I_n = \{x\}$$

This is useful in proving the existence of limits, roots, and fixed points.

51.0.31 Weierstraß Approximation Theorem

The *Weierstraß Approximation Theorem* states:

Every continuous function $f : [a, b] \rightarrow \mathbb{R}$ can be uniformly approximated by a polynomial $P(x)$, i.e., for every $\varepsilon > 0$ there exists a polynomial $P(x)$ such that:

$$|f(x) - P(x)| < \varepsilon \quad \forall x \in [a, b]$$

This theorem is foundational in numerical analysis and approximation theory.

51.0.32 Zero Sequence

A sequence (a_n) is called a *null sequence* or *zero sequence* if:

$$\lim_{n \rightarrow \infty} a_n = 0$$

Every zero sequence is convergent (to 0), and plays an essential role in convergence proofs and epsilon-delta arguments.

More formally, a sequence $\{a_n\}_{n=1}^{\infty}$ of real numbers is called a zero sequence if and only if for all $\varepsilon > 0$ there exists n_0 such that

$$|a_n| < \varepsilon, \forall n \geq n_0 \in \mathbb{N}$$

Note that $\{|a_n|\}_{n=1}^{\infty}$ is only a zero sequence if $\{a_n\}_{n=1}^{\infty}$ is a zero sequence.

Example:

Is $a_n = \frac{1}{n}$ a zero sequence?

Choose $n_0 = \frac{1}{\varepsilon}$ then

$$n \geq \frac{1}{\varepsilon} \implies \frac{1}{n} \leq \varepsilon, \forall n \in \mathbb{N}, n \geq n_0$$

Thus, it is a zero sequence

51.0.33 Comparison Criteria for Sequences

Given $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ with

$$|b_n| \leq |a_n|$$

For all $n \in \mathbb{N}$ with $n \geq n_0 \in \mathbb{N}$. If $a_n \rightarrow 0, n \rightarrow \infty$ then $b_n \rightarrow 0, n \rightarrow \infty$.

We also call a_n a majorant of b_n .

51.0.34 Properties of Zero Sequences

Given the zero sequences $\{a_n\}_{n=1}^{\infty}, \{b_n\}_{n=1}^{\infty}$ and the bounded sequence $\{c\}_{n=1}^{\infty}$ then

- $\{a_n\}_{n=1}^{\infty}$ is also bounded.
- $\{\alpha a_n + \beta b_n\}_{n=1}^{\infty}, \alpha, \beta \in \mathbb{R}$ is also a zero sequence.
- $\{\sqrt{|a_n|}\}_{n=1}^{\infty}$ is a zero sequence.
- $\{a_n \cdot c_n\}_{n=1}^{\infty}$ is a zero sequence.

51.0.35 The Monotony Principle

Every bounded monotonic sequence converges.

Proof:

Define $a := \sup\{a_n : n \in \mathbb{N}\}$. We know that our sequence is bounded and in this case it has the property that $a_n \leq a$.

We use the definition of convergence

$$\begin{aligned} |a_{n_0} - a| &< \varepsilon \\ a_{n_0} &> a - \varepsilon \end{aligned}$$

Using the approximation property of the supremum we get

$$a_n \geq a_{n_0} > a - \varepsilon, \forall n \geq n_0$$

Therefore, we have for each $\varepsilon > 0$ an $n_0 \in \mathbb{N}$

$$a - \varepsilon \leq a_{n_0} \leq a \forall n \geq n_0$$

Thus, our theorem is true.

□◻◻

- If (a_n) is non-decreasing and bounded above, then $\lim a_n$ exists and equals $\sup\{a_n\}$.
- If (a_n) is non-increasing and bounded below, then $\lim a_n = \inf\{a_n\}$.

To prove the convergence of a sequence via this principle we follow the next steps

1. Prove the Monotonicity by induction
2. Calculate the limit
3. Prove the limitedness via induction
4. Write the conclusion

Example:

$$a_{n+1} = \sqrt{2a_n - 3} + 1$$

Let us assume that the series is decreasing or $a_{n+1} < a_n$

$$\begin{aligned} a_{n+2} &< a_{n+1} \\ \sqrt{2a_{n+1} - 3} + 1 &< \sqrt{2a_n - 3} + 1 \\ \sqrt{2a_{n+1} - 3} &< \sqrt{2a_n - 3} \\ 2a_{n+1} - 3 &< 2a_n - 3 \\ a_{n+1} &< a_n \end{aligned}$$

□◻◻

Now let us calculate the limit

$$\lim_{n \rightarrow \infty} a_{n+1} = \lim_{n \rightarrow \infty} \sqrt{2a_n - 3} + 1$$

Because of the recursion we know that both a_{n+1} and a_n have the same value, and we call it a .

$$\begin{aligned} a &= \sqrt{2a - 3} + 1 \\ a &= 2 \end{aligned}$$

We only have to prove the limitedness now:

For $n = 0$ we have $a_1 = 4 > 2$ then let us assume that the a is never going below 2.

$$a_{n+1} = \sqrt{2a_n - 3} + 1 \geq \sqrt{2(2) - 3} + 1 = 2$$

By setting $a_n = 2$ we get the limit again

□◻◻

51.0.36 Operations on Sequences

Sum of Sequences

The sum of two sequences (a_n) and (b_n) is a new sequence (c_n) where each term c_n is the sum of the corresponding terms of (a_n) and (b_n) :

$$c_n = a_n + b_n, \quad \forall n \in \mathbb{N}.$$

If $\lim_{n \rightarrow \infty} a_n = A$ and $\lim_{n \rightarrow \infty} b_n = B$, then the limit of the sum sequence is:

$$\lim_{n \rightarrow \infty} (a_n + b_n) = A + B.$$

Multiplication of Sequences

The product of two sequences (a_n) and (b_n) is a new sequence (d_n) where each term d_n is the product of the corresponding terms of (a_n) and (b_n) :

$$d_n = a_n \cdot b_n, \quad \forall n \in \mathbb{N}.$$

If $\lim_{n \rightarrow \infty} a_n = A$ and $\lim_{n \rightarrow \infty} b_n = B$, then the limit of the product sequence is:

$$\lim_{n \rightarrow \infty} (a_n \cdot b_n) = A \cdot B.$$

Absolute Value of a Sequence

The absolute value of a sequence (a_n) is a new sequence (e_n) where each term e_n is the absolute value of the corresponding term of (a_n) :

$$e_n = |a_n|, \quad \forall n \in \mathbb{N}.$$

If $\lim_{n \rightarrow \infty} a_n = A$, then the limit of the absolute value sequence is:

$$\lim_{n \rightarrow \infty} |a_n| = |A|.$$

Conjugate of a Complex Sequence

If (a_n) is a sequence of complex numbers, where $a_n = x_n + iy_n$ with $x_n, y_n \in \mathbb{R}$, then the conjugate of (a_n) is a new sequence $(\overline{a_n})$ where each term $\overline{a_n}$ is the complex conjugate of a_n :

$$\overline{a_n} = x_n - iy_n, \quad \forall n \in \mathbb{N}.$$

If $\lim_{n \rightarrow \infty} a_n = A = X + iY$, then the limit of the conjugate sequence is:

$$\lim_{n \rightarrow \infty} \overline{a_n} = \overline{A} = X - iY.$$

Complex Limit (Real and Imaginary Parts)

For a sequence of complex numbers (a_n) where $a_n = x_n + iy_n$, the limit of the sequence $\lim_{n \rightarrow \infty} a_n = A = X + iY$ exists if and only if the limits of the real part sequence (x_n) and the imaginary part sequence (y_n) exist individually:

$$\lim_{n \rightarrow \infty} a_n = A \iff \left(\lim_{n \rightarrow \infty} \operatorname{Re}(a_n) = \operatorname{Re}(A) = X \quad \text{and} \quad \lim_{n \rightarrow \infty} \operatorname{Im}(a_n) = \operatorname{Im}(A) = Y \right).$$

This means we can analyze the convergence of a complex sequence by examining the convergence of its real and imaginary parts separately.

Asymptotic Equality

Two sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are said to be asymptotically equal, denoted by $a_n \sim b_n$, if

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1,$$

provided that $b_n \neq 0$ for all sufficiently large n . Asymptotic equality implies that the sequences behave similarly as n approaches infinity.

51.0.37 Boundness of a Function

A real valued function f is *bounded below* if

$$\forall x \in \mathcal{D}(f) f(x) \geq m$$

and *bounded above* if

$$\forall x \in \mathcal{D}(f) f(x) \leq m$$

51.0.38 Inverse function due to Monotony

Given a strictly monotonic function f . Then f has an inverse.

Proof:

Given $x_1 < x_2$ or $x_1 > x_2$ then due to the monotony

$$f(x_1) < f(x_2) \text{ or } f(x_1) > f(x_2)$$

Thus, $f(x_1) = f(x_2) \iff x_2 = x_1$

□◻◻

51.0.39 Domain Restriction

Given $f : A \rightarrow B$ and $C \subsetneq A$ then

$$g := f|_C : C \rightarrow B, x \mapsto f(x)$$

f with *restricted* domain. The case with $A \subsetneq C$ will be the *expansion*.

Chapter 52

Limits

Limits are a foundational concept in calculus and analysis. They describe the behavior of functions and sequences as the input approaches a particular value or infinity.

52.0.1 Definition of the Limit

Let f be a function defined near a point a . We say:

$$\lim_{x \rightarrow a} f(x) = L$$

if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < |x - a| < \delta \Rightarrow |f(x) - L| < \varepsilon$$

This is the precise ε - δ *definition* of a limit.

52.0.2 Alternative Definition

Given a neighborhood U of x_0 then $\dot{U} := U \setminus \{x_0\}$ the *pointed neighborhood* of x_0 . For $\delta > 0$ we defined

$$\dot{U}_\delta(x_0) = (x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$$

Using this we can then define the limit as:

Given $f : D \rightarrow \mathbb{R}$ with $x_0 \in D$ being an accumulation point. L is the limit if for all $\varepsilon > 0$ there is a $\delta > 0$ such that

$$x \in D \cap \dot{U}_\delta(x_0) \text{ holds } |f(x) - L| < \varepsilon$$

52.0.3 Limit Calculation Rules

Sum Rule: $\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$

Difference Rule: $\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$

Product Rule: $\lim_{x \rightarrow a} (f(x)g(x)) = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$

Quotient Rule: $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$

Power Rule: $\lim_{x \rightarrow a} f(x)^n = (\lim_{x \rightarrow a} f(x))^n$

Root Rule: $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$ (when defined)

Logarithm Rule: $\lim_{x \rightarrow a} \ln(f(x)) = \ln(\lim_{x \rightarrow a} f(x))$ (when defined)

e limit Rule: $\lim_{x \rightarrow 0} (1 + f(x))^{\frac{1}{f(x)}} = e$

Proof of the Sum Rule:

For any $\varepsilon > 0$, there exists a $\delta_1 > 0$ such that if $0 < |x - a| < \delta$ then $0 < |f(x) - L| < \varepsilon$. The same goes for $g(x)$ with its respective limit and delta. Now choose $\delta = \min\{\delta_1, \delta_2\}$ such that

$$|f(x) - L| < \frac{\varepsilon}{2}$$

$$|g(x) - K| < \frac{\varepsilon}{2}$$

This using now the triangle inequality we get

$$|f(x) + g(x) - (L + K)| \leq |f(x) - L| + |g(x) - K| < 2 \frac{\varepsilon}{2} = \varepsilon$$

□◻◻

Example:

Find the limit of $\lim_{x \rightarrow 0} \frac{e^x - 1}{x}$

Apply a change of variable

$$z = e^x - 1 \iff z + 1 = e^x$$

Use the logarithm

$$z + 1 = e^x \iff \ln(z + 1) = x$$

The trend for $x \rightarrow 0$ tells us that $z \rightarrow 0$ because $\lim_{x \rightarrow 0} z = \lim_{x \rightarrow 0} e^x - 1 = 0$

Our limit is now

$$\lim_{z \rightarrow 0} \frac{z}{\ln(z + 1)} = \lim_{z \rightarrow 0} \frac{1}{\frac{1}{z} \ln(z + 1)} = \lim_{z \rightarrow 0} \frac{1}{\ln\left((z + 1)^{\frac{1}{z}}\right)}$$

The limit of a logarithm is the limit of the logarithm, and we also use the quotient rule

$$\frac{1}{\ln\left(\lim_{z \rightarrow 0} (z + 1)^{\frac{1}{z}}\right) = \ln(e)} = \frac{1}{\ln e} = 1$$

52.0.4 Limits at Infinity

$$\lim_{n \rightarrow \infty} f(n) \iff \forall \varepsilon > 0, \exists N > 0 : \forall n > N : |a_n - L| < \varepsilon$$

with $a_n = f(n)$ as a sequence.

Some common limits at infinity are:

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0, \quad \lim_{n \rightarrow \infty} e^n = \infty, \quad \lim_{n \rightarrow \infty} \frac{1}{n^2} = 0$$

Example:

Prove that $\lim_{n \rightarrow \infty} \frac{2n+1}{3n+7} = \frac{2}{3}$

We start with the definition. Let $\varepsilon > 0$ be given. We need to find $N > 0$ such that for all $n > N$:

$$\begin{aligned} \left| \frac{2n+1}{3n+7} - \frac{2}{3} \right| &< \varepsilon \\ \left| \frac{(6n+3) - (6n+14)}{9n+21} \right| &< \varepsilon \\ \left| \frac{-11}{9n+21} \right| &\leq \frac{|-11|}{|9n|+|21|} = \frac{|-11|}{9n+21} < \frac{11}{9n} < \varepsilon \end{aligned}$$

Thus, we get

$$n > \frac{11}{9\varepsilon}$$

By the Archimedean property, there exists an integer N such that

$$N > \frac{11}{9\varepsilon}$$

Then for all $n > N$ we have

$$\left| \frac{2n+1}{3n+7} - \frac{2}{3} \right| < \varepsilon \text{ for all } n > N > \frac{11}{9\varepsilon}$$

Thus, $\varepsilon > \frac{11}{9n}$ which satisfies the definition of the limit.

□◻◻

52.0.5 Indeterminate Forms

Indeterminate forms arise in limits when the expression does not directly imply a limit value:

$$\frac{0}{0}, \quad \frac{\infty}{\infty}, \quad 0 \cdot \infty, \quad \infty - \infty, \quad 1^\infty, \quad \infty^0, \quad 0^0$$

52.0.6 Limit of the n-th Root of n

We will show that $\lim_{n \rightarrow \infty} \sqrt[n]{n} = 1$

Proof:

Let $n = (1 + (\sqrt[n]{n} - 1))^n$. This is valid substitution. Then

Now let use the binomial theorem $(1 + (\sqrt[n]{n} - 1))^n = (1 + b_n)^n$

$$\begin{aligned} (1 + b_n)^n &= \sum_{k=0}^n \binom{n}{k} b_n^k \\ &= 1 + b_n + \frac{n(n-1)}{2} b_n^2 + \dots \\ &> \frac{n(n-1)}{2} b_n^2 \end{aligned}$$

Then

$$\begin{aligned} (1 + b_n)^n &= n > \frac{n(n-1)}{2} b_n^2 \\ \sqrt{\frac{2}{n-1}} &> b_n \end{aligned}$$

Now if we take the limit

$$\lim_{n \rightarrow \infty} \sqrt{\frac{2}{n-1}} = 0 > \lim_{n \rightarrow \infty} b_n$$

Then

$$\lim_{n \rightarrow \infty} 1 + b_n = 1 + 0 = 1$$

ΩΕΘ

52.0.7 Limit of a Composite Function

If $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow L} g(x) = g(L)$ (i.e., g is continuous at L), then:

$$\lim_{x \rightarrow a} g(f(x)) = g\left(\lim_{x \rightarrow a} f(x)\right)$$

52.0.8 Right and Left Limits

For $(a - \delta, a)$ and $(a, a + \delta)$

$$\lim_{x \rightarrow a^-} f(x) = L_-, \quad \lim_{x \rightarrow a^+} f(x) = L_+$$

The two-sided limit $\lim_{x \rightarrow a} f(x)$ exists if and only if $L_- = L_+$.

52.0.9 L'Hôpital's Rule

If $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or $\pm\infty$, and $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ exists, then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

This rule is used to resolve indeterminate forms $\frac{0}{0}$ and $\frac{\infty}{\infty}$.

52.0.10 Limit of a fraction

For expressions like:

$$\lim_{n \rightarrow \infty} \left(\frac{a}{b}\right)^n$$

- If $\frac{a}{b} < 1$: limit is 0
- If $\frac{a}{b} > 1$: limit diverges to ∞

52.0.11 Limit due to Sequence

Given an accumulation point x_0 in the domain of f then

$$\lim_{x \rightarrow x_0} f(x) = L$$

If and only if for the sequence in the domain of f the following is true

$$\lim_{n \rightarrow \infty} x_n = x_0$$

Proof:

$\Rightarrow$ Assume $\lim_{x \rightarrow x_0} f(x) = L$. Let a_n be a sequence in the domain with $a_n \rightarrow x_0$ and $a_n \neq x_0$ we must show that $f(a_n) \rightarrow L$

Let $\varepsilon > 0$. There exists $\delta > 0$ such that if $0 < |x - x_0| < \delta$ then $|f(x) - L| < \varepsilon$. But since $a_n \rightarrow x_0$ then $\exists N : |a_n - x_0| < \delta, \forall n > N$. Thus, for all $n > N$, $|f(a_n) - L| < \varepsilon$ and so $f(a_n) \rightarrow L$.

$\Leftarrow$ Assume that for every sequence a_n in the domain which converges to x_0 $f(a_n) \rightarrow L$

Suppose for the sake of contradiction that $\lim_{x \rightarrow x_0} f(x) \neq L$

Then, there exists $\varepsilon > 0$ such that for all $\delta > 0$, $\exists x \in \mathcal{D}, 0 < |x - x_0| < \delta$ where $|f(x) - L| \geq \varepsilon$. Therefore, by letting $\delta_n = \frac{1}{n}$, $\exists a_n \in \mathcal{D}$ satisfying $0 < |a_n - x_0| < \frac{1}{n}$ and which $|f(a_n) - L| \geq \varepsilon$.

Then by contradiction $f(a_n)$ does not converge to L .

Hence, it must be that $\lim_{x \rightarrow x_0} f(x) = L$

□

52.0.12 Limit of a Recursive Sequence

Let (a_n) be defined recursively:

$$a_{n+1} = f(a_n)$$

If (a_n) converges to L and f is continuous, then:

$$\lim_{n \rightarrow \infty} a_n = L \Rightarrow L = f(L)$$

This is used to find fixed points of recursive definitions.

52.0.13 Important Limits

- $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$
- $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \frac{1}{2}$
- $\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$
- $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$
- $\lim_{n \rightarrow \infty} \frac{n!}{n^n} = 0$
- $\lim_{n \rightarrow \infty} \sqrt[n]{a} = 1$ (for $a > 0$)
- $\lim_{x \rightarrow 0} \frac{1}{\cos x} = 1$

52.0.14 Squeeze Theorem (Sandwich Theorem)

If $g(x) \leq f(x) \leq h(x)$ near a , and:

$$\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L \Rightarrow \lim_{x \rightarrow a} f(x) = L$$

Proof:

We know that $g(x) \leq f(x) \leq h(x)$ therefore, for $g(x)$ and $h(x)$ we can define the following inequalities.

$$\begin{aligned} |g(x)_{N_1} - L| &< \varepsilon \\ |h(x)_{N_2} - L| &< \varepsilon \end{aligned}$$

Then let us define $n = \max(N_1, N_2)$. And we can also write

$$\begin{aligned} -\varepsilon &< g(x)_n - L < \varepsilon \\ -\varepsilon &< h(x)_n - L < \varepsilon \end{aligned}$$

Now, we write the following because we are assuming they all have the same limit

$$\begin{aligned} -\varepsilon &< g(x)_n - L \leq f(x)_n - L \leq h(x)_n - L < \varepsilon \\ -\varepsilon &< f(x)_n - L < \varepsilon \end{aligned}$$

from our original inequality. Thus, there has been no contradiction.

□♦♦

Example:

$$\lim_{x \rightarrow 0} x^2 \cos\left(\frac{1}{x}\right) = 0$$

52.0.15 The Number e as a Limit

The number e is defined as:

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

More generally:

$$\lim_{n \rightarrow \infty} \left(1 + \frac{a}{bn^d}\right)^{n^d} = e^{\frac{a}{b}}$$

Example I:

$$\lim_{x \rightarrow \infty} \left(\frac{x^2 + 1}{x^2 + 2}\right)^{x^2}$$

$$\lim_{x \rightarrow \infty} \left(\left(\frac{x^2 + 1}{x^2 + 2}\right)^x\right)^x$$

$$\lim_{x \rightarrow \infty} \left(\left(\frac{1 + \frac{1}{x^2}}{1 + \frac{2}{x^2}}\right)^x\right)^x$$

$$\lim_{x \rightarrow \infty} \left(\frac{e}{e^2}\right)^x = \left(\frac{1}{e}\right)^x = 0$$

Example II:

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^{x^2}$$

$$\lim_{x \rightarrow \infty} \left(\left(1 + \frac{1}{x}\right)^{x^2}\right)^{\frac{x}{x^2}}$$

$$\lim_{x \rightarrow \infty} \left(\left(1 + \frac{1}{x} \right)^x \right)^x$$

$$\lim_{x \rightarrow \infty} \left(\lim_{n \rightarrow \infty} \left(1 + \frac{1}{x} \right)^x \right)^x$$

$$\lim_{x \rightarrow \infty} (e)^x = \infty$$

52.0.16 Limits of arctan

- $\lim_{x \rightarrow \infty} \arctan x = \frac{\pi}{2}$
- $\lim_{x \rightarrow -\infty} \arctan x = -\frac{\pi}{2}$
- $\arctan x$ is continuous and differentiable everywhere, with a horizontal asymptote at $y = \pm \frac{\pi}{2}$

52.0.17 Bolzano-Weierstraß Theorem

Every bounded, convergent sequence in $\mathbb{C}$ has a convergent subsequence. More precisely the sequence a has biggest and smallest accumulation point h^* and h_*

$$\forall \varepsilon > 0 : h_* - \varepsilon < a_n < h^* + \varepsilon \text{ for all-most all } n \in \mathbb{N}$$

$$h^* := \lim_{n \rightarrow \infty} \sup a_n$$

$$h_* := \lim_{n \rightarrow \infty} \inf a_n$$

52.0.18 Uniqueness of the Limit

Let $f : D \subset \mathbb{R} \rightarrow \mathbb{R}$, and suppose a is a limit point of D . Then: If $\lim_{x \rightarrow a} f(x) = L_1$ and $\lim_{x \rightarrow a} f(x) = L_2$, then $L_1 = L_2$.

In other words, a function can have at most one limit at a given point. This is the *uniqueness of limits*.

Proof:

Assume $L_1 \neq L_2$, and let $\varepsilon = \frac{|L_1 - L_2|}{2} > 0$.

- Since $\lim_{x \rightarrow a} f(x) = L_1$, there exists $\delta_1 > 0$ such that for all $x \in D$, $0 < |x - a| < \delta_1$ implies:

$$|f(x) - L_1| < \varepsilon$$

- Since $\lim_{x \rightarrow a} f(x) = L_2$, there exists $\delta_2 > 0$ such that for all $x \in D$, $0 < |x - a| < \delta_2$ implies:

$$|f(x) - L_2| < \varepsilon$$

Let $\delta = \min(\delta_1, \delta_2)$, and choose $x \in D$ such that $0 < |x - a| < \delta$.

Then:

$$|f(x) - L_1| < \varepsilon, \quad |f(x) - L_2| < \varepsilon$$

Using the triangle inequality:

$$|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2| \leq |f(x) - L_1| + |f(x) - L_2| < \varepsilon + \varepsilon = 2\varepsilon$$

Thus,

$$|L_1 - L_2| < |L_1 - L_2|$$

This contradiction tells us that, our assumption was false. Hence, $L_1 = L_2$. The limit is unique.

□◻◻

52.0.19 Every Convergent Sequence is a Cauchy Sequence

Let (a_n) be a sequence in $\mathbb{R}$. We say:

If (a_n) converges, then it is a Cauchy sequence.

A sequence (a_n) is a *Cauchy sequence* if:

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ such that } \forall n, m \geq N, \quad |a_n - a_m| < \varepsilon$$

Proof:

Assume $\lim_{n \rightarrow \infty} a_n = L$. Then:

- For every $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that:

$$\forall n \geq N, \quad |a_n - L| < \frac{\varepsilon}{2}$$

- Now, for any $n, m \geq N$, use the triangle inequality:

$$|a_n - a_m| = |a_n - L + L - a_m| \leq |a_n - L| + |a_m - L| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

Conclusion:

$$|a_n - a_m| < \varepsilon \text{ for all } n, m \geq N \Rightarrow (a_n) \text{ is Cauchy}$$

This completes the proof that every convergent sequence is Cauchy.

□◻◻

52.0.20 Limsup and Liminf

Given a sequence $\{a_k\}_{k=1}^{\infty}$ and the infinite sequences x_n and y_n defined as

$$x_n := \sup \{a_k \mid k \geq n\} \quad y_n := \inf \{a_k \mid k \geq n\}$$

We define *limes superior* and *limes inferior* as

$$a^* = \lim_{n \rightarrow \infty} \sup a_n := \lim_{n \rightarrow \infty} x_n$$

$$a_* = \lim_{n \rightarrow \infty} \inf a_n := \lim_{n \rightarrow \infty} y_n$$

52.0.21 Limit by Bounds

If the limit superior and inferior are equal then the limit exists.

Chapter 53

Convergence Criteria

In this section, we present various tools used to determine whether an infinite series or sequence converges. These criteria are essential for analyzing the behavior of sequences and series in analysis.

53.0.1 Quotient (Ratio) Test

Let $\sum a_n$ be a series with positive terms, and suppose the limit

$$L = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right|$$

exists.

- If $L < 1$, the series converges absolutely.
- If $L > 1$ or $L = \infty$, the series diverges.
- If $L = 1$, the test is inconclusive.

Proof of the Ratio Test (for $L < 1$)

Let $r < 1$ such that $L < r < 1$. Then for sufficiently large n , we have:

$$\left| \frac{a_{n+1}}{a_n} \right| \leq r \Rightarrow a_{n+1} \leq r a_n$$

By induction:

$$a_{n+k} \leq a_n r^k$$

So,

$$\sum_{n=N}^{\infty} a_n \leq a_N \sum_{k=0}^{\infty} r^k = \frac{a_N}{1-r}$$

Hence, the series converges by comparison with a geometric series.

53.0.2 Zero Sequence Criterion

If a series $\sum a_n$ converges, then:

$$\lim_{n \rightarrow \infty} a_n = 0$$

The converse is not true — the harmonic series is a classic counterexample.

53.0.3 Majorant (Comparison Test)

If $0 \leq a_n \leq b_n$ for all n and $\sum b_n$ converges, then $\sum a_n$ converges.

53.0.4 Minorant

If $0 \leq b_n \leq a_n$ and $\sum b_n$ diverges, then $\sum a_n$ diverges.

53.0.5 Root Test

Let $\sum a_n$ be a series with non-negative terms, and suppose:

$$L = \limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}$$

- If $L < 1$, the series converges absolutely.
- If $L > 1$, the series diverges.
- If $L = 1$, the test is inconclusive.

Proof of the Root Test (for $L < 1$)

Let r be such that $L < r < 1$. Then for large n :

$$\sqrt[n]{|a_n|} \leq r \Rightarrow |a_n| \leq r^n$$

Since $\sum r^n$ converges, by comparison $\sum |a_n|$ converges absolutely.

53.0.6 Leibniz Criterion (Alternating Series Test)

Let $\sum (-1)^n a_n$ be a series with:

- a_n decreasing: $a_{n+1} \leq a_n$
- $\lim a_n = 0$

Then the series converges.

Example:

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$$

53.0.7 Compactness Criterion

A sequence (x_n) in $\mathbb{R}^n$ has a convergent subsequence if and only if it is bounded. This is a consequence of the **Bolzano-Weierstraß theorem**, which is a form of sequential compactness in $\mathbb{R}^n$.

53.0.8 Cauchy Criterion I

A sequence (a_n) converges if and only if it is a *Cauchy sequence*:

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N}, \forall n, m \geq n_0 \Rightarrow |a_n - a_m| < \varepsilon$$

This is an internal (non-limit based) criterion for convergence, valid in complete metric spaces.

53.0.9 Cauchy Criterion II

A series $a_n = \sum_{i=0}^k x_i$ converges $\iff \forall d \in \mathbb{R}, d > 0 \exists N \in \mathbb{N} \forall n, m > N : |\sum_{i=0}^n x_i + \sum_{i=0}^m x_i|$

$$= \left| \sum_{i=0}^n x_i \right| < d$$

53.0.10 The Cauchy Product

Let $\sum a_n$, $\sum b_n$ be convergent series. The *Cauchy product* is:

$$\sum_{n=0}^{\infty} c_n, \quad \text{where } c_n = \sum_{k=0}^n a_k b_{n-k}$$

If $\sum a_n$ and $\sum b_n$ converge absolutely, then their Cauchy product converges absolutely and:

$$\sum c_n = \left(\sum a_n \right) \left(\sum b_n \right)$$

Cauchy Product Example

Prove that $e^x e^y = e^{x+y}$

Note that $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ and $e^y = \sum_{n=0}^{\infty} \frac{y^n}{n!}$

Therefore,

$$e^x e^y = \sum_{n=0}^{\infty} \frac{x^n}{n!} \sum_{n=0}^{\infty} \frac{y^n}{n!}$$

And because of the absolute convergence

$$e^x e^y = \sum_{n=0}^{\infty} \sum_{k=0}^n \frac{x^k}{k!} \frac{y^{n-k}}{(n-k)!}$$

Finally,

$$e^x e^y = \sum_{n=0}^{\infty} \sum_{k=0}^n x^k y^{n-k} \binom{n}{k} \frac{1}{n!} = \sum_{n=0}^{\infty} \frac{(x+y)^n}{n!}$$

QED

53.0.11 Geometric Series

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r} \quad \text{for } |r| < 1$$

If $|r| \geq 1$, the series diverges.

53.0.12 Harmonic Series

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

This series diverges, even though its terms approach zero.

Proof:

Group terms:

$$\left(\frac{1}{2} \right) + \left(\frac{1}{3} + \frac{1}{4} \right) + \left(\frac{1}{5} + \cdots + \frac{1}{8} \right) + \cdots > \sum_{k=1}^{\infty} \frac{1}{2}$$

The lower bound is a divergent series, hence the harmonic series diverges.

Chapter 54

Convergence Radius

Consider a power series of the form

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n,$$

where c_n are the coefficients, x is the variable, and a is the center of the series. The radius of convergence, denoted by R , is a non-negative real number (or ∞) such that the series converges if $|x-a| < R$ and diverges if $|x-a| > R$. In other words, the interval of convergence is $(a-R, a+R)$, possibly including one or both endpoints.

$$R = \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{|a_n|}}$$

or

$$R = \limsup_{n \rightarrow \infty} \frac{a_n}{a_{n+1}}$$

54.0.1 Test of the Borders

When the absolute value of the difference between x and a is equal to the radius of convergence (i.e., $|x-a| = R$), the convergence of the power series must be checked separately. This involves substituting the values $x = a \pm R$ into the series and determining whether the resulting series converges or diverges using other convergence tests (e.g., the comparison test, ratio test, alternating series test).

54.0.2 Approximating the Error

When a power series is used to approximate a function, it is often necessary to estimate the error in the approximation. If the power series converges, the error can be made arbitrarily small by including a sufficient number of terms. For an alternating series that satisfies the conditions of the Alternating Series Test, the error in approximating the sum by the first n terms is bounded by the absolute value of the $(n+1)$ -th term:

$$\left| \sum_{k=0}^{\infty} c_k(x-a)^k - \sum_{k=0}^n c_k(x-a)^k \right| \leq |c_{n+1}(x-a)^{n+1}|.$$

For other power series, techniques such as Taylor's Theorem with remainder can be used to estimate the error.

54.0.3 Complete Example

Find the convergence radius of

$$\sum_{n=1}^{\infty} \frac{x^n}{n2^n}$$

at $x = 0$.

Step 1: Extract the sequence a_n

The general form is

$$\sum_0^{\infty} a_n (x - x_0)^n$$

in our case

$$= \sum_{n=1}^{\infty} x^n \frac{1}{n2^n}$$

with $\frac{1}{n2^n}$ as a_n .

Step 2: Compute the radius

$$\begin{aligned} R &= \frac{1}{\limsup_{n \rightarrow \infty} \sqrt[n]{\frac{1}{n2^n}}} \\ &= \frac{1}{\limsup_{n \rightarrow \infty} \frac{\sqrt[n]{1}}{\sqrt[n]{n} \sqrt[n]{2^n}}} \\ &= \frac{1}{\frac{1}{2}} = 2 \end{aligned}$$

Step 3: Check Borders

The borders to check are $0 - 2$ to $0 + 2$.

For -2:

$$\sum_{n=1}^{\infty} \frac{(-2)^n}{n2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n (2)^n}{n2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$$

by Leibniz, it converges.

For 2:

$$\sum_{n=1}^{\infty} \frac{(2)^n}{n2^n} = \sum_{n=1}^{\infty} \frac{1}{n}$$

This is the harmonic series therefore it diverges.

Step 4: Write the interval

The final interval of convergence is $[-2, 2)$.

Chapter 55

Calculating Series Values

55.0.1 Telescoping Sums and Partial Fraction Decomposition

Telescoping sums are series where most of the terms cancel out, leaving only a few terms. Partial fraction decomposition is a technique used to rewrite rational functions as a sum of simpler fractions, which can often lead to telescoping sums.

Example: $\sum_{k=1}^{\infty} \frac{1}{4k^2-1}$

Consider the infinite series

$$S = \sum_{k=1}^{\infty} \frac{1}{4k^2-1}.$$

We can decompose the fraction using partial fraction decomposition:

$$\frac{1}{4k^2-1} = \frac{1}{(2k-1)(2k+1)} = \frac{A}{2k-1} + \frac{B}{2k+1}.$$

Multiplying both sides by $(2k-1)(2k+1)$ gives

$$1 = A(2k+1) + B(2k-1).$$

To solve for A and B , we can use the following values of k :

- For $k = \frac{1}{2}$: $1 = A(1+1) + B(0) \implies A = \frac{1}{2}$.
- For $k = -\frac{1}{2}$: $1 = A(0) + B(-1-1) \implies B = -\frac{1}{2}$.

Thus, we have

$$\frac{1}{4k^2-1} = \frac{1}{2} \left(\frac{1}{2k-1} - \frac{1}{2k+1} \right).$$

Now we can write the series as a telescoping sum:

$$S = \frac{1}{2} \sum_{k=1}^{\infty} \left(\frac{1}{2k-1} - \frac{1}{2k+1} \right).$$

Let S_n be the n -th partial sum:

$$\begin{aligned}
S_n &= \frac{1}{2} \left[\left(\frac{1}{1} - \frac{1}{3} \right) + \left(\frac{1}{3} - \frac{1}{5} \right) + \left(\frac{1}{5} - \frac{1}{7} \right) + \cdots + \left(\frac{1}{2n-1} - \frac{1}{2n+1} \right) \right] \\
&= \frac{1}{2} \left[1 - \frac{1}{2n+1} \right].
\end{aligned}$$

Taking the limit as n approaches infinity, we get

$$S = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{1}{2} \left(1 - \frac{1}{2n+1} \right) = \frac{1}{2}(1 - 0) = \frac{1}{2}.$$

Therefore, the value of the series is $\frac{1}{2}$.

Example II:

Now an example using the geometric series

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{4}{3^n} &= \sum_{n=0}^{\infty} 4 \left(\frac{1}{3} \right)^n \\
&= \frac{4}{1 - \frac{1}{3}}
\end{aligned}$$

Chapter 56

Transforming Functions into Power Series

Many functions can be represented as power series, which is useful for various applications in calculus and analysis. This section discusses some common techniques for transforming functions into power series.

56.0.1 Common Cases

Here are some common cases where functions can be easily transformed into power series:

$$\frac{1}{1-x}$$

The function $\frac{1}{1-x}$ has the well-known geometric series representation:

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n, \quad |x| < 1.$$

$$\frac{1}{1-(x-x_0)^n}$$

A slight generalization is:

$$\frac{1}{1-(x-x_0)^n} = \sum_{k=0}^{\infty} (x-x_0)^{nk}, \quad |x-x_0| < 1.$$

$$\frac{1}{1-d(x-x_0)^n}$$

For a constant d :

$$\frac{1}{1-d(x-x_0)^n} = \sum_{k=0}^{\infty} d^k (x-x_0)^{nk}, \quad |d(x-x_0)^n| < 1.$$

Example: $\frac{x+1}{3-x}$

Let's find the power series representation of the function $f(x) = \frac{x+1}{3-x}$.

Partial Fractions

First, we perform partial fraction decomposition. We can rewrite the function as:

$$\frac{x+1}{3-x} = \frac{-(3-x)+4}{3-x} = -1 + \frac{4}{3-x}.$$

Manipulating the Expression

Now, we manipulate the fraction to fit the form $\frac{1}{1-u}$:

$$\frac{4}{3-x} = \frac{4}{3(1-\frac{x}{3})} = \frac{4}{3} \cdot \frac{1}{1-\frac{x}{3}}.$$

Power Series Representation

Using the geometric series formula, we have

$$\frac{1}{1-\frac{x}{3}} = \sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n, \quad \left|\frac{x}{3}\right| < 1.$$

Thus,

$$\begin{aligned} \frac{x+1}{3-x} &= -1 + \frac{4}{3} \sum_{n=0}^{\infty} \left(\frac{x}{3}\right)^n \\ &= -1 + \frac{4}{3} \sum_{n=0}^{\infty} \frac{x^n}{3^n} \\ &= -1 + \sum_{n=0}^{\infty} \frac{4}{3^{n+1}} x^n, \quad |x| < 3. \end{aligned}$$

Therefore, the power series representation of the function is

$$\frac{x+1}{3-x} = -1 + \sum_{n=0}^{\infty} \frac{4}{3^{n+1}} x^n, \quad |x| < 3.$$

Chapter 57

Continuity

A real-valued function f is continuous if it is continuous for all points in its domain.

57.0.1 Definition of Continuity at a Point

A function f is *continuous* at a point x_0 if:

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon$$

Equivalently,

$$\lim_{x \rightarrow x_0} f(x) = f(x_0)$$

To solve continuity problems the general approach is to:

1. Write down the definition.
2. Substitute f for the concrete function into the definition.
3. Modify the expression with the concrete f $|f(x) - f(y)|$ in way that you get $|x - y|$ inside the expression.
4. Choose a δ if not given. On the one side it is often $\delta = |x - y|$ and then substitute into the expression. On the other side if δ is given make an approximation which is greater than the current expression and then simplify. In both cases after simplifying you should get a ε is the function is continuous.

Example:

$\varepsilon - \delta$ Exercise for $f(x) = \frac{1}{x}$ at $x_0 > 0$

We want to find δ such that:

$$|x - x_0| < \delta \Rightarrow \left| \frac{1}{x} - \frac{1}{x_0} \right| < \varepsilon$$

Start by rewriting:

$$\left| \frac{1}{x} - \frac{1}{x_0} \right| = \left| \frac{x_0 - x}{xx_0} \right| = \frac{|x - x_0|}{|x||x_0|}$$

Assume:

$$x \in \left[\frac{x_0}{2}, \frac{3x_0}{2} \right] \Rightarrow |x| \geq \frac{x_0}{2}$$

Then:

$$\left| \frac{1}{x} - \frac{1}{x_0} \right| \leq \frac{|x - x_0|}{\frac{x_0}{2} \cdot x_0} = \frac{2}{x_0^2} |x - x_0|$$

Choose:

$$\delta = \min \left\{ \frac{x_0}{2}, \frac{x_0^2 \varepsilon}{2} \right\}$$

Example:

Prove that

$$f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$$

is continuous at $x_0 = 0$.

Recall, f is continuous at $x_0 = 0$ if:

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon$$

We have $\varepsilon > 0$ and we want to find $\delta > 0$:

Let us assume $|x - 0| < \delta \iff \delta > |x|$ then

$$\begin{aligned} |f(x) - f(0)| &= |f(x) - 0| = \left| x^2 \sin\left(\frac{1}{x}\right) \right| \\ &\leq |x^2| \cdot \left| \sin\left(\frac{1}{x}\right) \right| \\ &\leq |x^2| \cdot 1 = x^2 < \varepsilon \\ &\Rightarrow x < \sqrt{\varepsilon} \end{aligned}$$

Choose $\delta = \sqrt{\varepsilon}$.

Let $\varepsilon > 0$ be given and choose $\delta = \sqrt{\varepsilon}$. Then for all x with $|x - 0| < \delta$ we have:

$$|f(x) - f(0)| < \varepsilon$$

□

57.0.2 Characterization using sequences

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and $x_0 \in (a, b)$.

The function f is continuous at x_0 if and only if for every sequence

$$\{x_n\}_{n \in \mathbb{N}}$$

with $x_n \in (a, b)$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} x_n = x_0$, it holds that

$$\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$$

57.0.3 Left and Right Continuity

Let $f : D \rightarrow \mathbb{R}$ be a function and $x_0 \in D$. The function f is called *left-continuous* at x_0 if f is continuous on $D \cap (-\infty, x_0]$. Analogously, f is *right-continuous* at x_0 if f is continuous on $D \cap [x_0, \infty)$.

57.0.4 Rules of Continuity

Let f and g be continuous at x_0 :

- $f + g$, $f - g$, $f \cdot g$ are continuous at x_0
- $\frac{f}{g}$ is continuous at x_0 if $g(x_0) \neq 0$
- Compositions: if g is continuous at x_0 , and f is continuous at $g(x_0)$, then $f \circ g$ is continuous at x_0

57.0.5 Continuity and Compactness

A function over an interval $[a, b]$ which is continuous is *bounded*.

Proof:

If f is not bounded, then $\forall n \in \mathbb{N}, \exists x_n \in [a, b]$ such that

$$|f(x_n)| > n$$

By the Bolzano-Weierstrass theorem, there exists a convergent subsequence $\{x_{n_k}\} \subset [a, b]$ such that

$$\lim_{k \rightarrow \infty} x_{n_k} = x^* \in [a, b]$$

By continuity of f :

$$\lim_{k \rightarrow \infty} f(x_{n_k}) = f(x^*) = \infty$$

But by construction $|f(x_{n_k})| > n_k \rightarrow \infty$ as $k \rightarrow \infty$, which is a contradiction to the $x^* \in \mathcal{D}(f)$

□□□

57.0.6 Extreme-Value Theorem

If $f : [a, b] \rightarrow \mathbb{R}$ is continuous on the closed interval $[a, b]$, then exist $x_m, x_M \in [a, b]$ such that

$$f(x_m) = \inf_{[a,b]} f, \quad f(x_M) = \sup_{[a,b]} f$$

Proof:

Let $m := \inf_{[a,b]} f$, $M := \sup_{[a,b]} f$ we know by the previous theorem that f is bounded, so $m, M \in \mathbb{R}$.

By the Bolzano-Weierstrass theorem, there exist convergent subsequences $\{x_{n_k}\}, \{x_{m_k}\} \subset [a, b]$ such that

$$\lim_{k \rightarrow \infty} f(x_{n_k}) = m, \quad \lim_{k \rightarrow \infty} f(x_{m_k}) = M$$

Since $[a, b]$ is closed, the limits of these subsequences $x_m, x_M \in [a, b]$. Then by continuity of f :

$$f(x_m) = \lim_{k \rightarrow \infty} f(x_{n_k}) = m, \quad f(x_M) = \lim_{k \rightarrow \infty} f(x_{m_k}) = M$$

□□□

57.0.7 Extreme Points

If for set $A \subseteq \mathbb{R}$ with $f : A \rightarrow \mathbb{R}$ holds that

$$f(x_m) = \inf_{x \in A} f(x) =: m \quad \text{or} \quad f(x_M) = \sup_{x \in A} f(x) =: M$$

then x_m is called a *point of minimum* and x_M a *point of maximum* of f on A . Also, $f(x_m)$ is called the *minimum* and $f(x_M)$ the *maximum* of f on A . These are the *extreme values*.

57.0.8 Middle Value Theorem

If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and $f(a) \neq f(b)$, then for every y_0 between $f(a)$ and $f(b)$ there exists $x_0 \in (a, b)$ such that $f(x_0) = y_0$.

57.0.9 Continuity of the Inverse by a decreasing Function

Given a monotonic decreasing continuous function f on $[a, b]$ with $f(a) = c$ and $f(b) = d$, then the inverse function $f^{-1} : [d, c] \rightarrow [a, b]$ is also continuous and monotonic.

Proof:

Continuity of the Root and Logarithm Function

$x \mapsto \sqrt[n]{x}$ is continuous on $[0, \infty)$

$x \mapsto \log_n x$ is continuous on $(0, \infty)$

57.0.10 Lipschitz Continuity

A function $f : D \subset \mathbb{R} \rightarrow \mathbb{R}$ is *Lipschitz continuous* if:

$$\exists L > 0 \text{ such that } |f(x) - f(y)| \leq L|x - y| \quad \forall x, y \in D$$

- Every Lipschitz continuous function is uniformly continuous.
- The smallest such L is called the *Lipschitz constant*.

57.0.11 Fixed Point

A *fixed point* of a function f is a point x^* such that:

$$f(x^*) = x^*$$

57.0.12 Fixed Point on Interval

Given a continuous function $f : [a, b] \rightarrow \mathbb{R}$ with $f([a, b]) \subseteq [a, b]$, then there exists at least one fixed point $x^* \in [a, b]$ such that $f(x^*) = x^*$.

57.0.13 Banach Fixed Point Theorem

Let (X, d) be a complete metric space, and $f : X \rightarrow X$ a *contraction mapping*, i.e.,

$$\exists L < 1 \text{ such that } d(f(x), f(y)) \leq L \cdot d(x, y)$$

Then:

- f has a unique fixed point $x^* \in X$, such that $f(x^*) = x^*$
- Iterating $x_{n+1} = f(x_n)$ converges to x^*

Interpretation and Meaning

The Banach Fixed Point Theorem ensures:

- Existence and uniqueness of solutions (fixed points)
- Convergence of approximation by iteration
- Powerful tool in numerical methods and differential equations

A Priori and A Posteriori Approximations

A priori estimate:

$$|x_n - x^*| \leq \frac{L^n}{1 - L} |x_1 - x_0| < \epsilon$$

A posteriori estimate:

$$|x_n - x^*| \leq \frac{L}{1 - L} |x_n - x_{n-1}| \epsilon$$

Steps for a Fixed Point Exercise

Given $f : [a, b] \rightarrow [a, b]$, to prove existence and convergence:

1. Show monotonicity: f is increasing or decreasing.
2. Check interval preservation: $f([a, b]) \subseteq [a, b]$
3. Check Lipschitz continuity: Find $L < 1$
4. Fixed point iteration: Choose x_0 , compute $x_{n+1} = f(x_n)$
5. Apply a priori or a posteriori bound

Example:

$$f(x) = \frac{1}{x} + 3 \text{ on } [2, 5]$$

Step 1: Monotonicity

In this case we will assume that it is monotonous.

Step 2: Domain check

$$x \in [2, 5] \Rightarrow \frac{1}{x} \in [0.2, 0.5] \Rightarrow f(x) \in [3.2, 3.5] \subset [2, 5]$$

Step 3: Lipschitz constant

$$f'(x) = -\frac{1}{x^2} \Rightarrow |f'(x)| \leq \frac{1}{2^2} = \frac{1}{4} < 1 \Rightarrow \text{Lipschitz with } L = \frac{1}{4} < 1$$

Step 4: Fixed point iteration

$$x_0 = 3, \quad x_1 = f(3) = \frac{1}{3} + 3 = \frac{10}{3}, \quad x_2 = f(x_1), \dots$$

Step 5: Approximations

A priori

$$|x_n - x^*| \leq \frac{L^n}{1 - L} |x_1 - x_0|$$

$$|x_n - x^*| \leq \frac{\frac{1}{4}^n}{1 - \frac{1}{4}} \left| \frac{10}{3} - 3 \right|$$

$$\begin{aligned}
|x_n - x^*| &\leq \frac{\left(\frac{1}{4}\right)^n}{1 - \frac{1}{4}} \cdot \left| \frac{10}{3} - 3 \right| \\
&= \frac{\left(\frac{1}{4}\right)^n}{\frac{3}{4}} \cdot \frac{1}{3} \\
&= \frac{4}{3} \cdot \left(\frac{1}{4}\right)^n \cdot \frac{1}{3} \\
&= \frac{4}{9} \cdot \left(\frac{1}{4}\right)^n
\end{aligned}$$

For $\varepsilon = 0, 1$

$$\begin{aligned}
\frac{4}{9} \cdot \left(\frac{1}{4}\right)^n &< \varepsilon \\
\left(\frac{1}{4}\right)^n &< \frac{9}{4}\varepsilon \\
\ln\left(\left(\frac{1}{4}\right)^n\right) &< \ln\left(\frac{9}{4}\varepsilon\right) \\
n \cdot \ln\left(\frac{1}{4}\right) &< \ln\left(\frac{9}{4}\varepsilon\right) \\
n &> \frac{\ln\left(\frac{9}{4}\varepsilon\right)}{\ln\left(\frac{1}{4}\right)} \\
n &> \frac{\ln\left(\frac{9}{4}\varepsilon\right)}{-\ln 4}
\end{aligned}$$

A posteriori

$$|x_n - x^*| \leq \frac{L}{1-L} |x_n - x_{n-1}| = \frac{\frac{1}{4}}{1 - \frac{1}{4}} |x_n - x_{n-1}| = \frac{1}{3} |x_n - x_{n-1}|$$

57.0.14 Uniform Continuity

A function $f : \mathcal{D} \subset \mathbb{R} \rightarrow \mathbb{R}$ is *uniformly continuous* if:

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } |x - y| < \delta \Rightarrow |f(x) - f(y)| < \varepsilon \quad \forall x, y \in \mathcal{D}$$

Example:

$f(x) = \frac{1}{x}$ is not uniformly continuous on $(0, 1)$

Let $\varepsilon = 1$. We want to find numbers in $(0, 1)$ for x, y such that $|x - y| < \delta$ but $|f(x) - f(y)| \geq 1$.

Choose $x \in (0, 1)$ with $x < \delta$ and let $y = \frac{x}{2}$. Then

$$|x - y| = \left| x - \frac{x}{2} \right| = \frac{x}{2} < \frac{\delta}{2} < \delta$$

But,

$$|f(x) - f(y)| = \left| \frac{1}{x} - \frac{1}{y} \right| = \left| \frac{1}{x} - \frac{2}{x} \right| = \left| -\frac{1}{x} \right| = \frac{1}{x} > 1$$

Since $\frac{1}{x} > 1$ for $x < 1$, we have a contraction and therefore f is not uniformly continuous on $(0, 1)$.

57.0.15 Boundedness by Uniform Continuity

If a function $f : (a, b) \rightarrow \mathbb{R}$ is uniformly continuous on the open interval (a, b) , then f is bounded.

57.0.16 Cauchy Sequence by Uniform Continuity

Given $\mathcal{D} \subseteq \mathbb{R}$ and a uniformly continuous function $f : \mathcal{D} \rightarrow \mathbb{R}$. Then for every Cauchy sequence $\{x_n\} \subseteq \mathcal{D}$, the sequence $\{f(x_n)\}$ is also a Cauchy sequence.

Chapter 58

Differentiation

Differentiation is a core concept in calculus. It describes the rate at which a quantity changes and provides tools for analyzing and modeling change in real-world phenomena.

58.0.1 Definition of the Derivative

The derivative of a function f at a point x is defined as the limit:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

or

$$f'(x) = \lim_{h \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

provided this limit exists.

Geometrically, $f'(a)$ is the slope of the tangent line to the graph of f at $x = a$.

58.0.2 Derivative Rules

Power Rule:

$$\frac{d}{dx}x^n = nx^{n-1}, \quad \text{for } n \in \mathbb{R}$$

Constant Rule:

$$\frac{d}{dx}c = 0, \quad \text{for constant } c$$

Constant Multiple Rule:

$$\frac{d}{dx}[c \cdot f(x)] = c \cdot f'(x)$$

Sum and Difference Rule:

$$\frac{d}{dx}[f(x) \pm g(x)] = f'(x) \pm g'(x)$$

Product Rule:

$$\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$$

Quotient Rule:

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$$

Chain Rule:

$$\frac{d}{dx}f(g(x)) = f'(g(x)) \cdot g'(x)$$

58.0.3 Extrema and Critical Points Test

A function f has a critical point at $x = c$ if:

$$f'(c) = 0 \quad \text{or} \quad f'(c) \text{ does not exist}$$

To determine the nature of the critical point:

- *First Derivative Test:* Check sign changes of f' around c
- *Second Derivative Test:*

$$f''(c) > 0 \Rightarrow \text{local minimum} \quad f''(c) < 0 \Rightarrow \text{local maximum}$$

This works, because the second derivative is telling us about the concavity of the function. If it is positive then we will have function which is concave up a minimum; and if negative concave down a maximum.

58.0.4 Trigonometric Derivatives

$$\frac{d}{dx} \sin x = \cos x$$

$$\frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tan x = \sec^2 x$$

$$\frac{d}{dx} \cot x = -\csc^2 x$$

$$\frac{d}{dx} \sec x = \sec x \tan x$$

$$\frac{d}{dx} \csc x = -\csc x \cot x$$

58.0.5 Derivatives of Inverse Trigonometric Functions

$$\begin{aligned}\frac{d}{dx} \arcsin x &= \frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx} \arccos x &= -\frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx} \arctan x &= \frac{1}{1+x^2} \\ \frac{d}{dx} \operatorname{arccot} x &= -\frac{1}{1+x^2} \\ \frac{d}{dx} \operatorname{arcsec} x &= \frac{1}{|x|\sqrt{x^2-1}} \\ \frac{d}{dx} \operatorname{arccsc} x &= -\frac{1}{|x|\sqrt{x^2-1}}\end{aligned}$$

58.0.6 The Differential

The differential dy of a function $y = f(x)$ is defined as:

$$dy = f'(x) dx$$

This linear approximation estimates the change in y for a small change in x .

58.0.7 The Secant Equation and Graphical Meaning

The *secant line* through points $(x_0, f(x_0))$ and $(x_0 + h, f(x_0 + h))$ has slope:

$$\frac{f(x_0 + h) - f(x_0)}{h}$$

As $h \rightarrow 0$, the secant line approaches the tangent line. Graphically, this means:

$$\lim_{h \rightarrow 0} \text{slope of secant} = \text{slope of tangent} = f'(x) = \Delta_h$$

The secant line equation is given by:

$$L(x) = f(x_0) + \Delta_h f'(x_0)(x - x_0)$$

58.0.8 Optimization via the Derivative

To find local or global extrema:

1. Compute $f'(x)$
2. Solve $f'(x) = 0$ to find critical points
3. Use the first or second derivative test to classify
4. Evaluate endpoints if optimizing over a closed interval

Example I:

Maximize area $A = x(10 - 2x)$

$$A = 10x - 2x^2, \quad A' = 10 - 4x, \quad A' = 0 \Rightarrow x = 2.5$$

Example II:

Given a number $c > 0$, we want to factor it into two positive numbers x and y such that their product is c , and their sum $S = x + y$ is minimal.

Since $x \cdot y = c$, we can express one variable in terms of the other, e.g.,

$$y = \frac{c}{x}$$

Substituting into the sum:

$$S(x) = x + \frac{c}{x}$$

To minimize $S(x)$, we take the derivative:

$$S'(x) = 1 - \frac{c}{x^2}$$

Setting the derivative to zero gives:

$$1 - \frac{c}{x^2} = 0 \quad \Rightarrow \quad x^2 = c \quad \Rightarrow \quad x = \sqrt{c}$$

Then:

$$y = \frac{c}{x} = \sqrt{c}$$

So the minimal sum occurs when $x = y = \sqrt{c}$, and the minimum value of the sum is:

$$x + y = 2\sqrt{c}$$

Let $c = 25$. Then the two positive factors that minimize the sum are:

$$x = y = \sqrt{25} = 5$$

and the minimal sum is:

$$x + y = 5 + 5 = 10$$

58.0.9 Power Rule Derivation

We derive the power rule:

$$\frac{d}{dx}x^n = nx^{n-1}$$

for $n \in \mathbb{N}$ using the limit definition:

$$\frac{d}{dx}x^n = \lim_{h \rightarrow 0} \frac{(x+h)^n - x^n}{h}$$

Use the Binomial Theorem:

$$(x+h)^n = x^n + nx^{n-1}h + \cdots + h^n \Rightarrow \text{Only the linear term survives after dividing by } h$$

58.0.10 Implicit Differentiation

Given:

$$9x^2 + 4y^2 = 25$$

Differentiate both sides implicitly:

$$18x + 8y \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{18x}{8y} = -\frac{9x}{4y}$$

This method is used when y is not isolated and is defined implicitly.

Chapter 59

Integration

Integration is the reverse process of differentiation and is used to calculate areas, volumes, accumulated change, and more.

59.0.1 Definition (Riemann Integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. We define the *definite integral* as:

$$\int_a^b f(x) dx = \lim_{\|P\| \rightarrow 0} \sum_{i=1}^n f(\xi_i) \Delta x_i$$

Where $P = \{x_0, \dots, x_n\}$ is a partition, $\xi_i \in [x_{i-1}, x_i]$, and $\Delta x_i = x_i - x_{i-1}$.

59.0.2 Upper and Lower Sums

Let f be bounded on $[a, b]$ and P a partition:

$$\underline{S}(f, P) = \sum_{i=1}^n m_i \Delta x_i, \quad \overline{S}(f, P) = \sum_{i=1}^n M_i \Delta x_i$$

with:

$$m_i = \inf_{x \in [x_{i-1}, x_i]} f(x), \quad M_i = \sup_{x \in [x_{i-1}, x_i]} f(x)$$

If $\sup \underline{S} = \inf \overline{S}$, then f is Riemann integrable.

59.0.3 Concrete and Non-Concrete Integrals

- *Concrete*: Can be explicitly evaluated with anti-derivatives.
- *Non-concrete*: Cannot be integrated in elementary terms (e.g., $\int e^{-x^2} dx$).

59.0.4 Rules of Integration

- $\int 0 dx = C$
- $\int c dx = cx$
- $\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (n \neq -1)$
- $\int \frac{1}{x} dx = \ln |x| + C$
- $\int e^x dx = e^x + C$

- $\int \sin x \, dx = -\cos x + C$
- $\int \cos x \, dx = \sin x + C$
- Linearity: $\int (af + bg) \, dx = a \int f \, dx + b \int g \, dx$

59.0.5 Substitution Rule (U-Substitution)

If $u = g(x)$ and $f = F'$, then:

$$\int f(g(x))g'(x) \, dx = \int f(u) \, du$$

Example:

$$\int \frac{1}{(x-1)^4} \, dx, \quad u = x-1 \Rightarrow \int \frac{1}{u^4} \, du = -\frac{1}{3u^3} + C = -\frac{1}{3(x-1)^3} + C$$

59.0.6 Integration by Parts

$$\int u \, dv = uv - \int v \, du$$

ILATE: Use this order to choose u :

1. Inverse Trig
2. Logarithmic
3. Algebraic
4. Trigonometric
5. Exponential

Cyclic Integrals: Some integrals cycle back to the original (e.g., $e^x \sin(x)$).

Example:

$$\int e^x \sin(x) \, dx$$

Let:

$$u = \sin(x), \, dv = e^x \Rightarrow du = \cos(x) \, dx, \, v = e^x$$

$$\int e^x \sin(x) \, dx = e^x \sin(x) - \int e^x \cos(x) \, dx$$

$$u = \cos(x), \, dv = e^x \Rightarrow du = -\sin(x) \, dx, \, v = e^x$$

$$\begin{aligned} \int e^x \sin(x) \, dx &= e^x \sin(x) - \left[\cos(x)e^x - \int \sin(x)e^x \, dx \right] \\ \int e^x \sin(x) \, dx &= e^x \sin(x) - \cos(x)e^x - \int \sin(x)e^x \, dx \\ 2 \int e^x \sin(x) \, dx &= e^x \sin(x) - \cos(x)e^x \\ \int e^x \sin(x) \, dx &= \frac{e^x \sin(x) + \cos(x)e^x + C}{2} \end{aligned}$$

59.0.7 Integration by Partial Fractions

Applies to rational functions. Four cases:

- *Linear factors*

Denominator: $(x - a)(x - b)$

Decomposition:

$$\frac{P(x)}{(x - a)(x - b)} = \frac{A}{x - a} + \frac{B}{x - b}$$

- *Repeated linear factors*

Denominator: $(x - a)^n$

Decomposition:

$$\frac{P(x)}{(x - a)^n} = \frac{A_1}{x - a} + \frac{A_2}{(x - a)^2} + \cdots + \frac{A_n}{(x - a)^n}$$

- *Irreducible quadratic factors*

Denominator: $(x^2 + px + q)$, where $x^2 + px + q$ cannot be factored over the reals

Decomposition:

$$\frac{P(x)}{x^2 + px + q} = \frac{Ax + B}{x^2 + px + q}$$

- *Repeated irreducible quadratic factors*

Denominator: $(x^2 + px + q)^n$

Decomposition:

$$\frac{P(x)}{(x^2 + px + q)^n} = \frac{A_1x + B_1}{x^2 + px + q} + \frac{A_2x + B_2}{(x^2 + px + q)^2} + \cdots + \frac{A_nx + B_n}{(x^2 + px + q)^n}$$

Example:

$$\int \frac{1}{x^2 - 1} dx = \int \left(\frac{1}{2(x - 1)} - \frac{1}{2(x + 1)} \right) dx = \frac{1}{2} \ln \left| \frac{x - 1}{x + 1} \right| + C$$

59.0.8 Trigonometric Substitution

Trigonometric substitution is useful for evaluating integrals involving square roots of quadratic expressions.

- $\sqrt{a^2 - x^2}$: use $x = a \sin \theta$
- $\sqrt{a^2 + x^2}$: use $x = a \tan \theta$
- $\sqrt{x^2 - a^2}$: use $x = a \sec \theta$

Example I:

$$\int \frac{1}{\sqrt{4 - x^2}} dx$$

We use: $x = 2 \sin \theta \Rightarrow dx = 2 \cos \theta d\theta$

$$\sqrt{4 - x^2} = \sqrt{4 - 4 \sin^2 \theta} = \sqrt{4 \cos^2 \theta} = 2 \cos \theta$$

Substitute:

$$\int \frac{1}{\sqrt{4 - x^2}} dx = \int \frac{1}{2 \cos \theta} \cdot 2 \cos \theta d\theta = \int d\theta = \theta + C$$

Return to x using:

$$x = 2 \sin \theta \Rightarrow \theta = \arcsin \left(\frac{x}{2} \right) \Rightarrow \int \frac{1}{\sqrt{4 - x^2}} dx = \arcsin \left(\frac{x}{2} \right) + C$$

Example II:

$$\int \frac{x^3}{\sqrt{x^2+4}} dx$$

Use: $x = 2 \tan \theta \Rightarrow dx = 2 \sec^2 \theta d\theta$

$$\sqrt{x^2+4} = \sqrt{4 \tan^2 \theta + 4} = \sqrt{4 \sec^2 \theta} = 2 \sec \theta$$

Substitute:

$$\int \frac{x^3}{\sqrt{x^2+4}} dx = \int \frac{(2 \tan \theta)^3}{2 \sec \theta} \cdot 2 \sec^2 \theta d\theta = \int \frac{8 \tan^3 \theta}{\sec \theta} \cdot 2 \sec^2 \theta d\theta$$

Simplify:

$$\begin{aligned} 8 \int \tan^2 \theta \sec \theta \tan \theta dx &= 8 \int (\sec^2 \theta - 1) \sec \theta \tan \theta dx \\ &= 8 \int (\sec^2 \theta \sec \theta - \sec \theta) \tan \theta dx \\ &= 8 \left(\int \sec^2 \theta \sec \theta \tan \theta dx - \int \sec \theta \tan \theta dx \right) \\ &= 8 \left(\int \sec^2 \theta \sec \theta dx - \int \sec \theta \tan \theta dx \right) \end{aligned}$$

For the first one let $u = \sec^3 \theta$ and $du = 3 \sec^2 \theta d\theta$

$$\int u du = \frac{\sec^3 \theta}{3} + c$$

For the second one

$$\int \sec \theta \tan \theta = \sec \theta + c$$

Together

$$8 \left(\frac{\sec^3 \theta}{3} - \sec \theta + c \right)$$

Then solve the integral using trigonometric identities and reduction formulas. Return to x using:

$$\tan \theta = \frac{x}{2} \quad \sec \theta = \frac{\sqrt{x^2+4}}{2}$$

This leaves us with

$$\frac{8}{3} \left(\frac{\sqrt{x^2+4}}{2} \right)^3 - 8 \left(\frac{\sqrt{x^2+4}}{2} \right) + c$$

Example III:

$$\int \frac{x^3}{\sqrt{x^4-4}} dx$$

Use: $x = 2 \sec \theta, \quad dx = 2 \sec \theta \tan \theta d\theta$

Then

$$\int \frac{(2 \sec \theta)^2 2 \sec \theta \tan \theta d\theta}{2 \tan \theta} = \int 8 \sec^4 \theta d\theta$$

$$8 \int (\tan^2 \theta + 1) \sec^2 \theta = 8 \left(\int \tan^2 \theta \sec^2 \theta + \int \sec^2 \theta \right)$$

Now let $u = \tan \theta$ and $du = \sec^2 \theta d\theta$

$$8 \left(\int u du + \int \sec^2 \theta d\theta \right) = \frac{8}{3} \tan^3 \theta + 8 \tan \theta + c$$

Now we return to x via our triangle:

$$\tan \theta = \frac{\sqrt{x^2 + 4}}{2}$$

$$\frac{8}{3} \left(\frac{\sqrt{x^2 + 4}}{2} \right)^3 + 8 \frac{\sqrt{x^2 + 4}}{2} + c$$

59.0.9 Average Value of a Function

$$\bar{f} = \frac{1}{b-a} \int_a^b f(x) dx$$

59.0.10 Mean Value Theorem for Integrals

If f is continuous on $[a, b]$, then:

$$\exists c \in [a, b] \text{ such that } \int_a^b f(x) dx = f(c)(b-a)$$

59.0.11 Length, Area, Volume Formulas

Arc length:

$$L = \int_a^b \sqrt{1 + f'(x)^2} dx$$

Area between curves:

$$A = \int_a^b |f(x) - g(x)| dx$$

Volume (disk):

$$V = \pi \int_a^b f(x)^2 dx$$

Shell:

$$V = 2\pi \int_a^b x f(x) \sqrt{1 + f'(x)^2} dx$$

59.0.12 Differentiation of Integrals with Variable Bounds

If:

$$F(x) = \int_{a(x)}^{b(x)} f(t) dt$$

Then:

$$F'(x) = f(b(x)) \cdot b'(x) - f(a(x)) \cdot a'(x)$$

59.0.13 Parameter-Dependent Integrals

Let $F(p) = \int_a^b f(x, p) dx$. If f is continuous and differentiable in p , then:

$$\frac{d}{dp} \int_a^b f(x, p) dx = \int_a^b \frac{\partial f}{\partial p}(x, p) dx$$

59.0.14 Leibniz Rule for Differentiating Under the Integral Sign

$$\frac{\partial F}{\partial x} \int_{a(x)}^{b(x)} f(x, t) dx = f(x, b(x)) \cdot b'(x) - f(x, a(x)) \cdot a'(x) + \int_{a(x)}^{b(x)} \frac{\partial f}{\partial t}(x, t) dt$$

59.0.15 Improper Integrals with infinite Bounds

$$\int_a^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_a^R f(x) dx$$

The value when the bounds look like $\int_{-\infty}^\infty f(x) dx$ is called the *Cauchy-Main-Value*.

$$\int_{-\infty}^\infty f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^c f(x) dx + \int_c^R f(x) dx$$

59.0.16 Improper Integrals with Discontinuity

These integrals have some kind of asymptote between the bounds we want to evaluate. More concrete:

$$\int_a^b f(x) dx \text{ improper if } f \text{ has an infinite discontinuity on } [a, b]$$

To tackle this type of problems we just like before use a limit to approach the discontinuity without reaching it.

$$\int_a^b dx = \lim_{\varepsilon \rightarrow 0} \int_a^{c-\varepsilon} f(x) dx + \int_{c+\varepsilon}^b f(x) dx$$

Here c is the value of the discontinuity.

Example:

$$\int_0^1 \frac{1}{2\sqrt{x}}$$

Discontinuity at $c = 0$

$$\begin{aligned}
\int_0^1 \frac{1}{2\sqrt{x}} &= \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon}^1 \frac{1}{2\sqrt{x}} dx \\
&= \lim_{\varepsilon \rightarrow 0} \sqrt{x} \Big|_{\varepsilon}^1 \\
&= 1 - \lim_{\varepsilon \rightarrow 0} \sqrt{\varepsilon} = 1
\end{aligned}$$

59.0.17 Convergence Criteria for Improper Integrals

- *Majorant:* If $0 \leq f(x) \leq g(x)$ and $\int g(x) dx$ converges, then so does $\int f(x) dx$.
- *Minorant:* If $f(x) \geq g(x) \geq 0$ and $\int g(x) dx$ diverges, then $\int f(x) dx$ diverges.

59.0.18 Integral Test for Series

Let $f(n) = a_n$ where f is continuous, decreasing, and positive for $n \geq N$:

$$\sum_{n=N}^{\infty} a_n \text{ converges} \iff \int_N^{\infty} f(x) dx \text{ converges}$$

Example:

$$\sum \frac{1}{n^p} \text{ converges} \iff p > 1$$

59.0.19 Integrals with Absolute Value

When given an integral with an absolute value in it, we have to take a look at the function to see how it behaves when x is negative and then split the integral into two or more integrals where the border contain the critical point.

Example:

Integrate

$$\int_{-\infty}^{\infty} e^{-|x|} dx$$

This is function will be split at zero, and we will get

$$\begin{aligned}
&\int_{-\infty}^{\infty} e^{-|x|} dx \\
&\lim_{R \rightarrow \infty} \int_{-R}^0 e^x dx + \int_0^R e^{-x} dx \\
&\lim_{R \rightarrow \infty} 1 - e^{-R} + -e^{-R} + 1 = 2
\end{aligned}$$

Chapter 60

Important Theorems of Single Variable Calculus

60.0.1 Intermediate Value Theorem

Let f be a continuous function on the closed interval $[a, b]$. If $f(a) \neq f(b)$ and N is any number between $f(a)$ and $f(b)$, then there exists $c \in (a, b)$ such that $f(c) = N$.

Proof:

Without loss of generality, assume $f(a) < N < f(b)$. Define:

$$S = \{x \in [a, b] \mid f(x) < N\}$$

Since $a \in S$, S is non-empty and bounded above by b . Let $c = \sup S$. We want to show that $f(c) = N$ by satisfying the inequalities $f(c) \geq N$ and $f(c) \leq N$.

If $f(c) < N$, by the continuity of f , there exists $\delta > 0$ such that for all $x \in (c, c + \delta)$:

$$f(x) < N$$


This contradicts the definition of c as the supremum of S .

If $f(c) > N$, again by the continuity of f , there exists $\delta > 0$ such that for all $x \in (c - \delta, c)$:

$$f(x) > N$$

This contradicts the definition of c as the supremum of S . Hence, we must have $f(c) = N$.

□◻◻



60.0.2 Extreme Value Theorem

If f is continuous on the closed interval $[a, b]$, then f attains a maximum and a minimum value on $[a, b]$. That is, there exist $c, d \in [a, b]$ such that:

$$f(c) \leq f(x) \leq f(d) \quad \text{for all } x \in [a, b]$$

Proof:

Since f is continuous on the compact interval $[a, b]$, the image $f([a, b]) \subset \mathbb{R}$ is also compact. A compact subset of $\mathbb{R}$ is closed and bounded, and hence attains its supremum and infimum.

Therefore, there exist $c, d \in [a, b]$ such that:

$$f(c) = \min\{f(x) : x \in [a, b]\}, \quad f(d) = \max\{f(x) : x \in [a, b]\}$$

□◻

60.0.3 Rolle's Theorem


Let f be continuous on $[a, b]$, differentiable on (a, b) , and suppose $f(a) = f(b)$. Then, there exists $c \in (a, b)$ such that $f'(c) = 0$.

Proof:

By the Extreme Value Theorem, f attains a maximum or minimum at some point $c \in [a, b]$.

If $c \in (a, b)$, then by Fermat's theorem, $f'(c) = 0$. If the maximum or minimum occurs at a or b , since $f(a) = f(b)$, f must be constant, so $f'(x) = 0$ everywhere on (a, b) , and in particular for some $c \in (a, b)$.

□◻



60.0.4 Mean Value Theorem

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then there exists $c \in (a, b)$ such that:

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Proof:

Define the auxiliary function:

$$g(x) = f(x) - \left(\frac{f(b) - f(a)}{b - a} \right) (x - a)$$


Then $g(a) = f(a)$, $g(b) = f(b) - m(b - a) = f(a) \Rightarrow g(a) = g(b)$

Apply Rolle's theorem to $g(x)$: Since $g(a) = g(b)$, and g is continuous and differentiable, there exists $c \in (a, b)$ such that $g'(c) = 0$

Then:

$$g'(x) = f'(x) - \left(\frac{f(b) - f(a)}{b - a} \right) \Rightarrow f'(c) = \frac{f(b) - f(a)}{b - a}$$

□◻◻



Chapter 61

Coordinate Systems

Coordinate systems provide different ways of describing points in space, adapted to the geometry of the problem. Below are the most common systems used in multi-variable calculus.

61.0.1 Cartesian Coordinates

The *Cartesian coordinate system* uses mutually orthogonal axes. A point in $\mathbb{R}^2$ is given by:

$$(x, y) \quad \text{and in } \mathbb{R}^3 : \quad (x, y, z)$$

This is the most natural system for rectangular domains and is the basis for most definitions in vector calculus. The coordinate axes are orthogonal and equidistant.

61.0.2 Polar Coordinates

In polar coordinates, a point $(x, y) \in \mathbb{R}^2$ is described by:

$$r \geq 0 \text{ (radius from origin),} \quad \varphi \in [0, 2\pi) \text{ (angle from x-axis)}$$

Transformations:

$$x = r \cos \varphi, \quad y = r \sin \varphi$$

This comes from our right triangle with $r = 1$ where

$$\cos \varphi = \frac{x}{r} \quad \sin \varphi = \frac{y}{r}$$

Inverse:

$$r = \sqrt{x^2 + y^2}, \quad \varphi = \arctan \left(\frac{y}{x} \right)$$

This system is useful for circular or radial symmetry. The Jacobian determinant is:

$$|J| = r dr d\theta$$

61.0.3 Cylindrical Coordinates

Cylindrical coordinates extend polar coordinates by adding a height z component:

$$(r, \varphi, z) \quad \text{where } r \geq 0, \varphi \in [0, 2\pi), z \in \mathbb{R}$$

Transformations:

$$x = r \cos \varphi, \quad y = r \sin \varphi, \quad z = z$$

Inverse:

$$r = \sqrt{x^2 + y^2}, \quad \varphi = \arctan\left(\frac{y}{x}\right), \quad z = z$$

This coordinate system is ideal for objects with rotational symmetry about the z -axis. The Jacobian determinant is:

$$|J| = r dr d\theta dz$$

61.0.4 Spherical Coordinates

In spherical coordinates, a point $(x, y, z) \in \mathbb{R}^3$ is represented using:

$$\rho \geq 0 \text{ (radial distance), } \theta \in [0, \pi] \text{ (inclination), } \varphi \in [0, 2\pi) \text{ (azimuth)}$$

Transformations:

$$x = \rho \sin \theta \cos \varphi, \quad y = \rho \sin \theta \sin \varphi, \quad z = \rho \cos \theta$$

Inverse:

$$\rho = \sqrt{x^2 + y^2 + z^2}, \quad \theta = \arccos\left(\frac{z}{\rho}\right), \quad \varphi = \arctan\left(\frac{y}{x}\right)$$

Spherical coordinates are useful for spherical symmetry (e.g., in gravitational or electric fields). The Jacobian determinant is:

$$|J| = \rho^2 \sin \theta d\rho d\varphi d\theta$$

Each coordinate system is tailored to specific geometries and greatly simplifies integrals when used appropriately. Coordinate transformations require careful application of the Jacobian determinant to account for stretching or distortion of space.

Chapter 62

Parameterized Curves

62.0.1 Definition

A *parameterized curve* is a mathematical object where each point on the curve is described as a function of one or more parameters, typically denoted by t . Formally, a parameterized curve in $\mathbb{R}^n$ is a continuous function:

$$\mathbf{r}(t) = (x(t), y(t), z(t), \dots)$$

Where t ranges over some interval $I \subseteq \mathbb{R}$, and each component function (e.g., $x(t)$, $y(t)$) is real-valued.

62.0.2 Purpose

The purpose of parameterizing a curve is to provide a flexible and detailed way of describing geometric objects, particularly when they cannot be easily expressed as a simple function $y = f(x)$. Parameterizations allow us to:

- Describe curves that loop, cross themselves, or have vertical segments.
- Compute quantities like arc length, tangent vectors, and curvature.
- Easily model motion along a curve, where t can represent time.

62.0.3 From parametric to explicit form

This process is rather simple because in the case where it is possible to change from one form to the other we only need to write the parameter in terms of one of the variables and then substitute.

62.0.4 How to Parameterize a Function

To parameterize a curve or a function:

1. Introduce a new parameter t , often representing time or distance along the curve.
2. Express the coordinates (x, y) (or (x, y, z)) as functions of t .
3. Ensure that as t varies over a chosen interval, the set of points $(x(t), y(t))$ traces the desired curve.

Sometimes, parameterizations are natural (e.g., circles using trigonometric functions), while in other cases, one must carefully design a suitable parameterization based on the curve's properties.

Example:

Parameterizing a Circle:

Consider the unit circle defined by the equation:

$$x^2 + y^2 = 1$$

A natural parameterization uses the trigonometric functions sine and cosine:

$$x(t) = \cos(t), \quad y(t) = \sin(t)$$

For $t \in [0, 2\pi]$.

As t varies from 0 to 2π , the point $(x(t), y(t))$ traces out the entire circle exactly once.

62.0.5 Arclength and t Parameter

Recall that the arclength of a curve in 3 dimensions is

$$\int_a^b \sqrt{f'(t)^2 + g'(t)^2 + h'(t)^2} dt,$$

which is norm of our tangent vector.

$$\int_a^b \sqrt{|\vec{v}(t)|^2} dt$$

Arclength Parameter

This is given by

$$s(t) = \int_a^t \sqrt{|\vec{\tau}(t)|^2} dt,$$

where τ is the length parameter.

This parameter is intrinsic to the curve, and represents one choice because there only one valid interpretation of the length. The downside is that this is hard to compute.

The t Parameter

For the t parameter, we can say that this is intrinsic to the object along the curve, it can be interpreted in multiple ways therefore, many choices and is easy to compute.

Chapter 63

Multi-variable Functions

63.0.1 Open Sets

Let $\|\cdot\|$ be a norm on $\mathbb{R}^n$. The ε -neighborhood of a point $\vec{x}_0 \in \mathbb{R}^n$ is defined as:

$$U_\varepsilon(\vec{x}_0) := \{\vec{x} \in \mathbb{R}^n \mid \|\vec{x} - \vec{x}_0\| < \varepsilon\}$$

A point $\vec{x}_0 \in D \subseteq \mathbb{R}^n$ is called an *interior point* of D if there exists $\varepsilon > 0$ such that $U_\varepsilon(\vec{x}_0) \subseteq D$. A set D is called *open* if all its points are interior points.

Although ε -neighborhoods depend on the chosen norm, due to the equivalence of norms in $\mathbb{R}^n$, openness is norm-independent.

63.0.2 Closed Sets

Let $D \subseteq \mathbb{R}^n$ and $\|\cdot\|$ a norm. A point $\vec{x}_0$ is called an *accumulation point* (or *limit point*) of D if for every $\varepsilon > 0$, the neighborhood $U_\varepsilon(\vec{x}_0)$ contains a point $\vec{x} \neq \vec{x}_0$ such that $\vec{x} \in D$.

A set $D \subseteq \mathbb{R}^n$ is called *closed* if it contains all its accumulation points.

63.0.3 Boundedness and Order

A set $D \subseteq \mathbb{R}^n$ is called *bounded* if there exists a real number $M > 0$ such that:

$$\|\vec{x}\| < M \quad \text{for all } \vec{x} \in D$$

If no such bound M exists, the set is called *unbounded*.

63.0.4 Sequences

Let $(\vec{x}_n) \subseteq \mathbb{R}^n$ be a sequence of vectors. We say that $\vec{x}_n \rightarrow \vec{x} \in \mathbb{R}^n$ as $n \rightarrow \infty$ if for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that:

$$\|\vec{x}_n - \vec{x}\| < \varepsilon \quad \text{for all } n > n_0$$

The sequence $(\vec{x}_n)$ is called *convergent* if such $\vec{x}$ exists.

The sequence $(\vec{x}_n)$ is called a *Cauchy sequence* if for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that:

$$\|\vec{x}_n - \vec{x}_m\| < \varepsilon \quad \text{for all } n, m > n_0$$

Example I:

$$\lim_{n \rightarrow \infty} \vec{X}_n = \lim_{n \rightarrow \infty} \begin{bmatrix} \frac{1}{n} \\ 1 + \frac{1}{n^2} \\ \frac{\sin n}{n} \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

63.0.5 Different Norms

From linear algebra it is known what a *Norm* is and its properties:

Let V be a vector space over $\mathbb{R}$ or $\mathbb{C}$. A function $\|\cdot\| : V \rightarrow \mathbb{R}$ is called a norm if for all $x, y \in V$ and all scalars α , the following properties hold:

1. *Non-negativity*: $\|x\| \geq 0$, and $\|x\| = 0$ if and only if $x = 0$.
2. *Absolute homogeneity*: $\|\alpha x\| = |\alpha| \cdot \|x\|$.
3. *Triangle inequality*: $\|x + y\| \leq \|x\| + \|y\|$.

Commonly used norms in $\mathbb{R}^n$ (or $\mathbb{C}^n$) include:

- *Sum Norm* (also known as the ℓ_1 norm):

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- *Max Norm* (also known as the ℓ_∞ norm):

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$$

- *Euclidean Norm* (also known as the ℓ_2 norm):

$$\|x\|_2 = \left(\sum_{i=1}^n |x_i|^2 \right)^{1/2}$$

Example II:

Given is

$$\vec{X}_n = \begin{bmatrix} 5 - \frac{10}{n} \\ \frac{1}{n} \end{bmatrix}$$

we want to find n_0 for $\varepsilon = 0, 1$

$$\lim_{n \rightarrow \infty} \vec{X}_n = \begin{bmatrix} 5 \\ 0 \end{bmatrix} = \vec{X}_0$$

Now use the definition for convergence and *Euclidean Norm*

$$\|\vec{X}_n - \vec{X}_0\| < \varepsilon$$

$$\left\| \begin{bmatrix} -\frac{10}{n} \\ \frac{1}{n} \end{bmatrix} \right\|$$

$$\frac{\sqrt{101}}{n} < 0, 1$$

$$n > 0, 1\sqrt{101} = 100, 5$$

Therefore, $n_0 = 101$.

Equivalence of Norms

In finite-dimensional vector spaces, all norms are equivalent. That is, if $\|\cdot\|_a$ and $\|\cdot\|_b$ are two norms on a finite-dimensional vector space V , then there exist constants $c, C > 0$ such that for all $x \in V$,

$$c\|x\|_a \leq \|x\|_b \leq C\|x\|_a$$

This implies that while different norms may measure length differently, they all induce the same topology and convergence behavior in finite-dimensional spaces.

Equivalence in the Reals

In the $\mathbb{R}^n$ all norms are equivalent and

$$\begin{aligned}\|\vec{x}\|_\infty &\leq \|\vec{x}\|_2 \leq \sqrt{n}\|\vec{x}\|_\infty \\ \|\vec{x}\|_\infty &\leq \|\vec{x}\|_1 \leq n\|\vec{x}\|_\infty \\ \|\vec{x}\|_2 &\leq \|\vec{x}\|_1 \leq n\|\vec{x}\|_2\end{aligned}$$

63.0.6 Accumulation Point

A vector $\vec{x}_0 \in \mathbb{R}^n$ is called an *accumulation point* of a set $D \subseteq \mathbb{R}^n$ if for every $\varepsilon > 0$, the neighborhood $U_\varepsilon(\vec{x}_0)$ contains a point $\vec{x} \in D \setminus \{\vec{x}_0\}$.

In the context of sequences, an accumulation point of a sequence $(\vec{x}_n)$ is a point $\vec{x} \in \mathbb{R}^n$ for which there exists a subsequence $(\vec{x}_{n_k})$ such that $\vec{x}_{n_k} \rightarrow \vec{x}$.

63.0.7 Bolzano–Weierstraß Theorem

Every infinite, bounded sequence in $\mathbb{R}^n$ has at least one accumulation point.

- Every bounded infinite sequence has at least one convergent subsequence.
- A bounded sequence is *convergent* if and only if it has exactly one accumulation point.
- Every Cauchy sequence in $\mathbb{R}^n$ is convergent (since $\mathbb{R}^n$ is complete).

63.0.8 Limits

A function $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ has the limit $g \in \mathbb{R}$ at a point $\vec{X}_0 \in U$ if:

$$\lim_{\vec{X} \rightarrow \vec{X}_0} f(\vec{X}) = g$$

means that for every sequence $(\vec{X}_n) \subset U$ converging to $\vec{X}_0$, we have:

$$\lim_{n \rightarrow \infty} f(\vec{X}_n) = g$$

The convergence must hold along all possible paths to the point $\vec{X}_0$, making the multi-variable limit path-independent and unique (if it exists).

Example:

$$\vec{X}_n = \begin{bmatrix} \frac{1}{n} \\ 1 + \frac{1}{n^2} \\ \frac{\sin(n)}{n} \end{bmatrix}$$

The limit towards infinity is then

$$\lim_{n \rightarrow \infty} \begin{bmatrix} \frac{1}{n} \\ 1 + \frac{1}{n^2} \\ \frac{\sin(n)}{n} \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

63.0.9 Partial Functions

For a function $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, a *partial function* is a restriction of f to a variable by fixing all others.

Examples in $\mathbb{R}^3$:

$$f(x_1, x_2, x_3) = 5x_1 + x_2x_3$$

Fixing values:

$$x_2 = 3, \quad x_3 = 5 \Rightarrow f_1(x_1) = 5x_1 + 15$$

$$x_1 = 2, \quad x_3 = 5 \Rightarrow f_2(x_2) = 10 + 5x_2x_1 = 2, \quad x_2 = 3 \Rightarrow f_3(x_3) = 10 + 3x_3$$

Each f_i reduces the multi-variable function to a single-variable slice (cross-section), allowing analysis along coordinate axes.

63.0.10 Continuity

Let $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, and $\vec{X}_0 \in U$. The function f is said to be *continuous at $\vec{X}_0$* if:

$$\lim_{\vec{X} \rightarrow \vec{X}_0} f(\vec{X}) = f(\vec{X}_0)$$

This means that for any sequence $(\vec{X}_n) \rightarrow \vec{X}_0$, we have:

$$\lim_{n \rightarrow \infty} f(\vec{X}_n) = f(\vec{X}_0)$$

To check if a multi-variable function is continuous it is enough to prove that it is continuous towards the origin

$$\begin{aligned} \lim_{\vec{X} \rightarrow \vec{X}_0} f(\vec{X}) &= f(\vec{X}_0) \\ \iff \lim_{\vec{X} \rightarrow \vec{X}_0} (f(\vec{X}) - f(\vec{X}_0)) &= 0 \\ \iff \lim_{\vec{h} \rightarrow 0} (f(\vec{X}_0 + \vec{h}) - f(\vec{X}_0)) &= 0 \\ \iff \lim_{\vec{h} \rightarrow 0} g(\vec{h}) &= 0 \end{aligned}$$

If we are checking if a function is continuous and get an *indeterminate form* there are three main approaches.

1. *Factorizing*: Either normal factorization, adding a clever zero or multiplying by a clever one.
2. *Change of Coordinates*: Sometimes changing the coordinate system allows us to simplify the expression.
3. *Different Paths*: Trying different paths can be pretty straightforward. If we get two different values for the same point across different paths then the function is not continuous.

Example:

Given is $\frac{x^3+2yx^2+y^2x+2y^3}{x+2y}$ for $(x, y) \neq (0, 0)$ and 0 for $(0, 0)$.

$$\begin{aligned}\frac{x^3+2yx^2+y^2x+2y^3}{x+2y} &= \frac{x^2(x+2y)+y(x+2y)}{x+2y} \\ &= \frac{(x+2y)(x^2+y^2)}{x+2y} \\ &= x^2+y^2\end{aligned}$$

Now by taking the limit we see that

$$\lim_{(x,y) \rightarrow (0,0)} x^2+y^2 = 0$$

Thus, the function is continuous.

Example:

Check for continuity towards the origin of $f(x, y) = \frac{xy^2}{x^2+y^4}$.

For the path $x = y$ and $y = 0$

$$f(x, 0) = \frac{x \cdot 0}{x^2 + 0} = 0$$

For the path $x = y^2$

$$f(y^2, y) = \frac{y^4}{2y^4} = \frac{1}{2}$$

Because we got two different values the function is not continuous.

Continuity in Polar Coordinates for 2D

Let $\vec{X} = (x, y)$ be written in polar coordinates:

$$x = r \cos \varphi, \quad y = r \sin \varphi$$

Then:

$$\vec{X}_n \rightarrow (0, 0) \iff r_n \rightarrow 0$$

This alternative approach helps evaluate limits and continuity at the origin using radial convergence.

Example:

Given is

$$f(xy) = \begin{cases} \frac{xy}{x^2+y^2} & \text{for } (x, y) \neq (0, 0) \\ 0 & \text{else} \end{cases}$$

Let $x = r \cos(\theta)$ and $y = r \sin(\theta)$

$$\begin{aligned}
& \lim_{r \rightarrow 0} \frac{r \cos(\theta) r \sin(\theta)}{r^2 \cos^2(\theta) + r^2 \sin^2(\theta)} \\
&= \lim_{r \rightarrow 0} \frac{r^2 \cos(\theta) \sin(\theta)}{r^2 (\cos^2(\theta) + \sin^2(\theta))} \\
&= \lim_{r \rightarrow 0} \frac{\cos(\theta) \sin(\theta)}{1}
\end{aligned}$$

Because of the dependence on θ which can give us for $\theta = \frac{\pi}{4}$ $\frac{1}{2}$ this functions is not continuous.

Substitution

When dealing with multi-variable functions we can sometimes use *substitution* for converting our function into a single variable function to use *L'Hospital's Rule*.

Example:

Given is $\frac{e^{x^2+y^2}-1}{x^2+y^2}$.

See, that we have repeated terms we can substitute. $p^2 = x^2 + y^2$.

$$\frac{e^{p^2} - 1}{p^2}$$

Now differentiate

$$\frac{2pe^{p^2}}{2p} = e^{p^2}$$

Now take the limit

$$\lim_{p \rightarrow 0} e^{p^2} = 1$$

63.0.11 Uniform and Lipschitz Continuity

Uniform Continuity:

A function $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is called *uniformly continuous* on D if:

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } \|\vec{X} - \vec{X}_0\| < \delta \Rightarrow |f(\vec{X}) - f(\vec{X}_0)| < \varepsilon$$

The key feature is that δ depends only on ε , not on the points $\vec{X}, \vec{X}_0$ themselves. If f is continuous on a compact set (closed and bounded), then f is uniformly continuous.

Lipschitz Continuity:

A function $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is *Lipschitz continuous* if there exists a constant $L > 0$ such that:

$$|f(\vec{X}) - f(\vec{Y})| \leq L \|\vec{X} - \vec{Y}\| \quad \text{for all } \vec{X}, \vec{Y} \in D$$

If $L < 1$, then f is called a *contraction*.

63.0.12 Epsilon–Delta Criterion

Let $f : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, and $\vec{X}_0 \in D$. The function f is continuous at $\vec{X}_0$ if:

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ such that } \|\vec{X} - \vec{X}_0\| < \delta \Rightarrow |f(\vec{X}) - f(\vec{X}_0)| < \varepsilon$$

This generalizes the familiar epsilon-delta criterion from single-variable calculus to multi-variable settings using norms.

63.0.13 Fixpoints

A point $\vec{X}_0 \in D \subseteq \mathbb{R}^n$ is called a *fix-point* of a function $\vec{\varphi} : D \rightarrow \mathbb{R}^n$ if:

$$\vec{\varphi}(\vec{X}_0) = \vec{X}_0$$

That is, applying the function does not change the point — it maps to itself.

63.0.14 Banach Fixed Point Theorem

Let $\vec{\varphi} : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a *contraction mapping*, i.e., there exists a constant $L < 1$ such that:

$$\|\vec{\varphi}(\vec{X}) - \vec{\varphi}(\vec{Y})\| \leq L \|\vec{X} - \vec{Y}\| \quad \text{for all } \vec{X}, \vec{Y} \in D$$

Then:

- There exists a unique fix-point $\vec{X}^* \in D$ such that $\vec{\varphi}(\vec{X}^*) = \vec{X}^*$
- Iteratively defined sequences $\vec{X}_{n+1} = \vec{\varphi}(\vec{X}_n)$ converge to the fix-point

This result is fundamental in nonlinear analysis and iterative numerical methods.

63.0.15 Vector fields

A *vector field* is a function $\vec{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Also, just a function with vectors as inputs and outputs.

$$\vec{f}(x_1, x_2, \dots, x_n) = A_1(x_1, \dots, x_n)\vec{e}_1 + \dots + A_n(x_1, \dots, x_n)\vec{e}_n$$

We call a vector field *continuous/differentiable* if all functions

$$A_1, A_2, \dots, A_n$$

are continuous/differentiable.

As a side node, the *gradient* of a multi-variable function is a vector field.

Chapter 64

Multi-variable Differentiation

64.0.1 Partial Derivatives

Let $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ and $\vec{X}_0 = (x_1^{(0)}, \dots, x_n^{(0)}) \in U$. Then f is *partially differentiable with respect to x_i* at $\vec{X}_0$ if the limit:

$$\frac{\partial f}{\partial x_i}(\vec{X}_0) := \lim_{h \rightarrow 0} \frac{f(x_1^{(0)}, \dots, x_i^{(0)} + h, \dots, x_n^{(0)}) - f(\vec{X}_0)}{h}$$

exists. This derivative measures the rate of change of f in the direction of the x_i -axis while keeping all other variables fixed.

A function is (*partially*) *differentiable* at $\vec{X}_0$ if all partial derivatives $\frac{\partial f}{\partial x_i}(\vec{X}_0)$ exist.

64.0.2 Differentiability

We call a function f differentiable at $(x_1, x_2, \dots, x_n)$ if and only if:

$$f(x_1 + \Delta x_1, x_2 + \Delta x_2, \dots, x_n + \Delta x_n) - f(x_1, x_2, \dots, x_n) =$$

$$\frac{\partial f(x_1, x_2, \dots, x_n)}{\partial x_1} \Delta x_1 + \frac{\partial f(x_1, x_2, \dots, x_n)}{\partial x_2} \Delta x_2 + \dots + E_1(\Delta x_1) + E_2(\Delta x_2) + \dots + E_n(\Delta x_n)$$

where $\lim_{\Delta x_1 \rightarrow 0} \frac{E_1(\Delta x_1)}{\Delta x_1} = \lim_{\Delta x_2 \rightarrow 0} \frac{E_2(\Delta x_2)}{\Delta x_2} = \dots = 0$

64.0.3 The Gradient

Let $f : U \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable at $\vec{X}_0 = (x_1^{(0)}, \dots, x_n^{(0)})$. The *gradient* of f at $\vec{X}_0$ is:

$$\nabla f(\vec{X}_0) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(\vec{X}_0) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\vec{X}_0) \end{bmatrix}$$

The angle of the maximal slope is $\phi = \frac{\pi}{2}$. This comes from the fact that the directional derivative which is defined as $\langle \nabla f, \vec{v} \rangle = \|\nabla f\| \|\vec{v}\| \cos(\phi)$ is orthogonal to the gradient. And for smallest slope $\phi = \frac{3\pi}{4}$. Also, the *gradient* forms a vector field.

Formally, we have a vector d^* which points in the direction of the max slope defined by

$$d^* = \arg_{\|\vec{d}\|=1} \max D_{\vec{v}} f(\vec{x})$$

where $\vec{v}$ is a unit vector.

The directional derivative written as the inner product with the gradient gives us

$$d^* = \arg_{\|d\|=1} \max \langle \nabla f, \vec{v} \rangle$$

Then using the definition of the dot product

$$d^* = \arg_{\|d\|=1} \max \|\nabla f\| \|\vec{v}\| \cos(\phi)$$

Finally, because the magnitude of $\vec{v}$ is one and to maximize cosine we set $\phi = \frac{\pi}{2}$

$$d^* = \|\nabla f\|$$

Therefore, the magnitude of the gradient is the maximal slope.

Origin of the formula

Suppose along the curve $r(t) = x_1(t)\hat{v}_1 + x_2(t)\hat{v}_2 + \dots + x_n(t)\hat{v}_n$ that $f(x_1(t), \dots, x_n(t)) = C$

$$\frac{\partial}{\partial t} f(x_1(t), \dots, x_n(t)) = \frac{\partial}{\partial t} C$$

$$\frac{\partial f}{\partial x_1} \frac{dx_1}{dt} + \dots + \frac{\partial f}{\partial x_n} \frac{dx_n}{dt} = 0$$

$$\left\langle \left(\frac{\partial f}{\partial x_1} \hat{v}_1 + \dots + \frac{\partial f}{\partial x_n} \hat{v}_n \right), \left(\frac{dx_1}{dt} \hat{v}_1 + \dots + \frac{dx_n}{dt} \hat{v}_n \right) \right\rangle = 0$$

From the formula we see that: ∇f is the normal and $\frac{d\vec{r}}{dt}$ the tangent vector of the curve

Ωℰℑ

Gradient Operations:

- $\nabla(f + g) = \nabla f + \nabla g$
- $\nabla(\lambda f) = \lambda \nabla f$ for $\lambda \in \mathbb{R}$
- $\nabla(fg) = f \nabla g + g \nabla f$

The $\frac{\text{grad } f}{\|\text{grad } f\|}$ points in the direction of the steepest increase of f and $-\frac{\text{grad } f}{\|\text{grad } f\|}$ in the lowest increase.

64.0.4 The Tangent Plane

The tangent plane to a differentiable surface $z = f(x, y)$ at the point (x_0, y_0) is the plane that best approximates the surface near that point. It is given by the linearization of f :

$$T(x, y) = f(x_0, y_0) + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

This plane touches the graph of the function at a single point and shares its slope in both x - and y -directions.

It can also be written in the parameterized form:

$$T(x, y) = \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ 0 \\ f_x(x_0, y_0) \end{bmatrix} + \mu \begin{bmatrix} 0 \\ 1 \\ f_y(x_0, y_0) \end{bmatrix}$$

Generalization

$$\begin{aligned}T(\vec{X}) &= f(\vec{x}_0) + \sum_{i=1}^n f_{x_i}(\vec{x}_0)(x_i - x_i^0) \\&= f(\vec{x}_0) + \langle \nabla f, (\vec{x} - \vec{x}_0) \rangle\end{aligned}$$

Example: $f(x, y) = 4x^2y + xy^2 + 1$

Compute partial derivatives:

$$f_x(x, y) = 8xy + y^2, \quad f_y(x, y) = 4x^2 + 2xy$$

At the point $(1, 1)$:

$$f(1, 1) = 4(1)^2(1) + 1(1)^2 + 1 = 6, \quad f_x(1, 1) = 8(1)(1) + 1 = 9, \quad f_y(1, 1) = 4(1)^2 + 2(1)(1) = 6$$

Thus, the tangent plane is:

$$T(x, y) = 6 + 9(x - 1) + 6(y - 1) \Rightarrow T(x, y) = 9x + 6y - 9$$

64.0.5 The Directional Derivative

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable at $\vec{x}_0$, and let $\vec{v} \in \mathbb{R}^n$ be a direction vector with $\|\vec{v}\| = 1$. The *directional derivative* of f at $\vec{x}_0$ in the direction $\vec{v}$ is defined as:

$$D_{\vec{v}}f(\vec{x}_0) := \lim_{h \rightarrow 0} \frac{f(\vec{x}_0 + h\vec{v}) - f(\vec{x}_0)}{h}$$

Formula:

$$D_{\vec{v}}f(\vec{x}_0) = \langle \nabla f(\vec{x}_0), \vec{v} \rangle$$

Schwarz Inequality (Cauchy-Schwarz):

$$|\langle \vec{a}, \vec{b} \rangle| \leq \|\vec{a}\| \cdot \|\vec{b}\|$$

This implies:

$$|D_{\vec{v}}f(\vec{x}_0)| \leq \|\nabla f(\vec{x}_0)\|$$

Derivation of the Directional Derivative Formula

Let $f(x, y)$ be differentiable, and let $\vec{v} = (e_1, e_2)$ be a direction vector. We aim to derive the formula for the directional derivative $D_{\vec{v}}f(x_0, y_0)$.

We begin by considering the tangent plane to the surface $z = f(x, y)$ at the point $(x_0, y_0, f(x_0, y_0))$. This plane is spanned by the vectors:

$$\vec{u}_1 = \begin{bmatrix} 1 \\ 0 \\ f_x(x_0, y_0) \end{bmatrix}, \quad \vec{u}_2 = \begin{bmatrix} 0 \\ 1 \\ f_y(x_0, y_0) \end{bmatrix}$$

These correspond to directional derivatives along the x - and y -axes, respectively.

Now consider the direction vector $\vec{v} = (e_1, e_2)$. Lift it into 3D space (onto the tangent plane) as:

$$\vec{u}_3 = \begin{bmatrix} e_1 \\ e_2 \\ D_{\vec{v}}f \end{bmatrix}$$

Since all three vectors lie in the same plane, the determinant of the matrix formed by these vectors as columns must vanish:

$$\det(\vec{u}_1, \vec{u}_2, \vec{u}_3) = 0$$

Explicitly:

$$\det \begin{bmatrix} 1 & 0 & e_1 \\ 0 & 1 & e_2 \\ f_x(x_0, y_0) & f_y(x_0, y_0) & D_{\vec{v}}f \end{bmatrix} = 0$$

Expanding the determinant gives:

$$f_x(x_0, y_0) \cdot e_2 - f_y(x_0, y_0) \cdot e_1 + D_{\vec{v}}f = 0$$

Solving for $D_{\vec{v}}f$, we obtain:

$$D_{\vec{v}}f = f_x(x_0, y_0) \cdot e_1 + f_y(x_0, y_0) \cdot e_2$$

Vector notation:

$$D_{\vec{v}}f = \langle \nabla f(x_0, y_0), \vec{v} \rangle$$

This is the desired result: the directional derivative equals the dot product of the gradient of f and the direction vector $\vec{v}$.

64.0.6 The Total Differential

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable at $\vec{x}_0$, then the *total differential* of f at $\vec{x}_0$ is the linear approximation:

$$df = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(\vec{x}_0) dx_i = \langle \nabla f(\vec{x}_0), d\vec{x} \rangle$$

This expresses how small changes in the input variables propagate into changes in the function values.

64.0.7 The Chain Rule

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be differentiable, and suppose $\vec{x} = \vec{x}(t) \in \mathbb{R}^n$ is a differentiable path. Then:

$$\frac{d}{dt}f(\vec{x}(t)) = \left\langle \nabla f(\vec{x}(t)), \frac{d\vec{x}(t)}{dt} \right\rangle$$

This is the chain rule in vector form.

Or to be more explicit, the small change in t makes a change in $x_1, x_2, \dots$ and the addition of all this changes sum up to the total change in the original function.

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x_1} \frac{\partial x_1}{\partial t} + \dots + \frac{\partial z}{\partial x_n} \frac{\partial x_n}{\partial t}$$

64.0.8 Absolute and Relative Error

Given an approximate value $\tilde{x}$ for the exact value x :

- *Absolute error:* $\Delta z_{\max} \leq |f_{x_1}| |\Delta x_1| + \dots + |f_{x_n}| |\Delta x_n|$
- *Relative error:* $\varepsilon = \frac{\Delta z_{\max}}{z}$ with dependence on the error of the inputs $\frac{\Delta x}{x}$ etc.

In multi-variable contexts, similar formulas apply using vector norms.

Absolute Error Example

We consider the function:

$$z = \sqrt{x^2 + y^2}$$

with the measured values:

$$x = 4 \text{ cm}, \quad y = 3 \text{ cm}$$

Both x and y are measured with a precision of $\Delta x = \Delta y = 0,1 \text{ cm}$. The side length calculated is $z = 5 \text{ cm}$. We now determine the maximum possible absolute error in z .

Step 1: Partial derivatives of z :

$$\frac{\partial z}{\partial x} = \frac{x}{\sqrt{x^2 + y^2}}, \quad \frac{\partial z}{\partial y} = \frac{y}{\sqrt{x^2 + y^2}}$$

Step 2: Evaluate at the given point:

$$\frac{\partial z}{\partial x}(4, 3) = \frac{4}{5}, \quad \frac{\partial z}{\partial y}(4, 3) = \frac{3}{5}$$

Step 3: Use the absolute error formula:

$$\begin{aligned} \Delta z &\leq \left| \frac{\partial z}{\partial x} \right| \cdot \Delta x + \left| \frac{\partial z}{\partial y} \right| \cdot \Delta y \\ \Delta z &\leq \frac{4}{5} \cdot 0,1 + \frac{3}{5} \cdot 0,1 = \frac{7}{5} \cdot 0,1 = 0,14 \text{ cm} \end{aligned}$$

Conclusion:

The result for z is accurate to within 0,14cm.

Relative Error Example

The relative error of a function $z = f(x_1, \dots, x_n)$ is defined by:

$$\varepsilon = \frac{\Delta z_{\max}}{z}$$

Assuming z depends on variables with known relative measurement errors, and the function is differentiable, we estimate:

$$\varepsilon = \frac{\left| \frac{\partial f}{\partial x_1} \right| \cdot \Delta x_1 + \dots + \left| \frac{\partial f}{\partial x_n} \right| \cdot \Delta x_n}{f(x_1, \dots, x_n)}$$

or more directly using relative errors:

$$\varepsilon \approx \left| \frac{\Delta x_1}{x_1} \right| + \dots + \left| \frac{\Delta x_n}{x_n} \right|$$

Example: Volume of a Cuboid

Let:

$$z = f(a, b, c) = a \cdot b \cdot c$$

Assume the side lengths $a = b = c$ are measured with a uniform relative error $\varepsilon_{\text{in}} = \frac{\Delta a}{a} = \frac{\Delta b}{b} = \frac{\Delta c}{c} = \varepsilon$

Then the partial derivatives are:

$$\frac{\partial z}{\partial a} = bc, \quad \frac{\partial z}{\partial b} = ac, \quad \frac{\partial z}{\partial c} = ab$$

Applying the relative error formula:

$$\frac{\Delta z_{\text{max}}}{z} \leq \frac{bc \cdot \Delta a + ac \cdot \Delta b + ab \cdot \Delta c}{abc} = \frac{\Delta a}{a} + \frac{\Delta b}{b} + \frac{\Delta c}{c} = 3\varepsilon$$

Target accuracy:

If the final result z must be accurate within $\varepsilon_z = 0,01$ (i.e., 1), then we must satisfy:

$$3\varepsilon \leq 0,01 \Rightarrow \varepsilon \leq \frac{0,01}{3} = 0,0033 \quad (\text{or } 0,33\%)$$

Conclusion:

Each input measurement must be made with a maximum relative error of 0,33% to ensure the output $z = abc$ is accurate to within 1 percent.

64.0.9 Implicit Differentiation

If a function $F(x, y) = 0$ defines y implicitly as a function of x , and F is differentiable, then:

$$\frac{dz}{dx} = -\frac{F_x}{F_z}$$

This generalizes to higher dimensions using the total differential and the implicit function theorem.

Example:

Given is $x^2 + y^2 + z^2 = 1$ differentiate with respect to x . See z or the target function as function in terms of the other variables. In this case $z(x, y)$.

$$\frac{d}{dx}x^2 + y^2 + z^2 - 1 = 0$$

$$2x + 2z \frac{dz}{dx} = 0$$

$$\frac{dz}{dx} = -\frac{x}{z}$$

Example:

Given the implicitly defined function:

$$yx^2 + \frac{x}{y} + x + y - 4 = 0$$

We are asked to compute the derivative $y'(1)$, that is, the derivative of y with respect to y at the point $(1, 1)$.

Step 1: Differentiate implicitly

Differentiate both sides of the equation with respect to x , treating $y = y(x)$ as a function of x :

$$\begin{aligned}\frac{d}{dx}(yx^2) + \frac{d}{dx}\left(\frac{x}{y}\right) + \frac{d}{dx}(x) + \frac{d}{dx}(y) - \frac{d}{dx}(4) &= 0 \\ y'x^2 + y \cdot 2x + \frac{y - xy'}{y^2} + 1 + y' &= 0\end{aligned}$$

Collect terms involving y' . Group all y' terms on one side:

$$y'x^2 - \frac{xy'}{y^2} + y' + 2xy + \frac{1}{y} + 1 = 0$$

Factor y' :

$$y' \left(x^2 - \frac{x}{y^2} + 1 \right) = - \left(2xy + \frac{1}{y} + 1 \right)$$

Evaluate at $x = 1, y = 1$

Substitute $x = 1, y = 1$:

$$\begin{aligned}y'(1 - 1 + 1) &= -(2 + 1 + 1) \\ y'(1) &= -4 \\ y' &= -4\end{aligned}$$

Final Answer

$$y'(1) = -4$$

64.0.10 Schwarz's Theorem (Clairaut's Theorem)

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be twice continuously differentiable. Then for any $i \neq j$, the mixed partial derivatives satisfy:

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$$

That is, the order of partial differentiation does not matter if all second derivatives are continuous.

64.0.11 The Jacobian

Let $\vec{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$, where

$$\vec{f}(\vec{x}) = \begin{bmatrix} f_1(x_1, \dots, x_n) \\ \vdots \\ f_m(x_1, \dots, x_n) \end{bmatrix}$$

The *Jacobian matrix* of $\vec{f}$ is the $m \times n$ matrix:

$$J_{\vec{f}}(\vec{x}) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

In the case $n = m$, the *Jacobian determinant* $\det J_{\vec{f}}(\vec{x})$ describes local invertibility and orientation.

64.0.12 Taylor's Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be k -times continuously differentiable at $\vec{x}_0$. Then the Taylor polynomial of degree 2 around $\vec{x}_0$ is:

$$f(\vec{x}) \approx f(\vec{x}_0) + \langle \nabla f(\vec{x}_0), \vec{x} - \vec{x}_0 \rangle + \frac{1}{2}(\vec{x} - \vec{x}_0)^T H_f(\vec{x}_0)(\vec{x} - \vec{x}_0) + \cdots,$$

where $H_f(\vec{x}_0)$ is the *Hessian matrix* of second partial derivatives.

Example:

Let $f(x, y) = e^y \sin(x + 2y)$. We expand around the point $(a, b) = (0, 0)$.

First, compute the necessary derivatives at $(0, 0)$:

$$\begin{aligned} f(0, 0) &= \sin(0) = 0 \\ f_x &= e^y \cos(x + 2y) \Rightarrow f_x(0, 0) = 1 \\ f_y &= 2ye^y \cos(x + 2y) + e^y \sin(x + 2y) \Rightarrow f_y(0, 0) = 0 + 2 = 2 \\ f_{xx} &= -e^y \sin(x + 2y) \Rightarrow f_{xx}(0, 0) = 0 \\ f_{xy} &= f_{yx} = 2e^y \sin(x + 2y) + e^y \cos(x + 2y) \Rightarrow f_{xy}(0, 0) = 3 \\ f_{yy} &= -3e^y \sin(x + 2y) + 4e^y \cos(x + 2y) \Rightarrow f_{yy}(0, 0) = 0 + 4 = 4 \end{aligned}$$

Now, the second order Taylor polynomial at $(0, 0)$ is:

$$\begin{aligned} f(x, y) &\approx 0 + 1 \cdot x + 2 \cdot y \\ &\quad + \frac{1}{2} \cdot 0 \cdot x^2 + 3 \cdot xy + \frac{1}{2} \cdot 4 \cdot y^2 \\ &= x + 2y + 3xy + 2y^2 \end{aligned}$$

64.0.13 Relative Extrema

To determine local extrema for $f : \mathbb{R}^n \rightarrow \mathbb{R}$, follow this procedure:

Step 1: Find critical points by solving:

$$\nabla f(\vec{x}) = 0$$

Step 2: Compute the Hessian matrix:

$$H_f(\vec{x}) = \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)$$

Step 3: Analyze the Hessian at critical points:

- If H is positive definite $f_{xx}(x_0, y_0) > 0, \det(H) > 0 \implies$ local minimum
- If H is negative definite $f_{xx}(x_0, y_0) < 0, \det(H) > 0 \implies$ local maximum
- If H has mixed signs (indefinite) $\det(H) < 0 \implies$ saddle point
- If $\det(H) = 0 \implies$ Next derivative decides

The determinant is needed as part of the test because, while the second derivative tells us about the concavity, the determinant helps us determine if all the concavities given by the axis second derivatives with respect to $x_1, x_2, \dots, x_n$ and the mixed partials are up, down or neither. If the signs are mixed then it is a saddle.

Example: $f(x, y) = x^2 + y^2$

$$\nabla f = \begin{bmatrix} 2x \\ 2y \end{bmatrix} \Rightarrow \text{Critical point at } (0, 0)$$

$$H_f = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \Rightarrow \text{positive definite} \Rightarrow (0, 0) \text{ is a local minimum}$$

64.0.14 Finding Candidate Points for Zeros While Seeking Relative Extrema

To find local minima or maxima of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we first identify *critical points*, which are the candidates for relative extrema. These are the points where the gradient of f vanishes or does not exist.

Step 1: Compute the Gradient

The gradient of f , denoted ∇f , collects all the first partial derivatives:

$$\nabla f(x_1, \dots, x_n) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{bmatrix}$$

Step 2: Solve the System $\nabla f = 0$

Find all points $\vec{x}_0 \in \mathbb{R}^n$ such that:

$$\nabla f(\vec{x}_0) = \vec{0}$$

This is a nonlinear system of n equations in n variables. The solutions $\vec{x}_0$ are the *candidate points for extrema*, and also referred to as the **stationary points** or *critical points* of the function.

You use the following techniques to find the points:

- Factorization
- Addition of rows
- Substitution

There are more but which one to use depends on the kind of problem.

Step 3: Analyze the Nature of the Critical Points

Once all critical points $\vec{x}_0$ satisfying $\nabla f(\vec{x}_0) = 0$ are found, analyze each using the *second derivative test* involving the Hessian matrix H_f :

$$H_f(\vec{x}) = \left(\frac{\partial^2 f}{\partial x_i \partial x_j} \right)$$

Evaluate H_f at each critical point and use the eigenvalues or definiteness to classify:

- If $f_{xx}(x_0, y_0) > 0$ and $D > 0$, then f has a *local minimum* at (x_0, y_0) .
- If $f_{xx}(x_0, y_0) < 0$ and $D > 0$, then f has a *local maximum* at (x_0, y_0) .
- If $D < 0$, then (x_0, y_0) is a *saddle point*.
- If $D = 0$, the *second derivative test is inconclusive*; higher-order derivatives must be considered.

Note: Non-differentiable Points

If the function f is not differentiable at some point $\vec{x}_0$, but defined there, this point may also be a candidate for extremum and should be examined separately using limit-based analysis or directional behavior.

64.0.15 Lagrange Multipliers

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be the function to optimize, subject to the constraint $g(\vec{x}) = 0$, where $g : \mathbb{R}^n \rightarrow \mathbb{R}$.

If we look at the constraint intersection with the original function we are going to notice that along the intersection there are maxima and minima with their respective tangent lines and gradients that are normal to them. We also are going to see that both the gradient of f and the gradient of g are just scalar version of each other so:

This leads to the *Lagrange system*:

$$\begin{cases} \nabla f(\vec{x}) = \lambda \nabla g(\vec{x}) \\ g(\vec{x}) = 0 \end{cases}$$

For more conditions we can use $g_1, \dots, g_n$ and $\lambda_1, \dots, \lambda_n$

$$\begin{cases} \nabla f(\vec{x}) = \lambda_1 \nabla g_1(\vec{x}) + \dots + \lambda_n \nabla g_n(\vec{x}) \\ g_1(\vec{x}) = 0 \\ \vdots \\ g_n(\vec{x}) = 0 \end{cases}$$

We can also compact these equations in the *Lagrangian*

$$\mathcal{L}(x, y, \dots, \lambda_1, \dots, \lambda_n) = f(x, y, \dots) + \lambda_1 g_1(x, y, \dots) + \dots + \lambda_n g_n(x, y, \dots)$$

Determinant method

Sometimes it is a good way to solve a problem of this topic is to:

1. Build the *Lagrangian*
2. $\nabla L = 0$ and maybe compute the determinant to generate another equation to solve
3. Get rid of λ via factorization, substitution, addition of rows, etc.
4. Solve the system without λ

The Bordered Hessian

The Bordered Hessian matrix is:

$$H_B = \begin{bmatrix} 0 & (\nabla g(x_1, \dots, x_n))^T \\ \nabla g(x_1, \dots, x_n) & \nabla^2 \mathcal{L}(x, \dots, x_n, \lambda_1, \dots, \lambda_n) \end{bmatrix}$$

Where:

- $\nabla g(x_1, \dots, x_n)$ is the $m \times n$ Jacobian matrix of the constraints.
- $\nabla^2 \mathcal{L}(x, \dots, x_n, \lambda_1, \dots, \lambda_n)$ is the $n \times n$ Hessian of the Lagrangian with respect to x .

Example I:

Find the point on the circle $x^2 + y^2 = 4$ that is the closest to the point $(3, 4)$.

The function $f(x, y)$ is this case the euclidean norm:

$$f(x, y) = \sqrt{(x-3)^2 + (y-4)^2}$$

subject to the constraint:

$$g(x, y) = x^2 + y^2 - 4 = 0$$

The gradient of the constraint is:

$$\nabla g = \begin{bmatrix} 2x \\ 2y \end{bmatrix}$$

Since $\nabla g \neq 0$, the rank condition is fulfilled, and no further critical points need to be analyzed.

Step 1: Build the Lagrangian

$$\mathcal{L}(x, y, \lambda) = f(x, y) + \lambda g(x, y)$$

In this case we can use a property of the root the function to our advantage. Notice that the minimum of the root function is the minimum of the term under root therefore, we can simplify some work by writing $f(x, y)$ without the root.

$$\mathcal{L}(x, y, \lambda) = (x-3)^2 + (y-4)^2 + \lambda(x^2 + y^2 - 4)$$

Step 2: Compute the necessary conditions

$$\begin{aligned} \text{I} \quad \frac{\partial \mathcal{L}}{\partial x} &= 2(x-3) + 2\lambda x = 0 \\ \text{II} \quad \frac{\partial \mathcal{L}}{\partial y} &= 2(y-4) + 2\lambda y = 0 \\ \text{III} \quad \frac{\partial \mathcal{L}}{\partial \lambda} &= x^2 + y^2 - 4 = 0 \end{aligned}$$

Now, this is the difficult part. The best approach in this case is to use substitution to with a top-bottom approach:

$$x = \frac{3}{1+\lambda}$$

$$y = \frac{4}{1+\lambda}$$

Solving for λ is also an option but in these kinds of systems of equations where both the partial with respect to x and y are very similar it is a good idea to solve for them because they tend to have a similar form when solving for them.

$$\begin{aligned}
\left(\frac{3}{1+\lambda}\right)^2 + \left(\frac{4}{1+\lambda}\right)^2 - 4 &= 0 \\
\frac{9}{(1+\lambda)^2} + \frac{16}{(1+\lambda)^2} - 4 &= 0 \\
\frac{25}{(1+\lambda)^2} &= 4 \\
25 &= 4(1+\lambda)^2 \\
\sqrt{\frac{25}{4}} &= 1+\lambda \\
\lambda &= -1 \pm \frac{5}{2}
\end{aligned}$$

$$\lambda_1 = \frac{-7}{2} \quad \lambda_2 = \frac{3}{2}$$

Now, let us substitute backwards for λ_1 :

$$x = \frac{3}{1 - \frac{7}{2}} = \frac{6}{-5}$$

$$y = \frac{4}{1 - \frac{7}{2}} = \frac{8}{-5}$$

Now, for λ_2 :

$$x = \frac{3}{1 + \frac{3}{2}} = \frac{6}{5}$$

$$y = \frac{4}{1 + \frac{3}{2}} = \frac{8}{5}$$

Step 3: Hessian Test with Constraint

We construct the Hessian:

$$H = \begin{bmatrix} 2+2\lambda & 0 & 2x \\ 0 & 2+2\lambda & 2 \\ 2x & 2 & 0 \end{bmatrix}$$

For λ_1 we get $\det(H) = \frac{244}{5} > 0$, a minimum and for λ_2 $\det(H) = \frac{-244}{5}$ a maximum.

Example II:

Optimize the distance to the origin $f(x, y, z)$ with respect to the constraints $g_1(x, y, z) = z^2 - x^2 - y^2 = 0$ and $g_2(x, y, z) = x - 2z - 3 = 0$

$$\begin{aligned}
\nabla f &= \lambda \nabla g_1 + \mu \nabla g_2 \\
g_1 &= 0 \\
g_2 &= 0
\end{aligned}$$

We know that $f(x, y, z) = \sqrt{x^2 + y^2 + z^2}$. Like with other minimization problems we can just find the minimum of the part under root thus, $f(x, y, z) = x^2 + y^2 + z^2$. Now we can put the build our system of equations

$$\begin{aligned}
\text{I } 2x &= \lambda(-2x) + \mu \\
\text{II } 2y &= \lambda(-2y) \\
\text{III } 2z &= \lambda(2z) + \mu(-2) \\
\text{IV } z^2 &= x^2 + y^2 \\
\text{V } x &= 2z + 3
\end{aligned}$$

For the equation **II** we can already see that the equation is equal to zero only if $\lambda = -1$ or $y = 0$. Thus, we have got our first numbers to work with.

Case $\lambda = -1$:

$$\begin{aligned}
\text{I } 2x &= -1(-2x) + \mu \implies \mu = 0 \\
\text{III } 2z &= -1(2z) + 0(-2) \implies z = 0 \\
\text{IV } 0^2 &= x^2 + y^2 \implies x = y = 0 \\
\text{V } 0 &= 0 + 3 \implies 0 = 3
\end{aligned}$$

With the last contradiction we know that the whole case $\lambda = -1$ is not valid.

Case $y = 0$:

$$\begin{aligned}
\text{IV } z^2 &= x^2 + y^2 \implies z^2 = x^2 \implies z = \pm x \\
\text{V } x &= 2x + 3 \implies x = -3(z = x) \\
\text{V } x &= -2x + 3 \implies x = 1(z = -x)
\end{aligned}$$

Now with this new information we can get two different points $P_1(1, 0, -1)$ and $P_2(-3, 0, -3)$. We do not need to write the Hessian in this case because with geometric intuition the problem can be solved.

$$f(P_1) = 1^2 + 0^2 + (-1)^2 = 2$$

So the distance is $\sqrt{2}$.

$$f(P_2) = -3^2 + 0^2 - 3^2 = 18$$

Thus, the distance is $3\sqrt{2}$.

Our final Answer is that P_1 is the closest point to the intersection of the two planes with the function f .

64.0.16 The Tangent Vector

Let $\vec{X}(t) \in \mathbb{R}^n$ be a differentiable, parameterized curve. The *tangent vector* at the point $\vec{X}(t_0)$ is given by:

$$\vec{X}'(t_0) = \begin{bmatrix} x'_1(t_0) \\ x'_2(t_0) \\ \vdots \\ x'_n(t_0) \end{bmatrix}$$

This vector points in the direction in which the curve is moving at t_0 , and its magnitude corresponds to the instantaneous speed. The unit tangent vector is:

$$\vec{X}(t) = \frac{\vec{X}'(t)}{\|\vec{X}'(t)\|}$$

The speed of the vector in $\mathbb{R}^2$ is given by:

$$\vec{X}(t) = \begin{bmatrix} t \\ f(t) \end{bmatrix}$$

and the tangent is:

$$T(t) = \begin{bmatrix} t \\ f(t) \end{bmatrix} + \lambda \begin{bmatrix} 1 \\ f'(t) \end{bmatrix}$$

Derivation of the Tangent Vector Formula

Given a position vector function $\vec{r}(t)$, the tangent vector to the curve described by $\vec{r}(t)$ is obtained by taking the derivative of $\vec{r}(t)$ with respect to time. This derivative is defined as the following limit:

$$\frac{d\vec{r}}{dt} = \lim_{\Delta t \rightarrow 0} \frac{\vec{r}(t + \Delta t) - \vec{r}(t)}{\Delta t}$$

Assuming that $\vec{r}(t)$ is expressed in terms of the canonical basis vectors $\hat{i}, \hat{j}, \hat{k}$, we can write:

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$$

Then, the increment becomes:

$$\vec{r}(t + \Delta t) = x(t + \Delta t)\hat{i} + y(t + \Delta t)\hat{j} + z(t + \Delta t)\hat{k}$$

So the difference in the numerator of the derivative is:

$$\begin{aligned} \vec{r}(t + \Delta t) - \vec{r}(t) &= [x(t + \Delta t) - x(t)]\hat{i} \\ &\quad + [y(t + \Delta t) - y(t)]\hat{j} \\ &\quad + [z(t + \Delta t) - z(t)]\hat{k} \end{aligned}$$

Dividing by Δt and taking the limit:

$$\begin{aligned} \frac{d\vec{r}}{dt} &= \lim_{\Delta t \rightarrow 0} \frac{\vec{r}(t + \Delta t) - \vec{r}(t)}{\Delta t} \\ &= \left(\lim_{\Delta t \rightarrow 0} \frac{x(t + \Delta t) - x(t)}{\Delta t} \right) \hat{i} \\ &\quad + \left(\lim_{\Delta t \rightarrow 0} \frac{y(t + \Delta t) - y(t)}{\Delta t} \right) \hat{j} \\ &\quad + \left(\lim_{\Delta t \rightarrow 0} \frac{z(t + \Delta t) - z(t)}{\Delta t} \right) \hat{k} \\ &= \frac{dx}{dt} \hat{i} + \frac{dy}{dt} \hat{j} + \frac{dz}{dt} \hat{k} \end{aligned}$$

This vector $\frac{d\vec{r}}{dt}$ points in the direction of motion and is tangent to the curve at each point t .

64.0.17 Linear Regression

Linear Regression is can be done using the gradient and the data points of a data set, instead of using matrices

Steps:

The dimension used for this case is 2 for simplicity

1. *Construct the Square error function: $SE(\alpha, \beta) = \sum_{i=0}^n (\alpha x_i + \beta - y_i)^2$*
2. *Take the gradient and set it to zero:*

$$\begin{bmatrix} \sum_{i=0}^n 2x_i(\alpha x_i + \beta - y_i) \\ 2 \sum_{i=0}^n \alpha x_i + \beta y_i \end{bmatrix} = 0$$

3. *Solve for Alpha and Beta:*

$$\alpha = \frac{\sum_i x_i y_i - \sum_i x_i \sum_i y_i}{\sum_i x_i^2 - (\sum_i x_i)^2}$$

$$\beta = \frac{1}{n} \sum_i y_i - \alpha \frac{1}{n} \sum_i x_i$$

Chapter 65

Multidimensional Integrals

65.0.1 Arclength of a Curve

We will use the famous technique of approximating the length/area/volume of something by dividing it in to equally smaller section that are straight lines in this case. In other cases they will be prisms, squares, etc. Now if we look at the difference between two points in our 3D space, for this Example we notice that there are three differences Δi Δj Δk . Now let us take the length of the vector $\sqrt{(\Delta i)^2 + (\Delta j)^2 + (\Delta k)^2}$. The total Arclength is going to be the sum of all of these sectors

$$L = \sum_{i=1}^n \sqrt{(\Delta i)^2 + (\Delta j)^2 + (\Delta k)^2}$$

if we use a clever one $\left(\frac{\Delta t}{\Delta t}\right)^2$ we get

$$L = \sum_{i=1}^n \sqrt{\left(\frac{\Delta i}{\Delta t}\right)^2 + \left(\frac{\Delta j}{\Delta t}\right)^2 + \left(\frac{\Delta k}{\Delta t}\right)^2} \Delta t$$

Now if we take the limit we get

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{\left(\frac{\Delta i}{\Delta t}\right)^2 + \left(\frac{\Delta j}{\Delta t}\right)^2 + \left(\frac{\Delta k}{\Delta t}\right)^2} \Delta t$$

which is

$$L = \int_a^b \sqrt{\left(\frac{\Delta i}{\Delta t}\right)^2 + \left(\frac{\Delta j}{\Delta t}\right)^2 + \left(\frac{\Delta k}{\Delta t}\right)^2} dt$$

$$L = \int_a^b \sqrt{(f'(t))^2 + (g'(t))^2 + (h'(t))^2} dt$$

For differentiable function on the interval $t \in [a; b]$

65.0.2 Line Integrals

Over Vector Field

Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a *vector field*, and let γ be a smooth curve parametrized by $\vec{r}(t)$, $t \in [a, b]$. The *line integral* of F along γ is:

$$\int_{\gamma} \left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle dt = \int_a^b \left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}}{dt} \right\rangle dt = \vec{F}(\vec{r}(b)) - \vec{F}(\vec{r}(a))$$

In the context of physics this integral measures the work done by the field $\vec{F}$ along the path γ , such as force over distance.

In compact form

$$\int_{\gamma} \langle \vec{F}, d\vec{r} \rangle$$

Example: Line Integral over a Vector Field

Let $f(x, y) = 2x + 2y$ be our vector field.

Let the curve c be the quarter-circle of radius 1 from $(1, 0)$ to $(0, 1)$, parametrized by:

$$\vec{r}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \cos(t) \\ \sin(t) \end{bmatrix} \quad t \in \left[0, \frac{\pi}{2}\right]$$

We compute the line integral of $\vec{F}$ along c :

$$\int_C \left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle dt = \int_0^{\frac{\pi}{2}} \left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle dt$$

First, compute $\vec{F}(\vec{r}(t))$:

$$\vec{F}(\vec{r}(t)) = \begin{bmatrix} 2 \cos(t) \\ 2 \sin(t) \end{bmatrix}$$

Then the tangent vector of $\vec{r}(t)$:

$$\frac{d\vec{r}(t)}{dt} = \begin{bmatrix} -\sin(t) \\ \cos(t) \end{bmatrix}$$

Now compute the inner product:

$$\left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle = -2 \cos(t) \sin(t) + \cos(t) 2 \sin(t) = 0$$

Therefore, the line integral is:

$$\begin{aligned} & \int_C \left\langle \vec{F}, \frac{d\vec{r}}{dt} \right\rangle dt \\ &= \int_0^{\frac{\pi}{2}} 0 dt \\ &= 0 \end{aligned}$$

The line integral of the gradient vector field $f(x, y) = 2x + 2y$ over this curve is zero.


Over Scalar field

Let $f(x_1, x_2, \dots, x_n)$ be a scalar field defined on a smooth curve c , which is parametrized by a vector function.

$$\vec{r}(t) = (x_1(t), x_2(t), \dots, x_n(t)), \quad t \in [a, b]$$

Then, the line integral of f over c is given by:

$$\int_C f(x_1, x_2, \dots, x_n) ds = \int_a^b f(\vec{r}(t)) \left\| \frac{\vec{r}(t)}{dt} \right\| dt$$



This formula comes from approximating the area of the squares under the curve with height $f(x_k, y_k)$ and the arclength Δs_k , we will simplify to three dimensions for the sake of simplicity, but like seen before this works for n dimensions. $A_k = f(x_k, y_k) \Delta s_k$, where $\Delta s_k = \sqrt{(\Delta x_k)^2 + (\Delta y_k)^2}$. Now by using our formula for the arclength we already know, we get $\left\| \frac{\vec{r}(t)}{dt} \right\| dt = ds$.

Example: Scalar Field

Compute the line integral of $f(x, y) = x^2 + y^2$ over the straight line segment going from $(1, 1)$ to $(3, 3)$

This time our curve is $y = x$ so the parametrized curve is $\begin{cases} x(t) = t \\ y(t) = t \end{cases}$ this left us with:

$$\int_1^3 2t^2 \sqrt{2} dt = \frac{52\sqrt{2}}{3}$$

65.0.3 Line Integrals with respect to a specific variable

We are going to take another look at line integrals but from the perspective of just one variable and how this affects the formulas we know. For the next explanation we will use only functions of two variables to keep the examples simple.

Over Scalar Field with respect to x or y

Imagining how this works is not specially difficult. Imagine taking a line integral where we do not move along the y -axis and only care about the plane xz . Here our line integral only changes in the fact that in the norm of our curve the component that was responsible for y will be set 0 and the ds changes to the variable we are integrating with respect to. In this case dx .

$$\int_c f(x, y) dx = \int_c f(g(t), h(t)) \sqrt{g'(t)^2} dt$$

Over Vector Field with respect to x or y

In the case of a field the story is different in the fact that now our output vectors change depending on the variable we are choosing to left out.

Now, given a field $\vec{F}(x, y) = M(x, y)\mathbf{i} + N(x, y)\mathbf{j}$ and the curve $\vec{r}(t) = g(t)\mathbf{i} + h(t)\mathbf{j}$. The line integral of $\vec{F}$ along C is

$$\int_{\gamma} \left\langle \nabla \vec{F}, \frac{\vec{r}(t)}{dt} \right\rangle dt$$
$$\int_a^b M(x, y)g'(t)dt + \int_a^b N(x, y)h'(t)dt$$

Here $g'(t)dt = dx$ and $h'(t)dt = dy$

$$\int_a^b Mdx + \int_a^b Ndy$$

These two integrals give us both the integral with respect to x and y .

Example:

Curve C is a portion of the parabola $y = x^2$ that goes through the points $(1,1)$ to $(2,4)$

Compute: $\int_a^b \frac{x}{y} dy$

Step 1: Parameterize C

The trick for this cases where the curve is the graph of a function is to set $x = t$ and then solve for y in terms of x .

In this case we get

$$\vec{r}(t) = t\mathbf{i} + t^2\mathbf{j} \text{ with } 1 \leq t \leq 2$$

Step 2: Substitute into the integral

$$h'(t)dt = 2tdt$$

$$\int_a^b \frac{t}{t^2} 2tdt = \int_1^2 \frac{1}{t} 2tdt = 2$$

65.0.4 Flow and Circulation

For velocity $\vec{F} = M(x, y)\mathbf{i} + N(x, y)\mathbf{j}$ and a smooth curve C parameterized by $\vec{r}(t)$

$$Flow = \int_C \langle \vec{F}, \vec{T} \rangle ds$$

If the curve C is closed, then it is called *Circulation*.

$$Circulation = \oint_C \langle \vec{F}, \vec{T} \rangle ds$$

Flow is a measurement of the degree of which a curve goes along the vector field.

Circulation Density

The *circulation density* of a vector field $\vec{F}(x, y) = M(x, y)\hat{i} + N(x, y)\hat{j}$ is given by

$$CD = \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right)$$

It derives from imagining having a square as our curve where from our starting point (x, y) we move $(x + \Delta x, y)$, $(x + \Delta x, y + \Delta y)$, $(x, y + \Delta y)$ generating four different curves. Therefore our total circulation is given by

$$\oint_C \langle \vec{F}, d\vec{r} \rangle = \sum_{i=1}^4 \int_{C_i} \langle \vec{F}, \vec{T} \rangle d\vec{r}$$

The dot products are given by

- Bottom: $\langle \vec{F}(x, y) \Delta x \hat{i} \rangle = M(x, y) \Delta x$
- Right: $\langle \vec{F}(x + \Delta x, y) \Delta y \hat{j} \rangle = N(x + \Delta x, y) \Delta y$
- Top: $\langle \vec{F}(x, y + \Delta y) - \Delta x \hat{i} \rangle = -M(x, y + \Delta y) \Delta x$
- Left: $\langle \vec{F}(x, y) - \Delta y \hat{j} \rangle = N(x, y) \Delta y$

Then the Top minus Bottom is $-(M(x, y + \Delta y) - M(x, y) \Delta x)$ and notice that inside the parenthesis we have something similar to the partial

$$\frac{\partial M}{\partial y} = \lim_{\Delta y \rightarrow 0} \frac{M(x, y + \Delta y) - M(x, y)}{\Delta y}$$

Thus, by multiplying a clever one $\frac{\Delta y}{\Delta y}$ we get $-\frac{\partial M}{\partial y} \Delta y \Delta x$. Similarly, we get for the Right minus left case $\frac{\partial N}{\partial x} \Delta y \Delta x$

Finally by putting all together

$$\oint_C \langle \vec{F}, d\vec{r} \rangle = -\frac{\partial M}{\partial y} \Delta y \Delta x + \frac{\partial N}{\partial x} \Delta y \Delta x = \left(-\frac{\partial M}{\partial y} + \frac{\partial N}{\partial x} \right) \Delta y \Delta x$$

$\Delta y \Delta x$ is the area of the rectangle.

65.0.5 Green's Theorem Circulation-Curl Form

For C , a smooth, simple, closed curve enclosing R , $\vec{F} = M\hat{i} + N\hat{j}$ having M, N continuous first partials in an open region containing R

$$\oint_C \langle \vec{F}, d\vec{r} \rangle = \int \int_R \left(-\frac{\partial M}{\partial y} + \frac{\partial N}{\partial x} \right) dx dy$$

65.0.6 Flux

For a continuous field $\vec{F} = M(x, y)\hat{i} + N(x, y)\hat{j}$, a smooth curve with no intersections C and $\vec{n}$ the outward normal

$$Flux = \oint_C \langle \vec{F}, \vec{n} \rangle ds$$

The derivation of the formula for actually computing the *flux* goes as follows: First, we find ourselves is 2 dimensions and we want to find a normal vector to the curve. We can pretend that we are in 3 dimensions and define a vector

$$\vec{k} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

This gives us $\vec{n} = \vec{T} \times \vec{k}$. Now either by using the determinant trick or formally computing the cross product we get

$$\vec{T} \times \vec{k} = \vec{n} = \begin{bmatrix} \frac{dy}{ds} \\ \frac{dx}{ds} \\ 0 \end{bmatrix}$$

The final formula is then

$$Flux = \oint_C \langle \vec{F}, \vec{n} \rangle ds = \oint_C \left\langle \vec{F}, \begin{bmatrix} \frac{dy}{ds} \\ \frac{dx}{ds} \\ 0 \end{bmatrix} \right\rangle ds = \int_a^b M dy - \int_a^b N dx$$

One extra note about the derivation is that if the curve goes counter clockwise we use $\vec{n} = \vec{T} \times \vec{k}$ but if it goes clockwise $\vec{n} = \vec{k} \times \vec{T}$ because of the right-hand-rule.

65.0.7 Flux Density

Just like for the *Circulation Density* we will use the small square to examine the divergence in a small region. Following a similar process with the formula for flux we can get a similar result.

$$FD = \left(-\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} \right)$$

65.0.8 Green's Theorem Divergence-Flux Form

For C , a smooth, simple, closed curve enclosing R , $\vec{F} = M\hat{i} + N\hat{j}$ having M, N continuous first partials in an open region containing R

$$\oint_C \langle \vec{F}, \vec{n} \rangle ds = \iint_R \left(\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} \right) dxdy$$

65.0.9 Conservative Vectors Fields

A field $\vec{F}$ is *conservative* on an open domain if

$$\int_C \langle \vec{F}, d\vec{r} \rangle$$

is the same for all paths C between points A and B in the domain. In other words it only depends on the endpoints.

65.0.10 The Potential Function

A vector field $\vec{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called *conservative* if there exists a scalar function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ such that:

$$\vec{F} = \nabla f$$

The function f is then called a *potential function* of $\vec{F}$ and also

$$\int_C \langle \vec{F}, d\vec{r} \rangle = f(B) - f(A)$$

Proof:

$$\int_C \langle \vec{F}, d\vec{r} \rangle = \int_a^b \left\langle \vec{F}(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle dt$$

By assumption

$$= \int_a^b \left\langle \nabla f(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle dt$$

This is like the chain rule. Therefore,

$$\begin{aligned} \left\langle \nabla f(\vec{r}(t)), \frac{d\vec{r}(t)}{dt} \right\rangle &= \frac{\partial f}{\partial x_1} \frac{\partial x_1}{\partial t} + \cdots + \frac{\partial f}{\partial x_n} \frac{\partial x_n}{\partial t} = \frac{d}{dt} f(\vec{r}(t)) \\ &= \int_a^b \frac{d}{dt} \phi(\vec{r}(t)) dt \end{aligned}$$

By the fundamental theorem of calculus

$$= f(B) - f(A)$$

□

Check for conservative vector fields

Using the definition $\vec{F} = \nabla f$ and by writing

$$\begin{aligned} \vec{F} &= f_1(x_1, x_2, \dots) \hat{i}_1 + f_2(x_1, x_2, \dots) \hat{i}_2 + \cdots + f_n(x_1, x_2, \dots) \hat{i}_n \\ &= \frac{\partial f}{\partial x_1} \hat{i}_1 + \frac{\partial f}{\partial x_2} \hat{i}_2 + \cdots + \frac{\partial f}{\partial x_n} \hat{i}_n \end{aligned}$$

We get the following and reducing to 3 variables and dimensions for the sake of simplicity

$$\frac{\partial f_1}{\partial y} = \frac{\partial}{\partial y} \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} \frac{\partial f}{\partial y} = \frac{\partial f_2}{\partial x}$$

We can also keep making pairs like this, but checking one is enough.

Finding the Potential Function of a Vector Field

To find the potential function $f(x, y)$ of a vector field $\vec{F}(x, y) = (P(x, y), Q(x, y))$, we must check if the field is conservative and then integrate accordingly. A vector field is conservative if there exists a scalar potential function f such that:

$$\vec{F} = \nabla f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix}$$

Given:

$$\vec{F}(x, y) = \begin{bmatrix} 2xy + e^x \\ x^2 + \frac{1}{2\sqrt{y}} \end{bmatrix}$$

Let $P(x, y) = 2xy + e^x$, and $Q(x, y) = x^2 + \frac{1}{2\sqrt{y}}$.

First, check if the field is conservative by verifying:

$$\frac{\partial P}{\partial y} = \frac{\partial}{\partial y}(2xy + e^x) = 2x$$

$$\frac{\partial Q}{\partial x} = \frac{\partial}{\partial x}\left(x^2 + \frac{1}{2\sqrt{y}}\right) = 2x$$

Since $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$, the field is conservative.

Step 1: Integrate $P(x, y)$ with respect to x :

$$f(x, y) = \int (2xy + e^x) dx = x^2y + e^x + h(y)$$

Here, $h(y)$ is a function of y only.

Step 2: Differentiate $f(x, y)$ with respect to y :

$$\frac{\partial f}{\partial y} = x^2 + h'(y)$$

Set this equal to $Q(x, y) = x^2 + \frac{1}{2\sqrt{y}}$, so:

$$x^2 + h'(y) = x^2 + \frac{1}{2\sqrt{y}} \Rightarrow h'(y) = \frac{1}{2\sqrt{y}}$$

Step 3: Integrate to find $h(y)$:

$$h(y) = \int \frac{1}{2\sqrt{y}} dy = \sqrt{y} + C$$

Therefore, the potential function is:

$$f(x, y) = x^2y + e^x + \sqrt{y} + C$$

65.0.11 Double Integral over a Rectangular Region

Let $A = [x_0, x_1] \times [y_0, y_1]$ be a rectangular region in $\mathbb{R}^2$. The double integral of a function $f(x, y)$ over A is:

$$\iint_A f(x, y) dA = \int_{x_0}^{x_1} \int_{y_0}^{y_1} f(x, y) dy dx$$

For rectangular domains, the order of integration does not affect the result:

$$\int_{x_0}^{x_1} \int_{y_0}^{y_1} f(x, y) dy dx = \int_{y_0}^{y_1} \int_{x_0}^{x_1} f(x, y) dx dy$$

65.0.12 Steps for finding the Region of Integration

When Integrals are not defined over regions defined from constants then finding the bounds of integration can become a tricky task depending on the region we want to integrate.

The general process for setting up iterated integrals over a non-rectangular region is as follows:

1. **Sketch the region if possible:** Begin by sketching the region of integration. A visual representation helps you understand the boundaries, the shape of the region, and the relationships between variables.
2. **Choose the order of integration:** Decide the order in which you'll integrate the variables. Remember that integration proceeds from the inside out, so the inner integral corresponds to the variable you integrate first.
3. **Determine the bounds for each variable:** For the chosen order of integration, identify the limits of integration for each variable. The bounds for the inner variable are often functions of the outer variable. For example, if the region is described by the inequality $x + y < 1$ and you choose to integrate with respect to y first, you must express y 's bounds in terms of x . This means solving the inequality for y :

$$y < 1 - x \quad \Rightarrow \quad y \in [c(x), 1 - x]$$

where $c(x)$ is the lower bound for y , possibly another function of x . Then the full double integral would look like:

$$\int_a^b \int_{c(x)}^{1-x} f(x, y) dy dx$$

Repeat this process for all variables involved. If you've sketched the region, you can often find these bounds by "slicing" the region—holding one variable constant and observing how the others vary within the region.

65.0.13 Integration over Curvilinear Domains

If the region $A \subset \mathbb{R}^2$ is not rectangular, it is divided into subregions ΔA_k , and the integral is defined as the limit:

$$\iint_A f(x, y) dA = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k, y_k) \Delta A_k$$


In practice, for regions bounded by curves, we integrate in the form:

$$\int_a^b \int_{u(x)}^{o(x)} f(x, y) dy dx$$

Here, $y \in [u(x), o(x)]$ describes the vertical bounds as functions of x .

Example:

Compute the volume under $f(x, y) = 1 + x + y^2$ bounded by x -axis, $x = 1$ and y -axis $y = \sqrt{x}$



First we are going to imagine slicing the area under this sector vertically along the x -axis from 0 up to 1 which is the volume of an Area under a curve that is represented by going along the y -axis.

$$\begin{aligned}
 & \int_0^1 \int_0^{\sqrt{x}} 1 + x + y^2 dy dx \\
 & \int_0^1 \left[y + xy + \frac{y^3}{3} \right]_0^{\sqrt{x}} dx \\
 & \int_0^1 \left[\sqrt{x} + x\sqrt{x} + \frac{\sqrt{x}^3}{3} \right]_0^{\sqrt{x}} dx \\
 & = \left[\frac{2}{3}x^{\frac{3}{2}} + \frac{4}{3}\frac{2}{5}x^{\frac{5}{2}} \right]_0^1 = \frac{6}{5}
 \end{aligned}$$

If we choose the other way then we have an area with a fix y in the inside that goes from y^2 to 1 with respect to x .

$$\int_0^1 \int_{y^2}^1 1 + x + y^2 dx dy$$

65.0.14 Changing the order of integration to solve tricky integrals


Given $\int_0^8 \int_{\sqrt[3]{x}}^2 \frac{1}{y^4+1} dy dx$ which is very hard to integrate, let us change the order to make it easier.

$$\int_0^2 \int_0^{y^3} \frac{1}{y^4+1} dx dy$$

The steps are the same as in the previous Example.

1. Identify the area you want to integrate, what are the bounds respective to x and y
2. Write the same expression without bounds and change the order of the differentials
3. Imagine slicing the area underneath vertically for x and horizontally for y with their respective bounds corresponding to the perspective.
4. Write the new integral with the opposite of order of integration

Visualization of the area underneath the surface



Here the original bounds are: x goes from 0 to 8 and y goes from $\sqrt[3]{x}$ to 2. As you can see in the diagram by slicing vertically the value of x is fixed, and we move towards the upper-bound of y which is 2. Therefore, the smallest x can get is 0 and the biggest is 2. While for y by slicing vertically we get that x goes from 0 to $\sqrt[3]{x}$, but we do not want to leave x , so we solve for y and get y^3 . Now we can put all of this together and integrate.

In general changing the order of integration is only about using the logic to set the bounds of integration, but starting from a different perspective.

65.0.15 The Jacobian

The Jacobian matrix is a fundamental concept in multi-variable calculus. The Jacobian is the best approximation of a linear transformation for a point.

For the intuition of why, we use partial derivatives, just think about how we use them as this tiny changes in the component of original basis vectors.

Given a vector-valued function

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad f(x_1, \dots, x_n) = \begin{bmatrix} f_1(x_1, \dots, x_n) \\ f_2(x_1, \dots, x_n) \\ \vdots \\ f_m(x_1, \dots, x_n) \end{bmatrix},$$

the *Jacobian matrix* of f is the $m \times n$ matrix of all first-order partial derivatives:

$$J_f(x) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} & \dots & \frac{\partial f_1}{\partial x_n} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} & \dots & \frac{\partial f_2}{\partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \frac{\partial f_m}{\partial x_2} & \dots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}.$$

Local linearity

While performing non-linear transformations on a space. We can see that in a tiny neighborhood around a point, the transformation appears to be linear. And the matrix that encodes that transformation is the *Jacobian*.

For intuition, just imagine that in the tiny neighborhood the transformation is composed of small steps in the $x_1, \dots, x_n$ directions which give of the partial derivatives that represent this small steps.

Determinant

When $m = n$, i.e., the function maps from $\mathbb{R}^n$ to $\mathbb{R}^n$, the Jacobian matrix is square. In this case, the *Jacobian determinant* is defined as:

$$\det(J_f(x)),$$

and provides important information about the local behavior of the function. For instance:

- If $\det(J_f(x)) \neq 0$, the function is locally invertible at x by the inverse function theorem.
- The sign and magnitude of the determinant gives us information about orientation and volume scaling of the transformation.

When we change our systems of coordinates it is important that our differential make sense in the original system we were. Because of that, we take the determinant with respect to the new coordinates to tell us the scaling factor that we have to use for our result to make sense.

Use Cases:

The Jacobian matrix and its determinant appear in many areas, including:

- **Change of variables in integrals:** The absolute value of the Jacobian determinant is used to adjust the measure when changing coordinates in multiple integrals.
- **Nonlinear optimization:** The Jacobian is crucial for computing gradients and performing techniques like Newton's method in multiple dimensions.
- **Differential equations:** Stability analysis of dynamical systems often involves evaluating the Jacobian at equilibrium points.

List of Common Jacobians

- *Cartesian Coordinates:*
 - Definition: The position of a point is given by (x, y, z) in a 3D space along orthogonal axes.
 - Jacobian:

$$J_{\text{Cartesian}} = 1dx$$

- *Polar Coordinates (2D):*
 - Definition: The position of a point in a plane is given by (r, θ) , where r is the radial distance and θ is the angle from the positive x -axis.
 - Jacobian:

$$J_{\text{Polar}} = r dr d\theta$$

- *Cylindrical Coordinates:*
 - Definition: The position of a point in 3D space is given by (r, θ, z) , where (r, θ) are the polar coordinates in the xy -plane and z is the height along the z -axis.
 - Jacobian:

$$J_{\text{Cylindrical}} = r dr d\theta dz$$

- *Spherical Coordinates:*
 - Definition: The position of a point in 3D space is given by (ρ, θ, ϕ) , where:
 - * ρ is the distance from the origin,
 - * θ is the azimuthal angle (in the xy -plane from the positive x -axis),

* ϕ is the polar angle (from the positive z -axis).

– Jacobian:

$$J_{\text{Spherical}} = \rho^2 \sin(\phi) d\rho d\theta d\phi$$

65.0.16 Integration in Polar Coordinates

For radially symmetric regions or functions, switching to polar coordinates simplifies the computation.

Transformation:

$$x = r \cos \varphi, \quad y = r \sin \varphi$$

The Jacobian of the transformation gives the area element:

$$dA = r dr d\varphi$$

Integral:

$$\iint_D f(x, y) dA = \int_{\varphi_0}^{\varphi_1} \int_0^{R(\varphi)} f(r \cos \varphi, r \sin \varphi) r dr d\varphi$$

Origin of the Area

When we divide our circle in the sections we get both a difference in then angle and the radius which generate another area. The difference $\Delta A = \text{Area of big wedge} - \text{Area of small wedge}$

Radius: $r_k + \frac{\Delta r}{2}$ for the big wedge and $r_k - \frac{\Delta r}{2}$ for the small wedge

Angle: $\frac{\Delta \theta}{2\pi}$ which is the angle of the fraction of the circle because the formula for the area is $r^2\pi$ and the whole circle would be 2π

Final Formula for the difference in the Area

$$\begin{aligned} \Delta A &= \frac{\Delta \theta}{2\pi} \left(\left[r_k + \frac{\Delta r}{2} \right]^2 - \left[r_k - \frac{\Delta r}{2} \right]^2 \right) \\ &= r_k \Delta r \Delta \theta \end{aligned}$$

Which multiplied by the function value and by taking the limit of these sections we get that the volume:

$$V = \int_{\theta_1}^{\theta_2} \int_{r_1(\theta)}^{r_2(\theta)} f(r, \theta) r \Delta r \Delta \theta$$

Example:

$f(r, \theta) = r$ above cardioid $r = 1 - \sin \theta$

Then

$$\begin{aligned} V &= \int_0^{2\pi} \int_0^{1-\sin \theta} r r dr d\theta \\ &= \int_0^{2\pi} \frac{r^3}{3} \Big|_0^{1-\sin \theta} d\theta = \int_0^{2\pi} \frac{(1-\sin \theta)^3}{3} d\theta = \frac{5\pi}{3} \end{aligned}$$

65.0.17 Improper Integrals over Unbounded Regions

If the domain of integration is unbounded (e.g., the entire plane $\mathbb{R}^2$), the double integral is defined via a limit process:

$$\iint_{\mathbb{R}^2} f(x, y) dx dy := \lim_{M_1, M_2 \rightarrow \infty} \lim_{N_1, N_2 \rightarrow \infty} \int_{M_1}^{M_2} \int_{N_1}^{N_2} f(x, y) dy dx$$

Often, using polar coordinates simplifies such cases, reducing multiple limits to one:

$$\iint_{\mathbb{R}^2} f(x, y) dx dy = \int_0^{2\pi} \int_0^\infty f(r \cos \varphi, r \sin \varphi) r dr d\varphi$$

Gaussian Integral

We are going to find the integral of the *unbounded integral* e^{-x^2} using a combination of techniques.

$$\begin{aligned} S &= \int_{-\infty}^{\infty} e^{-x^2} dx \\ S^2 &= \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \\ S^2 &= \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-y^2} dy \right) \end{aligned}$$

Now let us translate to polar coordinates

$$\begin{aligned} S^2 &= \left(\int_0^{2\pi} \int_0^\infty r e^{-(r \cos \theta)^2} r e^{-(r \sin \theta)^2} dr d\theta \right) \\ S^2 &= \left(\int_0^{2\pi} \int_0^\infty r e^{-(r \cos \theta)^2 + (r \sin \theta)^2} dr d\theta \right) \\ S^2 &= \left(\int_0^{2\pi} \int_0^\infty r e^{-(r^2)} dr d\theta \right) \\ S^2 &= \left(\int_0^{2\pi} \int_0^\infty \frac{1}{2} e^{-s} ds d\theta \right) \\ S^2 &= \left(\int_0^{2\pi} d\theta \right) \\ S^2 &= \pi \\ S &= \sqrt{\pi} \end{aligned}$$

ΩℰΩ

65.0.18 Triple Integrals

The triple integral allows the computation of volume or mass in three-dimensional space.

Let $V \subset \mathbb{R}^3$ be a bounded region, then:

$$\iiint_V f(x, y, z) dV = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k, y_k, z_k) \Delta V_k,$$

where ΔV_k are small sub-volumes approximating V .

In practice, evaluate:

$$\iiint_V f(x, y, z) \, dz \, dy \, dx,$$


with limits determined by the geometry of the volume. The order of integration may be rearranged to simplify computation.

$$\int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} \int_{f_1(x,y)}^{f_2(x,y)} dz \, dy \, dx$$

Triple Integrals in Cartesian Coordinates

In this short section we, will look at an example of how to calculate the volume enclosed in between two functions using triple integrals.

Given are the functions $f_1(x, y) = x^2 + y^2$ and $f_2(x, y) = 3 - x^2 - y^2$



Let us find the intersection

$$x^2 + y^2 = 3 - x^2 - y^2$$

$$x^2 + y^2 = \frac{3}{2}$$

Now we have to find the limits of integration for

$$\int_{x=a}^{x=b} \int_{y=g_1(x)}^{y=g_2(x)} \int_{f_1(x,y)}^{f_2(x,y)} dz \, dy \, dx,$$

using our constraint and our original functions.

$$\int_{x=a}^{x=b} \int_{y=-\sqrt{\frac{3}{2}-x^2}}^{\sqrt{\frac{3}{2}-x^2}} \int_{x^2+y^2}^{3-x^2-y^2} dz \, dy \, dx$$

Here z means the height which is just the output of the original functions.

The bounds of y just by looking at our constraint and solving for it. And x only depends on the concrete limit of integration.

Coordinate Transformation and the Jacobian Determinant

Let a coordinate transformation be defined by:

$$x = x(u, v, w), \quad y = y(u, v, w), \quad z = z(u, v, w)$$

Then the triple integral transforms as:

$$\iiint_{(x,y,z)} f(x, y, z) \, dx \, dy \, dz = \iiint_{(u,v,w)} f(x(u, v, w), y(u, v, w), z(u, v, w)) \cdot |\det J| \, du \, dv \, dw$$

where J is the *Jacobian matrix* of the transformation:

$$J = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{bmatrix}$$

Example: Transformation to Cylindrical Coordinates

The standard cylindrical coordinate transformation is:

$$x = r \cos \varphi, \quad y = r \sin \varphi, \quad z = z$$

The Jacobian determinant is:

$$|\det J| = \begin{vmatrix} \cos \varphi & -r \sin \varphi & 0 \\ \sin \varphi & r \cos \varphi & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$$

Thus, the triple integral in cylindrical coordinates becomes:

$$\iiint f(x, y, z) \, dx \, dy \, dz = \iiint f(r \cos \varphi, r \sin \varphi, z) \, r \, dr \, d\varphi \, dz$$

Calculating the center of mass with triple integrals

We can make use of triple integration to calculate the center of mass of a body with respect to x , y or z with the following formulas:

$$\begin{aligned} x_s &= \iiint_V x p(x, z, y) \, dz \, dy \, dx \\ y_s &= \iiint_V y p(x, z, y) \, dz \, dy \, dx \\ z_s &= \iiint_V z p(x, z, y) \, dz \, dy \, dx \end{aligned}$$

Where $p(x, y, z)$ is our mass function.

65.0.19 Surfaces

In vector calculus sometimes, we want to examine the surface generated by a function. An approach to analyze this surfaces is via parameterization, and another is using explicit and implicit forms.

Explicit and Implicit Forms

For a 3d surfaces the *Implicit form* is given by

$$F(x, y, z) = c$$

and the *Explicit* by

$$z = f(x, y)$$

Parametric Form

It is given by

$$\vec{r}(u, v) = f(u, v)\hat{i} + g(u, v)\hat{j} + h(u, v)\hat{k}$$

where $a \leq u \leq b$ and $c \leq v \leq d$.

Example:

For the example we take the cone formed by the function $z = \sqrt{x^2 + y^2}$ which is in the explicit form.

The implicit form is then $0 = z - \sqrt{x^2 + y^2}$ and the parametric form can be described very naturally by

$$\vec{r}(r, \theta) = r \cos(\theta)\hat{i} + r \sin(\theta)\hat{j} + r\hat{k}$$

using something similar to cylindrical coordinates with $r = \sqrt{x^2 + y^2}$.

Surface Area Formula

Continuing with the surfaces in 3d we can compute the area using the following formula

$$A = \iint_D \left\| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right\| du dv.$$

Notice that we are using the fact that the length of a cross product is the area of the parallelogram formed by vectors it is orthogonal to. Using this on a small scale with the partial of u and v and then adding up results in the formula.

Example I: Area of the sphere

We want to compute the surface area of a sphere of radius a and show that it equals $4\pi a^2$. Let us proceed step by step.

The equation of a sphere of radius a centered at the origin is given implicitly by

$$x^2 + y^2 + z^2 = a^2.$$

To parametrize the sphere, one possible way is to solve for z :

$$z = \pm \sqrt{a^2 - x^2 - y^2}, \quad x^2 + y^2 \leq a^2.$$

This representation works, but it splits the sphere into two surfaces (upper and lower hemispheres). This makes the parametrization less convenient for computing surface area.

A more natural parametrization comes from spherical coordinates. Any point on the sphere can be written as

$$\begin{cases} x = a \sin \theta \cos \varphi, \\ y = a \sin \theta \sin \varphi, \\ z = a \cos \theta, \end{cases}$$

where the parameters range over

$$0 \leq \theta \leq \pi, \quad 0 \leq \varphi < 2\pi.$$

Here, θ is the polar angle (measured from the z -axis) and φ is the azimuthal angle in the xy -plane.

The surface area of a parametrized surface $\vec{r}(u, v)$ is given by

$$A = \iint_D \left\| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right\| du dv.$$

For the parametrization $\vec{r}(\theta, \varphi) = (a \sin \theta \cos \varphi, a \sin \theta \sin \varphi, a \cos \theta)$, we compute:

$$\frac{\partial \vec{r}}{\partial \theta} = (a \cos \theta \cos \varphi, a \cos \theta \sin \varphi, -a \sin \theta),$$

$$\frac{\partial \vec{r}}{\partial \varphi} = (-a \sin \theta \sin \varphi, a \sin \theta \cos \varphi, 0).$$

Their cross product is

$$\frac{\partial \vec{r}}{\partial \theta} \times \frac{\partial \vec{r}}{\partial \varphi} = (a^2 \sin^2 \theta \cos \varphi, a^2 \sin^2 \theta \sin \varphi, a^2 \sin \theta \cos \theta).$$

The magnitude is

$$\left\| \frac{\partial \vec{r}}{\partial \theta} \times \frac{\partial \vec{r}}{\partial \varphi} \right\| = a^2 \sin \theta.$$

Therefore, the surface area is

$$A = \int_0^{2\pi} \int_0^\pi a^2 \sin \theta d\theta d\varphi.$$

Compute the integrals:

$$A = a^2 \left(\int_0^{2\pi} d\varphi \right) \left(\int_0^\pi \sin \theta d\theta \right).$$

$$A = a^2(2\pi)(2) = 4\pi a^2.$$

Thus, the surface area of the sphere is

$$4\pi a^2$$

□◻◻

Example II:

Compute the surface area of the intersection of the plane $z = -x$ and the cylinder $x^2 + y^2 = 4$.

We will use polar coordinates because the intersection is circular.

$$\vec{r}(r, \theta) = \begin{bmatrix} r \cos(\theta) \\ r \sin(\theta) \\ -r \cos(\theta) \end{bmatrix}$$

Take the partials

$$\vec{r}_r = \begin{bmatrix} \cos(\theta) \\ \sin(\theta) \\ -\cos(\theta) \end{bmatrix}$$

$$\vec{r}_\theta = \begin{bmatrix} -r \sin(\theta) \\ r \cos(\theta) \\ r \sin(\theta) \end{bmatrix}$$

Compute the cross product:

$$\vec{r}_r \times \vec{r}_\theta = \begin{bmatrix} r \\ 0 \\ r \end{bmatrix}, \quad \|\vec{r}_r \times \vec{r}_\theta\| = \sqrt{r^2 + r^2} = \sqrt{2} r$$

Now integrate over the disk $0 \leq r \leq 2$, $0 \leq \theta \leq 2\pi$:

$$\text{Area} = \int_0^{2\pi} \int_0^2 \sqrt{2} r \, dr \, d\theta = \int_0^{2\pi} \left[\frac{\sqrt{2}}{2} r^2 \right]_0^2 d\theta = \int_0^{2\pi} 2\sqrt{2} \, d\theta = 4\pi\sqrt{2}.$$

The final result is

$$4\pi\sqrt{2}$$

65.0.20 Implicit Surface Area

Given a smooth, continuous surface $F(x, y, z) = c$ where $\langle \nabla F, \vec{p} \rangle \neq 0$ the surface area is given by

$$SA = \iint_R \frac{\|\nabla F\|}{|\langle \nabla F, \vec{p} \rangle|} dA$$

$\vec{p}$ can be $\hat{i}$, $\hat{j}$ or $\hat{k}$.

65.0.21 Explicit Surface Area

Given a smooth, continuous surface $F(x, y, z) = f(x, y) - z = 0$ where $\nabla F = f_x \hat{i} + f_y \hat{j} - \hat{k}$, $|\langle \nabla F, \hat{k} \rangle| = |-1| = 1$ and $\|\nabla F\| = \sqrt{f_x^2 + f_y^2 + 1}$

$$SA = \iint_R \sqrt{f_x^2 + f_y^2 + 1} dA$$

65.0.22 Surface Integrals

We define a function $G(x, y, z)$ define on a smooth surface S with an area $d\sigma$ whose integral is

$$SI = \iint_R G(x, y, z) d\sigma$$

The function G associates each point on the surface to a value in a higher dimension.

Surface integral in different forms:

Parametric

$$SI = \iint_R G(f(u, v), g(u, v), h(u, v)) \left\| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right\| du dv.$$

Explicit

$$SI = \iint_R G(x, y, z) \sqrt{f_x^2 + f_y^2 + 1} dA$$

Implicit

$$SI = \iint_R G(x, y, f(x, y)) \frac{\|\nabla F\|}{|\langle \nabla F, \vec{p} \rangle|} dA$$

65.0.23 The Orientation of a Surface

A smooth surface is *orientable* if there exists a continuous field of unit normal vectors to the surface.

65.0.24 The Flux of a Surface

For a continuous field $\vec{F}$ on a smooth surface S oriented with normal unit vectors $\vec{n}$,

$$Flux = \iint_S \langle \vec{F}, \vec{n} \rangle d\sigma$$

where σ is the surface area in the corresponding form. And the normal unit $\vec{n}$ is for parametric forms

$$\vec{n} = \frac{\frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v}}{\left\| \frac{\partial \vec{r}}{\partial u} \times \frac{\partial \vec{r}}{\partial v} \right\|}$$

and for the implicit surfaces

$$\vec{n} = \frac{\nabla g}{\|\nabla g\|}$$

65.0.25 Curl

Let $\vec{F} = (F_1, F_2, F_3) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a vector field.

$$\text{curl } \vec{F} = \nabla \times \vec{F} = \begin{bmatrix} \frac{\partial F_3}{\partial x_2} - \frac{\partial F_2}{\partial x_3} \\ \frac{\partial F_1}{\partial x_3} - \frac{\partial F_3}{\partial x_1} \\ \frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2} \end{bmatrix}$$

These describe how the field spreads or rotates around a point. As a side not, the curl of the gradient of f is 0 if f has continuous mixed second derivatives.

There exists a connection between the curl and conservative vector fields. Notice that the components of the curl vector are the conditions similar to the conditions for a vector field to be conservative. Therefore, a conservative vector field has no curl because if the conditions are true then the curl is the zero-vector.

A point where $\text{curl } \vec{f} \neq 0$ is called a whirlpool.

65.0.26 Stokes's Theorem

For S a piecewise smooth oriented surface with piecewise smooth boundary C and $\vec{F}$ a field with continuous first partials for each component on an open region containing S . The circulation around the curve is equal to

$$C = \oint_C \langle \vec{F}, d\vec{r} \rangle = \iint_S \langle (\nabla \times \vec{F}), \vec{n} \rangle d\sigma$$

65.0.27 Divergence

We define *divergence* as

$$\text{div } \vec{F} = \langle \nabla, \vec{F} \rangle = \frac{\partial F_1}{\partial x_1} + \frac{\partial F_2}{\partial x_2} + \frac{\partial F_3}{\partial x_3}$$

A point where $\text{div } \vec{f} > 0$ is called a source.

65.0.28 The Divergence Theorem

For S a piecewise smooth oriented, closed surface with piecewise smooth boundary C and $\vec{F}$ a field with continuous first partials for each component on an open region containing S . The flux around the curve is equal to

$$\iint_S \langle \vec{F}, \vec{n} \rangle ds = \iiint_D \langle \nabla, \vec{F} \rangle dV$$

65.0.29 Unified View of Vector Calculus

Integrating a differential operator acting on a field over a domain is the same as adding the field components along the boundary.

Formally:

The integral of a differential form over a boundary $\partial\Omega$ of some orientable manifold Ω is equal to the integral of its exterior derivative over the whole of Ω .

$$\int_{\partial\Omega} w = \int_{\Omega} dw$$

Chapter 66

Differential Equations

A Differential Equation is an equation that relates a function (the dependent variable) with a variable or multiple variables (the independent variables) and its derivative.

They can be categorized in *Ordinary* and *Partial* Differential Equations.

66.0.1 Ordinary Differential Equations

An equation of the form

$$y^n = f(x, y, x', y', \dots, y^{n-1})$$

is called *Ordinary Explicit Differential Equations of order n*

If $y^n = 0$ then it is *implicit*

If the exponent of the dependent variable is 1, and it is build like a linear equation then it is categorized as *Linear*

66.0.2 Initial Value Problems and Solutions

The specification of an explicit differential equation and the values

$$x_0, \quad y(x_0) = y_0, \quad y'(x_0) = y_1, \quad \dots, \quad y^{(n-1)}(x_0) = y_{n-1}$$

is called an *initial value problem (IVP)*.

An n -times differentiable function $y(x)$ that satisfies the explicit differential equation is called the *general solution* of the differential equation. If it also satisfies the conditions of the initial value problem from the previous definition, it is called the *particular solution* of the IVP.

66.0.3 Volterra Integral

The function $y(x)$ is a solution of the IVP $y' = f(x, y)$, $y(x_0) = y_0$ if and only if

$$y(x) = y_0 + \int_{x_0}^x f(t, y(t)) dt$$

Proof:

$$\begin{aligned} \int_{x_0}^x f(t, y(t)) dt &= \int_{x_0}^x y' \\ y(x) &= y(x_0) \end{aligned}$$

The other way would be

$$\left(\int_{x_0}^x f(t, y(t)) dt + y_0 \right)' = f(x, y(x)) = y'$$

QED

66.0.4 The Picard-Lindelöf Iteration Method

We can approximate the desired solution using the Volterra integral equation as follows. Start with an arbitrary initial function, for example a constant — preferably the initial value:

$$g_0(x) = y_0$$

Then form the next approximation to the desired function $y(x)$ as:

$$g_{n+1}(x) = y_0 + \int_{x_0}^x f(t, g_n(t)) dt$$

66.0.5 Integration case

To solve a ODE of the form $y' = f(x)$ it is as easy as integrating $f(x)$.

66.0.6 Separation of Variables

An equation $F(x, y, y', \dots) = 0$ is called an equation of *separable variables* if it can be written in one of the following forms:

- $\frac{dy}{dx} = f(x)g(y)$
- $\frac{dy}{dx} = \frac{f(x)}{g(y)}$
- $\frac{dy}{dx} = \frac{g(y)}{f(x)}$

Steps to solve a ODE of separable variables

1. Solve for the derivative
2. Check that it is indeed an ODE of separable variables
3. Separate the variables
4. Integrate with respect to the independent variable

Example:

Given is $(y - 2)y' = 3x - 5$

It is an ODE of separable variable

$$y' = \frac{3x - 5}{y - 2}$$

Now we separate the variables and Integrate

$$\int (y - 2)y' dx = \int 3x - 5 dx$$

Note that $y' = \frac{dy}{dx}$ and $\frac{dy}{dx} dx = dy$ (only as a convenient fiction in reality we are changing the variables) thus,

$$\int (y-2)dy = \int 3x-5dx$$

$$\frac{y^2}{2} - 2y + c = \frac{3x^2}{2} - 5x + c$$

The solution we get is an implicit solution. To make it look nice solve for y if you can.

Understanding the fiction

The previous fiction we used can explain in the following manner.

Given $f(y)\frac{dy}{dx}dx = \int f(y)dy$. We want to prove that this equality holds via the *chain rule* $\frac{du}{dx} = \frac{du}{dy}\frac{dy}{dx}$.

$$\begin{aligned}\frac{d}{dx}\left(\int f(y)dy\right) &= \frac{d}{dx}\left(\int f(y)dy\right)\frac{dy}{dx} \\ &= f(y)dy\frac{dy}{dx} \\ \implies \int f(y)\frac{dy}{dx} &= \int f(y)dy\end{aligned}$$

QED

66.0.7 Geometry of ODE

You can imagine the plot of an ODE as a *Slope field* where the slope (derivative) of a function obeys a certain condition, this being the differential equation $\frac{dy}{dx} = \text{expr}$.

A point on this field is called an *Initial Condition* and the curve that goes through that point in the slope field is called an *Integral curve* which is a solution to the differential equation with respect to that initial condition.

66.0.8 Existence and Uniqueness

Let $f(x, y)$ be continuous on a rectangle $[a, b] \times \mathbb{R}$. Then the differential equation

$$y' = f(x, y)$$

has at least one solution. A unique solution is only guaranteed under additional conditions.

A function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is called *Lipschitz continuous with respect to y* if there exists a constant $L > 0$ such that

$$|f(x, y_1) - f(x, y_2)| \leq L|y_1 - y_2| \quad \text{for all } (x, y_1), (x, y_2) \in D.$$

Let $x_0 \in [a, b]$, and let $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous and bounded, and Lipschitz continuous in y . Then the initial value problem

$$y' = f(x, y), \quad y(x_0) = y_0$$

has a unique solution on $[a, b]$.

Or in easier words: If f and $\frac{\partial f}{\partial y}$ are continuous near (x_0, y_0) then there is a unique solution on an interval $\alpha < x_0 < \beta$ to the IVP

$$y' = f(x, y), \quad y(x_0) = y_0$$

66.0.9 How to deal with initial conditions

After solving a differential equation we may have the problem that certain initial conditions were set at the start, thus, making our general solution not specific.

Solving this problem is pretty easy, because we just have to solve for the constant terms in our general solution with respect to our initial conditions. Like for example $y(0) = 1$ we would plug 0 for all x 's in our general solution the whole equation will be set equal to 1.

66.0.10 Solving DE with Substitution

In some cases a complex function can be simplified via a substitution.

Case I

$$y' = f(ax + by(x) + c)$$

With a substitution like $z(x) = ax + by(x) + c$ the right-hand side will become $f(z)$ and for the left-hand side we get $y(x) = \frac{z(x) - ax - c}{b}$.

Now we can substitute in the original equation, and we get

$$y'(x) = \frac{1}{b}(z'(x) - a) = f(z)$$

$$z'(x) = a + bf(z)$$

Now we can integrate and find the solution and then return to our original variables.

Example:

$$y' = (x + y(x))^2$$

Here $a = 1$, $b = 1$, $c = 0$ and $f(z) = z^2$

1. Substitute

$$z(x) = x + y(x)$$

$$y(x) = z(x) - x$$

$$y'(x) = z'(x) - 1$$

2. Plug in the DE

$$y' = (x + y(x))^2$$

$$z'(x) - 1 = z(x)^2$$

3. Integrate

$$z'(x) = 1 + z(x)^2$$

$$\frac{z'(x)}{1 + z(x)^2}$$

$$\int \frac{1}{1 + z^2} dz = \int 1 dx$$

$$\arctan(z) = x + c$$

$$z = \tan(x + c)$$

4. Return to the original variables

$$x + y(x) = \tan(x + c)$$

$$y(x) = \tan(x + c) - x$$

Case II

$$y' = f\left(\frac{y}{x}\right)$$

For this to be valid $z(x) = \frac{y}{x}$ our $y(x)$ needs to look like $y(x) = z(x)x$ and so $y' = z(x) + z'(x)x$.

Now we substitute in the differential equation using $f(z) = z(x) + z'(x)x$

$$z'(x) = \frac{f(z) - z(x)}{x}$$

Now we can integrate and then return to our original variables. $y(x) = z(x)x$, where $z(x)$ is our result of the integration.

Example:

$$y' = \frac{y(x)}{x} + 1$$

This is converted to $y' = f(z) = z + 1$

1. Substitute

$$z(x) = \frac{y(x)}{x}$$

$$y(x) = z(x)x$$

$$y'(x) = z(x) + z'(x)x$$

2. Plug in the DE

$$z(x) + z'(x)x = z(x) + 1$$

3. Integrate

$$z'(x)x = 1$$

$$\int z' dz = \int \frac{1}{x} dx$$

$$z = (\ln(x) + c)x$$

66.0.11 Linear Differential Equations

An ODE is called linear if all occurrences of the dependent variable and its derivatives have an exponent of at most 1.

$$a_n(x)y^n + a_{n-1}y^{n-1} + \cdots + a_1(x)y' + a_0(x)y = b(x)$$

If such an Equation has also the property $b(x) = 0$ it is called *Homogeneous*

How to solve a Linear Differential Equation

In this case we will focus on first Order ODE.

Write it in the standard form $y' + p(x)y = f(x)$

Now use *Integrating Factor Method*

66.0.12 Integrating Factor Method

We are going to multiply our expression by function $r(x)$

$$r(x)y' + r(x)p(x)y = r(x)f(x)$$

Now we would like to write the left side as $\frac{d}{dx}yr(x)$ because now if we were to integrate it would take us to the expression in the right.

When we evaluate this new expression we get that

$$\frac{d}{dx}yr(x) = y'r(x) + r'(x)y$$

which is not quite what we actually have, but it is close. Now note that we can say

$$r'(x) = r(x)p(x)$$

This is a differential equation of separable variables thus, we can

$$\frac{r'(x)}{r(x)} = p(x)$$

$$\int \frac{r'(x)}{r(x)} dx = \int p(x) dx$$

$$\ln(r(x)) = \int p(x) dx$$

after using e

$$r(x) = e^{\int p(x) dx}$$

Now we have found an $r(x)$ and now that we know that such a function exists we can return to our initial problem and say

$$\frac{d}{dx}r(x)y = r(x)f(x)$$

$$y = \frac{1}{r(x)} \int r(x)f(x) dx$$

and

$$r(x) = e^{\int p(x) dx}$$

To summarize, given $y' + f(x)y = g(x)$ the steps are:

1. Find $c(x) = e^{\int f(x) dx}$.

2. Multiply the original expression with $c(x)$.

$$y'c(x) + f(x)c(x)y = g(x)c(x)$$

3. Use the power rule of differentiation to simplify.

$$y'c(x) + f(x)c(x)y = \frac{d}{dx}(c(x)y)$$

4. Integrate.

$$\begin{aligned}\int \frac{d}{dx}(c(x)y) dx &= \int g(x)c(x) dx \\ c(x)y &= \int g(x)c(x) dx \\ y &= \frac{\int g(x)c(x) dx}{c(x)}\end{aligned}$$

Example:

Given are $y' + 4y = e^{-x}$ and $y(0) = \frac{4}{3}$.

Here $4 = p(x)$ and $e^{-x} = f(x)$.

Now let us use the formula for $r(x) = e^{\int p(x)}$. This gives us $r(x) = e^{\int 4dx} = e^{4x}$.

Now recall what we saw earlier

$$e^{4x}y' + e^{4x}4y = e^{4x}e^{-x} = e^{3x}$$

Our left side is just $\frac{d}{dx}e^{4x}y$

Now let us complete the exercise with our last formula by integrating both sides

$$\begin{aligned}\frac{d}{dx}e^{4x}y &= e^{3x} \\ y &= \frac{1}{e^{4x}} \int e^{3x} \\ y &= \frac{1}{e^{4x}} \frac{1}{3} e^{3x} + c\end{aligned}$$

Which is our general solution

To find our specific solution we just plug 0 in to our general solution that gives us $\frac{1}{3} + c = \frac{4}{3}$ thus, $c = 1$

66.0.13 Homogeneous Differential Equations

A differential equation

$$M(x, y)dx + N(x, y)dy = 0$$

is considered *Homogeneous* if and only if $M(x)$ and $N(x)$ are *homogeneous*, and they have the same degree.

The equation $y' + f(x)y = 0$ is called *homogeneous linear differential equation of first order*.

Homogeneous Equation

A function $f(x, y)$ is called *homogeneous* if and only if it can be written

$$f(xt, yt) = t^n f(x, y)$$

Example:

$$\begin{aligned}f(x, y) &= x^3 + y^3 \\f(tx, ty) &= (tx)^3 + (ty)^3 \\t^3(x^3 + y^3) &= t^3 f(x, y)\end{aligned}$$

It is important to note that the key is that each term have a total degree of n . For example: $x^2y + x3y^2$

Homogeneous Functions Theorem

If $f(x, y)$ is homogeneous of degree 0 in x and y then f is a function of $\frac{y}{x}$

Or $f(x, y, y')$ is homogeneous if it can be written as $y' = f(\frac{x}{y})$ or $y' = f(\frac{y}{x})$

66.0.14 Solving Homogeneous DE I

1. Write in the form $M(x, y)dx + N(x, y)dy = 0$
2. Check if it is homogeneous
3. Change the variables to $y = ux$ or $x = uy$ depending on the situation
4. Solve using the method of separable variables method
5. Rewrite the answer in terms of the original variables again

Example:

Give is the function $(x - y)dx = -xdy$

Step 1. We write it in the desired form $(x - y)dx + xdy = 0$

Step 2. We notice that both are of degree 1. This time we skip the rigorous check.

Step 3. We are going to choose the simpler function to do the change of variable depending on the *differential!*

The simpler function is xdy thus, we are to change y

$$\begin{aligned}y &= ux \\dy &= ux' + xu'\end{aligned}$$

and

$$u = \frac{y}{x}$$

Now substitute

$$\begin{aligned}(x - ux)dx + x(udx + xdu) &= 0 \\xdx - uxdx + xudx + x^2du &= 0 \\(x - ux + xu)dx + x^2du &= 0 \\xdx + x^2du &= 0 \\xdx &= -x^2du \\dx &= -xdu\end{aligned}$$

Step 4. Now we can integrate both sides

$$\int \frac{1}{x} dx = \int du$$

$$\ln x + c = -u + c$$

Step 5. Now we use $u = \frac{y}{x}$ to return to our original variables

$$\ln x + c = -\frac{y}{x} + c$$

$$-x \ln x - xc = y$$

66.0.15 Solving Homogeneous DE I

Another method for solving this kind of equations $y' + f(x)y = 0$ is to follow the next steps.

Step 1: Separate the variables

$$\frac{dy}{dx} = -f(x)y$$

Step 2: Use the separation of variables

$$\int \frac{\frac{dy}{dx}}{y} = \int -f(x) dx$$

$$\int \frac{dy}{y} = \int -f(x) dx$$

$$\ln |y| = \int -f(x) dx$$

$$y = Ce^{\int -f(x) dx}$$

This method works more directly.

66.0.16 Inhomogeneous Equation

The equation $y' + f(x)y = g(x)$ is called *linear inhomogeneous differential equation of first order* $g(x)$ is called the *distortion function*. The correspondent IVP is called *linear IVP*.

66.0.17 Solutions of LDE

Every linear IVP with continuous functions $f(x)$ and $g(x)$ plus a bounded $f(x)$ has exactly one solution.

Proof:

$$y' = -f(x)y + g(x) \text{ therefore, } |-f(x)y_1 + g(x) - (-f(x)y_2 + g(x))| = |f(x)||y_1 - y_2|$$

$$\leq M|y_1 - y_2|$$

Therefore, the function is Lipschitz continuous, thus, a solution is possible.

□◻◻

Variation of the Constants

Recall that for the homogeneous case $y' + f(x)y = 0$ case we use $y = ce^{\int -f(x)dx}$, but now we have $y' + f(x)y = g(x)$, thus, let us use the following approach

$$y = c(x)e^{\int -f(x)dx}$$

Now use the product rule to derivate

$$\begin{aligned}y' &= c'(x)e^{\int -f(x)dx} + c(x)e^{\int -f(x)dx}(-f(x)) \\&= c'(x)e^{\int -f(x)dx} - f(x)y\end{aligned}$$

thus,

$$y' + f(x)y = c'(x)e^{\int -f(x)dx}$$

therefore, $g(x) = c'(x)e^{\int -f(x)dx}$

Then we can get a solution for $c(x)$ from

$$\begin{aligned}g(x) &= c'(x)e^{\int -f(x)dx} \\g(x) &= c'(x)e^{-\int f(x)dx} \\c'(x) &= \frac{g(x)}{e^{-\int f(x)dx}} \\c'(x) &= g(x)e^{\int f(x)dx}\end{aligned}$$

Example:

Given is $y' - \cos(x)y = xe^{\sin(x)}$ with $y(0) = 3$.

1. Solve as if it were homogeneous

$$\begin{aligned}y' - \cos(x)y &= 0 \\y' &= \cos(x)y \\\frac{y'}{y} &= \cos(x) \\\int \frac{1}{y}dy &= \int \cos(x)dx \\\ln |y| + c &= \sin(x) + c \\y &= e^{\sin(x)}c\end{aligned}$$

2. Use $y = c(x)e^{\sin(x)}$

$$\begin{aligned}y' &= c(x)\cos(x)e^{\sin(x)} + c'(x)e^{\sin(x)} \\y' &= \cos(x)y + c'(x)e^{\sin(x)} \\y' - \cos(x)y &= c'(x)e^{\sin(x)}\end{aligned}$$

3. Compare with the original equation

$$y' - \cos(x)y = xe^{\sin(x)}$$

$$y' - \cos(x)y = c'(x)e^{\sin(x)}$$

4. Solve for $c'(x)$

$$xe^{\sin(x)} = c'(x)e^{\sin(x)}$$

$$x = c'(x)$$

5. Integrate

$$\int x dx = \int c'(x) dx$$

$$\frac{x^2}{2} + c = c(x)$$

6. Substitute in our approach $y = c(x)e^{\sin(x)}$ from earlier

$$y = \left(\frac{x^2}{2} + c \right) e^{\sin(x)}$$

7. Solve for c using our initial condition if asked

$$y(0) = 3$$

$$3 = \left(\frac{0^2}{2} + c \right) e^{\sin(0)}$$

$$3 = c$$

66.0.18 Existence and Uniqueness Differential Equations

For the IVP

$$y^n + p_{n-1}y^{n-1} + \cdots + p_0(x)y = f(x)$$

$$y(a) = b_0, y'(a) = b_1, \dots, y^{n-1}(a) = b_{n-1}$$

If all p_i and f are continuous on the interval I about a , then there exists a unique solution on I .

66.0.19 Theorem of Superposition and the General Solution

Suppose $y_1, \dots, y_n$ solve

$$y^n + p_{n-1}y^{n-1} + \cdots + p_0(x)y = 0$$

Then $y = c_1y_1 + \cdots + c_ny_n$ also solves the equation, and it is the *General Solution* if and only if the *Wronskian* $W(t) \neq 0$ for some t_0 .

66.0.20 Undetermined Coefficients

Given an equation $y' + ay = g(x)$ notice that a does not depend on x . The homogeneous equation to this case will be $y' + ay = 0$ with a solution y_h and some partial solution y_p . Now remember from *Linear Algebra* the *general* = *particular* + *homogeneous* more precisely $y = y_p + y_h$ therefore, adding a particular solution to a homogeneous does not change the general solution of our DE.

Now let us substitute in our differential equation.

$$\begin{aligned}(y_h + y_p)' + a(y_h + y_p) &= y_h' + y_p' + ay_h + ay_p \\ &= (y_h' + ay_h) + (y_p' + ay_p) = 0 + g(x) = g(x)\end{aligned}$$

Steps for solving Constant Coefficients problems

1. Find the *homogeneous* solution for $y' + ay = 0$. y_h is going to have a free parameter.
2. Find a partial solution for $y' + ay = g(x)$
3. Use $y = y_h + y_p$
4. If you have an IVP solve for the free parameter in y_h

Now we have to find a way to solve for y_p and y_h

Find the partial solution

We will use the method called based on guessing the type of the function on the right side.

Example:

Given is the following IVP: $y' + 2y = 2x + 13$ with $y(0) = 8$

By looking carefully we see that on the right-hand side we just have a normal polynomial, thus, we can substitute $y = bx + c$ and derivate $y' = b$. In the case for when we have higher order derivative we would differentiate for each of them.

Now substitute

$$b + 2(bx + c) = 2x + 13$$

$$2bx + b + 2c = 2x + 13$$

Now by comparing the Coefficients with the other side

$$2b = 2$$

$$b + 2c = 13$$

We get: $b = 1$, $c = 6$. And we have found a *particular* solution for our DE $y_p = x + 6$

Now we have to find a *homogeneous* solution with the formula we already know $y_h = ce^{\int f(x)dx}$ in our case $y_h = ce^{\int 2dx} = ce^{-2x}$

$$y = y_h + y_p = ce^{-2x} + x + 6$$

Now considering the condition $y(0) = 8$, this gives:

$$y(0) = c + 6 = 8 \Rightarrow c = 2$$

and we obtain the particular solution:

$$y = 2e^{-2x} + x + 6$$

Table of reference

Type of Forcing Function	Disturbance Function $g(x)$	Approach for y_p
Constant	k_0	c_0
Linear	$k_0 + k_1x$	$c_0 + c_1x$
Polynomial	$\sum_{i=0}^n k_i x^i$	$\sum_{i=0}^n c_i x^i$
Exponential	ke^{bx} , $b \neq a$ ke^{ax}	$c_0 e^{bx}$ $c_0 x e^{ax}$
Trigonometric	$k \sin(bx) + l \cos(bx)$	$c_0 \sin(bx) + c_1 \cos(bx)$

It is important to point out that sometimes it is necessary to combine different types of functions to get the solution.

Example:

$$y'' - 2x'3y = t^2 + 3e^{-t} \cos(4t)$$

Here the approach would be

$$y_p = (A + Bt + Ct^2) + D(e^{-t} \cos(4t)) + E(e^{-t} \sin(4t))$$

The best way is to tackle down each of the types of functions separately.

An alternative to way of using this method is to

1. Find the homogeneous solution.
2. Multiply both sides of the equation by it.
3. Integrate

This will give you the exact same result and is faster.

66.0.21 Bernoulli Equation

This is used for equations of the form

$$y' + P(x)y = Q(x)y^n$$

We can not use the methods we already know for this equation because it is linear but with a clever substitution we can turn it into a linear differential equation.

Let us use a substitution

$$u = y^{1-n}$$

then

$$u' = (1-n)y^{-n}y'$$

Now this looks kinda similar to the original equation multiply by y^{-n} that looks like

$$y^{-n}y' + P(x)y^{-n}y = Q(x)y^{-n}yn$$

$$y^{-n}y' + P(x)y^{1-n} = Q(x)$$

Notice that $y^{-n} = \frac{u'}{(1-n)y'}$, thus, after substitution

$$\frac{1}{1-n}u' + P(x)u = Q(x)$$

Now our equation is *linear*, and we can use the *integrating factor* method.

Example:

Given is $y' - 5y = \frac{-5}{2}xy^3$.

$$u = y^{-2} \text{ and } u' = -2y^{-3}y'$$

$$y^{-3} = \frac{u'}{-2y'}$$

And our equation after multiplying by y^{-3}

$$y^{-3}y' - 5yy^{-3} = -\frac{5}{2}x$$

$$y^{-3}y' - 5y^{-2} = -\frac{5}{2}x$$

$$\frac{u'}{-2y'}y' - 5u = -\frac{5}{2}x$$

Now we can use the *Integrating Factor Method*. First bring to the standard form. $y' + p(x)y = f(x)$

$$u' + 10u = 5x$$

Remember that $r(x) = e^{\int p(x)dx} = e^{\int 10dx} = e^{10x}$ therefore, we have

$$e^{10x}u' + 10ue^{10x} = e^{10x}5x$$

Now let us integrate

$$\frac{d}{dx}(e^{10x}u) = e^{10x}5x$$

$$Ce^{10x}u = \int e^{10x}5xdx = 5x\frac{e^{10x}}{10} - \int e^{10x}5dx$$

$$Ce^{10x}u = \int e^{10x}5xdx = 5x\frac{e^{10x}}{10} - \frac{5}{10}\int e^{10x}dx$$

$$Ce^{10x}u = \frac{xe^{10x}}{2} - \frac{5}{100}e^{10x} + c$$

$$Ce^{10x} = \frac{x}{2} - \frac{1}{20} + \frac{c}{e^{10x}} = y^{-2}$$

66.0.22 Autonomous Equations

An *Autonomous Differential Equation* only depends on the dependent variable y . **Example:** $\frac{dy}{dt} = (1+y)(1-y)$.

Equilibrium Solutions

Values where $f(y) = 0$ are *Equilibrium Solutions*.

Asymptotically stable

An equilibrium solution $y = a$ is *asymptotically stable* if solutions that start near a tend towards a as $t \rightarrow \infty$.

Asymptotically unstable

An equilibrium solution $y = a$ is *asymptotically unstable* if solutions that start near a leave as $t \rightarrow \infty$.

66.0.23 Linear Inhomogeneous DEs with Non-Constant Coefficients

The method of *variation of constants* is often used alongside the *superposition principle*. In this method, the integration constant in the function $c(x)$ is omitted as it is part of the particular solution.

The superposition principle also holds for linear inhomogeneous differential equations with non-constant coefficients:

$$\begin{aligned}y'_p + f(x)y_p &= g(x) \quad (\text{particular solution}) \\y'_h + f(x)y_h &= 0 \quad (\text{homogeneous solution})\end{aligned}$$

For the total solution $y = y_h + y_p$, we get:

$$\begin{aligned}y' + f(x)y &= (y_h + y_p)' + f(x)(y_h + y_p) \\&= y'_h + y'_p + f(x)y_h + f(x)y_p \\&= g(x)\end{aligned}$$

Thus, the solution can be decomposed into the general solution of the homogeneous DE and a particular solution based on the form of the right-hand side.

Example:

$$y' = -\frac{y}{x} + 1$$

The homogeneous DE is $y' = -\frac{y}{x}$ which can be solved by separating the variables. $y_h = \frac{c}{x}$

Now let use $y_p = \frac{c(x)}{x}$.

$$y'_p = \frac{c'(x)x - c(x)}{x^2}$$

Let us break down the fraction to find where to put our homogeneous solution we have already found

$$\frac{c'(x)}{x} - \frac{c(x)}{x^2} = \frac{c'(x)}{x} + \frac{1}{x} \frac{c}{x} = \frac{c'(x)}{x} + \frac{1}{x} y$$

and we get

$$y'_p = -\frac{y}{x} + \frac{c'(x)}{x}$$

Here we compare with the inhomogeneous part of the original equation and get

$$\frac{c'(x)}{x} = 1 \implies c'(x) = x$$

then $c(x)$ is equal to $\frac{x^2}{2}$ and therefore, $y_p = \frac{x^2}{x} \frac{1}{x} = \frac{x}{2}$

As our final result we get $y = \frac{c}{x} + \frac{x}{2}$

66.0.24 Power Series Approach

Another method to solve a certain type of differential equation is to find the function given an initial conditions. This can be better understood with an example.

$$y' = y$$

$$y(0) = 1$$

We say $y = \sum_{n=0}^{\infty} a_n x^n$ and $y(0) = a_0 = 1$.

Now let us derivate you y

$$y = \sum_{n=0}^{\infty} a_n x^n = a_0 x^0 + a_1 x^1 + \dots = 1 + a_1 x^1 + a_2 x^2 + \dots$$

$$y' = 0 + a_1 + 2a_2 x + 3a_3 x^2 + \dots$$

Thus, we can rewrite the sum as

$$y' = \sum_{n=1}^{\infty} a_n n x^{n-1}$$

Starting from 1 because 0 does not contribute to the sum. After shifting the index we get

$$y' = \sum_{n=0}^{\infty} a_{n+1} (n+1) x^n$$

Now we build our original DE with our new definitions for y' and y

$$\sum_{n=0}^{\infty} a_{n+1} (n+1) x^n = \sum_{n=0}^{\infty} a_n x^n$$

If we compare the coefficients we get

$$a_n = a_{n+1} (n+1)$$

$$\frac{a_n}{(n+1)} = a_{n+1}$$

after iterating backwards we get

$$a_{n+1} = \frac{a_n}{n+1} = \frac{\frac{a_{n-1}}{(n-1+1)}}{n+1} = \frac{\frac{a_{n-2}}{\frac{(n-1)}{(n-1+1)}}}{n+1}$$

$$a_n = \frac{a_0}{(n+1)!} = \frac{1}{(n+1)!} = 1$$

therefore, $a_n = \frac{1}{n!}$ which gives us the solution

$$\sum_{n=0}^{\infty} \frac{1}{n!} x^n = e^x$$

Theorem for the Power Series of DE

For a differential equation of the form:

$$A(x)y^n + \cdots + B(x)y' + C(x)y = 0$$

more specific after dividing by $A(X)$

$$y^n + \cdots + P(x)y' + Q(x)y = 0,$$

with $x = a$ as an *Ordinary Point* if $\dots, P(x)$ and $Q(x)$ are *Analytic* at $x = a$. Otherwise, a *Singular Point*.

Then this equation has n linearly independent solutions of the form

$$y(x) = \sum_{n=0}^{\infty} c_n (x - a)^n$$

The radius of convergence is at least as large as distance to the nearest *Singular Point*.

66.0.25 Exact Differential Equations

We will take a short look at functions with two inputs via implicit differentiation $F(x, y(x)) = 0$. With $\frac{\partial F}{\partial x} = P$ and $\frac{\partial F}{\partial y} = Q$.

$$P(x, y)dx + Q(x, y)dy = 0$$

after dividing by dx

$$P(x, y) + Q(x, y)y' = 0$$

We define this kind of differential equations as *exact*. And the $P_y = Q_x$ as *Integrability-Condition*.

By finding the *potential* we can solve this kind of differential equations.

Example:

Step 1: Solve for y'

$$\begin{aligned}(12xy + 3)dx + 6x^2dy &= 0 \\ (12xy + 3) + 6x^2y' &= 0 \\ y' &= -\frac{12xy + 3}{6x^2}\end{aligned}$$

Step 2: Test the Integrability-Condition

$$\begin{aligned}P(x, y) &= 12xy + 3 \\ Q(x, y) &= 6x^2 \\ P_y &= 12x\end{aligned}$$

$$Q_x = 12x$$

Step 3: Integrate P with respect to x and add $h(y)$

$$\begin{aligned} F(x, y) &= \int P(x, y) dx = \int (12xy + 3) dx \\ &= 6x^2y + 3x + h(y) \end{aligned}$$

Step 4: Differentiate F with respect to y and set equal to Q

$$\frac{\partial}{\partial y} (6x^2y + 3x + h(y)) = 6x^2 + c'(y)$$

Step 5: Integrate with respect to y

$$\begin{aligned} 6x^2 + c'(y) &= 6x^2 \\ c'(y) &= 0 \\ \int c'(y) &= \int 0 = C \end{aligned}$$

But $C = 0$ because, C in Q is zero. Therefore, $F(x, y) = 6x^2y + 3x$.

Euler Multipliers

Sometimes a seemingly *exact* differential equations is not possible to solve do the partial derivatives of $P(x, y)$ and $Q(x, y)$ not being equal. An approach to this problem are the so-called *Euler Multipliers*, which are some factor $\mu(x)$ or $\mu(y)$ that when found. Can help solve the differential equation.

$$P(x, y)\mu(x) + Q(x, y)y'\mu(x) = 0$$

or

$$P(x, y)\mu(y) + Q(x, y)y'\mu(y) = 0$$

The steps for solving this kind of problems are

1. Choose either $\mu(x)$ or $\mu(y)$ and multiply the whole equation by it.
2. Find the new partial derivatives of P and Q .
3. Set both partial equals and solve for μ in the newly created differential equation using the necessary method. Also make sure that the solution has no zeroes. This is necessary because otherwise, we would have multiplied our original equation times zero.
4. Substitute μ with the solution. And if now both partials are equal then you can now solve the exact differential equation. If not try with another μ using the other variable. And if this also does not work then the differential equation can not be solved. More specific, if $\mu(x)$ depends on y and $\mu(y)$ on x then there is no way to make the differential equation exact.

Example:

$$3y^2 + 2xyy' = 0, x > 0$$

Here we have $P(x, y) = 3y^2$ and $Q(x, y) = 2xy$, whose partial derivatives are different

$$\frac{\partial P}{\partial y} = 6y \neq 2y = \frac{\partial Q}{\partial x}$$

Now we try to use a multiplier $\mu(x)$.

$$P_2(x, y) = 3y^2\mu(x) \text{ and } Q_2(x, y) = 2xy\mu(x)$$

Now by taking the partials again

$$\begin{aligned}\frac{\partial P_2}{\partial y} &= 6y\mu(x) \\ \frac{\partial Q_2}{\partial x} &= 2y(\mu(x) + x\mu'(x))\end{aligned}$$

Now by setting them equal $6y\mu(x) = 2y(\mu(x) + x\mu'(x))$ we can solve a new differential equation to find $\mu(x)$ and $\mu'(x)$.

$$\begin{aligned}6y\mu(x) &= 2y(\mu(x) + x\mu'(x)) \\ 3\mu(x) &= \mu(x) + x\mu'(x) \\ 2\mu(x) &= x\mu'(x) \\ 2\mu &= x \frac{d\mu}{dx} \\ \int \frac{2}{x} dx &= \int \frac{d\mu}{\mu} \\ 2 \ln(x) &= \ln(\mu) \\ \mu &= x^2\end{aligned}$$

Now finally we check if now the partials are equal, which in this case is true.

$$\begin{aligned}\frac{\partial P_2}{\partial y} &= 6yx^2 \\ \frac{\partial Q_2}{\partial x} &= 6yx^2\end{aligned}$$

Now the equation can be solved like normal.

66.0.26 Constant Coefficients ODE

Consider the second-order linear differential equation with constant coefficients:

$$ay'' + by' + cy = 0,$$

where $a, b, c \in \mathbb{R}$, and $a \neq 0$.

We look for solutions of the form $y = e^{rt}$. Substituting into the equation:

$$\begin{aligned}y &= e^{rt}, \\ y' &= re^{rt} \\ y'' &= r^2e^{rt}\end{aligned}$$

Substituting into the original equation:

$$ar^2e^{rt} + bre^{rt} + ce^{rt} = 0$$

Factoring out e^{rt} (which is never zero):

$$e^{rt}(ar^2 + br + c) = 0 \quad \Rightarrow \quad ar^2 + br + c = 0.$$

This is a quadratic equation in r . Solving using the quadratic formula:

$$r = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Suppose there are two linearly independent solution y_1 and y_2 to

$$y'' + p(x)y' + q(x)y = 0,$$

then the general solution is $y = c_1y_1 + c_2y_2$.

We analyze the nature of the roots r based on the discriminant $D = b^2 - 4ac$. Also, for dealing with initial conditions of the form $y^n = a, \dots$ we only have to plug the values into our solutions $y_h, y'_h, \dots$ to get the constants.

Case 1: Two Distinct Real Roots ($D > 0$)

There not much not to just plug the solutions of the root.

$$y = c_1e^{r_1t} + c_2e^{r_2t}$$

Case 2: Repeated Real Root ($D = 0$)

For this case we need to use a trick.

$$y = c_1e^{rt} + c_2te^{rt}$$

Note that this could be contra intuitive but te^{rt} is a solution and is linearly independent. For more repeated roots you can use higher powers of t $t, t^2, t^3, \dots$

Case 3: Complex Conjugate Roots ($D < 0$)

When the discriminant $D = b^2 - 4ac < 0$, the roots of the characteristic equation are complex:

$$r = \alpha \pm i\beta,$$

where $\alpha = -\frac{b}{2a}$ and $\beta = \frac{\sqrt{4ac-b^2}}{2a}$.

The general solution to the differential equation is:

$$y(t) = c_1e^{(\alpha+i\beta)t} + c_2e^{(\alpha-i\beta)t}.$$

To express this in real form, we use Euler's formula:

$$e^{i\beta t} = \cos(\beta t) + i\sin(\beta t).$$

Now rewrite each exponential term:

$$e^{(\alpha+i\beta)t} = e^{\alpha t} \cdot e^{i\beta t} = e^{\alpha t}(\cos(\beta t) + i \sin(\beta t)),$$

$$e^{(\alpha-i\beta)t} = e^{\alpha t} \cdot e^{-i\beta t} = e^{\alpha t}(\cos(\beta t) - i \sin(\beta t)).$$

Now to get rid of the imaginary part we can do the following trick

$$\frac{y_1 + y_2}{2} = e^{\alpha t} \cos \beta t$$

$$\frac{y_1 - y_2}{2i} = e^{\alpha t} \sin \beta t$$

This might feel wrong, but these are in fact linear combinations of our original solution, and they are also linearly independent, because of that we can just use them for our general solution.

$$y = c_1 e^{\alpha t} \cos \beta t + c_2 e^{\alpha t} \sin \beta t$$

Example:

Given is $y''' + y' = 0$.

$$y = e^{rt}$$

$$r^3 + r = 0$$

$$r(r^2 + 1) = 0$$

$$r(r + i)(r - i) = 0$$

Now $r_1 = 0$ $r_2 = -i$ $r_3 = i$

$$y = c_1 e^{0t} + c_2 e^{-it} + c_3 e^{it}$$

$$y = c_1 + c_2 \cos t + c_3 \sin t$$

Inhomogeneous Case

For the inhomogeneous case we would apply the method of the *Undetermined Coefficients*. But differentiating two times or more depending on the order of the differential equation and add the homogeneous solution to the solution we get from that.

Example:

Solve the differential equation:

$$y' + \frac{1}{x}y = \sin(x)$$

The integrating factor is:

$$\mu(x) = e^{\int \frac{1}{x} dx} = e^{\ln |x|} = |x|$$

Assuming $x > 0$, we take $\mu(x) = x$. Multiply through by the integrating factor:

$$xy' + y = x \sin(x)$$

The left-hand side is the derivative of xy :

$$\frac{d}{dx}(xy) = x \sin(x)$$

$$\int \frac{d}{dx}(xy) dx = \int x \sin(x) dx$$

Use integration by parts on the right:

$$\int x \sin(x) dx = -x \cos(x) + \int \cos(x) dx = -x \cos(x) + \sin(x)$$

So:

$$xy = -x \cos(x) + \sin(x) + C$$

$$y = -\cos(x) + \frac{\sin(x)}{x} + \frac{C}{x}$$

66.0.27 The Wronskian

Let $y_1, y_2, \dots, y_n$ be n functions that are at least $(n-1)$ -times differentiable on some interval. The *Wronskian* of these functions is defined as:

$$W(y_1, y_2, \dots, y_n)(x) = \begin{vmatrix} y_1(x) & y_2(x) & \cdots & y_n(x) \\ y_1'(x) & y_2'(x) & \cdots & y_n'(x) \\ \vdots & \vdots & \ddots & \vdots \\ y_1^{(n-1)}(x) & y_2^{(n-1)}(x) & \cdots & y_n^{(n-1)}(x) \end{vmatrix}$$

If $W(A) \neq 0$ then they are linearly independent.

66.0.28 Linearity Property

Let $y_p^{(1)}$ be a particular solution of

$$y'' + ay' + by = g_1(x)$$

and $y_p^{(2)}$ a particular solution of

$$y'' + ay' + by = g_2(x),$$

then

$$y_p = y_p^{(1)} + y_p^{(2)}$$

is a particular solution of

$$y'' + ay' + by = g_1(x) + g_2(x).$$

66.0.29 Variation of Parameters

For a linear differential equation of the form

$$y'' + p(x)y' + q(x)y = f(x)$$

The homogeneous equation has a solution of the linearly independent y_1 and y_2 of the form $y = u_1y_1 + u_2y_2$. We can guess the functions y_1 and y_2 .

So, for finding a solution of these types of equations we use the following formulas.

$$u_1 = - \int \frac{y_2 g}{y_1 y_2' - y_2 y_1'} dx$$

$$u_2 = - \int \frac{y_1 g}{y_1 y_2' - y_2 y_1'} dx$$

These formulas come from:

Differentiating two times:

$$y = u_1 y_1 + u_2 y_2$$

$$y' = u_1 y_1' + y_1 u_1' + u_2 y_2' + y_2 u_2'$$

Here we can add another constraint that $y_1 u_1' + y_2 u_2' = 0$. This allows us to simplify the expression to:

$$y' = u_1 y_1' + u_2 y_2'$$

Now we can derivate one more time:

$$y'' = u_1 y_1'' + y_1' u_1' + u_2 y_2'' + y_2' u_2'$$

Substitute in the original equation

$$y'' + p(x)y' + q(x)y = f(x)$$

$$u_1 y_1'' + y_1' u_1' + u_2 y_2'' + y_2' u_2' + p(x)(u_1 y_1' + u_2 y_2') + q(x)(u_1 y_1 + u_2 y_2) = g(x)$$

$$u_1 y_1'' + y_1' u_1' + u_2 y_2'' + y_2' u_2' + p(x)(u_1 y_1') + p(x)(u_2 y_2') + q(x)(u_1 y_1) + q(x)(u_2 y_2) = g(x)$$

$$u_1 y_1'' + y_1' u_1' + u_2 y_2'' + y_2' u_2' + p(x)(u_1 y_1') + p(x)(u_2 y_2') + q(x)(u_1 y_1) + q(x)(u_2 y_2) = g(x)$$

With non-generic function a lot of the terms would cancel out. After that we add multiply by y_1 one of the rows and y_2 so that we can cancel them in a row operation. Which will left us with only two integrals to solve.

66.0.30 Which method to use depending on the type of differential equations

First Order

- Separable $\Rightarrow$ Separation of Variables.
- Form $y' = f(ax + by(x) + c) \Rightarrow$ Substitution $z = ax + by(x) + c$ and $f(z)$.
- Form $y' = \frac{y}{x} \Rightarrow$ Substitution $z = \frac{y}{x}$
- Linear $\Rightarrow$ Integrating Factor or Variation of Constants.
- Bernoulli (Non-linear) $\Rightarrow$ Bernoulli Substitution.
- Autonomous $\Rightarrow$ Equilibrium Analysis.
- Non-homogeneous $\Rightarrow$ Undetermined Coefficients or integrating factor.

Second Order and higher

- Constant Coefficients $\Rightarrow$ Guess e^{rx} .
- Non-homogeneous $\Rightarrow$ Undetermined Coefficients.
- Linear $\Rightarrow$ Series Solutions.

Exact

Using the method for exact differential equations.

66.0.31 Linear Systems of Differential Equations

Let us start by looking at a system of first order homogeneous differential equations

$$\begin{aligned}y_1' &= ay_1 + by_2 \\ y_2' &= cy_1 + dy_2\end{aligned}$$

This can be written as

$$\frac{d}{dt} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

$$Ax = \vec{y}'$$

Then if we use the guess $y(t) = \vec{v}e^{\lambda t}$ and differentiate with respect to t we get

$$\lambda \vec{v}e^{\lambda t} = A\vec{v}e^{\lambda t}$$

$$\lambda \vec{v} = A\vec{v},$$

which is our *Eigenvalue* problem. This can be solved via

$$\det(A - \lambda I) = 0$$

$$\lambda^2 - (a + c)\lambda + ad - bc = 0$$

And this solution can lead to the three cases we already know: distinct real roots, complex Conjugate roots and repeated roots.

We can also use substitution to solve for one of the y_i , which are functions of x like

$$\begin{aligned}
y_1' &= ay_1 + by_2 \\
y_2' &= cy_1 + dy_2 \\
by_2 &= y_1' - ay_1 \\
y_2 &= \frac{y_1' - ay_1}{b}
\end{aligned}$$

Then we can differentiate our first equation with respect to x

$$\begin{aligned}
by_2 &= y_1' - ay_1 \\
by_2' &= y_1'' - ay_1' \\
&= b(cy_2 + dy_1) \\
&= bcy_2 + bdy_1 \\
&= c(y_1' - ay_1) + bdy_1 \\
y_1'' - (a + c)y_1' + (ac - bd)y_1 + bdy_1 &= 0
\end{aligned}$$

Example:

We are given the system of differential equations:

$$\begin{cases} y' = y + z \\ z' = -2y + 3z \end{cases}$$

We write this system in matrix form:

$$u' = Au, \quad \text{where } u = \begin{bmatrix} y \\ z \end{bmatrix}, \quad A = \begin{bmatrix} 1 & 1 \\ -2 & 3 \end{bmatrix}$$

To solve, we find the eigenvalues of A by solving the characteristic equation:

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ -2 & 3 - \lambda \end{vmatrix} = (1 - \lambda)(3 - \lambda) + 2 = \lambda^2 - 4\lambda + 5 = 0$$

Solving the quadratic equation:

$$\lambda = \frac{4 \pm \sqrt{(-4)^2 - 4(1)(5)}}{2} = \frac{4 \pm \sqrt{-4}}{2} = \frac{4 \pm 2i}{2} = 2 \pm i$$

Now we find an eigenvector corresponding to $\lambda = 2 + i$ by solving:

$$(A - (2 + i)I)v = 0$$

$$A - (2 + i)I = \begin{bmatrix} 1 - (2 + i) & 1 \\ -2 & 3 - (2 + i) \end{bmatrix} = \begin{bmatrix} -1 - i & 1 \\ -2 & 1 - i \end{bmatrix}$$

From the first row of the system:

$$(-1 - i)v_1 + v_2 = 0 \Rightarrow v_2 = (1 + i)v_1$$

So, an eigenvector corresponding to $\lambda = 2 + i$ is:

$$v = \begin{bmatrix} 1 \\ 1+i \end{bmatrix}$$

After repeating the previous computation for $\lambda = 2 - i$ we get the complex solution to the system:

$$u(t) = c_1 e^{(2+i)t} \begin{bmatrix} 1 \\ 1+i \end{bmatrix} + c_2 e^{(2-i)t} \begin{bmatrix} 1 \\ 1-i \end{bmatrix}$$

Chapter 67

Combinatorics

67.0.1 Three Principles of Combinatorics

These principles are *Addition*, *Multiplication* and *Inclusion-Exclusion*.

67.0.2 The Basic Counting Principle

For n outcomes of event A and m of B there are $n \cdot m$ outcomes for event A and B .

In combinatorics we can model a lot of problems as having n elements from which we can take one at the time a certain number of times, and then we take into consideration if we allow repetition or the order matters or not.

67.0.3 Permutation

Number of ways of ordering n distinct elements. In other words the number of tuples of n distinct elements. Of course the order matters.

$$n!$$

Example:

67.0.4 Permutation with repetition

Number of ways of ordering n non-distinct elements. Here $m_1, \dots, m_n$ is the number a specific item is repeated in the original set. By dividing by the respective amount of the repeated non-distinct element, we eliminate the permutations, where the identical element swap their place.

$$\frac{n!}{m_1! \cdots m_n!}$$

This formula comes from the *multinomial coefficient* given by

$$\binom{n}{n_1 n_2 \dots n_m} = \frac{n!}{n_1! \cdots n_m!},$$

which comes from

$$\binom{n}{n_1} \binom{n - n_1}{n_2} \binom{n - n_1 - n_2}{n_3} \cdots \binom{n - n_1 - \cdots - n_{m-1}}{n_m}.$$

This is also a kind of *partition problem*, because it can be formulated as if you have n different elements, and you want to divide them into m bins with capacities $m_1, m_2, \dots, m_m$ how many possible ways are there for these partitions.

In this case the order matters but only for the different elements.

Example:

67.0.5 Combination with repetition

We can think of it as having n types of objects, and we want to take with repetition until we fill our k slots.

$$\binom{n+k-1}{n-1}$$

For understanding, we have n types of objects and because we want a combination of those objects the order does not matter. Thus, we can put them together as contiguous groups. We have k slots and n types thus we can separate our groups with $n-1$ separators $|$. The question becomes how many ways are there to put $n-1$ separator into $n+k-1$ slots. Note that we have $n+k-1$ positions if we count the actual slots and the separator slots in between.

$$*|***|**, \quad n=6, \quad k=2, \quad \text{positions in between} = 7$$

Example:

67.0.6 Variation

Ways of put n objects in k slots with repetition. Here the order does not matter.

$$n^k$$

Example:

67.0.7 Variation without repetition

Ways of put n objects in k slots without repetition. Number of tuples with k elements which can be selected from n elements, without repetition. The order matters.

$$\frac{n!}{(n-k)!}$$

Example:

67.0.8 Combination without repetition

Ways of choose k objects of n elements. Or put k elements from n into k slots where the order does not matter.

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Note that like in the variation we use $\frac{n!}{(n-k)!}$ to limit our slots to k . Then for the k objects we choose we can arrange them in $k!$ ways but because the order does not matter we divide by $k!$ to get rid of the repeated combinations

Example:

67.0.9 Multinomial Coefficients

When using the binomial coefficient repeatedly one after the other we can abbreviate this consecutive uses as

$$\binom{n}{k_1, \dots, k_n}$$

This is like from n choose k_1 then from $n - k_1$ choose k_2 and so on.

Multinomial Theorem

Given $x_1, \dots, x_p \in \mathbb{R}$ and $\mathbb{N}$ then

$$(x_1 + \dots + x_p)^n = \sum_{k_1 + \dots + k_p = n} \binom{n}{k_1, \dots, k_p} x_1^{k_1} \dots x_p^{k_p}$$

67.0.10 Example Problems

Defected Machines

A new delivery with 10 machines from which 2 are not working properly have to be tested. For that purpose a man in charge selects 5 of the machines randomly, and then he proceeds to make a decision based on the number of faulty machines found in the delivery. If among the 5 samples only 1 machine does not work it will be accepted otherwise it will be rejected. What is the probability of a delivery being accepted?

Answer:

Total cases: $\binom{10}{5}$

$$\mathbb{P}(\text{accepted}) = \frac{\binom{8}{5}}{\binom{10}{5}} + \frac{\binom{2}{1}\binom{8}{4}}{\binom{10}{5}}$$

$\binom{8}{5}$ represents no faulty machine in the choice, $\binom{2}{1}\binom{8}{4}$ represents only one machine does not work.

67.0.11 Disarray

Number of permutation in which no object ends in the same initial spot. Also, known as the *sub-factorial*.

$$!n = n! \sum_{k=0}^n \left(\frac{(-1)^k}{k!} \right)$$

67.0.12 Bell Numbers

Number of ways of grouping n objects in an arbitrary number of slots.

$$B(N) = \sum_{k=0}^{N-1} \binom{N}{k} B(K),$$

with $B(1) = 1$ and $B(2) = 2$.

67.0.13 Ramanujan Numbers

The same as the Bell Numbers but, all objects are equal. As an example the number of ways to decompose a number.

$$R(N) \approx \frac{1}{4N\sqrt{3}} e^{2\pi\sqrt{\frac{N}{6}}}$$

67.0.14 Stirling Numbers I

Ways of permuting N items with K exchanges.

$$\begin{bmatrix} N+1 \\ K \end{bmatrix} = N \begin{bmatrix} N \\ K \end{bmatrix} + \begin{bmatrix} N \\ K-1 \end{bmatrix}$$

67.0.15 Stirling Numbers II

Ways of grouping n items in k slots.

$$\left\{ \begin{matrix} N \\ K \end{matrix} \right\} = \frac{1}{K!} \sum_{j=0}^K (-1)^{K-j} \binom{K}{j} j^N$$

67.0.16 Lah Numbers

Ways of building K lists with a set N elements.

$$\begin{bmatrix} N \\ K \end{bmatrix} = \binom{N-1}{K-1} \frac{N!}{K!} = \sum_{j=0}^K \begin{bmatrix} N \\ K \end{bmatrix} \left\{ \begin{matrix} N \\ j \end{matrix} \right\}$$

67.0.17 Euler's Numbers

Number of permutations of a set of N elements of different size in which K elements are bigger than their previous element.

$$\left\langle \begin{matrix} N \\ K \end{matrix} \right\rangle = \sum_{j=0}^K (-1)^j \binom{N+1}{j} (K+1-j)^N$$

67.0.18 Catalan Numbers

These have different applications like the number of ways of the triangulations of a polygon or the number of trees with N leaves.

$$\mathfrak{C}(N) = \binom{2N}{N} - \binom{2N}{N+1}$$

or

$$\mathfrak{C}(N) = \sum_{k=1}^n (k-1) \mathfrak{C}(N-k).$$

Chapter 68

Probability

68.0.1 Basics

A probability experiment is considered ideal if and only the following properties:

- We know the possible outcomes.
- It can be repeated multiple times under the same conditions.
- It will be executed under strict conditions.

68.0.2 The Birthday Paradox

This is a small question that says: What is the probability of at least two people having their birthday on the same day in a group of x persons.

$$\mathbb{P}(x) = 1 - \prod_{n=0}^{x-1} \frac{365 - n}{365}$$

This formula gives us the probability of total minus all persons having it on different days. Which what we are looking for.

68.0.3 The Sample Space

The set of all possible outcomes of an experiment is denoted as Ω and its cardinality is the number of possible outcomes.

For experiments with Ω as the result set of one repetition then we define

$$\Omega = \Omega \times \cdots \times \Omega$$

As the *sample space* of the experiment.

Finally, each subset of Ω is called an *outcome*, each subset with only one element in it is called an *elemental outcome*. The set of all possible outcomes is $\mathcal{P}(\Omega)$

68.0.4 Laplace Distribution and Experiment

A *Laplace distribution* is a probability experiment with finite outcomes where each elemental outcome has the same probability of occurring.

$$\mathbb{P}(\omega) = \frac{\omega}{|\Omega|}$$

Then an experiment with this property is considered a *Laplace experiment*.

$$\mathbb{P}(E) = \frac{E}{|\Omega|}$$

68.0.5 Sigma Algebra

Given a non-empty set Ω and $\mathcal{A} \subseteq \mathcal{P}(\Omega)$ a set of subsets of Ω . $\mathcal{A}$ is a σ -Algebra if and only if

- I. $\Omega \in \mathcal{A}$
- II. $A \in \mathcal{A} \implies A^c \in \mathcal{A}$
- III. $A_1, A_2, A_3, \dots \in \mathcal{A} \implies \bigcup_{n \in \mathbb{N}} A_n \in \mathcal{A}$

The Power-set is a Sigma Algebra

This proof is pretty straight forward.

For the first axiom, we know that $\Omega \in \mathcal{P}$.

For the second axiom, if A is a subset of the power-set then $\Omega \setminus A$ is also in the power-set.

And lastly, for a sequence of sets in the power-set their union is also in the power-set, because the power-set contains all subsets of Ω .

□□□

Properties of Sigma Algebras

If $\mathcal{A}$ is a sigma algebra then

- $\emptyset \in \mathcal{A}$
- $A_1, A_2, \dots \in \mathcal{A} \implies \bigcap_{n \in \mathbb{N}} A_n \in \mathcal{A}$
- $A, B \in \mathcal{A} \implies A \setminus B \in \mathcal{A}$

Proof:

For the first, because $\mathcal{A}$ is a subset of the power-set of Ω and the empty set is always a subset of a collection, this is trivial.

Given a number of sets in $\mathcal{A}$, using the laws of De Morgan we get

$$\bigcap_{n \in \mathbb{N}} A_n = \bigcup_{n \in \mathbb{N}} A_n^c$$

and we know that the complements are part of $\mathcal{A}$ by definition.

For the last part, $A \setminus B = A \cap B^c$, because $B^c \in \mathcal{A}$ we can apply De Morgan $(A^c \cup (B^c)^c)^c = (A^c \cup B)^c$. Now because this is the union of two sets in the sigma algebra, and we know by definition that their union is also in the set.

□□□

68.0.6 Probability Measurement

Given a non-empty set Ω and $\mathcal{A} \subseteq \mathcal{P}$ a sigma algebra. We define the function $\mathbb{P} : \mathcal{A} \rightarrow \mathbb{R}$ as a *probability measurement* if

- I. $\forall A \in \mathcal{A} : \mathbb{P}(A) \geq 0$
- II. $\mathbb{P}(\Omega) = 1$
- III. $A_n \in \mathcal{A}, n \in \mathbb{N}$ disjunctive in pairs $\implies \mathbb{P}(\bigcup_{n \in \mathbb{N}} A_n) = \sum_{n \in \mathbb{N}} \mathbb{P}(A_n)$

Also, the triplet $\Omega, \mathcal{A}, \mathbb{P}$ is called a *probability space*.

Probability space of the Power-set

We want to prove that the probability space with the power-set of Ω and $\mathbb{P}(A) = \frac{|A|}{|\Omega|}$ is in fact a probability space

- $\mathbb{P}(A) = \frac{|A|}{|\Omega|}$ is always greater or equal to zero.
- $\mathbb{P}(\Omega) = \frac{|\Omega|}{|\Omega|} = 1$
- If all of our subsets in the power-set are pairwise disjunctive this mean that their intersection is the empty-set. In other wise there therefore, their independent of each other which means that their probabilities are also independent.

Discrete Probability Spaces

A probability space $\Omega, \mathcal{P}(\Omega), \mathbb{P}$ is considered *discrete* if Ω is finite or infinite countable. In that case $\Omega = \{\omega_i : i \in I\}$ for $I = \{1, \dots, n\}$

For $A \in \mathcal{A}$

$$\mathbb{P}(A) = \mathbb{P}\left(\bigcup_{i \in I: \omega_i \in A} \{\omega_i\}\right) = \sum_{i \in I: \omega_i \in A} \mathbb{P}(\{\omega_i\}) = \sum_{i \in I: \omega_i \in A} p(\omega_i)$$

The function $\mathbb{P}(\{\omega_i\}) = p(\omega_i)$ is called a *probability function*

To test if specific function or distribution is actually a probability one we have to check.

- $\forall p(\omega) \geq 0 \mid \omega \in \Omega$
- $\sum_{\omega \in \Omega} p(\omega) = 1$

Properties of Probability-Measurements

Given $(\Omega, \mathcal{A}, \mathbb{P})$ and $A, B \in \mathcal{A}$

- $\mathbb{P}(\emptyset) = 0$
- $A_1, \dots, A_n \in \mathcal{A}$ pairwise disjoint: $\mathbb{P}(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n \mathbb{P}(A_i)$
- $\mathbb{P}(B \setminus A) = \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$
- $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$
- If $A \subset B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$
- $0 \leq \mathbb{P}(A) \leq 1$
- $A_1, \dots, A_n \in \mathcal{A}$: $\mathbb{P}(\bigcup_{i=1}^n A_i) \leq \sum_{i=1}^n \mathbb{P}(A_i)$
- $A_1, A_2, \dots \in \mathcal{A}$: $\mathbb{P}(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \mathbb{P}(A_i)$

Proof:

- Define $A_n = \emptyset$ and by the properties of the probability spaces we know that $A_n \in \mathcal{A}, n \in \mathbb{N}$ disjunctive in pairs $\implies \mathbb{P}(\bigcup_{n \in \mathbb{N}} A_n) = \sum_{n \in \mathbb{N}} \mathbb{P}(A_n)$

Therefore, $\mathbb{P}(\emptyset) = \sum_{i=1}^n \mathbb{P}(A_i)$ and the only real number with the property $x + x = x$ is 0.

- By definition
- $B = (A \cap B) \cup B \setminus A$ and by the definition of the addition of probabilities

$$\mathbb{P}(B) = \mathbb{P}(B \setminus A) + \mathbb{P}(A \cap B)$$

- Using the addition property and by writing the union as $(A \setminus B) \cup (B \setminus A) \cup (A \cap B)$ we get that

$$\mathbb{P}(A \cup B) = \mathbb{P}(B) - \mathbb{P}(A \cap B) + \mathbb{P}(A) - \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B)$$

which results in $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

- $\Omega = A^c \cup A$ and $1 = \mathbb{P}(\Omega)$, thus by the addition $\mathbb{P}(A) + \mathbb{P}(A^c) = 1$
- Given that if $A = B$ then the equality holds the case where A is a subset of B comes from the fact that it is a strict subset of the other than its cardinality is smaller.
- By the definition that $\mathbb{P}(\Omega) = 1$ and the property $\mathbb{P}(\emptyset) = 0$ we are given an upper and lower bound for our probability distribution. Therefore, every subset of Ω will have a probability less or equal to it. For the lower bound the logic is that it has no other subset with fewer elements than it, but it is part of sets with more elements.
- Let us define $B_n = A_n \setminus \bigcap_{i=1}^{n-1} A_i$ which are pairwise disjoint.

$$\mathbb{P}\left(\bigcup A_n\right) = \mathbb{P}\left(\bigcup B_n\right) = \sum \mathbb{P}(B_i) \leq \sum \mathbb{P}(A_i)$$

Due to the fact that the B_i are pairwise disjoint.

- This is practically the same but for an infinite case.

□◻◻

68.0.7 Inclusion-Exclusion Formula for Two Non-Disjoint Sets

Let $|A \cap B| = k$, then:

$$|A \cup B| = |A| + |B| - |A \cap B| = n + m - k \quad (\text{since we do not count the intersection twice})$$

68.0.8 Inclusion-Exclusion Formula for Three Non-Disjoint Sets

$$\begin{aligned} |A \cup B \cup C| &= |(A \cup B) \cup C| \\ &= |A \cup B| + |C| - |(A \cup B) \cap C| \\ &= |A| + |B| - |A \cap B| + |C| - |(A \cap C) \cup (B \cap C)| \\ &= |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C| \end{aligned}$$

68.0.9 General Formula for the Cardinality of the union of sets

$$\left| \bigcup_{i=1}^n M_i \right| = \sum_{I \subseteq \{1, \dots, n\}, I \neq \emptyset} (-1)^{|I|-1} \left| \bigcap_{i \in I} M_i \right|$$

68.0.10 Stochastic Independence

Two results $A, B \in \mathcal{A}$ are considered *independent* if

$$\mathbb{P}(A \mid B) = \mathbb{P}(A)$$

$$\mathbb{P}(B \mid A) = \mathbb{P}(B)$$

then

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$$

Example:

When throwing a die two times is the probability of throwing during the first throw an even number and when throwing the second die the sum gives us an even number.

$$\Omega = \{1, \dots, 6\}^2 \quad A = \{2, 4, 6\} \times \{1, \dots, 6\} \quad B = \{2, 4, 6\} \cup \{1, 3, 5\}^2$$

Therefore,

$$\mathbb{P}(A) = \frac{|A|}{|\Omega|} = \frac{18}{36} = \frac{1}{2}$$

$$\mathbb{P}(B) = \frac{|B|}{|\Omega|} = \frac{18}{36} = \frac{1}{2}$$

And

$$\mathbb{P}(A \cap B) = \frac{|\{2, 4, 6\}^2|}{|\Omega|} = \frac{1}{4}$$

This is just $\mathbb{P}(A) \cdot \mathbb{P}(B)$. Therefore, they are independent.

□

Independence for multiple results

$$\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2) \cdot \dots \cdot \mathbb{P}(A_n)$$

For all pairs A_i, A_j where $i \leq j$

Independence in Laplace Distribution

Give a Laplace distribution $\mathbb{P} = \frac{|A|}{|\Omega|}$ with two independent results A and B .

$$\frac{|A \cap B|}{|A|} = \frac{|B|}{|\Omega|}$$

This means that B is a probable in A as in the greater picture.

Complement Property

Given the probability space $(\Omega, \mathcal{A}, \mathbb{P})$ and $A, B \in \mathcal{A}$ as two independent results. Then A^c and B are also independent.

Proof:

The sets $A \cap B$ and $A^c \cap B$ are disjoint and their union is

$$B = (A \cap B) \cup (A^c \cap B)$$

If we calculate the probability

$$\mathbb{P}(B) = \mathbb{P}(A \cap B) + \mathbb{P}(A^c \cap B)$$

This gives us

$$\mathbb{P}(A^c \cap B) = \mathbb{P}(B) - \mathbb{P}(A \cap B)$$

Using the Independence of A and B

$$\mathbb{P}(A^c \cap B) = \mathbb{P}(B) - \mathbb{P}(A)\mathbb{P}(B) = \mathbb{P}(B)(1 - \mathbb{P}(A))$$

$$\mathbb{P}(B) = \mathbb{P}(B)(1 - \mathbb{P}(A)) = \mathbb{P}(B) \cdot \mathbb{P}(A^c)$$

□♦♦

Independence of Blocks

Given $A_1, \dots, A_n$ independent events, divided into two blocks $A_1, \dots, A_k$ and $A_{k+1}, \dots, A_n$. Let B and C be events resulted of the union, intersection or complement of the first two blocks. Then this results are independent

68.0.11 The Product Space

Given two or more probability spaces with their respective tuples of two independent experiments. We define their combined probability space as the *product space* where

$$\Omega := \Omega_1 \times \dots \times \Omega_n$$

and

$$p(\omega_1, \dots, \omega_n) = p(\omega_1) \cdot \dots \cdot p(\omega_n)$$

An example is throwing two dices at the same time.

68.0.12 The Product Space of Discrete Probability Spaces

Given the product space $(\Omega, \mathcal{P}(\Omega), \mathbb{P})$ of the spaces $(\Omega_i, \mathcal{P}(\Omega_i), \mathbb{P}_i)$ and $A_i \in \mathcal{P}(\Omega_i)$. We define

$$\begin{aligned} A'_i &:= \{\omega \in \Omega : \omega_i \in A_i\} \\ &= \{(w_1, \dots, w_n) : w_j \in \Omega_j, \text{ for } j \neq i \text{ and } w_i \in A_i\} \\ &= \Omega_1 \times \dots \times \Omega_{i-1} \times A_i \times \Omega_{i+1} \times \dots \times \Omega_n \end{aligned}$$

For $i = 1, \dots, n$. Then it holds that $\mathbb{P}(A'_i) = \mathbb{P}_i(A_i)$ and the events A'_i are independent.

68.0.13 Bernoulli-Experiments

An experiment where $\Omega = \{0, 1\}$, $\mathbb{P}(\{1\}) = p$ and $\mathbb{P}(\{0\}) = 1 - p$ for $p \in (0, 1)$ is called a *Bernoulli-Experiment*.

Bernoulli Schema

If we perform a Bernoulli-Experiment n -times with a success probability of p . Where $\mathbb{P}(A_i) = p$. We define

$$C_k = \bigcup_{I \in \{1, \dots, n\} \wedge |I|=k} \left(\bigcap_{i \in I} A_i \cap \bigcap_{i \notin I} A_i^c \right)$$

For the probability we get

$$\mathbb{P}(C_k) = \sum_{I \in \{1, \dots, n\} \wedge |I|=k} \mathbb{P} \left(\bigcap_{i \in I} A_i \cap \bigcap_{i \notin I} A_i^c \right) = \sum_{I \in \{1, \dots, n\} \wedge |I|=k} \prod_{i \in I} \mathbb{P}(A_i) = \prod_{i \notin I} \mathbb{P}(A_i^c) = \binom{n}{k} p^k (1-p)^{n-k}$$

68.0.14 Conditional Probability

Given the events A and B where $\mathbb{P}(A) > 0$. The *conditional probability* of B under the condition A is defined as

$$\mathbb{P}(B \mid A) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(A)}$$

Example:

In words, what is the probability of B if we limit ourself only to the case where A happens. Also what is the ratio between $A \cap B$ and A . From this definition we can also derive these two statements which are also equivalent.

$$\mathbb{P}(B \mid A) = \mathbb{P}(B) \iff A \text{ and } B \text{ are independent}$$

Proof:

$$\begin{aligned} \mathbb{P}(B \cap A) &= \mathbb{P}(A)\mathbb{P}(B) = \mathbb{P}(A \cap B)\mathbb{P}(A) \\ &= \mathbb{P}(B) \end{aligned}$$

□◻◻

Conditional Probability of a Subset

Given the events A and B where $\mathbb{P}(A) > 0$. If $A \subset B$ then $\mathbb{P}(B \mid A) = 1$

Proof:

$A \subset B \implies A \cap B = A$ therefore

$$\mathbb{P}(B \mid A) = \frac{\mathbb{P}(A)}{\mathbb{P}(A)} = 1$$

□◻◻

Probability Measurement of Conditional Probability

Given the probability space $(\Omega, \mathcal{A}, \mathbb{P})$ and $A \in \mathcal{A}$ where $\mathbb{P}(A) > 0$. The map $B \rightarrow \mathbb{P}(B \mid A)$ is a probability measurement with the following properties.

- $\forall \in \mathcal{A} : \mathbb{P}(B \mid A) \geq 0$
- $\mathbb{P}(\Omega \mid A) = 1$
- $B_n \in \mathcal{A}, n \in \mathbb{N}$ pairwise disjoint $\implies \mathbb{P}(\bigcup_{n \in \mathbb{N}} B_n \mid A) = \sum_{n \in \mathbb{N}} \mathbb{P}(B_n \mid A)$

We also get

$$\mathbb{P}(\emptyset \mid A) = 0$$

$$\mathbb{P}(B^c \mid A) = 1 - \mathbb{P}(B \mid A)$$

Proof:

- Because our sets do not have negative cardinalities and the lowest we can reach is the probability of the empty-set we can not get below 0.
- Similar to another proof we get $\frac{\mathbb{P}(A)}{\mathbb{P}(A)}$
- By the definition of probability space.

□◻◻

Multiplication Theorem

Given $A_1, \dots, A_n$ events with $\mathbb{P}(A_1 \cap \dots \cap A_{n-1}) \neq 0$. Then

$$\mathbb{P}(A_1 \cap \dots \cap A_n) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2 \mid A_1) \cdot \dots \cdot \mathbb{P}(A_n \mid A_1 \cap \dots \cap A_{n-1})$$

Theorem of the Total Probability

Given our standard probability space, the disjoint partitions of Ω , $B_1, \dots, B_n$ where $B_1 \cup \dots \cup B_n = \Omega$, $B_i \cap B_j = \emptyset$ for $i \neq j$ and $\mathbb{P}(B_i) > 0$ then

$$\mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A \cap B_i) = \sum_{i=1}^n \mathbb{P}(A \mid B_i) \mathbb{P}(B_i) \forall A \in \mathcal{A}$$

Example:

You are given three bags

- Bag A has 2 red balls and 3 green balls
- Bag B has 3 red balls and 2 green balls
- Bag C has 1 red ball and 4 green balls

What is the probability of a ball being red?

The answer is given by

$$\mathbb{P}(R) = \mathbb{P}(R \cap A) + \mathbb{P}(R \cap B) + \mathbb{P}(R \cap C)$$

Which gives us

$$\mathbb{P}(R) = \mathbb{P}(R \mid A) \mathbb{P}(A) + \mathbb{P}(R \mid B) \mathbb{P}(B) + \mathbb{P}(R \mid C) \mathbb{P}(C)$$

Finally we plug in the individual numbers

$$\mathbb{P}(R) = \frac{1}{3} \left(\frac{2}{5} + \frac{3}{2} + \frac{1}{4} \right) = \frac{6}{15}$$

68.0.15 Multistage Random Experiments

Let $\Omega_1, \dots, \Omega_n$ be discrete sample spaces of n sub-experiments, and let $\Omega = \Omega_1 \times \dots \times \Omega_n$. Furthermore, let an initial distribution be given by the probability function p_1 with

$$\sum_{\omega_1 \in \Omega_1} p_1(\omega_1) = 1$$

and transition probabilities p_j for $j = 2, \dots, n$ satisfying

$$\sum_{\omega_j \in \Omega_j} p_j(\omega_j \mid \omega_1, \dots, \omega_{j-1}) = 1,$$

for $(\omega_1, \dots, \omega_{j-1}) \in \Omega_1 \times \dots \times \Omega_{j-1}$.

This is used in the case when we model our experiment as a tree of probabilities.

For the multistage random experiment, the following two *path rules* apply:

1. Let $(\omega_1, \dots, \omega_n) \in \Omega$. Then we have:

$$p(\omega_1, \dots, \omega_n) = p_1(\omega_1) \cdot p_2(\omega_2 \mid \omega_1) \cdot \dots \cdot p_n(\omega_n \mid \omega_1, \dots, \omega_{n-1})$$

2. For $A \subset \Omega$, it holds that

$$\mathbb{P}(A) = \sum_{(\omega_1, \dots, \omega_n) \in A} p(\omega_1, \dots, \omega_n)$$

Note that for independent sub-experiments, we obtain the product space with

$$p(\omega_1, \dots, \omega_n) = p_1(\omega_1) \cdot p_2(\omega_2) \cdot \dots \cdot p_n(\omega_n).$$

68.0.16 Bayes Theorem

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space. Furthermore, let $B_1, \dots, B_n$ be a disjoint partition of the sample space Ω with $\mathbb{P}(B_i) > 0$ for all $i = 1, \dots, n$. Then, for every event $A \in \mathcal{A}$, the following holds:

$$\mathbb{P}(B_i | A) = \frac{\mathbb{P}(A | B_i) \mathbb{P}(B_i)}{\mathbb{P}(A)} = \frac{\mathbb{P}(A | B_i) \mathbb{P}(B_i)}{\sum_{k=1}^n \mathbb{P}(A | B_k) \mathbb{P}(B_k)}.$$

In words and by changing A with E of evidence and B with H of hypothesis, we want to know how much the actual evidence changes our initial belief. The process can be described as: you have a hypothesis, you have observed some evidence and finally, you want to know the probability that your hypothesis is true given the observed evidence.

$$\mathbb{P}(H | E) = \frac{\mathbb{P}(E | H) \mathbb{P}(H)}{\mathbb{P}(H) \mathbb{P}(E | H) + \mathbb{P}(\neg H) \mathbb{P}(E | \neg H)}$$

- $\mathbb{P}(H)$ is called the *prior*. Our initial hypothesis, without seeing the evidence.
- $\mathbb{P}(E | H)$ is called the *likelihood*. This is the probability that we will see our hypothesis reflected on the evidence given that our hypothesis is true.
- $\mathbb{P}(E | \neg H)$ represents the probability that our hypothesis will be reflected on the sample given our hypothesis is false.

With this knowledge, breaking down the formula is very intuitive

- $\mathbb{P}(H | E)$ is the *posterior*. It describes the following question: How often is H true among the cases where H is true. probability
- $\mathbb{P}(H) \mathbb{P}(E | H)$ is the proportion of our hypothesis in the sample being true times the actual proportion of the fitting samples inside our hypothesis.
- $\mathbb{P}(H) \mathbb{P}(E | H) + \mathbb{P}(\neg H) \mathbb{P}(E | \neg H)$ consist of our expression above plus the proportion of the counter hypothesis times the proportion of the representatives of our hypothesis which will show up in the evidence in the counter group.

Example:

A disease affects 1 percent of the population, and you were tested positive with a test which has a 99.9 percent accuracy. What is the probability for you truly being infected?

$$\mathbb{P}(I | +) = \frac{\mathbb{P}(I) \mathbb{P}(+ | I)}{\mathbb{P}(I) \mathbb{P}(+ | I) + \mathbb{P}(\neg I) \mathbb{P}(+ | \neg I)}$$

Our numbers are:

- $\mathbb{P}(I) = 0.001$
- $\mathbb{P}(+ | I) = 0.99$
- $\mathbb{P}(\neg I) = 0.999$
- $\mathbb{P}(+ | \neg I) = 0.1$

Thus, our formula gives us a 9 percent chance of actually being infected.

68.0.17 Law of Large Numbers

Given a sequence of independent random variables $X_1, X_2, \dots, X_n$ with the same mean μ and the same σ^2 . The sample mean $Z_n = \frac{1}{n} \sum_{i=1}^n X_i$ will converge to μ as $n \rightarrow \infty$.

Formally $P(|Z_n - \mu| > \epsilon)_{n \rightarrow \infty} \rightarrow 0$.

Proof:

Since our random variables are independent

$$\begin{aligned} E[Z_n] &= \frac{1}{n} \sum_{i=1}^n E[X_i] = \mu \\ \text{var}[Z_n] &= \frac{1}{n} \sum_{i=1}^n \text{var}[X_i] = \frac{\sigma^2}{n} \end{aligned}$$

as $n \rightarrow \infty$, $\text{var}(Z_n) \rightarrow 0$.

Formally,

$$\begin{aligned} \mathbb{P}(|Z_n - \mu| \geq \epsilon) &\leq \frac{\text{var}(Z_n)}{\epsilon^2} \\ &= \frac{\sigma^2}{n\epsilon^2} \rightarrow 0 \text{ as } n \text{ goes to infinity} \end{aligned}$$

68.0.18 Strong Law of large Numbers

Given $X_1, X_2, \dots$ independent random variables with the same distribution and $|E[X_1]| < \infty$ then

$$Z_n \rightarrow E[X_1]$$

where $Z_n = \frac{1}{n} \sum_{i=1}^n X_i$

68.0.19 Central Limit Theorem

Given the same conditions of the *Law of large Numbers*. The sample mean

$$Z_n = \frac{1}{n} \sum_{i=1}^n X_i \sim \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$$

or equivalently

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\sqrt{n} \frac{Z_n - \mu}{\sigma} \leq x\right) = \Phi(x)$$

Example:

Approximate the probability of a tricked coin landing heads with $n = 10000$ throws and the number of heads between $4900 < X < 5100$.

Our variable is binomial distributed

$$\mu = E[X] = np = 5000$$

$$\sigma = 50$$

Then for

$$\mathbb{P}\left(Z_1 \leq \sqrt{n} \frac{Z_n - \mu}{\sigma} \leq Z_2\right)$$

with $Z_1 = \frac{4900-5000}{50}$, $Z_2 = \frac{5100-5000}{50}$ We get

$$\Phi(Z_2) - \Phi(Z_1) = 0.9545$$

68.0.20 Approximating the Binomial Distribution with the Normal Distribution

Given a binomial distribution with parameters n and p . For large n the distribution can be approximated by a normal distribution with parameters $\mu = np$ and $\sigma = \sqrt{np(1-p)}$.

Example:

Given $n = 75$ and $p = 0.6$ approximate the probability of getting between $\mathbb{P}(X \geq 52)$

$\mu = np = 45$, $\sigma^2 = np(1-p) = 18$

$$Z = \frac{X - \mu}{\sigma} = \frac{X - 45}{\sqrt{18}}$$

Therefore,

$$\begin{aligned} & \mathbb{P}\left(Z \geq \frac{52 - \mu}{\sqrt{18}}\right) \\ &= 1 - \Phi\left(\frac{52 - 45}{\sqrt{18}}\right) = 0.0582 \end{aligned}$$

For $\mathbb{P}(50 \leq X \leq 52)$ we have

$$\mathbb{P}\left(\frac{50 - \mu}{\sqrt{18}} \leq Z \leq \frac{52 - \mu}{\sqrt{18}}\right) = \Phi\left(\frac{52 - 45}{\sqrt{18}}\right) - \Phi\left(\frac{50 - 45}{\sqrt{18}}\right)$$

68.0.21 Theorem of Moivre-Laplace

Given $X_1, X_2, \dots, X_n$ Bernoulli distributed random variables and $S_n = \sum_{i=1}^n X_i$ then the number of successes is given by

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{S_n - np}{\sqrt{np(1-p)}} \leq x\right) = \Phi(x)$$

For the case that we have a region with two bounds $a < S_n < b$ we get

$$\mathbb{P}\left(\frac{S_n - np}{\sqrt{np(1-p)}} \leq Z_2\right) - \mathbb{P}\left(\frac{S_n - np}{\sqrt{np(1-p)}} \leq Z_1\right)$$

If normal approximation is used for a discrete distribution we have to apply the continuity correction.

$$\mathbb{P}(a < S_n < b) \approx \Phi\left(\frac{b + 0.5 - np}{\sqrt{np(1-p)}}\right) - \Phi\left(\frac{a - 0.5 - np}{\sqrt{np(1-p)}}\right)$$

Continuity Correction

An approximation is useful if $np(1-p) \geq 9$

Example:

Approximate the probability of getting between $4900 < X < 5100$ heads when throwing a tricked coin $n = 10000$ times with $p = 0.5$.

$$\mathbb{P}\left(\frac{S_n - np}{\sqrt{np(1-p)}} \leq Z_2\right) - \mathbb{P}\left(\frac{S_n - np}{\sqrt{np(1-p)}} \leq Z_1\right)$$

with $Z_1 = \frac{4900-5000}{50}$, $Z_2 = \frac{5100-5000}{50}$

$$\Phi(Z_2) - \Phi(Z_1) = 0.9545$$

The complete formula for this case looks like this

$$\mathbb{P}(4900 < S_n < 5100) \approx \Phi\left(\frac{5100 + 0.5 - np}{\sqrt{np(1-p)}}\right) - \Phi\left(\frac{4900 - 0.5 - np}{\sqrt{np(1-p)}}\right)$$

Chapter 69

Single Variable Distributions

69.0.1 Random Variables

Given a probability space $(\Omega, \mathcal{A}, \mathbb{P})$, we call a function $f : \Omega \rightarrow \mathbb{R}$ *measurable* if for all $z \in \mathbb{R}$ the following applies

$$f^{-1}((-\infty, z]) = \{\omega \in \Omega : f(\omega) \leq z\} \in \mathcal{A}$$

This measurable function $X : \Omega \rightarrow \mathbb{R}$ is a *random variable*.

Note that for discrete probability spaces $\{\omega : X(\omega) \leq z\} \subseteq \Omega \in \mathcal{P}$

Also, it is considered discrete if it has only a finite or an infinite countable number of values.

Finally, we define the probability for random variables as

$$\mathbb{P}(X = k) := \mathbb{P}(\{X = k\})$$

Operations with Random Variables

Given two R.V. X, Y and some number $\alpha \in \mathbb{R}$, then

- $X + Y$
- $X - Y$
- $X \cdot Y$
- $\max(X, Y)$
- $\min(X, Y)$
- αX or αY

are also R.V.

Example:

A company has fixed costs of 250 euros per month. It produces toys for a price of 6 euros with a production price of 3 euros. On average, it sells 1000, $\mu = 1000$ toys with a $\sigma = \sqrt{500}$. Determine the random variables of the costs, the sales volume and the profit.

Costs: $X_c = 250 + 3X$ with $\mu_c = 250 + 3\mu = 3250$ and $\sigma^2 = 9 \cdot 500 = 1500$

Sales volume: $X_s = 6X$ with $\mu_c = 6\mu = 6000$ and $\sigma^2 = 6 \cdot 500 = 3000$

Profit: $X_p = X_s - X_c = 250 + 6X - 3X$ with $\mu_s = 3X - 250 = 2750$ and $\sigma^2 = 3 \cdot 500 = 1500$

69.0.2 Indicator Function

Given $A \subseteq \Omega$, we define the *indicator function* as

$$\mathbb{1}_A(\omega) = \begin{cases} 1, & \text{if } \omega \in A \\ 0, & \text{if } \omega \notin A \end{cases}$$

We also get the following rules:

- $\mathbb{1}_\emptyset = 0$
- $\mathbb{1}_A^2 = \mathbb{1}_A$
- $\mathbb{1}_{A^c} = 1 - \mathbb{1}_A$
- $\mathbb{1}_{A \cap B} = \mathbb{1}_A \mathbb{1}_B$
- $\mathbb{1}_{A \cup B} = \mathbb{1}_A + \mathbb{1}_B - \mathbb{1}_{A \cap B}$
- $\mathbb{1}_{A+B} = \mathbb{1}_A + \mathbb{1}_B$

Indicator Sum

Given the events $A_1, \dots, A_n \in \Omega$. We define the random variable

$$X := \sum_{j=1}^n \mathbb{1}_{A_j}$$

as the *indicator sum* or *counting variable*.

The realization of X gives us how much A_j have occurred.

$$\{X = k\} = \sum_{T \subseteq \{1, \dots, n\}: |T|=k} \left(\bigcap_{j \in T} A_j \cap \bigcap_{j \notin T} A_j^c \right), k \in \{0, 1, \dots, n\}$$

69.0.3 Distributions

Given our probability space and a R.V. X . The probability measurement $\mathbb{P}_X$ to $\mathbb{R}$ defined by

$$\mathbb{P}_X(A) := \mathbb{P}(\{\omega : X(\omega) \in A\}), \text{ for } A \subseteq \mathbb{R}$$

is called as the *distribution* of X . Also, we write $\mathbb{P}(X \in A)$ as an abbreviation for $\mathbb{P}(\{\omega : X(\omega) \in A\})$.

69.0.4 Distribution Function

Given a probability space, a probability variable and a distribution $\mathbb{P}_X$ we define the function

$$\mathbb{F}_X(x) = \mathbb{P}(X \leq x) = \mathbb{P}(\{\omega : X(\omega) \leq x\})$$

as the *distribution function*.

Properties of Distribution Functions

- $\mathbb{F}_X$ is monotonic growing
- It is right-sided continuous
- $\lim_{x \rightarrow -\infty} \mathbb{F}_X(x) = 0$ and $\lim_{x \rightarrow \infty} \mathbb{F}_X(x) = 1$

69.0.5 Cumulative Probability Calculations

- $\mathbb{P}(X = a) = \mathbb{P}(X = a)$
- $\mathbb{P}(X \leq a) = \mathbb{P}(X \leq a)$
- $\mathbb{P}(X < a) = \mathbb{P}(X \leq a - 1)$
- $\mathbb{P}(X > a) = 1 - \mathbb{P}(X \leq a)$
- $\mathbb{P}(X \geq a) = 1 - \mathbb{P}(X \leq -1)$
- $\mathbb{P}(a \leq X \leq b) = \mathbb{P}(X \leq b) - \mathbb{P}(X \leq a)$
- $\mathbb{P}(|X - a| < b) = \mathbb{P}(-b < |X - a| < b) = \mathbb{P}(b) - \mathbb{P}(-b)$

69.0.6 The Expected Value for Discrete Random Variables

Given $\Omega, \mathcal{A}, \mathbb{P}$, a probability space and X a discrete random variable. We define the *expected result* as

$$\mathbb{E}[X] = \sum_{x \in X(\Omega)} x \mathbb{P}(X = x)$$

which exists only if

$$\sum_{x \in X(\Omega)} |x| \mathbb{P}(X = x) < \infty$$

Where x is an event $\mathbb{P}(X = x)$ its respective probability. It represents the mean of the random variable.

Transformation of the Expected Value

Given $u : X(\Omega) \rightarrow \mathbb{R}$ where $\sum_{x \in X(\Omega)} |u(x)|p(x) < \infty$ then

$$\mathbb{E}[u(X)] = \sum_{x \in X(\Omega)} u(x) \mathbb{P}(X = x)$$

This represents the expectation of the function.

69.0.7 Triangle Inequality for Expected Values

$$|\mathbb{E}[X]| \leq \mathbb{E}[|X|]$$

Proof:

$$\begin{aligned} |\mathbb{E}[X]| &= \left| \sum_{x \in X(\Omega)} x \mathbb{P}(X = x) \right| \leq \sum_{x \in X(\Omega)} |x| \mathbb{P}(X = x) \\ &= \sum_{x \in X(\Omega)} |x| \mathbb{P}(X = x) = \mathbb{E}[|X|] \end{aligned}$$

□◻◻

Properties of Expected Values

Given two random variables X and Y , if both have an expected value then

- $\mathbb{E}[\alpha X] = \alpha \mathbb{E}[X]$
- $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- $\mathbb{E}[b] = b$
- $X \leq Y \Rightarrow \mathbb{E}[X] \leq \mathbb{E}[Y]$

69.0.8 Variance

Given a random variable X for which $E[(X - E[X])^2]$ exists. Then we define the *variance* of X as

$$\text{var}(X) = E[(X - E[X])^2] = \sum_{x \in X} (x - E[X])^2 \mathbb{P}(X = x)$$

And its root is the *standard deviation*.

Properties of the Variance

Let X be a (discrete) random variable for which the variance exists. Then the following hold:

- $\text{var}(aX + b) = a^2 \text{var}(X)$, for $a, b \in \mathbb{R}$
- $\text{var}(X) = E[X^2] - (E[X])^2$
- For $a \in \mathbb{R}$, we have $E[(X - a)^2] = \text{var}(X) + (E[X] - a)^2$, and in particular, $E[(X - a)^2] \geq \text{var}(X)$. Equality holds if and only if $a = E[X]$.

Example:

Given $E(X) = 1.2$ and $\mathbb{P}(X = 0) = 0.16$, $\mathbb{P}(X = 1) = 0.48$, $\mathbb{P}(X = 2) = 0.36$.

$$E[X^2] = 0^2(0.16) + 1^2(0.48) + 2^2(0.36) = 1.92$$

$$\text{var}(X) = E[(X - E[X])^2] = (0 - 1.2)^2(0.16) + (1 - 1.2)^2(0.48) + (2 - 1.2)^2(0.36)$$

69.0.9 Moments

For a random variable X , we define k . *moment* as $E[X^k]$

Markov Inequality

Given a random variable X , $a \in \mathbb{R}$ and $h : D \rightarrow [0, \infty)$ a monotonic growing function with the properties $h(a) > 0$ and $X(\Omega) \subseteq D$ then

$$\mathbb{P}(X \geq a) \leq \frac{E[h(X)]}{h(a)}$$

Simpler,

$$\mathbb{P}(X \geq a) \leq \frac{E[X]}{a}$$

There is a special case for $Y = |X|$, $a > 0$ and $h(x) = x$

$$\mathbb{P}(X \geq a) \leq \frac{E[h(X)]}{a}$$

Chebychev Inequality

Given a random variable X with a variance and $a > 0$ then

$$\mathbb{P}(|X - E[X]| \geq a) \leq \frac{\text{var}(X)}{a^2}$$

This states that the probability of X is at least a away of its mean can not be too big.

69.0.10 Probability Density

A Lebesgue-integrable function $f : \mathbb{R} \rightarrow [0, \infty)$ is called the (probability) *density* or *density function* of the random variable X or of the distribution $\mathbb{P}_X$ if, for all $a, b \in \mathbb{R}$ with $a \leq b$, the following holds:

$$\mathbb{P}(a < X \leq b) = \mathbb{P}_X((a, b]) = \int_a^b f(x) dx.$$

Distributions that possess a probability density are called *continuous distributions*.

Properties of Continuous Distributions

- $\mathbb{F}_X$ is continuous at x if and only if $\mathbb{P}(X = x) = 0$.

-

$$\int_{-\infty}^{\infty} f_X(x) dx = \mathbb{F}_X(\infty) = \lim_{x \rightarrow \infty} P(X \leq x) = 1.$$

- If X is a continuous random variable, then F_X is continuous.

How to Test if a Function works as a Density

To test if given a function it can be used as a density functions we test:

- $f(x) \geq 0 \forall x \in \mathbb{R}$
- $\int_{-\infty}^{\infty} f(x) dx = 1$

Zero Property

Given a continuous random variable X then

For $a, b \in \mathbb{R}$

$$\mathbb{P}(a \leq X \leq b) = F_X(b) - F_X(a) = \int_a^b f_X(x) dx = \mathbb{P}(a < X \leq b)$$

Also $A \in \mathcal{B}(\mathbb{R}) : \mathbb{P}(X \in A) = \int_A f_X(x) dx$

And $A \in \mathcal{B}(\mathbb{R})$ (discrete) : $A = \{a_i\}_{i \in I}, I \in \mathbb{N} :$

$$\mathbb{P}(X \in A) = \sum_{i \in I} \mathbb{P}(X = a_i) = 0$$

69.0.11 Expected Value for Continuous Random Variables

Given a continuous random variable X with a density f_X . The *expected value* of X exists if $\int |x| f_X(x) dx, \infty$. In that case we define it as

$$\mathbb{E}[X] = \int x f_X(x) dx$$

We also get all of its properties like for the discrete case.

Transformation Lemma

Given a continuous random variable X with a density f and a measurable function $u : \mathbb{R} \rightarrow \mathbb{R}$ with $\int |u(x)| f(x) dx < \infty$ then the following applies

$$\mathbb{E}[u(X)] = \int u(x) f(x) dx$$

69.0.12 Transformation Theorem for Densities

Given a random variable X with values in the open interval $I \subseteq \mathbb{R}$ and the density f_X . If $J \subseteq \mathbb{R}$ is an open interval and $u : I \rightarrow J$ is a diffeomorphism then $Y = u(X)$ has the density

$$f_Y(y) = f_X(u^{-1}(y))|(u^{-1})'(y)|$$

For $y \in J$ and $f_Y(y) = 0$ for $y \in \mathbb{R} \setminus J$.

69.0.13 Quantile

Let X be a random variable with distribution F . For $p \in (0, 1)$, a value q_p is called a *p-quantile* of X (or of F) if:

$$\mathbb{P}(X \leq q_p) \geq p \text{ and } \mathbb{P}(q_p \leq X) \geq 1 - p,$$

or equivalently,

$$F(q_p) \geq p \text{ and } \lim_{x \nearrow q_p} F(x) \leq p.$$

In general, the p -quantile is not unique. If F is strictly increasing and continuous, then q_p is unique. If q_p is not unique, we can make it “unique” by defining the smallest quantile as

$$F^{-1}(p) = \inf\{x \in \mathbb{R} : F(x) \geq p\}.$$

69.0.14 The Moment Generating Function

The n -th moment of a distribution are given by $E[X^n]$. We also know that the first moment is the mean and the variance is closely related to the second moment. The moments *uniquely* determine the distribution.

The *moment generating function* is defined as

$$M(t) = \begin{cases} \sum_x e^{tx} \mathbb{P}(X = x) & \text{if } X \text{ discrete} \\ \int_{-\infty}^{\infty} e^{tx} \mathbb{P}(X = x) dx & \text{if } X \text{ continuous} \end{cases}$$

$M(t)$ uniquely determines the cumulative probability distribution of $\mathbb{P}(X = x)$

Derivative of the Moment Generating Function

The n -th derivative of the *moment generating function* has the following property

$$M^n(0) = E[X^n]$$

Proof (Informal):

We will handle only the continuous case, but the same holds for the discrete case

$$\begin{aligned} \frac{d}{dt} \int_{-\infty}^{\infty} e^{tx} \mathbb{P}(X = x) dx &= \int_{-\infty}^{\infty} x e^{tx} \mathbb{P}(X = x) dx \\ \frac{d}{dt} \int_{-\infty}^{\infty} x e^{tx} \mathbb{P}(X = x) dx &= \int_{-\infty}^{\infty} x^2 e^{tx} \mathbb{P}(X = x) dx \end{aligned}$$

And the pattern repeats. This is just given us the definition of each of the moments.

◻◻◻

The Additive Property of the Moment Generating Function

Let $X_1, X_2, \dots, X_n$ be random variables with their corresponding moment generating functions. Let $Z = \sum_i X_i$ then

$$M_Z(t) = \prod_i M_{X_i}(t)$$

69.0.15 Discrete Uniform Distribution

A random variable $X \in \{1, 2, \dots, n\}$ with $\mathbb{P}(X = i) = \frac{1}{n}$ for $i = 1, \dots, n$ is uniformly distributed. The distribution $\mathbb{P}_X$ is called the *discrete uniform distribution*. Noted as $\mathcal{U}(n)$

$$\mathbb{P}(X = i) = \begin{cases} \frac{1}{n}, & i = 1, \dots, n \\ 0, & \text{else} \end{cases}$$

The expected value is given by

$$\mathbb{E}(X) = \frac{n+1}{2}$$

and the standard deviation is given by

$$\sigma = \frac{n^2 - 1}{12}$$

Finally, its cumulative distribution is given by

$$\mathbb{F}_X(x) = \sum_{i=1}^x \frac{1}{n}$$

69.0.16 Continuous Uniform Distribution

A random variable $X \in [a, b]$ with $\mathbb{P}(X = i) = \frac{1}{b-a}$ for $i \in [a, b]$ is uniformly distributed. The distribution is called the *continuous uniform distribution*. Noted as $\mathcal{U}(a, b)$

$$\mathbb{P}(X = i) = \begin{cases} \frac{1}{b-a}, & i \in [a, b] \\ 0, & \text{else} \end{cases}$$

The expected value is given by

$$\mathbb{E}(X) = \frac{a+b}{2}$$

and the standard deviation is given by

$$\sigma = \frac{(b-a)^2}{12}$$

Finally, its cumulative distribution is given by

$$\mathbb{F}_X(x) = \begin{cases} 0, & x < a \\ \frac{x-a}{b-a}, & x \in [a, b] \\ 1, & x > b \end{cases}$$

69.0.17 Bernoulli Distribution

The Bernoulli distribution is used modeling for binary events with only one trial, noted as $\mathcal{B}(p)$

$$\mathbb{P}(X = x) = \begin{cases} p, & x = 1 \\ 1 - p, & x = 0 \end{cases}$$

The expected value, variance and cumulative distribution function are give by

$$\mathbb{E}(X) = p$$

$$\text{var}(X) = p(1 - p)$$

$$\mathbb{F}_X = \begin{cases} 0, & x < 0 \\ 1 - p, & 0 \leq x < 1 \\ 1, & x \geq 1 \end{cases}$$

69.0.18 Binomial Distribution

From the Bernoulli distribution we can derive the *binomial distribution* $\text{Bin}(n, p)$ as having a number of consecutive independent Bernoulli experiments.

This distribution is for discrete binary experiments. Either an event occurs or it does not.

$$\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

With

$$\mathbb{E}[X] = np$$

$$\text{var}(X) = \mathbb{E}[X](1 - p)$$

The cumulative distribution function is given by

$$\mathbb{F}_X = \sum_{i=0}^k \mathbb{P}(X = i)$$

If we let $n \rightarrow \infty$ with np being not too small then the limit of this distribution becomes the normal distribution.

69.0.19 Normal Distribution

The *normal distribution* $\mathcal{N}(\mu, \sigma^2)$ is the limit of the *binomial distribution* under certain conditions, and it is given by

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2} e^{\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}}$$

It also has

$$\mathbb{E}[X] = np = \mu$$

$$\text{var}(X) = E[X](1 - p)$$

To compute its cumulative probability we need to use special tables, due to the fact that the function can not be analytically integrated. To get a specific we use tables which pre-calculated values computed numerically using the so called Φ function. This function is used in the following manner:

We plug the values in the expression, our expected value and the standard deviation

$$Z = \frac{X - \mu}{\sigma}$$

$\Phi(Z)$ you use the portion before the comma as the row and after the comma as the column to read the value from.

Standard Normal Distribution

The *standard* normal distribution has the parameters $\mu = 0$ and $\sigma^2 = \text{var} = 1$.

To transform a normal distributed random variable X you can define a new random variable

$$Z = \frac{X - \mu}{\sigma}$$

Example I:

Given $X \sim N(10, 4)$. What is $\mathbb{P}(8 \leq X \leq 12)$?

We define $Z = \frac{X-10}{2}$. Then we get

$$\mathbb{P}(8 \leq X \leq 12) = \mathbb{P}\left(\frac{8-10}{2} \leq Z \leq \frac{12-10}{2}\right) = \mathbb{P}(-1 \leq Z \leq 1)$$

Using the standard normal distribution table we get

$$\mathbb{P}(-1 \leq Z \leq 1) = \mathbb{F}_Z(1) - \mathbb{F}_Z(-1) = 0.8413 - 0.1587 = 0.6826$$

Example II:

The diameter of mass-produced balls is normally distributed. The mean and standard deviation can be estimated using the following experiment:

Of two sieves, sieve A has holes with a diameter of 69 mm, and sieve B has holes with a diameter of 72 mm. It is observed that 30.85 percent of the balls pass through sieve A, while 15.87 percent of the balls are retained by sieve B.

Determine estimates for the mean and standard deviation of the ball diameters based on these experimental results.

First we have:

$$\mathbb{P}(X \leq 69) = 0.3085 \quad \mathbb{P}(X \geq 72) = 0.1587 \implies \mathbb{P}(X \leq 72) = 0.8413$$

If we look at the Z -table we get the z -values -0.5 and 1

This means

$$\frac{69 - \mu}{\sigma} = -0.5 \quad \frac{72 - \mu}{\sigma} = 1$$

Now we can solve the system of equations and the results will be:

$$\mu = 70 \quad \sigma = 2$$

69.0.20 Geometric Distribution

A random variable $X \in \mathbb{N}_0$ with $\mathbb{P}(X = n) = p(1 - p)^n$ for some $p \in (0, 1)$ and $n \in \mathbb{N}_0$ is geometrically distributed. The distribution $\mathbb{P}_X$ is called the *geometric distribution*.

Similar to the binomial distribution, the *geometric* $\text{Geo}(n, p)$ distribution is used for an experiment with two outcomes, and we want to know the probability of $k - 1$ non-successes until a success happens

$$\mathbb{P}(X = k) = (1 - p)^{k-1}p$$

We also get

$$\mathbb{E}[X] = \mu = \frac{1 - p}{p}$$

$$\text{var}(X) = \frac{1 - p}{p^2}$$

The cumulative distribution function is given by

$$\mathbb{F}_X = \begin{cases} 0, & x < 1 \\ 1 - (p - 1)^{\lfloor x \rfloor}, & x \geq 1 \end{cases}$$

69.0.21 Poisson Distribution

A random variable $X \in \mathbb{N}_0$ with

$$\mathbb{P}(X = n) = \frac{\lambda^n}{n!} e^{-\lambda}, \quad \text{for some } \lambda > 0,$$

We also define the distribution function

$$\mathbb{F}_X(n) = \sum_{i=0}^n \frac{\lambda^i}{i!} e^{-\lambda}$$

is Poisson distributed. The distribution $\mathbb{P}_X$ is called the *Poisson distribution* $\text{Poi}(\lambda)$. In simple words, it is the number of events x , in a specific interval. This is a limit of the binomial distribution for large n and small p .

We also get our expected result and variance

$$\mathbb{E}[X] = \lambda = np$$

and

$$\text{var}(X) = \lambda$$

The derivation is the following, let $\lambda = np$

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \binom{n}{k} p^k (1-p)^{n-k} \\
&= \frac{n!}{k!(n-k)!} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\
&= \frac{n(n-1) \dots (n-k+1)}{k!} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \\
&= \frac{n(n-1) \dots (n-k+1)}{k!} \frac{\lambda^k}{n^k} \left(1 - \frac{\lambda}{n}\right)^{n-k} \\
&= \frac{\lambda^k}{k!} \frac{n(n-1) \dots (n-k+1)}{n^k} \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-k} \\
&= \frac{\lambda^k}{k!} \cdot 1 \cdot e^{-\lambda} \cdot 1 \\
&= \frac{\lambda^k}{k!} \cdot e^{-\lambda}
\end{aligned}$$

69.0.22 Hyper-geometric Distribution

Similar to the binomial distribution, but we keep track of the changes after each choice. In other words, binomial distribution without repetition. We notate this as $\text{HG}(M, N, n)$.

$$\mathbb{P}(X = k) = \frac{\binom{M}{k} \binom{N-M}{n-k}}{\binom{N}{n}}$$

Nomenclature:

- N : Total number of elements
- M : Elements with the trait A
- $N - M$: Elements without the trait A
- k : Number to elements to take
- n : Number of iterations

Each part of the formula has a different meaning. $\binom{N}{n}$ represents the total number of ways to choose the elements from the total. $\binom{M}{k}$ gives us the ways of choose k elements with the trait from its total M and finally, $\binom{N-M}{n-k}$ represents the remaining ways of choosing elements we are not interested in.

We also get

$$\mathbb{E}[X] = n \frac{M}{N}$$

$$\text{var}(X) = \mathbb{E}[X] \left(1 - \frac{M}{N}\right) \left(\frac{N-n}{N-1}\right)$$

The cumulative distribution function is given by

$$\mathbb{F}_X(x) = \sum_{i=1}^x \mathbb{P}(X = i)$$

69.0.23 Exponential Distribution

We define the *exponential distribution* $\mathcal{E}(\lambda)$ as a real defined random variable, used mostly for the duration of time intervals.

$$f(n) = \begin{cases} \lambda e^{-\lambda n} & , n \geq 0 \\ 0 & , \text{ else} \end{cases}$$

Where *lambda* our *rate* parameter.

$$\lambda = \frac{1}{\text{avarage}}$$

We also define the distribution function

$$\mathbb{F}_X(X = n) = \int_{-\infty}^n \mathbb{P}(X = n) = \begin{cases} 1 - e^{-\lambda n} & , n \geq 0 \\ 0 & , \text{ else} \end{cases}$$

We also have

$$\mathbb{E}(X) = \frac{1}{\lambda}$$

$$\text{var}(X) = \frac{1}{\lambda^2}$$

Memory-less Property of the Exponential Distribution

This distribution is *memory-less*. This means that the future probabilities of an event are independent of the previous ones.

$$\mathbb{P}(X \geq x + y \mid X > y) = \mathbb{P}(X \geq x)$$

The Connection between the Exponential Distribution and the Poisson Process

A neat connection between these two concepts is that given an exponential distribution over some time t then the number of events which occurred during that interval is given by a Poisson distribution $\text{Poi}(\lambda t)$

Example:

Emails com at a rate of 5 per minute ($\lambda = 5$). What is the probability of waiting t until the next email?

The probability of getting an email after t waiting is given by the following exponential distribution

$$\mathbb{P}(T > t) = e^{-5t}$$

What is the probability of getting no emails in the next t minutes?

This is given by the following Poisson process

$$\mathbb{P}(k = 0) = \frac{5^0 e^{-5t}}{0!} = e^{-5t}$$

69.0.24 Gamma Distribution

This distribution comes directly from the connection between the exponential distribution and the Poisson process. Given events $w_0, \dots, w_n$ with an exponential distribution. The *gamma* distribution gives us the waiting time for the r -th event.

$$T_r = \sum_{k=0}^r w_k$$

We denote this as $T_r \sim \text{gamma}(r, \lambda)$. The probability density function is given by

$$\mathbb{P}(T_r > t) = \mathbb{P}(N_t \geq r - 1) = \sum_{k=0}^{r-1} \frac{e^{-\lambda t} (\lambda t)^k}{k!} = \dots = e^{-\lambda t} \frac{\lambda^r t^{r-1}}{(r-1)!}$$

with N_t is the number of Poisson events in t .

69.0.25 Chi-Square Distribution

The square of the *normal distribution* is called the *Chi-Square Distribution* noted as $\mathcal{N}(a\mu + b, a^2\sigma^2)$ or χ_n^2 with $n \in \mathbb{N}$ degrees of freedom.

Given a random variable X with density

$$f(x) = \frac{1}{2^{\frac{n}{2}} \Gamma(\frac{n}{2})} x^{\frac{n}{2}-1} e^{-\frac{x}{2}}$$

where Γ is the gamma function defined as

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$$

For the 1 degree of freedom we get

$$f(x) = \frac{1}{\sqrt{2\pi}} x^{-\frac{1}{2}} e^{-\frac{x}{2}}$$

We also have

$$\mathbb{E}[X] = n$$

$$\text{var}(X) = \mathbb{E}[X]2n$$

Derivation

The probability density function is reached by taking the cumulative density function of the normal distribution which is an Integral, square it, and then we take its derivative. We proceed as follows:

Given $X \sim \mathcal{N}(0, 1)$ we define $Y = X^2$. Then we get

$$F_Y(y) = \mathbb{P}(Y \leq y) = \mathbb{P}(-\sqrt{y} \leq X \leq \sqrt{y}) = F_X(\sqrt{y}) - F_X(-\sqrt{y})$$

This gives us

$\Phi(\sqrt{y}) - \Phi(-\sqrt{y})$ Now we take the derivative to get the density function

$$\begin{aligned}
f_Y(y) &= \frac{d}{dy} F_Y(y) = \frac{d}{dy} (\Phi(\sqrt{y}) - \Phi(-\sqrt{y})) \\
&= \Phi'(\sqrt{y}) \cdot \frac{1}{2\sqrt{y}} + \Phi'(-\sqrt{y}) \cdot \frac{1}{2\sqrt{y}} \\
&= y^{-\frac{1}{2}} f_X(\sqrt{y}) \\
&= \frac{1}{\sqrt{2\pi}} y^{-\frac{1}{2}} e^{-\frac{y}{2}}
\end{aligned}$$

69.0.26 Students t-Distribution

This is a distribution mostly used in statistics for hypothesis testing.

$$f_n(x) = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{n\pi}\Gamma\left(\frac{n}{2}\right)} \left(1 + \frac{x^2}{n}\right)^{-\frac{n+1}{2}}$$

This used to test the mean $\overline{X_n}$ against the true μ when σ^2 is unknown.

69.0.27 Activity Examples

Night guard

A night guard has 2 methods for finding the correct key out of 10. The first one A , consists in just testing each key without repetition. The second one B , consists on choosing one random key and then if it was not the correct one it just shuffles the keys and repeats the process.

What are the distributions of both the methods?

The distribution for A is the uniform with $n = 10$, for B it is the geometric distribution $P(X = k) = \left(\frac{9}{10}\right)^{k-1} \frac{1}{10}$

The guard is drunk $\frac{1}{3}$ of times. When it is drunk it uses method B , else A . What is the conditional probability of him being caught drunk during work after failing 8 times.

$$\mathbb{P}(\text{drunk} \mid X > 8) = \frac{\mathbb{P}(\text{drunk})\mathbb{P}(X > 8 \mid \text{not drunk})}{\mathbb{P}(X > 8)}$$

This is

$$\mathbb{P}(B \mid X \leq 8) = \frac{\frac{1}{3} \cdot \left(1 - \left(\frac{9}{10}\right)^{8-1} \frac{1}{10}\right)}{\frac{1}{3} \cdot \left(1 - \left(\frac{9}{10}\right)^{8-1} \frac{1}{10}\right) + \frac{2}{3} \cdot \left(1 - \frac{8}{10}\right)}$$

Chapter 70

Multivariable Distributions

Multivariable or *joint distributions* are a way of combining two or more random variables.

70.0.1 Measurability of Random Variables and Random Vectors

Let $X_1, X_2, \dots, X_n$ be random variables on $(\Omega, \mathcal{A}, P)$. Then each $X_i : \Omega \rightarrow \mathbb{R}$ is measurable.

Measurability in $\mathbb{R}$.

$$X_i^{-1}((-\infty, z]) \in \mathcal{A} \quad \text{for all } z \in \mathbb{R}$$

Alternative Representation.

Define $X : \Omega \rightarrow \mathbb{R}^n$ by

$$X(\omega) = (X_1(\omega), \dots, X_n(\omega))$$

It can be shown (from measure theory) that X is measurable.

Measurability in $\mathbb{R}^n$.

$$X^{-1}(\mathbb{R}) \in \mathcal{A}, \quad \text{where } \mathbb{R} = (a_1, b_1] \times (a_2, b_2] \times \dots \times (a_n, b_n]$$

for arbitrary $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$.

From this (by measure theory) it follows that for any set $A \subseteq \mathbb{R}^n$,

$$X^{-1}(A) \in \mathcal{A}$$

Hence, the probability

$$\mathbb{P}(X \in A) = \mathbb{P}(\{\omega \in \Omega : X(\omega) \in A\})$$

is well-defined for all $A \subseteq \mathbb{R}^n$.

70.0.2 Joint Distribution and Random Vectors

Let $X_1, \dots, X_n$ be random variables on a probability space $(\Omega, \mathcal{A}, P)$. Then P_X (respectively $P_{X_1, \dots, X_n}$) is defined as

$$\mathbb{P}_X(A) = P((X_1, \dots, X_n) \in A), \quad A \subseteq \mathbb{R}^n$$

and is called the *joint distribution* of $X_1, \dots, X_n$. The vector $X = (X_1, \dots, X_n)$ is called a *random vector*.

This is just the probability of n events happening simultaneously.

70.0.3 Joint Probability Function

Let $X_1, \dots, X_n$ be discrete random variables. The function p_X , defined as

$$\mathbb{P}_X(x) = \mathbb{P}(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

for $x \in X(\Omega) \subseteq \mathbb{R}^n$, is the *joint probability function* of $X_1, \dots, X_n$, or equivalently, of the random vector X .

As before, the distribution P_X is uniquely determined by the probability function p_X :

$$\mathbb{P}_X(A) = \mathbb{P}((X_1, \dots, X_n) \in A) = \sum_{x \in A} \mathbb{P}(X_1 = x_1, \dots, X_n = x_n) = \sum_{x \in A} p_X(x)$$

Furthermore, for probability functions $p : \mathbb{R}^n \rightarrow \mathbb{R}$, the following holds:

$$\begin{aligned} \sum_{(x_1, \dots, x_n) \in X(\Omega)} p(x_1, \dots, x_n) &= 1 \\ p(x_1, \dots, x_n) &\geq 0 \end{aligned}$$

Example:

Let X, Y be the numbers shown when rolling two dice. The joint probability function is

$$p_{X,Y}(i, j) = \frac{1}{36}$$

for $i, j \in \{1, 2, \dots, 6\}$. Their combined probability function is

$$p_{X,Y}(i, j) = \begin{cases} \frac{1}{6}, & \text{if } i = j \\ 0, & \text{otherwise.} \end{cases}$$

70.0.4 Marginal Distributions of Discrete Random Variables

Let X, Y be two discrete random variables. The *marginal probability functions* of X and Y are given by

$$p_X(x) = \sum_y p_{X,Y}(x, y) \quad \text{and} \quad p_Y(y) = \sum_x p_{X,Y}(x, y).$$

This means that we fix one variable and sum over all possible values of the other variable. For the 2d case, if we want a marginal distribution of X we sum over all rows for a fixed column.

Generalized, let $X_1, \dots, X_n$ be random variables and $1 \leq i_1 < \dots < i_k \leq n$.

The joint distribution of $X_{i_1}, \dots, X_{i_k}$ is a *k-dimensional marginal distribution* of $X_1, \dots, X_n$.

For discrete random variables, the joint probability function of $X_{i_1}, \dots, X_{i_k}$ is called the *marginal probability function*.

Example:

Let X_1, X_2 be the outcomes of two dice rolls, and let $Y = X_1 + X_2$. The joint distribution of X_1 and Y is given by:

$X_1 \backslash Y$	2	3	4	5	6	7	8	9	10	11	12	$\mathbb{P}(X_1 = x)$
1	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	0	0	0	0	0	$\frac{1}{6}$
2	0	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	0	0	0	0	$\frac{1}{6}$
3	0	0	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	0	0	0	$\frac{1}{6}$
4	0	0	0	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	0	0	$\frac{1}{6}$
5	0	0	0	0	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	0	$\frac{1}{6}$
6	0	0	0	0	0	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{6}$
$\mathbb{P}(Y = y)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	

In the last column and row we get the marginal distributions of X_1 and Y respectively.

Marginal Probability Function for Multiple Discrete Variables

Let $X_1, \dots, X_n$ be discrete random variables with joint probability function $p(x_1, \dots, x_n)$, and let $1 \leq i_1 < \dots < i_k \leq n$. For the *marginal probability function* of $X_{i_1}, \dots, X_{i_k}$, the following holds:

$$p_{X_{i_1}, \dots, X_{i_k}}(x_{i_1}, \dots, x_{i_k}) = \sum_{(x_j)_{j \notin \{i_1, \dots, i_k\}}} p(x_1, \dots, x_n)$$

This is just notation for summing over all variables that are not of interest.

70.0.5 Multivariate Expected Value for Discrete Random Variables

Let $X_1, \dots, X_n$ be discrete random variables with joint probability function p , and let $u : \mathbb{R}^n \rightarrow \mathbb{R}$. Then, if the expectation exists, the following holds:

$$\mathbb{E}[u(X_1, \dots, X_n)] = \sum_{(x_1, \dots, x_n) \in X(\Omega)} u(x_1, \dots, x_n) p(x_1, \dots, x_n)$$

Example:

Let X and Y be random variables with the following joint probability function p :

$Y \backslash X$	1	2	3
3	$\frac{1}{6}$	0	$\frac{1}{4}$
4	$\frac{1}{8}$	$\frac{1}{3}$	$\frac{1}{8}$

Let $u(X, Y) = xy$ then our expected value is given by:

$$\mathbb{E}[XY] = \sum_{(x,y) \in (X,Y)(\Omega)} xy p(x,y) = 1 \cdot 3 \cdot \frac{1}{6} + 2 \cdot 3 \cdot 0 + 3 \cdot 3 \cdot \frac{1}{4} + 1 \cdot 4 \cdot \frac{1}{8} + 2 \cdot 4 \cdot \frac{1}{3} + 3 \cdot 4 \cdot \frac{1}{8} \approx 7.42$$

Expected Value as a Special Case of the Transformation Formula

Let X, Y be discrete random variables with joint probability function p . With $u(x, y) = x$, the transformation formula gives:

$$\mathbb{E}[X] = \sum_{(x,y) \in (X,Y)(\Omega)} x p(x,y)$$

70.0.6 Joint Probability Density Function

An integrable function $f : \mathbb{R}^n \rightarrow [0, \infty)$ is called the *joint (probability) density* of the random variables $X_1, \dots, X_n$ if, for all rectangles

$$R = (a_1, b_1] \times \cdots \times (a_n, b_n] \subseteq \mathbb{R}^n,$$

the following holds:

$$\mathbb{P}((X_1, \dots, X_n) \in R) = \int_R f(x_1, \dots, x_n) dx_1 \cdots dx_n.$$

Example I:

Find the probability of $\mathbb{P}(X \leq \frac{2}{3}, Y \leq \frac{1}{2})$, with the constraint $\Delta = \{(x, y) \in [0, 1]^2 : 0 \leq y \leq x \leq 1\}$

$$\begin{aligned} \int_0^{\frac{2}{3}} \int_0^{\frac{1}{2}} f(x, y) dy dx &= \int_0^{\frac{2}{3}} \int_0^{\frac{1}{2}} \mathbb{1}_\Delta dy dx \\ &= 2 \int_0^{\frac{2}{3}} \int_0^{\frac{1}{2}} \mathbb{1}_{dy dx} \\ &= 2 \int_0^{\frac{2}{3}} \int_0^{\min(x, \frac{1}{2})} \mathbb{1}_{dy dx} \\ &= 2 \left(\int_0^{\frac{1}{2}} x dx + \int_{\frac{1}{2}}^{\frac{2}{3}} \frac{1}{2} dx \right) = \frac{5}{12} \end{aligned}$$

Example II:

Find the densities of X and Y

$$\int_0^1 \int_0^1 6 \cdot x^2 \cdot y dy dx = 1$$

Density of X :

$$f_X(x) = \int_0^1 6x^2 y dy = 6x^2 \left[\frac{y^2}{2} \right]_0^1 = 3x^2 \quad \text{for } 0 \leq x \leq 1$$

$$f_X(x) = \begin{cases} 3x^2 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{else} \end{cases}$$

Density of Y

$$f_Y(y) = \int_0^1 6x^2 y dx = 6y \left[\frac{x^3}{3} \right]_0^1 = 2y \quad \text{for } 0 \leq y \leq 1$$

$$f_Y(y) = \begin{cases} 2y & \text{if } 0 \leq y \leq 1 \\ 0 & \text{else} \end{cases}$$

70.0.7 Multivariate Expected Value for Continuous Random Variables

Let $X_1, \dots, X_n$ be random variables with joint density $f(x_1, \dots, x_n)$, or equivalently, with joint distribution $P_{X_1, \dots, X_n}$.

Let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a measurable function. Then, if the expectation exists, the following holds:

$$\mathbb{E}[u(X_1, \dots, X_n)] = \int_{\mathbb{R}^n} u(x_1, \dots, x_n) f(x_1, \dots, x_n) dx_1 \cdots dx_n$$

70.0.8 Uniform Distribution on a Measurable Set

Let $I \subseteq \mathbb{R}^n$ be a measurable set. Random variables $(X_1, \dots, X_n) \in \mathbb{R}^n$ with joint density

$$f(x_1, \dots, x_n) = \frac{1}{|I|} \mathbb{1}_I(x_1, \dots, x_n)$$

are called *uniformly distributed*, where $|I|$ denotes the volume (Lebesgue measure) of the set I . The distribution $P_{X_1, \dots, X_n}$ is called the *uniform distribution*.

70.0.9 Probability over Measurable Sets

Let $X_1, \dots, X_n$ be random variables with joint density $f(x_1, \dots, x_n)$. For every measurable set $R \subseteq \mathbb{R}^n$, the following holds:

$$\mathbb{P}((X_1, \dots, X_n) \in R) = \int_R f(x_1, \dots, x_n) dx_1 \cdots dx_n$$

70.0.10 Joint Distribution Function for multiple Variables

Let $X_1, \dots, X_n$ be random variables. The function $F : \mathbb{R}^n \rightarrow \mathbb{R}$, defined by

$$F(x_1, \dots, x_n) = \mathbb{P}(X_1 \leq x_1, \dots, X_n \leq x_n),$$

is called the *joint distribution function* of $X_1, \dots, X_n$.

70.0.11 Marginal Densities

Let $X_1, \dots, X_n$ be random variables with joint density $f(x_1, \dots, x_n)$. The density of X_1 is given by

$$f_{X_1}(x_1) = \int \cdots \int f(x_1, \dots, x_n) dx_2 \cdots dx_n$$

For random variables X, Y with joint density $f(x, y)$, the following holds:

$$f_X(x) = \int f(x, y) dy$$

$$f_Y(y) = \int f(x, y) dx$$

The idea for higher dimensions is to eliminate all variables we are not interested in.

70.0.12 Expectation with Respect to a Joint Density

Let $X_1, \dots, X_n$ be random variables with joint density $f(x_1, \dots, x_n)$, and let $u : \mathbb{R}^n \rightarrow \mathbb{R}$ be a measurable function. Then the following holds:

$$\mathbb{E}[u(X_1, \dots, X_n)] = \int u(x_1, \dots, x_n) f(x_1, \dots, x_n) dx_1 \cdots dx_n$$

70.0.13 Stochastic Independence of Random Variables

The random variables $X_1, \dots, X_n$ are *stochastically independent* if, for all intervals $I_1, \dots, I_n \subseteq \mathbb{R}$, the following holds:

$$\mathbb{P}(X_1 \in I_1, \dots, X_n \in I_n) = \prod_{k=1}^n \mathbb{P}(X_k \in I_k)$$

70.0.14 Independence of Densities

Let X, Y be discrete random variables with joint probability function $p_{X,Y}(x, y)$ and marginal probability functions $p_X(x)$ and $p_Y(y)$. Then X and Y are *independent if and only if*

$$p_{X,Y}(x, y) = p_X(x)p_Y(y)$$

Let X and Y be discrete random variables with joint probability function $p_{X,Y}(x, y)$ and marginal probability functions $p_X(x)$ and $p_Y(y)$. The random variables X and Y are said to be *independent* if and only if

$$p_{X,Y}(x, y) = p_X(x)p_Y(y)$$

Also, if $X_1, \dots, X_n$ are independent random variables and

$$f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R}$$

Then are $f_1(X_1), \dots, f_n(X_n)$ also independent random variables.

Example:

Given

$$\int_0^1 \int_0^1 6 \cdot x^2 \cdot y \, dy \, dx = 1$$

with

$$f_X(x) = \int_0^1 6x^2 y \, dy = 6x^2 \left[\frac{y^2}{2} \right]_0^1 = 3x^2 \quad \text{for } 0 \leq x \leq 1$$

$$f_Y(y) = \int_0^1 6x^2 y \, dx = 6y \left[\frac{x^3}{3} \right]_0^1 = 2y \quad \text{for } 0 \leq y \leq 1$$

We test $f_{X,Y}(x, y) \stackrel{?}{=} f_X(x) \cdot f_Y(y)$

$$f_X(x) \cdot f_Y(y) = 3x^2 \cdot 2y = 6x^2 y = f_{X,Y}(x, y)$$

70.0.15 Variance for Independent Random Variables

Given the independent random variables $X_1, \dots, X_n$ then

$$\text{var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{var}(X_i)$$

Proof:

For $i \neq j$ we have due to independence

$$\mathbb{E}[(X_i - \mathbb{E}[X_i])(X_j - \mathbb{E}[X_j])] = \mathbb{E}[X_i - \mathbb{E}[X_i]] \cdot \mathbb{E}[X_j - \mathbb{E}[X_j]] = 0$$

This implies that

$$\begin{aligned} \text{var} \left(\sum_{i=1}^n X_i \right) &= \mathbb{E} \left[\left(\sum_{i=1}^n X_i - \mathbb{E}[X_i] \right)^2 \right] \\ &= \mathbb{E} \left[\sum_{i=1}^n (X_i - \mathbb{E}[X_i])^2 + \sum_{i \neq j} (X_i - \mathbb{E}[X_i])(X_j - \mathbb{E}[X_j]) \right] \\ &= \sum_{i=1}^n \mathbb{E}[(X_i - \mathbb{E}[X_i])^2] + \sum_{i \neq j} \mathbb{E}[(X_i - \mathbb{E}[X_i])(X_j - \mathbb{E}[X_j])] \\ &= \sum_{i=1}^n \text{var}(X_i) \end{aligned}$$

70.0.16 Conditional Distribution of Discrete Random Variables

Let X, Y be discrete random variables and $x \in \mathbb{R}$ such that $\mathbb{P}(X = x) > 0$. The mapping

$$A \mapsto \mathbb{P}(Y \in A \mid X = x),$$

for measurable $A \subseteq \mathbb{R}$, defines a probability distribution on $\mathbb{R}$ and is called the *conditional distribution of Y given $X = x$* .

The corresponding *conditional probability function* of Y given $X = x$ is defined by

$$p_{Y|X}(y \mid x) = \mathbb{P}(Y = y \mid X = x) = \frac{p_{X,Y}(x, y)}{p_X(x)}.$$

Two conditional discrete random variables X and Y are independent if the conditional probability of Y given $X = x$ is not dependent on x i.e.

Proof:

$$X, Y \text{ independent} \iff p_{X,Y}(x, y) = p_X(x)p_Y(y)$$

$$p_{Y|X}(y|x) = \frac{p_{X,Y}(x, y)}{p_X(x)} = \frac{p_X(x)p_Y(y)}{p_X(x)} = p_Y(y)$$

Otherwise, if $p_{Y|X}(y \mid x) = \mathbb{P}(Y = y \mid X = x)$ is not dependent on x then

$$p_{Y|X}(y|x) = \mathbb{P}(Y = y) = p_Y(y)$$

Then $p_{X,Y}(x, y) = p_X(x)p_{Y|X}(y \mid x) = p_X(x)p_Y(y)$

70.0.17 Conditional Density for Continuous Random Variables

Let X, Y be continuous random variables with joint density $f(x, y)$. The conditional density $f_{Y|X}(y \mid x)$ of Y given $X = x$ is defined by

$$f_{Y|X}(y \mid x) = \begin{cases} \frac{f(x, y)}{f_X(x)}, & \text{if } f_X(x) > 0, \\ 0, & \text{if } f_X(x) = 0. \end{cases}$$

The conditional distribution of Y given $X = x$ is the distribution with density $f_{Y|X}(y | x)$. Thus, for any measurable set $A \subseteq \mathbb{R}$,

$$\mathbb{P}(Y \in A | X = x) = \int_A f_{Y|X}(y | x) dy.$$

70.0.18 Covariance and Correlation of Random Variables

Let X, Y be random variables with existing variances. The *covariance*, joint dependence, is defined by

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = \sum_{i=1}^n \sum_{j=1}^m (x_i - E[X])(y_j - E[Y]) \mathbb{P}(X = x_i, Y = y_j)$$

If the variances are positive, the *correlation coefficient* $\rho_{X,Y}$ is defined as

$$\rho_{X,Y} = \frac{\text{cov}(X, Y)}{\sqrt{\text{var}(X) \text{var}(Y)}}.$$

If $\rho_{X,Y} = 0$, the random variables X and Y are called *uncorrelated*.

Properties of Covariance

- $\text{cov}(X, X) = \text{var}(X)$
- $\text{cov}(X, Y) = E[XY] - E[X]E[Y]$
- $\text{cov}(X_1 + X_2, Y) = \text{cov}(X_1, Y) + \text{cov}(X_2, Y)$
- $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2\text{cov}(X, Y)$
- If X and Y are independent, then $\text{cov}(X, Y) = 0$

Variance of the Sum of Random Variance

Given the random variables $X_1, \dots, X_n$ then

$$\text{var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{var}(X_i) + 2 \sum_{i \neq j} \text{cov}(X_i, X_j)$$

If the random variables are independent then

$$\text{var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{var}(X_i)$$

Special case: $n = 2$: $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2\text{cov}(X, Y)$

Cauchy-Schwarz Inequality for Random Variables

Let X, Y be random variables with $E[X^2]$ and $E[Y^2]$ finite. Then the following holds:

$$|E[X, Y]| \leq \sqrt{E[X^2]E[Y^2]}$$

Following

$$|\text{cov}(X, Y)| \leq \sqrt{\text{var}(X)\text{var}(Y)}$$

Correlation between minus and plus one

Let X, Y be random variables with existing variances. Then the following holds:

$$\rho_{X,Y} = 1 \iff Y = aX + b \text{ with } a > 0$$

$$\rho_{X,Y} = -1 \iff Y = aX + b \text{ with } a < 0$$

70.0.19 Multivariable Normal Distribution

Given $X_1, \dots, X_n$ independent $\mathcal{N}(0, 1)$ the density function of their joint distribution is given by

$$f(x_1, \dots, x_n) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}} e^{(-\frac{1}{2}x_i^2)} = \frac{1}{(2\pi)^{n/2}} e^{(-\frac{1}{2}X^T \cdot X)} \sim \mathcal{N}(\vec{0}, I_n)$$

Thus, for $\mu \neq \vec{0} \in \mathbb{R}^n$ and $\Sigma \in \mathbb{R}^{n \times n}$ positive definite,

$$f(x_1, \dots, x_n) = \frac{1}{(2\pi)^{\frac{d}{2}} |\Sigma|^{\frac{1}{2}}} e^{-\frac{1}{2}(X-\mu)^T \Sigma^{-1} (X-\mu)} \sim \mathcal{N}(\mu, \Sigma)$$

With d being the dimension of the space and $|\Sigma|$ being the determinant of the covariance matrix

$$\Sigma = \begin{bmatrix} \text{cov}(X_1, X_1) & \cdots & \text{cov}(X_1, X_n) \\ \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \cdots & \text{cov}(X_n, X_n) \end{bmatrix}$$

where $\mu_i = E[X_i]$ and $(X - \mu)^T \Sigma^{-1} (X - \mu)$ is a generalization of $\frac{(x-\mu)^2}{\sigma^2}$.

Computing Expected Values and Variances of Linear Combinations

Expected Value of Linear Combinations For random variables $X_1, \dots, X_n$ (not necessarily independent) with $E[X_i] = \mu_i$ and constants $a_1, \dots, a_n, b \in \mathbb{R}$, the expected value of a linear combination is always:

$$E \left[b + \sum_{i=1}^n a_i X_i \right] = b + \sum_{i=1}^n a_i \mu_i$$

This holds regardless of independence or correlation structure.

Variance of Linear Combinations The variance formula depends on whether the variables are independent:

Case 1: Independent Variables

If $X_1, \dots, X_n$ are mutually independent with $\text{var}(X_i) = \sigma_i^2$, then:

$$\text{var} \left(\sum_{i=1}^n a_i X_i \right) = \sum_{i=1}^n a_i^2 \sigma_i^2$$

Case 2: Dependent Variables (General Case)

If $X_1, \dots, X_n$ are not independent, we must account for covariances:

$$\text{var} \left(\sum_{i=1}^n a_i X_i \right) = \sum_{i=1}^n a_i^2 \sigma_i^2 + 2 \sum_{i < j} a_i a_j \text{cov}(X_i, X_j)$$

Equivalently, in matrix notation with $a = (a_1, \dots, a_n)^T$:

$$\text{var} \left(\sum_{i=1}^n a_i X_i \right) = a^T \Sigma a$$

Special Cases:

- If $\text{cov}(X_i, X_j) = 0$ for all $i \neq j$, we recover the independent case
- If $a_i = 1$ for all i (simple sum), then:

$$\text{var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \sigma_i^2 + 2 \sum_{i < j} \text{cov}(X_i, X_j)$$

- Adding a constant b does not affect variance: $\text{var}(b + Y) = \text{var}(Y)$

70.0.20 Example: Bivariate Normal Distribution

Consider the random vector

$$X = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim N_2(\mu, \Sigma),$$

with mean vector

$$\mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix},$$

and covariance matrix

$$\Sigma = \begin{pmatrix} \sigma_1^2 & \rho \sigma_1 \sigma_2 \\ \rho \sigma_1 \sigma_2 & \sigma_2^2 \end{pmatrix},$$

where $-1 < \rho < 1$ is the correlation coefficient.

Density:

The density of X is

$$f_X(x) = \frac{1}{2\pi|\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right),$$

where

$$\Sigma = \begin{pmatrix} \sigma_1^2 & \rho \sigma_1 \sigma_2 \\ \rho \sigma_1 \sigma_2 & \sigma_2^2 \end{pmatrix}$$

$$|\Sigma| = \sigma_1^2 \sigma_2^2 (1 - \rho^2),$$

and

$$\Sigma^{-1} = \frac{1}{\sigma_1^2 \sigma_2^2 (1 - \rho^2)} \begin{pmatrix} \sigma_2^2 & -\rho \sigma_1 \sigma_2 \\ -\rho \sigma_1 \sigma_2 & \sigma_1^2 \end{pmatrix}.$$

Detailed Example: Spare Parts Delivery

Consider a delivery containing:

- One box of ball bearings: $X_1 \sim \mathcal{N}(125, 1)$
- Two boxes of gears: $Y_1, Y_2 \sim \mathcal{N}(84, 4)$
- Three boxes of screws: $Z_1, Z_2, Z_3 \sim \mathcal{N}(65, 3)$

All variables are independent. The total weight is:

$$G = X_1 + Y_1 + Y_2 + Z_1 + Z_2 + Z_3$$

Step 1: Expected Total Weight

Using linearity of expectation:

$$\begin{aligned} E[G] &= E[X_1] + E[Y_1] + E[Y_2] + E[Z_1] + E[Z_2] + E[Z_3] \\ &= 125 + 84 + 84 + 65 + 65 + 65 \\ &= 488 \text{ kg} \end{aligned}$$

Step 2: Variance of Total Weight

Since all variables are independent:

$$\begin{aligned} \text{var}(G) &= \text{var}(X_1) + \text{var}(Y_1) + \text{var}(Y_2) + \text{var}(Z_1) + \text{var}(Z_2) + \text{var}(Z_3) \\ &= 1 + 4 + 4 + 3 + 3 + 3 \\ &= 18 \text{ kg}^2 \end{aligned}$$

Therefore, $G \sim \mathcal{N}(488, 18)$ with standard deviation $\sigma_G = \sqrt{18} = 3\sqrt{2} \approx 4.243 \text{ kg}$.

Step 3: Probability Calculations

Question: What is the probability that the delivery weighs more than 495 kg?

Standardize: $Z = \frac{G-488}{\sqrt{18}} \sim \mathcal{N}(0, 1)$

$$\begin{aligned} P(G > 495) &= P\left(\frac{G - 488}{\sqrt{18}} > \frac{495 - 488}{\sqrt{18}}\right) \\ &= P\left(Z > \frac{7}{\sqrt{18}}\right) \\ &= P(Z > 1.650) \\ &= 1 - \Phi(1.650) \\ &\approx 1 - 0.9505 = 0.0495 \end{aligned}$$

Thus, there is approximately a 4.95% probability that the delivery weighs more than 495 kg.

Alternative Scenario: Dependent Variables

Suppose instead that Y_1 and Y_2 are correlated with $\text{cov}(Y_1, Y_2) = 2$ (perhaps they're from the same production batch). Then:

$$\begin{aligned} \text{var}(G) &= 1 + 4 + 4 + 3 + 3 + 3 + 2\text{cov}(Y_1, Y_2) \\ &= 18 + 2(2) \\ &= 22 \text{ kg}^2 \end{aligned}$$

This increases the standard deviation to $\sqrt{22} \approx 4.690 \text{ kg}$, making extreme values more likely.

Independence of Random Variables under the Joint Normal Distribution

Given $X_1, \dots, X_n$ random variables with a joint normal distribution. If $\text{cov}(X_i, X_j) = 0$ for all $i \neq j$, then they are mutually independent.

Important: This equivalence between zero correlation and independence is special to the multivariate normal distribution and does not hold in general.

Linear Combinations of the Joint Normal Distribution

Given $X_1, \dots, X_n$ random variables with a joint normal distribution $\mathcal{N}(\mu, \Sigma)$. For any sequence of coefficients $a_1, \dots, a_n \in \mathbb{R}$, the linear combination $\sum_{i=1}^n a_i X_i$ is also normally distributed.

Specifically, if $X \sim \mathcal{N}(\mu, \Sigma)$ and $a \in \mathbb{R}^n$, then:

$$a^T X \sim \mathcal{N}(a^T \mu, a^T \Sigma a)$$

If the random variables are independent with $X_i \sim \mathcal{N}(\mu_i, \sigma_i^2)$, then:

$$b + \sum_{i=1}^n a_i X_i \sim \mathcal{N}\left(b + \sum_{i=1}^n a_i \mu_i, \sum_{i=1}^n a_i^2 \sigma_i^2\right)$$

Chapter 71

Statistics

In *probability*, we had models, and we assumed that have its ideal form. In *Statistics*, we try to estimate the model from samples. We will approximate its parameters and test hypothesis about the model.

71.0.1 Basic Nomenclature

- *Population*: Set of all objects about which we make a statement.
- *Sample*: Finite set of the Population.
- *Sample Perimeter*: Number of the samples. Like the number of participants.
- *Trait*: Special characteristic we are looking for. They can be discrete, continuous, non-numerical (can be converted to numerical).
- *Trait-quantifier*: Amount of correspondence to a trait.

71.0.2 Random Sampling

A *random sample* of size n is a sequence of n random variables $X_1, X_2, \dots, X_n$ which are independent and identically distributed (i.i.d.) with the same distribution as the random variable X .

Population Statistics

Let $x_1, x_2, \dots, x_N$ be the elements of the population.

- The *population mean* (or expectation) of X is defined as

$$\mu = \frac{1}{N} \sum_{i=1}^N x_i$$

- The *population variance* of X is defined as

$$\sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2$$

- The *population standard deviation* of X is defined as

$$\sigma = \sqrt{\sigma^2}$$

71.0.3 The 5 Number Summary

In the *descriptive statistics* we call the *5 number summary* of a data set as the list of number consisting on the: minimum, lower quantile, median, upper quantile and the maximum.

This numbers can be plotted in a box-plot

71.0.4 Sample Statistics

Let $X_1, X_2, \dots, X_n$ be a random sample of size n .

- The *sample mean* is defined as

$$\overline{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

it has the property that $\overline{X}_n \sim \mathcal{N}(\mu, \frac{\sigma^2}{n})$

- The *sample variance* is defined as

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \overline{X}_n)^2$$

- The *sample standard deviation* is defined as

$$s = \sqrt{s^2}$$

71.0.5 Finite n-Correction

$$\text{var}(\overline{X}_n) = \frac{\sigma^2}{n} \cdot \frac{n-1}{N-1}$$

for very large N .

Proof:

$$\begin{aligned} \text{var}(\overline{X}) &= \text{cov}(\overline{X}, \overline{X}) \\ &= \text{cov} \left(\frac{1}{N} \sum_{i=1}^n \overline{X}_i, \frac{1}{N} \sum_{j=1}^n \overline{X}_j \right) \\ &= \frac{1}{n^2} \left(\sum_{i=1}^n \sum_{j=1}^n \text{cov}(\overline{X}_i, \overline{X}_j) \right) \\ &= \frac{1}{n^2} \left(\sum_{i=1}^n \text{cov}(\overline{X}_i, \overline{X}_j) + \sum_{i=1}^n \text{cov}(\overline{X}_i, \overline{X}_j) \right) \\ &= \frac{1}{n^2} n \sigma^2 - \frac{n(n-1)}{N-1} \sigma^2 \\ &= \frac{\sigma^2}{n} \cdot \frac{n-1}{N-1} \end{aligned}$$

ΩεΩ

71.0.6 Frequency

Given a trait X with some quantifiers $(a_i)_{i \in I}$ for a discrete set I . Given a sample of observations $x_1, x_2, \dots, x_n$ of X . The *absolute frequency* h_i of a_i is defined as

$$h_i = \sum_{j=1}^n \mathbb{1}_{x_j=a_i} i \in I$$

The *relative frequency* is defined as

$$r_i = \frac{h_i}{n}, i \in I$$

71.0.7 Order Statistic

Given $X_1, \dots, X_n$ random variables. The random variables $X_{(1)}, \dots, X_{(n)}$ are called *order Statistic* of the variables $X_1, \dots, X_n$ if for each $\omega \in \Omega$ the following holds:

$$X_{(1)}(\omega) \leq X_{(2)}(\omega) \leq \dots \leq X_{(n)}(\omega)$$

and for each i there exist k_i such that $X_{(i)}(\omega) = X_{k_i}(\omega)$.

Note

$$X_{(1)} = \min\{X_1, \dots, X_n\}$$

$$X_{(n)} = \max\{X_1, \dots, X_n\}$$

Example:

Given 5, 1, 3, 4, 2 we get

$$x_{(1)} = 1, x_{(2)} = 2, \dots, x_{(5)} = 5$$

71.0.8 Median

Given $X_1, \dots, X_n$ random variables with their correspondent order statistic. Their *empiric median* is defined as

$$\text{med}(X_1, \dots, X_n) = \begin{cases} X_{(\frac{n+1}{2})} & n \text{ is odd} \\ \frac{1}{2} (X_{X_{n/2}} + X_{X_{n/2+1}}) & n \text{ is even} \end{cases}$$

The median is equivariant for addition

$$\text{med}(X_1 + a, \dots, X_n + a) = \text{med}(X_1, X_2, \dots, X_n) + a$$

Example:

Given 3.80, 4.33, 4.96, 3.71, 3.40. What is the median?

Order statistic: 3.40, 3.71, 3.80, 4.33, 4.96

Median: $n = 5$ thus, $\frac{n+1}{2} = 3$ our median is $\text{med}(X) = X_{(3)} = 3.80$

71.0.9 Empirical Quantile

Given $X_1, \dots, X_n$ random variables with their correspondent order statistic. The *empirical p-quantile* is defined as

$$q_p = \begin{cases} X_{[n \cdot p + 1]} & \text{if } np \notin \mathbb{N} \\ \frac{1}{2}(X_{np} + X_{np+1}) & \text{if } np \in \mathbb{N} \end{cases}$$

Note

$$\text{med}(X_1, \dots, X_n) = q_{0.5}$$

71.0.10 Modus

The most recurrent value in a set of samples is called the *modus*.

Given $x_1, \dots, x_n$ samples of discrete classes $c_1, \dots, c_n$. Define h_i as the recurrence of the class c_i in the sample set.

$$h_i = |\{x_j \in c_i : j = 1, \dots, n\}|$$

for $i = 1, \dots, d$. Then is the class c_k a modus if $h_k = \max_{i=1}^d h_i$

71.0.11 Empirical Variance

Let $X_1, \dots, X_n$ be random variables with mean $\bar{X}_n$. The *empirical variance* is defined as

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

s is called the *empirical standard deviation*.

As a side note, if our samples are distributed uniformly then

$$\text{var}(X) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x}_n)^2$$

71.0.12 Sigma Rules

Given $X \sim \mathcal{N}(\mu, \sigma^2)$

- $\mathbb{P}(\mu - \sigma \leq x \leq \mu + \sigma) \approx 68,3\%$
- $\mathbb{P}(\mu - 2\sigma \leq x \leq \mu + 2\sigma) \approx 95,4\%$
- $\mathbb{P}(\mu - 3\sigma \leq x \leq \mu + 3\sigma) \approx 99,7\%$

71.0.13 Absolute Deviation and Range

Let $X_1, \dots, X_n$ be random variables with mean $\bar{X}_n$, minimum $X_{(1)}$ and maximum $X_{(n)}$.

The range *absolute deviation* is defined as

$$\text{MAD} = \frac{1}{n} \sum_{i=1}^n |X_i - \bar{X}_n|$$

The sample *range* is defined as

$$r = X_{(n)} - X_{(1)} = \max\{X_1, \dots, X_n\} - \min\{X_1, \dots, X_n\}.$$

71.0.14 Interquartile Range

Let $X_1, \dots, X_n$ be random variables with 25% and 75% quantiles $q_{0.25}$ and $q_{0.75}$. The *interquartile range* is defined as

$$\text{IQR} = q_{0.75} - q_{0.25}.$$

71.0.15 Coefficient of Variation

Let $X_1, \dots, X_n$ be random variables with mean $\bar{X}_n$ and empirical variance s^2 . The *empirical coefficient of variation* is defined as

$$V = \frac{s}{\bar{X}_n}$$

71.0.16 Covariance

Covariance represents how much two values are dependent on each other

$$\text{cov}(X, Y) = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)(Y_i - \bar{Y}_n)$$

If the result is greater than zero they are 'moving' in the same direction and the slope is positive, less 'opposite' with negative slope and equal to zero means that they are independent, and the slope is zero.

71.0.17 Pearson Correlation Coefficient

The *pearson correlation coefficient* describes the degree of which two features are linearly correlated, and it is given by:

$$\rho_p = \frac{\text{cov}(X, Y)}{\sigma(X)\sigma(Y)}, \quad -1 < r < 1.$$

The closer the result the closer the relationship between the values. Zero means independence.

Classification of the values:

$$\rho_p \begin{cases} 1 & \text{the line has a positive slope} \\ -1 & \text{the line has a negative slope} \\ 0 & \text{the line is constant} \end{cases}$$

The individual interpretation of this values and the slope of the line is dependent on the context.

Invariance Under Linear Transformation

Given random variables X, Y , we can transform them linearly

$$\tilde{X} = a_x X - b_x, \quad \tilde{Y} = a_y Y + b_y$$

And get the following result when applying the pearson correlation coefficient

$$\rho_p(\tilde{X}, \tilde{Y}) = |\rho_p(X, Y)|$$

71.0.18 Spearman Correlation Coefficient

Given an order statistic $x_{(1)} \leq x_{(2)} \leq \dots \leq x_{(n)}$ from the sample $x_1, \dots, x_n$, we define the *rank* as:

$$\text{rank}(x_i) = k, \quad x_i = x_{(k)}.$$

We now define the *Spearman correlation coefficient* as

$$\rho_s = \frac{\sum_{i=1}^n \left(\text{rank}(x_i) - \frac{n+1}{2} \right) \left(\text{rank}(y_i) - \frac{n+1}{2} \right)}{\sqrt{\sum_{i=1}^n \left(\text{rank}(x_i) - \frac{n+1}{2} \right)^2 \sum_{i=1}^n \left(\text{rank}(y_i) - \frac{n+1}{2} \right)^2}}$$

This coefficient is used to measure non-linear, monotone correlation.

Invariance Strictly Monotone Transformations

Given random variables X, Y , we can transform them using strictly monotone functions

$$\tilde{X} = g(X), \quad \tilde{Y} = h(Y).$$

The following holds for the Spearman correlation coefficient

$$\rho_s(\tilde{X}, \tilde{Y}) = \begin{cases} \rho_s(X, Y), & \text{if both } g \text{ and } h \text{ are either decreasing or increasing} \\ -\rho_s(X, Y), & \text{else} \end{cases}$$

71.0.19 Empirical Distribution Function

Let $X_1, \dots, X_n$ be random variables. The function

$$F_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{X_i \leq x}$$

is the *empirical distribution function*

$$F_n(a_i) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{X_j \leq a_i}$$

If the trait are sorted $a_1 < a_2 < \dots$ then

$$F_n(a_i) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{X_j \leq a_i} = \sum_{k=1}^i \frac{1}{n} \sum_{j=1}^n \mathbb{1}_{x_j = a_k} = \sum_{k=1}^i r_k$$

where r_k is the relative frequency of k

Example:

Given the samples 3, 2, 4, 3, 1, 4, 3 we have the traits 1, 2, 3, 4 with relative frequencies

- $r_1 = \frac{1}{7}$
- $r_2 = \frac{1}{7}$
- $r_3 = \frac{3}{7}$
- $r_4 = \frac{2}{7}$

Then the empirical distribution function is given by

- $F_n(1) = r_1 = \frac{1}{7}$

- $F_n(2) = r_1 + r_2 = \frac{2}{7}$
- $F_n(3) = r_1 + r_2 + r_3 = \frac{5}{7}$
- $F_n(4) = r_1 + r_2 + r_3 + r_4 = 1$

71.0.20 Absolute Frequency sum

$$nF_n(a_i) = n \sum_{k=1}^i r_k = \sum_{k=1}^i h_k$$

where h_k is the absolute frequency of k

71.0.21 Grouped Data

When dealing with a lot of data sometimes it is easier to put it into *classes* $A_1, \dots, A_k$ which represent a certain range $[a, b], [a, b)$, etc. with a corresponding *absolute frequency* h_i , *relative frequency* r_i , *class middle* $\frac{a_{i+1} - a_i}{2} = \alpha_i$.

For the next definitions, the following example table will be used.

i	Duration	h_i	r_i	F_i
1	[0, 2)	1650	0.3300	0.3300
2	[2, 4)	1111	0.2222	0.5522
3	[4, 6)	720	0.1440	0.6962
4	[6, 8)	508	0.1016	0.7978
5	[8, 10)	332	0.0664	0.8642
6	[10, 15)	427	0.0854	0.9496
7	[15, 30]	252	0.0504	1.0000

$$F_i = \sum_{k=1}^i r_k$$

Empirical Distribution Function

$$F_n(x) \approx \mathbb{P}(X \leq x)$$

Mean

$$\bar{x} \approx \frac{1}{n} \sum_{i=1}^k h_i \alpha_i = \sum_{i=1}^k r_i \alpha_i$$

Empirical Variance

$$s^2 \approx \frac{1}{n-1} \sum_{i=1}^k h_i (\alpha_i - \bar{x}_n)^2$$

Empirical Quantile

For the quantile q_p find $F_i \geq p$ and $F_{i-1} < p$ this will give you a class $A(a_i, a_{i+1}]$. Then plug into the formula.

$$q_p = a_i + \frac{p - F_{i-1}}{F_i - F_{i-1}} (a_{i+1} - a_i)$$

Example:

$i = 2$, $A_2 = [2, 4)$ with $q = 0.5$

$$2 + \frac{0.5 - 0.33}{0.5522 - 0.33}(4 - 2) = 3.53$$

Mode Class

With $A_i = \text{interval}(a_i, a_{i+1})$

$$A_i \text{ with } h_i = \max_{j=1}^k h_j$$

71.0.22 Samples via Random Variables

Let $\mathcal{X} \subset \mathbb{R}^n$ and $X : \Omega \rightarrow \mathcal{X}$ be a random variable. A *sample* is a realization of $X = (X_1, \dots, X_n)^T$, i.e., $X(\omega)$ for some $\omega \in \Omega$.

- The set $\mathcal{X}$ is called the *sample set*.
- The samples $x_1, \dots, x_n$ correspond to the coordinates of the vector

$$\begin{bmatrix} X_1(\omega) \\ \vdots \\ X_n(\omega) \end{bmatrix} = X(\omega)$$

71.0.23 Statistical Model

Let $\mathcal{X} \subset \mathbb{R}^n$ and $X : \Omega \rightarrow \mathcal{X}$ be a random variable.

- A *statistical model* is a family of probability measures P on $\mathcal{X}$.
- If $P = \{P_\theta : \theta \in \Theta\}$, then Θ is called the parameter space of P , and $\theta \in \Theta$ is referred to as a parameter.

Example:

n -fold coin toss

- Sample space: $\mathcal{X} = \{0, 1\}^n$
- $X_i \sim \text{Ber}(p)$ (with values in $\{0, 1\}$)
- $X = (X_1, \dots, X_n)^T$ is a random variable in $\{0, 1\}^n$
- Sample: $X(\omega) = (x_1, \dots, x_n) \in \{0, 1\}^n$
- Informally: “Let $x_1, \dots, x_n$ be a sample...”
- Marginal distribution of the independent tosses: $\mathbb{P}_p(X_1 = 1) = p$
- Joint distribution: $\mathbb{P}_p^n$
- Statistical model for the coin toss:

$$P = \{\mathbb{P}_p^n : p \in (0, 1)\}$$

71.0.24 Estimators

Let $X : \Omega \rightarrow \mathcal{X}$ be a random variable with statistical model $P = \{P_\theta : \theta \in \Theta\}$.

1. A mapping $T_n : \mathcal{X} \rightarrow \Theta$, or the random variable $T_n(X)$, is called an *estimator* for θ .
2. An estimate is a realization of the estimator, i.e., $T_n(X)(\omega)$ for some $\omega \in \Omega$.
3. If for the true distribution of X we have $\mathbb{P}_X = \mathbb{P}_{\theta_0}$, then θ_0 is called the true parameter.

In simpler terms, an estimator is a function that takes in our samples and gives us an estimate of the parameter we are looking for.

71.0.25 Consistence of an Estimate

If $\hat{\theta}_n$ is an estimate of θ based on the sample size n . Then $\hat{\theta}_n$ is consistent if

$$\mathbb{P}(|\hat{\theta}_n - \theta| < \varepsilon) \rightarrow 1 \quad \text{as } n \rightarrow \infty$$

This means that $\hat{\theta}_n$ is an *unbiased estimate*.

71.0.26 Distortion

Let $X : \Omega \rightarrow \mathcal{X}$ be a random variable with statistical model $P = \{P_\theta : \theta \in \Theta\}$. For $\theta \in \Theta$, let E_θ denote the expectation with respect to P_θ , and let var_θ denote the corresponding variance.

- The *bias* of an estimator T_n is

$$\text{Bias}_{T_n}(\theta) = E_\theta[T_n(X)] - \theta$$

- An estimator is called *unbiased* if $\text{Bias}_{T_n}(\theta) = 0$ for all $\theta \in \Theta$, i.e.,

$$E_\theta[T_n(X)] = \theta$$

- A sequence of estimators $(T_n)_{n \in \mathbb{Z}}$ is called consistent if for all $\varepsilon > 0$ and all $\theta \in \Theta$,

$$\lim_{n \rightarrow \infty} \mathbb{P}_\theta(|T_n(X) - \theta| > \varepsilon) = 0$$

71.0.27 Method of Moments Estimators

Let $X : \Omega \rightarrow \mathcal{X}$ be a random variable with $X = (X_1, \dots, X_n)^T$ for independent, identically distributed random variables $X_1, \dots, X_n$.

The *method of moments* estimates the target distribution parameters based on the moments of the sample. The first moment is the expectation, the second moment is the variance, etc.

Based on the first and second moments, we have the following *unbiased and consistent estimators*:

Given $X_1, \dots, X_n$ random variables from a population with mean μ and variance σ^2 . Then for some $i \in 1, \dots, n$

$$E[X_i] = \mu$$

$$\text{var}[X_i] = \sigma^2$$

$$\sigma[X_i] = \sigma$$

71.0.28 Expected Value, Standard Deviation and Variance of the Sample Mean

$$E[\overline{X_n}] = \mu$$

$$\sigma(\overline{X_n}) = \frac{\sigma}{\sqrt{n}}$$

$$\text{var}(\overline{X_n}) = \frac{\sigma^2}{n}$$

Errors in the Method of Moments

Approximating the error when estimating the mean with the sample mean or variance can be challenging depending on the size of the population and the kind distribution of the data.

Sometimes we can use our methods for unbiased and consistent estimators, but they might not be accurate enough.

Another method is bootstrapping, which consists of resampling with replacement from the sample and calculating the estimator

71.0.29 Sample Variance as Estimator

The *sample variance* is given by

$$\hat{\sigma}_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = \frac{1}{n} \sum_{i=1}^n (X_i^2) - \bar{X}_n^2$$

And

$$E[\hat{\sigma}^2] = \frac{\sigma^2}{n} \cdot \frac{N(n-1)}{N-1}$$

Proof:

$$E[X_i^2] = \text{var}(X_i) + E[X_i]^2 = \sigma^2 + \mu^2$$

From this fact we can derive that

$$\begin{aligned} E[\hat{\sigma}^2] &= E \left[\frac{1}{n} \sum_{i=1}^n X_i^2 - \bar{X}_n^2 \right] \\ &= \frac{1}{n} \sum_{i=1}^n E[X_i^2 - \bar{X}_n^2] \\ &= \frac{1}{n} \sum_{i=1}^n [(\text{var}(X_i) + E[X_i]^2 - \text{var}(\bar{X}_n) - E[\bar{X}_n]^2)] \quad \text{apply the n-correction result} \\ &= \frac{1}{n} \sum_{i=1}^n \left[\sigma^2 + \mu^2 - \frac{\sigma^2}{n} \left(1 - \frac{n-1}{N-1} \right) - \mu^2 \right] \\ &= \frac{\sigma^2}{n} \left[n - \left(1 - \frac{n-1}{N-1} \right) \right] \\ &= \frac{\sigma^2}{n} \cdot \frac{N(n-1)}{N-1} \end{aligned}$$

ΩΕΩ

71.0.30 Normal Approximation to Distribution of the Sample Mean

Let $X_1, \dots, X_n$ be random variables from a population with mean μ and variance σ^2 . The *sample mean* $\bar{X}_n$ is approximately normally distributed with mean μ and variance $\frac{\sigma^2}{n}$ for large n .

Note, that this fact comes from the *Central Limit Theorem* which only applies for independent random variables. In our case because our population is very large we can assume independence due to the covariance between samples being very small.

This approximation gives us the ability to calculate probabilities about the sample mean using the normal distribution.

$$\mathbb{P}\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq z\right) \approx \Phi(z)$$

71.0.31 Linear Regression

Given samples $(x_1, y_1), \dots, (x_n, y_n)$ of random variables X and Y . The *linear regression* of Y on X is an approximation using the linear function

$$Y = aX + b$$

where we minimize *the sum of squared errors*

$$Q(a, b) = \sum_{i=1}^n (y_i - (ax_i + b))^2$$

by differentiating with respect to a and b .

$$\frac{\partial Q}{\partial a} = \sum_{i=1}^n ax_i^2 + bx_i - y_i x_i = 0$$

$$\frac{\partial Q}{\partial b} = \sum_{i=1}^n ax_i + b - y_i = 0$$

We proceed by defining the following terms

$$S = \sum_{i=1}^n x_i^2$$

$$R = \sum_{i=1}^n y_i x_i$$

Then

$$\frac{\partial Q}{\partial a} = \sum_{i=1}^n ax_i^2 + bx_i - y_i x_i = aS + b\overline{X_n} - R$$

$$\frac{\partial Q}{\partial b} = \sum_{i=1}^n ax_i + b - y_i = a\overline{X_n} + b - \overline{Y_n}$$

The Hessian is given then by

$$H = \begin{bmatrix} S & n\overline{X_n} \\ n\overline{X_n} & n \end{bmatrix}$$

With $\det(H_f) = n \sum_{i=1}^n x_i^2 - (\sum_{i=1}^n x_i)^2 > 0$

In general, this comes from

$$\begin{bmatrix} S & n\overline{X_n} \\ n\overline{X_n} & n \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} R \\ n\overline{Y_n} \end{bmatrix}$$

by solving for a and b we get using the inverse of the Hessian

$$\begin{bmatrix} a \\ b \end{bmatrix} = \frac{1}{nS - n^2\overline{X_n}^2} \begin{bmatrix} n & -n\overline{X_n} \\ -n\overline{X_n} & S \end{bmatrix} \begin{bmatrix} R \\ n\overline{Y_n} \end{bmatrix}$$

Thus, our square error is minimized when $S > 0, \det(H_f) > 0$

With a simple calculation we get

$$b = \frac{s_{XY}}{s_X^2} \quad a = \overline{Y_n} - b\overline{X_n}$$

where the *variance*

$$s_X^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \overline{X_n})^2$$

and the *covariance*

$$s_{XY} = \frac{1}{n-1} \sum_{i=1}^n (x_i - \overline{X_n})(y_i - \overline{Y_n})$$

Simplified Linear Regression

Ordinary least square with $y = b_0 + b_1x + \dots$ or for the simple case $\hat{y} = mx + b$.

- $m = b_1 = \frac{\sum_{i=1}^n (X_i - \overline{X_n})(Y_i - \overline{Y_n})}{\sum_{i=1}^n (X_i - \overline{X_n})^2}$
- $b = b_0 = \overline{Y_n} - b_1\overline{X_n}$

71.0.32 Monte Carlo Sampling

When we can not estimate the error of our estimators analytically, we can use *Monte Carlo Sampling* by assuming that our estimator $\hat{\theta}$ is true. Then we create simulated data from our estimated parameter, and we repeat the process k times to examine how our estimator behaves across the different simulations.

This not really done on paper, but with computers.

71.0.33 Likelihood

The *Likelihood* represents how likely a certain parameter θ is given in some **independent** data. We can change the target parameter or parameters and see how the likelihood changes. If the likelihood increases the parameter is more probable given the data.

For example, if we have data which we assume to be normally distributed, we can calculate the likelihood of different μ and σ values given the data. The higher the likelihood the more probable that parameter is given in the data.

Example:

A coin has been flipped 4 times as we observed the following data: H H H H. Our intuition tells us that the coin is biased; thus we want to find the probability $p \in [0, 1]$ of getting heads, with counter probability $1 - p$ of getting tails.

Our sequence is given by

$$\mathbb{P}(X = HHHH) = p^4$$

As we increase p the likelihood of observing our data increases. Thus, the maximum likelihood is achieved when $p = 1$.

71.0.34 Maximum Likelihood

Let $X = (X_1, \dots, X_n)^T : \Omega \rightarrow \mathcal{X}$ be independent random variables with statistical model $P = \{P_\theta^n : \theta \in \Theta\}$. Furthermore, let $x_1, \dots, x_n$ be a sample.

The function $L : \Theta \rightarrow [0, 1]$, defined by

$$L(\theta) = \begin{cases} \mathbb{P}_\theta(X_1 = x_1) \cdots \mathbb{P}_\theta(X_n = x_n) & \text{for disc. random vars.} \\ f_\theta(x_1) \cdots f_\theta(x_n) & \text{for cont. random vars.} \end{cases}$$

is called the *likelihood function*.

The parameter $\hat{\theta}$ that maximizes $L(\theta)$ is called the *maximum-likelihood estimate*.

The mapping $\hat{\theta}(x_1, \dots, x_n)$ is called the *maximum-likelihood estimator*.

The idea is that given some data we want to find the parameter θ that allows us to achieve the highest probability of observing the data in the ideal probability distribution.

Log-Likelihood-Estimator

The likelihood function can be pretty challenging to optimize, therefore, the less complicated *Log-Likelihood-function* is often used for maximization problem.

$$l(\theta) = \begin{cases} \sum_{i=1}^n \log(\mathbb{P}_\theta(X_i = x_i)) & \text{for disc. random vars.} \\ \sum_{i=1}^n \log(f_\theta(x_i)) & \text{for cont. random vars.} \end{cases}$$

Example:

Given $X_1, \dots, X_n \sim \text{Poisson}(\lambda)$

$$f(x_1, \dots, x_n | \lambda) = \prod_{i=1}^n \frac{\lambda^{x_i} e^{-\lambda}}{x_i!}$$

We can calculate the log-likelihood function with $\theta = \lambda$ as

$$\begin{aligned} l(\lambda) &= \prod_{i=1}^n \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \\ &\iff \ln \left(\prod_{i=1}^n \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \right) \\ &= \sum_{i=1}^n \ln \left(\frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \right) \\ &= \sum_{i=1}^n \ln(\lambda^{x_i} e^{-\lambda}) - \ln(x_i!) \\ &= \sum_{i=1}^n \ln(\lambda^{x_i}) + \ln(e^{-\lambda}) - \ln(x_i!) \\ &= \sum_{i=1}^n x_i \ln(\lambda) - \lambda - \ln(x_i!) \end{aligned}$$

Now we can derive with respect to λ and set to zero to find the maximum.

$$\frac{dl(\lambda)}{d\lambda} = \sum_{i=1}^n \frac{x_i}{\lambda} - 1 = 0$$

The second derivative is

$$\frac{d^2 l(\lambda)}{d\lambda^2} = \sum_{i=1}^n -\frac{x_i}{\lambda^2} < 0$$

Always less than zero, thus we have a maximum.

The first derivative gives us

$$\begin{aligned}\sum_{i=1}^n \frac{x_i}{\lambda} - 1 &= 0 \\ \frac{1}{\lambda} \sum_{i=1}^n [x_i] - n &= 0 \\ \frac{1}{\lambda} \sum_{i=1}^n [x_i] &= n \\ \lambda &= \frac{1}{n} \sum_{i=1}^n x_i = \overline{X_n}\end{aligned}$$

We get the sample mean as the maximum likelihood estimator for λ in a Poisson distribution.

Properties of the Maximum Likelihood Estimator

Given

$$l(\theta) = \sum \ln(\mathbb{P}(x_i | \theta))$$

where $\hat{\theta} = \operatorname{argmax}_{\theta} l(\theta)$. We get the following properties

- $\hat{\theta}$ is consistent, $\hat{\theta} \rightarrow \theta_{\text{true}}$ for $n \rightarrow \infty$
- $\hat{\theta} \sim \mathcal{N}\left(\theta_{\text{true}}, \frac{1}{nI(\theta_{\text{true}})}\right)$ where $I(\theta) = \mathbb{E}\left[\left(\frac{\partial}{\partial \theta} \ln(\mathbb{P}(x | \theta))\right)^2\right] = -\mathbb{E}\left[\frac{\partial^2}{\partial \theta^2} \ln(\mathbb{P}(x | \theta))\right]$
- *Crammer-Rao Inequality*: Given $X_1, \dots, X_n$ i.i.d. random variables with the unbiased estimator $\hat{\theta} = \hat{\theta}(X_1, \dots, X_n)$

$$\operatorname{var}(\hat{\theta}) \geq \frac{1}{nI(\theta)}$$

for all $\mathbb{P}(X | \theta)$ smooth.

Bayesian Maximum A posteriori Estimation

Recall the MLE but, we model the probability for the observations of the samples as $\mathbb{P}(x_i | \theta)$ and treat them as real conditional probabilities instead as a notational trick for probability of x with the parameter θ then we get

$$\mathbb{P}(\theta | x) = \frac{\mathbb{P}(x | \theta)\mathbb{P}(\theta)}{\mathbb{P}(x)}$$

If we optimize θ we get

$$MAP = \max \log(\mathbb{P}(x | \theta)\mathbb{P}(\theta))$$

the *maximum a posteriori estimation*.

71.0.35 Confidence Interval

A *confidence interval* p for μ is a random interval centered at $\overline{X}_n$ that contains the true mean μ with probability p .

$$\mathbb{P}(-\epsilon \leq \overline{X}_n - \mu \leq \epsilon) = p = 1 - \alpha$$

Given $\Phi(z) = 1 - \alpha$

If we standardize the expression we get

$$\mathbb{P}\left(-Z_{\frac{\alpha}{2}} \leq \frac{\overline{X}_n - \mu}{\sigma/\sqrt{n}} \leq Z_{\frac{\alpha}{2}}\right) = p = 1 - \alpha$$

This allows us to find the confidence interval given an α by: Finding the critical value $Z_{\frac{\alpha}{2}}$ from the Z -table.

This is done by looking for $1 - \frac{\alpha}{2}$ in the body of the table and finding the corresponding z value.

In general

$$(x, y) = \overline{X}_n \pm Z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}$$

Example:

Given *Confidence* = 0.95, $\alpha = 0.05$, $\overline{X}_n = 8978814.2$

The critical probability is $1 - \alpha/2 = 0.975$ which leads us to look at the Z -table it gives us 1.96.

Next we have $\sigma = 12471455.74$ and $n = 15$

giving the result

$$(2667401.97, 15290226.43) = \overline{X}_n \pm Z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}$$

71.0.36 Confidence Interval Formal Definition

Let $X = (X_1, \dots, X_n)^T : \Omega \rightarrow \mathcal{X}$ be a random variable with statistical model $P = \{\mathbb{P}_\theta^n : \theta \in \Theta\}$. Furthermore, let $x_1, \dots, x_n$ be a sample and let $\alpha \in [0, 1]$.

A mapping I that assigns to each sample $x = (x_1, \dots, x_n)^T \in X$ a subset $I(x) \subset \Theta$ is called a $(1 - \alpha)$ -confidence region for θ if

$$\mathbb{P}_\theta(\theta \in I(X_1, \dots, X_n)) \geq 1 - \alpha$$

for all $\theta \in \Theta$.

If $\Theta \subset \mathbb{R}$ and $I(x)$ is an interval for all $x \in X$, then I is called a $(1 - \alpha)$ -*confidence interval*.

71.0.37 t_{n-1}

The t_{n-1} distribution is used when the standard deviation σ is not known. It is defined as

$$t_{n-1} = \frac{\overline{X}_n - \mu}{s_n/\sqrt{n}}$$

This distribution is symmetric to the y -axis and approximates the standard normal distribution for large n .

71.0.38 Confidence Intervals for the Expected Value and Standard Deviation of Normal Distributions

Confidence Interval for μ with known σ :

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$. We can build a confidence interval for μ as follows:

$$\left[\overline{X}_n - \frac{q\sigma}{\sqrt{n}}, \overline{X}_n + \frac{q\sigma}{\sqrt{n}} \right]$$

where q is the $(1 - \alpha/2)$ -quantile of the standard normal distribution. This is computed using the formula

$$q = \Phi^{-1}(1 - \alpha/2)$$

If σ^2 is not known, we can use the s_n^2

Confidence Interval for μ with unknown σ :

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$. We can build a confidence interval for μ as follows:

$$\left[\overline{X}_n - \frac{t_{n-1, \alpha/2} s_n}{\sqrt{n}}, \overline{X}_n + \frac{t_{n-1, \alpha/2} s_n}{\sqrt{n}} \right]$$

where $t_{n-1, \alpha/2}$ is the $(1 - \alpha/2)$ -quantile of the t-distribution with $n - 1$ degrees of freedom.

Confidence Interval for σ with known μ :

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$. We can build a confidence interval for σ as follows:

$$\left[\frac{\sum_{i=1}^n (X_i - \mu)^2}{\chi_{n-1, 1-\alpha/2}^2}, \frac{\sum_{i=1}^n (X_i - \mu)^2}{\chi_{n, \alpha/2}^2} \right]$$

Confidence Interval for σ with unknown μ :

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$. We can build a confidence interval for σ as follows:

$$\left[\frac{(n-1)s_n^2}{\chi_{n-1, 1-\alpha/2}^2}, \frac{(n-1)s_n^2}{\chi_{n-1, \alpha/2}^2} \right]$$

71.0.39 Degrees of Freedom

The *degrees of freedom* of a random variable is the number of independent values or quantities which can be assigned to a statistical distribution.

For example, if we have n samples, and we know the mean, then we have $n - 1$ degrees of freedom because the last value is determined by the mean and the other $n - 1$ values.

71.0.40 Hypothesis Testing

Given $X : \Omega \rightarrow \mathcal{X}$ be random variable with statistical model $P = \{\mathbb{P}_\theta : \theta \in \Theta\}$. A subset Θ_i of Θ with $H_1 \in \Theta_i$ is called a *hypothesis*.

The *null-hypothesis* is denoted by $H_0 : \theta \in \Theta_0$ and the *alternative hypothesis* by $H_1 : \theta \in \Theta_1$. The *hypothesis test* is a decision rule to decide whether to reject H_0 or not based on the sample.

Example:

We want to test if a coin is fair.

- $\Theta = [0, 1]$

- $\Theta_0 = \{0.5\}, H_0 : p = 0.5$
- $\Theta_1 = [0, 1] \setminus \{0.5\}, H_1 : p \neq 0.5$

71.0.41 Test Statistic

Given $X : \Omega \rightarrow \mathcal{X}$ be random variable with statistical model $P = \{\mathbb{P}_\theta : \theta \in \Theta\}$. A *test statistic* is a mapping $\varphi : \mathcal{X} \rightarrow \{0, 1\}$ where $\varphi(x) = 1$ means that we reject H_0 and $\varphi(x) = 0$ means that we do not reject H_0 .

Right Assertions

- *Type I Assertion:* Accepting H_0 when it is true.
- *Type II Assertion:* Rejecting H_0 when it is false.

Error Types

- *Type I Error:* Rejecting H_0 when it is true. The probability of this error is called the *significance level* α .
- *Type II Error:* Not rejecting H_0 when it is false. The probability of this error is called β .

We can sadly not minimize both errors at the same time, therefore, we choose a significance level α and try to minimize β .

71.0.42 Quality, Power and Level

Let $X : \Omega \rightarrow \mathcal{X}$ be random variable with statistical model $P = \{\mathbb{P}_\theta : \theta \in \Theta\}$. For the test $\varphi : \mathcal{X} \rightarrow \{0, 1\}$ of the hypothesis $H_0 : \theta \in \Theta_0$ against $H_1 : \theta \in \Theta_1$:

Quality Function

The function $\beta_\varphi : \Theta \rightarrow [0, 1]$ defined by

$$\beta_\varphi(\theta) = \mathbb{P}_\theta(\varphi(X) = 1)$$

is called the *quality function* of the test φ .

Properties of the quality function:

- The probability of a Type I error is given by $\beta_\varphi(\theta) = \mathbb{P}_\theta(\varphi(X) = 1)$ for $\theta \in \Theta_0$
- The probability of a Type II error is given by $1 - \beta_\varphi(\theta) = \mathbb{P}_\theta(\varphi(X) = 0)$ for $\theta \in \Theta_1$

Significance Level

For $\alpha \in [0, 1]$ the test φ is called a *significance level* α test if

$$\sup_{\theta \in \Theta_0} \beta_\varphi(\theta) \leq \alpha$$

$\sup_{\theta \in \Theta_0} \beta_\varphi(\theta)$ is called the *level* of the test

Power

For $\theta \in \Theta_1$ is $\beta_\varphi(\theta)$ called the *power* of the test φ at θ .

Note the following:

- In practice, we choose a significance level α and look for a test φ with level $\leq \alpha$ that maximizes the power $\beta_\varphi(\theta)$ for $\theta \in \Theta_1$.
- A good test should have for $n \rightarrow \infty$, $\beta_\varphi(\theta) \rightarrow 1$

Ideal Matrix

	H_0 true	H_0 false
Reject H_0	$1 - \beta_\varphi(\theta) \geq 1 - \alpha$	$1 - \beta_\varphi(\theta)$
Do not reject H_0	$\beta_\varphi(\theta) \leq \alpha$	$\beta_\varphi(\theta)$

71.0.43 Critical Value

Given $X : \Omega \rightarrow \mathcal{X}$ be random variable with statistical model $P = \{\mathbb{P}_\theta : \theta \in \Theta\}$. For the test $H_0 : \theta \in \Theta_0$ vs. $H_1 : \theta \in \Theta_1$ and test φ with level α . A value $c_\alpha \in \mathbb{R}$ is called a *critical value* if

$$\varphi(x) = \begin{cases} 1 & \text{if } t(x) > c_\alpha \\ 0 & \text{else} \end{cases}$$

71.0.44 Test-Statistic

Given $X : \Omega \rightarrow \mathcal{X}$ be random variable with statistical model $P = \{\mathbb{P}_\theta : \theta \in \Theta\}$. $t : \mathcal{X} \rightarrow \mathbb{R}$ is called a *test-statistic* for the test $H_0 : \theta \in \Theta_0$ vs. $H_1 : \theta \in \Theta_1$ and c_α the critical value for the choice rule

$$\text{Reject } H_0 \text{ if } t(x) > c_\alpha$$

of test φ with level α .

71.0.45 p-Value

The *p-value* of a test-statistic t for the observation x is defined as

$$p = \sup_{\theta \in \Theta_0} \mathbb{P}_\theta(t(X) \geq t(x)) = 1 - \mathbb{P}_\theta(t(X) < t(x))$$

where θ is a parameter under the null-hypothesis H_0 , $t(x)$ is the value of the test-statistic for the observation x and X is the random variable.

The p-value gives us the probability of observing a test-statistic value as extreme or more extreme than the one observed, assuming that the null hypothesis H_0 is true.

- A small p-value means the observed data would be rare under H_0
- A large p-value means the data are consistent with H_0

Example:

We use the number of heads in n coin tosses to test if the coin is fair. $X \sim \text{Bin}(n, p)$

Our hypotheses are

- $H_0 : p = 0.5$
- $H_1 : p \neq 0.5$

We define our test φ as follows:

$$\varphi(x) = \begin{cases} 1 & \text{if } |X - \frac{n}{2}| > k \\ 0 & \text{else} \end{cases}$$

$\frac{n}{2}$ is the expected value for our random variable X under H_0 .

Our quality function is given by

$$\begin{aligned}
\beta_\varphi(p) &= \mathbb{P}_p(\varphi(X) = 1) = \mathbb{P}_p(|X - \frac{n}{2}| > k) \\
&= 1 - \mathbb{P}_p(|X - \frac{n}{2}| \leq k) \\
&= 1 - \mathbb{P}_p(\frac{n}{2} - k \leq X \leq \frac{n}{2} + k) \\
&= \sum_{i=\lfloor \frac{n}{2} \rfloor - k}^{\lfloor \frac{n}{2} \rfloor + k} \binom{n}{i} p^i (1-p)^{n-i}
\end{aligned}$$

To find the level of the test we compute for $n = 100$ and $k = 13$, and get $\beta_\varphi(\frac{1}{2}) = 0.0066$. Which is less than 0.05 thus our test is a level 0.05 test.

In summary, we have:

- **Significance Level:** $\alpha = 0.01$
- **Sample:** $x = (x_1, \dots, x_{100}) \in \{0, 1\}^{100}$
- **Test-Statistic:** $t(x) = \left| \sum_{i=1}^{100} x_i - 50 \right|$
- **Critical Value:** $c_\alpha = 13$ If we get heads 33 times then H_0 will be rejected since $t(x) = |33 - 50| = 17 > 13 = c_\alpha$
- **p-Value:**

$$p = \mathbb{P}_{1/2}(t(X) \geq 17) = \frac{1}{2^{100}} \sum_{i=0}^{33} \binom{100}{i} + \sum_{i=67}^{100} \binom{100}{i} \approx 0.0087$$

Due to $p = 0.0087 < \alpha = 0.01$ our observation seems to be rare under H_0 thus we reject H_0 .

71.0.46 Standard Error

The *standard error* (SE) of a statistic is the standard deviation of its sampling distribution or an estimate of that standard deviation. It is often used to measure the precision of a sample mean by estimating the variability between sample means that one would obtain if multiple samples were drawn from the same population.

$$SE = \frac{\sigma}{\sqrt{n}}$$

71.0.47 General Procedure for Hypothesis Testing

You are given a sample $X = (X_1, \dots, X_n)$ and you want to test some hypothesis about the distribution of the data. Mostly, you will have n , the sample size, α the significance level and some information about the distribution like the mean or variance.

The general procedure is as follows:

- Compute the missing information about the distribution (mean, variance, standard deviation, etc.) as needed.
- State the null and alternative hypothesis H_0 and H_1 . This will influence the test statistic and rejection region.
- Determine the test statistic and the rejection region based on the hypotheses and significance level (Z-test, Chi-Test, t-Test, etc.).
- Compute the test statistic value from the sample data.
- Using the significance level α and the test statistic distribution, find the critical value(s) for the rejection region.

- Compute the p-value for the test statistic value if needed.
- Compute the quality function if needed.

We will take a look at the kind of tests that can be performed.

71.0.48 Hypothesis Testing on Normal Distributed Random Variables

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$ i.i.d. random variables. We would want to test the following hypotheses:

- $H_0 : \mu = \mu_0$ vs. $H_1 : \mu \neq \mu_0$ (two-sided)
- $H_0 : \mu \leq \mu_0$ vs. $H_1 : \mu > \mu_0$ (right-sided)
- $H_0 : \mu \geq \mu_0$ vs. $H_1 : \mu < \mu_0$ (left-sided)

If σ^2 is known we use the *Gauss-Test*, otherwise we use the *t-Test*.

Gauss-Test (Z-Test)

Given $X = (X_1, \dots, X_n) \in \mathcal{X}$. We denote with z_α the α -quantile of $\mathcal{N}(0, 1)$. We also have the test-statistic

$$Z = \frac{\overline{X_n} - \mu_0}{\frac{\sigma}{\sqrt{n}}}$$

if $\mu = \mu_0 : Z \sim \mathcal{N}(0, 1)$

Test Type	Rejection Region	Critical Value	p-Value
Two-Sided	$ Z > z_{1-\alpha/2}$	$z_{1-\alpha/2}$	$2(1 - \Phi(z))$
Right-Sided	$Z > z_{1-\alpha}$	$z_{1-\alpha}$	$1 - \Phi(z)$
Left-Sided	$Z < -z_{1-\alpha}$	$-z_{1-\alpha}$	$\Phi(z)$

Use Case: This test is used when we want to know if the mean of a normal distribution is equal to some value μ_0 when the variance σ^2 is known.

Error Types in Gauss-Test

[h!]	Error Type I	Error Type II
$H_0 : \mu \leq \mu_0$	$1 - \Phi\left(z_{1-\alpha} + \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right)$	$\Phi\left(z_{1-\alpha} \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right)$
$H_0 : \mu \geq \mu_0$	$\Phi\left(z_\alpha + \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right)$	$1 - \Phi\left(z_\alpha + \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right)$
$H_0 : \mu = \mu_0$	$2 - \xi$	$\xi - 1$

with $\xi = \Phi\left(z_{1-\frac{\alpha}{2}} + \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right) + \Phi\left(z_{1-\frac{\alpha}{2}} - \sqrt{n} \frac{\mu_0 - \mu}{\sigma}\right)$

Example:

Given a sample of size $n = 400$ $\mu_0 = 1.74m$ and $\sigma_0^2 = 0.1m$ from height measurement of men in a city. We get a sample mean of $\overline{X_n} = 1.72m$. We want to test if the mean height is less than $1.74m$ with a significance level of $\alpha = 0.05$.

Following hypotheses are stated:

- $H_0 : \mu \geq 1.74m$
- $H_1 : \mu < 1.74m$

We compute the test-statistic

$$Z = \frac{\overline{X_n} - \mu_0}{\frac{\sigma}{\sqrt{n}}} = \frac{1.72 - 1.74}{\frac{\sqrt{0.1}}{\sqrt{400}}} = -4$$

The critical value is given by $z_{0.05} = -1.645$. Since $-4 < -1.645$ we reject H_0 and conclude that the mean height is less than $1.74m$.

t-Test

Given $X = (X_1, \dots, X_n) \in \mathcal{X}$. We denote with $t_{n-1, \alpha}$ the α -quantile of the t_{n-1} distribution. We also have the test-statistic

$$T = \sqrt{n} \frac{\overline{X_n} - \mu_0}{s_n}$$

if $\mu = \mu_0 : T \sim t_{n-1}$

[h!] Test Type	Rejection Region	Critical Value	p-Value
Two-Sided	$ T > t_{n-1, 1-\alpha/2}$	$t_{n-1, 1-\alpha/2}$	$2(1 - F_{t_{n-1}}(t))$
Right-Sided	$T > t_{n-1, 1-\alpha}$	$t_{n-1, 1-\alpha}$	$1 - F_{t_{n-1}}(t)$
Left-Sided	$T < -t_{n-1, 1-\alpha}$	$-t_{n-1, 1-\alpha}$	$F_{t_{n-1}}(t)$

Use Case: This test is used when we want to know if the mean of a normal distribution is equal to some value μ_0 when the variance σ^2 is unknown.

Example:

Given a sample of size $n = 25$ $\mu_0 = 100$, and we observed the following data:

[h!] 102	98	96	100	101
99	97	100	102	100
98	97	96	101	99
100	98	97	101	99
100	102	98	97	101

We want to test if the mean is different from 100 with a significance level of $\alpha = 0.05$.

We compute the sample mean $\overline{X_n} = 99.32$ and the sample standard deviation $s_n = 1.683$.

The test-statistic gives us

$$T = \sqrt{n} \frac{\overline{X_n} - \mu_0}{s_n} = \sqrt{25} \frac{99.32 - 100}{1.683} = -1.618$$

The critical value is given by $t_{24, 0.975} = 2.064$. Since $|-1.618| < 2.064$ we do not reject H_0 and conclude that the mean is not significantly different from 100.

71.0.49 Between-Sample t-Tests

Given two independent samples $X_1, \dots, X_n \sim \mathcal{N}(\mu_X, \sigma_X^2)$ and $Y_1, \dots, Y_m \sim \mathcal{N}(\mu_Y, \sigma_Y^2)$.

Define $D_i := X_i - Y_i \sim \mathcal{N}(\mu_X - \mu_Y, \sigma_{X+Y}^2)$.

Use Case: The null hypothesis is given by $H_0 : \mu_X = \mu_Y$ vs. $H_1 : \mu_X \neq \mu_Y$.

We get the test-statistic

$$T = \frac{\overline{X_n} - \overline{Y_m}}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}} \sim \mathcal{N}\left(\frac{\mu_X - \mu_Y}{\sqrt{\frac{\sigma_X^2}{n} + \frac{\sigma_Y^2}{m}}}, 1\right)$$

We reject H_0 if $|T| > z_{1-\alpha/2}$.

Case with Unknown Variances

If the σ_X^2 and σ_Y^2 are unknown we use the test-statistic

$$T = \frac{\overline{X_n} - \overline{Y_m}}{\sqrt{\frac{m+n}{mn}} s^2}$$

with

$$s^2 = \frac{1}{n+m-2}((n-1)s_X^2 + (m-1)s_Y^2)$$

where s_X^2 and s_Y^2 are the sample variances of X and Y .

Use Case: Reject H_0 if $|T| > t_{n+m-2, 1-\alpha/2}$.

71.0.50 Chi-Square-Test

Given $X_1, \dots, X_n \sim \mathcal{N}(\mu, \sigma^2)$ i.i.d. random variables.

Use Case: We want to test $H_0 : \sigma^2 = \sigma_0^2$ vs. $H_1 : \sigma^2 \neq \sigma_0^2$.

If μ is known we use the test-statistic

$$T_0 := \frac{1}{\sigma_0^2} \sum_{i=1}^n (X_i - \mu)^2 \sim \chi_n^2$$

Reject H_0 if $|T_0| > \chi_{n, 1-\alpha/2}^2$ or $|T_0| < \chi_{n, \alpha/2}^2$.

If μ is unknown we use the test-statistic

$$T_1 := \frac{(n-1) \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2}{\sigma_0^2} \sim \chi_{n-1}^2$$

Reject H_0 if $|T_1| > \chi_{n-1, 1-\alpha/2}^2$ or $|T_1| < \chi_{n-1, \alpha/2}^2$.

Example:

Given $n = 18$, $\bar{x} = 109$, $s = 13.5$, $\mu = 100$ and $\sigma = 15$. We want to test if the standard deviation is less than 15 with a significance level of $\alpha = 0.05$.

We state the hypotheses:

- $H_0 : \sigma^2 \geq 15^2$
- $H_1 : \sigma^2 < 15^2$

We compute the test-statistic

$$T_1 = \frac{(n-1)s^2}{\sigma_0^2} = \frac{17 \cdot 13.5^2}{15^2} = 16.38$$

The critical value is given by $\chi_{17, 0.05}^2 = 8.672$. Since $16.38 > 8.672$ we do not reject H_0 and conclude that the standard deviation is not significantly less than 15.

71.0.51 Bayesian Inference

The big idea is

$$\mathbb{P}(X | \theta) \rightarrow \mathbb{P}(\theta | X) = \frac{\mathbb{P}(X | \theta)\mathbb{P}(\theta)}{\mathbb{P}(X)}$$

where $\mathbb{P}(X | \theta)$ is the *likelihood* and $\mathbb{P}(\theta)$ is the *prior*.

We want to create derive a statistical model for some collected data with its corresponding parameters using prior beliefs.

Bayesian Update

Iterative *Bayesian Update Process*

$$\mathbb{P}_{k+1}(\theta \mid X) = \frac{\mathbb{P}_k(X \mid \theta)\mathbb{P}_k(\theta)}{\mathbb{P}_k(X)}$$

with $\mathbb{P}_{k+1} = \mathbb{P}_k(\theta \mid X)$ and its base value is our defined prior.

Chapter 72

Information Theory

Information theory is a branch of applied mathematics and electrical engineering involving the quantification of information.

72.0.1 Surprise

The *surprise* of an event $X = x$ is defined as

$$\text{Surprise}(x) = S(x) = \ln \left(\frac{1}{P(X = x)} \right)$$

72.0.2 Entropy

Entropy is defined as the expected surprise of a random variable X

$$\text{Entropy}(X) = H(X) = - \sum_{x \in X} P(X = x) S(x) = - \sum_{x \in X} P(X = x) \ln(P(X = x))$$

Or in a simpler way

$$\text{Entropy}(X) = \sum p(x) \ln \left(\frac{1}{p(x)} \right)$$

72.0.3 Mutual Information

Mutual Information measures how much information one random variable X contains about another random variable Y . It is defined as

$$I(X; Y) = \sum_{x \in X} \sum_{y \in Y} P(X = x, Y = y) \ln \left[\frac{P(X = x, Y = y)}{P(X = x)P(Y = y)} \right]$$

If one of the random variables is continuous then we can make bins for different range of values and then proceed to work with the bins as the discrete classes.

72.0.4 Cross Entropy

Cross Entropy is a measure of the difference between two probability distributions for a given random variable or set of events.

$$H(P, Q) = - \sum_{x \in X} P(X = x) \ln(Q(X = x))$$

where P is the true distribution and Q is the estimated distribution.

When there is no discrepancy between both distributions, the cross entropy is equal to the entropy of P .

It also has the following properties:

- *Lower Bound:*

$$H(P, Q) \geq H(P)$$

- *Asymmetry:*

$$H(P, Q) \neq H(Q, P)$$

72.0.5 KL-Divergence

Kullback-Leibler Divergence is a measure of how one probability distribution Q diverges from a second expected probability distribution P .

$$D_{KL}(P||Q) = \sum_{x \in X} P(X = x) \ln \left(\frac{P(X = x)}{Q(X = x)} \right)$$

It measures surprise from using Q to approximate P . The reverse of the cross entropy, where we use P to approximate Q .

72.0.6 Information Gain

Information Gain measures how much information a feature A gives us about the class label Y . It is defined as

$$IG(S, A) = H(S) - \sum_{v \in \text{values}(A)} \frac{|S_v|}{|S|} H(S_v)$$

where $H(S)$ is the entropy of the original set S , S_v is the subset of S for which feature A has value v , and $|S|$ and $|S_v|$ are the sizes of sets S and S_v respectively.

In the context of Trees we can think about information gain as

$$IG = H(\text{parent}) - \sum_i w_i H(\text{children}_i)$$

where w_i is the proportion of samples in child i relative to the parent node.

Example:

Suppose we have a dataset S with 10 samples, where 6 belong to class A and 4 belong to class B. The entropy of the original set S is calculated as:

$$H(S) = - \left(\frac{6}{10} \log_2 \left(\frac{6}{10} \right) + \frac{4}{10} \log_2 \left(\frac{4}{10} \right) \right) \approx 0.970$$

Now, let's say we split the dataset based on a feature A that can take two values: v_1 and v_2 . After the split, we have:

- Subset S_{v_1} with 4 samples (3 from class A and 1 from class B) - Subset S_{v_2} with 6 samples (3 from class A and 3 from class B)

The entropy for each subset is calculated as:

$$H(S_{v_1}) = - \left(\frac{3}{4} \log_2 \left(\frac{3}{4} \right) + \frac{1}{4} \log_2 \left(\frac{1}{4} \right) \right) \approx 0.811$$

$$H(S_{v_2}) = - \left(\frac{3}{6} \log_2 \left(\frac{3}{6} \right) + \frac{3}{6} \log_2 \left(\frac{3}{6} \right) \right) = 1.0$$

The weighted average entropy after the split is:

$$\sum_{v \in \text{values}(A)} \frac{|S_v|}{|S|} H(S_v) = \frac{4}{10} \cdot H(S_{v_1}) + \frac{6}{10} \cdot H(S_{v_2}) \approx 0.4 \cdot 0.811 + 0.6 \cdot 1.0 \approx 0.924$$

Therefore, the information gain from splitting on feature A is:

$$IG(S, A) = H(S) - \sum_{v \in \text{values}(A)} \frac{|S_v|}{|S|} H(S_v) \approx 0.970 - 0.924 = 0.046$$

72.0.7 Gini Impurity

Gini Impurity is a measure of how often a randomly chosen element from the set would be incorrectly labeled if it was randomly labeled according to the distribution of labels in the subset. It is calculated as

$$Gini(S) = 1 - \sum_{i=1}^C p_i^2$$

where C is the number of classes and p_i is the proportion of instances in class i in the set S .

The closer the Gini impurity is to 0, the more pure the set is.

Chapter 73

Machine Learning

Machine Learning is an area of computer science which focuses on making accurate predictions based on already gathered data and mathematical models.

73.0.1 Basic Terminology

- Historical Data: Refers to the data on which we are going to base our model.
- Feature Vector X : This some vector whose components represent the different numeric values of the characteristics of some object.
- Function f : is called *target function* $f : \mathcal{X} \rightarrow \mathcal{Y}$ which makes a perfect approximation
- Label y : Is some image of one $x \in \mathcal{X}$
- Data Sample D : this is a set of pairs (x_i, y_i)

Our goal is to find a function $g : \mathcal{X} \rightarrow \mathcal{Y}$ which approximates f

73.0.2 Solution Strategy

- The learning algorithm $\mathcal{A}$ uses data to select, from a set of candidate functions, the function g that best approximates the target function f .
- $\mathcal{H}$: Set of candidate functions (hypothesis set). (Example: linear functions)
- $g \in \mathcal{H}$ is also called the *final hypothesis*, which the algorithm selects at the end.

73.0.3 Central Learning Problems in Machine Learning

Supervised Learning

- Sought: Unknown target function.
- Data: Input–output pairs.
- Approximate the unknown target function f_{target} with a hypothesis h such that $h \approx f_{\text{target}}$.

Unsupervised Learning

- Sought: Unknown function that describes the data.
- Data: Only input data (no labels).
- Find a function f that describes the structure or distribution of the data well.

Reinforcement Learning

- Sought: Unknown strategy S that maximizes a reward.
- Data: (1) Input (2) Possible output (3) Evaluation of output (reward).

- Find a function f that approximates the optimal strategy S^* that maximizes the expected reward.

73.0.4 The Perceptron

The *perceptron* is our first model in machine learning. It consists on having a separate vector $w \in \mathbb{R}^d$ which consist of weights which are going to be associated with the components of $x \in \mathbb{R}^d$. We also define $y \in \{-1, 1\}$, and some value b to work as our border between the categories of y . We classify our prediction based on the result

$$\sum_{i=1}^d w_i x_i > b \text{ accept}$$

$$\sum_{i=1}^d w_i x_i < b \text{ reject}$$

More precisely

$$\begin{aligned} \sum_{i=1}^d w_i x_i + 1 \cdot b &> 0 \\ \sum_{i=1}^d w_i x_i + x_0 \cdot w_0 &> 0 \mid x_0 = 1 \quad w_0 = b \\ \sum_{i=0}^d w_i x_i &> 0 \\ w^T x &> 0 \end{aligned}$$

Finally, to get our respective y we define the following function

$$\text{sign}(w^T x) = \begin{cases} 1, & \sum_{i=1}^d w_i x_i > 0 \\ -1, & \sum_{i=1}^d w_i x_i < 0 \end{cases}$$


And we get our final perceptron equation

$$h(x) = \text{sign}(w^T x)$$

For the case $d = 2$, our vector w defines a *line*.

$$\begin{aligned} w_0 \cdot 1 + w_1 x_1 + w_2 x_2 &= 0 \\ x_2 &= -\frac{w_1}{w_2} x_1 - \frac{w_0}{w_2} \\ x_2 &= mx_1 + c \end{aligned}$$

Here a visualization



Note that y is not used for the visualization, as it is just using in the sign function to as for the classification.

The Perceptron Learning Algorithm

The *PLA* modifies w until the data are linearly separated, as long it is possible. It can be the case that in the current space it is not possible. Formally, if there exists a w^* such that $h(x_i) = y_i, \forall (x_i, y_i) \in \mathcal{D}$.

It goes as follows:

- *Data:* (x_i, y_i) with x being a vector $i = 1, \dots, N$.
- *Assumption:* The data are linearly separable, i.e., we can separate the data into their classes using a line ($d = 2$) or a hyperplane ($d > 2$).

Let $t = 0, 1, 2, \dots$ be the current iteration step, and let $w^{(t)}$ denote the vector w at iteration step t . Set $t = 0$ and $w^{(0)} = 0$.

1. Use the perceptron with $w^{(t)}$ and classify all data $\mathbf{x}_i, i = 1, \dots, N$. The labels predicted by the perceptron are denoted by $y_{i,\text{pred}}$.
2. Select any pair (x_j, y_j) that was misclassified (i.e., $y_{j,\text{pred}} \neq y_j$), and denote it by $(x^{(t)}, y^{(t)})$. If no such pair exists, terminate.
3. Update the weight vector w using the following rule:

$$w^{(t+1)} = w^{(t)} + y^{(t)}x^{(t)}$$

This just means taking the current w and adding the vector x times the corresponding label y which was expected.

4. Increase t by 1 and return to step 1.
5. After we got our final weights, we get our line or hyperplane by manipulating our expression $\sum_{i=0}^d x_i w_i = 0$

In Sample Errors

The *in-sample-error* is the mean of the amount of samples which were incorrectly labeled.

$$E_{in}(h) = \frac{1}{N} \sum_{i=1}^N W(h(x_i) \neq f(x_i))$$

where

$$W(\text{Expr}) = \begin{cases} 1 & \text{Expr} = \text{True} \\ 0 & \text{Expr} = \text{False} \end{cases}$$

It is the median of the errors made on the training data.

Out of Sample Errors

This refers to all errors which can occur due to the lack of data. Obviously we do not have all possible samples which will exist.

$$E_{out}(h) = P[h(x) \neq f(x)]$$

These errors have a randomness factor, or in other words an unknown probability distribution. They are harder to calculate due to their nature.

73.0.5 Pocket Algorithm

The *pocket algorithm* is an extension of the perceptron learning algorithm to handle non-linear separable data-sets. This is achieved by setting a maximum number of iterations while also keeping track of the vector with the lowest in-sample error value.

1. Initialize w and set $w_{\text{best}} = w$.
2. Compute the current in-sample error $E_{\text{best}} = E_{\text{in}}(w)$.
3. For each iteration:
 - (a) If a point (x_i, y_i) is misclassified, update:

$$w \leftarrow w + y_i \tilde{x}_i$$

- (b) Compute the new error $E_{\text{in}}(w)$.
- (c) If $E_{\text{in}}(w) < E_{\text{best}}$, then:

$$w_{\text{best}} \leftarrow w, \quad E_{\text{best}} \leftarrow E_{\text{in}}(w)$$

4. After all iterations, output w_{best} .

We use the following formula for error.

$$E_{\text{in}}(w) = \frac{1}{N} \sum_{i=1}^N \mathbb{I}(\text{sign}(w^T \tilde{x}_i) \neq y_i)$$

73.0.6 Preparing the Data Set

The data set $\mathcal{D}$ we split into training and test data to minimize our margin of error. If a model is trained only used the same data again, and again it will become less reliable with new data which differs from its training set. Making out of sample errors difficult to spot.

Another way of improving the quality of the data set is to add a certain degree of noise to our sample data. Thus, converting y into a probability-variable dependent on x . Therefore, $f \rightarrow P(y | x)$.

73.0.7 Linear Models Types of Supervised Learning

We distinguish between three types of supervised learning, which differ in their target function.

We also define our linear model as

$$\theta(s(x)) = \theta(\langle w, x \rangle)$$

This just means that we handle our data with linear functions.

Use cases:

1. *Classification*: discrete classes. $\theta(\cdot) = \text{sign}(\cdot)$
2. *Regression*: real numbers. $\theta(\cdot) = \text{id}(\cdot) = \cdot$
3. *Logistic Regression*: probabilities. $\theta(s) = \frac{e^s}{1+e^s}$

73.0.8 Classes of Classifications

The most common cases we encounter when we want to find our specific categories are:

- *Linear Separation* Data can be separated into classes via a line or hyperplane.
- *Non-Linear* Due to noise in the data.
- *Non-Linear* Due to a non-linear target function.

73.0.9 Linear Regression

From linear algebra we already know that the way of doing *linear regression* is via the method of the least squares.

$$w = (X^T X)^{-1} X^T y = X^\dagger y$$

The error-kinds for our model are going to be


$$E_{out}(h) = \mathbb{E}[(h(x) - y)^2]$$

$$E_{in}(h) = \frac{1}{N} \sum_{n=1}^N (h(x_n) - y_n)^2$$

Our hypothesis set $\mathcal{H}$ is composed of hypotheses of the form $h(x) = \langle x, w \rangle, x_0 = 1$. Our goal is to find some vector w such that E_{in} is minimal $w_{lin} = \arg \min_{w \in \mathbb{R}^d} E_{in}(w)$

$$E_{in}(w) = \frac{1}{N} \sum_{n=1}^N (\langle w, x \rangle - y_n)^2$$

and $w_{lin} := X^\dagger y$.



73.0.10 Non-Linear Transformations

The two algorithms covered until now have the condition that both of them are linear, but what if our data can not be linearly separated. We can modify the input variable x_i in a non-linear way to be able to separate the data linearly in another space and then apply our algorithms.

For this purpose, we define the new space $\mathcal{Z}$ whose elements are $z_n = \phi(x_n)$, where $\phi : \mathcal{X} \rightarrow \mathcal{Z}$ is a non-linear transformation. We also define $\hat{g}(z) = \text{sign}(\langle \hat{w}, z \rangle)$. Finally, our hypothesis set $\mathcal{H}_\phi$ contains functions of the form $h(x) = \hat{h}(\phi(x))$.

This can become very complex but for 3d or 2d cases with data visualization a non-linear transformation can be found using the mathematical technique of looking with your own eyes.

73.0.11 Approximation and Generalization

When trying to approximate the target function f we encounter the problem that our data may not give us the exact function because of the lack of data. To want to minimize the in-sample error but also make out approximation as general as possible.

Approximation: Small in-sample-error

Generalization: $E_{\text{out}} \approx E_{\text{in}}$

73.0.12 Bias and Variance

Fitting the training data well, but making poor predictions, is called the *bias-variance* trade-off.

The *bias* is a measurement of the difference between all the possible data predicted via our final hypothesis g the target function f . In other words, how much our approximations deviates from the actual perfect prediction.

The *variance* measures how much a final hypothesis g gotten from one specific data-set differ from one another.

There are two extreme cases: one is when the hypothesis-set has only one function, therefore we have zero variance and a big bias; and the other case is when the set is large and different data-sets produce similar functions, therefore we get a near zero bias but a high variance.

With this in mind we now define

$$E_{\text{out}} = \text{Bias} + \text{Variance}$$

Example:

Our target is $\sin(\pi x)$, and our hypothesis-sets are $\mathcal{H}_0$ the set of all horizontal lines $h(x) = b$ and $\mathcal{H}_1$ the set of all lines $h(x) = ax + b$. The training-sets are two data-points of the interval $[-1, +1]$.

73.0.13 Learning Curves

Learning curves plot E_{in} and E_{out} depending on the number of the data-points.

73.0.14 Overfitting

Overfitting refers to the situation when a model adapts itself too much to a given data-set. Precisely, it is the process in which hypotheses h continuously reduce the in-sample error, but leading to a bigger output-of-sample error.

We define *overfitting* for a hypothesis h , if for another hypothesis h' the following is true

$$E_{\text{in}}(h) < E_{\text{in}}(h') \text{ but } E_{\text{out}}(h) > E_{\text{out}}(h')$$

In summary, noise in the data (stochastic or deterministic) have a negative effect on the learning process and can lead to overfitting. To address this problem we will use *regularization*.

73.0.15 Underfitting

Underfitting refers to the case when the model has not been trained with enough data to make accurate approximations.

73.0.16 The Augmented Error

We define the *augmented error* as

$$E_{\text{aug}}(w) = E_{\text{in}}(w) + \frac{\lambda}{N} w^T w$$

λ is the parameter for how much we should regularize, $w^T w$ is the punish-term based on the weight of the vector and N is the number of data points.

73.0.17 Regularization

Regularization is the process of making the algorithm less sensible to data out of the sample our in other words, helps the learning algorithm to minimize the output-of-sample error. This is accomplished by different methods like adding error terms, special procedures etc.

In general, with regularization we want to decrease the complexity of our approximations to the target function to minimize overfitting.

Ridge Regression

This is a regularization technique used specially in linear models to prevent overfitting by discouraging the model from assigning excessively large values to its parameters.

The regularization using $\Omega(h) = w^T w$ we call is also called *weight decay* or *ridge regression*.

Example:

This example uses linear regression. Therefore,

$$\begin{aligned} E_{\text{in}}(h) &= \frac{1}{N} \sum_{n=1}^N (h(\phi(x_n)) - y_n)^2 \\ E_{\text{in}}(h) &= \frac{1}{N} \sum_{n=1}^N (w^T z_n - y_n)^2 \\ E_{\text{in}}(h) &= \frac{1}{N} (Zw - y)^T (Zw - y) \end{aligned}$$

And

$$Z := \begin{bmatrix} z_1^T \\ z_2^T \\ \vdots \\ z_n^T \end{bmatrix}$$

Our augmented error is given by

$$\begin{aligned} E_{\text{aug}}(w) &= E_{\text{in}} + \frac{\lambda}{N} w^T w \\ &= \frac{1}{N} (Zw - y_n)^T (Zw - y) + \frac{\lambda}{N} w^T w \end{aligned}$$

To minimize it we want to set its gradient to 0 which gives us

$$\begin{aligned} \nabla_w E_{\text{aug}}(w) = 0 &\implies Z^T (Zw - y) + \lambda w = 0 \\ Z^T Zw - Z^T y + \lambda w &= 0 \\ (Z^T Z - Z^T I)w + Z^T y &= 0 \\ w_{\text{reg}} &= (Z^T Z + \lambda I)^{-1} Z^T y \end{aligned}$$

Finally, as a remainder the formula without regularization

$$w_{\text{lin}} = (Z^T Z)^{-1} Z^T y$$

73.0.18 Validation

Validation refers to process of evaluating our algorithms and models based on specific parameters and procedures.

The hypothesis gained from a model which is going to be validated is marked as $\overline{g}_n$.

Quote from the statistical Learn-theory

The more often you use a data-set for training a model, taking decisions, etc. The hard is to approximate the capacity of the data-set to approximate the out-of-sample error.

73.0.19 Hold-Out-Validation

Data points are randomly shuffled into training and test data.

73.0.20 Cross Validation

Cross Validation is a technique to estimate the out-of-sample error. It consists on splitting the data-set into V different sub-sets of equal size called *folds*. Then, we train our model V times, each time using $V - 1$ disjoint sub-sets for training and the remaining one for testing. The final out-of-sample error is given by the average of all the out-of-sample errors gotten from each iteration.

We can apply this technique to different models and select the one with the lowest out-of-sample error.

The nomenclature:

- Number of data points in the training set $N_{\text{train}} = \frac{V-1}{V} N$
- Number of data points in the validation set $N_{\text{val}} = \frac{1}{V} N$
- $e_n := E_{\text{val}}(\overline{g}_n) = e(\overline{g}_n(x_n), y_n)$
- Cross Validation Error $E_{V\text{-fold CV}} = \frac{1}{V} \sum_{n=1}^V E_v$

73.0.21 Leave One Out Cross Validation

This is cross validation but with leaving always one data point out, instead of a fold.

The nomenclature:

- $e_n := E_{\text{val}}(\bar{g}_n) = e(\bar{g}_n(x_n), y_n)$
- *Cross Validation Error* $E_{\text{CV}} \frac{1}{N} \sum_{n=1}^N e_n$

73.0.22 Confusion Matrix

A *confusion matrix* is a table that is often used to describe the performance of a classification model (or "classifier") on a set of test data for which the true values are known. It allows the visualization of the performance of an algorithm.

For example, in a binary classification problem, the confusion matrix is a 2x2 table that summarizes the counts of true positive TP , true negative TN , false positive FP , and false negative FN predictions made by the model.

73.0.23 Sensitivity and Specificity

Sensitivity measures the proportion of actual positives in the data that are correctly identified as such (true positive rate). It is calculated as:

$$\text{Sensitivity} = \frac{TP}{TP + FN}$$

The denominator represents the sum of the column corresponding to actual positive cases, while the numerator represents the true positive cases.

Specificity measures the proportion of actual negatives in the data that are correctly identified as such (true negative rate). It is calculated as:

$$\text{Specificity} = \frac{TN}{TN + FP}$$

The numerator is the sum of all cells which are not in row or column of positive cases, while the denominator is the sum of the row corresponding to actual negative cases plus our previous numerator.

73.0.24 Learning Principles

Ockham

The simplest model which approximates the data is also the more plausible.

Corrupted Samples

If the sample are corrupted, biased or contains errors then the model will also not be neutral and will give false approximations.

Data Snooping

If a data-set affects the learning process then its capacity of approximating the out-of-sample error was reduced.

73.0.25 Logistic Regression

Logistic Regression is a statistical model that in its basic form uses a logistic function to model a binary dependent variable. This model returns a probability value which can be mapped to two or more discrete classes.

To achieve the desired result we apply a *linear regression* and then apply the *logistic function* to the result to squish it between 0 and 1.

The logistic function is defined as

$$\theta(s) = \frac{e^s}{1 + e^s} = \frac{1}{1 + e^{-s}}$$

called the *sigmoid function* with the property

$$\theta(-s) = 1 - \theta(s)$$

Note that $h(x_i) = \theta(w^T x_i + b) = \frac{1}{1 + e^{-(w^T x_i + b)}}$

Based on the output of the logistic function we classify our data in the respective classes. For example, if the output is greater than 0.5 we classify it as class 1, otherwise as class -1.

where $h_\theta(x^i)$ is the hypothesis function, y^i is the true label for the i -th data point x^i , and N is the total number of data points.

Maximum Likelihood

We can not optimize the sigmoid function directly; thus, we use the method of maximum likelihood.

$$L = (\theta) = \prod_{n=1}^N h_\theta(x^i)$$

We take that expression and modify it into

$$L = (\theta) = \prod_{n=1}^N (h_\theta(x^i))^{y^i} (1 - h_\theta(x^i))^{1-y^i}$$

Taking the logarithm of the likelihood function gives us the log-likelihood:

$$\log L(w, b) = -\frac{1}{N} \sum_{i=1}^N [y^i \log(h_\theta(x^i)) + (1 - y^i) \log(1 - h_\theta(x^i))]$$

And this is the function we want to maximize. Note, that this is the *cross entropy loss function*

Gradient of the Cross Entropy Loss Function

Using the cross entropy we can define our error for logistic regression. By taking the gradient

$$\log L(w, b) = -\frac{1}{N} \sum_{i=1}^n [y^i \log(h_\theta(x^i)) + (1 - y^i) \log(1 - h_\theta(x^i))]$$

We get

$$\frac{\partial \log L(w, b)}{\partial w} = \frac{1}{N} \sum_{i=1}^N (h_\theta(x^i) - y^i) x^i$$

$$\frac{\partial \log L(w, b)}{\partial b} = \frac{1}{N} \sum_{i=1}^N (h_\theta(x^i) - y^i)$$

In vector form

$$\nabla_w \log L(w, b) = \frac{1}{N} X^T (h_\theta(X) - y)$$

$$\nabla_b \log L(w, b) = \frac{1}{N} \sum_{i=1}^N (h_\theta(x^i) - y^i)$$

Logistic Regression Algorithm

1. Initialize w and b (e.g., to zeros or small random values).
2. For each iteration until convergence or for a fixed number of iterations:
 - (a) Compute the predictions for all training examples:

$$y_{\text{pred}} = h_\theta(x^i) = \frac{1}{1 + e^{-(w^T x^i + b)}}$$

- (b) Compute the gradients:

$$\nabla_w \log L(w, b) = \frac{1}{N} X^T (h_\theta(X) - y)$$

$$\nabla_b \log L(w, b) = \frac{1}{N} \sum_{i=1}^N (h_\theta(x^i) - y^i)$$

- (c) Update the parameters using gradient descent:

$$w \leftarrow w - \alpha \nabla_w \log L(w, b)$$

$$b \leftarrow b - \alpha \nabla_b \log L(w, b)$$

3. After all iterations, output the learned parameters w and b .

73.0.26 Trees

Trees are used mainly for *regression trees* or classification *decision trees*. They are mostly not an individual learning model but are just an extra layer.

Formal definition: Given a feature vector $x \in X$ in a d -dimensional feature space with $R_1, R_2, \dots, R_M$ being disjoint regions *hyperquadrants* in the feature space.

The hypothesis $h(x)$ of a tree is given by the formula

$$h(x) = \sum_{m=2}^M c_m \mathbb{1}_m(x)$$

With c_m being the coefficient given as

- *Regression*: c_m is the mean of all labels in R_m .
- *Classification*: c_m is the multiplicity label of all data points in the training set R_m .

The regions can be defined by *binary splitting*.

73.0.27 Regressions Trees

A *regression tree* divides the feature space into disjoint sub-spaces in such a way that it brings the value of all labels y_i for $x_i \in R_m$ as close as possible $\overline{y_{R_m}} = c_m$.

The following is again an optimization problem: Find the partition in M hyperquadrants, such that the sum of the square distance is minimal.

$$\text{Minimize}_{M, R_1, \dots, R_M} \sum_{m=1}^M \sum_{x_i \in R_m} (y_i - \overline{y_{R_m}})^2$$

with

$$\overline{y_{R_m}} = \frac{1}{N_m} \sum_{x_i \in R_m} y_i$$

and

$$N_m = |\{x_i \in R_m\}|$$

Recursive Binary Splitting

We previously state an optimization problem, which sadly can not be solved efficiently, instead we will approach it with *recursive binary splitting* an algorithm which goes as follows

1. Given R_0 a hyperquadrant with training data (x_i, y_i) . Given a feature vector $x_i = (x_{i1}, \dots, x_{id})$. Also, given $j \in \{1, \dots, d\}$ as the indexes of the feature and $s \in \mathbb{R}$ a bias term.
2. Define the regions

$$R_1(j, s) = \{x \mid (x)_j < s, x \in R_0\}$$

$$R_2(j, s) = \{x \mid (x)_j \geq s, x \in R_0\}$$

Find (j, s) which minimizes the following

$$\theta = \min \left(\sum_{i: x_i \in R_1(j, s)} (y_i - \overline{y_{R_1}})^2 + \sum_{i: x_i \in R_2(j, s)} (y_i - \overline{y_{R_2}})^2 \right)$$

3. For the next step repeat step one, $R_0 = R_1$ or $R_0 = R_2$ depending on the smallest, else terminate.

The termination criteria are for example

- Each leaf which has to be divided has a minimal number of data points contained.
- The tree can have maximal depth.
- The tree size is limited.

73.0.28 Classification Trees

We can take everything from the regression tree except the cost function. For *classification trees* the cost function is given as the label of the specific classes itself.

Nomenclature

- C is the number of classes in the classification problem.
- $p(i)$ is the probability of the class relative to the complete set.

73.0.29 Gini Impurity

Represents the mean probability for a data point being misclassified. $p(i) = p_i$

This probability that a data point was misclassified is given by

$$\sum_{k \neq i}^C p_k = 1 - p_i$$

The mean is then

$$E_{\text{Gini}} = \sum_i^C p_i(1 - p_i) = \sum_i^C p_i - \sum_i^C p_i^2 = 1 - \sum_i^C p_i^2$$

73.0.30 Information Gain in Machine Learning

Information gain is a measure commonly used in decision tree algorithms such as ID3, C4.5, and CART to determine which feature should be used to split the dataset at each step. It quantifies how much uncertainty about the class labels is reduced after splitting the data based on a particular feature.

Variance

For regression problems, we can use variance as a measure of impurity. The variance of a set of values is defined as:

$$\text{Var}(X) = \frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})^2$$

where $\bar{x}$ is the mean of the values in set X .

Covariance

The covariance between two random variables X and Y is defined as:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y]$$

Or the

$$\text{cov} = \frac{1}{N} \sum_{i=1}^N \prod_{j=1}^n \vec{x}_j$$

where $\vec{x}_j = (x_{j1}, x_{j2}, \dots, x_{jn})^T$ is a feature vector.

Correlation

The correlation coefficient between two random variables X and Y is defined as:

$$\rho_{X,Y} = \frac{\text{cov}(X, Y)}{\sigma_X \sigma_Y}$$

where σ_X and σ_Y are the standard deviations of X and Y , respectively.

Covariance Matrix

The covariance matrix for a set of n random variables $X_1, X_2, \dots, X_n$ is defined as:

$$\Sigma = \begin{bmatrix} \text{cov}(X_1, X_1) & \text{cov}(X_1, X_2) & \dots & \text{cov}(X_1, X_n) \\ \text{cov}(X_2, X_1) & \text{cov}(X_2, X_2) & \dots & \text{cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \text{cov}(X_n, X_2) & \dots & \text{cov}(X_n, X_n) \end{bmatrix}$$

Information content

For the occurrence of some event k with probability p_k we define

$$I_k := -\log_2 p_k$$

For the intuition, the rarer an event occurs the more information it provides.

Entropy

Entropy is the mean information content of an event in a probability experiment with a random variable X

$$H(X) := E[I] = -\sum_{k=1}^C p_k \log_2 p_k$$

where p_i is the proportion of samples belonging to class i , and k is the total number of classes. High entropy indicates a mixture of classes, while low entropy indicates more homogeneous data.

For a multivariate case with two random variables we can calculate the conditional entropy

$$H(X | Y = y_i) := E[I] = -\sum_i p(x_i | y_i) \log_2 p(x_i | y_i)$$

The *conditional entropy* is defined as

$$H(X | Y) = -\sum_i p(y_i) H(X | Y = y_i)$$

Definition of Information Gain

The reduction of the information content of an event X by knowing Y is defined as:

$$IG(X, Y) = H(X) - H(X | Y)$$

Intuition

A feature produces high information gain if the resulting subsets are relatively pure, meaning they have low entropy. Conversely, features that do not help distinguish between classes result in low information gain.

Purpose

Information gain guides decision tree algorithms by selecting features that maximally reduce uncertainty at each split. This helps create more efficient, interpretable, and accurate decision trees.

73.0.31 CART Algorithm

The *CART* (Classification and Regression Trees) algorithm is a popular decision tree algorithm used for both classification and regression tasks. The CART algorithm constructs binary trees by recursively splitting the data based on feature values to minimize impurity or variance.

The steps of the CART algorithm are as follows:

1. **Select the Best Split:** For each feature, evaluate all possible split points and calculate the impurity (e.g., Gini impurity for classification or variance for regression) for each split. Choose the feature and split point that results in the lowest impurity.
2. **Create Child Nodes:** Split the dataset into two subsets based on the selected feature and split point, creating left and right child nodes.
3. **Repeat Recursively:** For each child node, repeat steps 1 and 2 until a stopping criterion is met (e.g., maximum tree depth, minimum number of samples in a node, or no further reduction in impurity).
4. **Assign Class Labels or Values:** For classification tasks, assign the most common class label to each leaf node. For regression tasks, assign the mean value of the target variable to each leaf node.

73.0.32 Variance Reduction

In regression tasks, the CART algorithm aims to minimize the variance within each node after a split. The variance reduction is calculated as follows:

$$\text{Variance Reduction} = \text{Var}(R) - \left(\frac{N_{\text{left}}}{N} \text{Var}(R_{\text{left}}) + \frac{N_{\text{right}}}{N} \text{Var}(R_{\text{right}}) \right)$$

with

$$\text{var}(Y) = \frac{1}{N-1} \sum_{i=1}^N (y_i - \bar{y})^2$$

73.0.33 Random Forest

The *random forest* algorithm is an ensemble learning method that combines multiple decision trees to improve predictive performance and reduce overfitting.

The steps to create a random forest are as follows:

1. **Bootstrap Sampling:** For each tree in the forest, a bootstrap sample is created by randomly sampling the training data with replacement. This means that some data points may be included multiple times, while others may not be included at all.
2. **Feature Selection:** At each split in the decision tree, a random subset of features is selected. This randomness helps to decorrelate the trees in the forest, making the ensemble more robust.
3. **Tree Construction:** Each decision tree is grown to its maximum depth without pruning. The splitting criteria (e.g., Gini impurity or information gain) are used to determine the best splits based on the selected features.
4. **Aggregation:** For classification tasks, the final prediction is made by majority voting among all trees in the forest. For regression tasks, the predictions from all trees are averaged to obtain the final output.

73.0.34 Quantifying the Differences between Features

Examples:

- Euclidean Distance $d(x, y) = \|x - y\|$
- Cosine Similarity $\text{sim}(x, y) = \frac{x \cdot y}{\|x\| \|y\|}$

- Jaccard Similarity $\text{sim}(A, B) = \frac{|A \cap B|}{|A \cup B|} \in [0, 1]$

73.0.35 Nearest Neighbors

Given a data-set $\mathcal{D} = \{(x_1, y_1), \dots, (x_N, y_N)\}$ and a new point x we want to predict its label y . We choose a *distance metric* $d(x, x')$ to quantify the difference between two feature vectors.

To predict the label of x we find the k nearest neighbors of x in the data-set $\mathcal{D}$ based on the distance metric d .

Feature vectors in the training set are noted with its distance to x . $x_{[1]}$ symbolizes the nearest neighbor of x , $x_{[2]}$ the second nearest and so on

$$d(x, x_{[1]}) \leq d(x, x_{[2]}) \leq \dots d(x, x_{[N]})$$

Nearest Neighbor Model

The final hypothesis is given by $g(x) = y_{[1]}(x)$

73.0.36 k-Nearest Neighbors

This is a generalization of the NN model.

Given an integer $k \geq 1$ and $y \in \{-1, 1\}$ the *k-nearest neighbor* model (binary classification) is given by

$$g(x) = \text{sign} \left(\sum_{i=1}^k y_{[i]}(x) \right)$$

Note that for $k = 1$ we get the previous NN model.

k-Nearest Neighbor Regression

The KNN regression is given by:

- The mean of the labels of the k nearest neighbor
- Final hypothesis $\frac{1}{k} \sum_{i=1}^k y_{[i]}(x)$

73.0.37 Support Vector Machine

As starting point we take the perceptron we already defined, but this time with b , the bias term as an independent term

$$y_k = \text{sign}(w^T x + b), y_k \in \{-1, 1\}$$

A *support vector machine* is a perceptron whose decision boundary or *margin* has the maximum margin to all data-points.

If some data points lie on the margin then

$$\begin{aligned} (I) w^T x_i + b &= 0 \\ (II) w^T x_j + b &= 0 \end{aligned}$$

By performing $2II - I$ we get the vector lying on the hyperplane

$$w^T (x_j - x_i) = 0$$

We also know that w^T is normal to the hyperplane.

The name *support vector classifier* comes from the fact that the observations on the edge and within the soft margin are called *support vectors*.

Rescaling of the Signal

The term inside our *sign* function, the signal, can be rescaled without changing the result of the classification. Due to the property of the sign function $\text{sign}(\alpha s) = \text{sign}(s)$ for any $\alpha > 0$.

$$y_k = \text{sign}(w^T x_k + b) \implies y_k(w^T x_k + b) > 0$$

The behavior of the rescaling acts differently on if the data point is in the margin or not.

$$y_k(w^T x_k + b) = 1 \quad \text{for points in the margin}$$

$$y_k(w^T x_k + b) > 1 \quad \text{for points outside the margin}$$

Width of the Margin

Given two points x_1, x_2 lying on the margin with different labels. Therefore,

$$\begin{aligned} (I) w^T x_1 + b &= 1 \\ (II) w^T x_2 + b &= -1 \end{aligned}$$

Then we get

$$w^T(x_1 - x_2) = 2$$

The *width* of the margin r is given by the projection of the vector $x_1 - x_2$ onto the normal vector $\frac{w}{\|w\|}$

$$r = \frac{w^T(x_1 - x_2)}{\|w\|} = \frac{2}{\|w\|}$$

Therefore, to maximize the width of the margin we have to minimize $\frac{1}{2}\|w\|$.

73.0.38 Hard Margin SVM

A *hard margin SVM*: is a perceptron with maximal margin and no misclassifications.

Like previously stated, to maximize the width $r = \frac{2}{\|w\|}$ of the margin we have to minimize $\frac{1}{2}\|w\|$.

Formally, we want to solve the following optimization problem

Minimize: $\frac{1}{2}\|w\|^2$ under N constraints

Maximize: $y_n(w^T x_n + b) \geq 1, (n = 1, \dots, N)$

We will skip the derivation of the final solution, which is done via *Karush-Kuhn-Tucker conditions*. Now, we have a *dual optimization problem*

$$\max \left(-\frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m x_n^T x_m + \sum_{n=1}^N \alpha_n \right)$$

under the condition

$$\sum_{n=1}^N \alpha_n y_n = 0$$

In the primary problem we had

$$y_n = \text{sign}(w^T x_n + b)$$

Now in the dual problem we have

$$y_i = \text{sign} \left(\sum_{n=1}^N \alpha_n y_n x_n^T x_i + b \right)$$

with α being our solution vector with dimension N .

Some components of α will be zero, and the others positive, due to it being *sparse*. Therefore

$$y_i = \text{sign} \left(\sum_{k \in \mathcal{S}} \alpha_k y_k x_k^T x_i + b \right)$$

with $\mathcal{S} = \{j \mid \alpha_j \neq 0, j = 1, \dots, N\}$.

Finally, the data points x_k with $\alpha_k \neq 0$ are called *support vectors* and the $\alpha_k \neq 0$ the *support values*.

73.0.39 Soft Margin Support Vector Machine

The idea of *soft margin SVM* is to allow misclassifications but penalize them in the optimization problem. To get the best soft margin we can use cross validation.

Formally, we want to solve the following optimization problem:

Minimize: $\frac{1}{2}w^T w + C \sum_n \xi_n$ under N constraints

In other words, maximize $\frac{1}{2}w^T w$, the width of the margin, minimize $\sum_n \xi_n$, the total error.

With the constraint $y_n(w^T x_n + b) \geq 1 - \xi_n, (n = 1, \dots, N)$

$$\xi_n = \max(0, 1 - y_n(w^T x_n + b))$$

Thus,

$$\frac{1}{2}w^T w + C \sum_n \max(0, 1 - y_n(w^T x_n + b))$$

with C is our regularization parameter.

Our *dual optimization problem* is given by

$$\max \left(-\frac{1}{2} \sum_{n=1}^N \sum_{m=1}^N y_n y_m \alpha_n \alpha_m x_n^T x_m + \sum_{n=1}^N \alpha_n \right)$$

under the conditions

$$\sum_{n=1}^N \alpha_n y_n = 0, 0 \leq \alpha_n \leq C \forall n$$

Also,

$$y_i = \text{sign} \left(\sum_{n=1}^N \alpha_n y_n x_n^T x_i + b \right)$$

73.0.40 Non-Linear Support Vector Machines

We will allow non-linear feature Transformations $x \rightarrow \Phi(x)$

$$x_n^T x_i \rightarrow \Phi(x_n)^T \Phi(x_i)$$

Now our y_i becomes

$$y_i = \text{sign} \left(\sum_{n=1}^N \alpha_n y_n \Phi(x_n)^T \Phi(x_i) + b \right)$$

Let us also $K := \Phi(x_n)^T \Phi(x_i)$

Kernel Trick

The *kernel trick* allows us to compute the inner product in the transformed feature space without explicitly performing the transformation. For that we use so-called *kernel functions*, K to use as inner products in the feature space. We are basically using mathematical elegance to use high dimensional spaces without paying the computational cost.

The kernel functions must satisfy the following properties *Mercer's Theorem*:

K is a *kernel* if the kernel matrix K is symmetric and positive semi-definite for any set of input data points.

$$x^T K x \geq 0 \quad K = K^T$$

Common Kernel Functions:

- *Linear*: $k(x, y) = x^T y$
- *Polynomial*: $k(x, y) = (x^T y + c)$
- *RBF*: $k(x, y) = \exp(-\gamma \|x - y\|^2)$
- *Sigmoid*: $k(x, y) = \tanh(\gamma x^T y + c)$

We can also give parameters α, β for $k(\beta + \alpha x^T, y)$

Example:

Given $\Phi(x_1, x_2) = (1, \sqrt{2}x_1, \sqrt{2}x_2, x_1^2, \sqrt{2}x_1x_2, x_2^2)$ we get for

$$\langle \Phi(x), \Phi(y) \rangle = 1^2 + \sqrt{2}x_1\sqrt{2}y_1 + \sqrt{2}x_2\sqrt{2}y_2 + x_1^2y_1^2 + \sqrt{2}x_1x_2\sqrt{2}y_1y_2 + x_2^2y_2^2$$

The following kernel by simplifying

$$\langle \Phi(x), \Phi(y) \rangle = 1 + 2x^T y + (x^T y)^2 = (1 + x^T y)^2 = k(x, y)$$

Generalized $(d + x^T y)^2$

73.0.41 The Radial Kernel

The *radial basis function (RBF) kernel* is defined as

$$K(x, y) = \exp \left(-\frac{\|x - y\|^2}{2\sigma^2} \right)$$

where $\sigma > 0$ is a parameter that controls the width of the Gaussian function, and σ is the standard deviation of the Gaussian distribution.

The infinite-dimensional property comes from the fact that the RBF kernel can be expressed as an infinite series expansion,

$$K(x, y) = \sum_{n=0}^{\infty} \frac{1}{n!} \left(-\frac{\|x - y\|^2}{2\sigma^2} \right)^n$$

73.0.42 Least Squares Support Vector Machine

This is a simplified version of the soft margin SVN.

Formally,

Primary Optimization Problem:

Minimize: $\frac{1}{2}w^T w + \frac{\gamma}{2} \sum_n e_n^2$ under N constraints.

$$y_n(w^T x_n + b) = 1 - e_n,$$

with

$$e_n^2 = y_n^2(y_n - (w^T x_n + b))^2 = (y_n - (w^T x_n + b))^2$$

The Lagrangian we get is

$$\mathcal{L} = \frac{1}{2}w^T w + \frac{\gamma}{2} \sum_n e_n^2 - \sum_{n=1}^N \alpha_n [y_n(w^T x_n + b) - 1 + e_n]$$

Apply $\nabla_{w,b,e,\alpha} \mathcal{L} = 0$

$$\begin{aligned} \text{I} \quad \frac{\partial \mathcal{L}}{\partial w} &= w - \sum_{k=1}^N \alpha_k y_k x_k = 0 \iff w = \sum_{k=1}^N \alpha_k y_k x_k \\ \text{II} \quad \frac{\partial \mathcal{L}}{\partial b} &= - \sum_{k=1}^N \alpha_k y_k = 0 \iff y^T \vec{\alpha} = 0 \\ \text{III} \quad \frac{\partial \mathcal{L}}{\partial e} &= \gamma e_k - \alpha_k = 0 \iff \alpha_k = \gamma e_k \iff e = \frac{1}{\gamma} \vec{\alpha} \\ \text{IV} \quad \frac{\partial \mathcal{L}}{\partial \alpha_k} &= -y_k(w^T x_k + b) + 1 - e_k = 0 \iff y_k(w^T x_k + b) + e_k = 1 \forall k \end{aligned}$$

Now let us combine our results

$$\begin{aligned} \text{I} \quad w &= \sum_{k=1}^N \alpha_k y_k x_k \\ \text{II} \quad y^T \vec{\alpha} &= 0 \\ \text{III} \quad e &= \frac{1}{\gamma} \vec{\alpha} \\ \text{IV} \quad y_k(w^T x_k + b) + e_k &= 1 \forall k \end{aligned}$$

Substitute I and III in IV

$$\begin{aligned}
y_k(w^T x_k + b) + e_k &= 1 \forall k \\
\iff y_k \left(\sum_{j=1}^N \alpha_j y_j x_j^T x_k + b \right) + \frac{1}{\gamma} \alpha_k &= 1 \forall k \\
\iff y_k b + \sum_{j=1}^N \alpha_j y_j x_j^T x_k y_k + \frac{1}{\gamma} \alpha_k &= 1 \forall k
\end{aligned}$$

Now replace the dot product with the kernel function

$$\iff \forall yb + (yy^T \cdot K)\vec{a} + \frac{1}{\gamma} I\vec{a} = 1$$

The only results left to use are II and V which we can combine into

$$A = \begin{bmatrix} 0 & y^T \\ y & \Omega + \frac{1}{\gamma} I \end{bmatrix}$$

After this process we get

$$y_i = \text{sign}(w^T x_i + b) \implies \text{sign} \left(\sum_{k=1}^N \alpha_k y_k x_k^T x_i + b \right)$$

We can apply our kernel function and get

$$y_i = \text{sign} \left(\sum_{k=1}^N \alpha_k y_k K(x_k, x_i) + b \right)$$

Least Square Support Vector Machine Implementation

1. Implement the kernel function, typically

$$K(x_k, x_i) = \exp \left(-\frac{\|x_k - x_i\|^2}{2\sigma^2} \right)$$

2. Compute the $n \times n$ matrix $\Omega = yy^T \cdot K$ with y being the label vector and K the kernel matrix.
3. Find the matrix

$$A = \begin{bmatrix} 0 & y^T \\ y & \Omega + \frac{1}{\gamma} I \end{bmatrix}$$

4. Invert A and get b and $\vec{a}$

$$\begin{bmatrix} b \\ \vec{a} \end{bmatrix} = A^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

5. Predict the labels

$$y_i = \text{sign} \left(\sum_{k=1}^N \alpha_k y_k K(x_k, x_i) + b \right)$$

73.0.43 Dimension Reduction Principal Component Analysis

It refers to the reduction of the features for the learning process.

PCA (Principal Component Analysis) finds new axes — called components — for the data, such that the data show the largest possible variance along these axes. In other words, PCA looks for directions in which the data vary the most. Axes along which the data have little, or almost no variance can usually be ignored, because they contain little useful information.

The goal is to identify the directions (principal components) along which the variation in the data is maximal. Then by projecting our data onto these new axis we reduce the dimensionality of our data while retaining most of its variance. Hence, we can apply other machine learning algorithms more efficiently.

These new axis are generated via the orthogonal projection of the vectors.

The big picture is that

1. We have N feature vectors with D features each.
2. We want to reduce the number of features from D to $M < D$
3. We have a $D \times D$ covariance matrix S
4. We compute the eigenvalues and eigenvectors of S
5. We sort the eigenvalues with their corresponding eigenvectors in decreasing order and take the first M eigenvectors
6. These eigenvectors form the new basis for the feature space.

Average Vector

Given N features vectors with D features. The *average vector* is given by

$$\bar{x} = \frac{1}{N} \sum_{n=1}^N x_n$$

PCA Formula I

Given a matrix X where each row represents a feature vector. We can perform the principal component analysis via the following steps:

- Compute the average vector $\bar{x} = \frac{1}{N} \sum_{n=1}^N x_n$
- Get the *average matrix* by computing $\bar{X} = \vec{1} \cdot \bar{x}^T$
- Subtract the mean $B = X - \bar{X}$
- Compute the covariance matrix $cov = B^T B$
- Compute the eigenvectors and values of cov . We can write them as $cov V = V D$ where V is the matrix of the eigenvectors and D the matrix made of the eigenvalues in the main diagonal.
- The principal components are given as the columns of the matrix $T = B V$. Or if we are programming it from scratch sort the indices and then take the first k eigenvectors.
- Transform the original n -dimensional data into k dimensions. Projections with the dot product can be achieved by using the eigenvector matrix to transform our vectors. $X V^T$ is our new, transformed data.

PCA Formula II

We look for the direction u_1 such that the variance of the data at that axis becomes maximal. With the constraint

$$u_1^T u_1 = 1 \iff 1 - u_1^T u_1 = 0$$

The projection of the features vectors on u_1 is given by

$$u_1^T x_n$$

The variance of the projection is given by

$$\begin{aligned}\text{Var}(u_1^T x_n) &= \frac{1}{N} \sum_{n=1}^N (u_1^T x_n - u_1^T \bar{x})^2 \\ &= u_1^T S u_1\end{aligned}$$

With the covariance matrix

$$S = \frac{1}{N} \sum_{n=1}^N (x_n - \bar{x})(x_n - \bar{x})^T$$

We are going to of course use *Lagrange Multipliers* for that

$$\mathcal{L}(u_1, \lambda_1) = u_1^T S u_1 + \lambda_1 [1 - u_1^T u_1]$$

Take the gradient and set it to 0

$$\frac{\partial \mathcal{L}}{\partial \lambda_1} = u_1^T u_1 - 1 = 0$$

$$\frac{\partial \mathcal{L}}{\partial u_1} = S u_1 - \lambda_1 u_1 = 0 \iff S u_1 \lambda_1 u_1 \iff u_1^T S u_1 = \lambda_1$$

We can do this because u_1 is an eigenvector of S .

The previous process was only for one direction (the principal component).

Now we need to find more directions u_j such that they are orthogonal to each other and u_1 such that the variance is maximized.

We do this with induction

Given $u_1, \dots, u_M$ eigenvectors of S with the corresponding eigenvalues $\lambda_1 \geq \dots \geq \lambda_M$

We want to find u_{M+1} such that

- The Variance in that direction is maximal
- u_{M+1} is orthogonal to $u_1, \dots, u_M$
- u_{M+1} is normalized $u_{M+1}^T u_{M+1} = 1$

This is again a Lagrange problem

$$\mathcal{L}(u_{M+1}, \lambda_{M+1}, \eta_i) = u_{M+1}^T S u_{M+1} + \lambda_{M+1} [1 - u_{M+1}^T u_{M+1}] + \sum_{i=1}^M \eta_i u_{M+1}^T u_i$$

With the following partial derivatives

$$\frac{\partial \mathcal{L}}{\partial \lambda_{M+1}} = 1 - u_{M+1}^T u_{M+1} = 0$$

$$\frac{\partial \mathcal{L}}{\partial \eta_j} = u_{M+1}^T u_j = 0$$

$$\frac{\partial \mathcal{L}}{\partial u_{M+1}} = 2Su_{M+1} + 2\lambda_{M+1}u_{M+1} + \sum_{i=1}^M \eta_i u_i = 0$$

We know substitute

$$\frac{\partial \mathcal{L}}{\partial u_{M+1}} = 2u_j^T Su_{M+1} + 2\lambda_{M+1}u_j^T u_{M+1} + \sum_{i=1}^M \eta_i u_j^T u_i = 0$$

From that

$$\begin{aligned} 2u_j^T Su_{M+1} + \eta_j &= 0 \text{ apply the transpose} \\ 2u_{M+1}^T S^T u_j + \eta_j &= 0 \text{ remember that } S = S^T \\ 2u_{M+1}^T S u_j + \eta_j &= 2u_{M+1}^T \lambda_j u_j + \eta_j = 0 \end{aligned}$$

To find λ_{M+1} :

$$2Su_{M+1}^T - 2\lambda_{M+1}u_{M+1} = 0$$

This means that u_{M+1} is also an eigenvector of S .

Effect of the Mean

The *mean* (or center of mass) of the data influences the PCA coordinates:

- If the data have a mean that is not at the origin of the coordinate system, then after applying PCA, the transformed data will typically have offsets in the new coordinate system. These offsets can make interpretation more difficult.
- However, the PCA directions (the principal component axes) do not change if the data's mean changes.
- This is because the covariance matrix — which PCA is based on — is invariant to changes in the mean.
- Some software implementations of PCA do not use the covariance matrix directly. In those cases, the resulting PCA might depend on the original mean of the data.
- As a recommendation, always center each feature (variable) by subtracting its mean before applying PCA. This ensures that the data have a mean of 0 and that the PCA results are interpretable and consistent.

$$\hat{x}_i = x_i - \bar{x}$$

Z Scoring

Given $\bar{X}$ as the mean and σ as the standard deviation of X . The standardize feature Z (z-score) is given by

$$Z = \frac{X - \bar{X}}{\sigma}$$

In practice, data are often organized in matrices where each column represents a feature. In this case, standardize each feature by transforming every column separately so that it has mean 0 and variance 1 (i.e., standard deviation 1).

Recap of PCA

The basic assumption behind dimensionality reduction with PCA is: the directions with the largest variance contain the most information about the data.

Dimensionality reduction with PCA means that we discard those PCA components (directions) that represent only a small amount of variance.

PCA gives us new axes defined by direction vectors $u_1, \dots, u_D$ (the principal components).

Projecting a standardized data point x_1 onto these direction vectors gives the components of the PCA coordinates of that data point.

$$z_i = \begin{bmatrix} u_1^T \\ u_2^T \\ \vdots \end{bmatrix} x_i$$

Proportion of Variance

Variance of j -th component

$$\text{Var}(Z_j) = \text{Var}(u_j^T x) = u_j^T S u_j = u_j^T \lambda_j u_j = \lambda_j$$

Total Variance

$$\text{Var}_{\text{Total}} = \sum_{j=1}^D \text{Var}(Z_j) = \sum_{j=1}^D \lambda_j$$

Proportion of Variance

$$\text{PVE}(j) = \frac{\text{Var}(Z_j)}{\text{Var}_{\text{Total}}}$$

73.0.44 Kernel PCA

This is an extension of PCA to non-linear feature spaces via the kernel trick.

Reformulation of the PCA with Non-Linear Features

The covariance matrix $S = \frac{1}{N} \sum_{n=1}^N (x_n - \bar{x})(x_n - \bar{x})^T$ can be reformulated with $x_i \rightarrow \Phi(x_i)$ to

$$S = \frac{1}{N} \sum_{n=1}^N (\Phi(x_n) - \overline{\Phi(x)})(\Phi(x_n) - \overline{\Phi(x)})^T$$

Assume that the features vectors are centered in the feature space, i.e. $\overline{\Phi(x)} = 0$. Then we have

$$S = \frac{1}{N} \sum_{n=1}^N \Phi(x_n) \Phi(x_n)^T$$

Eigenvector as Linear Combination of the transformed Features

$$\begin{aligned}
Su_j &= \lambda_j u_j \\
Su_j &= \frac{1}{N} \sum_{n=1}^N \Phi(x_n) \Phi(x_n)^T u_j \\
&= \frac{1}{\lambda_j N} \sum_{n=1}^N \Phi(x_n) (\Phi(x_n)^T u_j) \\
&= \frac{1}{\lambda_j N} \sum_{n=1}^N \Phi(x_n)^T u_j (\Phi(x_n)) \\
&= \sum_{n=1}^N \left(\frac{1}{\lambda_j N} \Phi(x_n)^T u_j \right) \Phi(x_n) \\
&= \sum_{n=1}^N a_{ji} \Phi(x_i)
\end{aligned}$$

with $u_j = \sum_l \alpha_{jl} \Phi(x_l)$

Because we can express the eigenvectors as a linear combination of the transformed features, we can find the coefficients α_{jl} instead of u_j .

Find the Eigenvectors

$$\begin{aligned}
\frac{1}{N} \sum_i \Phi(x_i) \Phi(x_i)^T \sum_l \alpha_{jl} \Phi(x_l) &= \lambda_j \sum_l \alpha_{jl} \Phi(x_l) \\
\frac{1}{N} \sum_i \Phi(x_i) \left(\sum_l \alpha_{jl} \Phi(x_i)^T \Phi(x_l) \right) &= \lambda_j \sum_l \alpha_{jl} \Phi(x_l) \\
\frac{1}{N} \sum_i \Phi(x_i) \left(\sum_l \alpha_{jl} K(x_i, x_l) \right) &= \lambda_j \sum_l \alpha_{jl} K(x_i, x_l) \\
\frac{1}{N} K K \vec{\alpha}_j &= \lambda_j K \vec{\alpha}_j \\
K \vec{\alpha}_j &= N \lambda_j K \vec{\alpha}_j
\end{aligned}$$

Therefore, we have to solve the following eigenvalue problem

$$K \vec{\alpha}_j = N \lambda_j \vec{\alpha}_j$$

Which we reformulate to

$$K \vec{\alpha}_j = \hat{\lambda}_j \vec{\alpha}_j$$

The coefficients α_j are the components of the eigenvectors of the kernel matrix K .

Normalization of the Eigenvectors

To normalize the eigenvectors u_j we have to impose the condition

$$u_j = \sum_l \alpha_{jl} \Phi(x_l) \implies \hat{u} = \sum_l \hat{\alpha}_{jl} \Phi(x_l) = \hat{u}_j^T \hat{u}_j^T = 1 \forall j$$

This gives us

$$\sum_k \sum_l \hat{\alpha}_{jk} \hat{\alpha}_{jl} K(x_k, x_l) = 1 \forall j$$

Which can be reformulated to

$$\vec{\alpha}_j^T \hat{\lambda}_j \hat{\alpha}_j = 1 \forall j \implies \|\hat{\alpha}_j\| = \frac{1}{\sqrt{\hat{\lambda}_j}}$$

With

$$\vec{\alpha}_j \rightarrow \hat{\alpha}_j = \frac{\vec{\alpha}_j}{\sqrt{\hat{\lambda}_j}} \frac{\vec{\alpha}_j}{\|\vec{\alpha}_j\|}$$

Projection onto the KPCA Axis

$$\begin{aligned} \Phi(x)^T u_j &= \Phi(x)^T \sum_l \alpha_{jl} \Phi(x_l) \\ &= \sum_l \alpha_{jl} \Phi(x)^T \Phi(x_l) \\ &= \sum_l \alpha_{jl} K(x, x_l) \end{aligned}$$

Centering of the Features

We assumed that the features were centered in the transformed feature space. Now we want to normalize them.

Given

$$\hat{\Phi}(x_i) = \Phi(x_i) - \frac{1}{N} \sum_l \Phi(x_l)$$

and the kernel function

$$\begin{aligned} K(x_i, x_j) &= \Phi(x_i)^T \Phi(x_j) \\ &= \left(\Phi(x_i) - \frac{1}{N} \sum_l \Phi(x_l) \right)^T \left(\Phi(x_j) - \frac{1}{N} \sum_l \Phi(x_l) \right) \\ &= K(x_i, x_j) - \frac{1}{N} \sum_l K(x_i, x_l) - \frac{1}{N} \sum_k K(x_k, x_j) + \frac{1}{N^2} \sum_{k,l} K(x_k, x_l) \end{aligned}$$

Simplified

$$\hat{K} = K - 1_N K - K 1_N + 1_N K 1_N$$

where 1_N is the $N \times N$ matrix with all entries equal to $\frac{1}{N}$.

73.0.45 Kernel PCA Algorithm

Now we can summarize the KPCA algorithm

1. Choose a kernel function $K(x, y)$ and compute the kernel matrix K with entries $K_{ij} = K(x_i, x_j)$.
2. Center the kernel matrix to obtain $\hat{K}$ using $\hat{K} = K - 1_N K - K 1_N + 1_N K 1_N$
3. Solve the eigenvalue problem $\hat{K} \vec{\alpha}_j = \hat{\lambda}_j \vec{\alpha}_j$ to find the eigenvalues $\hat{\lambda}_j$ and eigenvectors $\vec{\alpha}_j$.
4. Normalize the eigenvectors $\vec{\alpha}_j \rightarrow \frac{1}{\sqrt{\hat{\lambda}_j}} \frac{\vec{\alpha}_j}{\|\vec{\alpha}_j\|}$
5. Project the data points onto the KPCA axes using

$$\Phi(x)^T \hat{u}_j = \sum_l \hat{\alpha}_{jl} K(x, x_l)$$

73.0.46 Linear Discriminant Analysis

The goal of the *linear discriminant analysis* is to find directions $u_1, u_2, \dots, u_k$ in the classes such they can be best separated.

For the direction u_1 with the projection of some data point x_i onto that direction defined as

$$\hat{x}_i = u_1^T x_i$$

Let

$$m^{(k)} = \frac{1}{N_k} \sum_{n \in \mathcal{C}_k} x_n$$

be the *mean vector* of class k .

The projection onto u_1 is given by

$$\hat{m}^{(k)} = u_1^T m^{(k)}$$

Within-Class Scatter

For quantifying the separation of the points projected onto u_1 we define the following the *within-class scatter* as

$$\begin{aligned} s_w^2 &:= \sum_{n \in \mathcal{C}_k} (\hat{x}_n - \hat{m}^{(k)})^2 \\ &= \sum_{n \in \mathcal{C}_k} (u_1^T x_n - u_1^T m^{(k)})^2 \\ &= u_1^T \left[\sum_{n \in \mathcal{C}_k} (x_n - m^{(k)})(x_n - m^{(k)})^T \right] u_1 \\ &= u_1^T S_{wk} u_1 \end{aligned}$$

The *total scatter* for two classes is given by

$$\begin{aligned} s_1^2 + s_2^2 &= u_1^T (S_{w1} + S_{w2}) u_1 \\ &= u_1^T S_W u_1 \end{aligned}$$

For n -classes we have

$$S_W = \sum_{k=1}^n S_{wk} = \sum_{k=1}^n \sum_{n \in \mathcal{C}_k} (x_n - m^{(k)})(x_n - m^{(k)})^T$$

Between-Class Scatter

The quantification of the separation between the middle points, called the *between-classes-scatter* is given by

$$\begin{aligned} (\hat{m}^{(1)} - \hat{m}^{(2)})^2 &= (u_1^T m^{(1)} - u_1^T m^{(2)})^2 \\ &= u_1^T (m^{(1)} - m^{(2)})(m^{(1)} - m^{(2)})^T u_1 \\ &= u_1^T S_b u_1 \end{aligned}$$

$$S_b = (m^{(1)} - m^{(2)})(m^{(1)} - m^{(2)})^T$$

where $m^{(1)}$ and $m^{(2)}$ are the mean vectors of class 1 and class 2.

Fisher Discriminant

The *Fisher Discriminant* is the ratio of the between-class scatter and the within-class scatter.

$$J(u_1) = \frac{u_1^T S_b u_1}{u_1^T S_W u_1} = \frac{(u_1^T (m^{(1)} - m^{(2)})(m^{(1)} - m^{(2)})^T u_1)}{u_1^T S_W u_1}$$

We want to maximize $J(u_1)$. This means to find the direction u_1 such that the between-class scatter is maximized and the within-class scatter is minimized. To accomplish this we take the gradient and set it to zero.

$$\begin{aligned} \frac{dJ(u_1)}{du_1} &= u_1^T S_W u_1 \frac{d}{du_1} (u_1^T S_b u_1) - u_1^T S_b u_1 \frac{d}{du_1} (u_1^T S_W u_1) \\ &= 2S_b u_1 (u_1^T S_W u_1) - 2S_W u_1 (u_1^T S_b u_1) = 0 \quad \text{divide by } 2(u_1^T S_W u_1) \\ &\iff S_b u_1 = \frac{u_1^T S_b u_1}{u_1^T S_W u_1} S_W u_1 \\ &\iff S_b u_1 = J(u_1) S_W u_1 \\ &\iff (S_W^{-1} S_b) u_1 = J(u_1) u_1 \end{aligned}$$

We arrive at an eigenvalue problem where u_1 is the eigenvector of $S_W^{-1} S_b$ with the corresponding eigenvalue $J(u_1)$.

Linear Discriminant Algorithm

1. Compute the within-class scatter matrix S_W and the between-class scatter matrix S_b .
2. Solve the eigenvalue problem $(S_W^{-1} S_b)u = \lambda u$ to find the eigenvalues λ and eigenvectors u .
3. Sort the eigenvectors by their corresponding eigenvalues in descending order.
4. Select the top k eigenvectors to form a projection matrix W .
5. Project the original data points onto the new subspace using $z_i = W^T x_i$.

73.0.47 Naive Bayes

The *naive Bayes* is a probabilistic classifier based on Bayes' Theorem with strong (naive) independence assumptions between the features.

Recall Bayes' theorem as

$$\mathbb{P}(A | B) = \frac{\mathbb{P}(B | A) \cdot \mathbb{P}(A)}{\mathbb{P}(B)}$$

For our case we will use

$$\mathbb{P}(y | X) = \frac{\mathbb{P}(X | y) \cdot \mathbb{P}(y)}{\mathbb{P}(X)}$$

$$\mathbb{P}(y | X) = \frac{(\prod_{i=1}^n \mathbb{P}(x_i | y)) \cdot \mathbb{P}(y)}{\mathbb{P}(X)}$$

where $X = (x_1, x_2, \dots, x_n)$ and with the assumption that the features are independent.

We want to select the class with the highest posterior probability

$$y = \operatorname{argmax}_y \mathbb{P}(y | X) = \operatorname{argmax}_y \frac{(\prod_{i=1}^n \mathbb{P}(x_i | y)) \cdot \mathbb{P}(y)}{\mathbb{P}(X)}$$

$$y = \operatorname{argmax}_y \prod_{i=1}^n \mathbb{P}(x_i | y) \mathbb{P}(y)$$

$$y = \operatorname{argmax}_y \sum_{i=1}^n \ln(\mathbb{P}(x_i | y)) + \ln(\mathbb{P}(y))$$

Nomenclature:

- $\mathbb{P}(y)$ is the proper probability, which in this case is the frequency of each class
- $\mathbb{P}(x_i | y)$ is the class conditional probability which is model with a normal distribution

$$\mathbb{P}(x_i | y) = \frac{1}{\sqrt{2\pi\sigma_y^2}} \exp\left(-\frac{(x_i - \mu_y)^2}{2\sigma_y^2}\right)$$

Algorithm

1. Calculate the mean, variance and prior (frequency) for each class
2. Calculate the posterior for each class with

$$y = \operatorname{argmax}_y \sum_{i=1}^n \ln(\mathbb{P}(x_i | y)) + \ln(\mathbb{P}(y))$$

and the normal distribution formula

3. Choose the class with the highest probability

73.0.48 KMeans

KMeans is an unsupervised learning iterative process that clusters the data set into k different clusters. Each sample is assigned to the cluster with the nearest mean, and then the means, *centroids*, and clusters are updated until there is no more change or the maximum number of iterations is reached.

Intra-Cluster Variance

Given $C_1, \dots, C_n$ clusters with centroids $\mu_1, \dots, \mu_n$. The *intra-cluster variance* is given by

$$J = \frac{1}{|C_i|} \sum_{i=1}^k \sum_{x_j \in C_i} \|x_j - \mu_i\|^2$$

The goal of the KMeans algorithm is to minimize J .

Centroid

The *centroid* of a cluster C_i is given by

$$\mu_i = \frac{1}{|C_i|} \sum_{x_j \in C_i} x_j$$

Algorithm

1. Initialize cluster centers randomly
2. Repeat until converged or max iterations are reached:
 - Update cluster labels: Assign points to the nearest centroid
 - Update centroids: Set center to the mean of each cluster

The distance measure can be chosen by the developer.

Important Note on KMeans

- KMeans can converge to local minima depending on the initial centroids.
- The results can vary between different runs with different initialization.

Elbow Heuristic

The *elbow heuristic* consists of plotting the explained variance as a function of the number of clusters and looking for an "elbow" point where the rate of decrease sharply changes.

- For $1, \dots, K$ determine the clustering and the intra-cluster variance $J(k)$
- Visualize $J(k)$ against k
- Choose k at the "elbow" point where the decrease in $J(k)$ starts to slow down significantly.

73.0.49 Scaling

Feature scaling is a technique used to standardize the range of independent variables or features of data. In machine learning, it is crucial because many algorithms are sensitive to the scale of the input data.

Common methods for feature scaling include:

- *Min-Max Scaling (Normalization)*: This technique scales the data to a fixed range, usually 0 to 1. The formula is:

$$X' = \frac{X - X_{min}}{X_{max} - X_{min}}$$

By adding $(b - a)$ to the formula we can scale it to any range $[a, b]$:

$$X' = a + \frac{(X - X_{min})(b - a)}{X_{max} - X_{min}}$$

- *Standardization (Z-score Normalization)*: This method transforms the data to have a mean of 0 and a standard deviation of 1. The formula is:

$$Z = \frac{X - \mu}{\sigma}$$

where μ is the mean and σ is the standard deviation of the feature.

- *Robust Scaling*: This method uses the median and the interquartile range (IQR) for scaling, making it robust to outliers. The formula is:

$$X' = \frac{X - \text{median}}{\text{IQR}}$$

Chapter 74

Numerical Mathematics

Numerical mathematics is the study of algorithms for the problems of continuous mathematics (Lloyd Trefethen, 1992).

74.1 Floating Point Numbers

A *floating point number* of a basis b has the representation

$$z = v \cdot m \cdot b^e,$$

where

- $b \in \mathbb{N} \setminus \{1\}$.
- $v \in \{-1, 1\}$.
- m is the *mantis*, a b -adic fraction

$$m = m_{-k} \cdot \dots \cdot m_{-1} m_0 m_1 \cdot \dots \cdot m_l = m_{-k} b^k + \dots + m_l b^{-l} = \sum_{i=-k}^l m_i b^{-i}$$

with $m_i \in \{0, 1, \dots, b-1\}$.

- $e \in \mathbb{N}$.

In this representation are numbers not really well differentiable, hence 0 can not be represented properly.

Example:

$$b = 2, \quad 10.11 \cdot 2^1 = (1 \cdot 2^1 + 0 \cdot 2^0 + 1 \cdot 2^{-1} + 1 \cdot 2^{-2}) \cdot 2 = \frac{11}{2}.$$

74.1.1 Normalized Representation

The *normalized representation* of floating point numbers is given by

$$\pm(m_0 \cdot m_1 \cdot \dots \cdot m_l) b^e = \pm \left(\sum_{i=0}^l m_i b^{-i} \right) b^e, m_0 \neq 0$$

is for numbers $\neq 0$ well defined.

74.1.2 Machine Precision

Given a floating point number in a basis b with mantis length l

$$\varepsilon_M = \frac{1}{2}b^{-l}$$

is called the *machine precision*.

74.1.3 Interpretation

In the floating point system numbers do not really represent one single number x but a range of numbers close to that x in the Reals. This combined with also rounding errors generated by trade-offs in range precision lead to the well-known errors when dealing with floating point numbers.

74.2 Errors

Almost all problems in the numerical analysis can be described as zero's problem with a function g , input x and output y ; $g(x, y) = 0$

For example linear system of equations $Ax = b$ is equivalent to

$$g(x, y) = 0, \quad x = (A, b), \quad g(x, y) = Ay - b$$

Criteria For Problems in the N.A.

The 3 criteria for problems in the numerical analysis are:

1. A solution should exists. $g(x, y) = 0 \iff y = f(x)$.
2. For each x there should be exactly one y .
3. There should be a continuous dependence of y on x , i.e. $x' \rightarrow x \implies y' \rightarrow y$. Also, small changes in the input mean small changes in the output.

74.2.1 Well- and Ill-Posed Problems


Given a problem $g(x, y) = 0$ for which a solution: exists, is unique and continuous dependence are defined, so is the problem *well-posed*, if not it is *ill-posed*.

74.2.2 Error Measurement

Given a well-posed problem with x and

$$y = f(x) \quad f : X \rightarrow V,$$

where X and Y are vector spaces with metrics $\|\cdot\|_X$ and $\|\cdot\|_Y$. We denote the following scenarios:



- y is the exact solution.
- $\tilde{x}$ is the floating point representation of x .
- $\hat{y}$ is the result of solution via exact arithmetic with the exact data of x .
- $\tilde{y}$ is the floating point solution obtained by using $\tilde{x}$.
- f is the original function.
- $\hat{f}$ is the discrete (finite), simplification of the problem with which we replace f .
- $\tilde{f}$ is the floating point implementation of $\hat{f}$.

Example:

Given

$$f(x) = e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}, \quad x = \frac{5}{3}$$

With the exact solution $y = e^{\frac{5}{3}} \approx 5.29449$.

We use the substitute problem

$$\hat{f}(x) = \sum_{k=0}^4 \frac{x^k}{k!} = \frac{10009}{1944} \approx 5.14866$$

Using a three decimals places for FP we get $\tilde{x} = 1.67$ and hence, $\tilde{y} = 5.16$.

74.2.3 Error Types

Given y and $\tilde{y}$ we get the following error types:

- $e = \tilde{y} - y$ is the *error*.
- $e_a = \|\tilde{y} - y\|_Y$ is the *absolute error*, actual amount of deviation.
- $e_r = \frac{\|\tilde{y} - y\|_Y}{\|y\|_Y}$, for $\|y\|_Y \neq 0$ is the *relative error*, percentage of deviation.

The relative error is in praxis more important for analyzing numerical problems, but due to its definition when the absolute error is small it can lead to unnecessary precision constraints. For this reason for practical applications, it is preferred to use a combination of both.

Example:

Given is $t_2 = -p + \sqrt{p^2 + q}$.

First,

$$x = (p, q), \quad y = f(x), \quad f(x_1, x_2) = -x_1 + \sqrt{x_1^2 + x_2}, \quad f : \mathbb{R}^2 \rightarrow \mathbb{R}$$

In this case $\hat{f} = f$, hence we do not need a substitute problem. We use for the metric $\|y\|_Y = |y|$. For $p = 10, q = 1$ we get

$$y = \sqrt{101} - 10 \approx 0.04988, \quad \tilde{y} = 0,$$

with

$$\begin{aligned}
e &= \tilde{y} - y \approx -0.4988, \\
e_a &= |\tilde{y} - y| \approx 0.4988, \\
e_r &= \frac{|\tilde{y} - y|}{|y|} = 1
\end{aligned}$$

74.2.4 Error Criteria

Given $\varepsilon_a, \varepsilon_r$ as bounds for the absolute and relative errors. We try to find an approximation $\tilde{y}$ of y such that:

$$\|\tilde{y} - y\|_Y \leq \varepsilon_a + \|y\|_Y \varepsilon_r.$$

We call this a *mixed error criterion*.

Note that if $\|y\|_Y$ is small, then ε_a is dominant and

$$e_a = \|\tilde{y} - y\|_Y \lesssim \varepsilon_a.$$

Else, then the dominant term is ε_r resulting in

$$\|\tilde{y} - y\|_Y \lesssim \|y\|_Y \varepsilon_r$$

i.e

$$e_r = \frac{\|\tilde{y} - y\|_Y}{\|y\|_Y} \lesssim \varepsilon_r$$

Example:

Let us take the last example and add $\varepsilon_a = \varepsilon_r = 10^{-2}$.

$$y = \sqrt{101} - 10 \approx 0.04988, \quad \tilde{y} = 0$$

Then,

$$\|\tilde{y} - y\| \approx 0.04988, \quad \varepsilon_a + |y|\varepsilon_r \approx 1.04988 \cdot 10^{-2}$$

Therefore,

$$\|\tilde{y} - y\|_Y > \|y\|_Y \varepsilon_r.$$

Hence, we reject the approximation $\tilde{y}$.

74.2.5 Error Kinds

Starting from the error $\tilde{y} - y = \tilde{f}(\tilde{x}) - f(x)$, let us expand by adding zero.

$$\tilde{y} - y = \tilde{f}(\tilde{x}) - +\hat{f}(\tilde{x}) + \hat{f}(\tilde{x}) - f(\tilde{x}) + f(\tilde{x}) - f(x)$$

From this we define the following error kinds:

- *Input Error:* $f(\tilde{x}) - f(x)$. Caused by the conversion to the floating point format. It describes the sensitivity of the original problem in correlation to changes in the input data. Can not be really influenced, perspective highly depends on our choices.

- *Procedural Error*: $\hat{f}(\tilde{x}) - f(\tilde{x})$. Caused by the quality of the substitute problem. It gives a hint about the quality of the substitute problem. Can be reduced by using a better substitute problem.
- *Accumulation Error*: $\tilde{f}(\tilde{x}) - \hat{f}(\tilde{x})$. Accumulation of the rounding errors with FP. Can be handled by taking the properties of FP into consideration by ordering our operations accordingly and/or using more stable representations for decimal numbers.

In practice, all error kinds should be considered.

74.2.6 Input Error

Due to our problem being well-posed, we know that

$$\|\tilde{x} - x\|_X \implies \|f(\tilde{x}) - f(x)\|_Y < \varepsilon$$

We want to study how the the conversion to the floating point format affects the results and how we can solve our problems with if possible small divergences.

Condition

Given $f : X \rightarrow Y$. The smallest constants c_a, c_r such that:

$$\|f(x') - f(x)\|_Y \leq c_a \|x' - x\|_X \iff c_a = \lim_{\Delta x \rightarrow 0} \frac{\|\Delta y\|}{\|\Delta x\|}, \quad c_a = \|J_f(x)\|$$

$$\frac{\|f(x') - f(x)\|_Y}{\|f(x)\|} \leq c_r \frac{\|x' - x\|_X}{\|x\|_X} \iff c_r = \frac{\|x\| \|f'(x)\|}{\|f(x)\|}, \quad c_r = \frac{\|x\|}{\|f(x)\|} \|J_f(x)\|$$

with $\|x' - x\|_X \leq \delta$; are called *absolute condition* and *relative condition* of f .

The interpretation of c_a is closely related to the Lipschitz continuity, meaning that a small variation in the input data due to conversion problems does not affect the results drastically.

For c_r , it is useful for when a users wants to limit the relative error by some amount ε_r , then

$$c_r \leq \frac{\varepsilon_r}{\varepsilon_M} \implies e_r(y) \leq \varepsilon_r$$

For c_a it can be deduced that it is telling that f is a Lipschitz continuous function with a smallest Lipschitz-constant c_a . Hence,

$$\left| \frac{f'(x) - f(x)}{x' - x} \right| \leq c_a$$

and for $x' \rightarrow x$, we get $|f'(x)| \leq c_a$.

Interpretation

- $c \approx 1$, well-conditioned.
- $c \gg 1$, ill-conditioned.
- $c = \infty$, ill-posed.

Approximations

For a continuous, differentiable f the following approximations are valid:

$$c_a \leq \max_{|\xi-x| \leq \delta} |f'(\xi)|, \quad k_r \leq \frac{|x|}{|f(x)|} \max_{|\xi-x| \leq \delta} |f'(\xi)|$$

where $\xi \in \{z \mid z = x + t(x' - x), t \in [0, 1]\}$ from the mean value theorem

$$\frac{f(x') - f(x)}{x' - x} = f'(\xi)$$

Example:

Given $f(x) = 1 + \sqrt{x-1}$ for $x > 1$.

The function is differentiable

$$f'(x) = \frac{1}{2\sqrt{x-1}}.$$

$|\xi - x| \leq \delta$ means $\xi \in [x - \delta, x + \delta]$, therefore:

$$\max_{|\xi-x| \leq \delta} \frac{1}{2\sqrt{\xi-1}} = \frac{1}{2\sqrt{x-\delta-1}}$$

For $x = 2, \delta = 10^{-8}$ we get

$$\max_{|\xi-x| \leq \delta} |f'(\xi)| = \frac{1}{2\sqrt{1-10^{-8}}} \leq \frac{1}{2\sqrt{\frac{1}{2}}} = \sqrt{\frac{1}{2}} < 1$$

Hence $c_a < 1$ and $c_r < \frac{|2|}{|f(2)|} = 1$.

Error Propagation Formulas of The Differential Error Analysis

The approximations

$$c_a \approx |f'(x)|, \quad c_r \approx \frac{|x|}{|f(x)|} |f'(x)|$$

are considered the *error propagation formulas of the error analysis*.

For the **general case** $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ we take the generalized mean value theorem

$$f(x') = f(x) + \left(\int_0^1 f'((x + t(x' - x))) dt \right) (x' - x),$$

where

$$f'(x) = \begin{bmatrix} \partial_{x_1} f_1(x) & \cdots & \partial_{x_n} f_1(x) \\ \vdots & \ddots & \vdots \\ \partial_{x_1} f_m(x) & \cdots & \partial_{x_n} f_m(x) \end{bmatrix}$$

We apply the metric

$$\begin{aligned} \|f(x') - f(x)\|_X &\leq \left\| \int_0^1 f'((x + t(x' - x))) dt \right\|_{X,Y} \|x' - x\|_X \\ &\leq \max_{\xi \in [x-\delta, x+\delta]} \|f'(\xi)\|_{X,Y} \|x' - x\|_X \end{aligned}$$

For all x' holds that $\|x' - x\|_X \leq \delta$, then

$$c_a \leq \max_{\|\xi - x\|_X \leq \delta} \|f'(\xi)\|_{X,Y}$$

and

$$c_r \leq \frac{\|x\|_X}{\|f(x)\|_Y} \max_{\|\xi - x\|_X \leq \delta} \|f'(\xi)\|_{X,Y}$$

Example:

We observe $f : \mathbb{R}^2 \rightarrow \mathbb{R} \mapsto x_1 + x_2$ with the metrics

$$\|x\|_X = \|x\|_\infty = \max(\|x_1\|, \|x_2\|), \quad \|y\|_Y = \|y\|_\infty = |y|$$

For the Jacobian we get

$$f'(x) = [\partial_{x_1} f(x), \partial_{x_2} f(x)] = [1, 1]$$

Due to our chosen vector metrics we use the infinite metric for matrices

$$\|f'(x)\|_{X,Y} = \|f'(x)\|_\infty = \|[1, 1]\| = 2$$

Therefore, $c_a \approx 2$ and $c_r \approx 2 \frac{\|x\|_\infty}{|f(x)|}$

74.2.7 Procedural Error

In contrast to the input error, this is much simpler and depends on the class of problem. For a linear system of equations using just matrix decomposition the difference $\hat{f}(\tilde{x}) - f(\tilde{x}) = 0$, while for an infinite sum it can vary depending on the number of terms use of the infinite sum.

74.2.8 Accumulation Error

This error occurs due to the floating point arithmetic not being associative for multiplication and addition. In contrast, they mostly preserve the commutativity. They are a commutative *manga* with identity for each operation $(F, +), (F, \cdot)$.

74.3 Error Analysis For Linear Systems

We have square system $Ax = b$ as a the well-posed problem

$$f(A, b) = A^{-1}b = x.$$

We analyze the problem with the condition, i.e the sensibility of x by changes in the input, and denote this as $A\tilde{x} = \tilde{b}$. Then

$$\begin{aligned}\Delta A &= \tilde{A} - A \\ \Delta b &= \tilde{b} - b \\ \Delta x &= \tilde{x} - x\end{aligned}$$

Under this conditions we can write

$$\begin{aligned}x - \tilde{x} &= A^{-1}b - A^{-1}\tilde{b} \\ \Delta x &= A^{-1}(\Delta b)\end{aligned}$$

Hence, if we take the metric and divide by $\|x\|$

$$\begin{aligned}\frac{\|\Delta x\|}{\|x\|} &= \frac{\|A^{-1}(\Delta b)\|}{\|x\|} \\ &\leq \frac{\|b\|}{\|x\|} \|A^{-1}\| \frac{\|\Delta b\|}{\|b\|} \\ &= \frac{\|Ax\|}{\|x\|} \|A^{-1}\| \frac{\|\Delta b\|}{\|b\|} \\ &\leq \frac{\|A\|\|x\|}{\|x\|} \|A^{-1}\| \frac{\|\Delta b\|}{\|b\|} \\ &= \|A\| \|A^{-1}\| \frac{\|\Delta b\|}{\|b\|}\end{aligned}$$

We used two important facts; first, $Ax = b$ and second, $\|Ax\| \leq \|A\|\|x\|$.

74.3.1 Condition of A Matrix

Given a metric $\|\cdot\|$ on $\mathbb{R}^n$ with A being regular. We define

$$\kappa(A) = \|A\| \|A^{-1}\|$$

as the *condition number* of the matrix.

It has the following properties:

- κ depends highly on the metric.
- If $\alpha\|B\|' \leq \|B\| \leq \beta\|B\|'$, then $\alpha^2\kappa'(B) \leq \kappa(B) \leq \beta^2\kappa'(B)$.
- $\kappa(A) \geq 1 \implies \|I\| = \|AA^{-1}\| \leq \|A\| \|A^{-1}\| = \kappa(A)$.
- $\kappa(A) = \kappa(A^{-1})$.
- $\kappa(AB) \leq \kappa(A)\kappa(B)$.

Extended Condition

Our condition is until now only defined for regular square matrices, which means that they are invertible. We will expand this concept in the following way.

Start with

$$\|A^{-1}\| = \sup_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \sup_{z \neq 0} \frac{\|z\|}{\|Az\|}$$

because $x = Az$ and because our matrices is bijective map we assume that we never get zero vector, but all other real numbers.

$$\begin{aligned}\sup_{z \neq 0} \frac{\|z\|}{\|x\|} &= \sup_{z \neq 0} \frac{1}{\left\|A \frac{z}{\|z\|}\right\|} \\ &= \sup_{\|z\|=1} \frac{1}{\|Az\|} \\ &= \max_{\|z\|=1} \frac{1}{\|Az\|} = \frac{1}{\min_{\|z\|=1} \|Az\|}\end{aligned}$$

By the fact that we are in a compact space we assume the equality of the supremum and maximum and now we can define the condition for any real matrix as follows:

$$\kappa(A) = \|A\| \|A^{-1}\| = \frac{\max_{\|y\|=1} \|Ay\|}{\min_{\|z\|=1} \|Az\|}$$

74.3.2 Inequality For Error Analysis in Linear Systems

Given a metric $\|\cdot\|$ on $\mathbb{R}^n$ with A being regular and $\|\Delta A\| < \frac{1}{\|A^{-1}\|}$, where $\|\cdot\|$ is an induced metric. Then, $\tilde{A}$ is regular and the followings holds:

$$\frac{\|\Delta x\|}{\|x\|} \leq \frac{\kappa(A)}{1 - \kappa(A) \frac{\|\Delta A\|}{\|A\|}} \left(\frac{\|\Delta A\|}{\|A\|} - \frac{\|\Delta b\|}{\|b\|} \right),$$

This the case for an error of type $(A + \Delta A)(x + \Delta x) = (b + \Delta b)$.

There are also the cases

- $A(x + \Delta x) = (b + \Delta b)$:

$$\frac{\|\Delta x\|}{\|x\|} \leq \kappa(A) \frac{\|\Delta b\|}{\|b\|}$$

- $(A + \Delta A)(x + \Delta x) = b$:

$$\frac{\|\Delta x\|}{\|x\|} \leq \frac{\kappa(A)}{1 - \kappa(A)} \frac{\|\Delta b\|}{\|b\|}$$

Example:

Given are

$$A = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix} \quad A^{-1} = \begin{bmatrix} -0.4 & 0.2 \\ 0.2 & 0.1 \end{bmatrix} \quad b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \|\Delta b\|_1 = 10^{-5}, \quad \|\Delta A\|_1 = 0$$

We are looking for an upper bound for $\|\Delta x\|_1$.

In this case we are handling the case $A(x + \Delta x) = (b + \Delta b)$:

$$\begin{aligned}\kappa(A) &= \|A\|_1 \cdot \|A^{-1}\|_1 = \frac{42}{10} \\ \|b\|_1 &= 8 \\ \|x\|_1 &= 2\end{aligned}$$

Resulting into

$$\begin{aligned}\frac{\|\Delta x\|}{\|x\|} &\leq \kappa(A) \frac{\|\Delta b\|}{\|b\|} \\ &\leq \frac{42}{10} \cdot \frac{10^{-5}}{8} 2 = 105 \cdot 10^{-5}\end{aligned}$$

74.3.3 Prager Oettli

For $x \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$ is

$$|x| = [|x_1|, \dots, |x_n|]^T, \quad |A| = (|a_{i,j}|)_{i,j}.$$

If $\det(A) \neq 0$ and $\tilde{x}$ is an approximation of $Ax = b$ with *residuum* $\tilde{r} = b - A\tilde{x}$ and

$$B \in \mathbb{R}^{n \times n}, c \in \mathbb{R}^n, b_{i,j} \geq 0, c_i \geq 0, \forall i, j$$

with

$$|\tilde{r}| \leq B|\tilde{x}| + c$$

Then there are $|\Delta A| \leq B$, $(|\Delta a_{i,j}| \leq b_{i,j}, \forall i, j)$ and $|\Delta b| \leq c$, such that $\tilde{x}$ becomes the exact solution of

$$(A + \Delta A)\tilde{x} = b + \Delta b$$

This is used when for example when we have system where we know A and b have a certain degree of noise B and c . This means that our system $Ax = b$ has a level of uncertainty. Accordingly, we can not expect an exact solution, but we can limit ourselves to a solution in a certain degree of uncertainty.

If we know B and c then we know that $\tilde{x}$ is the exact solution of distorted problem. If they are also component-wise small then $\tilde{x}$ is a good approximation of our original problem.

Example:

$$A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}, \quad b = \begin{bmatrix} -8 \\ 19 \end{bmatrix}$$

With the exact solution $x = [1, 10]^T$.

The coefficients for A and B have a noise of

$$|\Delta A| \leq \frac{1}{10} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = B, \quad |\Delta b| \leq \frac{1}{10} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = c$$

We assume that we have two different approximations $\tilde{x}_1 = [1, 9]^T$ and $\tilde{x}_2 = [0, 10]^T$.

We apply our formula for $\tilde{x}_1$

$$\tilde{r} = b - A\tilde{x}_1 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

Also,

$$|\tilde{r}| = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Continuing,

$$B|\tilde{x}_1| + c = \begin{bmatrix} 1.2 \\ 2 \end{bmatrix}$$

Therefore,

$$|\tilde{r}| = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \leq \begin{bmatrix} 1.2 \\ 2 \end{bmatrix} = B|\tilde{x}_1| + c$$

Hence, we can accept $\tilde{x}_1$ because of $|\Delta A| \leq B$ and $|\Delta b| \leq c$ with $(A = \Delta A)\tilde{x}_1 = b + \Delta b$.

For $\tilde{x}_2$ the reader can complete the example and note that the key inequality $|\tilde{r}| \leq B|\tilde{x}_1| + c$ does not hold.

74.3.4 QR-Decomposition For Numerical Mathematics

Givens Algorithm

This algorithm uses rotations for the elimination of the column elements.

Given a column vector $u = (u_1, \dots, u_n)^T \in \mathbb{R}^n$ where $w = \sqrt{u_1^2 + u_i^2}, i > 1$.

Set $w = 0, Q_{i,1} = I$ for $w \neq 0$

$$Q_{i,1} = \begin{bmatrix} c & & & s & & \\ & 1 & & & & \\ & & \ddots & & & \\ & -s & & 1 & & \\ & & & & \ddots & \end{bmatrix}, c = \pm \frac{u_1}{w}, s = \pm \frac{u_i}{w}$$

Then $Q_{i,1}$ is orthonormal and

$$Q_{i,1}u = \begin{bmatrix} \pm w \\ u_2 \\ \vdots \\ u_{i-1} \\ \vdots \\ 0 \\ u_{i+1} \\ \vdots \\ u_n \end{bmatrix}$$

i.e u_i will be eliminated and nothing changes besides u_1, u_i .

Proceeding, is we use the positive sign we get $s = [a_{1,1}, \dots, a_{n,1}]^T$ of A .

$$s_1^{(2)} = Q_{2,1}s_1 = \begin{bmatrix} \sqrt{a_{1,1}^2 + a_{2,1}^2} \\ 0 \\ a_{3,1} \\ \vdots \\ a_{n,1} \end{bmatrix}$$

Analogously, we can eliminate the other elements of the column

$$s_1^{(3)} = Q_{3,1}s_1^{(2)} = \begin{bmatrix} \sqrt{a_{1,1}^2 + a_{2,1}^2 + a_{3,1}^2} \\ 0 \\ 0 \\ a_{4,1} \\ \vdots \\ a_{n,1} \end{bmatrix} \dots s_1^{(n)} = Q_{n,1}s_1^{(n-1)} = \begin{bmatrix} \|s_1\|_2 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

In total we have $Q_1 = Q_{n,1} \cdot \dots \cdot Q_{2,1}$ for the first column; and $Q_1 s_1 = \|s_1\| e_1$.

We can continue with the second column and so on, giving us

$$Q_{n-1} \cdot \dots \cdot Q_1 A = R$$

By the orthonormality of the Q_i we have

$$A = Q_1^T \cdot \dots \cdot Q_{n-1}^T R = QR$$

Hence, there exists an orthonormal Q and an upper triangular R such that $A = QR$.

□◻◻

Algorithm

We have our system $Ax = b$ in $\mathbb{R}^{n \times n}$ and are going to store the Q and R in the same matrix. We start with $Q = I$ and $R = A$.

1. For each element in the current column, we compute

$$s = \frac{a_{i,j}}{\sqrt{a_{i,j}^2 + a_{k,j}^2}}, \quad c = \frac{a_{k,j}}{\sqrt{a_{i,j}^2 + a_{k,j}^2}}$$

with $i < k$ and k being ...

2. Create the matrix $Q_{i,j}$ with the following structure:

$$Q_{i,j} = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & c & s & \\ & & -s & c & \\ & & & & \ddots \end{bmatrix}$$

the position of c and s is at the i -th and k -th row and column.

3. Multiply $Q_{i,j}$ with A and store at the position of the generated zero the value s and at the position of c the value c .
4. Continue with the next element in the column until all elements are eliminated. Note that we applying the multiplication for the new A , we handle the places where we have stored the $\rho_{i-1,j-1}$ as if they were zero, because they are not used for the next steps.
5. Then, continue with the next column in the new sub-matrix and so on until we have an upper triangular matrix.
6. Now, compute $y = Q^T b = Q_{n-1} \cdot \dots \cdot Q_1 b$. One matrix a a time. We use the stored coefficient $\rho_{i,j} = s$ to then compute $s = \sqrt{1 - s^2}$ for the multiplication.
7. Finally, solve $Rx = Q_{n-1} \cdot \dots \cdot Q_1 Ax = y$ for x using back substitution.

It is important to note that this algorithm is four times more expensive than the LU -Decomposition, but it is more stable.

Example:

Find the QR -Decomposition of A and solve $Ax = b$.

$$A = \begin{bmatrix} 12 & -40 & 46 \\ -9 & 30 & 153 \\ 20 & 100 & 135 \end{bmatrix}, b = \begin{bmatrix} -16 \\ 12 \\ 140 \end{bmatrix}$$

We start by eliminating $a_{1,2}$:

$$c = \frac{12}{\sqrt{12^2 - (-9)^2}} = \frac{4}{5}, \quad s = \frac{-9}{\sqrt{12^2 - (-9)^2}} = -\frac{3}{5}$$

Therefore

$$Q_{2,1} = \begin{bmatrix} \frac{4}{5} & -\frac{3}{5} & 0 \\ \frac{3}{5} & \frac{4}{5} & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Then

$$Q_{2,1}A = \begin{bmatrix} 15 & -50 & -55 \\ 0 & 0 & 150 \\ 20 & 100 & 135 \end{bmatrix}$$

We store $\rho_{2,1} = s = -\frac{3}{5}$ at the generated 0 at $a_{2,1}$.

$$\begin{bmatrix} 15 & -50 & -55 \\ -\frac{3}{5} & 0 & 150 \\ 20 & 100 & 135 \end{bmatrix}$$

Now, we eliminate the element $a_{3,1}$:

$$c = \frac{15}{\sqrt{15^2 + 20^2}} = \frac{3}{5}, \quad s = \frac{20}{\sqrt{15^2 + 20^2}} = \frac{4}{5}$$

Now we get

$$Q_{3,1} = \begin{bmatrix} \frac{3}{5} & 0 & \frac{4}{5} \\ 0 & 1 & 0 \\ -\frac{4}{5} & 0 & \frac{3}{5} \end{bmatrix}$$

Applying the multiplication to our previously computed matrix $Q_{2,1}A$

$$Q_{3,1}Q_{2,1}A = \begin{bmatrix} 25 & 50 & 75 \\ 0 & 0 & 150 \\ 0 & 100 & 125 \end{bmatrix}$$

We store $\rho_{3,1} = s = \frac{4}{5}$ at $a_{3,1}$.

$$\begin{bmatrix} 25 & 50 & 75 \\ -\frac{3}{5} & 0 & 150 \\ \frac{4}{5} & 100 & 125 \end{bmatrix}$$

Continuing with $a_{3,2}$:

$$c = \frac{0}{\sqrt{0^2 + 100^2}} = 0, \quad s = \frac{100}{\sqrt{0^2 + 100^2}} = 1$$

$$Q_{3,2} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

Multiply

$$Q_{3,2}Q_{3,1}Q_{2,1}A = \begin{bmatrix} 25 & 50 & 75 \\ 0 & 100 & 125 \\ 0 & 0 & -150 \end{bmatrix}$$

Now, we store $\rho_{3,1} = s = 1$ at $a_{3,2}$

$$\begin{bmatrix} 25 & 50 & 75 \\ -\frac{3}{5} & 100 & 125 \\ \frac{4}{5} & 1 & -150 \end{bmatrix}$$

Now we solve $Qy = b \iff Q^T b = y$ for $Rx = y$ using

$$Q = (Q_{3,2}Q_{3,1}Q_{2,1})^T = \frac{1}{25} \begin{bmatrix} 12 & -16 & -15 \\ -9 & 12 & -20 \\ 20 & 15 & 0 \end{bmatrix}$$

and

$$R = Q_{3,2}Q_{3,1}Q_{2,1}A = 25 \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & -6 \end{bmatrix}$$

For each of the matrices in $Q^T b$ we reconstruct the the $Q_{i,j}$ matrix and perform the multiplication

Using $\rho_{2,1} = -\frac{3}{5}, c = \sqrt{1 - s^2} = \frac{4}{5}$:

$$Q_{2,1}b = \begin{bmatrix} \frac{4}{5} & -\frac{3}{5} & 0 \\ \frac{3}{5} & \frac{4}{5} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -16 \\ 12 \\ 140 \end{bmatrix} = \begin{bmatrix} -20 \\ 0 \\ 140 \end{bmatrix}$$

Then repeat for the others, and apply the multiply the new b to the reconstructed matrix and we get

$$y = \begin{bmatrix} 100 \\ 100 \\ 0 \end{bmatrix}$$

Finally, for $Rx = y$ we have

$$25 \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & -6 \end{bmatrix} x = y$$

which gives us $x = [2, 1, 0]^T$.

74.3.5 Householder Algorithm

The *Householder algorithm* similarly to Givens algorithm will give us a decomposition of a matrix A such that $A = QR$ with Q being orthonormal and R being upper triangular. The difference relies on the method of the transformations which instead of being rotations, they are reflections.

Householder Transformation

Given a vector $v \in \mathbb{R}^n$ we denote

$$H_v = I - 2 \frac{vv^T}{v^T v}$$

as the *Householder transformation* for v . Equivalently

$$H_v = I - 2 \frac{vv^T}{\|v\|_2^2}.$$

If v is normalized, then $H_v = I - 2vv^T$, which is a reflection across the hyperplane orthogonal to v .

Example:

Given $v = [1, 2]^T$ we have

$$H_v = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - 2 \frac{\begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \end{bmatrix}}{\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - 2 \frac{1}{5} \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} -3 & -4 \\ -4 & 3 \end{bmatrix}$$

74.3.6 Effect of a Householder Transformation

Given $x \in \mathbb{R}^n$ when multiplying H_k , which is the k -th Householder transformation, with x we get

$$H_k x = H_k \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k-1} \\ x_k \\ x_{k+1} \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{k-1} \\ ? \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

The vector to be used for the transformation defined as

$$v = x \pm \|x\|_2 e_k$$

The sign is not really important, but it is usually chosen to avoid cancellation errors.

The k -th Householder transformation is then defined as $\tilde{x}$ where all elements before the k -th element are zero, i.e. if we have to eliminate all elements below the first element, then we have $k = 1$ and $\tilde{x} = x$, but if we have to eliminate all elements below the second element, then we have $k = 2$ and $\tilde{x} = [0, x_2, x_3, \dots, x_n]^T$ and so on. Note that the resulting transformation will be still be used with x .

Algorithm

Given $A \in \mathbb{R}^{n \times n}$ we start with $Q = I$ and $R = A$. Similar to the LU -Decomposition we will start with the first column and then focus on the sub-matrix and so on.

1. Eliminate the elements below the pivot element $a_{i,i}$ by applying the Householder transformation H_i to the i -th column of the current matrix.
2. Multiply H_i with the current matrix and store the resulting matrix which will be used for the next step.
3. Continue with the next column and so on until we have an upper triangular matrix. The result should be $H_k \cdot H_{k-1} \cdot \dots \cdot H_1 \cdot A = R$ and $Q = H_1 \cdot H_2 \cdot \dots \cdot H_k$.

Note that the Householder matrices are self-inverse, i.e. $H_i^{-1} = H_i$ and therefore $A = QR$.

Example

Given is

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

We start with the first column and compute the Householder transformation for the first column. We have

$$v = \begin{bmatrix} 1 \\ 3 \end{bmatrix} + \sqrt{10} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + \sqrt{10} \\ 3 \end{bmatrix}$$

Then, we compute the Householder transformation

$$H_1 = I - 2 \frac{vv^T}{v^T v} = \begin{bmatrix} -\frac{1}{3} & -\frac{3}{5} \\ -\frac{3}{5} & \frac{1}{5} \end{bmatrix}$$

And we apply it to A

$$H_1 A = \begin{bmatrix} -\frac{1}{3} & -\frac{3}{5} \\ -\frac{3}{5} & \frac{1}{5} \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} -\sqrt{10} & -\frac{4}{5}\sqrt{10} \\ 0 & \frac{2}{5}\sqrt{10} \end{bmatrix} = R$$

Now, we have an upper triangular matrix and we can compute Q as H_1 and R as $H_1 A$, and we have $A = QR$.

74.4 Iterative Methods

An *iterative method*

$$x^{(k+1)} = \phi_k(x^{(k)}, x^{(k-1)}, \dots, x^{(0)})$$

is convergent if $x^{(k)} \rightarrow x$ for all $x^{(0)} \in \mathbb{R}^n$. It also has the *fixed point property* $x = \phi(x) \iff Ax = b$.

The *error* $e^{(k)}$ and the *residuum* $r^{(k)}$ are defined as

$$e^{(k)} = x - x^{(k)}, \quad r^{(k)} = b - Ax^{(k)}$$

If for the computation of $x^{(k+1)}$ only the direct predecessor $x^{(k)}$ uses, then

$$x^{(k+1)} = \phi(x^{(k)})$$

the process is considered with *one step* $\phi_k : \mathbb{R}^n \rightarrow \mathbb{R}^n$, else is considered with *multi step* $\phi_k : (\mathbb{R}^n)^{k+1} \rightarrow \mathbb{R}^n$.

If $\phi_k = \phi$ for all k then the process is called *stationary*, else it is called *non-stationary*.

74.4.1 Single Step Stationary Iteration Method

A *single step stationary iteration method* is given by

$$x^{(k+1)} = \phi(x^{(k)}) = Bx + c,$$

where B is the *iteration matrix* and c is the *iteration vector*.

74.4.2 Error of Single Step Stationary Iteration Method

If the single step stationary iteration method has the fixed point property, then for $e^{(k)} = x - x^{(k)}$, $r^{(k)} = b - Ax^{(k)}$ we have

$$Ae^{(k)} = r^{(k)}$$

Proof:

$$Ae^{(k)} = A(x - x^{(k)}) = Ax - Ax^{(k)} = b - Ax^{(k)} = r^{(k)}$$

□◻◻

74.4.3 Convergence

Analyzing our regular linear system $Ax = b$ with the single step iteration

$$x^{(k+1)} = \phi(x^{(k)}) = Bx^{(k)} + c$$

with the fixed point property $x = \phi(x) = Bx + c \iff Ax = b$ we get

For the error $e^{(k)} = x^{(k)} - x$ we have

$$e^{(k+1)} = x^{(k+1)} - x = Bx^{(k)} + c - (Bx + c) = Be^{(k)}$$

Therefore,

$$e^{(k)} = B^k e^{(0)}$$

Then

$$e^{(k)} \rightarrow 0 \quad \text{as} \quad k \rightarrow \infty$$

for an arbitrary $e^{(0)}$ if and only if $B^k \rightarrow 0$ as $k \rightarrow \infty$.

74.4.4 The Spectral Radius

The *spectral radius* of a matrix B is defined as

$$\rho(B) = \max\{|\lambda| \mid \lambda \in \sigma(B)\},$$

where $\sigma(B)$ is the *spectrum* of B , i.e. the set of all eigenvalues of B .

Properties of the Spectral Radius

- $\rho(B) \leq \|B\|_I$ for all induced metrics.
- $\forall \varepsilon > 0$ there exists an induced metric $\|\cdot\|_I$ from a vector metric $\|\cdot\|_V$ such that:

$$\|B\|_I = \sup_{w \neq 0} \frac{\|Bw\|_w}{\|w\|_V}$$

with $\|B\|_I \leq \rho(B) + \varepsilon$.

74.4.5 Gelfand

For any matrix metric $\|\cdot\|$ the following holds:

$$\rho(B) = \lim_{k \rightarrow \infty} \sqrt[k]{\|B^k\|}.$$

Also for an induced matrix metric $\|\cdot\|_I$ $\|B\|_I^{\frac{1}{k}}$ converges towards $\rho(B)$.

74.4.6 Splitting Method For Linear Systems

Given a regular matrix $A \in \mathbb{R}^{n \times n}$ in a system $Ax = b$, we can split A into

$$A = M - N$$

where M is regular and N is arbitrary. Then, we can write

$$Mx = Nx + b$$

and we can define the iteration

$$x^{(k+1)} = M^{-1}Nx^{(k)} + M^{-1}b,$$

and we set $x^{(0)} = \vec{0} \in \mathbb{R}^n$ as the initial guess.

Example:

Given the system

$$\begin{bmatrix} 4 & -1 & 0 & 0 \\ -1 & 4 & -1 & 0 \\ 0 & -1 & 4 & -1 \\ 0 & 0 & -1 & 3 \end{bmatrix} x = \begin{bmatrix} 15 \\ 10 \\ 10 \\ 10 \end{bmatrix}$$

Then we can split A into

$$M = \begin{bmatrix} 4 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}, \quad N = \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & -1 & 0 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix}$$

Which results in the iteration

$$x^{(k+1)} = \begin{bmatrix} \frac{1}{4} & 0 & 0 & 0 \\ 0 & \frac{1}{4} & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & -1 & 0 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} \frac{1}{4} & 0 & 0 & 0 \\ 0 & \frac{1}{4} & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} 15 \\ 10 \\ 10 \\ 10 \end{bmatrix} = \begin{bmatrix} 3.75 \\ 2.5 \\ 2.5 \\ 3.333\dots \end{bmatrix}$$

We can then continue the process using our resulting vector as the new $x^{(k)}$ and we will get closer to the solution of the system.

Triple Splitting

We can also define the splitting

$$A = D - L - U$$

where D is the diagonal of A , L is the lower triangular part of A and U is the upper triangular part of A . We then derive from that

$$\begin{aligned} Ax = b &\iff (D - L - U)x = b \\ &\iff Dx = (L + U)x + b \end{aligned}$$

The iteration is then given by

$$x^{(k+1)} = D^{-1}(L + U)x^{(k)} + D^{-1}b$$

74.4.7 The Residuum As Stopping Criterion

The *residuum* is defined as

$$r^{(k)} = b - Ax^{(k)}$$

When $\|r^{(k)}\| \leq \varepsilon \|b\|$ for a given $\varepsilon > 0$ then we can stop the iteration because we are close enough to the solution.

74.4.8 Convergence For Semi Positive Definite Matrices

Given semi positive definite matrices A, M ; if $2M - A$ is also semi positive definite, then

$$\phi(x) = Bx + d, \quad B = I - M^{-1}A, \quad d = M^{-1}b$$

converges and also $\|B\|_A = \|B\|_M = \rho(B) < 1$.

Also, the splitting $M + M^T - A$ is also spd. and $\rho(B) \leq \|B\|_A < 1$.

74.4.9 Strict Diagonal-dominant

A is strictly diagonal-dominant if

$$|a_{i,i}| > \sum_{j \neq i} |a_{i,j}|, \quad \forall i = 1, \dots, n$$

74.4.10 The Jacobi Method

The Jacobi method is a single step stationary iteration method with the splitting $A = D - (L + U)$ and the iteration:

$$x^{(k+1)} = D^{-1}(L + U)x^{(k)} + D^{-1}b$$

Example:

Given the system

$$\begin{bmatrix} 4 & -1 & 0 & 0 \\ -1 & 4 & -1 & 0 \\ 0 & -1 & 4 & -1 \\ 0 & 0 & -1 & 3 \end{bmatrix} b = \begin{bmatrix} 15 \\ 10 \\ 10 \\ 10 \end{bmatrix}$$

We can apply the Jacobi method with the splitting $A = D - (L + U)$ where

$$D = \begin{bmatrix} 4 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}, \quad L + U = \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & -1 & 0 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix}$$

Then, we can compute the iteration

$$x^{(k+1)} = \begin{bmatrix} \frac{1}{4} & 0 & 0 & 0 \\ 0 & \frac{1}{4} & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & -1 & 0 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} \frac{1}{4} & 0 & 0 & 0 \\ 0 & \frac{1}{4} & 0 & 0 \\ 0 & 0 & \frac{1}{4} & 0 \\ 0 & 0 & 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} 15 \\ 10 \\ 10 \\ 10 \end{bmatrix}$$

Giving us

$$x^{(1)} = \begin{bmatrix} 3.75 \\ 2.5 \\ 2.5 \\ 3.333 \dots \end{bmatrix}$$

74.4.11 Gauss-Seidel Method

The Gauss-Seidel method is a single step stationary iteration method with the splitting $A = (D - L) - U$ and the iteration

$$x^{(k+1)} = (D - L)^{-1} U x^{(k)} + (D - L)^{-1} b$$

74.4.12 After Iteration

This process refers to using an iterative method after using a mathematical algorithm for solving a certain problem to mitigate rounding errors. In our case we will present such a process with the LU -Decomposition.

While performing the LU -Decomposition we will get $\tilde{L}\tilde{U} = M$ which are the L, U matrices, but with rounding errors. We use the stationary linear iteration to improve the solution $\tilde{x}$ for which $A\tilde{x} = b$.

1. Perform the LU -Decomposition to get the triangular matrices and the approximate solution with errors.
2. Set $x_0 = \tilde{x}$
3. Repeat the following steps as long $\frac{\|p_k\|}{\|x_{k+1}\|} < \varepsilon$:
 - $r_k = b - Ax_k$.
 - $p_k = M^{-1}r_k$ in other words $\tilde{L}\tilde{U}p_k = r_k$. Also

```
p_k = np.linalg.solve(A, r_k)
```
 - $x_{k+1} = x_k + p_k$.

74.4.13 Gradient Descent

Given our stationary iteration

$$x^{(k+1)} = x^{(k)} + M^{-1}r^{(k)}$$

We add the parameter $\alpha^{(k)}$

$$x^{(k+1)} = x^{(k)} + \alpha^{(k)}M^{-1}r^{(k)}$$

With the goal in each step to find an $\alpha^{(k)}$ which gives us the "best" $x^{(k+1)}$. This means

$$\begin{aligned} e^{(k+1)} &= x - x^{(k+1)} \\ &= x - x^{(k)} + \alpha^{(k)}M^{-1}r^{(k)} \\ &= e^{(k)} - \alpha^{(k)}p^{(k)} \end{aligned}$$

Geometrically, $e^{(k+1)}$ is on a straight line determined by $e^{(k)}$ in the direction $p^{(k)}$. This means that $\|e^{(k+1)}\|$ will be minimal when $p^{(k)}$ is orthogonal to $e^{(k+1)}$

$$\begin{aligned} \langle e^{(k+1)}, p^{(k)} \rangle &= \langle p^{(k)}, e^{(k)} - \alpha^{(k)}p^{(k)} \rangle \\ &= \langle p^{(k)}, e^{(k)} \rangle - \alpha^{(k)} \langle p^{(k)}, p^{(k)} \rangle \\ \alpha^{(k)} &= \frac{\langle p^{(k)}, e^{(k)} \rangle}{\langle p^{(k)}, p^{(k)} \rangle} \end{aligned}$$

Hence, we have constructed the following iterative algorithm

$$\begin{aligned} r^{(k)} &= b - Ax^{(k)} \\ p^{(k)} &= M^{-1}r^{(k)} \\ \alpha^{(k)} &= \frac{\langle p^{(k)}, e^{(k)} \rangle}{\langle p^{(k)}, p^{(k)} \rangle} \\ x^{(k+1)} &= x^{(k)} + \alpha^{(k)}p^{(k)} \end{aligned}$$

The problem is that technically we do not know the exact solution x , but this can be solved using a suited inner product with metric

$$\|x\| = \|x\|_A = \sqrt{x^T A x}$$

Giving us

$$\alpha^{(k)} = \frac{\langle p^{(k)}, e^{(k)} \rangle_A}{\langle p^{(k)}, p^{(k)} \rangle_A} = \frac{\langle p^{(k)}, Ae^{(k)} \rangle}{\langle p^{(k)}, Ap^{(k)} \rangle} = \frac{\langle p^{(k)}, r^{(k)} \rangle}{\langle p^{(k)}, Ap^{(k)} \rangle}$$

74.5 Iterative Methods For Eigenvalue Problems

74.5.1 The Vector Iteration

Given a normal matrix $A \in \mathbb{C}^{n \times n}$, which is unitarily diagonalizable, i.e. there exists an orthonormal basis of eigenvectors, $u_1, \dots, u_n$.

Each vector $x \in \mathbb{C}^n$ can be written as $x = \sum_i x_i u_i$ and

$$Ax = \sum_i x_i \lambda_i u_i, \quad A^k x = \sum_i x_i \lambda_i^k u_i$$

If $|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|$ then the dominant eigenvalue is λ_1 . This means that by applying A to x multiple times we get closer to the dominant eigenvector u_1 and the dominant eigenvalue λ_1 .

Algorithm

1. Choose a $x^{(0)}$ such that $\|x^{(0)}\|_2 = 1$ and $\langle x^{(0)}, u_1 \rangle \neq 0$.
2. Repeat the following steps until convergence:

$$\begin{aligned} y^{(k+1)} &= Ax^{(k)} \\ x^{(k+1)} &= \frac{y^{(k+1)}}{\|y^{(k+1)}\|_2} \\ \mu^{(k+1)} &= \langle x^{(k+1)}, Ax^{(k+1)} \rangle = \frac{\langle x^{(k+1)}, y^{(k+1)} \rangle}{\|x^{(k)}\|_2^2} \end{aligned}$$

74.5.2 The Jacobi Method

The *Jacobi method* is an iterative method for computing the eigenvalues of a symmetric matrix A by applying a sequence of orthogonal transformations to A to make it diagonal.

The eigenvalues of A are then given by the diagonal elements of the resulting matrix.

The derivation will come from applying orthonormal matrices repeatedly to our original matrix.

$$A^{(0)} = A, \quad A^{(k)} = Q_k^T A^{(k-1)} Q_k$$

Using our Givens-Rotations

$$Q_k = \begin{bmatrix} 1 & & & & & & \\ & \ddots & & & & & \\ & & c & & s & & \\ & & & 1 & & & \\ & & & & \ddots & & \\ & & -s & & & c & \\ & & & & & 1 & \\ & & & & & & \ddots & \\ & & & & & & & 1 \end{bmatrix}$$

where

$$c = \cos(\alpha), \quad s = \sin(\alpha), \quad \alpha = \begin{cases} \frac{\pi}{4} \\ \frac{1}{2} \arctan\left(\frac{2a_{i_0, j_0}}{a_{j_0, j_0} - a_{i_0, i_0}}\right) \end{cases} \quad a_{i_0, i_0} = a_{j_0, j_0}$$

Note the following coordinates when reading the matrix from top to bottom, left to right:

$$\begin{array}{rcl} c & \text{at} & i_0, i_0 \\ s & \text{at} & i_0, j_0 \\ c & \text{at} & j_0, j_0 \\ -s & \text{at} & j_0, i_0 \end{array}$$

And we use the iteration

$$Q^{(0)} = I, \quad Q^{(k)} = Q^{(k-1)} Q_k$$

Stop Criterion

We define the square sum of the diagonal elements of the matrices $A^{(k)}$ as

$$N(A^{(k)}) = \sum_{i \neq j} \left(a_{i,j}^{(k)} \right)^2.$$

It holds that

$$N(A^{(k+1)}) \leq \alpha N(A^{(k)}), \quad \alpha \in [0, 1)$$

converges *strictly monotonously decreasing* towards zero. Thus, this is our convergence criteria given an upper bound.

74.5.3 The QR Method

We take a matrix $A \in \mathbb{R}^{n \times n}$ with n start vectors whose dimension is also n .

Choose $X^{(0)} = [x_1^{(0)}, \dots, x_n^{(0)}]$ where the vectors form an orthonormal basis

For $k = 0, 1, \dots$ repeat the following steps:

$$\begin{aligned} Y^{(k+1)} &= AX^{(k)} \\ Q^{(\tilde{k}+1)} R^{(k+1)} &= Y^{(k+1)} \\ X^{(k+1)} &= Q^{(\tilde{k}+1)} \\ T^{(k+1)} &= X^{(k)T} Y^{(k+1)} = X^{(k)T} AX^{(k)} \end{aligned}$$

Then for

$$\begin{aligned} T^{(k)} &= X^{(k-1)T} AX^{(k-1)} \\ &= X^{(k-1)T} Y^{(k)} \\ &= X^{(k-1)T} Q^{(\tilde{k})} R^{(k)} \\ &= (X^{(k-1)T} X^{(k)}) R^{(k)} \\ &= Q^{(k)} R^{(k)} \end{aligned}$$

We can also use $Q^{(k)} = X^{(k-1)T} X^{(k)}$. And because the vector of $X^{(k)}$ are orthonormal we also have that

$$\begin{aligned}
T^{(k+1)} &= X^{(k)T} A X^{(k)} \\
&= X^{(k)T} A X^{(k-1)} X^{(k-1)T} X^{(k)} \\
&= X^{(k)T} Y^{(k)} X^{(k-1)T} X^{(k)} \\
&= X^{(k)T} Q^{(k)} R^{(k)} X^{(k-1)T} X^{(k)} \\
&= X^{(k)T} X^{(k)} R^{(k)} X^{(k-1)T} X^{(k)} \\
&= R^{(k)} (X^{(k-1)T} X^{(k)}) \\
&= R^{(k)} Q^{(k)}
\end{aligned}$$

This means that our iteration can be simplified to

$$A^{(k)} = Q^{(k)} R^{(k)}, \quad A^{(k+1)} = R^{(k)} Q^{(k)}$$

Because A is in the real Schur-form $A = QTQ^T$, under certain circumstances we get:

$$A^{(k)} \xrightarrow{k \rightarrow \infty} T, \quad \hat{Q}^{(k)} \xrightarrow{k \rightarrow \infty} T$$

with $\hat{Q}^{(k)} = Q^{(0)} \dots Q^{(k-1)}$.

74.6 Zeros Problems

When dealing with different kinds of situations and models, most problems can be interpreted as looking for the zeros of a non-linear polynomial. For this there are plenty of theory and methods.

74.6.1 The Bisection Algorithm

Given a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$, with $a_0, b_0 \in \mathbb{R}$ such that

$$f(a_0) < 0, f(b_0) > 0 \quad \text{or} \quad f(a_0) > 0, f(b_0) < 0$$

this is $f(a_0)f(b_0) < 0$, then there must be a zero in between (Middle Value Theorem).

We define our iteration as

$$x_k = \frac{a_k + b_k}{2}$$

and apply the following rules:

If $f(x_k)f(a_k) < 0$ then

$$a_{k+1} = a_k, \quad b_{k+1} = x_k$$

Else

$$a_{k+1} = x_k, \quad b_{k+1} = b_k$$

then we repeat the process. The termination criteria is the number of iterations set for the computation.

74.6.2 The Regula-Falsi Algorithm

This is a variation of the *bisection algorithm* which improves the convergence velocity of the algorithm.

We start as before:

Given a continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$, with $a_0, b_0 \in \mathbb{R}$ such that

$$f(a_0) < 0, f(b_0) > 0 \quad \text{or} \quad f(a_0) > 0, f(b_0) < 0$$

this is $f(a_0)f(b_0) < 0$ implies there must be a zero in between

We define our iteration as

$$x_k = \frac{a_k + b_k}{2}$$

Now we use the information of $(a_k, f(a_k))$ and $(b_k, f(b_k))$ to define a line

$$s(x) = f(a_k) + \frac{f(b_k) - f(a_k)}{b_k - a_k}(x - a_k)$$

We set it to $s(x) = 0 = \dots$ which gives us

$$s(x_k) = 0 \iff x_k = -\frac{b_k - a_k}{f(b_k) - f(a_k)}f(a_k) = \frac{a_k f(b_k) - b_k f(a_k)}{b_k - a_k}$$

We used the point at which our secant crosses the x-axis as our new estimation giving us the new iteration rules:

$$x_k = \frac{a_k f(b_k) - b_k f(a_k)}{b_k - a_k}$$

If $f(x_k)f(a_k) < 0$ then

$$a_{k+1} = a_k, \quad b_{k+1} = x_k$$

Else

$$a_{k+1} = x_k, \quad b_{k+1} = b_k$$

74.6.3 The Secant Algorithm

This is again a slight modification of the previous algorithm given x_{-1}, x_0 .

$$x_{k+1} = x_k - \frac{f(x_k)}{\frac{f(x_k) - f(x_{k-1})}{x_k - x_{k-1}}}$$

This time we want to find a root and for that we define different secants which will approximate us to that specific point.

74.6.4 The Newton Algorithm

Takin the Taylor Series of our function f and only considering the first linear term we get

$$f(y) = f(x_k) + (y - x_k)f'(x_k), \quad y = x_k - \frac{f(x_k)}{f'(x_k)}$$

The *Newton algorithm* defines the following iteration:

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}$$

If we would stop at the quadratic term we would get:

$$x_{k+1} = x_k - \frac{f'(x_k) \pm \sqrt{(f'(x_k))^2 - 2f(x_k)f''(x_k)}}{f''(x_k)}.$$

The geometric interpretation is that we are approximating the zero by the tangent at the point $(x_k, f(x_k))$.

74.6.5 Local Convergence

Given $x_k \in X \subseteq \mathbb{R}^n$, we define *local convergence* for the iteration $x_{k+1} = \Phi(x_k)$ as

$$\lim_{k \rightarrow \infty} x_k = x \quad \forall x_0 \in X.$$

74.6.6 Contraction

Φ is considered a contraction according to $\|\cdot\|$ on $X \subseteq \mathbb{R}^n$ if

$$\Phi : X \rightarrow X$$

and there is an $\alpha < 1$ independent from x, y such that

$$\|\Phi(x) - \Phi(y)\| \leq \alpha \|x - y\| \quad \forall x, y \in X$$

74.6.7 The Fix Point Theorem of Banach

Given a closed set $X \subseteq \mathbb{R}^n$ with the contraction $\Phi : X \rightarrow X$ and

$$\|\Phi(x) - \Phi(y)\| \leq \alpha \|x - y\| \quad \forall x, y \in X$$

such that $\alpha < 1$ is independent. Then Φ has exactly one fix-point $x = \Phi(x)$ in X . Our *Picard iteration* converges for all $x_0 \in X$ towards x . Also the following approximations hold:

$$\|x - x_k\| \leq \frac{\alpha^k}{1 - \alpha} \|x_1 - x_0\| \quad \text{a-priori}$$

$$\|x - x_k\| \leq \frac{\alpha}{1 - \alpha} \|x_k - x_{k-1}\| \quad \text{a-posteriori}$$

74.6.8 Linear Convergence

For the iteration $\Phi(x_k) = x_{k+1}$, we define *linear convergence* as

$$\|e_{k+1}\| \leq \|e_k\| \quad \forall k, \quad 0 \leq c < 1.$$

The iteration is linearly convergent if such constant c exists.

74.6.9 Convergence With Order m

Convergence With Order m is given for an iteration $\Phi(x_k) = x_{k+1}$ if, $\Phi(x)$ is linearly convergent and $0 \leq c$ is independent from k and

$$\lim_{k \rightarrow \infty} \|e_k\| = 0 \quad \text{and} \quad \|e_{k+1}\| \leq c \|e_k\|^m \quad \forall k$$

74.6.10 Convergene Of The Differentiable Iteration

If given an $m \geq 2$, a fix-point x of $\Phi : \mathbb{R} \rightarrow \mathbb{R}$ such that Φ is m -times differentiable on x with the property

$$\Phi'(x) = \Phi''(x)' = \dots = \Phi^{(m-1)}(x) = 0,$$

then there exists a closed interval $X = [x - \delta, x + \delta]$, $\delta > 0$ such that the iteration

$$x_{k+1} = \Phi(x_k)$$

for all $x_0 \in X$ with at least order m converges.

Chapter 75

Computability

75.0.1 Church Turing Thesis

The intuitive computable functions are exactly all functions which can be computed by a *Turing Machine*. In other words every computational model can be expressed as such.

75.0.2 Textual Substitution

For a given condition of the form $\{E[x := R]\}$ we perform the substitution in the following way

$$(2x + y)[x := 4y] = 8y + y = 9y$$

If there are multiple replacements of the form $[\dots], \dots, [\dots]$ we perform those from left to right.

75.0.3 Leibniz Inference Rule

To clarify the notation we will be using from now looks like fraction but it means that assuming the statement above is true then the following (bellow) is also true.

$$\frac{X = Y}{E[z := X] = E[z := Y]}$$

75.0.4 Correctness of Algorithms

The *correctness* of algorithm is determined by the sum of the following properties the *partial correctness*, this means the algorithm at each step does the right thing, and *termination*, which like the name means that the algorithm terminates.

Partial Correctness

If before running P the condition φ is true, then after the termination P should ψ be true; these are the pre- and post-conditions.

Example:

$$\langle x = 0 \rangle x := x + 1 \langle x > 0 \rangle$$

Hoare Calculus

To read the following expression which look like fraction but are not, use the following:

- *Above*: if this is true
- *Below*: then this is also true

Assignment Rule

If $\mathbf{x}$ is a variable, $\mathbf{t}$ is an expression without side effects $\varphi\mathbf{x}/\mathbf{t}$ is ψ with all $\mathbf{x}$ substituted for $\mathbf{t}$.

$$\langle \varphi[\mathbf{x}/\mathbf{t}] \rangle \quad \mathbf{x} = \mathbf{t}; \quad \langle \psi \rangle$$

Note that this rule only has the part below, this means that this rule has no precondition.

Example I:

$$\langle \mathbf{y} = 5 \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} = \mathbf{y} \rangle$$

Example II:

$$\langle 5 > 4 \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} > 4 \rangle$$

Consequence Rule (Strong pre-condition)

Here α is a stronger condition than ϕ . This way our condition ϕ has an extra layer of protection.

$$\frac{\langle \varphi \rangle \quad P \quad \langle \psi \rangle \quad \alpha \implies \varphi}{\langle \alpha \rangle \quad P \quad \langle \psi \rangle}$$

Example:

$$\frac{\langle 5 = 5 \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} = 5 \rangle \quad \text{true} \implies 5 = 5}{\langle \text{true} \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} = 5 \rangle}$$

In code

```
<true>
<5=5>
x=5;
<x=5>
```

Consequence Rule (Weak post-condition)

This let us make the post-condition weaker.

$$\frac{\langle \varphi \rangle \quad P \quad \langle \psi \rangle \quad \psi \implies \beta}{\langle \varphi \rangle \quad P \quad \langle \beta \rangle}$$

Example:

$$\frac{\langle 5 = 5 \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} = 5 \rangle \quad \mathbf{x} = 5 \implies \mathbf{x} \geq 5}{\langle \text{true} \rangle \quad \mathbf{x} = 5; \quad \langle \mathbf{x} \geq 5 \rangle}$$

Note that two guards can only be consecutive if from the one above the one below is implied

Sequence Rule

This rule allows us to compress our conditions.

$$\frac{\langle \psi \rangle \quad P \quad \langle \psi \rangle \quad \langle \psi \rangle \quad Q \quad \langle \beta \rangle}{\langle \psi \rangle \quad PQ \quad \langle \beta \rangle}$$

Example:

$$\langle \text{true} \rangle \quad \mathbf{x} = 5; \quad \text{res} = \mathbf{x} * \mathbf{x} + 6; \quad \langle \text{res} = 31 \rangle$$

Condition Rule I

This rule enforces that no matter in which branch of the program is taken the post-condition is true.

$$\frac{\langle \phi \wedge B \rangle P \langle \psi \rangle \quad \phi \wedge \neg B \implies \psi}{\langle \phi \rangle \text{ if } (B) \{P\} \langle \psi \rangle}$$

Example:

```
<true>
<y=y>
res=y
<res=y>
```

```
if_(x_<_y)_ {
  res_ = y_ \land x_ <_ y_
  x_ = max(x, y)
  res_ = x
  res_ = max(x, y)
}
<res_ = max(x, y)>
```

Condition Rule II

Similar to the previous case, we want to make sure that independent of the branch we have a correct post-condition.

$$\frac{\langle \phi \wedge B \rangle P \langle \psi \rangle \quad \langle \phi \wedge \neg B \rangle Q \langle \psi \rangle}{\langle \phi \rangle \text{ if } (B) \text{ else } \{Q\} \langle \psi \rangle}$$

Example:

```
<true>
<x=x>
res=x
<res=x>
```

```
if_(x_<_0)_ {
  true_ \land x_ <_ 0_
  -x_ = |x|
  res_ = -x
  res_ = |x|
}_else_{
  true_ \land -x_ <_ 0_
  x_ = |x|
  res_ = x
  res_ = |x|
}
<res_ = |x|>
```

Loop Rule

In this rule we look for some *invariant* ϕ which before and after the loop is true.

$$\frac{\langle \phi \wedge B \rangle P \langle \phi \rangle}{\langle \phi \rangle \text{ while } (B) \{Q\} \langle \phi \wedge \neg B \rangle}$$

Note that from the guard before the loop $\langle \alpha \rangle$, we have $\alpha \implies \phi$ Finding the invariant can be tricky, but it commonly involves finding an integer expression, also some formula which holds for the entire loop.

Example:

In this case the invariant is $!res = n!$

```

UUUU<true>
UUUUiU=Un;
UUUUresU=U1;
UUUU<iU=UnU\landresU=U1>
UUUU<i!UresU=Un!>
UUUUwhile(iU>U1){
UUUU<i!UresU=Un!\landiU>U1>
UUUU<(iU-U1)!UresUiU=Un!>
UUUUUUUUresU=UresU*Ui;
UUUU<(iU-U1)!UresU=Un!>
UUUUUUUUiU=UiU-U1;
UUUU<i!UresU=Un!>
UUUU}
UUUU<\phiU\landU\negiU>U1>
UUUU<resU=Un!>

```

75.0.5 Termination

For each loop **while** (B){ P } we have to find some **int** expression V called the *variant* such that

$$B \implies V \geq 0 \text{ and } \langle V = m \wedge B \rangle P \langle V < m \rangle$$

In simpler words, we look for an integer expression which continuously decreases.

Example:

```

UUUU<iU=UmU\landiU>U1>
UUUUwhile(iU>Uq){
UUUU<iU-U1<UmU>
UUUUUUUUresU=UresU*Ui;
UUUU<iU-1<UmU>
UUUUUUUUiU=UiU-U1;;
UUUU<iU<UmU>
UUUU}

```

75.1 Random Access Machine

The *Random Access Machine*, *RAM* for short, is a mathematical model used in the analysis of algorithm. It helps us tackle the following questions when dealing with algorithms.

- *Correctness*: Does the algorithm compute what we want.
- *Termination*: Does the algorithm ever ends.
- *Speed*: How long does the algorithm need to execute.
- *Memory Consumption*: How much memory does the algorithm need.

The RAM consists of:

- Program memory (read-only access)
- Instruction counter
- Main memory (read and write)
- Memory cell 0 as accumulator
- Memory cells hold integers
 - No size limitation!
- Input and output tape
 - Arbitrarily many integers
 - No random access!

75.2 Complexity Theory

Asymptotic notations describe how algorithms perform as input size (n) grows.

They help analyze:

- *Time Complexity*: How runtime grows.
- *Space Complexity*: How memory usage grows.

75.2.1 Definitions

To clarify, all of these are sets of functions.

Big O (O)

- Upper Bound
- Describes the *worst-case* growth rate.
- $f(n) \in O(g(n))$ means $\exists c, n_0$ such that $\forall n > n_0, f(n) \leq c \cdot g(n)$. In other words, at some point our function f is always going to be less than g .

Omega (Ω)

- Lower Bound
- Describes the *best-case* growth rate.
- $f(n) \in \Omega(g(n))$ means $\exists c, n_0$ such that $\forall n > n_0, f(n) \geq c \cdot g(n)$. In other words, at some point our function f is always going to be greater than g .

Theta (Θ)

- Tight Bound
- Describes the *average-case* or exact growth rate.
- $f(n) \in \Theta(g(n))$ means $f(n) \in O(g(n))$ and $f(n) \in \Omega(g(n))$. Means, at some point our function f is always going to be in between two bounds.

Example I:

$$f(n) = 4n^2 + 8n + 15 \in O(n^2)$$

$$\begin{aligned} f(n) &\leq cn^2 \\ 4n^2 + 8n + 15 &\leq cn^2 \\ 4n^2 + 8n + 15 &\leq 4n^2 + 8n^2 + 15n^2 = 27n^2 \end{aligned}$$

Thus, we have found a constant $c = 27$, from which our function always less is then $g = cn^2$. Therefore, $f \in O(g)$.

Ω ∈ Θ

Example II:

$$n^3 \notin O(n^2)$$

Suppose the opposite, then

$$n^3 \in O(n^2) \implies n^3 \leq cn^2 \implies n \leq c \forall n \geq N, c \in R$$

Choose $n = \lceil n_0 + c + 1 \rceil$ which is a natural number.

$$\lceil n_0 + c + 1 \rceil \leq c$$

This is a contradiction. Therefore, $n^3 \notin O(n^2)$.

QED

75.2.2 Properties of the Complexity Classes

Linearity:

If $g(n) = \alpha f(n) + \beta$ where $\alpha, \beta \in R^+$ and $f \in \Omega(1)$ then

$$g \in O(f)$$

Addition:

$$f + g \in O(\max(f, g)) = \begin{cases} O(g) & \text{if } f \in O(g) \\ O(f) & \text{if } g \in O(f) \end{cases}$$

Multiplication:

$$a \in O(f) \wedge b \in O(g) \implies ab \in O(fg)$$

Limit:

If the limit

$$\lim_{n \rightarrow \infty} \frac{g(n)}{f(n)}$$

exists, then $g \in O(f)$. The reverse does not count.

75.2.3 Master's Theorem


Help us find the time complexity of divide-and-conquer recurrences:

$$T(n) = aT\left(\left\lceil \frac{b}{n} \right\rceil\right) + c(n^d)$$

Where:

- $a \geq 1$ is the number of sub-problems
- $b > 1$ is the factor by which the problem size shrinks
- $c(n^d), d \geq 1$ is the cost of the work outside recursion (e.g., combining results)

Imagine a tree generated by each time we divide the problem until some point where solving it becomes trivial $\Theta(1)$. This tree is going to have a depth of $\log_b n$ which are the times we can divide the problem and a max width of $n^{\log_b a}$ which is the number of leafs at the bottom.



Work at each level: $n, n, n, \dots, n \Rightarrow \log n$ levels

Total work: $n \log n$

Therefore, our total work is going to be

$$\begin{aligned}
 T(n) &= c(n) + ac(n/b) + a^2c(b/n^2) + \dots + \Theta(n^{\log_b a}) \\
 &= \Theta(n^{\log_b a}) + \sum_{j=0}^{\log_b(n)-1} a^j c(n/b^j)
 \end{aligned}$$

Cases

When looking at the tree generated by the divide-and-conquer we can get three cases:

$$T(n) \in \begin{cases} \text{The problem gets easier} & O(n^d) \text{ if } a < b^d \\ \text{The hardness is evenly distributed} & O(n^d \log n) \text{ if } a = b^d \\ \text{The problem get harder} & O(n^{\log_b a}) \text{ if } a > b^d \end{cases}$$

Example:

$$T(n) = 4T\left(\frac{n}{2}\right) + c(n)$$

Here $a = 4$, $b = 2$ and $d = 1$. Since $d < \log_b a$ $1 < \log_2 4 = 2$

$$T(n) \in O(n^{\log_b a}) = O(n^2)$$

75.2.4 How To Identify Time Complexity of a Function

The process can be summarized into 5 steps:

1. Write the definition of the bound we are looking for.
2. Identify the most dominant term in $f(n)$. The one that grows the faster.
3. Set up the inequality and solve for c .
4. Take the limit of $\frac{f(n)}{g(n)}$.
5. Choose a c using the knowledge of the limit and find n_0 .

Example:

Find the time complexity $O(n)$, the constant, at the start term for the inequality of $\sqrt{4n}$.

Step 1: Definition

$$f(n) \in O(g(n)) \iff f(n) \leq cg(n)$$

Step 2: Dominant Term

The dominant term is $\sqrt{n}$.

Step 3: Inequality

We are going to assume that it is part of $O(\sqrt{n})$

$$\sqrt{n} \leq c\sqrt{n}$$

Step 4: Limit

$$\lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{n}} = 1 \leq c$$

Step 5: Choose c

Let us choose $c = 2$ which gives us $n_0 = 1$.

75.2.5 How to Find Time Complexity In Code**1. Count Primitive Operations**

- Additions, multiplications, comparisons, etc.

2. Identify Loops

- Single loop $\Rightarrow O(n)$
- Nested loop $\Rightarrow$ Multiply the complexities

3. Check Recursive Patterns

- Use recurrence relations (often solve with Master's Theorem)

4. Ignore Constants

- Drop lower-order terms and constants (e.g., $O(n^2 + 3n + 5) \Rightarrow O(n^2)$)

Examples**Example I: Simple Loop**

```
for i in range(n):
    print(i)
```

$\Rightarrow O(n)$

Example II: Nested Loop

```
for i in range(n):
    for j in range(n):
        print(i, j)
```

$\Rightarrow O(n^2)$

75.2.6 Common Time Complexities

Complexity	Name	Example Algorithms
$O(1)$	Constant	Array access
$O(\log n)$	Logarithmic	Binary search
$O(n)$	Linear	Linear search
$O(n \log n)$	Linearithmic	Merge Sort, Heap Sort
$O(n^2)$	Quadratic	Bubble Sort, Selection Sort
$O(n^3)$	Cubic	Naive matrix multiplication
$O(2^n)$	Exponential	Recursive Fibonacci
$O(n!)$	Factorial	Traveling Salesman (brute force)

75.3 State Machines and Languages

75.3.1 Languages and Alphabets

A *language* L is a subset of words of an *alphabet* Σ which is a finite non-empty set of symbols which always includes the ε -word, the empty word.

Concatenation

An operation which can be done with members of an alphabet is the so called *concatenation* which is like basic addition. Therefore, it is associative and when added ε , the zero of alphabets it does not affect the words or its length.

75.3.2 Inductive Definitions

In the field of languages and alphabets an inductive function $f : \Sigma^* \rightarrow X$ will be defined in the following way.

1. *The base case:* $f(\varepsilon)$
2. *The induction step:* where the actual behavior of the function is defined. Helper functions can also be defined using the same structure.

It also has the following properties

- $|\varepsilon| = 0$
- For $w = v + a$ then $|w| = |v| + 1$ for $a \neq \varepsilon$

Example:

Define a function which count the occurrences of ue in a word with letters of the alphabet $\Sigma = \{a, \dots, z\}$.

$$\begin{aligned} f(\varepsilon) &= 0 \\ f(w \cdot x) &= \begin{cases} f(w) & \text{if } x \neq e \\ \text{help}(w) & \text{if } x = e \end{cases} \end{aligned}$$

The auxiliary function $\text{help} : \Sigma^* \rightarrow \mathbb{N}$ is defined as:

$$\text{help}(w \cdot x) = \begin{cases} f(w) + 1 & \text{if } x = u \\ f(w) & \text{otherwise} \end{cases}$$

That is, $\text{help}(w) = f(w) + 1$ if the last letter of w is u , and $f(w)$ otherwise.

Head and Tail

Head returns the first symbol of a word and is defined by

$$\begin{aligned} \text{head}(\varepsilon) &= \varepsilon \\ \text{head}(\varepsilon + a) &= a \\ \text{head}(v + a) &= \text{head}(v) \end{aligned}$$

Tail removes the first symbol of a word and is defined by

$$\begin{aligned} \text{tail}(\varepsilon) &= \varepsilon \\ \text{tail}(\varepsilon + a) &= \varepsilon \\ \text{tail}(v + a) &= \text{tail}(v) + a \end{aligned}$$

75.3.3 Fixes

Let Σ be an alphabet and let $v, w \in \Sigma^*$ be words.

- v is called a *substring* of w iff $\exists s, t \in \Sigma^* : w = svt$.
- v is called a *suffix* of w iff $\exists s \in \Sigma^* : w = sv$ - we write $v \sqsupseteq w$.
- v is called a *prefix* of w iff $\exists t \in \Sigma^* : w = vt$ - we write $v \sqsubseteq w$.
- v is called a *proper substring* (*prefix/suffix*) of w iff $v \neq \varepsilon \wedge v \neq w$ and v is a substring (prefix/suffix) of w . We then write $v \sqsubset w$ or $v \sqsubset w$.

ε is prefix suffix and substring, of all words.

75.3.4 The Canonical Order

Let $\Sigma = \{a_1, \dots, a_n\}$ be an alphabet with $n \geq 1$, and let $< \subseteq \Sigma \times \Sigma$ be an order on Σ with $a_1 < a_2 < \dots < a_n$.

We define the *canonical order* on Σ^* as follows:

$$u < v \quad \text{iff} \quad |u| < |v| \quad \vee \quad (|u| = |v| \wedge u = w \cdot a_i \cdot u' \wedge v = w \cdot a_j \cdot v' \wedge i < j)$$

for arbitrary $w, u, u', v, v' \in \Sigma^*$.

75.3.5 Languages

Let Σ be an alphabet. A *language* L over Σ is a subset of Σ^* ($L \subseteq \Sigma^*$).

$$\mathcal{P}(\Sigma^*) = \{L \mid L \subseteq \Sigma^*\},$$

the power set of Σ^* , is the set of all languages over Σ .

Furthermore, $L_\emptyset = \emptyset$ and $L_\varepsilon = \{\varepsilon\}$.

Another parts of a language is the *syntax* which defines the words in the language, and the *semantic* which determines the meaning of words.

75.3.6 Pumping Lemma

For a language L which can be recognized by a state machine, there is a natural number n such that for each word in L with length greater than or equal to n , the word w can be decomposed into three subwords x, y, z .

- $|xy| \leq n$
- $y \neq \varepsilon$
- $xy^iz \in L$ for all $i \geq 0$

Example:

Let $L = \{a^n b^n \mid n \geq 0\}$. This language is **not** regular, and the Pumping Lemma can be used to prove this by contradiction.

75.3.7 Algorithm

Given two alphabet Σ_1, Σ_2 and $\mathcal{D} \subseteq \Sigma_1^*$. An *algorithm* is a total function

$$A : \mathcal{D} \rightarrow \Sigma_2^*$$

Two algorithms are equivalent if and only if

$$\forall x \in \mathcal{D} : A(x) = B(x)$$

75.3.8 The Decision Problem

The *decision problem* $\langle \Sigma, L \rangle$ for a given alphabet Σ and a language $L \subseteq \Sigma^*$ is to decide, for each $w \in \Sigma^*$, whether

$$w \in L \quad \text{or} \quad w \notin L.$$

Therefore, one also speaks of the *word problem* $\langle \Sigma, L \rangle$.

An algorithm $A: \Sigma^* \rightarrow \{0, 1\}$ *solves* the decision problem $\langle \Sigma, L \rangle$ if, for all $w \in \Sigma^*$, we have

$$A(w) = \begin{cases} 1 & \text{if } w \in L, \\ 0 & \text{if } w \notin L. \end{cases}$$

We also say: A *recognizes* L .

75.3.9 Recursive Languages

Let Σ be an alphabet and let $L \subseteq \Sigma^*$ be a language. L is called *recursive* if and only if there exists an algorithm A that solves the decision problem $\langle \Sigma, L \rangle$.

75.3.10 Enumeration Problem

Let Σ be an alphabet, $L \subseteq \Sigma^*$ a language, and $n \in \mathbb{N}$. The *enumeration problem* $\langle \Sigma, L, n \rangle$ for L consists of listing the canonical first n words of L .

An algorithm $A: \mathbb{N} \rightarrow \mathcal{P}(L)$ is an *enumeration algorithm* for L if, for every input $n \in \mathbb{N}$, A outputs the canonical first n words $w_0, w_1, \dots, w_{n-1}$ of the language L .

75.3.11 Grammars

Let

$$G = (V, \Sigma, R, S)$$

where:

- V is a finite set of *variables* (also called non-terminal symbols).
- Σ is a finite set of *terminal symbols*, such that $V \cap \Sigma = \emptyset$.
- R is a finite set of *production rules* of the form

$$A \rightarrow \alpha,$$

where $A \in V$ and $\alpha \in (V \cup \Sigma)^*$.

- $S \in V$ is the **start symbol**.

Thus, a grammar G defines a *formal language* $L(G)$ as:

$$L(G) = \{ w \in \Sigma^* \mid S \Rightarrow^* w \}$$

where $\Rightarrow^*$ denotes the reflexive and transitive closure of the derivation relation defined by R .

Context Free Grammars

75.4 State Machines

A *state machine* (or *finite state machine*, FSM) is a computational model used to design algorithms, systems, or software by breaking them down into a finite number of *states*, *transitions*, *inputs*, and *outputs*. It behaves according to a current state and can transition to other states based on input conditions.


State machines are widely used in fields like software engineering, digital electronics, linguistics, and artificial intelligence to model dynamic systems.

75.4.1 Deterministic Finite Automaton (DFA)

DFA's are defined by a quintuple $(Q, \Sigma, \delta, q_0, F)$ which consists of Q states, an input alphabet Σ , a transition function $\delta : Q \times \Sigma \rightarrow Q$, a start state $q_0 \in Q$ and a set of terminations states $F \subset Q$.

This kind of automaton has the particularity that each state has *exactly one transition* for each possible input. Therefore, *deterministic*.

Example:



Here our language is

$$L = \{w \in \{a, b\}^* \mid \text{the sequence of transitions on } w \text{ leads to state } q_3\}$$

75.4.2 Non-Deterministic Finite Automaton (NFA)


An NFA is a state machine defined by a 5-tuple $(Q, \Sigma, \delta, q_0, F)$, where: Q is a finite set of states, Σ is the input alphabet, $\delta : Q \times \Sigma \rightarrow \mathcal{P}(Q)$ is the transition function (mapping to a set of states), $q_0 \in Q$ is the start state and $F \subset Q$ is the set of accepting states.

Unlike a DFA, an NFA allows:

- Multiple possible next states for a given input.
- No transition at all.

Example:

State	a	b
q_0	$\{q_0, q_1\}$	$\{q_0\}$
q_1	$\{q_2\}$	$\{q_1\}$
q_2	$\{q_3\}$	$\emptyset$
q_3	$\{q_3\}$	$\{q_3\}$



Here our language is still:

$$L = \{w \in \{a, b\}^* \mid \text{some computation path leads to } q_3\}$$

75.4.3 Epsilon-NFA (ε -NFA)

An ε -NFA is a nondeterministic finite automaton that allows transitions on the empty string ε , meaning the automaton can change states without consuming input.

It is defined as a 5-tuple $(Q, \Sigma, \delta, q_0, F)$, with the transition function:


$$\delta : Q \times (\Sigma \cup \{\varepsilon\}) \rightarrow \mathcal{P}(Q)$$

This means the automaton have the following properties:

- It can transition normally on $a \in \Sigma$ or move to other states "for free" via ε -transitions.
- *Epsilon-Closure*: It is the set of all states reachable via ε of a particular state plus itself.

Example:

State	a	b	ε
q_0	$\emptyset$	$\{q_0\}$	$\{q_1\}$
q_1	$\{q_2\}$	$\emptyset$	$\emptyset$
q_2	$\emptyset$	$\{q_3\}$	$\emptyset$
q_3	$\{q_3\}$	$\{q_3\}$	$\emptyset$



The language is:

$$L = \{w \in \{a, b\}^* \mid w \text{ contains the substring } ab\}$$

via the path:


$$q_0 \xrightarrow{\varepsilon} q_1 \xrightarrow{a} q_2 \xrightarrow{b} q_3$$

75.4.4 How to Construct a State Machine

To construct a state machine it is necessary that an alphabet and a language is given. Then the process revolves around identifying which words are a part of the language and how they can be expressed as a list of states and transition.

Example:

Given is the alphabet $\Sigma = \{a, b, c\}$ and the language $L = \{w \in \Sigma^* \mid w = a(bc)^m a^n \text{ and } \}$. Construct a DFA.



75.4.5 Regular Expressions

Regular expressions (regex) are a practical application of finite automaton (DFA/NFA). They define search patterns for strings and are processed internally using state machines.

Example:

Regex: $a(b|c)^*d$

- *Start State*: Awaiting input.
- *Transition on 'a'*: Move to a state expecting b or c .
- *Loop State*: Repeats b or c any number of times.
- *Final Transition*: Ends with a d .

Regular expressions describe sets of strings using a small set of operations:


- *Concatenation* ($R.S$): A string from R followed by a string from S .
- *Union* ($R + S$): A string from either R or S .
- *Kleene star* (R^*): Zero or more repetitions of strings from R .

Special symbols:

- $\emptyset$: Matches no string.
- ε : Matches the empty string.
- Literal characters (e.g., a, b, c): Match themselves.

Below are ε -NFAs (nondeterministic automaton with ε -transitions) for some regular expressions:


$R + E$




$R.E$



R^*



Example: $(a+b)^*$



75.4.6 State Machine Conversions and Minimization

Non-Deterministic Finite Automaton (NFA) to Deterministic Finite Automaton (DFA)


The process for the conversion is the so called *Power-Set-Construction-Method* or *Subset-Construction-Method*.

Steps:

1. Write the transition table of the given automaton.
2. Write a similar table with the transition as columns.
3. Start with q_0 and find which states can be reached with the transitions. If from our current state we can reach more than one state with our current transition, write the possible states as a set, then Proceed to write the set as the next entry to check (write in the next row).
4. We terminate the algorithm when we after evaluating our states we only get states we have already visited.

Example:

Convert the NFA to a DFA




Step 1: Transition Table

State	0	1
q_0	$\{q_0, q_1\}$	q_0
q_1	$\emptyset$	q_2
q_2	$\emptyset$	$\emptyset$

Step 2: Table of new states

State	0	1
q_0	$\{q_0, q_1\}$	q_0
$\{q_0, q_1\}$	$\{\{q_0, q_1\}, \emptyset\}$	$\{q_0, q_2\}$
$\{q_0, q_2\}$	$\{\{q_0, q_1\}, \emptyset\}$	$\{q_0, \emptyset\}$

Step 3: Construct DFA



Epsilon-NFA to DFA


The process is very similar to the previous one. The catch is that we have to keep in mind that our initial state is not only q_0 but also all other from q_0 reachable states via epsilon.

Steps:

1. Write a transition table.
2. Write the new initial state which is going to be a set of states with the original initial state and all states reachable via ε from it. This also includes states reachable via multiple epsilon transitions. This is the epsilon closure of the initial state.
3. Find the states reachable via the first transition of this epsilon closure.
4. Then find the epsilon closure of the previously found set of states. This becomes a new state in the DFA.
5. Repeat the process with each new created state and transition until you get no new epsilon closure.

Example:

Convert to DFA



Step 1: Transition Table

State	a	b	c	ε
p	$\emptyset$	q	r	$\{q, r\}$
q	p	r	p, q	$\emptyset$
r	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$

Step 2: Epsilon Closure and Transitions


The initial state is $A = \varepsilon(p) = \{p, q, r\}$.

Using the transition a we get the set $\{p\}$ with $\varepsilon(p) = \{p, q, r\}$. This state is just A again. Using the transition b we get the set $\{q, r\}$ with $B = \varepsilon(\{q, r\}) = \{q, r\}$. And finally with c we get the set $\{p, q, r\}$ with $\varepsilon(\{p, q, r\}) = \{p, q, r\}$.

Repeating this process with the rest of state will give us:

State	a	b	c
$\{p, q, r\}$	$\{p, q, r\}$	$\{q, r\}$	$\{p, q, r\}$
$\{q, r\}$	$\{p, q, r\}$	r	$\{p, q, r\}$
r	$\emptyset$	$\emptyset$	$\emptyset$

Step 3: Construct the Automaton




75.4.7 Epsilon-NFA to NFA

Here the key is to eliminate the *varepsilon* transitions from the automaton. It can be done with the following steps:

1. Write a table empty transition table of the automaton but without the transition epsilon. And then all epsilon closures of each state in a list
2. For state and transition. Take ε -closure of the state, find where do these states go with the current transition and the take their epsilon closure. The final closure goes into to the corresponding transition column of the current state.
3. Look for new final states. A final state will be a state that can be reached only using epsilon transitions.
4. Finally, write the new automaton using the information from the transition table with the new final states. It is important to note that the automaton is not minimal.

Example:

Transform to a nondeterministic automaton



Step 1: Set up table and closures

State	0	1
q_0		
q_1		
q_2		
q_3		

ϵ -Closures

$$\begin{aligned} q_0 &= \{q_0, q_1, q_2, q_3\} \\ q_1 &= \{q_1, q_2, q_3\} \\ q_2 &= \{q_2, q_3\} \\ q_3 &= \{q_3\} \end{aligned}$$


Step 2: Filling the table using the transition and closures

State	0	1
q_0	q_3	q_0
q_1	q_3	q_0
q_2	$\emptyset$	q_0
q_3	$\emptyset$	$\emptyset$

Step 3: Identify Final States

Here every state is a final state of the every state contains the final state in the epsilon closure.

Step 4: Draw the automaton



NFA to Regex

Converting a regular expression into a NFA is not really difficult, but the other direction is not so trivial. The most common approach to tackle this type of conversion is via a system of equations.

You are given some NFA with n states and m transitions. Set your system as follows:

1. Write an equation L_i for each state q_i . Each equation L_i represents the set of strings (language) accepted when starting from state q_i . These equations will be defined recursively in terms of transitions to other states.
2. For each L_i , describe all transitions from state q_i : For every possible outgoing transition from q_i to some q_j on symbol(s) w , add a term wL_j to the equation. If q_i is an accepting state, also include ϵ (the empty string).

Example: If from state 1 there is a transition $a \rightarrow 1$ and $b \rightarrow 2$, and state 1 is accepting, then

$$L_1 = aL_1 \cup bL_2 \cup \epsilon$$

3. Rewrite the equations using regular expression notation. Replace set notation with regex syntax: $\cup \rightarrow +$, concatenation is written without operator (e.g., aL_1) and replace sets of symbols (e.g., $\{a, b\}$) with grouped regex (e.g., $(a + b)$)

Example:

$$L_1 = aL_1 + bL_2 + \epsilon$$

4. Substitute equations recursively to reduce the system. There are two cases:

- If an equation L_i does not reference itself (i.e., no recursion), substitute it into the other equations directly.
- If an equation L_i does reference itself (e.g., $L_i = rL_i + s$), apply *Arden's Lemma*:

$$\text{If } L = YL \cup X, \text{ then } L = Y^*X$$

Example:

- Write for each state the equation L_i .
- Now treat each state as a start state and look for the sets of words this state accepts and lead to a terminating state then combine them with $\cup$. If the current state is also a terminating state then ε must also be added. Example: $L_1 = \{ab\}L_1 \cup bL_2$.

A trick is to write the table with the transition and for each row (state) you write $L_{row} = transition_1 L_{state_x} + transition_2 L_{state_y} + \dots$. Where the transition is a set of letters of the alphabet.

- Repeat step 2 for all L_i and then replace the set of words with regular expressions and the $\cup$ with $+$ Example: $L_1 = (a + b)L_1 + bL_2$.
- Start substituting equations into other equations. Here there are two cases. Either, the equation $L_x, \dots$ depends only on other equations different from itself or $L_x, \dots$ also depends on itself which causes infinite recursion. For the first case the problem solves itself by just doing the substituting until we have only one equation which is going to be our regex, but for the other case we will use *Arden's Lemma*.

$$L = (a + b)L + b \Rightarrow L = (a + b)^*b$$

Continue substituting until all equations are resolved. The expression for the initial state gives the regular expression for the NFA.

Arden's Lemma

Given X, Y as formal language with $\varepsilon \notin Y$. It applies that:

$$L = YL \cup X \text{ has exactly one solution: } L = Y^*X$$

And the special case $X = \emptyset$

$$L = YL + \emptyset = Y^*\emptyset = \emptyset$$

Proof:

To show $L = Y^*X$ is a solution and more specific the only solution.

$$\begin{aligned} L &= YL \cup X \\ Y^*X &= YY^*X \cup X \end{aligned} \text{ This shows that it is indeed a solution}$$

For the second part we can argue that if L is a solution for the equation then

$$Y^*X \subseteq L \implies X \cup YX \cup YYX \cup \dots \subseteq L$$

Here we always took longer words which are also part of the language and united them. For the second part we are going to use shorter words.

Assume that

$$L \subsetneq Y^*X$$


Then there is the shortest word w as the word in L which is not in Y^*X . But because $L = YL \cup X$ w must be either a word in X which is not possible because of $w \notin Y^*X$; or YL . This implies that w can be decomposed into uv with $u \in Y, u \neq \varepsilon, v \in L$. But this is a contradiction then w would be a word of the form Y^*X which not true, and this would imply that there is an even shorter word for the shortest word and so on. Therefore

$$L \subseteq Y^*X$$

□◻◻

Example:

Convert to Regex



Step 1: Initial Equations

$$L_1 = \{a, b\}L_1 \cup aL_2 \cup bL_3$$

$$L_2 = \varepsilon \cup bL_2 \cup bL_4$$

$$L_3 = \{a, b\}L_1 \cup \varepsilon \cup aL_4$$

$$L_4 = aL_2$$

Step 2: Regex Equations

$$L_1 = (a + b)L_1 + aL_2 + bL_3$$

$$L_2 = \varepsilon + bL_2 + bL_4$$

$$L_3 = (a + b)L_1 + \varepsilon + aL_4$$

$$L_4 = aL_2$$

Step 3: Substitution

L_4 in L_3 and L_2

$$L_3 = (a + b)L_1 + \varepsilon + aaL_2$$

$$L_2 = \varepsilon + bL_2 + baL_2$$

Now we can solve L_2 using the lemma.

$$\begin{aligned}
L_2 &= \varepsilon + bL_2 + baL_2 \\
&= \varepsilon + (b + ba)L_2 \\
&= (b + ba)^* \varepsilon \\
&= (b + ba)^*
\end{aligned}$$

L_2 in L_3 and L_1

$$\begin{aligned}
L_1 &= (a + b)L_1 + a(b + ba)^* + bL_3 \\
L_3 &= (a + b)L_1 + \varepsilon + aa(b + ba)^*
\end{aligned}$$

L_3 in L_1

$$\begin{aligned}
L_1 &= (a + b)L_1 + a(b + ba)^* + b((a + b)L_1 + \varepsilon + aa(b + ba)^*) \\
&= (a + b)L_1 + a(b + ba)^* + b(a + b)bL_1 + b + baa(b + ba)^* \\
&= (a + b)L_1 + b(a + b)L_1 + a(b + ba)^* + b + baa(b + ba)^* \\
&= ((a + b) + b(a + b))L_1 + a(b + ba)^* + b + baa(b + ba)^* \\
&= ((a + b) + b(a + b))L_1 + a(b + ba)^* + b + baa(b + ba)^* \\
&= ((a + b) + b(a + b))^*(a(b + ba)^* + b + baa(b + ba)^*)
\end{aligned}$$

This is the answer but we can simplify. Here $((a + b) + b(a + b))$ is a word of $(a + b)^*$

$$\begin{aligned}
L_1 &= ((a + b) + b(a + b))^*(a(b + ba)^* + b + baa(b + ba)^*) \\
&= (a + b)^*(a(b + ba)^* + b + baa(b + ba)^*)
\end{aligned}$$

75.4.8 Minimizing State Machines


Minimization aims to create an equivalent DFA with the *fewest number of states*. One widely used method is the *table-filling algorithm*.

Table Method for DFA Minimization

Steps:

1. Put all non-accepting states in an equivalence class α , and the accepting states in another called Ω . Write a table with the following columns: state, transition 1, transitions 2,..., transition n. On the first row put *alpha* and in the state column the states of the class. Then do the same with Ω . You can also optionally make a table of transition for the states.
2. Start evaluating each state with each transition, and write the equivalence class reached by that transition. This means that you have to write the name of the set (class) the reached state is in. For example if we evaluate a with transition 0, and it leads to b and b is in the same set as a then for the row a with the transition 0 we write the name of that set.
3. For each state in the current class which has a tuple of classes different from the rest create a new class which is going to be included in the next iteration of the table. Then repeat step 2.
4. Terminate the process if no more new class tuple appear.
5. Build a new DFA with merged equivalent states and updated transitions.

Example:



Step 1 and 2: Table

Class	State	0	1
α	A		
	B		
	C		
	D		
Ω	E		

Step 3: Evaluation

First iteration

Class	State	0	1
α	A	α	α
	B	α	α
	C	α	α
	D	α	Ω
Ω	E	α	α


Second iteration

Class	State	0	1
α	A	α	α
	B	α	β
	C	α	α
β	D	α	Ω
Ω	E	α	α

Third and final iteration

Class	State	0	1
α	A	γ	α
	C	γ	α
β	D	γ	Ω
γ	B	γ	β
Ω	E	γ	α

State 4: Draw the automaton



75.5 Regex

Regex is powerful tool for searching pattern inside a text which can be represented as state machines. In which our regular expression defines the transitions between the states.

75.5.1 Basic Characters

- `a` : Matches character 'a' exactly.
- `abc` : Matches the string "abc" exactly.

75.5.2 Meta Characters (Special Characters)

- `.` : Matches any single character except a newline.
- `\` : Escape character (used to escape special characters, e.g., `\.` to match a literal period).
- `^` : Matches the start of a string.
- `$` : Matches the end of a string.
- `|` : Alternation (logical OR, e.g., `cat|dog` matches either "cat" or "dog").

75.5.3 Character Classes

- `[abc]` : Matches any one of 'a', 'b', or 'c'.
- `[^abc]` : Matches any character except 'a', 'b', or 'c'.
- `[a-z]` : Matches any lowercase letter.
- `[A-Z]` : Matches any uppercase letter.
- `[0-9]` : Matches any digit.
- `[a-zA-Z0-9]` : Matches any alphanumeric character.
- `.` : Matches any character except newline.
- `\d` : Matches any digit (equivalent to `[0-9]`).
- `\D` : Matches any non-digit character.
- `\w` : Matches any word character (letters, digits, underscore).
- `\W` : Matches any non-word character.
- `\s` : Matches any whitespace character (spaces, tabs, newlines).
- `\S` : Matches any non-whitespace character.

75.5.4 Quantifiers

- `*` : Matches 0 or more occurrences of the preceding character (e.g., `a*` matches "", "a", "aa", etc.).
- `+` : Matches 1 or more occurrences (e.g., `a+` matches "a", "aa", etc.).
- `?` : Matches 0 or 1 occurrence (e.g., `a?` matches "" or "a").
- `{n}` : Matches exactly `n` occurrences (e.g., `a{3}` matches "aaa").
- `{n,}` : Matches at least `n` occurrences (e.g., `a{2,}` matches "aa", "aaa", etc.).
- `{n,m}` : Matches between `n` and `m` occurrences (e.g., `a{2,4}` matches "aa", "aaa", or "aaaa").

75.5.5 Grouping and Capturing

- `(abc)` : Capturing group, treats "abc" as a single unit.
- `(?:abc)` : Non-capturing group, groups "abc" without capturing.
- `\n` : Refers to the *n*th captured group (e.g., `(\w)\1` matches doubled letters like "oo").

75.5.6 Assertions and Anchors

- `^` : Matches the start of a string.
- `$` : Matches the end of a string.
- `\b` : Matches a word boundary (e.g., `\bword\b` matches "word" but not "sword").
- `\B` : Matches a non-word boundary.
- `(?=...)` : Positive look ahead (e.g., `a(=b)` matches 'a' only if followed by 'b').
- `(?!...)` : Negative look ahead (e.g., `a(?!b)` matches 'a' only if NOT followed by 'b').
- `(?<=...)` : Positive look behind (e.g., `(?<=a)b` matches 'b' only if preceded by 'a').
- `(?<!=...)` : Negative look behind (e.g., `(?<!=a)b` matches 'b' only if NOT preceded by 'a').

75.5.7 Advanced Constructs

- `(?(condition)yes|no)` : Conditional expression.
- `(?P<name>...)` : Named capturing group (e.g., `(?P<word>\w+)`).
- `(?P=name)` : Refers to a named capturing group.
- `(?#...)` : Comment within regex for readability.
- `(*UTF8)` : Enforces UTF-8 mode for international text processing.
- `(?>...)` : Atomic group (prevents backtracking within the group).

75.5.8 Examples

Example: Email validation

```
^[a-zA-Z0-9._%+-]+@[a-zA-Z0-9.-]+\.[a-zA-Z]{2,}$
```

Matches valid email addresses.

Example: Phone number validation

```
^\(?\d{3}\)?[-.]?\d{3}[-.]?\d{4}$
```

Matches US phone numbers like (123) 456-7890, 123-456-7890, or 123.456.7890.

Example: URL validation

```
^https?:\/\/(www\.)?[a-zA-Z0-9.-]+\.[a-zA-Z]{2,}(\S*)?$
```

Matches URLs like `http://example.com` or `https://www.example.com/path`.

Example: Hex color code validation

```
^#[A-Fa-f0-9]{6}|[A-Fa-f0-9]{3}$
```

Matches hex color codes like `#FFF` or `#A1B2C3`.

Example: Date format (YYYY-MM-DD) validation

```
^\d{4}-\d{2}-\d{2}$
```

Match dates formatted as `2024-03-01`.

75.6 Graph Theory

A *Graph* $G = \langle V, E \rangle$ consists of a finite set of vertices V and a relation $E \subseteq V \times V$ over the set. Depending on the traits of E it can be categorized into

- *Undirected*: E is symmetric.
- *Directed*: E is not symmetric.
- *Complete*: $E = V \times V$ for directed and $|E| \approx |V|(|V| - 1)/2$ for undirected graphs.
- *Loop-free*: $\forall (v, v) \in V \times V : (v, v) \notin E$
- *Weighted*: $G = \langle V, E, \phi \rangle$, where $\phi : E \rightarrow R$ and R is an arbitrary set. It mostly refers to adding a cost to each edge.
- *Dense*: $|E| \approx |V|^2$ for directed and $|E| \approx |V|(|V| - 1)/2$ for undirected graphs.
- *Sparse*: $|E| \ll |V|^2$ for directed and $|E| \ll |V|(|V| - 1)/2$ for undirected graphs.

In computer science the density of a graph is a main criteria when modeling a graph.

75.6.1 Connected Graphs

A graph $G = (V, E)$ undirected, then it is called *connected* if for all pairs $(v_i, v_j) \subseteq V \times V$ there exists a path from v_i to v_j . We define E_S the symmetric relation.

$$(v_1, v_2) \in E \iff (v_1, v_2) \in E_S \wedge (v_2, v_1) \in E_S$$

If G is directed then it is connected if and only if its undirected version is connected.

75.6.2 Paths

Given a graph G , we call a *path* a set of vertices if

$$\forall i \in \{1, \dots, k-1\} : (w_i, w_{i+1}) \in E$$

The most common types of paths are

- *Circle*: $w_1 = w_k$
- *Eulerian Circuit*: Each edge will be visited one time. $\forall (s, t) \in E : \exists_1 i : s = w_i \wedge t = w_{i+1}$
- *Hamiltonian Circuit*: Each vertex will be visited one time $\forall v \in V : \exists_1 i : v = w_i$
- *Loop-free*: $\forall (v, v) \in V \times V : (v, v) \notin E$

Path Cost

We define the *path cost* for a weighted graph as

$$\phi(p) = \sum_{i=1}^{k-1} \phi(w_i, w_{i+1})$$

75.6.3 The Adjacency Matrix

For a graph G with vertices $V = \{v_1, \dots, v_n\}$, we define the *adjacency matrix* $A = (a_{ij})$ as a $n \times n$ matrix with

$$a_{ij} = \begin{cases} 1 & \text{if } (v_i, v_j) \in E \\ 0 & \text{else} \end{cases}$$

75.6.4 The Relax Operation

In graph theory when referring to the action of updating the cost of visiting a vertex in a weighed graph we call this operation *relaxation*.

$$\text{relaz}(u, v, \text{weight}) = \begin{cases} d[u] + \text{weight}(u, v) & \text{if } d[u] + \text{weight}(u, v) < d[v] \\ d[u] & \text{else} \end{cases}$$