

Functional Analysis

Alex Rutar*
University of Waterloo

Fall 2019[†]

*arutar@uwaterloo.ca

[†]Last updated: September 13, 2019

Contents

Chapter I	Fundamentals of Functional Analysis	
1	Basic Elements of Functional Analysis	1
2	Linear operators and linear functionals	5
3	Axiom of Choice and the Hahn-Banach Theorem	8

I. Fundamentals of Functional Analysis

1 BASIC ELEMENTS OF FUNCTIONAL ANALYSIS

Throughout, we denote by \mathbb{F} either the field \mathbb{R} or the field \mathbb{C} .

BANACH SPACES

Definition. Let X be a vector space over \mathbb{F} . A **norm** is a functional $\|\cdot\| : X \rightarrow \mathbb{R}$ such that it is

- (*non-negative*) $\|x\| \geq 0$ for any $x \in X$
- (*non-degenerate*) $\|x\| = 0$ if and only if $x = 0$
- (*subadditivity*) $\|x + y\| \leq \|x\| + \|y\|$ for $x, y \in X$
- (*$|\cdot|$ -homogeneity*) $\|\alpha x\| = |\alpha| \|x\|$ for $\alpha \in \mathbb{F}$, $x \in X$.

We call the pair $(X, \|\cdot\|)$ a **normed vector space**. Furthermore, we say that $(X, \|\cdot\|)$ is a **Banach space** provided that X is complete with respect to the metric $\rho(x, y) = \|x - y\|$.

Example. (i) $(\mathbb{F}, |\cdot|)$ is a Banach space.

(ii) $(\mathbb{F}^n, \|\cdot\|_p)$, $x = (x_j)_{j=1}^n$,

$$\|x\|_p = \begin{cases} \left(\sum_{j=1}^n |x_j|^p \right)^{1/p} & 1 \leq p < \infty \\ \max_{j=1, \dots, n} |x_j| & p = \infty \end{cases}$$

(iii) Consider the space

$$L_p^{\mathbb{F}} = \left\{ f : [0, 1] \rightarrow \mathbb{F} \mid f \text{ is Lebesgue measurable, } \left(\int_0^1 |f|^p \right)^{1/p} < \infty \right\} \Big/ \sim_{\text{a.e.}}$$

where $1 \leq p < \infty$.

(iv) $L_{\infty}^{\mathbb{F}}[0, 1]$, $\|f\|_{\infty} = \text{ess sup}_{t \in [0, 1]} |f(t)|$.

(v) Let (X, d) be a metric space. Then

$$C_b^{\mathbb{F}}(X) = \{ f : X \rightarrow \mathbb{F} \mid f \text{ is continuous and bounded } \}, \quad \|f\|_{\infty} = \sup_{x \in X} |f(x)|$$

is a Banach space.

Here is a more interesting example:

Example. Let (X, d) be a metric space. We define the space of Lipschitz functions

$$\text{Lip}^{\mathbb{F}}(X, d) = \left\{ f : X \rightarrow \mathbb{F} \left| f \text{ is bounded, } L(f) = \sup_{\substack{x, y \in X \\ x \neq y}} \frac{|f(x) - f(y)|}{d(x, y)} < \infty \right. \right\}$$

We note that for $f : X \rightarrow \mathbb{F}$ that

$$f \in \text{Lip}^{\mathbb{F}}(X, d) \Leftrightarrow \text{there is } L \geq 0 \text{ s.t. } |f(x) - f(y)| \leq Ld(x, y) \text{ for all } x, y \in X \quad (1.1)$$

It is easy to verify that $L(f) = \min\{L \geq 0 : (1.1) \text{ holds for } f\}$. It is an easy exercise to see that $\text{Lip}^{\mathbb{F}}$ is a vector space, and that $L : \text{Lip}^{\mathbb{F}}(X, d) \rightarrow \mathbb{R}$ is a **semi-norm** (non-negative, subadditive, $|\cdot|$ -homogeneous). However, we do not have non-degeneracy (for example, constants are taken to 0). We define the Lipschitz norm

$$\|f\|_{\text{Lip}} = \|f\|_{\infty} + L(f)$$

1.1 Proposition. $(\text{Lip}^{\mathbb{F}}(X, d), \|\cdot\|_{\text{Lip}})$ is a Banach space.

PROOF Let $(f_n)_{n=1}^{\infty}$ be a Cauchy sequence in $(\text{Lip}^{\mathbb{F}}(X, d), \|\cdot\|_{\text{Lip}})$. Since $\|\cdot\|_{\infty} \leq \|\cdot\|_{\text{Lip}}$ on $\text{Lip}^{\mathbb{F}}(X, d)$, we see that $(f_n)_{n=1}^{\infty}$ is uniformly Cauchy (and bounded), and hence there is $f = \lim_{n \rightarrow \infty} f_n$ in $C_b^{\mathbb{F}}(X)$, where the limit is taken with respect to $\|\cdot\|_{\infty}$, since $(C_b^{\mathbb{F}}(X), \|\cdot\|_{\infty})$ is a Banach space. If $x, y \in X$, then

$$\begin{aligned} |f(x) - f(y)| &= \lim_{n \rightarrow \infty} |f_n(x) - f_n(y)| \leq \sup_{n \in \mathbb{N}} |f_n(x) - f_n(y)| \\ &\leq \sup_{n \in \mathbb{N}} L(f_n) d(x, y) \leq \sup_{n \in \mathbb{N}} \|f_n\|_{\text{Lip}} d(x, y) \end{aligned}$$

Since Cauchy sequences are bounded, we see that $|f(x) - f(y)| \leq Ld(x, y)$, where $L = \sup_{n \in \mathbb{N}} \|f_n\|_{\text{Lip}} < \infty$. Thus by (1.1), $f \in \text{Lip}^{\mathbb{F}}(X, d)$. Exercise: one may verify that $\|f - f_n\|_{\text{Lip}} \rightarrow 0$. \blacksquare

Another collection of basic examples are given by the sequence spaces. We can define

$$\ell_1^{\mathbb{F}} = \left\{ x = (x_j)_{j=1}^{\infty} \in \mathbb{F}^{\mathbb{N}} \left| \|x\|_1 = \sum_{j=1}^{\infty} |x_j| < \infty \right. \right\}$$

It is easy to see that $(\ell_1, \|\cdot\|_1)$ is a normed vector space.

For $1 < p < \infty$, and write

$$\ell_p^{\mathbb{F}} = \left\{ x \in \mathbb{F}^{\mathbb{N}} \left| \|x\|_p = \left(\sum_{j=1}^{\infty} |x_j|^p \right)^{1/p} < \infty \right. \right\}$$

Note that $0 \in \ell_p$, $\alpha \in \mathbb{F}$, $\alpha x \in \ell_p$ if $x \in \ell_p$. Let $q = p/(p-1)$ so that $1/p + 1/q = 1$. Then q is called the **conjugate index**. We have

1.2 Proposition. (Young's Inequality) If $a, b \geq 0$ in \mathbb{R} , then $ab \leq a^p/p + b^q/q$, with equality only if $a^p = b^q$.

and

1.3 Proposition. (Hölder's Inequality) If $x \in \ell_p$ and $y \in \ell_q$, then $xy = (x_i y_i)_{i=1}^\infty \in \ell_1$, with

$$\sum_{i=1}^{\infty} |x_i y_i| \leq \|x\|_p \|y\|_q$$

with equality exactly when $\text{sgn}(x_i y_i) = \text{sgn}(x_k y_k)$ for all $j, k \in \mathbb{N}$ where $x_i y_i \neq 0 \neq x_k y_k$, and $|x|^p = (|x_j|^p)_{j=1}^\infty$ and $|y|^q$ are linearly dependent in ℓ_1 .

and finally

1.4 Proposition. (Minkowski's Inequality) If $x, y \in \ell_p$, then $\|x + y\|_p \leq \|x\|_p + \|y\|_p$ with equality exactly when one of x or y is a non-negative scalar combination of the other.

REVIEW OF TOPOLOGY

Let X denote a non-empty set, and $\mathcal{P}(X)$ denote the power set of X .

Definition. A **topology** on a set X is a set τ of subsets of X such that

- (i) $\emptyset, X \in \tau$
- (ii) If $U_\alpha \in \tau$ for all $\alpha \in A$, then $\bigcup_{\alpha \in A} U_\alpha \in \tau$.
- (iii) If $n \in \mathbb{N}$ and $U_i \in \tau$ for each $1 \leq i \leq n$, then $\bigcap_{i=1}^n U_i \in \tau$.

The sets $U \in \tau$ are called the **open sets** in X , and sets of the form $X \setminus U$ for some open set U are called the **closed sets** in X . The pair (X, τ) is called a **topological space**.

The metric topology on a metric space (X, d) is the topology

$$\tau_d = \{ U \subseteq X \mid \text{for each } x_0 \in U, \text{ there is } \delta = \delta(x_0) \text{ s.t. } B_\delta(x_0) \subseteq U \}$$

Example. (i) Given two metrics d, ρ on X , we say that $d \sim \rho$ if and only if there are $c, C > 0$ such that

$$cd(x, y) \leq \rho(x, y) \leq Cd(x, y) \text{ for any } x, y \in X$$

Note that $d \sim \rho$ implies that $\tau_d = \tau_\rho$, but the reverse implication is not true. An example of this are the metrics on $X = \mathbb{R}$ given by $d(x, y)$ and $\rho(x, y) = \frac{|x-y|}{1+|x-y|}$. Then $d \sim \rho$ but $\tau_d = \tau_\rho$.

(ii) "Sorgenfrey line" Set $X = \mathbb{R}$, and consider

$$\sigma = \{ V \subseteq \mathbb{R} \mid \text{for any } s \in V, \text{ there is } \delta = \delta(s) > 0 \text{ s.t. } [s, s + \delta) \subseteq V \}$$

It is an exercise to verify that $\tau_{|\cdot|} \subsetneq \sigma$. We say that σ is **finer** than $\tau_{|\cdot|}$.

(iii) Relative topology: let (X, τ) be a topological space, and $\emptyset \neq A \subseteq X$. Then we can define a topology $\tau|_A = \{U \cap A : U \in \tau\}$.

Definition. Let (X, τ) and (Y, σ) be topological spaces, and $f : X \rightarrow Y$. We say that f is $(\tau - \sigma)$ -**continuous** at x_0 in X if,

- given $V \in \sigma$ such that $f(x_0) \in V$, then there exists $U \in \tau$ such that $x_0 \in U$ and $f(U) \subseteq V$.

We say that f is $(\tau - \sigma)$ -continuous if it is continuous at each x_0 in X .

SPACE OF BOUNDED CONTINUOUS FUNCTIONS INTO A NORMED SPACE

Let $(Y, \|\cdot\|)$ denote a normed space. We let $\tau_{\|\cdot\|}$ denote the topology given by the metric $\rho(x, y) = \|x - y\|$. Let (X, τ) denote any topological space. Then we write

$$C_b^Y(X) = \left\{ f : X \rightarrow Y \mid f \text{ is bounded and } \tau - \tau_{\|\cdot\|} \text{ - continuous} \right\}$$

With pointwise operations, we see that $C_b^Y(X)$ is a vector space. We also define for $f \in C_b^Y(X)$, $\|f\|_\infty = \sup\{\|f(x)\| : x \in X\}$, making $(C_b^Y(X), \|\cdot\|_\infty)$ a normed vector space.

1.5 Theorem. *If $(Y, \|\cdot\|)$ is a Banach space, then $(C_b^Y(X), \|\cdot\|_\infty)$ is a Banach space.*

PROOF Let $(f_n)_{n=1}^\infty$ be a Cauchy sequence in $(C_b^Y(X), \|\cdot\|_\infty)$. Then for any $x \in X$, we have that $(f_n(x))_{n=1}^\infty$ is Cauchy in $(Y, \|\cdot\|)$ since $\|f_n(x) - f_m(x)\| \leq \|f_n - f_m\|_\infty$, and hence admits a limit $f(x)$. In particular, $x \mapsto f(x)$ defines a function from X to Y . We shall fix $x_0 \in X$ and show that f is continuous at x_0 . Given $\epsilon > 0$, we let

- n_1 be so $n, m \geq n_1$ so that $\|f_n - f_m\|_\infty < \epsilon/4$.
- n_2 be so $n \geq n_2$ so that $\|f_n(x_0) - f(x_0)\| < \epsilon/4$.
- $N = \max\{n_1, n_2\}$.
- $U \in \tau, x_0 \in U$ such that $f_N(U) \subseteq B_{\epsilon/4}(f(x_0)) \subset Y$.

Then for $x \in U$, we let n_x be so $n_x \geq n_1$ and $n \geq n_x$, so that $\|f_{n_x}(x) - f(x)\| < \epsilon/4$. We then have

$$\begin{aligned} \|f(x) - f(x_0)\| &\leq \|f(x) - f_{n_x}(x)\| + \|f_{n_x}(x) - f_N(x)\| + \|f_N(x) - f_N(x_0)\| + \|f_N(x_0) - f(x_0)\| \\ &< \frac{\epsilon}{4} + \|f_{n_x} - f_N\|_\infty + \frac{\epsilon}{4} + \frac{\epsilon}{4} < \epsilon \end{aligned}$$

in other words that $f(U) \subseteq B_\epsilon(f(x_0))$.

Now let us check that $\|f\|_\infty < \infty$. Since $|\|f_n\|_\infty - \|f_m\|_\infty| \leq \|f_n - f_m\|_\infty$, so $(\|f_n\|_\infty)_{n=1}^\infty \subseteq \mathbb{R}$ is Cauchy, hence bounded. If $x \in X$, then

$$\|f(x)\| = \lim_{n \rightarrow \infty} \|f_n(x)\| \leq \sup_{n \in \mathbb{N}} \|f_n(x)\| \leq \sup_{n \in \mathbb{N}} \|f_n\|_\infty < \infty$$

so $\|f\|_\infty = \sup_{x \in X} \|f(x)\| < \infty$.

Notice that if ϵ, n_1 are as above, and further x_0, N are as above, we have for $n \geq n_1$

$$\|f_n(x_0) - f(x_0)\| \leq \|f_n(x_0) - f_N(x_0)\| + \|f_N(x_0) - f(x_0)\| < \frac{\epsilon}{2}$$

so $\|f_n - f\|_\infty = \sup_{x_0 \in X} \|f_n(x_0) - f(x_0)\| \leq \epsilon/2 < \epsilon$. This is uniform since n_1 is chosen uniformly in X . ■

1.6 Corollary. $(C_b^\mathbb{F}(X), \|\cdot\|_\infty)$ is a Banach space.

Let's first note the following general principle: let $(X, d), (Y, \rho)$ be metric spaces, where (X, d) is complete. If $\psi : X \rightarrow Y$ is a $(d - \rho)$ -isometry, then $(\psi(X), \rho|_{\psi(X)})$ is a complete metric space.

Example. (i) Let T be a non-empty set and let

$$\ell_\infty(T) = \left\{ x = (x_t)_{t \in T} \in \mathbb{F}^T \mid \|x\|_\infty < \infty \right\}$$

With pointwise operations, $(\ell_\infty, \|\cdot\|_\infty)$ is a normed space. In fact, it is a Banach space. Let us note that

$$f \mapsto (f(t))_{t \in T} : C_b(T, \mathcal{P}(T)) \rightarrow \ell_\infty(T)$$

is a surjective linear isometry, and the result follows.

- (ii) Let $c = \{x \in \ell_\infty \mid \lim_{n \rightarrow \infty} x_n \text{ exists}\}$. Then $(c, \|\cdot\|_\infty)$ is a Banach space. Consider the topological space given by $\omega = \mathbb{N} \cup \{\infty\}$, with topology

$$\tau_\omega = \mathcal{P}(\mathbb{N}) \cup \bigcup_{n \in \mathbb{N}} \{k \in \mathbb{N} : k \geq n\}$$

The map $f \mapsto (f(n))_{n=1}^\infty : C_b(\omega) \rightarrow c$ is a linear surjective isometry.

- (iii) $c_0 = \{x \in \mathbb{F}^\mathbb{N} \mid \lim_{n \rightarrow \infty} x_n = 0\} \subseteq c \subseteq \ell_\infty$.

1.7 Lemma. *If $x_0 \in X$ where (X, τ) is a topological space, then*

$$\mathcal{I}(x_0) = \{f \in C_b(X) \mid f(x_0) = 0\}$$

is closed, hence complete, subspace of $C_b(X)$.

PROOF If $(f_n)_{n=1}^\infty \subseteq \mathcal{I}(x_0)$ and $f = \lim_{n \rightarrow \infty} f_n$ with respect to $\|\cdot\|_\infty$ in $C_b(X)$, then $f(x_0) = \lim_{n \rightarrow \infty} f_n(x_0) = 0$. Thus $f \in \mathcal{I}(x_0)$, and closed subsets of complete spaces are themselves complete. ■

Now, $f \mapsto (f(n))_{n=1}^\infty : \mathcal{I}(\infty) \rightarrow c_0$ is a (linear) surjective isometry.

- (iv) Consider the Sorgenfity line (\mathbb{R}, σ) : verify that

$$c_b(\mathbb{R}, \sigma) = \left\{ f : \mathbb{R} \rightarrow \mathbb{F} \mid f \text{ is bounded and } \lim_{t \rightarrow t_0^+} f(t) = f(t_0) \text{ for } t \in \mathbb{R} \right\}$$

2 LINEAR OPERATORS AND LINEAR FUNCTIONALS

Let X, Y be vector spaces. We let $\mathcal{L}(X, Y) = \{S : X \rightarrow Y \mid S \text{ is linear}\}$; this is itself a vector space with pointwise operations. Let $(X, \|\cdot\|)$ be a normed space. We denote

$$D(X) = \{x \in X : \|x\| < 1\}$$

$$S(X) = \{x \in X : \|x\| = 1\}$$

$$B(X) = \{x \in X : \|x\| \leq 1\}$$

(Yes, this notation is confusion. No, I didn't choose it.)

2.1 Proposition. *If X, Y are normed spaces and $S \in \mathcal{L}(X, Y)$, then the following are equivalent:*

- (i) S is continuous
- (ii) S is continuous at some $x_0 \in X$
- (iii) $\|S\| = \sup_{x \in D(X)} \|Sx\| < \infty$.

Moreover, in this case, we have

$$\begin{aligned} \|S\| &= \min\{L > 0 : \|Sx\| \leq L\|x\| \text{ for } x \in X\} \\ &= \sup_{x \in S(X)} \|Sx\| = \sup_{x \in B(X)} \|Sx\| \end{aligned}$$

PROOF ($i \Rightarrow ii$) Obvious

($ii \Rightarrow iii$) Note that

$$Sx_0 + D(Y) = \{Sx_0 + y : y \in D(Y)\} = \{y \in Y : \|Sx_0 - y'\| < 1\}$$

is a neighbourhood of Sx_0 . By the definition of metric continuity, there is $\delta > 0$ such that

$$x_0 + \delta D(X) = \{x_0 + \delta x : x \in D(X)\} = \{x' \in X : \|x_0 - x'\| < \delta\}$$

such that

$$Sx_0 + \delta S(D(X)) = S(x_0 + \delta D(X)) \subseteq Sx_0 + D(Y)$$

which implies that $\delta S(D(X)) \subseteq D(Y)$ and $S(D(X)) \subseteq D(Y)/\delta$, in other words that $\|Sx\| \leq 1/\delta$ for $x \in D(X)$.

($iii \Rightarrow i$) If $x \in X$ and $\epsilon > 0$, then

$$\|Sx\| = (\|x\| + \epsilon) \left\| S \left(\frac{1}{\|x\| + \epsilon} \|x\| \right) \right\| \leq (\|x\| + \epsilon) \|S\|$$

Then, letting $\epsilon \rightarrow 0^+$, we see that

$$\|Sx\| \leq \|x\| \|S\| = \|S\| \|x\|$$

If $x, x' \in X$, then $\|Sx - Sx'\| \leq \|S\| \|x - x'\|$ is S is Lipschitz, hence continuous.

To complete the proof, the content of (iii) implies (i) tell us that the Lipschitz constant $L(S) \leq \|S\|$. Furthermore, if $\|x\| = 1$, the preceding proof gives us that $\|S\|_{S(X)}$.

Conversely,

$$\|S\| = \sup_{x \in D(X) \setminus \{0\}} \|Sx\| = \sup_{x \in D(X) \setminus \{0\}} \|x\| \left\| S \left(\frac{1}{\|x\|} x \right) \right\| \leq \sup_{x \in S(X)} \|Sx\|$$

The remaining equivalence is obvious. ■

We now let $\mathcal{B}(X, Y) = \{S \in \mathcal{L}(X, Y) \mid S \text{ is bounded}\}$. We will see that $\|\cdot\|$, above, defines a norm on $\mathcal{B}(X, Y)$.

2.2 Theorem. *If X, Y are normed spaces, then $(\mathcal{B}(X, Y), \|\cdot\|)$ is a normed space. Furthermore, if Y is a Banach spaces, then so to is $(\mathcal{B}(X, Y), \|\cdot\|)$.*

PROOF Define

$$\Gamma : \mathcal{B}(X, Y) \rightarrow C_b^Y(B(X))$$

given by $\Gamma(S) = S|_{B(X)}$. Then, by definition, Γ is linear, with

$$\|\Gamma(S)\|_\infty = \sup_{x \in B(X)} \|Sx\| = \|S\|$$

Thus $\|\cdot\|$ is a norm: if $S, T \in \mathcal{B}(X, Y)$, $\alpha \in \mathbb{F}$,

$$\begin{aligned} \|S + T\| &= \|\Gamma(S + T)\|_\infty = \|\Gamma(S) + \Gamma(T)\|_\infty \leq \|\Gamma(S)\|_\infty + \|\Gamma(T)\|_\infty = \|S\| + \|T\| \\ \|\alpha S\| &= \|\Gamma(\alpha S)\|_\infty = |\alpha| \|\Gamma(S)\|_\infty = |\alpha| \|S\|. \end{aligned}$$

Furthermore, $\Gamma : \mathcal{B}(X, Y) \rightarrow C_b^Y(\mathcal{B}(X))$ is an isometry.

Now suppose that Y is a Banach space. We will show that $\Gamma(\mathcal{B}(X, Y))$ is closed in $C_b^Y(\mathcal{B}(X))$, and hence $\mathcal{B}(X, Y) = \Gamma^{-1}(\Gamma(\mathcal{B}(X, Y)))$ is complete. Let $(S_n)_{n=1}^\infty \subset \mathcal{B}(X, Y)$ be $\|\cdot\|$ -Cauchy. Then $(\Gamma(S_n))_{n=1}^\infty$ is $\|\cdot\|_\infty$ -Cauchy in $C_b^Y(\mathcal{B}(X))$, and hence there is $f \in C_b^Y(\mathcal{B}(X))$ such that $\lim_{n \rightarrow \infty} \|\Gamma(S_n) - f\|_\infty = 0$. Then we let $S : X \rightarrow Y$ be given by

$$Sx = \begin{cases} \|x\| f\left(\frac{x}{\|x\|}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$$

If $x, x' \in X$ and $\alpha \in \mathbb{F}$ are all such that $x, x', x + \alpha x' \neq 0$, then

$$\begin{aligned} S(x + \alpha x') &= \|x + \alpha x'\| f\left(\frac{1}{x + \alpha x'}(x + \alpha x')\right) \\ &= \|x + \alpha x'\| \lim_{n \rightarrow \infty} S_n\left(\frac{1}{x + \alpha x'}(x + \alpha x')\right) \\ &= \lim_{n \rightarrow \infty} (S_n x + \alpha S_n x') = \lim_{n \rightarrow \infty} \left[\|x\| S_n\left(\frac{1}{\|x\|}x\right) + \alpha \|x'\| S_n\left(\frac{1}{\|x'\|}x'\right) \right] \\ &= \|x\| f\left(\frac{x}{\|x\|}\right) + \alpha \|x'\| f\left(\frac{x'}{\|x'\|}\right) \\ &= Sx + \alpha Sx' \end{aligned}$$

The above computation is easily performed if any of $x, x', x + \alpha x'$ are 0. Hence $S \in \mathcal{L}(X, Y)$. We see that S is continuous (say, at a point on $S(X)$), so $S \in \mathcal{B}(X, Y)$. Finally, as $S|_{\mathcal{B}(X)} = f = \lim_{n \rightarrow \infty} S_n|_{\mathcal{B}(X)}$ (with respect to the uniform norm), we have

$$\|S - S_n\| = \sup_{x \in \mathcal{B}(X)} \|(S - S_n)x\| = \|f - \Gamma(S_n)\|_\infty$$

goes to 0 as n goes to infinity. ■

Definition. Given a vector space X , let $X' = \mathcal{L}(X, \mathbb{F})$ denote the **algebraic dual**. If further X is a normed space, we let $X^* = \mathcal{B}(X, \mathbb{F})$ denote the (continuous) dual.

2.3 Corollary. If X is a normed spaces, then X^* is always a Banach space.

2.4 Theorem. Let for $x \in \ell_1$, $f_x : c_0 \rightarrow \mathbb{F}$ be given by $f_x(y) = \sum_{j=1}^\infty x_j y_j$. Then $f_x \in c_0^*$ with $\|f_x\| = \|x\|_1$. Furthermore, every element of c_0^* arises as above.

PROOF If $x \in \ell_1$ and $y \in c_0 \subseteq \ell_\infty$, then

$$\sum_{j=1}^\infty |x_j y_j| \leq \sum_{j=1}^\infty |x_j| \|y\|_\infty = \|x\|_1 \|y\|_\infty < \infty$$

so $f_x(y) = \sum_{j=1}^\infty x_j y_j$ is well-defined. It is obvious that f_x is linear: $f_x(y + \alpha y') = f_x(y) + \alpha f_x(y')$ for $y, y' \in c_0$ and $\alpha \in \mathbb{F}$. Also, $\|f_x\| \leq \|x\|_1$. We let $y^n = (\overline{\text{sgn } x_1}, \dots, \overline{\text{sgn } x_n}, 0, 0, \dots) \in c_0$, with $\|y^n\| = 1$. Then

$$\|f_x\| \geq |f_x(y^n)| = \sum_{j=1}^n x_j \overline{\text{sgn } x_j} = \sum_{j=1}^n |x_j|$$

so that $\|f_x\| \geq \|x\|_1$, and hence equality holds.

Now let $f \in c_0^*$, and write $e_n = (0, \dots, 0, 1, 0, 0, \dots) \in c_0$, and let $x_n = f(e_n)$. Then, let $y \in c_0$ and $y^n = (y_1, \dots, y_n, 0, 0, \dots)$ and we have

$$\|y - y^n\|_\infty = \sup_{j \geq n+1} |y_j|$$

which goes to 0 as n goes to infinity. Then since f is continuous, we have

$$f(y) = \lim_{n \rightarrow \infty} f(y^n) = \lim_{n \rightarrow \infty} \sum_{j=1}^n y_j x_j = \sum_{j=1}^{\infty} x_j y_j = f_x(y)$$

We use sequence $(y^n)_{n=1}^\infty$ as in $y^n \in c_0$, to see that

$$\sum_{j=1}^n |x_j| = |f(y^n)| \leq \|f\| < \infty$$

so $x \in \ell_1$. Thus $f = f_x$, as desired. ■

2.5 Corollary. $\ell_1 \cong c^*$ isometrically isomorphically.

PROOF For $y \in c$, let $L(y) = \lim_{n \rightarrow \infty} y_n$. Given $y \in c$, let $y^n = (y_1, \dots, y_n, L(y), L(y), \dots) \in c$. Notice that $\|y - y^n\|_\infty \rightarrow 0$ similarly as above.

We let $1 = (1, 1, \dots)$, and $1_n = (0, \dots, 0, 1, 1, \dots)$. If $m < n$, then $1_n - 1_m \in c_0$, so

$$|f(1_n) - f(1_m)| = |f_x(1_n - 1_m)| \leq \sum_{j=m+1}^n |x_j|$$

so that $(f(1_n))_{n=1}^\infty$ is Cauchy in \mathbb{F} . Let $x_0 = \lim_{n \rightarrow \infty} f(1_n)$. Let $\tilde{x} = (x_0, x_1, \dots) \in \ell_1$. Then letting $x_j = f(e_j)$, we see that

$$f(y) = \lim_{n \rightarrow \infty} f(y^n) = \sum_{j=1}^{\infty} x_j y_j + x_0 L(y)$$
■

Similarly as above, we may show that $\|f\| = \|\tilde{x}\|_1$.

Remark. We write $c_0^* \cong \ell_1$ isometrically.

2.6 Corollary. $(\ell_1, \|\cdot\|_1)$ is complete.

3 AXIOM OF CHOICE AND THE HAHN-BANACH THEOREM

Definition. Let S be a non-empty set. A **partial ordering** is a binary relation \leq on S which satisfies for $s, t, n \in S$,

- (i) (*reflexivity*) $s \leq s$
- (ii) (*transitivity*) $s \leq t, t \leq u$ implies $s \leq u$
- (iii) (*anti-symmetry*) $s \leq t, t \leq s$ implies $s = t$

We call the pair (S, \leq) a **partially ordered set**. We say that (S, \leq) is **totally ordered** if, given $s, t \in S$, at least one of $s \leq t$ or $t \leq s$ holds. We say that (S, \leq) is **well-ordered** if given any $\emptyset \neq S_0 \subseteq S$, there is some $s_0 \in S_0$ such that $s_0 \leq s$ for $s \in S_0$. A **chain** in a poset (S, \leq) is any $\emptyset \neq C \subseteq S$ such that $(S, \leq|_C)$ is totally ordered.

Example. (i) $X \neq \emptyset$, $(\mathcal{P}(X), \subseteq)$ is a poset
 (ii) (\mathbb{R}, \leq) is a totally ordered set
 (iii) (\mathbb{N}, \leq) , $(\omega = \mathbb{N} \cup \{\infty\}, \leq)$, are well-ordered sets.

3.1 Theorem. *The following are equivalent:*

- (i) (Axiom of Choice 1): For any $x \neq \emptyset$, there is a function $\gamma : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X$ such that $\gamma(A) \in A$ for each $A \in \mathcal{P}(X) \setminus \{\emptyset\}$.
- (ii) (Axiom of Choice 2): Given any $\{A_\lambda\}_{\lambda \in \Lambda}$ where $A_\lambda \neq \emptyset$ for each λ ,

$$\prod_{\lambda \in \Lambda} A_\lambda = \{(a_\lambda)_{\lambda \in \Lambda} : a_\lambda \in A_\lambda \text{ for each } \lambda\} \neq \emptyset$$

- (iii) (Zorn's Lemma): In a poset (S, \leq) , if each chain $C \subseteq S$ admits an upper bound in S , then (S, \leq) admits a maximal element.
- (iv) (Well-ordering principle): Any $S \neq \emptyset$ admits a well-ordering

PROOF Exercise. ■

Definition. Let X be a vector space (over k). A subset $S \subseteq X$ is called

- **linearly independent** if for any distinct $x_1, \dots, x_n \in S$, the equation $0 = \alpha_1 x_1 + \dots + \alpha_n x_n = 0$ where $\alpha_i \in k$ implies $\alpha_1 = \dots = \alpha_n = 0$.
- **spanning** if each $x \in X$ admits $x_i \in S$ and $\alpha_i \in k$ such that $x = \alpha_1 x_1 + \dots + \alpha_n x_n$.
- **Hamel basis** if it is both linearly independent and spanning

3.2 Proposition. *Any vector space X admits a Hamel basis.*

PROOF Let $\mathcal{L} = \{L \subseteq X : L \text{ is linearly independent}\}$. Then (\mathcal{L}, \subseteq) is a poset. Verify that for any chain $\mathcal{C} \subseteq \mathcal{L}$, that $U = \bigcup_{L \in \mathcal{C}} L \in \mathcal{L}$ and is an upper bound for \mathcal{C} . Apply Zorn to find a maximal element M in (\mathcal{L}, \subseteq) . Verify that M is spanning for X . ■

3.3 Corollary. *If X is an infinite dimensional normed space, then there exists $f \in X' \setminus X^*$.*

PROOF Our assumption provides $\{e_n\}_{n=1}^\infty$ which is linearly independent. By normalizing each element, we may and will suppose that each $\|e_n\| = 1$. Let

$$\text{span}\{e_n\}_{n=1}^\infty = \left\{ \sum_{j=1}^m \alpha_j e_{n_j} : m \in \mathbb{N}, \alpha_j \in \mathbb{F}, n_1 < \dots < n_m \right\}$$

and let B be any linearly independent set containing $\{e_n\}_{n=1}^\infty$. Define $f : X = \text{span } B \rightarrow \mathbb{F}$ be given for $x = \sum_{b \in B \setminus \{e_n\}_{n=1}^\infty} \alpha_b b + \sum_{j=1}^n \alpha_j e_{n_j}$ by $f(x) = \sum_{j=1}^n \alpha_j n_j$. The point is that $f(e_n) = n$ and $f(e) = 0$ for any other $e \in B$. Notice that

$$\|f\| = \sup_{x \in B(X)} |f(x)| \geq \sup_{n \in \mathbb{N}} |f(e_n)| = \sup_{n \in \mathbb{N}} n = \infty$$

■

Definition. Let X be a \mathbb{R} -vector space. A **sublinear functional** is any $\rho : X \rightarrow \mathbb{R}$ such that it satisfies

- (non-negative homogeneity) $\rho(tx) = t\rho(x)$ for $t \geq 0, x \in X$.
- (subadditivity) $\rho(x+y) \leq \rho(x) + \rho(y)$ for $x, y \in X$.

3.4 Theorem. (Hahn-Banach) Let X be a \mathbb{R} -vector space, $\rho : X \rightarrow \mathbb{R}$ a sublinear functional, $Y \subseteq X$ a subspace and $f \in Y'$ such that $f \leq \rho|_Y$. Then there exists $F \in X'$ such that $F|_Y = f$ and $F \leq \rho$ on X .

PROOF Given $x \in X \setminus Y$, we wish to find $c \in \mathbb{R}$ such that

$$f(y) + \alpha c \leq \rho(y + \alpha x)$$

for $y \in Y$ and $\alpha \in \mathbb{R}$. In this case, we let $F : \text{span } Y \cup \{x\} \rightarrow \mathbb{R}$ be given by $F(y + \alpha x) = f(y) + \alpha c$, and we have that F is linear and satisfies $F \leq \rho$ on $\text{span } Y \cup \{x\}$. To do this, let y_+, y_- in Y and observe that $f(y_+) + f(y_-) = f(y_+ + y_-) \leq \rho(y_+ + y_-) \leq \rho(y_+ + x) + \rho(y_- - x)$ so that $f(y_-) - \rho(y_- - x) \leq \rho(y_+ + x) - f(y_+)$. It thus follows that

$$\sup\{f(y) - \rho(y - x) : y \in Y\} \leq \inf\{\rho(y + x) - f(y) : y \in Y\}$$

so we may find $c \in \mathbb{R}$ for which

$$\sup\{f(y) - \rho(y - x) : y \in Y\} \leq c \leq \inf\{\rho(y + x) - f(y) : y \in Y\}$$

If $t > 0$, then for $y \in Y$,

$$c \leq \rho\left(\frac{1}{t}y + x\right) - f\left(\frac{1}{t}y\right) \Rightarrow tc \leq \rho(y + tx) - f(y) \Rightarrow f(y) + tc \leq \rho(y + tx)$$

and if $s > 0$, then for $y \in Y$,

$$f\left(\frac{1}{s}y\right) - \rho\left(\frac{1}{s}y - x\right) \leq c \Rightarrow sc \leq f(y) - \rho(y + sx) \Rightarrow f(y) - sc \leq \rho(y + sx)$$

Clearly, $f(y) + 0 \leq \rho(y + 0x)$. Hence, we have our desired inequality.

To be continued...

■