

# Boundary Reduction of Helmholtz Scattering Problem for exterior domains

Paul Escapil-Inchauspé

June 21, 2017

Let  $D$  be an open Lipschitz bounded domain,  $D^c := \mathbb{R}^d \setminus \overline{D}$  and  $\Gamma := \partial D$ ,  $d = 2, 3$  with exterior normal field  $\mathbf{n}$  pointing by convention toward the exterior domain. We use the superscript  $i = 1$  (resp.  $i = 2$ ) when referring to  $D^c$  (resp.  $D$ ). For a given scalar field  $U$ , we introduce the Dirichlet (resp. Neumann) traces on  $\Gamma$  as:

$$\gamma_0 U^i := U|_\Gamma \text{ and } \gamma_1 U := \langle \mathbf{n}, \nabla U|_\Gamma \rangle. \quad (1)$$

For a given  $\kappa$ , we define the Helmholtz equation operator  $L_\kappa : U \mapsto \Delta U + \kappa^2 U$ . For an index  $\beta \in \{0, 1\}$  us consider the following exterior boundary value problems  $(\mathbf{EP}_\beta)$ , which consist in finding the scattered field induced in response to a given incident field:

## Exterior Problems

Given  $\kappa > 0$  and  $U^{inc} \in H_{loc}^1(D^c)$  with  $L_\kappa U^{inc} = 0$  in  $D^c$ , find  $U := (U^{scat} + U^{inc}) \in H_{loc}^1(D^c)$  such that

$$\begin{cases} \Delta U + \kappa^2 U = 0 & \text{in } D^c, \\ \gamma_\beta U = 0 & \text{on } \Gamma, \\ \left| \frac{\partial}{\partial r} U^{scat} - i\kappa U^{scat} \right| = o(r^{\frac{1-d}{2}}) & \text{for } r \rightarrow \infty. \end{cases} \quad (2)$$

Let us introduce the boundary integral operators  $\mathcal{V}_\kappa, \mathcal{K}_\kappa, \mathcal{W}_\kappa, \mathcal{K}'_\kappa, \mathcal{I}$  defined on  $\Gamma$ . Then, we have the following propositions:

**Proposition 1.** *The sound-soft problem  $(\mathbf{EP}_0)$  is equivalent to each of the following BIE given on  $\Gamma$ :*

$$\begin{cases} \mathcal{V}_\kappa \gamma_1 U &= \gamma_0 U^{inc} \text{ and} \\ \mathcal{V}_\kappa \gamma_1 U^{scat} &= -(\mathcal{K}_\kappa - \frac{1}{2} \mathcal{I}) \gamma_0 U^{inc}. \end{cases} \quad (3)$$

**Proposition 2.** *The sound-hard problem  $(\mathbf{EP}_1)$  is equivalent to each of the following BIE given on  $\Gamma$ :*

$$\begin{cases} \mathcal{W}_\kappa \gamma_0 U &= \gamma_1 U^{inc} \text{ and} \\ \mathcal{W}_\kappa \gamma_0 U^{scat} &= (\mathcal{K}'_\kappa + \frac{1}{2} \mathcal{I}) \gamma_1 U^{inc}. \end{cases} \quad (4)$$