

Assignment 8: GLMs (Linear Regressios, ANOVA, & t-tests)

Haonan Pei

Spring 2026

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on generalized linear models.

Directions

1. Rename this file `<FirstLast>_A08_GLMs.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

!! AI generated code should NOT be used in this assignment

NOTE: For this assignment, you can skip tests for any of the assumptions required to run the models.

Set up your session

1. Set up your session. Check your working directory. Load the tidyverse, agricolae and other needed packages. Import the *raw* NTL-LTER raw data file for chemistry/physics (NTL-LTER_Lake_ChemistryPhysics_Raw.csv). Set date columns to date objects.
2. Build a ggplot theme and set it as your default theme.

```
#1
setwd(dir = "/home/guest/EDE_Spring2026")
getwd()
```

```
## [1] "/home/guest/EDE_Spring2026"
```

```
library(tidyverse)
```

```
## -- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
## v dplyr      1.1.4      v readr      2.1.5
## v forcats    1.0.0      v stringr    1.5.1
## v ggplot2    3.5.1      v tibble     3.2.1
## v lubridate  1.9.3      v tidyr      1.3.1
```

```
## v purrr      1.0.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()   masks stats::lag()
## i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become errors

#install.packages("agricolae")
library(agricolae)
NTL_data <- read_csv("Data/Raw/NTL-LTER_Lake_ChemistryPhysics_Raw.csv") %>%
  mutate(sampleddate = as.Date(sampleddate, format = "%m/%d/%Y"))

## Rows: 38614 Columns: 11
## -- Column specification -----
## Delimiter: ","
## chr (4): lakeid, lakename, sampleddate, comments
## dbl (7): year4, daynum, depth, temperature_C, dissolvedOxygen, irradianceWat...
##
## i Use 'spec()' to retrieve the full column specification for this data.
## i Specify the column types or set 'show_col_types = FALSE' to quiet this message.

#2
default_theme <- ggplot(NTL_data)
default_theme <- theme_bw() +
  theme(
    plot.title = element_text(size = 16, face = "bold"),
    axis.title = element_text(size = 14)
  )
theme_set(default_theme)
```

Simple regression

Our first research question is: Does mean lake temperature recorded during July change with depth across all lakes?

3. State the null and alternative hypotheses for this question: > Answer: H0: The mean lake temperature recorded during July does not change with depth across all lakes. Ha: The mean lake temperature recorded during July change with depth across all lakes.
4. Wrangle your NTL-LTER dataset with a pipe function so that the records meet the following criteria:
 - Only dates in July.
 - Only the columns: lakename, year4, daynum, depth, temperature_C
 - Only complete cases (i.e., remove NAs)
5. Visualize the relationship among the two continuous variables with a scatter plot of temperature by depth. Add a smoothed line showing the linear model, and limit temperature values from 0 to 35 °C. Make this plot look pretty and easy to read.

```
#4
NTL_data_filtered <- NTL_data %>%
  filter(month(sampleddate) == 7) %>%
```

```


select(lakename, year4, daynum, depth, temperature_C) %>%
drop_na()

#5
temp_depth_scatter <- ggplot(NTL_data_filtered, aes(x = depth, y = temperature_C)) +
  geom_point(color = "blue") +
  geom_smooth(method = "lm", color = "orange") +
  coord_cartesian(ylim = c(0, 35)) +
  labs(
    title = "Relationship Between Lake Temperature and Depth (July)",
    x = "Depth (m)",
    y = "Temperature (°C)"
  )

print(temp_depth_scatter)

## 'geom_smooth()' using formula = 'y ~ x'

```



6. Interpret the figure. What does it suggest with regards to the response of temperature to depth? Do the distribution of points suggest about anything about the linearity of this trend?

Answer: The figure suggests that lake temperature decreases as depth increases, indicating a negative relationship between temperature and depth. However, the temperature varies more at the shallow depths from 0 to 10 meters, showing that the relationship is not perfectly linear.

7. Perform a linear regression to test the relationship and display the results.

```
#7
temp_depth_lm <- lm(temperature_C ~ depth, data = NTL_data_filtered)
summary(temp_depth_lm)

##
## Call:
## lm(formula = temperature_C ~ depth, data = NTL_data_filtered)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.5173 -3.0192  0.0633  2.9365 13.5834
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  21.95597    0.06792   323.3  <2e-16 ***
## depth       -1.94621    0.01174  -165.8  <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.835 on 9726 degrees of freedom
## Multiple R-squared:  0.7387, Adjusted R-squared:  0.7387
## F-statistic: 2.75e+04 on 1 and 9726 DF, p-value: < 2.2e-16
```

8. Interpret your model results in words. Include how much of the variability in temperature is explained by changes in depth, the degrees of freedom on which this finding is based, and the statistical significance of the result. Also mention how much temperature is predicted to change for every 1m change in depth.

Answer: The linear regression results indicate a strong negative relationship between lake temperature and depth. Based on the adjusted r-squared value, the model explains 73.87% of the variability in temperature, indicating that the depth accounts for a large portion of the variation in temperature. The model has 1 degree of freedom for depth and 9726 degrees of freedom for the residual error. The p-value is $< 2.2e-16$, which suggests that the finding is extremely significant. The slope coefficient is equal to -1.9462 indicates that for every 1 meter increase in depth, lake temperature is predicted to decrease by approximately 1.95 Celsius degree.

Multiple regression

Let's tackle a similar question from a different approach. Here, we want to explore what might the best set of predictors for lake temperature in July across the monitoring period at the North Temperate Lakes LTER.

9. Run an AIC to determine what set of explanatory variables (year4, daynum, depth) is best suited to predict temperature.
10. Run a multiple regression on the recommended set of variables.

```
#9
NTLregression <- lm(temperature_C ~ year4 + daynum + depth, data = NTL_data_filtered)
best_regression <- step(NTLregression, direction = "both")
```

```
## Start: AIC=26065.53
## temperature_C ~ year4 + daynum + depth
##
##           Df Sum of Sq   RSS   AIC
## <none>                 141687 26066
## - year4    1         101 141788 26070
## - daynum   1        1237 142924 26148
## - depth    1       404475 546161 39189
```

```
summary(best_regression)
```

```
##
## Call:
## lm(formula = temperature_C ~ year4 + daynum + depth, data = NTL_data_filtered)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6536 -3.0000  0.0902  2.9658 13.6123
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) -8.575564   8.630715  -0.994  0.32044
## year4         0.011345   0.004299   2.639  0.00833 **
## daynum        0.039780   0.004317   9.215 < 2e-16 ***
## depth        -1.946437   0.011683 -166.611 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.817 on 9724 degrees of freedom
## Multiple R-squared:  0.7412, Adjusted R-squared:  0.7411
## F-statistic: 9283 on 3 and 9724 DF, p-value: < 2.2e-16
```

```
#10
multi_regression <- lm(formula = temperature_C ~ year4 + daynum + depth, data = NTL_data_filtered)
summary(multi_regression)
```

```
##
## Call:
## lm(formula = temperature_C ~ year4 + daynum + depth, data = NTL_data_filtered)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -9.6536 -3.0000  0.0902  2.9658 13.6123
##
## Coefficients:
##              Estimate Std. Error  t value Pr(>|t|)
## (Intercept) -8.575564   8.630715  -0.994  0.32044
## year4         0.011345   0.004299   2.639  0.00833 **
```

```
## daynum      0.039780   0.004317    9.215 < 2e-16 ***
## depth      -1.946437   0.011683  -166.611 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.817 on 9724 degrees of freedom
## Multiple R-squared:  0.7412, Adjusted R-squared:  0.7411
## F-statistic: 9283 on 3 and 9724 DF,  p-value: < 2.2e-16
```

11. What is the final set of explanatory variables that the AIC method suggests we use to predict temperature in our multiple regression? How much of the observed variance does this model explain? Is this an improvement over the model using only depth as the explanatory variable?

Answer: The final set of explanatory variables includes all three predictors: “year4”, “daynum”, and “depth”. The model explains approximately 74.11% of observed variance based on the adjusted r-squared value. Comparing the model using only depth (73.87%) as the explanatory variable, this new models lightly improves the ability to predict lake temperature.

Analysis of Variance

12. Now we want to see whether the different lakes have, on average, different temperatures in the month of July. Run an ANOVA test to complete this analysis. (No need to test assumptions of normality or similar variances.) Create two sets of models: one expressed as an ANOVA models and another expressed as a linear model (as done in our lessons).

```
#12
anova_test <- aov(temperature_C ~ lakename, data = NTL_data_filtered)
summary(anova_test)
```

```
##              Df Sum Sq Mean Sq F value Pr(>F)
## lakename      8  21642   2705.2     50 <2e-16 ***
## Residuals    9719 525813     54.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
lm_model <- lm(temperature_C ~ lakename, data = NTL_data_filtered)
anova(lm_model)
```

```
## Analysis of Variance Table
##
## Response: temperature_C
##              Df Sum Sq Mean Sq F value    Pr(>F)
## lakename      8  21642   2705.2  50.003 < 2.2e-16 ***
## Residuals    9719 525813     54.1
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```


13. Is there a significant difference in mean temperature among the lakes? Report your findings.

Answer: Yes. There is a statistically significant difference in mean temperature among the lakes. The ANOVA results show a significant effect of lakes on temperature (F value = 50, $p < 2.2e-16$). This indicates that the mean July temperature vary significantly across different lakes.

14. Create a graph that depicts temperature by depth, with a separate color for each lake. Add a `geom_smooth` (method = "lm", se = FALSE) for each lake. Make your points 50 % transparent. Adjust your y axis limits to go from 0 to 35 degrees. Clean up your graph to make it pretty.

```
#14.
ggplot(NTL_data_filtered, aes(x = depth, y = temperature_C, color = lakename)) +
  geom_smooth(method = "lm", se = FALSE) +
  geom_point(alpha = 0.5) +
  coord_cartesian(ylim = c(0, 35)) +
  labs(
    title = "Temperature vs Depth by Lake in July",
    x = "Depth (m)",
    y = "Temperature (°C)",
    color = "Lake"
  )
)
```

'geom_smooth()' using formula = 'y ~ x'



15. Use the Tukey's HSD test to determine which lakes have different means.

```
#15
```

```
tukey_test <- TukeyHSD(anova_test)
tukey_test
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = temperature_C ~ lakename, data = NTL_data_filtered)
##
## $lakename
##
```

	diff	lwr	upr	p adj
## Crampton Lake-Central Long Lake	-2.3145195	-4.7031913	0.0741524	0.0661566
## East Long Lake-Central Long Lake	-7.3987410	-9.5449411	-5.2525408	0.0000000
## Hummingbird Lake-Central Long Lake	-6.8931304	-9.8184178	-3.9678430	0.0000000
## Paul Lake-Central Long Lake	-3.8521506	-5.9170942	-1.7872070	0.0000003
## Peter Lake-Central Long Lake	-4.3501458	-6.4115874	-2.2887042	0.0000000
## Tuesday Lake-Central Long Lake	-6.5971805	-8.6971605	-4.4972005	0.0000000
## Ward Lake-Central Long Lake	-3.2077856	-6.1330730	-0.2824982	0.0193405
## West Long Lake-Central Long Lake	-6.0877513	-8.2268550	-3.9486475	0.0000000
## East Long Lake-Crampton Lake	-5.0842215	-6.5591700	-3.6092730	0.0000000
## Hummingbird Lake-Crampton Lake	-4.5786109	-7.0538088	-2.1034131	0.0000004
## Paul Lake-Crampton Lake	-1.5376312	-2.8916215	-0.1836408	0.0127491
## Peter Lake-Crampton Lake	-2.0356263	-3.3842699	-0.6869828	0.0000999
## Tuesday Lake-Crampton Lake	-4.2826611	-5.6895065	-2.8758157	0.0000000
## Ward Lake-Crampton Lake	-0.8932661	-3.3684639	1.5819317	0.9714459
## West Long Lake-Crampton Lake	-3.7732318	-5.2378351	-2.3086285	0.0000000
## Hummingbird Lake-East Long Lake	0.5056106	-1.7364925	2.7477137	0.9988050
## Paul Lake-East Long Lake	3.5465903	2.6900206	4.4031601	0.0000000
## Peter Lake-East Long Lake	3.0485952	2.2005025	3.8966879	0.0000000
## Tuesday Lake-East Long Lake	0.8015604	-0.1363286	1.7394495	0.1657485
## Ward Lake-East Long Lake	4.1909554	1.9488523	6.4330585	0.0000002
## West Long Lake-East Long Lake	1.3109897	0.2885003	2.3334791	0.0022805
## Paul Lake-Hummingbird Lake	3.0409798	0.8765299	5.2054296	0.0004495
## Peter Lake-Hummingbird Lake	2.5429846	0.3818755	4.7040937	0.0080666
## Tuesday Lake-Hummingbird Lake	0.2959499	-1.9019508	2.4938505	0.9999752
## Ward Lake-Hummingbird Lake	3.6853448	0.6889874	6.6817022	0.0043297
## West Long Lake-Hummingbird Lake	0.8053791	-1.4299320	3.0406903	0.9717297
## Peter Lake-Paul Lake	-0.4979952	-1.1120620	0.1160717	0.2241586
## Tuesday Lake-Paul Lake	-2.7450299	-3.4781416	-2.0119182	0.0000000
## Ward Lake-Paul Lake	0.6443651	-1.5200848	2.8088149	0.9916978
## West Long Lake-Paul Lake	-2.2356007	-3.0742314	-1.3969699	0.0000000
## Tuesday Lake-Peter Lake	-2.2470347	-2.9702236	-1.5238458	0.0000000
## Ward Lake-Peter Lake	1.1423602	-1.0187489	3.3034693	0.7827037
## West Long Lake-Peter Lake	-1.7376055	-2.5675759	-0.9076350	0.0000000
## Ward Lake-Tuesday Lake	3.3893950	1.1914943	5.5872956	0.0000609
## West Long Lake-Tuesday Lake	0.5094292	-0.4121051	1.4309636	0.7374387
## West Long Lake-Ward Lake	-2.8799657	-5.1152769	-0.6446546	0.0021080

16. From the findings above, which lakes have the same mean temperature, statistically speaking, as Peter Lake? Does any lake have a mean temperature that is statistically distinct from all the other lakes?

Answer: The Tukey's HSD test indicates that a pair of lakes that has adjusted p_value smaller than 0.05 is significantly distinct from each other. In this way, Paul Lake and Ward Lake

has statistically same mean temperature as Peter lakes (larger adjusted p_value than 0.05). Moreover, there is no a lake have a mean temperature that is statistically distinct from all the other lakes. Each lake at least has one non-significant comparison with another lake.

17. If we were just looking at Peter Lake and Paul Lake. What's another test we might explore to see whether they have distinct mean temperatures?

Answer: In order to comparing only Peter Lake and Paul Lake, it would be appropriate to explore the Two-Sample T-Test to determine whether their mean temperatures are significantly different since this is an approach that is appropriate when comparing the means of two independent groups.

18. Wrangle the July data to include only records for Crampton Lake and Ward Lake. Run the two-sample T-test on these data to determine whether their July temperature are same or different. What does the test say? Are the mean temperatures for the lakes equal? Does that match you answer for part 16?

```
NTL_t_test <- NTL_data_filtered %>%
  filter(lakename %in% c("Crampton Lake", "Ward Lake"))
t_test <- t.test(temperature_C ~ lakename, data = NTL_t_test)
t_test
```

```
##
## Welch Two Sample t-test
##
## data: temperature_C by lakename
## t = 1.1181, df = 200.37, p-value = 0.2649
## alternative hypothesis: true difference in means between group Crampton Lake and group Ward Lake is not equal to 0
## 95 percent confidence interval:
## -0.6821129 2.4686451
## sample estimates:
## mean in group Crampton Lake      mean in group Ward Lake
##           15.35189              14.45862
```

Answer: The test shows that the p-value is greater than 0.05 (0.2649), which means the mean July temperatures for these two lakes are statistically equal to each other. This result is also match my answer for part 16 (adjusted p_value = 0.9714).