

# Thermodynamics in VC210

Haotian Fu

University of Michigan–Shanghai Jiao Tong University Joint Institute

November 15, 2021

# Outline

## 1 Concept Review

- 1<sup>st</sup> Law
- 2<sup>nd</sup> Law
- 3<sup>rd</sup> Law
- 0<sup>th</sup> Law
- Practical Formula

## 2 Examples

# Outline

## 1 Concept Review

- 1<sup>st</sup> Law
- 2<sup>nd</sup> Law
- 3<sup>rd</sup> Law
- 0<sup>th</sup> Law
- Practical Formula

## 2 Examples

# First law of thermodynamics

$$U = Q + W \quad (1)$$

## Remarks:

- Differential form &  $U = f(T)$

# First law of thermodynamics

$$U = Q + W \quad (1)$$

## Remarks:

- Differential form &  $U = f(T)$
- $W$  - work, including expansion work and non-expansion work. However, we only discuss the expansion work here.

# First law of thermodynamics

$$U = Q + W \quad (1)$$

## Remarks:

- Differential form &  $U = f(T)$
- $W$  - work, including expansion work and non-expansion work. However, we only discuss the expansion work here.
- $Q$  - heat, can be modeled as  $Q_V$  and  $Q_p$

# Enthalpy

$$H = U + pV \quad (2)$$

## Remarks:

- Differential form &  $H = f(T)$ .

# Enthalpy

$$H = U + pV \quad (2)$$

## Remarks:

- Differential form &  $H = f(T)$ .
- Pay attention to the common name of enthalpys, such as,  $H_{vap}$ ,  $H_{fus}$ ,  $H_{sub}$ .



# Enthalpy

$$H = U + pV \quad (2)$$

## Remarks:

- Differential form &  $H = f(T)$ .
- Pay attention to the common name of enthalpys, such as,  $H_{vap}$ ,  $H_{fus}$ ,  $H_{sub}$ .
- $pV$  should be fully understood.

# Heat capacity

$$C = \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T} \quad (3)$$

## Remarks:

- Heat capacity & specific heat capacity.

# Heat capacity

$$C = \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T} \quad (3)$$

## Remarks:

- Heat capacity & specific heat capacity.
- $Q$  should be fully understood.

# Heat capacity

$$C = \lim_{\Delta T \rightarrow 0} \frac{\Delta Q}{\Delta T} \quad (3)$$

## Remarks:

- Heat capacity & specific heat capacity.
- $Q$  should be fully understood.
- Relationship with Kinetic Molecular Theory (KMT).

# KMT & Heat capacity<sup>1</sup>

	Atoms	Linear molecules	Nonlinear molecules
$C_{V,m}$	$\frac{3}{2}R$	$\frac{5}{2}R$	$3R$
$C_{P,m}$	$\frac{5}{2}R$	$\frac{7}{2}R$	$4R$

Figure:  $C_{V,m}$  and  $C_{P,m}$  of three types of molecules.

<sup>1</sup>You may refer to *Chemicle Principle* pp. 402

# The second law of thermodynamics

## Clausius Statement

Heat can never pass from a colder to a warmer body without some other change, connected therewith, occurring at the same time.

# The second law of thermodynamics

## Kelvin Statement

It is impossible for a self-acting machine, unaided by any external agency, to convey heat from one body to another at a higher temperature.

It is impossible, by means of inanimate material agency, to derive mechanical effect from any portion of matter by cooling it below the temperature of the coldest of the surrounding objects.

# Entropy

$$dS = \frac{\delta Q_R}{T} \quad (4)$$

## Remarks:

- The intuition of entropy.



# Entropy

$$dS = \frac{\delta Q_R}{T} \quad (4)$$

## Remarks:

- The intuition of entropy.
- Entropy criterion.

# Entropy

$$dS = \frac{\delta Q_R}{T} \quad (4)$$

## Remarks:

- The intuition of entropy.
- Entropy criterion.
- Calculation of entropy.

# Gibb's Free Energy

$$G = H - TS \quad (5)$$

- Differential form.

# Gibb's Free Energy

$$G = H - TS \quad (5)$$

- Differential form.
- Comparing with  $\Delta G = \Delta H - T\Delta S$

# Gibb's Free Energy

$$G = H - TS \quad (5)$$

- Differential form.
- Comparing with  $\Delta G = \Delta H - T\Delta S$
- Gibb's free energy criterion.

# Gibb's Free Energy

$$G = H - TS \quad (5)$$

- Differential form.
- Comparing with  $\Delta G = \Delta H - T\Delta S$
- Gibb's free energy criterion.
- Calculation of Gibb's free energy.

# The third law of thermodynamics

$$S \rightarrow c \quad \text{as} \quad T \rightarrow 0 \text{ K}$$

Especially,  $c = 0$  when the object is **perfect crystal**.

# The zeroth law of thermodynamics

## content

If two thermodynamic systems are each in thermal equilibrium with a third system, then they are in thermal equilibrium with each other.



# Practical Formula

$$Q_V = \Delta U \quad (6)$$

$$Q_p = \Delta H \quad (7)$$

$$C_V = \frac{\Delta U}{\Delta T} \quad (8)$$

$$C_p = \frac{\Delta H}{\Delta T} \quad (9)$$

$$\Delta S_{\text{surrounding}} = -\frac{\Delta H}{T} \quad (10)$$

$$S = k \ln W \quad (11)$$

# Practical Formula

phase change

$$\Delta S = \frac{Q_R}{T} \quad (12)$$

physical change with constant pressure

$$\Delta S = nC_{V,m} \ln \left( \frac{T_2}{T_1} \right) + nR \ln \left( \frac{V_2}{V_1} \right) \quad (13)$$

physical change with constant volume

$$\Delta S = nC_{p,m} \ln \left( \frac{T_2}{T_1} \right) + nR \ln \left( \frac{p_1}{p_2} \right) \quad (14)$$

# Practical Formula

Clausius–Clapeyron relation

$$\ln \left( \frac{p_2}{p_1} \right) = -\frac{\Delta H}{R} \left( \frac{1}{T_2} - \frac{1}{T_1} \right) \quad (15)$$

constant temperature

$$\Delta G = \int V dp \quad \text{since} \quad dG = -s dT + V dp \quad (16)$$

# Outline

## 1 Concept Review

- 1<sup>st</sup> Law
- 2<sup>nd</sup> Law
- 3<sup>rd</sup> Law
- 0<sup>th</sup> Law
- Practical Formula

## 2 Examples

# Bohn-Haber Cycle

The standard molar enthalpy of combustion of ethanol ( $\text{C}_2\text{H}_5\text{OH}$ ) and acetic acid ( $\text{CH}_3\text{COOH}$ ) is  $-136.50$  and  $-67.50$  kJ/mol.

The enthalpy of dissolution is  $11.71$  and  $1.46$  kJ/mol correspondingly. Calculate the enthalpy of oxidation of ethanol in water solution.

# Liquid-Solution

Liquid  $A$  and  $B$  form an ideal solution. The solution containing 1 mol  $A$  and 2 mol  $B$  at 343.2 K has a vapor pressure of 50.663 kPa.

If 3 mol  $A$  is added, the vapor pressure of the solution increases to 70.928 kPa. Calculate (1)  $p_A^*$  and  $p_B^*$  at 343.2 K; (2) the gaseous composition of the first mixed solution.