

UC DAVIS

STA 135

HW 4 SOLUTIONS

THESE SOLUTIONS ARE USED WITH PERMISSION FROM THE PUBLISHER. BY NO MEANS SHOULD THESE SOLUTIONS BE DISTRIBUTED OR SHARED IN ANY FORM. THEY ARE FOR PERSONAL USE ONLY.

6.2 Using a critical value $t_{n-1}(\alpha/2p) = t_{10}(0.0125) = 2.6338$,

	LOWER	UPPER
Bonferroni C. I.:	-20.57	1.85
	-2.97	29.52
Simultaneous C. I.:	-22.45	3.73
	-5.70	32.25

6.6 a) Treatment 2: Sample mean vector $\begin{bmatrix} 2 \\ 4 \end{bmatrix}$; sample covariance matrix $\begin{bmatrix} 1 & -3/2 \\ -3/2 & 3 \end{bmatrix}$

Treatment 3: Sample mean vector $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$; sample covariance matrix $\begin{bmatrix} 2 & -4/3 \\ -4/3 & 4/3 \end{bmatrix}$

$$S_{\text{pooled}} = \begin{bmatrix} 1.6 & -1.4 \\ -1.4 & 2 \end{bmatrix}$$

$$b) \quad T^2 = [2-3, 4-2] \left[\left(\frac{1}{3} + \frac{1}{4} \right) \begin{bmatrix} 1.6 & -1.4 \\ -1.4 & 2 \end{bmatrix} \right]^{-1} \begin{bmatrix} 2-3 \\ 4-2 \end{bmatrix} = 3.88$$

$$\frac{(n_1+n_2-2)p}{(n_1+n_2-p-1)} F_{p, n_1+n_2-p-1}(.01) = \frac{(5)2}{4} (18) = 45$$

Since $T^2 = 3.88 < 45$ do not reject $H_0: \mu_2 - \mu_3 = \underline{0}$ at the $\alpha = .01$ level.

c) 99% simultaneous confidence intervals:

$$\mu_{21} - \mu_{31}: (2-3) \pm \sqrt{45} \sqrt{\left(\frac{1}{3} + \frac{1}{4}\right) 1.6} = -1 \pm 6.5$$

$$\mu_{22} - \mu_{32}: 2 \pm 7.2$$

6.9 For any matrix C

$$\underline{\bar{d}} = \frac{1}{n} \sum \underline{d}_j = C \left(\frac{1}{n} \sum \underline{x}_j \right) = C \underline{\bar{x}}$$

$$\text{and } \underline{d}_j - \underline{\bar{d}} = C(\underline{x}_j - \underline{\bar{x}})$$

$$\text{so } S_d = \frac{1}{n-1} \sum (\underline{d}_j - \underline{\bar{d}})(\underline{d}_j - \underline{\bar{d}})' = C \left(\frac{1}{n-1} \sum (\underline{x}_j - \underline{\bar{x}})(\underline{x}_j - \underline{\bar{x}})' \right) C' = C S_C C'$$

Note: For 6.16, this is one solution for the specific contrast matrix, C , specified. There are many contrast matrices that could have been specified.

$$6.16 \quad H_0: C\mu = 0; H_1: C\mu \neq 0 \quad \text{where} \quad C = \begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

Summary statistics:

$$\bar{x} = \begin{bmatrix} 1906.1 \\ 1749.5 \\ 1509.1 \\ 1725.0 \end{bmatrix}; \quad S = \begin{bmatrix} 105625 & 94759 & 87249 & 94268 \\ & 101761 & 76166 & 81193 \\ & & 91809 & 90333 \\ & & & 104329 \end{bmatrix}$$

$$T^2 = n(\bar{C}\bar{x})'(CSC')^{-1}(\bar{C}\bar{x}) = 254.7$$

$$\frac{(n-1)(q-1)}{(n-q+1)} F_{q-1, n-q+1}(\alpha) = \frac{(30-1)(4-1)}{(30-4+1)} F_{3,27}(.05) = 9.54$$

Since $T^2 = 254.7 > 9.54$ we reject H_0 at $\alpha = .05$ level.


95% simultaneous confidence interval for "dynamic" versus "static" means $(\mu_1 + \mu_2) - (\mu_3 + \mu_4)$ is, with $c' = [1 \quad 1 \quad -1 \quad -1]$,

$$c'\bar{x} \pm \sqrt{\frac{(n-1)(q-1)}{(n-q+1)} F_{q-1, n-q+1}(\alpha)} \sqrt{\frac{c'Sc}{n}}$$

Copyright © 2012 Pearson Education, Inc. Publishing as Prentice Hall

$$= 421.5 \pm 174.5 \rightarrow (247.596)$$

Note: the QQ diagnostic plots on 6.18 are for reference and were not required to answer the question.



6.18 continued

mean vector for females:

X1BAR
4.9006593
4.6229089
3.9402858

mean vector for males:

X2BAR
4.7254436
4.4775738
3.7031858

SPOOLED 0.0187388 0.0140655 0.0165386
0.0140655 0.0113036 0.0127148
0.0165386 0.0127148 0.0158563

TSQ	CVTSQ	F	CVF	PVALUE
85.052001	8.833461	27.118029	2.8164658	4.355E-10

linear combination most responsible for rejection

of H_0 has coefficient vector:

COEFFVEC
-43.72677
-8.710687
67.546415

95% simultaneous CI for the difference

in female and male means

LOWER	UPPER
0.0577676	0.2926638
0.0541167	0.2365537
0.1290622	0.3451377

Bonferroni CI

LOWER	UPPER
0.0768599	0.2735714
0.0689451	0.2217252
0.1466248	0.3275751

6.19

$$a) \quad \bar{\underline{x}}_1 = \begin{bmatrix} 12.219 \\ 8.113 \\ 9.590 \end{bmatrix}; \quad \bar{\underline{x}}_2 = \begin{bmatrix} 10.106 \\ 10.762 \\ 18.168 \end{bmatrix};$$

$$S_1 = \begin{bmatrix} 223.0134 & 12.3664 & 2.9066 \\ & 17.5441 & 4.7731 \\ & & 13.9633 \end{bmatrix}$$

$$S_2 = \begin{bmatrix} 4.3623 & .7599 & 2.3621 \\ & 25.8512 & 7.6857 \\ & & 46.6543 \end{bmatrix};$$

$$S_{\text{pooled}} = \begin{bmatrix} 15.8112 & 7.8550 & 2.6959 \\ & 20.7458 & 5.8960 \\ & & 26.5750 \end{bmatrix}$$

$$\left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) S_{\text{pooled}} \right]^{-1} = \begin{bmatrix} 1.0939 & -.4084 & -.0203 \\ & .8745 & -.1525 \\ & & .5640 \end{bmatrix}$$

$$H_0: \underline{\mu}_1 - \underline{\mu}_2 = \underline{0}$$

$$\text{Since } T^2 = (\bar{\underline{x}}_1 - \bar{\underline{x}}_2)' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) S_{\text{pooled}} \right]^{-1} (\bar{\underline{x}}_1 - \bar{\underline{x}}_2) = 50.92$$

$$> \frac{(n_1 + n_2 - 2)p}{(n_1 + n_2 - p - 1)} F_{p, n_1 + n_2 - p - 1}(.01) = \frac{(57)(3)}{55} F_{3, 55}(.01) = 13.$$

we reject H_0 at the $\alpha = .01$ level. There is a difference in the (mean) cost vectors between gasoline trucks and diesel trucks.

$$b) \quad \hat{\underline{a}} = S_{\text{pooled}}^{-1} (\bar{\underline{x}}_1 - \bar{\underline{x}}_2) = \begin{bmatrix} 3.58 \\ -1.88 \\ -4.48 \end{bmatrix}$$

c) 99% simultaneous confidence intervals are:

$$\mu_{11} - \mu_{21}: 2.113 \pm 3.790$$

$$\mu_{12} - \mu_{22}: -2.650 \pm 4.341$$

$$\mu_{13} - \mu_{23}: -8.578 \pm 4.913$$

d) Assumption $t_1 = t_2$.

Since S_1 and S_2 are quite different, it may not be reasonable to pool. However, using "large sample" theory ($n_1 = 36$, $n_2 = 23$) we have, by Result 6.4,

$$(\bar{x}_1 - \bar{x}_2 - (\mu_1 - \mu_2))' \left[\frac{1}{n_1} S_1 + \frac{1}{n_2} S_2 \right]^{-1} (\bar{x}_1 - \bar{x}_2 - (\mu_1 - \mu_2)) \sim \chi_p^2$$

Since

$$(\bar{x}_1 - \bar{x}_2)' \left[\frac{1}{n_1} S_1 + \frac{1}{n_2} S_2 \right]^{-1} (\bar{x}_1 - \bar{x}_2) = 43.15 > \chi_3^2(.01) = 11.34$$

we reject $H_0: \mu_1 - \mu_2 = 0$ at the $\alpha = .01$ level. This is consistent with the result in part (a).

Note: part (d) was somewhat open-ended and not well defined. If students did not invoke Result 6.4, but repeated analysis with outliers removed and commented on it then that is okay.