

# Direct and Inverse Spectral Problems for 2-dimensional Hamiltonian Systems

Harald Woracek

Vienna University of Technology

Operator Model

oooo  
oooo

Examples

oo  
oo  
oooo

Case I:  $lc \leftrightarrow lc$

ooooo  
o  
oo

Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$

oo  
ooo  
oooooooo

Case III:  $lp \leftrightarrow lp$

oooo  
ooo  
oo

We are going to survey the spectral theory of 2-dimensional  
Hamiltonian Systems without a potential term.

Operator Model

oooo  
oooo

Examples

oo  
oo  
oo  
oooo

Case I:  $lc \leftrightarrow lc$

ooooo  
o  
oo

Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$

oo  
ooo  
oooooooo

Case III:  $lp \leftrightarrow lp$

ooooo  
ooo  
oo

We are going to survey the spectral theory of 2-dimensional Hamiltonian Systems without a potential term.

Many results are classical, dating back to the 1950's or 60's (for particular cases, even much earlier). Others are more recent, and taken from work of various authors.

We are going to survey the spectral theory of 2-dimensional Hamiltonian Systems without a potential term.

Many results are classical, dating back to the 1950's or 60's (for particular cases, even much earlier). Others are more recent, and taken from work of various authors.

We will also see a few results shown by myself (with different coauthors).

We are going to survey the spectral theory of 2-dimensional Hamiltonian Systems without a potential term.

Many results are classical, dating back to the 1950's or 60's (for particular cases, even much earlier). Others are more recent, and taken from work of various authors.

We will also see a few results shown by myself (with different coauthors).

These slides are available from my website

<http://asc.tuwien.ac.at/index.php?id=woracek>



Not all what is written on these slides is *strictly* correct.

We will occasionally neglect some technical difficulties and/or exeptional cases.

Each such instance will be clearly marked.

We consider  $2 \times 2$ -Hamiltonian systems without potential:

$$y'(t) = zJH(t)y(t), \quad t \in (s_-, s_+).$$

Here the *Hamiltonian*  $H$  shall be subject to

- $H(t) : (s_-, s_+) \rightarrow \mathbb{R}^{2 \times 2}$ ,
- $H(t) \geq 0$ ,  $t \in (s_-, s_+)$ ,
- $H \in L^1_{\text{loc}}(s_-, s_+)$ ,
- $H$  does not vanish identically on any set of positive measure,
- $z \in \mathbb{C}$  a parameter,
- $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ .

Operator Model

oooo  
oooo

Examples

oo  
oo  
oooo

Case I:  $lc \leftrightarrow lc$

ooooo  
o  
oo

Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$

oo  
ooo  
oooooooo

Case III:  $lp \leftrightarrow lp$

oooo  
ooo  
oo

This equation is the eigenvalue equation of a differential operator.  
We investigate the spectral theory of its selfadjoint realizations.



Operator Model

oooo  
oooo

Examples

oo  
oo  
oo  
ooooCase I:  $lc \leftrightarrow lc$ ooooo  
o  
ooCase II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$ oo  
ooo  
ooooooooCase III:  $lp \leftrightarrow lp$ oooo  
ooo  
oo

This equation is the eigenvalue equation of a differential operator. We investigate the spectral theory of its selfadjoint realizations.

**Direct Problems:** Given a Hamiltonian  $H$ , find information about spectral data of selfadjoint realizations.

This equation is the eigenvalue equation of a differential operator.  
We investigate the spectral theory of its selfadjoint realizations.

**Direct Problems:** Given a Hamiltonian  $H$ , find information about spectral data of selfadjoint realizations.

**Inverse Problems:**

This equation is the eigenvalue equation of a differential operator. We investigate the spectral theory of its selfadjoint realizations.

**Direct Problems:** Given a Hamiltonian  $H$ , find information about spectral data of selfadjoint realizations.

**Inverse Problems:**

- **Existence Theorems:** Given some spectral data, does there exist a Hamiltonian  $H$  which leads to this data.

This equation is the eigenvalue equation of a differential operator. We investigate the spectral theory of its selfadjoint realizations.

**Direct Problems:** Given a Hamiltonian  $H$ , find information about spectral data of selfadjoint realizations.

**Inverse Problems:**

- Existence Theorems: Given some spectral data, does there exist a Hamiltonian  $H$  which leads to this data.
- Uniqueness Theorems: Which spectral data obtained from some Hamiltonian determine this Hamiltonian uniquely.

# Outline

## Operator Model

Definition of  $L^2(H)$  and  $T_{\max}(H)$

Three Fundamental Cases

## Examples

The Schrödinger Equation

The String Equation

The Hamburger Moment Problem

Case I:  $lc \leftrightarrow lc$

Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$

Case III:  $lp \leftrightarrow lp$

## The Operator Model

Let  $H$  be a Hamiltonian on  $(s_-, s_+)$ , let  $(a, b) \subseteq (s_-, s_+)$  and  $\phi \in \mathbb{R}$ . Then  $(a, b)$  is  $H$ -indivisible of type  $\phi$ , if

$$H(t) = h(t) \cdot \begin{pmatrix} \cos \phi \\ \sin \phi \end{pmatrix} (\cos \phi, \sin \phi), \quad t \in (a, b),$$

with some scalar function  $h \in L^1_{\text{loc}}(a, b)$ .

## The Operator Model

Let  $H$  be a Hamiltonian on  $(s_-, s_+)$ , let  $(a, b) \subseteq (s_-, s_+)$  and  $\phi \in \mathbb{R}$ . Then  $(a, b)$  is *H-indivisible of type  $\phi$* , if

$$H(t) = h(t) \cdot \begin{pmatrix} \cos \phi \\ \sin \phi \end{pmatrix} (\cos \phi, \sin \phi), \quad t \in (a, b),$$

with some scalar function  $h \in L^1_{\text{loc}}(a, b)$ .

### Definition (The model space $L^2(H)$ )

The *model space*  $L^2(H)$  is the space of all  $f : (s_-, s_+) \rightarrow \mathbb{C}^2$  with

- $\|f\|_H^2 := \int_{s_-}^{s_+} f(t)^* H(t) f(t) dt < \infty$ .
- If  $(a, b) \subseteq (s_-, s_+)$  is indivisible of type  $\phi$ , then

$$(\cos \phi, \sin \phi) f(t) = \text{constant on } (a, b).$$

# The Operator Model

In the definition of  $L^2(H)$ , we tacitly understand that two functions  $f, g$  with

$$H(t)f(t) = H(t)g(t), \quad t \in (s_-, s_+) \text{ a.e.},$$

are identified.



## The Operator Model

In the definition of  $L^2(H)$ , we tacitly understand that two functions  $f, g$  with

$$H(t)f(t) = H(t)g(t), \quad t \in (s_-, s_+) \text{ a.e.},$$

are identified.

If endowed with the scalar product

$$(f, g)_H = \int_{s_-}^{s_+} g(t)^* H(t) f(t) dt, \quad f, g \in L^2(H),$$

the space  $L^2(H)$  becomes a Hilbert space.

# The Operator Model



Here we suppress some technical terms.

**Definition (The maximal operator  $T_{\max}(H)$ )**

The (graph of the) *maximal operator*  $T_{\max}(H)$  is

$$T_{\max}(H) = \left\{ (f; g) \in L^2(H) \times L^2(H) : \right. \\ \left. f \text{ is locally absolutely continuous and } f' = JHg \right\}$$

# The Operator Model



Here we suppress some technical terms.

**Definition** (The maximal operator  $T_{\max}(H)$ )

The (graph of the) *maximal operator*  $T_{\max}(H)$  is

$$T_{\max}(H) = \left\{ (f; g) \in L^2(H) \times L^2(H) : \right. \\ \left. f \text{ is locally absolutely continuous and } f' = JHg \right\}$$

The operator  $T_{\max}(H)$  is closed.

# The Operator Model

Definition (The minimal operator  $T_{\min}(H)$ )

The *minimal operator*  $T_{\min}(H)$  is  $T_{\min}(H) = T_{\max}(H)^*$ .

# The Operator Model

Definition (The minimal operator  $T_{\min}(H)$ )

The *minimal operator*  $T_{\min}(H)$  is  $T_{\min}(H) = T_{\max}(H)^*$ .

The operator  $T_{\min}(H)$  is closed and symmetric. It is either selfadjoint, or completely nonselfadjoint.

## Limit Circle vs. Limit Point Case

The spectral theory of  $T_{\max}(H)$  depends on the growth of  $H$  towards the endpoints  $s_-$  and  $s_+$ .

## Limit Circle vs. Limit Point Case

The spectral theory of  $T_{\max}(H)$  depends on the growth of  $H$  towards the endpoints  $s_-$  and  $s_+$ .

- $H$  is in *limit circle case* at  $s_-$ , if  $(x_0 \in (s_-, s_+))$

$$\int_{s_-}^{x_0} \operatorname{tr} H(t) dt < \infty \quad \left( \Leftrightarrow H \in L^1_{\text{loc}}([s_-, s_+)) \right).$$

- $H$  is in *limit point case* at  $s_-$ , if  $(x_0 \in (s_-, s_+))$

$$\int_{s_-}^{x_0} \operatorname{tr} H(t) dt = \infty.$$

## Limit Circle vs. Limit Point Case

The spectral theory of  $T_{\max}(H)$  depends on the growth of  $H$  towards the endpoints  $s_-$  and  $s_+$ .

- $H$  is in *limit circle case* at  $s_-$ , if  $(x_0 \in (s_-, s_+))$

$$\int_{s_-}^{x_0} \operatorname{tr} H(t) dt < \infty \quad \left( \Leftrightarrow H \in L_{\text{loc}}^1([s_-, s_+)) \right).$$

- $H$  is in *limit point case* at  $s_-$ , if  $(x_0 \in (s_-, s_+))$

$$\int_{s_-}^{x_0} \operatorname{tr} H(t) dt = \infty.$$

Similar: *limit circle case* at  $s_+$  and *limit point case* at  $s_+$ .



# The Three Fundamental Cases

We know the operator  $T_{\min}(H)$  is closed and symmetric.

## The Three Fundamental Cases

We know the operator  $T_{\min}(H)$  is closed and symmetric.

Its deficiency indices are always finite and equal.

## The Three Fundamental Cases

We know the operator  $T_{\min}(H)$  is closed and symmetric.

Its deficiency indices are always finite and equal.

- Case I,  $lc \leftrightarrow lc$ :  $(2, 2)$ .
- Case II,  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$ :  $(1, 1)$ .
- Case III,  $lp \leftrightarrow lp$ :  $(0, 0)$ .

## The Three Fundamental Cases

We know the operator  $T_{\min}(H)$  is closed and symmetric.

Its deficiency indices are always finite and equal.

- Case I,  $l_c \leftrightarrow l_c$ :  $(2, 2)$ .
- Case II,  $l_c \leftrightarrow l_p$  or  $l_p \leftrightarrow l_c$ :  $(1, 1)$ .
- Case III,  $l_p \leftrightarrow l_p$ :  $(0, 0)$ .

In Case III,  $T_{\min}(H) = T_{\max}(H)$ . Hence,  $T_{\min}(H)$  is selfadjoint and is the only selfadjoint realization.

In the Cases I and II, there are many different selfadjoint realizations.

## Boundary Values

If  $s_-$  is in limit circle case, each  $f = (f_1, f_2)^T \in \text{dom } T_{\max}(H)$  has a continuous extension to  $s_-$ . Similar for  $s_+$ .

## Boundary Values

If  $s_-$  is in limit circle case, each  $f = (f_1, f_2)^T \in \text{dom } T_{\max}(H)$  has a continuous extension to  $s_-$ . Similar for  $s_+$ .

Selfadjoint realizations can be described with boundary conditions.

## Boundary Values

If  $s_-$  is in limit circle case, each  $f = (f_1, f_2)^T \in \text{dom } T_{\max}(H)$  has a continuous extension to  $s_-$ . Similar for  $s_+$ .

Selfadjoint realizations can be described with boundary conditions.

- Assume Case lc ↔ lp. Then (for example)

$$A_D := \{(f; g) \in T_{\max}(H) : f_1(s_-) = 0\}$$

is a selfadjoint restriction of  $T_{\max}(H)$ .

## Boundary Values

If  $s_-$  is in limit circle case, each  $f = (f_1, f_2)^T \in \text{dom } T_{\max}(H)$  has a continuous extension to  $s_-$ . Similar for  $s_+$ .

Selfadjoint realizations can be described with boundary conditions.

- Assume Case  $lc \leftrightarrow lp$ . Then (for example)

$$A_D := \{(f; g) \in T_{\max}(H) : f_1(s_-) = 0\}$$

is a selfadjoint restriction of  $T_{\max}(H)$ .

- Assume Case  $lc \leftrightarrow lc$ . Then (for example) for each  $\tau \in \mathbb{R} \cup \{\infty\}$

$$A_{D,\tau} = \{(f; g) \in T_{\max}(H) : f_1(s_-) = 0, \tau f_1(s_+) + f_2(s_+) = 0\}$$

is a selfadjoint restrictions of  $T_{\max}(H)$ .



## Reparameterizations

Two Hamiltonians  $H_1$  on  $H_2$  defined on  $(s_-^1, s_+^1)$  and  $(s_-^2, s_+^2)$ , respectively, are *reparameterizations of each other*, if there exists

$$\phi : (s_-^2, s_+^2) \rightarrow (s_-^1, s_+^1)$$

such that

- $\phi$  is bijective and monotonically increasing,
- $\phi$  and  $\phi^{-1}$  are both absolutely continuous,
- $H_2(t) = H_1(\phi(t)) \cdot \phi'(t)$  for  $t \in (s_-^2, s_+^2)$ .

## Reparameterizations

Two Hamiltonians  $H_1$  on  $H_2$  defined on  $(s_-^1, s_+^1)$  and  $(s_-^2, s_+^2)$ , respectively, are *reparameterizations of each other*, if there exists

$$\phi : (s_-^2, s_+^2) \rightarrow (s_-^1, s_+^1)$$

such that

- $\phi$  is bijective and monotonically increasing,
- $\phi$  and  $\phi^{-1}$  are both absolutely continuous,
- $H_2(t) = H_1(\phi(t)) \cdot \phi'(t)$  for  $t \in (s_-^2, s_+^2)$ .

If  $H_1$  and  $H_2$  are reparameterizations of each other, their operator models are unitarily equivalent.

## Examples: 1. The Schrödinger Equation

Consider the equation  $(0 < T < \infty)$

$$-y''(t) + V(t)y(t) = zy(t), \quad t \in [0, T],$$

where the *potential*  $V(t)$  belongs to  $L^1([0, T])$ .

## Examples: 1. The Schrödinger Equation

Consider the equation  $(0 < T < \infty)$

$$-y''(t) + V(t)y(t) = zy(t), \quad t \in [0, T],$$

where the *potential*  $V(t)$  belongs to  $L^1([0, T])$ .

Let  $y_1$  and  $y_2$  be the solutions of  $-y''(t) + V(t)y(t) = 0$  with

$$y_1(0) = 0, y_1'(0) = 1, \quad y_2(0) = 1, y_2'(0) = 0,$$

and define

$$H(t) := \begin{pmatrix} y_1(t)^2 & y_1(t)y_2(t) \\ y_1(t)y_2(t) & y_2(t)^2 \end{pmatrix}, \quad t \in [0, T].$$

## Examples: 1. The Schrödinger Equation

Consider the equation  $(0 < T < \infty)$

$$-y''(t) + V(t)y(t) = zy(t), \quad t \in [0, T],$$

where the *potential*  $V(t)$  belongs to  $L^1([0, T])$ .

Let  $y_1$  and  $y_2$  be the solutions of  $-y''(t) + V(t)y(t) = 0$  with

$$y_1(0) = 0, y_1'(0) = 1, \quad y_2(0) = 1, y_2'(0) = 0,$$

and define

$$H(t) := \begin{pmatrix} y_1(t)^2 & y_1(t)y_2(t) \\ y_1(t)y_2(t) & y_2(t)^2 \end{pmatrix}, \quad t \in [0, T].$$

Then  $H$  is a Hamiltonian which is  $lc \leftrightarrow lc$ .

## Examples: 1. The Schrödinger Equation

A function  $y(t)$  solves the Schrödinger equation with potential  $V(t)$  and parameter  $z$ , if and only if the function

$$u(t) = \begin{pmatrix} y_1(t) & y_2(t) \\ y'_1(t) & y'_2(t) \end{pmatrix} \cdot \begin{pmatrix} y(t) \\ y'(t) \end{pmatrix}$$

solves the Hamiltonian system with Hamiltonian  $H(t)$  and parameter  $z$ .

## Examples: 1. The Schrödinger Equation

A function  $y(t)$  solves the Schrödinger equation with potential  $V(t)$  and parameter  $z$ , if and only if the function

$$u(t) = \begin{pmatrix} y_1(t) & y_2(t) \\ y_1'(t) & y_2'(t) \end{pmatrix} \cdot \begin{pmatrix} y(t) \\ y'(t) \end{pmatrix}$$

solves the Hamiltonian system with Hamiltonian  $H(t)$  and parameter  $z$ .

The operator models of the Schrödinger equation and of the Hamiltonian system are unitarily equivalent.

## Examples: 2. The String Equation



Here we ignore some technicalities.

Let  $L > 0$ , and  $\mu$  be a positive Borel measure on  $\mathbb{R}$  with  $\text{supp } \mu \subseteq [0, L]$  and  $\mu(\{L\}) = 0$ . Consider the integral equation boundary value problem with complex parameter  $z$ :

$$y'(t) + \int_{[0,t]} zy(u)d\mu(u) = 0, \quad y'(0-) = 0.$$



## Examples: 2. The String Equation



Here we ignore some technicalities.

Let  $L > 0$ , and  $\mu$  be a positive Borel measure on  $\mathbb{R}$  with  $\text{supp } \mu \subseteq [0, L]$  and  $\mu(\{L\}) = 0$ . Consider the integral equation boundary value problem with complex parameter  $z$ :

$$y'(t) + \int_{[0,t]} zy(u)d\mu(u) = 0, \quad y'(0-) = 0.$$

Set  $m(t) := \mu((-\infty, t))$ , and

$$\hat{m}(x) = \begin{cases} \inf \{t \geq 0 : x \leq m(t)\} , & x \in [0, m(L)] \\ L & , \quad x > m(L) \end{cases}$$

## Examples: 2. The String Equation



Here we ignore some technicalities.

Define

$$H(x) := \begin{pmatrix} \hat{m}(x)^2 & \hat{m}(x) \\ \hat{m}(x) & 1 \end{pmatrix}, \quad x \in (0, \infty).$$

## Examples: 2. The String Equation



Here we ignore some technicalities.

Define

$$H(x) := \begin{pmatrix} \hat{m}(x)^2 & \hat{m}(x) \\ \hat{m}(x) & 1 \end{pmatrix}, \quad x \in (0, \infty).$$

Then  $H$  is a Hamiltonian which is  $lc \leftrightarrow lp$ .

## Examples: 2. The String Equation



Here we ignore some technicalities.

Define

$$H(x) := \begin{pmatrix} \hat{m}(x)^2 & \hat{m}(x) \\ \hat{m}(x) & 1 \end{pmatrix}, \quad x \in (0, \infty).$$

Then  $H$  is a Hamiltonian which is  $lc \leftrightarrow lp$ .

The operator models of the string equation and of the Hamiltonian system with this Hamiltonian are unitarily equivalent.

## Examples: 3. The Hamburger Moment Problem

Let  $(s_n)_{n \geq 0}$  be a sequence of real numbers. Is this sequence the sequence of power moments of some positive Borel measure on the real line? That is, does there exist a positive Borel measure  $\mu$  on  $\mathbb{R}$  with

$$s_n = \int_{\mathbb{R}} t^n d\mu(t), \quad n \geq 0 \quad ?$$

## Examples: 3. The Hamburger Moment Problem

Let  $(s_n)_{n \geq 0}$  be a sequence of real numbers. Is this sequence the sequence of power moments of some positive Borel measure on the real line? That is, does there exist a positive Borel measure  $\mu$  on  $\mathbb{R}$  with

$$s_n = \int_{\mathbb{R}} t^n d\mu(t), \quad n \geq 0 \quad ?$$

The answer is yes, if and only if

$$\det [(s_{i+j})_{i,j=0}^N] \geq 0, \quad N \geq 0.$$

## Examples: 3. The Hamburger Moment Problem

Assume that  $D_N = \det[(s_{i+j})_{i,j=0}^N] \geq 0$ ,  $N \geq 0$ .

## Examples: 3. The Hamburger Moment Problem

Assume that  $D_N = \det[(s_{i+j})_{i,j=0}^N] \geq 0$ ,  $N \geq 0$ .

Either

- the solution of the Hamburger moment problem, i.e., the measure having  $(s_n)_{n \geq 0}$  as its moment sequence, is unique,

or

- there exist infinitely many such measures.



## Examples: 3. The Hamburger Moment Problem

Assume that  $D_N = \det[(s_{i+j})_{i,j=0}^N] \geq 0$ ,  $N \geq 0$ .

Either

- the solution of the Hamburger moment problem, i.e., the measure having  $(s_n)_{n \geq 0}$  as its moment sequence, is unique,

or

- there exist infinitely many such measures.

If  $D_N = 0$  for some  $N \geq 0$ , then the solution is unique and is a discrete measure with finitely many pointmasses.

## Examples: 3. The Hamburger Moment Problem

Assume that  $D_N = \det[(s_{i+j})_{i,j=0}^N] > 0$ ,  $N \geq 0$ .

## Examples: 3. The Hamburger Moment Problem

Assume that  $D_N = \det[(s_{i+j})_{i,j=0}^N] > 0$ ,  $N \geq 0$ .

Set

$$E_N = \det[(s_{i+j+1})_{i,j=0}^N], \quad C_N = \det[(s_{i+j-1})_{i,j=0}^N] \quad (s_{-1} = 0),$$

$$l_0 = 1, \quad l_N = (E_N^2 + C_N^2)(D_{N-1}D_N)^{-1}, \quad N \geq 1,$$

$$t_0 = 0, \quad t_N = \sum_{n=0}^{N-1} l_n, \quad n \geq 1, \quad T = \lim_{N \rightarrow \infty} t_n,$$

$$\theta_0 = \frac{\pi}{2}, \quad \theta_N = \begin{cases} \arctan\left(-\frac{E_N}{C_N}\right), & C_N \neq 0 \\ \frac{\pi}{2}, & C_N = 0 \end{cases}$$

## Examples: 3. The Hamburger Moment Problem

Define

$$H(t) = \begin{pmatrix} \cos \theta_N \\ \sin \theta_N \end{pmatrix} (\cos \theta_N, \sin \theta_N), \quad t \in [t_N, t_{N-1}), \quad N \geq 0.$$

## Examples: 3. The Hamburger Moment Problem

Define

$$H(t) = \begin{pmatrix} \cos \theta_N \\ \sin \theta_N \end{pmatrix} (\cos \theta_N, \sin \theta_N), \quad t \in [t_N, t_{N+1}), \quad N \geq 0.$$

Then  $H$  is a Hamiltonian which is

- $lc \leftrightarrow lp$ , if the solution is unique,
- $lc \leftrightarrow lc$ , if the solution is not unique.

## Examples: 3. The Hamburger Moment Problem

Define

$$H(t) = \begin{pmatrix} \cos \theta_N \\ \sin \theta_N \end{pmatrix} (\cos \theta_N, \sin \theta_N), \quad t \in [t_N, t_{N+1}), \quad N \geq 0.$$

Then  $H$  is a Hamiltonian which is

- $lc \leftrightarrow lp$ , if the solution is unique,
- $lc \leftrightarrow lc$ , if the solution is not unique.

The set of solutions of the Hamburger moment problem coincides with the set of all *spectral measures* of the Hamiltonian  $H$ .

## Case $lc \leftrightarrow lc$ . Spectral Measures

Denote by  $W(t, z) = (w_{ij}(t, z))_{i,j=1}^2$  the solution of

$$\frac{d}{dt}W(t, z)J = zW(t, z)H(t), \quad t \in [s_-, s_+], \quad W(s_-, z) = I.$$

## Case $lc \leftrightarrow lc$ . Spectral Measures

Denote by  $W(t, z) = (w_{ij}(t, z))_{i,j=1}^2$  the solution of

$$\frac{d}{dt}W(t, z)J = zW(t, z)H(t), \quad t \in [s_-, s_+], \quad W(s_-, z) = I.$$

Then, for each  $\tau \in \mathbb{R} \cup \{\infty\}$ , the function

$$q_{H,\tau}(z) = \frac{w_{11}(s_+, z)\tau + w_{12}(s_+, z)}{w_{21}(s_+, z)\tau + w_{22}(s_+, z)}$$

belongs to the *Nevanlinna class*, that is,

- $q_H$  is analytic in  $\mathbb{C} \setminus \mathbb{R}$  and  $q_H(\bar{z}) = \overline{q_H(z)}$ ,
- $\operatorname{Im} q_H(z) \geq 0$  for  $\operatorname{Im} z > 0$ .



## Case $lc \leftrightarrow lc$ . Spectral Measures

We can represent  $q_{H,\tau}$  as (*Herglotz integral representation*)

$$q_{H,\tau}(z) = a_{H,\tau} + b_{H,\tau}z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_{H,\tau}(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

with

- $a_{H,\tau} \in \mathbb{R}$ ,  $b_{H,\tau} \geq 0$ ,
- $\mu_H$  positive Borel measure with  $\int_{\mathbb{R}} \frac{d\mu_{H,\tau}(t)}{1+t^2} < \infty$ .

## Case $lc \leftrightarrow lc$ . Spectral Measures

We can represent  $q_{H,\tau}$  as (*Herglotz integral representation*)

$$q_{H,\tau}(z) = a_{H,\tau} + b_{H,\tau}z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_{H,\tau}(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

with

- $a_{H,\tau} \in \mathbb{R}$ ,  $b_{H,\tau} \geq 0$ ,
- $\mu_H$  positive Borel measure with  $\int_{\mathbb{R}} \frac{d\mu_{H,\tau}(t)}{1+t^2} < \infty$ .

The measure  $\mu_{H,\tau}$  can be computed from  $q_{H,\tau}$  by means of the Stieltjes Inversion Formula, and the constant  $b_{H,\tau}$  from the behaviour of  $q_{H,\tau}$  towards  $i\infty$ .

## Case $l_c \leftrightarrow l_c$ . Spectral Measures

We can represent  $q_{H,\tau}$  as (*Herglotz integral representation*)

$$q_{H,\tau}(z) = a_{H,\tau} + b_{H,\tau}z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_{H,\tau}(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

with

- $a_{H,\tau} \in \mathbb{R}$ ,  $b_{H,\tau} \geq 0$ ,
- $\mu_H$  positive Borel measure with  $\int_{\mathbb{R}} \frac{d\mu_{H,\tau}(t)}{1+t^2} < \infty$ .

The measure  $\mu_{H,\tau}$  can be computed from  $q_{H,\tau}$  by means of the Stieltjes Inversion Formula, and the constant  $b_{H,\tau}$  from the behaviour of  $q_{H,\tau}$  towards  $i\infty$ .

Each measure  $\mu_{H,\tau}$  obtained in this way is called a *spectral measure of  $H$* .

## Case $lc \leftrightarrow lc$ . Fourier Transforms



Here we ignore the exceptional case  $b_{H,\tau} > 0$  and technicalities.

### Theorem (Integral transforms, $lc \leftrightarrow lp$ )

Let  $\tau \in \mathbb{R} \cup \{\infty\}$ . A unitary map  $U_\tau : L^2(H) \rightarrow L^2(\mu_{H,\tau})$  is defined by

$$(U_\tau f)(x) = \int_{s_-}^{s_+} (w_{21}(t, x), w_{22}(t, x)) H(t) f(t) dt.$$

Its inverse  $U_\tau^{-1} : L^2(\mu_{H,\tau}) \rightarrow L^2(H)$  is given as

$$(U_\tau^{-1} F)(t) = \int_{-\infty}^{\infty} \begin{pmatrix} w_{21}(t, x) \\ w_{22}(t, x) \end{pmatrix} F(x) d\mu_{H,\tau}(x).$$

## Case $lc \leftrightarrow lc$ . Fourier Transforms



Here we ignore the exceptional case  $b_{H,\tau} > 0$  and technicalities.

### Theorem (Unitary equivalence)

Let  $\tau \in \mathbb{R} \cup \{\infty\}$ . The selfadjoint realization

$$A_{D,\tau} = \{(f; g) \in T_{\max}(H) : f_1(s_-) = 0, \tau f_1(s_+) + f_2(s_+) = 0\}$$

is unitarily equivalent to the multiplication operator  $M_x$  in  $L^2(\mu_{H,\tau})$  via  $U_\tau$ , that is,

$$U_\tau \circ A_{D,\tau} = M_x \circ U_\tau.$$

## Case $l_c \leftrightarrow l_c$ . A Direct Theorem

### Theorem (Direct Spectral Theorem)

- *Each selfadjoint realization has compact resolvents.*
- *The spectrum of each selfadjoint realization defined by separated boundary conditions is simple.*
- *Let  $(\lambda_n^+)$  and  $(\lambda_n^-)$  denote the sequences of positive and negative, respectively, eigenvalues of a selfadjoint realization arranged according to increasing modulus. Then*

$$\lim_{n \rightarrow \infty} \frac{n}{\lambda_n^+} = \lim_{n \rightarrow \infty} \frac{n}{\lambda_n^-} = \frac{1}{\pi} \int_{s_-}^{s_+} \sqrt{\det H(t)} dt.$$

## Case $lc \leftrightarrow lc$ . A Uniqueness Theorem

A Hamiltonian  $H$  is not uniquely determined by the spectrum of one of its selfadjoint realizations. It may happen that  $H_1$  and  $H_2$  are different (not reparameterizations of each other), and still

$$\sigma(A_{D,0}^1) = \sigma(A_{D,0}^2).$$

## Case $lc \leftrightarrow lc$ . A Uniqueness Theorem

A Hamiltonian  $H$  is not uniquely determined by the spectrum of one of its selfadjoint realizations. It may happen that  $H_1$  and  $H_2$  are different (not reparameterizations of each other), and still

$$\sigma(A_{D,0}^1) = \sigma(A_{D,0}^2).$$

### Theorem (Inverse Theorem / Uniqueness)

*Assume that two Hamiltonians  $H_1$  and  $H_2$  satisfy*

$$\sigma(A_{D,0}^1) = \sigma(A_{D,0}^2) \quad \text{and} \quad \sigma(A_{D,\infty}^1) = \sigma(A_{D,\infty}^2),$$

*then  $H_1$  and  $H_2$  are equal (up to a reparameterization).*



## Case $lc \leftrightarrow lc$ . Existence Theorems

### Theorem (Characterization of spectra)

Let  $(\lambda_n)$  be a sequence of pairwise different real numbers. Then there exists a Hamiltonian  $H$  in  $lc \leftrightarrow lc$  with  $\{\lambda_n\} = \sigma(A_{D,0})$ , if and only if all  $\lambda_n$  are nonzero, and

- the limits  $\lim \frac{n}{\lambda_n^+}$  and  $\lim \frac{n}{\lambda_n^-}$  exist in  $[0, \infty)$  and are equal, where  $(\lambda_n^+)$  and  $(\lambda_n^-)$  denote the sequences of positive and negative elements of  $(\lambda_n)$ ,
- $\lim_{R \rightarrow \infty} \sum_{|\lambda_n| \leq R} \frac{1}{\lambda_n}$  exists in  $\mathbb{R}$ ,
- $\sum_n \frac{1}{|\lambda_n|^2 |A'(\lambda_n)|} < \infty$  where  $A(z) = \lim_{R \rightarrow \infty} \prod_{|\lambda_n| \leq R} \left(1 - \frac{z}{\lambda_n}\right)$ .

## Case $lc \leftrightarrow lc$ . Existence Theorems

### Theorem (Characterization of pairs of spectra)

Let  $(\lambda_n)$  and  $(\mu_n)$  be two sequences of pairwise different real numbers. Then there exists a Hamiltonian  $H$  in  $lc \leftrightarrow lc$  with  $\{\lambda_n\} = \sigma(A_{D,0})$  and  $\{\mu_n\} = \sigma(A_{D,\infty})$ , if and only if all  $\lambda_n$  are nonzero, the point zero is among the  $\mu_n$ 's, and

- the sequences  $(\lambda_n)$  and  $(\mu_n)$  interlace,
- $\lim_{\lambda_n^+} \frac{n}{\lambda_n} = \lim_{\lambda_n^-} \frac{n}{\lambda_n} \in [0, \infty)$  and  $\lim_{R \rightarrow \infty} \sum_{|\lambda_n| \leq R} \frac{1}{\lambda_n}$  exists in  $\mathbb{R}$ ,
- $\sum_n \frac{1}{|\lambda_n|^2 |A'(\lambda_n) B(\lambda_n)|} < \infty$  where

$$A(z) = \lim_{R \rightarrow \infty} \prod_{|\lambda_n| \leq R} \left(1 - \frac{z}{\lambda_n}\right), \quad B(z) = z \lim_{R \rightarrow \infty} \prod_{0 < |\mu_n| \leq R} \left(1 - \frac{z}{\mu_n}\right).$$

Operator Model

oooo  
oooo

Examples

oo  
oo  
oo  
oooo

Case I:  $lc \leftrightarrow lc$

ooooo  
o  
oo

Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$

●o  
ooo  
oooooooo

Case III:  $lp \leftrightarrow lp$

ooooo  
ooo  
oo

## Case $lc \leftrightarrow lp$ . The Weyl Construction

The cases  $lc \leftrightarrow lp$  and  $lp \leftrightarrow lc$  are fully analogous. We confine attention to  $lc \leftrightarrow lp$ .

## Case $lc \leftrightarrow lp$ . The Weyl Construction

The cases  $lc \leftrightarrow lp$  and  $lp \leftrightarrow lc$  are fully analogous. We confine attention to  $lc \leftrightarrow lp$ .

### Theorem (Weyl coefficient. Existence)

Denote by  $W(t, z) = (w_{ij}(t, z))_{i,j=1}^2$  the solution of

$$\frac{d}{dt}W(t, z)J = zW(t, z)H(t), \quad t \in [s_-, s_+), \quad W(s_-, z) = I.$$

Then, for each  $\tau \in \mathbb{R} \cup \{\infty\}$  the limit

$$q_H(z) = \lim_{t \nearrow s_+} \frac{w_{11}(t, z)\tau + w_{12}(t, z)}{w_{21}(t, z)\tau + w_{22}(t, z)}$$

exists locally uniformly on  $\mathbb{C} \setminus \mathbb{R}$ . It does not depend on  $\tau$ .

## Case $lc \leftrightarrow lp$ . The Weyl Construction

### Theorem (Weyl coefficient. Properties)

*The function  $q_H$  belongs to the Nevanlinna class, that is,*

- *$q_H$  is analytic in  $\mathbb{C} \setminus \mathbb{R}$  and  $q_H(\bar{z}) = \overline{q_H(z)}$ ,*
- *$\operatorname{Im} q_H(z) \geq 0$  for  $\operatorname{Im} z > 0$ .*

## Case $lc \leftrightarrow lp$ . The Weyl Construction

### Theorem (Weyl coefficient. Properties)

The function  $q_H$  belongs to the Nevanlinna class, that is,

- $q_H$  is analytic in  $\mathbb{C} \setminus \mathbb{R}$  and  $q_H(\bar{z}) = \overline{q_H(z)}$ ,
- $\operatorname{Im} q_H(z) \geq 0$  for  $\operatorname{Im} z > 0$ .

We can therefore represent  $q_H$  as

$$q_H(z) = a_H + b_H z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_H(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

with

- $a_H \in \mathbb{R}$ ,  $b_H \geq 0$ ,
- $\mu_H$  positive Borel measure with  $\int_{\mathbb{R}} \frac{d\mu_H(t)}{1+t^2} < \infty$ .

## Case $lc \leftrightarrow lp$ . The Weyl Construction

### Theorem (Weyl coefficient. Properties)

The function  $q_H$  belongs to the Nevanlinna class, that is,

- $q_H$  is analytic in  $\mathbb{C} \setminus \mathbb{R}$  and  $q_H(\bar{z}) = \overline{q_H(z)}$ ,
- $\operatorname{Im} q_H(z) \geq 0$  for  $\operatorname{Im} z > 0$ .

We can therefore represent  $q_H$  as

$$q_H(z) = a_H + b_H z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) d\mu_H(t), \quad z \in \mathbb{C} \setminus \mathbb{R},$$

with

- $a_H \in \mathbb{R}$ ,  $b_H \geq 0$ ,
- $\mu_H$  positive Borel measure with  $\int_{\mathbb{R}} \frac{d\mu_H(t)}{1+t^2} < \infty$ .

The measure  $\mu_H$  obtained in this way is called the *spectral measure of  $H$* .

## Case $lc \leftrightarrow lp$ . A Direct Theorem



Here we ignore the exceptional case  $b_H > 0$  and technicalities.

### Theorem (Direct Spectral Theorem)

A unitary map  $U : L^2(H) \rightarrow L^2(\mu_H)$  is defined by

$$(Uf)(x) = \int_{s_-}^{s_+} (w_{21}(t, x), w_{22}(t, x)) H(t) f(t) dt.$$

It intertwines  $A_D$  and the multiplication operator  $M_x$  in  $L^2(\mu_H)$ :

$$U \circ A_D = M_x \circ U.$$

Its inverse  $U^{-1} : L^2(\mu_H) \rightarrow L^2(H)$  is given as

$$(U^{-1}F)(t) = \int_{-\infty}^{\infty} \begin{pmatrix} w_{21}(t, x) \\ w_{22}(t, x) \end{pmatrix} F(x) d\mu_H(x).$$



## Case $l_c \leftrightarrow l_p$ . The Inverse Theorem

The following Existence and Uniqueness Theorem is *the* major result in the spectral theory of Hamiltonian systems.

## Case $lc \leftrightarrow lp$ . The Inverse Theorem

The following Existence and Uniqueness Theorem is *the* major result in the spectral theory of Hamiltonian systems.

### Theorem (Inverse Spectral Theorem)

*Let a function  $q$  in the Nevanlinna class be given. Equivalently, let  $a \in \mathbb{R}$ ,  $b \geq 0$ , and a positive Borel measure  $\mu$  with  $\int_{\mathbb{R}} \frac{d\mu(t)}{1+t^2}$  be given.*

*Then there exists a (up to reparameterization) unique Hamiltonian in  $lc \leftrightarrow lp$  whose Weyl coefficient equals  $q$ .*

## Case $lc \leftrightarrow lp$ . A Local Uniqueness Theorem

### Theorem (Local uniqueness)

Let  $H_1$  and  $H_2$  be Hamiltonians defined on  $(s_-^1, s_+^1)$  and  $(s_-^2, s_+^2)$ , respectively. For  $a > 0$  set

$$s_a^j = \sup \left\{ t \in [s_-^j, s_+^j) : \int_{s_-^j}^t \sqrt{\det H_j(x)} dx < a \right\}, \quad j = 1, 2.$$

Then the following are equivalent.

- $H_1 \big|_{(s_-^1, s_a^1)}$  and  $H_2 \big|_{(s_-^2, s_a^2)}$  are reparameterizations of each other.
- $q_{H_1}(z) - q_{H_2}(z) = O((\operatorname{Im} z)^3 e^{-2a \operatorname{Im} z})$ ,  $z \hat{\rightarrow} i\infty$ .

## Case $lc \leftrightarrow lp$ . Semibounded Spectrum

### Theorem (Consequence of semibounded spectrum)

Let  $H$  be given with  $\inf \operatorname{supp} \mu_H > -\infty$ . Then there exist unique  $L \in (0, \infty]$  and  $\nu : [0, L) \rightarrow [0, +\infty)$ , such that

- $\nu$  is nondecreasing, right-continuous, and normalized by  $\nu(0) \in [0, \pi)$  and  $\nu(t) - \nu(t-) < \pi$ ,
- $H$  is (a reparameterization of)

$$H_\nu(x) = \begin{cases} \begin{pmatrix} [\cot \nu(t)]^2 & -\cot \nu(t) \\ -\cot \nu(t) & 1 \end{pmatrix} & \text{if } \nu(t) \notin \pi\mathbb{Z} \\ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} & \text{if } \nu(t) \in \pi\mathbb{Z} \end{cases}$$

## Case $l_c \leftrightarrow l_p$ . Semibounded Spectrum

The question converse to the above theorem is: Given  $\nu$  with the properties stated in the theorem. Is the spectral measure of  $H_\nu$  semibounded from below ?

## Case $l_c \leftrightarrow l_p$ . Semibounded Spectrum

The question converse to the above theorem is: Given  $\nu$  with the properties stated in the theorem. Is the spectral measure of  $H_\nu$  semibounded from below ?

The answer is *unknown*.

## Case $l_c \leftrightarrow l_p$ . Semibounded Spectrum

### Theorem (The case of finite negativity)

*The following are equivalent:*

- $H$  is (up to a reparameterization) equal to  $H_\nu$  with  $\nu$  being (in addition) bounded.
- $(-\infty, 0) \cap \text{supp } \mu_H$  is a finite set.

## Case $l_c \leftrightarrow l_p$ . Discrete Spectrum

It is a question of interest for which Hamiltonians their selfadjoint realizations have discrete spectrum (equivalently, have compact resolvents).



## Case $l_c \leftrightarrow l_p$ . Discrete Spectrum

It is a question of interest for which Hamiltonians their selfadjoint realizations have discrete spectrum (equivalently, have compact resolvents).

The answer is *unknown*.

## Case $l_c \leftrightarrow l_p$ . Discrete Spectrum

It is a question of interest for which Hamiltonians their selfadjoint realizations have discrete spectrum (equivalently, have compact resolvents).

The answer is *unknown*.

What is easy to see is that for each discrete subset  $M$  of  $\mathbb{R}$ , there exist (infinitely many) Hamiltonians  $H$  which are  $l_c \leftrightarrow l_p$  and such that

$$\sigma(A_D) = M.$$

## Case $lc \leftrightarrow lp$ . Discrete Spectrum

Write  $H(t) = (h_{ij}(t))_{i,j=1}^2$ , and set  $B(t) = \int_{s_-}^t h_{12}(x) dx$ .

For  $r > 0$  set

$$M_1(r) = \left\{ \lambda \in \mathbb{R} \setminus \{0\} : \limsup_{t \nearrow s_+} \left( \int_{s_-}^t h_{22}(x) e^{-2\lambda B(x)} dx \int_t^{s_+} h_{11}(x) e^{2\lambda B(x)} dx \right) \leq \frac{r}{\lambda^2} \right\},$$

$$M_2(r) = \left\{ \lambda \in \mathbb{R} \setminus \{0\} : \limsup_{t \nearrow s_+} \left( \int_{s_-}^t h_{11}(x) e^{2\lambda B(x)} dx \int_t^{s_+} h_{22}(x) e^{-2\lambda B(x)} dx \right) \leq \frac{r}{\lambda^2} \right\}.$$

## Case $l_c \leftrightarrow l_p$ . Discrete Spectrum



In the literature this theorem is only stated. We have not seen a proof.

### Theorem (Discreteness of spectrum)

If  $\mu_H$  is discrete, then

$$\mathbb{R} \setminus \{0\} \subseteq M_1(1) \cup M_2(1).$$

If there exist sequences  $\lambda_i \rightarrow +\infty$  and  $\mu_i \rightarrow -\infty$  with

$$\{\lambda_i : i \in \mathbb{N}\} \cup \{\mu_i : i \in \mathbb{N}\} \subseteq \bigcup_{r < \frac{1}{4}} (M_1(r) \cup M_2(r)),$$

then  $\mu_H$  is discrete.

## Case lc ↔ lp. Discrete Spectrum

### Theorem (The diagonal case)

Assume that  $H$  is diagonal, that is,

$$H(x) = \begin{pmatrix} h_1(x) & 0 \\ 0 & h_2(x) \end{pmatrix}, \quad x \in (s_-, s_+) \text{ a.e.}$$

Then  $\mu_H$  is discrete, if and only if either

$$\int_{s_-}^{s_+} h_1(x) dx < \infty \text{ and } \lim_{x \nearrow s_+} \left( \int_x^{s_+} h_1(t) dt \cdot \int_{s_-}^x h_2(t) dt \right) = 0,$$

or

$$\int_{s_-}^{s_+} h_2(x) dx < \infty \text{ and } \lim_{x \nearrow s_+} \left( \int_x^{s_+} h_2(t) dt \cdot \int_{s_-}^x h_1(t) dt \right) = 0.$$

## Case $l_c \leftrightarrow l_p$ . Hilbert-Schmidt Property

Contrasting discreteness of the spectrum, the property that the spectrum is discrete with square summable eigenvalues can be characterized in general.

## Case $l_c \leftrightarrow l_p$ . Hilbert-Schmidt Property

Contrasting discreteness of the spectrum, the property that the spectrum is discrete with square summable eigenvalues can be characterized in general.

### Theorem (Characterization of Hilbert-Schmidt class)

*The resolvents of selfadjoint realizations belong to the Hilbert-Schmidt class, if and only if there exists an angle  $\phi \in [0, \pi)$  such that (here  $\xi_\alpha = \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix}$ )*

- $\int_{s_-}^{s_+} \xi_\phi^T H(t) \xi_\phi dt < \infty,$
- $\int_{s_-}^{s_+} \left( \int_{s_-}^t \xi_{\phi+\frac{\pi}{2}}^T H(u) \xi_{\phi+\frac{\pi}{2}} du \right) \xi_\phi^T H(t) \xi_\phi dt < \infty.$

## Case $l_p \leftrightarrow l_p$ . Vector-Valued $L^2$ -Spaces

Let  $\Omega = (\Omega_{ij})_{i,j=1}^n$  be a positive  $n \times n$ -matrix valued Borel measure on  $\mathbb{R}$ , that is, a map of Borel sets to positive semidefinite  $n \times n$ -matrices which is  $\sigma$ -additive and satisfies  $\Omega(\emptyset) = 0$ .



## Case $lp \leftrightarrow lp$ . Vector-Valued $L^2$ -Spaces

Let  $\Omega = (\Omega_{ij})_{i,j=1}^n$  be a positive  $n \times n$ -matrix valued Borel measure on  $\mathbb{R}$ , that is, a map of Borel sets to positive semidefinite  $n \times n$ -matrices which is  $\sigma$ -additive and satisfies  $\Omega(\emptyset) = 0$ .

Set  $\rho(\Delta) = \text{tr } \Omega(\Delta)$ , then  $\rho$  is a finite positive Borel measure on  $\mathbb{R}$  and each entry  $\Omega_{ij}$  is absolutely continuous w.r.t.  $\rho$ . The *symmetric derivative* of  $\Omega$  w.r.t.  $\rho$  is

$$\frac{d\Omega}{d\rho}(x) = \lim_{\varepsilon \downarrow 0} \frac{\Omega((x - \varepsilon, x + \varepsilon))}{\rho((x - \varepsilon, x + \varepsilon))}, \quad x \in \mathbb{R} \text{ } \rho\text{-a.e.}$$

## Case $l_p \leftrightarrow l_p$ . Vector-Valued $L^2$ -Spaces

Let  $\Omega = (\Omega_{ij})_{i,j=1}^n$  be a positive  $n \times n$ -matrix valued Borel measure on  $\mathbb{R}$ , that is, a map of Borel sets to positive semidefinite  $n \times n$ -matrices which is  $\sigma$ -additive and satisfies  $\Omega(\emptyset) = 0$ .

Set  $\rho(\Delta) = \text{tr } \Omega(\Delta)$ , then  $\rho$  is a finite positive Borel measure on  $\mathbb{R}$  and each entry  $\Omega_{ij}$  is absolutely continuous w.r.t.  $\rho$ . The *symmetric derivative* of  $\Omega$  w.r.t.  $\rho$  is

$$\frac{d\Omega}{d\rho}(x) = \lim_{\varepsilon \downarrow 0} \frac{\Omega((x - \varepsilon, x + \varepsilon))}{\rho((x - \varepsilon, x + \varepsilon))}, \quad x \in \mathbb{R} \text{ } \rho\text{-a.e.}$$

$L^2(\Omega)$  is the space of all  $f : \mathbb{R} \rightarrow \mathbb{C}^n$  which are  $\rho$ -measurable and  $(f(x), \frac{d\Omega}{d\rho}(x)f(x))_{\mathbb{C}^n} \in L^1(\rho)$ . It is endowed with

$$(f, g)_{L^2(\Omega)} = \int_{\mathbb{R}} (f(x), \frac{d\Omega}{d\rho}(x)g(x))_{\mathbb{C}^n} d\rho(x).$$

## Case $lp \leftrightarrow lp$ . Matrix Weyl Function

Choose a point  $s_0 \in (s_-, s_+)$ . Then

- $H_+ = H|_{(s_0, s_+)}$  is a Hamiltonian  $lc \leftrightarrow lp$ .
- $H_- = H|_{(s_-, s_0)}$  is a Hamiltonian in  $lp \leftrightarrow lc$ ,

## Case $lp \leftrightarrow lp$ . Matrix Weyl Function

Choose a point  $s_0 \in (s_-, s_+)$ . Then

- $H_+ = H|_{(s_0, s_+)}$  is a Hamiltonian  $lc \leftrightarrow lp$ .
- $H_- = H|_{(s_-, s_0)}$  is a Hamiltonian in  $lp \leftrightarrow lc$ ,

The  $2 \times 2$ -matrix valued function

$$Q_H(z) = \frac{1}{q_{H_+}(z) + q_{H_-}(z)} \begin{pmatrix} q_{H_+}(z)q_{H_-}(z) & -q_{H_+}(z) \\ -q_{H_+}(z) & -1 \end{pmatrix}$$

belongs to the  $2 \times 2$ -*Nevanlinna class*, that is,

- $Q_H$  is analytic in  $\mathbb{C} \setminus \mathbb{R}$  and  $Q_H(\bar{z}) = Q_H(z)^*$ ,
- $\operatorname{Im} q_H(z) = \frac{1}{2i}(Q_H(z) - Q_H(z)^*)$  is positive semidefinite for each  $z$  with  $\operatorname{Im} z > 0$ .

## Case $lp \leftrightarrow lp$ . Matrix Weyl Function

We can represent  $Q_H$  as ( $z \in \mathbb{C} \setminus \mathbb{R}$ )

$$Q_H(z) = a_H + b_H z + \int_{\mathbb{R}} \left( \frac{1}{t-z} - \frac{t}{1+t^2} \right) \cdot (1+t^2) d\Omega_H(t),$$

where

- $a_H, b_H \in \mathbb{C}^{2 \times 2}$ , with  $a_H = a_H^*$  and  $b_H$  positive semidefinite,
- $\Omega_H$  positive  $2 \times 2$ -matrix valued Borel measure.

## Case $lp \leftrightarrow lp$ . The Titchmarsh-Kodaira formula



Here we ignore the exceptional case  $b_H \neq 0$ .

### Theorem (Unitary equivalence)

*The selfadjoint operator  $T_{\min}(H)$  is unitarily equivalent to the multiplication operator in the space  $L^2(\Omega_H)$ . That is, there exists a unitary operator  $U : L^2(H) \rightarrow L^2(\Omega_H)$  which intertwines  $T_{\min}(H)$  and  $M_x$ :*

$$U \circ T_{\min}(H) = M_x \circ U.$$

## Case $lp \leftrightarrow lp$ . The Titchmarsh-Kodaira formula



Here we ignore the exceptional case  $b_H \neq 0$ .

### Theorem (Unitary equivalence)

*The selfadjoint operator  $T_{\min}(H)$  is unitarily equivalent to the multiplication operator in the space  $L^2(\Omega_H)$ . That is, there exists a unitary operator  $U : L^2(H) \rightarrow L^2(\Omega_H)$  which intertwines  $T_{\min}(H)$  and  $M_x$ :*

$$U \circ T_{\min}(H) = M_x \circ U.$$

The action of  $U$  can again be described as an integral transform.

## Case $l_p \leftrightarrow l_p$ . The Titchmarsh-Kodaira formula



Here we ignore the exceptional case  $b_H \neq 0$ .

### Theorem (Unitary equivalence)

*The selfadjoint operator  $T_{\min}(H)$  is unitarily equivalent to the multiplication operator in the space  $L^2(\Omega_H)$ . That is, there exists a unitary operator  $U : L^2(H) \rightarrow L^2(\Omega_H)$  which intertwines  $T_{\min}(H)$  and  $M_x$ :*

$$U \circ T_{\min}(H) = M_x \circ U.$$

The action of  $U$  can again be described as an integral transform.

### Corollary

*The spectral multiplicity of  $T_{\min}(H)$  cannot exceed 2.*



Operator Model

○○○○  
○○○○

Examples

○○  
○○  
○○○○Case I:  $lc \leftrightarrow lc$ ○○○○○  
○  
○○Case II:  $lc \leftrightarrow lp$  or  $lp \leftrightarrow lc$ ○○  
○○○  
○○○○○○○○Case III:  $lp \leftrightarrow lp$ ○○○○  
●○○  
○○

## Case $lp \leftrightarrow lp$ . Local Spectral Multiplicity

Let  $E$  be the projection valued spectral measure of  $T_{\min}(H)$ , and let  $\sigma_1, \sigma_2$  be scalar positive Borel measures with  $\sigma_2 \ll \sigma_1 \sim E$  such that  $T_{\min}(H)$  is unitarily equivalent to the multiplication operator in  $L^2(\sigma_1) \oplus L^2(\sigma_2)$ .



Operator Model

○○○○  
○○○○

Examples

○○  
○○  
○○  
○○○○Case I:  $l_c \leftrightarrow l_c$ ○○○○○  
○  
○○Case II:  $l_c \leftrightarrow l_p$  or  $l_p \leftrightarrow l_c$ ○○  
○○○  
○○○○○○○○Case III:  $l_p \leftrightarrow l_p$ ○○○○○  
○●○  
○○

## Case $l_p \leftrightarrow l_p$ . Local Spectral Multiplicity

### Theorem

We have  $N_H(x) = \text{rank} \frac{d\Omega_H}{d\rho}(x)$  for  $x \in \mathbb{R}$ ,  $\rho$ -a.e. (notice  $\rho \sim E$ ).

## Case $l_p \leftrightarrow l_p$ . Local Spectral Multiplicity

### Theorem

We have  $N_H(x) = \text{rank} \frac{d\Omega_H}{d\rho}(x)$  for  $x \in \mathbb{R}$ ,  $\rho$ -a.e. (notice  $\rho \sim E$ ).

Denote by  $\lambda$  the Lebesgue measure, and set  $\mu := \mu_{H_+} + \mu_{H_-}$ .

## Case $lp \leftrightarrow lp$ . Local Spectral Multiplicity

### Theorem

We have  $N_H(x) = \text{rank } \frac{d\Omega_H}{d\rho}(x)$  for  $x \in \mathbb{R}$ ,  $\rho$ -a.e. (notice  $\rho \sim E$ ).

Denote by  $\lambda$  the Lebesgue measure, and set  $\mu := \mu_{H_+} + \mu_{H_-}$ .

Decompose

$$E = E_{ac} + E_s \quad \text{with} \quad E_{ac} \ll \lambda, \quad E_s \perp \lambda,$$

and further

$$E_s = E_{s,ac} + E_{s,s} \quad \text{with} \quad E_{s,ac} \ll \mu, \quad E_{s,s} \perp \mu.$$

## Case $l_p \leftrightarrow l_p$ . Local Spectral Multiplicity

### Theorem

We have  $N_H(x) = \text{rank } \frac{d\Omega_H}{d\rho}(x)$  for  $x \in \mathbb{R}$ ,  $\rho$ -a.e. (notice  $\rho \sim E$ ).

Denote by  $\lambda$  the Lebesgue measure, and set  $\mu := \mu_{H_+} + \mu_{H_-}$ .

Decompose

$$E = E_{ac} + E_s \quad \text{with} \quad E_{ac} \ll \lambda, \quad E_s \perp \lambda,$$

and further

$$E_s = E_{s,ac} + E_{s,s} \quad \text{with} \quad E_{s,ac} \ll \mu, \quad E_{s,s} \perp \mu.$$

Moreover, decompose

$$\mu = \mu_{ac} + \mu_s \quad \text{with} \quad \mu \ll \lambda, \quad \mu \perp \lambda.$$

## Case $l_p \leftrightarrow l_p$ . Local Spectral Multiplicity

Set  $(x \in \mathbb{R}, \mu\text{-a.e.})$

$$r(x) = \begin{cases} 2, & \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) \in (0, \infty] \\ 1, & \text{exactly one of } \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) \text{ is nonzero} \\ 0, & \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) = 0 \end{cases}$$

## Case $lp \leftrightarrow lp$ . Local Spectral Multiplicity

Set  $(x \in \mathbb{R}, \mu\text{-a.e.})$

$$r(x) = \begin{cases} 2, & \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) \in (0, \infty] \\ 1, & \text{exactly one of } \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) \text{ is nonzero} \\ 0, & \frac{d\mu_{H+}}{d\mu}(x), \frac{d\mu_{H-}}{d\mu}(x) = 0 \end{cases}$$

### Theorem (Computation of the multiplicity function)

We have

- $E_{ac} \sim \mu_{ac}$  and  $N_H(x) = r(x)$ ,  $E_{ac}\text{-a.e.}$
- $E_{s,ac} \sim \mathbb{1}_{\{r(x)=2\}} d\mu_s$  and  $N_H(x) = 1$ ,  $E_{s,ac}\text{-a.e.}$
- $N_H(x) = 1$ ,  $E_{s,s}\text{-a.e.}$



## Case $lp \leftrightarrow lp$ . Simple Spectrum

### Corollary

*The singular spectrum of  $T_{\min}(H)$  is always simple.*

## Case $lp \leftrightarrow lp$ . Simple Spectrum

### Corollary

*The singular spectrum of  $T_{\min}(H)$  is always simple.*

### Theorem (Characterization of simplicity)

*The operator  $T_{\min}(H)$  has simple spectrum if and only if the set*

$$\begin{aligned} & \{x \in \mathbb{R} : \lim_{\epsilon \downarrow 0} \operatorname{Im} q_{H_+}(x + i\epsilon) \text{ exists in } (0, \infty)\} \\ & \cap \{x \in \mathbb{R} : \lim_{\epsilon \downarrow 0} \operatorname{Im} q_{H_-}(-x + i\epsilon) \text{ exists in } (0, \infty)\} \end{aligned}$$

*has Lebesgue measure zero.*

## Case $lp \leftrightarrow lp$ . Simple Spectrum

An explicit sufficient condition for simplicity is:

### Theorem

Assume that  $H_+$  has the Hilbert-Schmidt property, i.e., that there exists  $\phi \in [0, \pi)$  with

- $\int_{s_0}^{s_+} \xi_\phi^T H(t) \xi_\phi dt < \infty,$
- $\int_{s_0}^{s_+} \left( \int_{s_0}^t \xi_{\phi+\frac{\pi}{2}}^T H(u) \xi_{\phi+\frac{\pi}{2}} du \right) \xi_\phi^T H(t) \xi_\phi dt < \infty.$

Then the spectrum of  $T_{\min}(H)$  is simple.