


# ANOVA


# Analysis of Variance

- In the previous section we used Z- and t-Distributions to answer the question *“What is the probability that two samples come from the same population?”*




# Analysis of Variance

- In this section we introduce a new distribution – the F-Distribution
- Used to answer the question  
*“What is the probability that two samples come from populations that have the same variance?”*



# Analysis of Variance

- In this section we introduce a new distribution – the F-Distribution
- Can also answer the question *“What is the probability that **three or more** samples come from the same population?”*




# ANOVA

## Analysis of Variance

# ANOVA


- In the previous section we tested two samples to see if they likely came from the same parent population.
- What if we had three (or more) samples?
- Could we do the same thing?



# ANOVA

- Our null hypothesis would look like:

$$H_0: \mu_A = \mu_B = \mu_C$$




# ANOVA

- We *could* test each pair:

$$H_0: \mu_A = \mu_B \quad \alpha = 0.05$$

$$H_0: \mu_A = \mu_C \quad \alpha = 0.05$$

$$H_0: \mu_B = \mu_C \quad \alpha = 0.05$$





# ANOVA

- The problem is, our overall confidence drops:

$$H_0: \mu_A = \mu_B$$

$$\alpha = 0.05$$

$$.95 \times .95 \times .95 = .857$$


$$H_0: \mu_A = \mu_C$$

$$\alpha = 0.05$$

*85.7% confidence level*


$$H_0: \mu_B = \mu_C$$

$$\alpha = 0.05$$



# ANOVA


- This is where ANOVA comes in!
- We compute an **F value**, and compare it to a critical value determined by our **degrees of freedom** (the number of groups, and the number of items in each group)



# ANOVA

Let's work with some data:

GroupA	GroupB	GroupC
37	62	50
60	27	63
52	69	58
43	64	54
40	43	49
52	54	52
55	44	53
39	31	43
39	49	65
23	57	43



# ANOVA

First calculate the sample means

Next calculate the overall mean

	GroupA	GroupB	GroupC
	37	62	50
	60	27	63
	52	69	58
	43	64	54
	40	43	49
	52	54	52
	55	44	53
	39	31	43
n	39	49	65
	23	57	43
$\mu_{A,B,C}$			
$\mu_{TOT}$			

# ANOVA

ANOVA considers two types of **variance**:

## Between Groups

how far group means stray  
from the total mean

## Within Groups

how far individual values stray  
from their respective group mean

# ANOVA

The F value we're trying to calculate is simply the ratio between these two variances!

$$F = \frac{\textit{VarianceBetweenGroups}}{\textit{VarianceWithinGroups}}$$

# ANOVA

Recall the equation for variance:

$$s^2 = \frac{\Sigma(x - \bar{x})^2}{n - 1} = \frac{SS}{df}$$

Here  $\Sigma(x - \bar{x})^2$  is the “sum of squares” *SS*  
and  $n - 1$  is the “degrees of freedom” *df*

# ANOVA

So the formula for the F value becomes:

$$F = \frac{\text{VarianceBetweenGroups}}{\text{VarianceWithinGroups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

SSG = Sum of Squares Groups

SSE = Sum of Squares Error

$df_{groups}$  = degrees of freedom (groups)

$df_{error}$  = degrees of freedom (error)



# ANOVA

$$SSG = 420$$

Sum of Squares Groups

$$(\mu_A - \mu_{TOT})^2 = (44 - 49)^2 = 25$$

$$(\mu_B - \mu_{TOT})^2 = (50 - 49)^2 = 1$$

$$(\mu_C - \mu_{TOT})^2 = (53 - 49)^2 = 16$$

---

42

Multiply by the number of items in each group:

$$42 \times 10 = 420$$

	GroupA	GroupB	GroupC
	37	62	50
	60	27	63
	52	69	58
	43	64	54
	40	43	49
	52	54	52
	55	44	53
	39	31	43
	39	49	65
	23	57	43
$\mu_{A,B,C}$	44	50	53
$\mu_{TOT}$	49		

# ANOVA

$$SSG = 420$$

$$df_{groups} = 2$$

## Degrees of Freedom Groups

$$df_{groups} = n_{groups} - 1$$

$$= 3 - 1$$

$$= 2$$

	GroupA	GroupB	GroupC
	37	62	50
	60	27	63
	52	69	58
	43	64	54
	40	43	49
	52	54	52
	55	44	53
	39	31	43
	39	49	65
$\mu_{A,B,C}$			
$\mu_{TOT}$	49		

# ANOVA

$$SSG = 420$$

$$df_{groups} = 2$$

$$SSE = 3300$$

## Sum of Squares Error

$(x_A - \mu_A)^2$	$(x_A - \mu_A)^2$	$(x_B - \mu_B)^2$	$(x_B - \mu_B)^2$	$(x_C - \mu_C)^2$	$(x_C - \mu_C)^2$
49	64	144	16	9	1
256	121	529	36	100	0
64	25	361	361	25	100
1	25	196	1	1	144
16	441	49	49	16	100
1062			1742		496

TOTAL

3300

$$(37 - 44)^2$$

$$= (-7)^2$$

$$= 49$$

GroupA	GroupB	GroupC
37	62	50
60	27	63
52	69	58
43	64	54
40	43	49
52	54	52
55	44	53
39	31	43
39	49	65
23	57	43
$\mu_{A,B,C}$	50	53
$\mu_{TOT}$	49	

# ANOVA

Degrees of Freedom Error

$$\begin{aligned}
 df_{error} &= (n_{rows} - 1) * n_{groups} \\
 &= (10 - 1) * 3 \\
 &= 27
 \end{aligned}$$

	GroupA	GroupB	GroupC
	37	62	50
	60	27	63
	52	69	58
	43	64	54
	40	43	49
	52	54	52
	55	44	53
	39	31	43
	39	49	65
	23	57	43
$\mu_{A,B,C}$	44	50	53
$\mu_{TOT}$	49		

# ANOVA

$$\begin{aligned}
 SSG &= 420 \\
 df_{groups} &= 2 \\
 SSE &= 3300 \\
 df_{error} &= 27
 \end{aligned}$$

Plug these into our formula:

$$F = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}} = \frac{\frac{420}{2}}{\frac{3300}{27}} = \frac{210}{122.2} = \mathbf{1.718}$$

GroupA	GroupB	GroupC
37	62	50
60	27	63
52	69	58
43	64	54
40	43	49
52	54	52
55	44	53
39	31	43
39	49	65
23	57	43

$\mu_{A,B,C}$

$\mu_{TOT}$

# ANOVA with Excel Data Analysis

	A	B	C	D	E		
1	Anova: Single Factor						
2							
3	SUMMARY						
4	Groups	Count	Sum	Average	Variance		
5	GroupA	10	440	44	118		
6	GroupB	10	500	50	193.5555556		
7	GroupC	10	530	53	55.11111111		
8							
9							
10	ANOVA						
11	Source of Variation	SS	df	MS	F	P-value	F crit
12	Between Groups	420	2	210	1.718181818	0.198430533	3.354130829
13	Within Groups	3300	27	122.2222			
14							
15	Total	3720	29				
16							

Data Analysis

Analysis Tools

- Anova: Single Factor
- Anova: Two-Factor With Replication
- Anova: Two-Factor Without Replication
- Correlation
- Covariance
- Descriptive Statistics
- Exponential Smoothing
- F-Test Two-Sample for Variances
- Fourier Analysis
- Histogram


OK

Cancel

Help

# F Distribution

# F-Distribution





# F-Distribution

Look up our critical value from an F-table


use a table set for

95% confidence

find numerator df

find denominator df

critical value = 3.35

 denominator df		F-Table Upper Tail Area of 0.05				
		Numerator df				
		1	2	3	4	5
25	25	4.24	3.39	2.99	2.76	2.60
26	26	4.23	3.37	2.98	2.74	2.59
27	27	4.21	3.35	2.96	2.73	2.57
28	28	4.20	3.34	2.95	2.71	2.56
29	29	4.18	3.33	2.93	2.70	2.55
30	30	4.17	3.32	2.92	2.69	2.53

# F-Scores in MS Excel

- In Microsoft Excel, the following function returns an F-score:

$\alpha$	df1	df2	Formula	Output Value
0.05	2	27	=FINV(A2,B2,C2)	3.3541308285292

# F-Scores in Python

```
>>> from scipy import stats  
>>> stats.f.ppf(1-.05,dfn=2,dfd=27)  
3.3541308285291986
```

# ANOVA


Recall our null hypothesis:

$$H_0: \mu_A = \mu_B = \mu_C$$

Since  $F$  is less than  $F_{\text{critical}}$

$$1.718 < 3.354$$

we fail to reject the  
null hypothesis!



# ANOVA Exercise #1

- In an effort to receive faster payment of invoices, a company introduces two discount plans
- One set of customers is given a 2% discount if they pay their invoice early
- Another set is offered a 1% discount
- A third set is not offered any incentive



# ANOVA Exercise #1

- The results are as follows:
- Using ANOVA, can we say that the offers result in faster payments?



2% disc	1% disc	no disc
11	21	14
16	15	11
9	23	18
14	10	16
10	16	21

# ANOVA Exercise #1

1. Calculate the means



	2% disc	1% disc	no disc
	11	21	14
	16	15	11
	9	23	18
	14	10	16
	10	16	21
$\mu_{2,1,0}$	12	17	16
$\mu_{TOT}$	15		

$$SSG = 70$$



# ANOVA Exercise #1

## 2. Find Sum of Squares Groups

$$(\mu_2 - \mu_{TOT})^2 = (12 - 15)^2 = 9$$

$$(\mu_1 - \mu_{TOT})^2 = (17 - 15)^2 = 4$$

$$(\mu_0 - \mu_{TOT})^2 = (16 - 15)^2 = 1$$

---

14

Multiply by the number of items in each group:

$$14 \times 5 = 70$$

2% disc	1% disc	no disc
11	21	14
16	15	11
9	23	18
14	10	16
10	16	21

$\mu_{2,1,0}$

$\mu_{TOT}$



# ANOVA Exercise #1

$$SSG = 70$$

$$df_{groups} = 2$$



## 3. Degrees of Freedom Groups

$$df_{groups} = n_{groups} - 1$$

$$= 3 - 1$$

$$= 2$$

2% disc	1% disc	no disc
11	21	14
16	15	11
9	23	18
14	10	16
10	16	21

$\mu_{2,1,0}$	
$\mu_{TOT}$	

# ANOVA Exercise #1

$$SSG = 70$$

$$df_{groups} = 2$$

$$SSE = 198$$



## 4. Sum of Squares Error

$(x_2 - \mu_2)^2$	$(x_1 - \mu_1)^2$	$(x_0 - \mu_0)^2$
1	16	4
16	4	25
9	36	4
4	49	0
4	1	25
<b>34</b>	<b>106</b>	<b>58</b>
<b>TOTAL</b>		<b>198</b>

	2% disc	1% disc	no disc
	11	21	14
	16	15	11
	9	23	18
	14	10	16
	10	16	21
$\mu_{2,1,0}$	12	17	16
$\mu_{TOT}$	15		

# ANOVA Exercise #1

$$\begin{aligned}SSG &= 70 \\df_{groups} &= 2 \\SSE &= 198 \\df_{error} &= 12\end{aligned}$$



## 5. Degrees of Freedom Error

$$\begin{aligned}df_{error} &= (n_{rows} - 1) * n_{groups} \\&= (5 - 1) * 3 \\&= 12\end{aligned}$$

	2% disc	1% disc	no disc
	11	21	14
	16	15	11
	9	23	18
	14	10	16
	10	16	21
$\mu_{2,1,0}$	12	17	16
$\mu_{TOT}$	15		

# ANOVA Exercise #1

$$\begin{aligned}SSG &= 70 \\df_{groups} &= 2 \\SSE &= 198 \\df_{error} &= 12\end{aligned}$$



6. Calculate F value:

$$F = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}} = \frac{\frac{70}{2}}{\frac{198}{12}} = \frac{35}{16.5} = \mathbf{2.121}$$

7. Look up  $F_{critical}$  : **3.885**

	2% disc	1% disc	no disc
	11	21	14
	16	15	11
	9	23	18
	14	10	16
	10	16	21
$\mu_{2,1,0}$	12	17	16
$\mu_{TOT}$	15		


# ANOVA Exercise #1

$$\begin{aligned}SSG &= 70 \\df_{groups} &= 2 \\SSE &= 198 \\df_{error} &= 12\end{aligned}$$




Since  $F$  falls to the left of  $F_{\text{critical}}$   
 $2.121 < 3.885$

we fail to reject the  
null hypothesis!



# ANOVA Exercise #1

We don't have enough to support the idea that our offers changed the average number of days that customers took to pay their invoices!



# Two-Way ANOVA

# One-Way vs Two-Way ANOVA

- In the previous examples we used one-way ANOVA to test one independent variable.
- For the invoice problem, the independent variable was the incentive offered.
- The dependent variable was the time it took to receive payment.



# One-Way vs Two-Way ANOVA

- Two-Way ANOVA lets us test two independent variables at the same time
- For the invoice example, we might also consider the amount due
- We would have 3 invoices for \$50, 3 for \$100, etc. and offer different incentives at each dollar amount.

# One-Way vs Two-Way ANOVA

- The resulting data might look like this:
- Here, each row or dollar amount is called a **block**.
- Essentially, we want to isolate and remove any variance contributed by the blocks, to better understand the variance in the groups.

	2% disc	1% disc	no disc
\$50	16	23	21
\$100	14	21	16
\$150	11	16	18
\$200	10	15	14
\$250	9	10	11

# One-Way vs Two-Way ANOVA

- So how do we do that?

	2% disc	1% disc	no disc
\$50	16	23	21
\$100	14	21	16
\$150	11	16	18
\$200	10	15	14
\$250	9	10	11

# Two-Way ANOVA

- The goal of ANOVA is to separate different aspects of the total variance.
- In the previous examples we had only

The diagram shows a data table for a Two-Way ANOVA. It has two columns for groups and one row for the total mean. The first column is labeled 'Group 1' (orange header) and the second 'Group 2' (green header). The first row contains the values 8 and 11. The second row contains 10 and 12. The third row contains 12 and 13. To the left of the first column is a blue box labeled  $\mu_{1,2}$  with a bracket connecting it to the values 10 and 12. To the right of the second column is a blue box labeled  $\mu_{TOT}$  with a bracket connecting it to the values 12 and 11. Below the table, a red bracket spans the width of the two columns.

Group 1	Group 2
8	11
10	12
12	13
$\mu_{1,2}$ 10	12 $\mu_{TOT}$ 11

**Sum of Squares Groups (SSG)**

» between groups


and **Sum of Squares Error (SSE)**

» within groups

# Two-Way ANOVA

- These two variances **SSG** and **SSE** add up to our total variance

Sum of Squares Total (SST)



Sum of Squares Groups (SSG)

» between groups

and Sum of Squares Error (SSE)

» within groups

# Two-Way ANOVA

- Now we'll look at variance between rows, or blocks

	Group 1	Group 2	
Block A	8	11	
Block B	10	12	
Block C	12	13	$\mu_{TOT}$
$\mu_{1,2}$	10	12	11

Sum of Squares Groups (SSG) » between groups  
and Sum of Squares Error (SSE) » within groups

# Two-Way ANOVA

- First calculate the block means

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

- Then calculate the  
**Sum of Squares Blocks (SSB)** » between blocks  
**Sum of Squares Groups (SSG)** » between groups  
and **Sum of Squares Error (SSE)** » within groups

# Two-Way ANOVA

- ANOVA still considers the relationship between the SSG and the SSE

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

Sum of Squares Blocks (SSB)

» between blocks

Sum of Squares Groups (SSG)

» between groups

and Sum of Squares Error (SSE)

» within groups



# Two-Way ANOVA

- By calculating the SSB, we remove some of the variance in SSE

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

Sum of Squares Blocks (SSB)

» between blocks

Sum of Squares Groups (SSG)

» between groups

and Sum of Squares Error (SSE)

» within groups

# Two-Way ANOVA

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

## Sum of Squares Groups (SSG)

$$(\mu_1 - \mu_{TOT})^2 = (10 - 11)^2 = 1$$

$$(\mu_2 - \mu_{TOT})^2 = (12 - 11)^2 = 1$$

2

multiply by the number  
of items in each group:  $2 \times 3 = 6$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

**SSG = 6**

# Two-Way ANOVA

## Sum of Squares Blocks (SSB)

$$(\mu_A - \mu_{TOT})^2 = (9.5 - 11)^2 = 2.25$$

$$(\mu_B - \mu_{TOT})^2 = (11 - 11)^2 = 0$$

$$(\mu_C - \mu_{TOT})^2 = (12.5 - 11)^2 = 2.25$$

4.5

multiply by the number  
of items in each block:

$$4.5 \times 2 = 9$$

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	
Block B	10	12	
Block C	12	13	
$\mu_{1,2}$			11

$SSG = 6$

$SSB = 9$

# Two-Way ANOVA

## Sum of Squares Total (SST)

$$\begin{aligned} & (8 - 11)^2 + (11 - 11)^2 + \\ & (10 - 11)^2 + (12 - 11)^2 + \\ & (12 - 11)^2 + (13 - 11)^2 = 16 \end{aligned}$$

no need to multiply since  
every item is represented

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{\text{groups}}}}{\frac{SSE}{df_{\text{error}}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

# Two-Way ANOVA

## Sum of Squares Error (SSE)

$$\begin{aligned}SSE &= SST - SSG - SSB \\ &= 16 - 6 - 9 = 1\end{aligned}$$

no need to multiply since we're working with totals already

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{\text{groups}}}}{\frac{SSE}{df_{\text{error}}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$SSG = 6$   
 $SSB = 9$   
 $SST = 16$   
 $SSE = 1$

# Two-Way ANOVA

So how do we calculate F?

Degrees of Freedom Groups is unchanged:

$$\begin{aligned} df_{groups} &= n_{groups} - 1 \\ &= 2 - 1 \\ &= 1 \end{aligned}$$

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

$$SSE = 1$$

$$df_{groups} = 1$$

# Two-Way ANOVA

So how do we calculate F?

Degrees of Freedom Error has changed:

$$\begin{aligned} df_{error} &= (n_{blocks} - 1)(n_{groups} - 1) \\ &= (3 - 1)(2 - 1) \\ &= 2 \end{aligned}$$

$$F = \frac{Var. Between Groups}{Var. Within Groups} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

$$SSE = 1$$

$$df_{groups} = 1$$

$$df_{error} = 2$$

# Two-Way ANOVA

So how do we calculate F?

$$F = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}} = \frac{\frac{6}{1}}{\frac{1}{2}} = 12$$

$$F = \frac{\frac{Var. Between Groups}{SSE}}{\frac{df_{groups}}{df_{error}}} = \frac{SSG}{SSE} \cdot \frac{df_{error}}{df_{groups}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

$$SSE = 1$$

$$df_{groups} = 1$$

$$df_{error} = 2$$



# Two-Way ANOVA

$F_{groups} = 12$  feels like a high value.

However, in a two-way ANOVA,  $F_{critical}$  is found for groups and blocks separately!

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

$$SSE = 1$$

$$df_{groups} = 1$$

$$df_{error} = 2$$

# Two-Way ANOVA

$F_{groups} = 12$  feels like a high value.

For groups, with 1df in the numerator and 2 df in the denominator,

$$F_{critical} = 18.5$$

$$F = \frac{\text{Var. Between Groups}}{\text{Var. Within Groups}} = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}}$$

	Group 1	Group 2	$\mu_{A,B,C}$
Block A	8	11	9.5
Block B	10	12	11
Block C	12	13	12.5
$\mu_{1,2}$	10	12	11

$$SSG = 6$$

$$SSB = 9$$

$$SST = 16$$

$$SSE = 1$$

$$df_{groups} = 1$$

$$df_{error} = 2$$

# ANOVA Exercise #2

- Let's go back to the invoice problem, and add a new independent variable
- Here each **block** represents an invoice amount
- The dependent variable is still days elapsed until payment

	2% disc	1% disc	no disc
\$50	16	23	21
\$100	14	21	16
\$150	11	16	18
\$200	10	15	14
\$250	9	10	11



# ANOVA Exercise #2

1. Calculate the group means,  
the block means,  
and the total mean



	2% disc	1% disc	no disc	$\mu_{\text{block}}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{\text{col}}$				15

# ANOVA Exercise #2

## 2. Sum of Squares Groups

$$(\mu_2 - \mu_{TOT})^2 = (12 - 15)^2 = 9$$

$$(\mu_1 - \mu_{TOT})^2 = (17 - 15)^2 = 4$$

$$(\mu_0 - \mu_{TOT})^2 = (16 - 15)^2 = 1$$

---

14

Multiply by the number of items in each group:

$$14 \times 5 = 70$$




	2% disc	1% disc	no disc	$\mu_{\text{block}}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{\text{col}}$				15

$$SSG = 70$$

# ANOVA Exercise #2

## 3. Degrees of Freedom Groups

$$\begin{aligned}df_{groups} &= n_{groups} - 1 \\&= 3 - 1 \\&= 2\end{aligned}$$



	2% disc	1% disc	no disc	$\mu_{block}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{col}$				15

$SSG = 70$

$df_{groups} = 2$

# ANOVA Exercise #2

## 4. Sum of Squares Blocks

$$(\mu_{50} - \mu_{TOT})^2 = (20 - 15)^2 = 25$$

$$(\mu_{100} - \mu_{TOT})^2 = (17 - 15)^2$$

$$(\mu_{200} - \mu_{TOT})^2 = (15 - 15)^2 = 0$$

$$(\mu_{200} - \mu_{TOT})^2 = (13 - 15)^2 = 4$$

$$(\mu_{250} - \mu_{TOT})^2 = (10 - 15)^2 = 25$$

$$\begin{matrix} TO \\ 58 \times 3 = 174 \end{matrix}$$

58

$SSG = 70$   
 $SSB = 174$

$df_{groups} = 2$


	2% disc	1% disc	no disc	$\mu_{block}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{col}$				15



# ANOVA Exercise #2

## 5. Sum of Squares Total

$(x_2 - \mu_{tot})^2$	$(x_1 - \mu_{tot})^2$	$(x_0 - \mu_{tot})^2$
1	64	36
1	36	1
16	1	9
25	0	1
36	25	16
<b>79</b>	<b>126</b>	<b>63</b>
<b>TOTAL</b>		<b>268</b>



	2% disc	1% disc	no disc	$\mu_{block}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{col}$				<b>15</b>

$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$


$$df_{groups} = 2$$



# ANOVA Exercise #2

## 6. Sum of Squares Error

$$\begin{aligned}SSE &= SST - SSG - SSB \\ &= 268 - 70 - 174 = 24\end{aligned}$$



	2% disc	1% disc	no disc	$\mu_{\text{block}}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{\text{col}}$				15

$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$


$$SSE = 24$$

$$df_{\text{groups}} = 2$$

# ANOVA Exercise #2

## 7. Degrees of Freedom Error

$$\begin{aligned}
 df_{error} &= (n_{blocks} - 1)(n_{groups} - 1) \\
 &= (5 - 1)(3 - 1) \\
 &= 8
 \end{aligned}$$



	2% disc	1% disc	no disc	$\mu_{block}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{col}$				15

$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$

$$SSE = 24$$


$$df_{groups} = 2$$

$$df_{error} = 8$$

# ANOVA Exercise #2

## 8. Calculate F

$$F = \frac{\frac{SSG}{df_{groups}}}{\frac{SSE}{df_{error}}} = \frac{\frac{70}{2}}{\frac{24}{8}} = \frac{35}{3} = \mathbf{11.67}$$



	2% disc	1% disc	no disc	$\mu_{block}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{col}$				15

$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$

$$SSE = 24$$

$$df_{groups} = 2$$

$$df_{error} = 8$$

$$\mathbf{F = 11.67}$$

# ANOVA Exercise #2

9. Find  $F_{\text{critical}}$

$$\alpha = 0.05$$

$$df_{\text{numerator}} = 2$$

$$df_{\text{denominator}} = 8$$

$$F_{\text{critical}} = 4.46$$



	2% disc	1% disc	no disc	$\mu_{\text{block}}$
\$50	16	23	21	
\$100	14	21	16	
\$150	11	16	18	
\$200	10	15	14	
\$250	9	10	11	
$\mu_{\text{col}}$				15

$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$

$$SSE = 24$$

$$df_{\text{groups}} = 2$$

$$df_{\text{error}} = 8$$


$$F = 11.67$$

$$F_{\text{critical}} = 4.46$$

# ANOVA Exercise #1

Since  $F$  falls to the right of  $F_{\text{critical}}$   
 $4.46 < 11.67$

we reject the null hypothesis!



$$SSG = 70$$

$$SSB = 174$$

$$SST = 268$$

$$SSE = 24$$

$$df_{\text{groups}} = 2$$


$$df_{\text{error}} = 8$$

$$F = 11.67$$

$$F_{\text{critical}} = 4.46$$

# 2-way ANOVA in Excel

	A	B	C	D	E	F	G	H	I	J	K	L
1	Anova: Two-Factor Without Replication											
2												
3	SUMMARY	Count	Sum	Average	Variance							
4	Row 1	3	60	20	13							
5	Row 2	3	51	17	13							
6	Row 3	3	45	15	13							
7	Row 4	3	39	13	7							
8	Row 5	3	30	10	1							
9												
10	Column 1	5	60	12	8.5							
11	Column 2	5	85	17	26.5							
12	Column 3	5	80	16	14.5							
13												
14												
15	ANOVA											
16	Source of Variation	SS	df	MS	F	P-value	F crit					
17	Rows	174	4	43.5	14.5	0.000974668	3.837853355					
18	Columns	70	2	35	11.666667	0.004249458	4.458970108					
19	Error	24	8	3								
20												
21	Total	268	14									



# Two-Way ANOVA with Replication

# Without vs With Replication

without replication

	GroupA	GroupB	GroupC
Block1	16	23	21
Block2	14	21	16
Block3	11	16	18
Block4	10	15	14
Block5	9	10	11
Block6	8	8	10

with replication

	GroupA	GroupB	GroupC
Block1	16	23	21
	14	21	16
	11	16	18
Block2	10	15	14
	9	10	11
	8	8	10

Samples have multiple values  
Samples have a mean value



# Two-Way ANOVA with Replication

- Introduces the concept of **sample means** and **sample variance**
- Introduces the concept of **interactions**

# Two-Way ANOVA with Replication

- As with our previous 2-way ANOVA, we consider two independent variables organized into groups and blocks
- We sample every block/group combination
- With replication, block/group samples have multiple measurements

# Two-Way ANOVA with Replication

- Consider an experiment that measures the height of plants
- We apply three types of fertilizer A, B & C – these are our Groups
- Plants are kept at two temperatures (warm & cold) – these are our Blocks
- We assign 3 plants to each sample



# Two-Way ANOVA

- First calculate the mean for each 3-item sample
- Calculate column means
- Calculate block means
- Calculate the overall mean

Fertilizer:				
	A	B	C	
Warm	13	21	18	Block Means
	14	19	15	
	12	17	15	
Cold	16	14	15	Means
	18	11	13	
	17	14	8	
Sample Means	13	19	16	
	17	13	12	
Column Means	15	16	14	15

# Two-Way ANOVA

- As before, calculate the Sum of Squares Blocks

$$(\textcolor{red}{16} - \textcolor{green}{15})^2 + (\textcolor{blue}{14} - \textcolor{green}{15})^2 = 2$$

$$\times 9 \text{ items per block} = \textcolor{black}{18}$$

Fertilizer:	A	B	C	
Warm	13	21	18	<b>16</b>
	14	19	15	
	12	17	15	
Cold	16	14	15	<b>14</b>
	18	11	13	
	17	14	8	

Block Means

Sample Means	13	19	16
	17	13	12

Column Means	<b>15</b>	<b>16</b>	<b>14</b>	<b>15</b>
--------------	-----------	-----------	-----------	-----------

$$\text{SSB} = 18$$

# Two-Way ANOVA

- As before, calculate the Sum of Squares Columns

$$\begin{aligned}
 & (15 - 15)^2 + (16 - 15)^2 + \\
 & (14 - 15)^2 = 2 \\
 & \times 6 \text{ items per column} = 12
 \end{aligned}$$

Fertilizer:	A	B	C		
Warm	13	21	18	16	Block
	14	19	15		
	12	17	15		
Cold	16	14	15	14	Means
	18	11	13		
	17	14	8		

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
			15

SSB = 18    SSC = 12

# Two-Way ANOVA

- As before, calculate the Degrees of Freedom Columns

$$df_{columns} = (3-1) = 2$$

Fertilizer:	A	B	C	
Warm	13	21	18	Block Means
	14	19	15	
	12	17	15	
Cold	16	14	15	Block Means
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
			15

$$SSB = 18 \quad SSC = 12 \quad df_{columns} = 2$$

# Two-Way ANOVA

- We have a new statistic: SS Interactions
- For each sample mean, subtract the matching block and column means, add back the overall mean, square the result

Fertilizer:	A	B	C		
Warm	13	21	18	16	Block Means
	14	19	15		
	12	17	15		
Cold	16	14	15	14	Block Means
	18	11	13		
	17	14	8		

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
			15

SSB = 18    SSC = 12     $df_{\text{columns}} = 2$



# Two-Way ANOVA

$$\begin{aligned}
 & (13 - 16 - 15 + 15)^2 + \\
 & (19 - 16 - 16 + 15)^2 + \\
 & (16 - 16 - 14 + 15)^2 + \\
 & (17 - 14 - 15 + 15)^2 + \\
 & (13 - 14 - 16 + 15)^2 + \\
 & (12 - 14 - 14 + 15)^2 = 28
 \end{aligned}$$

$\times 3 \text{ items per sample} = 84$

Fertilizer:	A	B	C	Block Means
Warm	13	21	18	
	14	19	15	
	12	17	15	
Cold	16	14	15	14
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
			15

SSB = 18    SSC = 12     $df_{\text{columns}} = 2$   
 SSI = 84

# Two-Way ANOVA

- Calculate the Sum of Squares Total

4	36	9	
1	16	0	
9	4	0	
1	1	0	
9	16	4	
4	1	49	164

Fertilizer:	A	B	C	
Warm	13	21	18	<b>16</b>
	14	19	15	
	12	17	15	
Cold	16	14	15	<b>14</b>
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12

Column Means	<b>15</b>	<b>16</b>	<b>14</b>	<b>15</b>
--------------	-----------	-----------	-----------	-----------

$$SSB = 18 \quad SSC = 12 \quad df_{\text{columns}} = 2$$

$$SSI = 84$$

$$SST = 164$$

# Two-Way ANOVA

- Calculate the Sum of Squares Error by subtracting the other values from the SST:  
 $164 - 18 - 12 - 84 = 50$

Fertilizer:	A	B	C	Block Means
Warm	13	21	18	
	14	19	15	
	12	17	15	
Cold	16	14	15	14
	18	11	13	
	17	14	8	

Sample Means	13	19	16	
	17	13	12	
Column Means	15	16	14	15

$$\begin{aligned}
 \text{SSB} &= 18 & \text{SSC} &= 12 & \text{df}_{\text{columns}} &= 2 \\
 \text{SSI} &= 84 & \text{SSE} &= 50 & & \\
 \text{SST} &= 164 & & & & 
 \end{aligned}$$

# Two-Way ANOVA

- Degrees of Freedom Error

$$blocks \times columns \times (items - 1)$$

$$= 2 \times 3 \times (3 - 1) = 12$$

Fertilizer:	A	B	C	
Warm	13	21	18	Block Means
	14	19	15	
	12	17	15	
Cold	16	14	15	Block Means
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
			15

$$SSB = 18 \quad SSC = 12 \quad df_{columns} = 2$$

$$SSI = 84 \quad SSE = 50 \quad df_{error} = 12$$

$$SST = 164$$

# Two-Way ANOVA

- Calculate F

$$F = \frac{\frac{SSC}{df_{columns}}}{\frac{SSE}{df_{error}}} = \frac{\frac{12}{2}}{\frac{50}{12}} = 1.44$$

Fertilizer:	A	B	C	
Warm	13	21	18	16
	14	19	15	
	12	17	15	
Cold	16	14	15	14
	18	11	13	
	17	14	8	
Sample Means	13	19	16	
Means	17	13	12	
Column Means	15	16	14	15

SSB = 18    SSC = 12     $df_{columns} = 2$   
 SSI = 84    SSE = 50     $df_{error} = 12$   
 SST = 164

# Two-Way ANOVA

$$F = 1.44$$

- Look up  $F_{\text{critical}}$

$$F_{(0.05, 2, 12)} = 3.885$$

Fertilizer:	A	B	C	Block Means
Warm	13	21	18	
	14	19	15	
	12	17	15	
Cold	16	14	15	
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14

15


$$SSB = 18 \quad SSC = 12 \quad df_{\text{columns}} = 2$$

$$SSI = 84 \quad SSE = 50 \quad df_{\text{error}} = 12$$

$$SST = 164$$

# Two-Way ANOVA

- A look at Interaction:



Fertilizer:	A	B	C	Block Means
Warm	13	21	18	
	14	19	15	
	12	17	15	
Cold	16	14	15	14
	18	11	13	
	17	14	8	

Sample Means	13	19	16
	17	13	12
Column Means	15	16	14
	15		

$$SSB = 18 \quad SSC = 12 \quad df_{\text{columns}} = 2$$

$$SSI = 84 \quad SSE = 50 \quad df_{\text{error}} = 12$$

$$SST = 164$$

# 2-way with Replication in Excel

	A	B	C	D	E	F	G	H	I	J	K	L
1	Anova: Two-Factor With Replication											
2												
3	SUMMARY	Fertilizer A	Fertilizer B	Fertilizer C	Total							
4	WARM											
5	Count	3	3	3	9							
6	Sum	39	57	48	144							
7	Average	13	19	16	16							
8	Variance	1	4	3	8.75							
9												
10	COLD											
11	Count	3	3	3	9							
12	Sum	51	39	36	126							
13	Average	17	13	12	14							
14	Variance	1	3	13	9.5							
15												
16	Total					ANOVA						
17	Count	6	6	6		Source of Variation	SS	df	MS	F	P-value	F crit
18	Sum	90	96	84		Sample	18	1	18	4.32	0.059785686	4.747225347
19	Average	15	16	14		Columns	12	2	6	1.44	0.275086887	3.885293835
20	Variance	5.6	13.6	11.2		Interaction	84	2	42	10.08	0.002698928	3.885293835
21						Within	50	12	4.16667			

