High Harmonic Generation in Laser Plasma Interaction

Kulwinder Kaur (2021PHS7190) Harikesh Kushwaha (2021PHS7181)

Adviser: Prof. Vikrant Saxena


Indian Institute of Technology, Delhi




Introduction

Interaction of light with matter at ultra high light intensity gives access to novel physical regimes which are barely, if at all, explored in lab.

- Intensity of $10^{23} W/cm^{-2}$ has been reached experimentally.¹
- QED at $I = 10^{25} W/cm^{-2}$. Schwinger field at $I = 10^{29} W/cm^{-2}$.
- Plasma is overdense if $\omega < \omega_p$.
- Harmonics are generated by interaction of laser with overdense plasma.³





¹Henri Vincenti 10.1103/physrevlett.123.105001

² Jin Woo Yoon et al 10.1364/OPTICA.420520

³R. Lichters et al 10 . 1063 / 1 . 871619

Summary of Work Done in Previous Semester

- Interaction of highly intense laser pulse with overdense and underdense plasma
- Change in effective critical density of plasma for relativistic laser pulse
- The oscillations of plasma surface.
 - Oscillations increases with increase in intensity
 - Surface oscillations have even harmonics
- Study of high harmonics generation in laser plasma interaction (normal incidence)
 - Only odd harmonics are generated
 - A resonance at $n_0/n_c=4$ is also observed
 - Increasing intensity and pulse duration increases number of harmonics
 - No effect of the envelopes

What Now?


- Effect of Super Gaussian (SG) envelopes on the generated high harmonics
- Oblique incident and p- and s-polarized laser
- Selection rule for p- and s-polarized laser

Simulation Details

We want to study the effect of super Gaussian envelope on the generated high harmonics. We performed some simulations in 1D3V. Here are some parameters:


- Particles per cell: 100
- Number of cells: 16000
- Pulse duration = 20τ ($\tau \approx 3.3 \text{fs}$)
- Simulation time = 40τ
- Wavelength $\lambda_I = 1 \mu m$
- Intensity of laser for $a_0 = 1$ is $I = 1.37 \times 10^{18} W/cm^2$

We also performed simulations with p- and s-polarized laser incidence at oblique angle.




Oblique Incidence: Transformations

 We follow Bourdier⁴ to make a transformation which lets us simulate oblique incidence in 1D.



⁴Bourdier, A. 10 . 1063 / 1.864355

p- and s- Polarized Laser: Selection Rule



Results: SG Envelope




Figure: The spectrum of SG envelopes with power 2,4,6, and 8 is shown in a sigle plot. A small increase in the peak amplitude is observed with increasing power.

7 / 10

Results: p-Polarized Laser




Figure: Spectrum of HHG for p-polarized light. Simulation parameters are $\alpha = \pi/6$, the denity is $n_0 = 7n_c$ and $a_0 = 4$

Results: s-Polarized Laser




Figure: Spectrum of HHG for s-polarized light. Simulation parameters are $\alpha=\pi/4$, the denity is $n_0=7n_c$ and $a_0=4$

Current Status and Future Plan of Work

Current Status

- A brief overview of HHG generation in laser plasma interaction
- SG envelopes has very little effect on the generated harmonics
- For p-polarized laser, even and odd p-polarized harmonics.
- For s-polarized laser, odd s-polarized hyarmonics and even p-polarized harmonics.

Future Plan of Work

- Study oblique incidence and polarization more regously.
- Do 2D simulations.
- Compare it with the 1D results.

Acknowledgement

We would like to extend our sincerest gratitude to Professor Vikrant Saxena for his unwavering support, patience, motivation, enthusiasm, and invaluable guidance.