

# Group 8 - GLM Exercises

November 10, 2015

## 1. A density that belongs in the one-parameter exponential family has the following (canonical) representation:

$$p(z) = e^{(q[z\theta + c(\theta)] + h(z, q))}$$

where  $\theta$  is called the natural parameter, and  $c$  and  $h$  are functions whose exact form depends on the particular density. For many members of this family,  $q > 0$ , is a further unknown parameter, often called a precision parameter.

1. The following densities belong to the exponential family. Identify  $\theta$ ,  $q$  and an appropriate  $c(\theta)$ , for each of them (take into account that for some  $z$  might be a transformation of  $t$ ):

**Normal**

$$\begin{aligned} t &\sim N(\mu, -q) = \exp\left\{\frac{-(t-\mu)^2}{2q}\right\} \\ &= \exp\left\{\frac{-(t^2 - 2\mu t + \mu^2)}{2q}\right\} \\ &= \exp\left\{-\frac{1}{2q}\left[t^2 - 2\mu t + \mu^2\right]\right\} \\ &= \exp\left\{-\frac{1}{q}\left[\frac{t^2}{2} - \mu t + \frac{\mu^2}{2}\right]\right\} \\ &= \exp\left\{\frac{1}{q}\left[-\frac{t^2}{2} + \mu t - \frac{\mu^2}{2}\right]\right\} \\ &= \exp\left\{\frac{1}{q}\left[\mu t - \frac{\mu^2}{2}\right] - \frac{t^2}{2q}\right\} \end{aligned}$$

**Result:**

$$\theta = \mu$$

$q = q$  (since  $q^{-1}$  is the variance parameter passed, than  $q$  the parameter is the precision and the  $q$  above is the variance for the normal distribution)

$$c(\theta) = \frac{\mu^2}{2}$$

**Bernoulli**

$$\begin{aligned} t &\sim \text{Bern}(n, p) = p^z(1-p)^{1-z} \\ &= \exp\left\{\log(p^z) + \log((1-p)^{1-z})\right\} \end{aligned}$$

$$\begin{aligned}
&= \exp \left\{ z \log(p) + (1-z) \log(1-p) \right\} \\
&= \exp \left\{ z \log(p) + \log(1-p) - z \log(1-p) \right\} \\
&= \exp \left\{ z \log\left(\frac{p}{1-p}\right) + \log(1-p) \right\}
\end{aligned}$$

**Result:**

$$\theta = \log\left(\frac{p}{1-p}\right)$$

$$q = 1$$

$$c(\theta) = -\log(1-p)$$

**Binomial**

$$\begin{aligned}
t &\sim \text{Bin}(n, p) = \binom{n}{k} p^k (1-p)^{n-k} \\
&= \exp \left\{ \log\left(\binom{n}{k}\right) + \log(p^k) + \log((1-p)^{n-k}) \right\} \\
&= \exp \left\{ \log\left(\binom{n}{k}\right) + k \log(p) + (n-k) \log(1-p) \right\} \\
&= \exp \left\{ \log\left(\binom{n}{k}\right) + k \log\left(\frac{p}{1-p}\right) + n \log(1-p) \right\} \\
&= \exp \left\{ n \left[ \frac{k}{n} \log\left(\frac{p}{1-p}\right) + \log(1-p) \right] + \log\left(\binom{n}{k}\right) \right\}
\end{aligned}$$

**Result:**

$$\theta = \log\left(\frac{p}{1-p}\right)$$

$$q = n$$

$$c(\theta) = \log\left(\binom{n}{k}\right)$$

**Poisson**

$$\begin{aligned}
t &\sim \text{Poisson}(\lambda) = \frac{\lambda^k e^{-\lambda}}{k!} \\
&= \exp \left\{ \log(\lambda^k e^{-\lambda}) - \log(k!) \right\} \\
&= \exp \left\{ k * \log(\lambda) - \lambda \log(e) - \log(k!) \right\} \\
&= \exp \left\{ k * \log(\lambda) - \lambda - \log(k!) \right\}
\end{aligned}$$

**Result:**

$$\theta = \log(\lambda)$$

$$q = 1$$

$$c(\theta) = -\lambda$$

**2. Identify the canonical link function for each of the models given above.**

**Normal**

$$\mu$$

**Bernoulli**

$$\text{logit}(\mu)$$

**Binomial**

$$\text{logit}(\mu)$$

**Poisson**

$$\log(\mu)$$

**3. Consider now the generalised linear model:**

$$t_n|x_n \sim \text{NDEF}(\theta(x_n, w), q\gamma_n) \text{ with}$$

$$\theta(x_n, w) = (c')^{-1}(g^{-1}(\phi(x_n)^T w)) =: f(\phi(x_n)^T w)$$

We know the log-likelihood of the NDEF family of distributions takes the form:

$$2\log(p(t|x, \gamma, w, q)) = 2q \sum_n \gamma_n \left[ t_n \theta(x_n, w) - c(\theta(x_n, w)) \right]$$

We will prove it is log-concave if it is the negative of a log-convex function, so will prove:

$$f(\theta(x_n, w), q\gamma_n) = -2q \sum_n \gamma_n \left[ t_n \theta(x_n, w) - c(\theta(x_n, w)) \right]$$

is log-convex, taking into account the link function constructed as:

$$g(y) = c'^{-1}(\theta) \Rightarrow \theta(x_n, w) = \phi(x_n)^T w$$

We will prove it is log-convex by showing the second derivative is positive semi-definite for all  $\theta(x_n, w)$

The first derivative of the negative log likelihood is:

$$f'(\theta(x_n, w), q\gamma_n) = -2q \left[ \sum_n \gamma_n t_n \phi(x_n)^T - \sum_n \gamma_n c'(\phi(x_n)^T w) \right]$$

$$f''(\theta(x_n, w), q\gamma_n) = 2q \sum_n \gamma_n c''(\phi(x_n)^T w)$$

We know that  $c''$  is the variance function and  $\gamma_n$  and  $q$  are positive by definition. So the second derivative is always positive and therefore:

$$x^T f'' x \geq 0 \text{ for all } x.$$

## 2. $R^2$ and deviance

**1. Show that for any linear regression model:**

$$-2\log p(t|X, w_{MLE}, q_{MLE}) = N \log e^T e + \text{const}$$

where “const” does not depend on  $M$  or  $X$ .

We know that:

$$w_{mle} = (\phi^T \phi)^{-1} \phi^T t$$

and

$$q_{mle} = \left( \frac{1}{N} e^T e \right)^{-1}$$

We use this to solve for the deviance of the maximum likelihood function:

$$\begin{aligned} -2\log p(t|X, w_{mle}, q_{mle}) &= -N\log(q_{mle}) + q_{mle} e^T e \\ -N\log(q_{mle}) + q_{mle} e^T e &= -N\log\left(\left(\frac{1}{N} e^T e\right)^{-1}\right) + \left(\frac{1}{N} e^T e\right)^{-1} e^T e \\ &= N\log\left(\frac{1}{N} e^T e\right) + N \\ &= N[\log(e^T e) - \log(N)] + N \\ &= N\log(e^T e) - N(\log(N) + 1) \\ &= N\log(e^T e) + \text{const} \end{aligned}$$

**2. Show that in the null model,**

$$w_{0,MLE} = \bar{t}$$

We know from **1**:

$$w_{mle} = (\phi^T \phi)^{-1} \phi^T t$$

In the case of the null model,  $\phi$  is a vector of ones so this becomes:

$$\frac{1}{N} \sum_{n=1}^N t_n = \bar{t}$$

**3. The null model is nested within the saturated model, and it corresponds to the special case where  $w_1 = \dots w_M = 0$ . Let  $D_0$  be the deviance of the null model and  $D_1$  be that of the saturated model. Show that:**

$$D_0 - D_1 = -N\log(1 - R^2)$$

**where  $R^2$  is the coefficient  $R^2$  for the saturated model.**

We know from **part 1** that:

$$-2\log p(t|X, w_{mle}, q_{mle}) = N\log(e^T e) + \text{const}$$

Which also what we equate as  $D_M$  for model  $M$ , so:

$$\begin{aligned} D_0 - D_1 &= N\log(e_0^T e_0) - N\log(e_1^T e_1) \\ &= N\log\left(\frac{e_0^T e_0}{e_1^T e_1}\right) \end{aligned}$$

We use the following properties:

$$e = t - \hat{t}, \text{ and,}$$

$$R^2 = 1 - \frac{\sum_i (t_i - \hat{t}_i)^2}{\sum_i (t_i - \bar{t})^2}$$

$$1 - R^2 = \frac{\sum_i (t_i - \hat{t}_i)^2}{\sum_i (t_i - \bar{t})^2}$$

To show that

$$= N\log\left(\frac{e_0^T e_0}{e_1^T e_1}\right)$$

$$\begin{aligned}
&= N \log \left( \frac{\sum_i (t_i - \bar{t})^2}{\sum_i (t_i - \hat{t}_i)^2} \right) \\
&= N \log(\sum_i (t_i - \bar{t})^2) - N \log(\sum_i (t_i - \hat{t}_i)^2) \\
&= -N \log \left( \frac{\sum_i (t_i - \hat{t}_i)^2}{\sum_i (t_i - \bar{t})^2} \right) \\
&= -N \log(1 - R^2)
\end{aligned}$$