

Group 8 - GLM Exercises

November 10, 2015

1. A density that belongs in the one-parameter exponential family has the following (canonical) representation:

$$p(z) = \dots$$

where θ is called the natural parameter, and c and h are functions whose exact form depends on the particular density. For many members of this family, $q > 0$, is a further unknown parameter, often called a precision parameter.

1. The following densities belong to the exponential family. Identify θ , q and an appropriate $c(\theta)$, for each of them (take into account that for some z might be a transformation of t):

Normal

[ADD ME]

Bernoulli

$$t \sim \text{Bern}(n, p) = p^z (1-p)^{1-z}$$

$$\exp \left\{ \log(p^z) + \log((1-p)^{1-z}) \right\}$$

$$\exp \left\{ z \log(p) + (1-z) \log(1-p) \right\}$$

$$\exp \left\{ z \log(p) + \log(1-p) - z \log(1-p) \right\}$$

$$\exp \left\{ z \log\left(\frac{p}{1-p}\right) + \log(1-p) \right\}$$

Result:

$$\theta = \log\left(\frac{p}{1-p}\right)$$

$$q = 1$$

$$c(\theta) = -\log(1-p)$$

Binomial

$$t \sim \text{Bin}(n, p) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$\exp \left\{ \log\left(\binom{n}{k}\right) + \log(p^k) + \log((1-p)^{n-k}) \right\}$$

$$\exp\left\{\log\binom{n}{k} + k\log(p) + (n-k)\log(1-p)\right\}$$

$$\exp\left\{\log\binom{n}{k} + k\log\left(\frac{p}{1-p}\right) + n\log(1-p)\right\}$$

$$\exp\left\{n\left[\frac{k}{n}\log\left(\frac{p}{1-p}\right) + \log(1-p)\right] + \log\binom{n}{k}\right\}$$

Result:

$$\theta = \log\left(\frac{p}{1-p}\right)$$

$$q = n$$

$$c(\theta) = \log\binom{n}{k}$$

Poisson

[ADD ME]

2. Identify the canonical link function for each of the models given above.

I JUST LOOKED THIS UP I'M NOT SURE IT'S CORRECT.

Normal

$$\mu$$

Bernoulli

$$\text{logit}(\mu)$$

Binomial

$$\text{logit}(\mu)$$

Poisson

$$\log(\mu)$$

3. Consider now the generalised linear model:

$$t_n|x_n \sim N(\theta(x_n, w), q\gamma_n) \text{ with}$$

$$\theta(x_n, w) = (c')^{-1}(g^{-1}(\phi(x_n)^T w)) =: f(\phi(x_n)^T w)$$

[ADD ME]