Started on	Tuesday, 12 November 2024, 7:02 AM
State	Finished
Completed on	Tuesday, 12 November 2024, 7:04 AM
Time taken	1 min 58 secs
Marks	1.00/1.00
Grade	**4.00** out of 4.00 (**100**%)

Question **1**

Correct

Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that A[j] - A[i] = k, i != j.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as 5 - 1 = 4

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```
#include <stdio.h>

int countPairs(int arr[], int n, int k) {
    int i = 0, j = 1;

    while (j < n) {
        int diff = arr[j] - arr[i];

        if (diff == k && i != j) {
            return 1; // Pair found
        } else if (diff < k) {
            j++; // Increase difference by m
        } else {
            i++; // Decrease difference by m
        }

        // Ensure pointers are not the same
        if (i == j) {
            j++;
        }
    }

    return 0; // No such pair found
}

int main() {
    int n, k;

    scanf("%d", &n);

    int arr[n];

    for (int i = 0; i < n; i++) {
        scanf("%d", &arr[i]);
    }


    scanf("%d", &k);

    int result = countPairs(arr, n, k);
```

```
    printf("%d\n", result);

    return 0;
}

```

	Input	Expected	Got	
✔	3 1 3 5 4	1	1	✔
✔	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✔
✔	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✔
✔	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✔

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

◄ 5-Pair with Difference-O(n^2)Time Complexity,O(1) Space Complexity

Jump to...