DEPENDENCE OF THE SIVERS TMD SHIFT ON QUARK LONGITUDINAL MOMENTUM FRACTION x FROM LATTICE QCD

BY

HARIPRASHAD RAVIKUMAR, B.Sc., M.Sc., M.S.

A dissertation submitted to the Graduate School
in partial fulfillment of the requirements
for the degree
PhD in Physics

Major: Physics

NEW MEXICO STATE UNIVERSITY
LAS CRUCES, NEW MEXICO
(Month 2026)

[If you are filing for copyright, place the notice here.
If not filing for copyright, delete this page.]

(Type in Candidates Name)

Candidate

(Type in Candidates Major)

Major

This Dissertation is approved on behalf of the faculty of New Mexico State University, and it is acceptable in quality and form for publication:

Delete the lines not needed for your committee

Approved by the thesis Committee:

(Type in Chairperson Name)

Chairperson

(Type in Committee member's name)

Committee Member

(Type in Committee member's name)

Committee Member

(Type in Committee member's name)

Committee Member

(Type in Committee member's name)

Committee Member

DEDICATION

I dedicate this .....

ACKNOWLEDGMENTS

I would like to thank my advisor Dr Michael Engelhardt, ...

VITA

2015–2018	B.Sc. in Physics, Dr. N.G.P. Arts and Science College, Coimbatore, Tamil Nadu, India
2018	Summer Research Intern, Indian Institute of Astrophysics (IIA), Bengaluru, Karnataka, India
2019–2021	M.Sc. in Physics, National Institute of Technology Jalandhar (NIT-J), Jalandhar, Punjab, India
2021–2024	M.S. in Physics, New Mexico State University, Las Cruces, New Mexico, USA
2026	Modeling and Simulation Summer Intern, Western Digital, San Jose, California, USA
2021–2026	Graduate Research/Teaching Assistant, New Mexico State University, Las Cruces, New Mexico, USA

PROFESSIONAL AND HONORARY SOCIETIES

Physics Graduate Student Organization, NMSU

American Physical Society

LATTICE-USQCD

FIELD OF STUDY

Major Field: Physics (Lattice Quantum Chromodynamics)

ABSTRACT

THE x-DEPENDENCE OF THE SIVERS TMD FROM LATTICE QCD

BY

HARIPRASHAD RAVIKUMAR, B.Sc., M.Sc., M.S.

Doctor of Philosophy

New Mexico State University

Las Cruces, New Mexico, 2022

Dr. Name I. Surname , Position

In this work we present... (abstract goes here)

Contents

LIST OF TABLES xiv

LIST OF FIGURES xxvi

1 INTRODUCTION AND MOTIVATION 1

1.1 From form factors to partons 1

1.2 Beyond one dimension 2

1.3 Spin and single-spin asymmetries 4

1.4 Experimental and phenomenological status 6

1.5 Lattice QCD and the x-dependence problem 7

1.6 This thesis 8

1.7 Outline 9

2 QCD AND HADRON STRUCTURE 12

2.1 Conventions 12

2.2 Quantum field theory and the path integral 13

2.3 Gauge invariance and QCD 14

2.4 Asymptotic freedom, confinement and chiral symmetry 16

2.5 Deep inelastic scattering and the parton model 17

2.6 Form factors, GPDs and the transverse structure of the nucleon 19

2.7 SIDIS, Drell-Yan and TMD factorization 21

2.8 The TMD correlator and gauge links 22

2.8.1 Process-dependent gauge links 22

2.8.2 Leading-twist TMDs 23

2.8.3 Time reversal and the sign change 24

2.9 Rapidity divergences, the soft factor and evolution 25

2.10 Transverse position space and the Sivers shift 26

2.11 From $b{\cdot}P$ to x . . . 27

2.12 TMDs from lattice QCD . . . 29

3 LATTICE QCD METHODOLOGY . . . 32

3.1 The Euclidean Lattice . . . 32

3.1.1 Euclidean Dirac matrices. . . . 32

3.2 Gauge Fields on the Lattice . . . 33

3.2.1 Continuum limit of the Wilson action. . . . 34

3.3 Fermions on the Lattice . . . 35

3.3.1 Naive discretization and doublers. . . . 35

3.3.2 Wilson fermions. . . . 36

3.3.3 Clover improvement. . . . 37

3.3.4 Other formulations. . . . 37

3.4 Path Integral, Wick Contractions and Quark Propagators . . . 37

3.4.1 Monte Carlo evaluation. . . . 38

3.4.2 Computing propagators. . . . 38

3.4.3 Example: the pion two-point function. . . . 38

3.5 The Nucleon Two-Point Function . . . 39

3.5.1 Interpolating operator and contraction. . . . 39

3.5.2 Spectral decomposition. . . . 39

3.6 Three-Point Functions of Local Operators . . . 40

3.6.1 Definition and spectral decomposition. . . . 40

3.6.2 Quark-level contraction and sequential propagators. . . . 41

3.6.3 Ratio and zero momentum transfer. . . . 41

3.7 Non-Local Staple-Link Operators . . . 42

3.7.1 Lattice operator and three-point function. . . . 42

3.7.2 Contraction. . . . 43

3.7.3 Ratio and matrix element. . . . 44

4 TMD CORRELATOR FOR LATTICE CALCULATIONS 46

4.1 Definition of TMDs 46

4.2 TMDs in Fourier space 48

4.3 Link geometry 48

4.4 Symmetry Constraints 50

4.5 Correlator Parametrization 52

4.6 Generalized Sivers shift from amplitudes 55

5 LATTICE DATA & EXTRACTING THE $x-$DEPENDENCE OF SIVERS SHIFT 59

5.1 The Sivers Shift as a Function of Momentum Fraction x 59

5.2 The Collins-Soper Parameter ($\hat{\zeta}$) 61

5.3 Kinematic Constraints to scan full $b \cdot P$ 61

5.3.1 Explicit Calculation for Hadron Momentum $P_1 = -1$ 61

5.4 Lattice Parameter Space and Off-Axis Links 62

5.4.1 Off-Axis $\boldsymbol{v}$ and Spatial Interpolation 64

5.4.2 Spatial Interpolation of Off-Axis Coordinates 69

5.5 Quark Separation Vectors ($\boldsymbol{b}$) and Amplitude Parity 72

5.5.1 Gauge Link Construction for $\boldsymbol{b}$ 73

5.5.2 Symmetry Improving Operation (Path Averaging) for $\boldsymbol{b}$ 76

5.6 Solving for Real and Imaginary Amplitudes $\widetilde{A}_{iB}^{\mathrm{Re}}$ and $\widetilde{A}_{iB}^{\mathrm{Im}}$ from correlator $\widetilde{\Phi}^{[\gamma^\mu]}_{\mathrm{unsubtr.}}$ 77

5.7 Constructing the unpolarized $\widetilde{A}_{2B}$ and Sivers $\widetilde{A}_{12B}$ Amplitudes 83

5.8 Transformation to the Lorentz-Invariant Variable Space 85

5.9 Numerical Results and Parity Constraints of the Extracted Amplitudes 86

5.9.1 Amplitude Profiles at $\hat{\zeta} = 0.090772$ ($P_L = -1$) 86

5.9.2 Amplitude Profiles at $\hat{\zeta} = 0.294367$ ($P_L = -2$) 89

5.9.3 Amplitude Profiles at $\hat{\zeta} = 0.539684$ ($P_L = -3$) 91

5.9.4 Amplitude Profiles at $\hat{\zeta} = 0.785756$ ($P_L = -4$) . . . 93

6 FITTING METHODOLOGIES AND STABILITY ANALYSIS . . . 97

6.1 Discretization Effects and Short-Distance Data Exclusion . . . 98

6.2 Failure of Simple Gaussian Model . . . 99

6.3 Physics-Informed Symbolic Regression via PySR . . . 100

6.3.1 Expression Search and Configuration . . . 101

6.3.2 The Physics-Informed Custom Loss Function . . . 101

6.3.3 Pareto Front of Discovered Models . . . 103

6.4 Systematic Exploration of the Power-Law Model . . . 107

6.5 Candidate Selection . . . 112

6.6 Simultaneous Fitting Strategy . . . 114

6.6.1 Simultaneous Fit with Full Covariance Matrix . . . 115

6.6.2 Simultaneous Fit Results for $\hat{\zeta} = 0.090772$ ($\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-1, 0, 0\}$) . . . 116

6.6.3 Simultaneous Fit Results for $\hat{\zeta} = 0.294367$ ($\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$) . . . 121

6.6.4 Simultaneous Fit Results for $\hat{\zeta} = 0.539684$ ($\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$) . . . 126

6.6.5 Simultaneous Fit Results for $\hat{\zeta} = 0.785756$ ($\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$) . . . 131

6.7 Test for Factorization in $(P \cdot b)$ and b^2 . . . 136

6.7.1 Ratio Test on the Lattice Data . . . 136

6.7.2 Rejection of Separable Longitudinal and Transverse Factorization . . . 139

7 EXTRAPOLATIONS TO THE PHYSICAL LIMIT . . . 144

7.1 Motivation for the $\hat{\zeta} \to \infty$ Extrapolation . . . 144

7.2 Extrapolation at fit-parameters level . . . 146

7.3 Testing Factorization in x and $|b_T|$ at the Physical Limit . . . 151

7.3.1 Quantifying the Distance from a Factorized Form . . . 151

7.3.2 Factorization Test for the Sivers Shift . . . 152

7.3.3 Factorization Test at the Amplitude Level . . . 154

8 CONCLUSION AND OUTLOOK 159

A QLUA: Gamma $\Gamma(n)$ 160

REFERENCES 161

List of Tables

1 Leading-twist quark TMDs, arranged by the polarization of the nucleon (rows) and of the quark (columns): unpolarized (U), longitudinally polarized (L) and transversely polarized (T). The Sivers function $f_{1T}^{\perp}$ and the Boer-Mulders function $h_1^{\perp}$ are naively T-odd. 24

2 Resulting Collins-Soper parameters ($\hat{\zeta}$) and the necessary geometric constraint on the staple vector components to ensure scan full $b \cdot P$. 62

3 Configuration parameters for the Lattice QCD non-local matrix element extraction. Our lattice size is of $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is of $a = 0.11403$ fm 63

4 Hyperparameter and operator configuration for the Physics-Informed `PySR` Symbolic Regression search. 101

5 Pareto front of analytical candidate models discovered by `PySR` for the real amplitude $\widetilde{A}_{2B}^{\mathrm{Re}}(\eta, b_L, b_T)$, ranked by mathematical complexity and loss. The optimal physics-informed models ($C = 14$ and $C = 16$) are highlighted. 103

6 Sequential Lorentzian fits for $\widetilde{A}_{12B}^{\mathrm{Re}}$ at individual η slices showing consistent χ^2/dof and stable parameters. 111

7 Sequential Lorentzian fits for $\widetilde{A}_{2B}^{\mathrm{Im}}$ at individual η slices showing consistent χ^2/dof. 112

8 Simultaneous Fit with common f: for $P_L = -1$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global χ^2/DoF $= 1.85(10)$ 116

9 Numerical Integration of Fourier Transforms: $P_L = -1, b_{min} \geq 3$ 119

10 Simultaneous Fit with common f: for $P_L = -2$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global χ^2/DoF $= 1.197(85)$ 121

11 Numerical Integration of Fourier Transforms: $P_L = -2, b_{min} \geq 3$ 124

12 Simultaneous Fit with common f: for $P_L = -3$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global χ^2/DoF $= 0.712(66)$ 126

13 Numerical Integration of Fourier Transforms: $P_L = -3, b_{min} \geq 3$ 129

14 Simultaneous Fit with common f: for $P_L = -4$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global χ^2/DoF $= 0.434(51)$ 131

15 Numerical Integration of Fourier Transforms: $P_L = -4, b_{min} \geq 3$ 134

16 All 34 $(P\cdot b, b^2)$ pairs available at $\eta|v|/a = 8$ and $P_L = -1$, where $P\cdot b = -P_L\, b_L$ and $b^2 = b_L^2 + b_T^2$. Grey entries fail the cut $|\boldsymbol{b}| \geq 3a$. Bold entries are the data used in the factorization test of Figure 75. 137

17 Origin of each grid value used in Figure 75. "Quad. (b_1^2, b_2^2, b_3^2)" means the parabola in b^2 through the data at those three points on the same $P \cdot b$ row; "lin. $[b_a^2, b_b^2]$" means linear interpolation in b^2 between the data at b_a^2 and b_b^2. Where a cell has two entries, the upper one gives the linear points (circles) and the lower one the quadratic points (triangles) of Figure 75. The pattern is the same for all three amplitudes. 137

18 Comparison of the four factorized longitudinal-transverse models across nucleon momenta P_L, with extracted parameters shown at $P_L = -1$. While models with power-law transverse envelopes describe $\widetilde{A}_{2B}^{\mathrm{Re}}$ passably, all four factorized ansatze fail on $\widetilde{A}_{2B}^{\mathrm{Im}}$ and $\widetilde{A}_{12B}^{\mathrm{Re}}$. 141

19 Numerical integration of the Fourier-transform integrals at $|b_T| = 3a \approx 0.34$ fm, evaluated using the extrapolated ($\hat{\zeta} \to \infty$) amplitudes of Fig. 81, over the full range and three restricted x-ranges isolating the quark, antiquark, and combined contributions. 147

20 Relative distance of the extrapolated Sivers shift $\langle k_y \rangle_{TU}(x, |b_T|)$ from its optimal factorized approximation $F(x)G(|b_T|)$, Eq. (7.9), over $|b_T| \in [0.3421, 0.9403]$ fm for four choices of x-window. 153

21 Relative distance of each extrapolated amplitude (and their combination $\widetilde{A}_{2B}^{[x]\mathrm{Re}} - i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$) from its optimal factorized approximation, over the same $|b_T| \in [0.3421, 0.9403]$ fm grid and x-windows as Table 20. 155

List of Figures

1 Relations between the distributions that describe the quark structure of the nucleon. Integrating the generalized TMDs (or their Fourier transforms, the Wigner distributions) over transverse position gives TMDs, and integrating over transverse momentum gives GPDs. The collinear PDFs are obtained from TMDs by integrating over $\boldsymbol{k}_{\mathrm{T}}$ and from GPDs in the forward limit. Moments of GPDs in x give the elastic form factors. 3

2 Schematic picture of the Sivers effect. For a nucleon moving into the page ($\boldsymbol{P}$ along $\hat{\boldsymbol{z}}$) with transverse spin $\boldsymbol{S}_{\mathrm{T}}$ along $\hat{\boldsymbol{y}}$, the second term of Eq. (1.1) becomes $-k_x f_{1T}^{\perp}/m_N$. A nonzero Sivers function therefore shifts the density of unpolarized quarks along $\hat{\boldsymbol{x}}$, perpendicular to both $\boldsymbol{P}$ and $\boldsymbol{S}_{\mathrm{T}}$ (solid circle), relative to the symmetric unpolarized distribution (dashed). The shift shown, toward $+\hat{\boldsymbol{x}}$, corresponds to $f_{1T}^{\perp} < 0$. 5

3 Handbag diagram for the leading contribution to the DIS hadronic tensor. The virtual photon scatters off a quark of momentum k with $k^+ = xP^+$, and the lower blob is the quark-quark correlator whose γ^+ projection gives the PDF $f_1(x)$. The dashed line indicates the sum over final states. 18

4 Staple-shaped gauge links connecting the quark fields at 0 and b in the TMD correlator, Eq. (2.37). (a) In SIDIS the link runs to $+\infty$ along the light-cone direction n of the outgoing quark, Eq. (2.38). (b) In Drell-Yan it runs to $-\infty$, Eq. (2.39). The dashed segment is the transverse link at infinity. In the lattice calculation the legs have finite length $\eta|v|$ along a space-like direction v, and the two cases correspond to $\eta v{\cdot}P \to +\infty$ and $\eta v{\cdot}P \to -\infty$ [1]. 23

5 Quark fields $\psi(x)$ and $\psi(x + a\hat{\mu})$ on the sites of a cubic spacetime lattice, the link variable $U_\mu(x)$ between them, and the plaquette $P_{\mu\nu}(y)$ of Eq. (3.6). The product $U_\mu(x)\psi(x + a\hat{\mu})$ transforms in the same way as $\psi(x)$. Reproduced from Ref. [2], Fig. 1.1. 34

6 Staple-shaped gauge link $\mathcal{U}[0, \eta v, \eta v + b, b]$. The staple points along the direction v, in general $v = (0, v_1, v_2, v_3)$, and the limit $\eta \to \infty$ extends it. 49

7 Directions of the spatial staple vectors $\boldsymbol{v}$ mapped in the $v_1 - v_3$ plane. Each vector corresponds to the requisite geometry to satisfy the $b \cdot P \neq 0$ constraint for the four sequential Collins-Soper parameters $\hat{\zeta}$. 63

8 Ideal continuous trajectory of the off-axis staple vector (solid blue) for $P_1 = -1$. Circular markers indicate target continuous coordinates for sequential values of η. Dashed lines illustrate the discrete, step-wise permutations of gauge links used to reach the surrounding bounding integer lattice sites. 65

9 Discrete gauge link configurations defining the interpolation bounds for $P_1 = -1$. Empty brackets $\{\}$ denote a zero-length origin link, while numerical arrays (e.g., $\{1, 3, 1\}$) give the sequence of directional steps. 65

10 Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -2$. 66

11 Discrete gauge link configurations defining the interpolation bounds for $P_1 = -2$. 66

12 Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -3$. 67

13 Discrete gauge link configurations defining the interpolation bounds for $P_1 = -3$. 67

14 Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -4$. 68

15 Discrete gauge link configurations defining the interpolation bounds for $P_1 = -4$. 68

16 Illustration of path degeneracy. Two distinct step-wise gauge link permutations (e.g., solid and dashed lines) terminate at the identical discrete lattice coordinate. We average the correlators from these equivalent paths to suppress local gauge noise before interpolation. 69

17 Quadratic interpolation of the real and imaginary parts of the correlation function for $\boldsymbol{b} = \{1, 2, 0\}$, $\eta = 7$, and $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = (-1, 0, 0)$. Red markers indicate the path-averaged lattice data points as a function of their deviation from the ideal constraint trajectory, $\Delta v_1 = (v_1^{\text{constraint}} - v_1^{\text{lattice}})/a$. The gray band indicates the quadratic fit ('InterpolationOrder $\to$ 2'), and the central blue marker represents the final extracted value at $\Delta v_1 = 0$, lying exactly on the required off-axis spatial trajectory. 71

18 The sampled operator separation vectors mapped in the spatial $b_1 - b_2$ plane (where $b_1 \equiv b_L$ and $b_2 \equiv b_T$). 72

19 Mapping of the spatial lattice vectors into the physical invariant Variable Space $(b\cdot P/(2\pi), |b|/a)$. The trajectories diverge based on the selected target $\hat{\zeta}$. 72

20 Coverage of the quark separation vectors across the simulated lattice volume and their corresponding physical Variable Space representation. Our lattice size is of $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is of $a = 0.11403$ fm 72

21 Discrete gauge link configurations for purely transverse quark separations ($\boldsymbol{b} = \{0, b_2, 0\}$). The numerical arrays show the sequential lattice steps (e.g., '2' represents a shift in the x_2 direction). Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm. 73

22 Gauge link sequences for $+b_1$ (positive longitudinal separation). Symmetric sets of step-wise gauge link paths required to access finite $\pm b \cdot P$ and isolate the real and imaginary components of the invariant amplitudes. Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm. 74

23 Mirrored gauge link sequences for $-b_1$ (negative longitudinal separation).Symmetric sets of step-wise gauge link paths required to access finite $\pm b \cdot P$ and isolate the real and imaginary components of the invariant amplitudes. Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm. 75

24 Illustration of path degeneracy for the quark separation vector $\boldsymbol{b}$. Distinct step-wise permutations (solid and dashed lines) terminating at the same Cartesian coordinate are averaged to suppress local gauge noise. 76

25 The full network of path-averaged $\boldsymbol{b}$ vectors in the $b_1/a - b_2/a$ plane. Solid red lines represent the ideal vectors, while dashed lines illustrate the underlying degenerate staircase paths calculated and averaged on the discrete lattice. 76

26 Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 87

27 Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 8

28 $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$, $\widetilde{A}^{\mathrm{Re}}_{12B}$ & $\widetilde{A}^{\mathrm{Im}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1, 0, 0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space. Notice the lack of discernible signal in $\widetilde{A}^{\mathrm{Im}}_{12B}$. 88

29 Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 89

30 Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 8

31 $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$, $\widetilde{A}^{\mathrm{Re}}_{12B}$ & $\widetilde{A}^{\mathrm{Im}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space. 90

32 Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 91

33 Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 9

34 $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, $\widetilde{A}_{12B}^{\mathrm{Re}}$ & $\widetilde{A}_{12B}^{\mathrm{Im}}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-3,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a=8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space. 92

35 Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-4,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 93

36 Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-4,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space. 9

37 $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, $\widetilde{A}_{12B}^{\mathrm{Re}}$ & $\widetilde{A}_{12B}^{\mathrm{Im}}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-4,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a=8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space. 94

38 The $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude at fixed momentum ($P=-1$) and fixed η (here $\eta=8$). The short-distance data ($|\boldsymbol{b}|<3a$) curve in the opposite direction to the long-distance data ($|\boldsymbol{b}|\geq 3a$). 98

39 The $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitude at fixed momentum ($P=-1$) and fixed η (here $\eta=8$). The short-distance data ($|\boldsymbol{b}|<3a$) curve in the opposite direction to the long-distance data ($|\boldsymbol{b}|\geq 3a$). 99

40 Failure of Gaussian model fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ (with $\alpha-\beta=-0.0427(27)$ and $\chi^2/\mathrm{dof}\approx 235$) and $\widetilde{A}_{12B}^{\mathrm{Re}}$ (with $\alpha-\beta=-0.022(25)$ and $\chi^2/\mathrm{dof}\approx 1.4$) amplitudes. 100

41 The top `PySR` analytical models ($C=11,14,16,17$) compared with the lattice data for $\widetilde{A}_{2B}^{\mathrm{Re}}$ at fixed staple extents $\eta=6,7,8,10$ (with $P_1=-1$ and $|\boldsymbol{b}|\geq 3a$). Each panel shows the amplitude as a function of the longitudinal separation b_L/a at fixed transverse separations b_T, with the four Pareto-front models overlaid. The two exponential-η models ($C=14$, solid; $C=16$, dashed) give the most balanced description of the data across all η values, while the pure power-law-η models ($C=11$, dotted; $C=17$, dot-dashed) deviate slightly and systematically at the ends of the η range. 105

42 Diagnostic straight-line test for the Gaussian hypothesis. If the amplitude followed a pure Gaussian profile $\widetilde{A}_{2B}^{\mathrm{Re}} \propto e^{-c\,b^2}$, the data would appear as a straight line on a logarithmic ordinate versus b^2 (red dashed line, best Gaussian fit with $c = 0.058$). The lattice data (black markers) curve upward away from the Gaussian line at large b^2, indicating a heavier-than-Gaussian tail, which the `PySR` power-law form $(1 + b^2)^{-1.39}$ (blue solid line) accurately captures. The curvature is direct, model-independent evidence that the spatial decay of the non-local quark bilinear follows an algebraic power law rather than a Gaussian exponential. 106

43 3D jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of 16 Gaussian ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta}\, \exp\big[-c\,(b_L P_L)^2 - d\,(b_L^2 + b_T^2)\big]$, in which the coefficients c and d are each independently allowed to be either constant or one of three η-dependent forms, $c \to c(\eta)$ and/or $d \to d(\eta)$. 107

44 3D Jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of power-law ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta}\, \big(1 + (c+d)b_L^2 + d b_T^2\big)^{-j}$, in which the coefficient c is either constant or one of several η-dependent forms, $c \to c(\eta)$. 108

45 3D jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of power-law ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta}\, \big(1 + (c+d)b_L^2 + d b_T^2\big)^{-j}$, in which the coefficient c takes further η-dependent forms, $c \to c(\eta)$, that yield wider Fourier transforms in x space. 109

46 Sequential Lorentzian fits for $\widetilde{A}_{2B}^{\mathrm{Re}}$ at individual η slices, holding the global power $j = 2$ and amplitude $a = 4$ constant. 110

47 The extracted parameter $c(\eta)$ overlaid with the candidate asymptotic models $c_1(\eta), \ldots, c_{10}(\eta)$. The extracted values approach an asymptote at large η. 110

48 The extracted parameter $d(\eta)$, which shows no visible dependence on the staple extent η. . . . 111

49 Lattice extracted parameters c (left) and d (right) for $\widetilde{A}_{12B}^{\mathrm{Re}}$ as a function of η. Both parameters show no visible dependence on η. 111

50 Lattice extracted parameters c (left) and d (right) for $\widetilde{A}_{2B}^{\mathrm{Im}}$ as a function of η. The data are noisier than for the real amplitudes, but consistent with the constant-parameter ansatz. 112

51 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 116

52 Fit for $\widetilde{A}_{2B}^{\mathrm{Im}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 117

53 Fit for $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 117

54 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$ & $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit. 118

55 x-dependence at $b^2 = 9a^2$ for $P_L = -1$. Top row: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, and $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$. Bottom row: $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, $\langle k_y\rangle_{TU}$, and $x\langle k_y\rangle_{TU}$. Bands show the Jackknife uncertainty. 120

56 The quantities of Figure 55 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -1$. 120

57 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 121

58 Fit for $\widetilde{A}_{2B}^{\mathrm{Im}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 122

59 Fit for $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 122

60 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$ & $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-2,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a=8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit. 123

61 x-dependence at $b^2=9a^2$ for $P_L=-2$. Top row: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, and $\tilde{f}_1^{(0)}=2\widetilde{A}_{2B}^{[x]}$. Bottom row: $\tilde{f}_{1T}^{\perp(1)}=-2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, $\langle k_y\rangle_{TU}$, and $x\langle k_y\rangle_{TU}$. Bands show the Jackknife uncertainty. 125

62 The quantities of Figure 61 as functions of x and $b^2/a^2\in[9,68]$ for $P_L=-2$. 125

63 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-3,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 126

64 Fit for $\widetilde{A}_{2B}^{\mathrm{Im}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-3,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 127

65 Fit for $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-3,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 127

66 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$ & $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-3,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a=8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit. 128

67 x-dependence at $b^2=9a^2$ for $P_L=-2$. Top row: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, and $\tilde{f}_1^{(0)}=2\widetilde{A}_{2B}^{[x]}$. Bottom row: $\tilde{f}_{1T}^{\perp(1)}=-2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, $\langle k_y\rangle_{TU}$, and $x\langle k_y\rangle_{TU}$. Bands show the Jackknife uncertainty. 130

68 The quantities of Figure 67 as functions of x and $b^2/a^2\in[9,68]$ for $P_L=-2$. 130

69 Fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi)=\{-4,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 131

70 Fit for $\widetilde{A}^{\mathrm{Im}}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 132

71 Fit for $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid. 132

72 Fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ & $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit. 133

73 x-dependence at $b^2 = 9a^2$ for $P_L = -2$. Top row: $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$, $i\widetilde{A}^{[x]\mathrm{Im}}_{2B}$, and $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$. Bottom row: $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$, $\langle k_y\rangle_{TU}$, and $x\langle k_y\rangle_{TU}$. Bands show the Jackknife uncertainty. 135

74 The quantities of Figure 73 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -2$. 135

75 Factorization ratio test at $\eta|v|/a = 8$, $P_L = -1$ ($\hat{\zeta} = 0.090772$) for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ and $\widetilde{A}^{\mathrm{Re}}_{12B}$. Top row: $T(P\cdot b, b^2)$ of Eq. (6.28), the ratio to the $P\cdot b = 4$ row, plotted against b^2. Bottom row: $B(P\cdot b, b^2)$ of Eq. (6.29), the ratio to $b^2 = 18$, plotted against $P\cdot b$. Under factorization, every set of points is flat. Filled circles use linear interpolation in b^2 on all three rows. Open triangles use quadratic interpolation on the $P\cdot b = 0$ and 4 rows and linear interpolation on the $P\cdot b = 3$ row (Table 17). A triangle is omitted where both schemes give the same value, which happens for every point that involves only the $P\cdot b = 3$ row and data points. Error bars are Jackknife errors and do not include any uncertainty from the interpolation. 138

76 Fits testing the separability hypothesis. Pairing an exponential decay in $(P\cdot b)$ with a Gaussian transverse profile gives a poor fit ($\chi^2/\mathrm{dof} \approx 4.98$). In models with a lower χ^2/dof, the coefficient c consistently fits to zero within uncertainties and is negative in each Jackknife set, so the data reject the separability hypothesis. 139

77 Fits testing the separability hypothesis with η-dependent c, written $c(\eta)$. In models with a lower χ^2/dof, the coefficient c is now positive. 140

78 Discrete $\hat{\zeta}$-slices of the reconstructed amplitudes $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$ and $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$ and the unpolarized and Sivers TMD combinations $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$ and $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, plotted against both x and $1/\hat{\zeta}$ at fixed transverse separation $|b_T| = 3a \approx 0.34$ fm. Each ribbon corresponds to one of the four simulated momenta $P_L = -1, -2, -3, -4$ (equivalently $1/\hat{\zeta} = 11.02,\ 3.40,\ 1.85,\ 1.27$). The systematic drift of every panel with $1/\hat{\zeta}$ motivates the extrapolation to the physical light-cone limit $\hat{\zeta} \to \infty$ performed below. 145

79 Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$ for the same four momenta and the same $|b_T|$ as Fig. 78, restricted to the quark region $x \geq 0$. 145

80 Extrapolation of the twelve simultaneous-fit parameters to the physical limit $\hat{\zeta} \to \infty$, using only the three higher momenta $P_L = -2, -3, -4$ (black points, plotted against $1/\hat{\zeta}$). The dashed red curve and shaded band show the fitted ansatz $c + d/\hat{\zeta}^2$ of Eq. 7.1, with each panel's Jackknife χ^2/dof noted in its title; the blue square at $1/\hat{\zeta} = 0$ marks the extrapolated physical-limit value. The parameters c_{re}, c_{im}, and c_{reA12B} are shown rescaled by $1/P_L^2$, and a_{im} by $1/P_L$, as described in the text. 147

81 Unpolarized and Sivers TMD amplitudes reconstructed from the extrapolated ($\hat{\zeta} \to \infty$) fit parameters of Fig. 80, evaluated as continuous functions of x at fixed transverse separation $|b_T| = 3a \approx 0.34$ fm: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, the full unpolarized combination $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$, and the Sivers combination $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$. Shaded bands are the Jackknife uncertainty propagated from the twelve extrapolated parameters. 148

82 Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$ built from the extrapolated amplitudes of Fig. 81, restricted to the quark region $x \geq 0$. Shaded bands are the Jackknife uncertainty. 148

83 The four extrapolated ($\hat{\zeta} \to \infty$) amplitude quantities of Fig. 81, shown as surfaces in x and $|b_T|$ (from $3a \approx 0.34$ fm to $\sqrt{68}\,a \approx 0.94$ fm). The joint x-$|b_T|$ dependence shows how the reconstructed amplitudes change with transverse quark separation at the physical limit. 149

84 The extrapolated Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$ of Fig. 82, shown as surfaces in $x \geq 0$ and $|b_T|$ over the same range. 149

85 Comparison, at fixed $|b_T| = 3a \approx 0.34$ fm, between the four amplitude quantities evaluated at each of the four simulated momenta ($P_L = -1, -2, -3, -4$, colored curves, each using its own unextrapolated fit parameters) and the extrapolated $\hat{\zeta} \to \infty$ result (black dashed curve), which uses only $P_L = -2, -3, -4$. The extrapolated curve differs from all three finite-momentum curves used in the extrapolation, including the one at the largest simulated momentum, so the fit-parameter extrapolation of Fig. 80 captures real momentum dependence and does not just reproduce the $P_L = -4$ result. 150

86 The same comparison as Fig. 85 for the Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$ ($x \geq 0$). 150

87 Left and center: the optimal separable factors $g(x)$ and $h(|b_T|)$ (Eqs. (7.7)-(7.8), normalized so mean$[g^2] = 1$) for the extrapolated Sivers shift, shown for all four x-windows of Table 20. Right: the reconstructed $g(x)h(|b_T|)$ (dashed) compared to the actual extrapolated Sivers shift (solid) at the smallest and largest fitted $|b_T|$, using the $x \in [0, 1]$ fit; the visible gap between the solid and dashed curves is the residual quantified in Table 20. 154

88 $R_b(x) = \langle k_y \rangle_{TU}(x, |b_T|)/\langle k_y \rangle_{TU}(x, 0.34 \text{ fm})$ for four values of $|b_T|$, as a function of x. Under exact factorization each curve would be flat. Instead, every curve rises from its $x \to 0$ value to a peak near $x \approx 0.1$ before decaying, so the $|b_T|$-dependence of the Sivers shift itself depends on x. 154

89 $R_x(|b_T|) = \langle k_y \rangle_{TU}(x, |b_T|)/\langle k_y \rangle_{TU}(0.3, |b_T|)$ for four values of x, as a function of $|b_T|$. The $x = 0.2$ curve changes by more than a factor of two across the fitted $|b_T|$ range instead of staying flat, so the x-dependence of the Sivers shift itself depends on $|b_T|$, consistent with Fig. 88. 155

90 The same ratio test as Fig. 88, $R_b(x)$, repeated at each finite momentum individually using its own raw (unextrapolated) fit parameters. None of the four panels shows a flat curve, so the non-factorization is a feature of the fitted amplitudes and not an artifact of the $\hat{\zeta} \to \infty$ extrapolation. 156

91 The same ratio test as Fig. 89, $R_x(|b_T|)$, repeated at each finite momentum individually. The departure from flatness grows with $|P_L|$, most visibly in the $x = 0.2$ curve, which is consistent with the parameter mismatch identified in the extrapolated j_X and d_X values above. 157

92 Optimal separable factor $g(x)$ for each of the four amplitude combinations of Table 21, for all four x-windows (normalized so mean$[g^2] = 1$). Each amplitude's $g(x)$ is dominated by a sharp rise as $x \to 0$, which steepens as the window narrows and reflects the Bessel-function behavior at small x. 158

93 Optimal separable factor $h(|b_T|)$ for each of the four amplitude combinations of Table 21, for all four x-windows. For all four amplitudes, $h(|b_T|)$ falls smoothly with increasing $|b_T|$, but its magnitude and slope depend on the x-window used to define the fit, which itself signals imperfect factorization. 159

94 Reconstructed $g(x)h(|b_T|)$ (dashed) against the actual extrapolated amplitude data (solid) at the smallest and largest fitted $|b_T|$, using the $x \in [0, 1]$ fit, for each of the four amplitude combinations of Table 21. The largest reconstruction gap appears near $x \to 0$ for every amplitude, consistent with the $x \in [0, 1]$ column of Table 21 having the largest relative distance in every row. 160

1 INTRODUCTION AND MOTIVATION

Protons and neutrons make up almost all of the visible mass in the universe, yet their internal structure is still only partly understood. We know the theory that governs them. Quantum Chromodynamics (QCD) describes quarks and gluons interacting through the exchange of color charge, and its Lagrangian fits on a single line. What we cannot do in general is solve it. At the distance scale of a nucleon, about one femtometer, the QCD coupling is large, perturbation theory does not converge, and the quarks and gluons are permanently confined inside hadrons. The properties of the nucleon therefore have to be extracted from experiment through well-defined observables or computed numerically from first principles.

This thesis is about one such observable, the Sivers function $f_{1T}^{\perp}$. It describes how the transverse momentum of an unpolarized quark is correlated with the transverse spin of the nucleon that contains it. We compute the Sivers function with lattice QCD and, unlike earlier lattice calculations, we resolve its dependence on the longitudinal momentum fraction x carried by the quark. This chapter explains where the Sivers function comes from, why it matters, what experiments have measured, and what a lattice calculation can add. It closes with a summary of the contributions of this work and an outline of the remaining chapters.

1.1 From form factors to partons

The first direct evidence that the proton has a finite size came from elastic electron scattering. In the 1950s Hofstadter and collaborators measured the angular distribution of electrons scattered off hydrogen and found that it fell below the prediction for a point charge. The deviation is described by form factors, which are Fourier transforms of the charge and magnetization distributions, and it corresponds to a proton radius of roughly 0.8 fm [3].

A decade later, the SLAC-MIT experiments pushed the electron energy high enough to break the proton apart. In this deep inelastic scattering (DIS) regime the cross section showed a surprising feature: at large momentum transfer Q^2 it depended, to a good approximation, only on the dimensionless ratio $x_B = Q^2/(2P\cdot q)$ and not on Q^2 separately [4, 5]. Bjorken had predicted this scaling behavior from current algebra [6]. Feynman gave it a simple physical meaning [7]: in a frame where the proton moves very fast, the virtual photon scatters

incoherently off pointlike constituents, the partons, and x_B is the fraction of the proton momentum carried by the struck parton. The observed relation between the two DIS structure functions, $F_2 = 2xF_1$, showed that the charged partons have spin 1/2 [8]. They were identified with the quarks that Gell-Mann and Zweig had introduced to organize the hadron spectrum [9, 10].

The parton picture raised a puzzle. If quarks are bound so tightly that no free quark has ever been observed, why do they behave as nearly free particles when struck hard? The answer came with non-Abelian gauge theory. Quarks carry a threefold color charge, and the color interaction is mediated by eight gluons that are themselves charged [11, 12]. Gross and Wilczek [13] and Politzer [14] showed that in such a theory the effective coupling decreases logarithmically at short distances. This property, asymptotic freedom, explains Bjorken scaling and its slow logarithmic violation. The same running makes the coupling grow at long distances, which is consistent with confinement. QCD therefore describes both regimes, and the strong coupling at long distances is the reason nucleon structure has to be studied with non-perturbative tools.

1.2 Beyond one dimension

The central objects of the parton model are the parton distribution functions (PDFs). The unpolarized quark PDF $f_1^q(x)$ gives the number density of quarks of flavor q carrying momentum fraction x in a fast-moving nucleon. PDFs are universal: the same functions describe DIS, the Drell-Yan production of lepton pairs in hadron collisions [15], jet production at the LHC, and many other processes. This universality rests on collinear factorization theorems, which separate a hard partonic cross section, calculable in perturbation theory, from the non-perturbative distributions. PDFs have been determined from global fits to experimental data with high precision over a wide range of x.

A PDF, however, is a one-dimensional picture. It integrates over all the transverse degrees of freedom of the parton, both its position and its momentum in the plane perpendicular to the nucleon's direction of motion. Two extensions restore this information. Generalized parton distributions (GPDs), which appear in exclusive processes such as deeply virtual Compton scattering, encode the correlation between x and the transverse position $\boldsymbol{b}_\mathrm{T}$ of the parton [16, 17, 18, 19]. At zero skewness, their Fourier transform in the momentum transfer gives impact-parameter distributions, which have a density interpretation [20]. Transverse momentum

dependent parton distributions (TMDs) instead describe the joint distribution in x and the intrinsic transverse momentum $\boldsymbol{k}_\mathrm{T}$ [21, 22, 23]. They appear in processes where a transverse momentum is measured that is small compared to the hard scale, such as semi-inclusive DIS (SIDIS) and Drell-Yan production at low transverse momentum of the lepton pair.

The idea that partons move transversely inside the nucleon is almost as old as the parton model. Cahn showed that intrinsic transverse momentum produces a $\cos\phi_h$ modulation of the SIDIS cross section [24]. Collins and Soper formulated the factorization of processes sensitive to transverse momentum [25, 21], and Collins, Soper and Sterman used it to resum large logarithms in the transverse momentum distribution of Drell-Yan pairs [26]. The complete set of TMDs at leading power was classified for Drell-Yan by Ralston and Soper [27] and for SIDIS by Kotzinian [28] and by Mulders and Tangerman [29, 22]. Their operator definitions, including the treatment of gauge links and rapidity divergences, reached their current form much later [30, 23, 31].

Both GPDs and TMDs can be obtained from a common parent, the generalized TMDs (GTMDs), whose Fourier transforms are quantum phase-space (Wigner) distributions in x, $\boldsymbol{b}_\mathrm{T}$ and $\boldsymbol{k}_\mathrm{T}$ [32, 33, 34]. Figure 1 sketches these relations. Together, GPDs and TMDs make up the program of nucleon "tomography", a central goal of the Electron-Ion Collider (EIC) now under construction [35, 36, 37].



Figure 1: Relations between the distributions that describe the quark structure of the nucleon. Integrating the generalized TMDs (or their Fourier transforms, the Wigner distributions) over transverse position gives TMDs, and integrating over transverse momentum gives GPDs. The collinear PDFs are obtained from TMDs by integrating over $\boldsymbol{k}_\mathrm{T}$ and from GPDs in the forward limit. Moments of GPDs in x give the elastic form factors.

1.3 Spin and single-spin asymmetries

Spin measurements have repeatedly shown that this picture is incomplete. In 1988 the European Muon Collaboration measured the polarized structure function g_1 of the proton and concluded that the quark spins account for only a small fraction of the proton spin [38]. The result, the "proton spin crisis", started a long program to understand how the spin of the nucleon is shared among quark spin, gluon spin, and orbital angular momentum [39, 18, 40]. Orbital motion requires transverse momentum, so TMDs are one natural place to look for it.

A second puzzle came from hadron collisions with a transversely polarized beam or target. Large transverse polarizations of Λ hyperons produced in unpolarized proton collisions were observed in the 1970s [41], and large left-right asymmetries in the production of pions off transversely polarized protons were measured by the E704 experiment at Fermilab [42]. Such transverse single-spin asymmetries (SSAs) were not expected. In collinear perturbative QCD, Kane, Pumplin and Repko showed that they are suppressed by the quark mass over the hard scale and by a power of α_s [43], which makes them far too small to explain the data.

In 1990 Sivers proposed a mechanism beyond the collinear framework [44]. If the transverse momentum distribution of quarks in a transversely polarized proton is not symmetric, but shifted in a direction perpendicular to the spin, then the produced hadrons inherit a preferred side, and a left-right asymmetry follows. The size of this effect is set by a new distribution function, today called the Sivers function $f_{1T}^{\perp}(x, \boldsymbol{k}_{\mathrm{T}}^2)$. For a nucleon moving along the z direction with transverse spin $\boldsymbol{S}_{\mathrm{T}}$, the number density of unpolarized quarks reads

$$f_{q/N^{\uparrow}}(x, \boldsymbol{k}_{\mathrm{T}}) = f_1^q(x, \boldsymbol{k}_{\mathrm{T}}^2) - \frac{\epsilon_{ij}\, k_{T,i}\, S_{T,j}}{m_N}\, f_{1T}^{\perp q}(x, \boldsymbol{k}_{\mathrm{T}}^2)\ , \tag{1.1}$$

with $\epsilon_{12} = +1$, so that $\epsilon_{ij} k_{T,i} S_{T,j} = (\hat{\boldsymbol{z}} \times \boldsymbol{k}_{\mathrm{T}}) \cdot \boldsymbol{S}_{\mathrm{T}}$. This is the "Trento" convention [45], and it is the form used throughout this thesis (compare Eq. (4.7)). Figure 2 illustrates the resulting distortion of the quark distribution in transverse momentum space.

The Sivers function soon ran into a theoretical objection. Collins argued in 1993 that time-reversal invariance of QCD forces it to vanish [46]. Under naive time reversal (time reversal applied to the operator definition without regard to the gauge link) the term $\epsilon_{ij} k_{T,i} S_{T,j}$ changes sign while the rest of the correlator does not, so the Sivers term had to be zero. Functions with this property are still called "naively T-odd". For almost ten



Figure 2: Schematic picture of the Sivers effect. For a nucleon moving into the page ($\boldsymbol{P}$ along $\hat{\boldsymbol{z}}$) with transverse spin $\boldsymbol{S}_{\rm T}$ along $\hat{\boldsymbol{y}}$, the second term of Eq. (1.1) becomes $-\,k_x f_{1T}^{\perp}/m_N$. A nonzero Sivers function therefore shifts the density of unpolarized quarks along $\hat{\boldsymbol{x}}$, perpendicular to both $\boldsymbol{P}$ and $\boldsymbol{S}_{\rm T}$ (solid circle), relative to the symmetric unpolarized distribution (dashed). The shift shown, toward $+\hat{\boldsymbol{x}}$, corresponds to $f_{1T}^{\perp} < 0$.

years the Sivers mechanism was considered ruled out.

The situation changed in 2002. Brodsky, Hwang and Schmidt computed the SIDIS single-spin asymmetry in a simple spectator model and found a nonzero, leading-power result [47]. The asymmetry came from the final-state interaction between the struck quark and the target remnant: a single gluon exchange provides the phase needed for a spin asymmetry. Collins then showed where the 1993 argument had gone wrong [48]. A gauge-invariant definition of a TMD requires a Wilson line, the gauge link, that connects the two quark fields in the correlator. The link resums the gluon exchanges between the struck quark and the remnant, and its path depends on the process. In SIDIS the struck quark leaves the nucleon, the interactions are in the final state, and the link points to the future. In Drell-Yan production the quark annihilates with an antiquark from the other hadron, the interactions are in the initial state, and the link points to the past [49, 50, 51]. Time reversal does not force the Sivers function to vanish. It exchanges the two link paths, and the result is a definite prediction,

$$f_{1T}^{\perp}(x, \boldsymbol{k}_{\rm T}^2)\Big|_{\rm SIDIS} = -f_{1T}^{\perp}(x, \boldsymbol{k}_{\rm T}^2)\Big|_{\rm DY} \,. \tag{1.2}$$

The Sivers function is therefore not universal in the usual sense. It is universal up to a sign that is fixed by the color flow of the process [48, 52]. Testing Eq. (1.2) is a direct test of the gauge structure of QCD and of TMD factorization. The same reasoning applies to the Boer-Mulders function $h_1^{\perp}$, which correlates the transverse spin of a quark with its transverse momentum in an unpolarized nucleon [53].

The Sivers function is connected to several other parts of hadron physics. Burkardt pointed out that the Sivers shift can be understood as the combination of a spatial distortion of the quark distribution in a transversely polarized nucleon, which is related to the GPD E and thus to the anomalous magnetic moments, and an attractive final-state interaction. This "chromodynamic lensing" picture predicts opposite signs for up and down quarks [54, 55]. Because the transverse momentum of the nucleon as a whole must vanish, the Sivers shifts of all quarks and gluons have to sum to zero (the Burkardt sum rule) [56]. The Sivers function also links to the collinear twist-3 quark-gluon correlation of Qiu and Sterman [57], which describes SSAs at large transverse momentum. The two descriptions agree in their overlap region [58]. Finally, the Sivers effect requires interference between amplitudes in which the quark carries different orbital angular momentum, which ties it to the spin question raised by the EMC.

1.4 Experimental and phenomenological status

In SIDIS, $\ell N^{\uparrow} \to \ell' h X$, the Sivers function produces a characteristic azimuthal modulation $\sin(\phi_h - \phi_S)$ in the angle between the transverse momentum of the detected hadron and the transverse spin of the target [59]. The HERMES experiment at DESY reported the first clear observation of a nonzero Sivers asymmetry for positive pions produced off a transversely polarized proton target [60], with a final analysis of the full data set published later [61]. The COMPASS experiment at CERN found asymmetries compatible with zero on a deuteron target [62], as expected if the up and down quark Sivers functions have opposite signs and nearly cancel in an isoscalar target, and clear nonzero asymmetries on a proton target [63]. Measurements on a polarized ^{3}He target at Jefferson Lab, which acts as an effective polarized neutron, added information on the flavor structure [64].

The sign change of Eq. (1.2) requires a Drell-Yan-like measurement. The STAR collaboration at RHIC measured the transverse single-spin asymmetry in $W^{\pm}$ and Z^0 production in polarized proton collisions [65], and COMPASS performed the first polarized Drell-Yan measurement with a pion beam on a transversely polarized proton target [66], together with a SIDIS analysis at matching hard scales [67]. Both results favor the predicted sign change, but the statistical significance is still limited.

Phenomenological analyses combine these data to extract the Sivers function. Early fits used the parton model with Gaussian transverse momentum dependence [68, 69]. Later analyses included TMD evolution at

increasing perturbative accuracy [70, 71, 72, 73], and a global analysis showed that the SSAs in SIDIS, Drell-Yan, e^+e^- annihilation and proton-proton collisions can be described with a common set of non-perturbative functions [74]. The extractions agree that the up quark Sivers function is negative and the down quark function is positive in SIDIS, consistent with lensing. They are much less certain about the shape. The data cover a limited range of x, and the x dependence and the $\boldsymbol{k}_{\mathrm{T}}$ dependence enter the fits through assumed functional forms, so the extracted shapes carry a model bias [75]. Sea quark and gluon Sivers functions are poorly constrained.

New data are coming. The 12 GeV upgrade of Jefferson Lab extends SIDIS measurements to high x with high luminosity [76], and the Electron-Ion Collider will measure the Sivers asymmetry over a wide kinematic range with polarized protons and light nuclei [36, 37]. A first-principles determination of the x dependence of the Sivers function would provide a benchmark for these measurements that does not depend on fit assumptions.

1.5 Lattice QCD and the x-dependence problem

Lattice QCD is the only known method that computes hadron matrix elements from the QCD Lagrangian with systematically improvable uncertainties [77, 78, 2]. Space-time is discretized on a four-dimensional Euclidean grid, the path integral is evaluated by Monte Carlo sampling, and the continuum theory is recovered as the lattice spacing goes to zero. Lattice calculations of nucleon form factors and of low Mellin moments of PDFs are well established [79, 80, 81].

Parton distributions are harder. They are defined through operators whose fields are separated along a light-like direction. Such separations cannot be represented at a single Euclidean time, so a lattice calculation cannot evaluate the defining matrix elements directly. The traditional way around this problem uses the operator product expansion: Mellin moments $\int dx\, x^{n-1} f(x)$ are matrix elements of local operators and can be computed on the lattice. In practice only the lowest few moments are accessible, because the reduced symmetry of the hypercubic lattice causes higher-moment operators to mix with lower-dimensional ones [79]. A few moments are not enough to reconstruct a function of x.

For TMDs the situation is different. The quark fields in a TMD correlator are separated in the transverse direction as well, and when the gauge link is taken slightly off the light cone, the whole operator can be boosted

to a frame where it lives at a single time. Musch, Hägler, Engelhardt, Negele and Schäfer turned this observation into a lattice method [82, 83, 1]. The TMD correlator is decomposed into Lorentz-invariant amplitudes, $\widetilde{A}_i$, that depend on scalar products such as b^2, $b{\cdot}P$ and the Collins-Soper parameter $\hat{\zeta}$. The amplitudes are computed in a convenient lattice frame and then used, by Lorentz invariance, in the frame that defines the TMDs. Soft factors and multiplicative renormalization constants cancel in suitable ratios of amplitudes, which yields well-defined observables such as the generalized Sivers shift. With staple-shaped gauge links this approach gave the first lattice results for the naively T-odd Sivers and Boer-Mulders shifts, including the sign change between SIDIS and Drell-Yan [1]. Later work extended these calculations to larger values of $\hat{\zeta}$, to the pion, and to different fermion discretizations [84, 85, 86]. A complementary line of research, large-momentum effective theory (LaMET), computes equal-time "quasi" distributions in highly boosted hadrons and matches them to light-cone distributions [87, 88, 89]. It has produced, among other results, lattice determinations of the Collins-Soper evolution kernel [90, 91, 92, 93].

The published Sivers shift calculations in the Lorentz-invariant approach are restricted to $b{\cdot}P = 0$. The variable $b{\cdot}P$ is Fourier conjugate to x, so setting it to zero amounts to integrating over x. These calculations therefore give the Sivers shift averaged over all quarks and antiquarks, not its dependence on the momentum fraction. The TMD Handbook, the community review written by the TMD Topical Collaboration, lists the extension to scans in $b{\cdot}P$ as a necessary next step for lattice TMD studies and shows only a preliminary exploration of it [75]. Taking that step is the subject of this thesis.

1.6 This thesis

We compute the isovector generalized Sivers shift $\langle k_y \rangle_{TU}$ in lattice QCD as a function of the momentum fraction x. The calculation uses staple-shaped gauge links and the Lorentz-invariant amplitude method, and it scans nonzero values of $b{\cdot}P$ by giving the quark separation b a component along the nucleon momentum. The main elements of the work are the following.

1. We work out the kinematic constraints that the staple direction must satisfy when $b{\cdot}P \neq 0$. These constraints require off-axis staples. We construct them on the lattice by combining lattice paths and

interpolating between the accessible directions, and we separate the b-even (real) and b-odd (imaginary) parts of the amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ (Chapter 5).

2. We map the lattice data, taken at discrete Cartesian separations, onto the Lorentz-invariant variables $b{\cdot}P$ and b^2, restricting the data to $|\boldsymbol{b}| \geq 3a$ to reduce discretization artifacts.

3. We show that a Gaussian description of the amplitudes, of the kind used in earlier lattice TMD studies [83], leads to amplitudes that are not integrable in $b{\cdot}P$, so that the Fourier transform to x does not exist. Using physics-informed symbolic regression [94], we find a power-law form, $(1 + b^2)^{-j}$ in the relevant variables, that decays properly at large separations and describes the data well. We then fit all amplitudes simultaneously with their full covariance (Chapter 6).

4. We Fourier transform the fitted amplitudes to obtain continuous functions of x, extrapolate them in the Collins-Soper parameter toward the light-cone limit $\hat{\zeta} \to \infty$, and test whether the result factorizes into separate functions of x and $|\boldsymbol{b}_{\mathrm{T}}|$ (Chapter 7).

1.7 Outline

Chapter 2 collects the theoretical background: our conventions, the path integral formulation of quantum field theory, QCD and its gauge symmetry, Wilson lines, asymptotic freedom and confinement, deep inelastic scattering and collinear PDFs, SIDIS and Drell-Yan kinematics, the definition of TMDs with gauge links, the sign change of the Sivers function, rapidity divergences and the soft factor, and the approaches used to compute TMDs on the lattice.

Chapter 3 describes lattice QCD as used in this work: the discretized gauge and fermion actions, two- and three-point correlation functions, non-local operators with staple-shaped gauge links, and the ratio method used to extract nucleon matrix elements.

Chapter 4 presents the theoretical framework of the calculation in detail. It defines the TMD correlator and its soft factor, the staple link geometry and the Collins-Soper parameter $\hat{\zeta}$, the symmetry constraints on the correlator, and the decomposition into the Lorentz-invariant amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ from which f_1 and $f_{1T}^{\perp}$ are reconstructed.

Chapter 5 explains how the x dependence is extracted. It covers the kinematic constraints for $b{\cdot}P \neq 0$, the off-axis staple directions and their interpolation, the solution of the linear system for the amplitudes, and the transformation of the lattice data to Lorentz-invariant variables.

Chapter 6 develops the fit models. It explains why a Gaussian ansatz fails, introduces the short-distance cut, describes the symbolic regression search, tests factorization in $b{\cdot}P$ and b^2, and presents the simultaneous correlated fits.

Chapter 7 extrapolates the results to the light-cone limit $\hat{\zeta} \to \infty$ and tests the factorization of the x-dependent Sivers shift in x and $|\boldsymbol{b}_\mathrm{T}|$.

Chapter 8 summarizes the results and discusses future directions.

2 QCD AND HADRON STRUCTURE

This chapter collects the theoretical background for the rest of the thesis. We start with conventions and with the path integral formulation of quantum field theory, which is the starting point of lattice QCD. We then introduce QCD, its gauge symmetry and the Wilson lines that make non-local operators gauge invariant, and we review asymptotic freedom and confinement. The second half of the chapter turns to hadron structure. We describe deep inelastic scattering and the collinear parton distributions, then semi-inclusive DIS and the Drell-Yan process, which lead to transverse momentum dependent distributions. We define TMDs with process-dependent gauge links, explain why the Sivers function changes sign between SIDIS and Drell-Yan, discuss rapidity divergences and the soft factor, and introduce the generalized Sivers shift computed in this work. The chapter ends with an overview of how TMDs can be accessed in lattice QCD. The detailed parametrization used in our calculation is given in Chapter 4, and the lattice methods in Chapter 3.

2.1 Conventions

We use natural units, $\hbar = c = 1$, and the Minkowski metric $g^{\mu\nu} = \mathrm{diag}(+1, -1, -1, -1)$. Light-cone components of a four-vector w are defined by

$$w^{\pm} = \frac{1}{\sqrt{2}} \left(w^0 \pm w^3 \right) , \tag{2.1}$$

and we write $w = (w^+, w^-, \boldsymbol{w}_{\mathrm{T}})$, where the bold transverse vector $\boldsymbol{w}_{\mathrm{T}} = (w^1, w^2)$ is a Euclidean two-vector. Scalar products read

$$a{\cdot}b = a^+ b^- + a^- b^+ - \boldsymbol{a}_{\mathrm{T}}{\cdot}\boldsymbol{b}_{\mathrm{T}} \ . \tag{2.2}$$

It is convenient to introduce the light-like vectors $\bar{n} = (1, 0, 0, 1)/\sqrt{2}$ and $n = (1, 0, 0, -1)/\sqrt{2}$, which satisfy $\bar{n}^2 = n^2 = 0$ and $\bar{n}{\cdot}n = 1$, so that $w^+ = n{\cdot}w$ and $w^- = \bar{n}{\cdot}w$. A nucleon of mass m_N moving along the positive z axis with no transverse momentum has

$$P = P^+ \, \bar{n} + \frac{m_N^2}{2P^+} \, n \ , \tag{2.3}$$

and its spin vector is

$$S = \Lambda \, \frac{P^+}{m_N} \, \bar{n} - \Lambda \, \frac{m_N}{2P^+} \, n + \boldsymbol{S}_{\mathrm{T}} \ , \tag{2.4}$$

where Λ is the helicity and $\boldsymbol{S}_{\mathrm{T}}$ the transverse spin, with $\Lambda^2 + \boldsymbol{S}_{\mathrm{T}}^2 = 1$ for a pure state. The Levi-Civita tensor is normalized as $\epsilon^{0123} = +1$, and the transverse tensor $\epsilon_{ij} \equiv \epsilon_{-+ij}$ has $\epsilon_{12} = +1$. We use $\gamma^{\pm} = (\gamma^0 \pm \gamma^3)/\sqrt{2}$, $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ and $\sigma^{\mu\nu} = \frac{i}{2}[\gamma^\mu, \gamma^\nu]$. Nucleon states are normalized covariantly, $\langle P', S'|P, S\rangle = 2E_P\,(2\pi)^3\delta^3(\boldsymbol{P} - \boldsymbol{P'})\,\delta_{SS'}$. These conventions follow Ref. [1]. The relations between Minkowski and Euclidean gamma matrices used in the lattice analysis are listed in Section 5.6.

In this convention a parton with light-cone momentum fraction x has $k^+ = xP^+$. Some references, including the TMD Handbook [75], place the antiquark field at b and the quark field at the origin, which reverses the sign of b relative to our definitions. We always write the operator as $\bar{q}(0)\,\Gamma\,\mathcal{U}\,q(b)$, as in Refs. [83, 1].

2.2 Quantum field theory and the path integral

A quantum field theory is fully specified by its correlation functions, the vacuum expectation values of time-ordered products of field operators. For a scalar field ϕ with action $S[\phi] = \int d^4x\,\mathcal{L}(\phi, \partial\phi)$, the path integral expresses them as [95]

$$\langle\Omega|\,T\{\phi(x_1)\cdots\phi(x_n)\}\,|\Omega\rangle = \frac{\int \mathcal{D}\phi\;\phi(x_1)\cdots\phi(x_n)\;e^{iS[\phi]}}{\int \mathcal{D}\phi\;e^{iS[\phi]}}\;. \tag{2.5}$$

The oscillating weight e^{iS} makes this expression hard to define and unsuitable for numerical evaluation. Both problems are removed by a rotation to imaginary time. Setting $x^0 = -ix_4$ with real x_4, the action turns into $iS \to -S_E$, where S_E is the Euclidean action, and correlation functions become

$$\langle\mathcal{O}_1\cdots\mathcal{O}_n\rangle_E = \frac{1}{Z}\int \mathcal{D}\phi\;\mathcal{O}_1\cdots\mathcal{O}_n\;e^{-S_E[\phi]}\;, \qquad Z = \int \mathcal{D}\phi\;e^{-S_E[\phi]}\;. \tag{2.6}$$

For a bosonic theory with a real, bounded-below action, e^{-S_E}/Z is a probability distribution over field configurations. This is what allows Monte Carlo methods to be used once space-time is discretized, as described in Chapter 3.

The Euclidean formulation has a simple connection to the spectrum of the Hamiltonian H. For an operator $\mathcal{O}$ with Euclidean time dependence $\mathcal{O}(\tau) = e^{H\tau}\mathcal{O}(0)e^{-H\tau}$, inserting a complete set of energy eigenstates gives

$$\langle\Omega|\,\mathcal{O}(\tau)\,\mathcal{O}^\dagger(0)\,|\Omega\rangle = \sum_n \left|\langle\Omega|\mathcal{O}|n\rangle\right|^2 e^{-(E_n - E_\Omega)\tau}\;. \tag{2.7}$$

At large Euclidean time, the lowest state with nonzero overlap dominates. Hadron masses and matrix elements are extracted from lattice correlation functions in exactly this way (Chapter 3).

Fermions are described by anticommuting Grassmann fields ψ and $\bar{\psi}$. For an action bilinear in the fermion fields, $S_F = \int d^4x\, \bar{\psi}\, M\, \psi$, the Grassmann integral can be done exactly,

$$\int \mathcal{D}\bar{\psi}\, \mathcal{D}\psi\; e^{-\bar{\psi} M \psi} = \det M \;, \qquad \langle \psi(x)\, \bar{\psi}(y) \rangle = M^{-1}(x,y) \;. \tag{2.8}$$

In QCD, M is the Dirac operator in a background gluon field, so integrating out the quarks leaves a path integral over gluon fields only, weighted by $\det M$ times the exponential of the gauge action. Correlation functions of quark operators become products of quark propagators M^{-1} averaged over gluon configurations.

2.3 Gauge invariance and QCD

QCD is a non-Abelian gauge theory with gauge group $SU(3)$ [11, 12, 13, 14]. Each quark flavor f is a Dirac spinor carrying a color index in the fundamental representation. The gluon field $A_\mu = A_\mu^a t^a$ takes values in the Lie algebra, where the generators $t^a = \lambda^a/2$, $a = 1, \ldots, 8$, are built from the Gell-Mann matrices λ^a and satisfy

$$[t^a, t^b] = i f^{abc}\, t^c \;, \qquad \mathrm{tr}\left(t^a t^b\right) = \frac{1}{2}\delta^{ab} \;, \tag{2.9}$$

with the $SU(3)$ structure constants f^{abc}. The QCD Lagrangian density is

$$\mathcal{L}_{\mathrm{QCD}} = \sum_f \bar{q}_f(x)\, (i\gamma^\mu D_\mu - m_f)\, q_f(x) - \frac{1}{4} F^a_{\mu\nu}(x) F^{a\,\mu\nu}(x) \;, \tag{2.10}$$

with the covariant derivative and field strength tensor

$$D_\mu = \partial_\mu - i g\, A^a_\mu\, t^a \;, \tag{2.11}$$

$$F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g\, f^{abc} A^b_\mu A^c_\nu \;. \tag{2.12}$$

Here g is the gauge coupling and m_f the quark mass of flavor f. The last term in $F^a_{\mu\nu}$ makes the gluons interact with each other. The photon of quantum electrodynamics carries no charge and has no such self-interaction.

The Lagrangian is invariant under local $SU(3)$ transformations $V(x) = \exp\left(i\alpha^a(x) t^a\right)$,

$$\begin{aligned} q(x) &\to V(x)\, q(x) \;, \qquad \bar{q}(x) \to \bar{q}(x)\, V^\dagger(x) \;, \\ A_\mu(x) &\to V(x) A_\mu(x) V^\dagger(x) - \frac{i}{g}\big(\partial_\mu V(x)\big) V^\dagger(x) \;, \end{aligned} \tag{2.13}$$

under which $D_\mu q \to V D_\mu q$ and $F_{\mu\nu} \to V F_{\mu\nu} V^\dagger$. Physical observables must be gauge invariant.

Gauge invariance has an immediate consequence for the operators that define parton distributions. A local bilinear such as $\bar{q}(x)\Gamma q(x)$ is gauge invariant, but a non-local bilinear $\bar{q}(x)\Gamma q(y)$ with $x \neq y$ is not, because the two fields transform with V at different points. The fields have to be connected by a Wilson line, the parallel transporter of color along a path. For a path $\mathcal{C}$ that starts at x and ends at y we define

$$\mathcal{U}[\mathcal{C}] = \mathcal{P} \exp\left(-ig \int_{\mathcal{C}} dz^\mu \, A_\mu(z)\right), \tag{2.14}$$

where the path ordering $\mathcal{P}$ places fields at points closer to the start of the path to the left. This Wilson line transforms as

$$\mathcal{U}[\mathcal{C}] \to V(x)\,\mathcal{U}[\mathcal{C}]\,V^\dagger(y) \;, \tag{2.15}$$

so the operator

$$\bar{q}(x)\,\Gamma\,\mathcal{U}[\mathcal{C}]\,q(y) \tag{2.16}$$

is gauge invariant for any path from x to y and any Dirac matrix Γ. For a path made of straight segments through the points $x, z_1, \ldots, z_n, y$ we write $\mathcal{U}[x, z_1, \ldots, z_n, y] = \mathcal{U}[x, z_1]\,\mathcal{U}[z_1, z_2]\cdots\mathcal{U}[z_n, y]$, the notation used for the staple-shaped links in Eq. (4.11). The operator depends on the path, not only on the endpoints. Which path is appropriate cannot be decided by gauge invariance alone; for parton distributions it is fixed by the scattering process, as we explain in Section 2.8.

A Wilson line along a closed path is a Wilson loop, $W(\mathcal{C}) = \frac{1}{3}\mathrm{tr}\,\mathcal{U}[\mathcal{C}]$, which is gauge invariant by itself. Wilson loops are the basic objects of Wilson's lattice formulation [77]. On the lattice, the gluon field is represented by link variables $U_\mu(x)$, which are the Wilson lines of single lattice links. The non-local operators of Eq. (2.16) are then products of link variables along a path on the lattice, which makes them straightforward to evaluate.

The smallest Wilson loop on a lattice with spacing a is the plaquette $U_{\mu\nu}(x)$, the product of the four links around an elementary square in the $\mu\nu$ plane. For small a it can be written as $U_{\mu\nu}(x) = \exp\big(\pm iga^2 F_{\mu\nu}(x) + \mathcal{O}(a^3)\big)$, with the sign fixed by the orientation of the loop, and expanding the exponential gives

$$1 - \frac{1}{3}\,\mathrm{Re}\,\mathrm{tr}\,U_{\mu\nu}(x) = \frac{g^2 a^4}{6}\,\mathrm{tr}\big(F_{\mu\nu}(x)F_{\mu\nu}(x)\big) + \mathcal{O}(a^5) \qquad (\text{no sum over } \mu,\nu)\;. \tag{2.17}$$

Summing this expression over all plaquettes reproduces the Euclidean Yang-Mills action in the continuum limit. This is the origin of the Wilson gauge action used in Chapter 3.

2.4 Asymptotic freedom, confinement and chiral symmetry

Quantum corrections make the QCD coupling depend on the momentum scale μ at which it is defined. With $\alpha_s = g^2/(4\pi)$, the renormalization group equation reads

$$\mu^2 \frac{d\alpha_s}{d\mu^2} = \beta(\alpha_s) = -\left(b_0\,\alpha_s^2 + b_1\,\alpha_s^3 + \cdots\right), \qquad b_0 = \frac{33 - 2n_f}{12\pi}\ , \tag{2.18}$$

where n_f is the number of quark flavors lighter than μ. For $n_f \leq 16$ the coefficient b_0 is positive, and at leading order

$$\alpha_s(\mu^2) = \frac{1}{b_0\,\ln\left(\mu^2/\Lambda^2_{\rm QCD}\right)}\ . \tag{2.19}$$

The coupling decreases logarithmically at large μ. This is asymptotic freedom [13, 14], and it is why processes with a large momentum transfer can be computed in perturbation theory. At the Z boson mass the coupling is $\alpha_s(M_Z) = 0.1180(9)$ [96]. The scale $\Lambda_{\rm QCD}$, a few hundred MeV, is generated dynamically; the classical Lagrangian with massless quarks contains no dimensionful parameter. As μ approaches $\Lambda_{\rm QCD}$ the coupling becomes of order one and the perturbative expansion breaks down.

At long distances QCD confines color. No isolated quark or gluon has been observed, and the hadron spectrum consists of color-singlet states. On the lattice, confinement shows up as an area law for large Wilson loops. For a rectangular loop of spatial extent R and temporal extent T,

$$\langle W(R,T)\rangle \xrightarrow{T\to\infty} e^{-V(R)\,T}\ , \qquad V(R) \approx \sigma R \quad (R \text{ large})\ , \tag{2.20}$$

where $V(R)$ is the potential between a static quark and antiquark and σ is the string tension [77, 97]. The same kind of Wilson line self-interaction appears in the staple-shaped links of TMD operators, where it produces a dependence on the staple length that has to be removed by forming ratios (Section 2.9).

The third non-perturbative feature is the spontaneous breaking of chiral symmetry. For massless u and d quarks the Lagrangian is invariant under separate $SU(2)$ rotations of left- and right-handed quark fields. The QCD vacuum has a nonzero quark condensate $\langle\bar{q}q\rangle$, which breaks this symmetry down to the isospin subgroup.

The pions are the corresponding pseudo-Goldstone bosons, light because the quark masses are small, and the Gell-Mann-Oakes-Renner relation $f_\pi^2 m_\pi^2 = -(m_u+m_d)\langle\bar{q}q\rangle$ links their mass to the condensate. Chiral symmetry breaking also explains why the nucleon, with a mass of about 0.94 GeV, is much heavier than the sum of its valence quark masses. Its mass, like its internal structure, comes from the non-perturbative dynamics of QCD.

These three properties set the problem this thesis addresses. Asymptotic freedom makes hard scattering processes calculable, so that experiments can be interpreted in terms of parton distributions. Confinement and chiral symmetry breaking make the distributions themselves non-perturbative, so that computing them requires a method that does not rely on a small coupling. Lattice QCD is such a method.

2.5 Deep inelastic scattering and the parton model

In deep inelastic lepton-nucleon scattering, $\ell(l) + N(P) \to \ell(l') + X$, the lepton exchanges a virtual photon of momentum $q = l - l'$ with the nucleon, and the final hadronic state X is not observed. The standard kinematic variables are

$$Q^2 = -q^2 \,, \qquad x_B = \frac{Q^2}{2P\cdot q} \,, \qquad y = \frac{P\cdot q}{P\cdot l} \,. \tag{2.21}$$

The hadronic part of the cross section is contained in the hadronic tensor,

$$W^{\mu\nu} = \frac{1}{4\pi}\int d^4\xi \; e^{iq\cdot\xi} \, \langle P| \left[J^\mu(\xi), J^\nu(0)\right] |P\rangle \,, \tag{2.22}$$

where J^μ is the electromagnetic current. Current conservation, parity and Lorentz invariance allow two structure functions for an unpolarized target,

$$W^{\mu\nu} = \left(-g^{\mu\nu} + \frac{q^\mu q^\nu}{q^2}\right) F_1(x_B, Q^2) + \left(P^\mu - \frac{P\cdot q}{q^2}\, q^\mu\right)\left(P^\nu - \frac{P\cdot q}{q^2}\, q^\nu\right) \frac{F_2(x_B, Q^2)}{P\cdot q} \,, \tag{2.23}$$

and, neglecting the nucleon mass, the cross section is

$$\frac{d^2\sigma}{dx_B\, dQ^2} = \frac{4\pi\alpha^2}{x_B\, Q^4}\left[(1-y)\, F_2(x_B, Q^2) + x_B\, y^2\, F_1(x_B, Q^2)\right] . \tag{2.24}$$

In the parton model the photon scatters elastically off a quark that carries a fraction x of the nucleon's light-cone momentum. Momentum conservation at the photon vertex then sets $x = x_B$ up to corrections suppressed by $1/Q^2$, and the structure functions become

$$F_2(x_B) = 2x_B F_1(x_B) = x_B \sum_q e_q^2 \left[f_1^q(x_B) + f_1^{\bar{q}}(x_B)\right] , \tag{2.25}$$

where e_q is the quark charge in units of the proton charge and $f_1^q(x)$ is the unpolarized PDF of quark flavor q [7, 6, 8]. The first equality, the Callan-Gross relation, reflects the spin 1/2 of the quarks. The corresponding "handbag" diagram is shown in Figure 3.



Figure 3: Handbag diagram for the leading contribution to the DIS hadronic tensor. The virtual photon scatters off a quark of momentum k with $k^+ = xP^+$, and the lower blob is the quark-quark correlator whose γ^+ projection gives the PDF $f_1(x)$. The dashed line indicates the sum over final states.

QCD turns the parton model into a controlled approximation. The collinear factorization theorem states that, up to power corrections in $1/Q$, the structure functions are convolutions of perturbatively calculable coefficient functions with PDFs defined as matrix elements of quark and gluon operators [21, 23]. For the unpolarized quark distribution this definition is

$$f_1^q(x) = \frac{1}{2}\int \frac{db^-}{2\pi}\; e^{ixP^+b^-}\,\langle P|\,\bar q(0)\,\gamma^+\,\mathcal{U}[0,nb^-]\,q(nb^-)\,|P\rangle\;. \tag{2.26}$$

The quark fields are separated along the light-cone direction n, and the straight Wilson line along n between them makes the operator gauge invariant. In light-cone gauge $A^+ = 0$ this Wilson line is equal to one, and $f_1^q(x)$ has the interpretation of a number density of quarks with plus-momentum xP^+ [23, 98]. Eq. (2.26) has support on $-1 \le x \le 1$. Positive x corresponds to quarks, and negative x to antiquarks, with $f_1^{\bar q}(x) = -f_1^q(-x)$ for $x > 0$.

Replacing γ^+ in Eq. (2.26) by $\gamma^+\gamma_5$ or by $i\sigma^{i+}\gamma_5$ defines the helicity distribution $g_1(x)$ of quarks in a longitudinally polarized nucleon and the transversity distribution $h_1(x)$ of transversely polarized quarks in a transversely polarized nucleon. Their integrals over x are local matrix elements. For the isovector combination $u-d$ they give the nucleon axial charge, $\int dx\, g_1^{u-d}(x) = g_A$, and the tensor charge, $\int dx\, h_1^{u-d}(x) = g_T$. These three collinear distributions reappear as the $\boldsymbol{k}_\mathrm{T}$-integrals of TMDs in Section 2.8.

The operator in Eq. (2.26) contains ultraviolet divergences, and its renormalization introduces a scale μ.

The resulting dependence of $f_1(x,\mu)$ on μ is governed by the DGLAP equations [99, 100, 101],

$$\frac{df_1^q(x,\mu)}{d\ln\mu^2} = \frac{\alpha_s(\mu^2)}{2\pi}\int_x^1 \frac{dz}{z}\left[P_{qq}(z)\, f_1^q\Big(\frac{x}{z},\mu\Big) + P_{qg}(z)\, f_1^g\Big(\frac{x}{z},\mu\Big)\right], \tag{2.27}$$

with perturbatively known splitting functions $P_{ij}(z)$. Evolution in μ is perturbative, but the distributions at any one scale are not.

For lattice QCD, an important property of Eq. (2.26) is its relation to local operators. Expanding the non-local operator around $b^- = 0$ produces the Mellin moments,

$$\int_{-1}^1 dx\; x^{n-1} f_1^q(x) = \langle x^{n-1}\rangle_q\;, \qquad \langle P|\,\mathcal{O}_q^{\{\mu_1\cdots\mu_n\}}\,|P\rangle = 2\,\langle x^{n-1}\rangle_q\; P^{\{\mu_1}\cdots P^{\mu_n\}}\;, \tag{2.28}$$

where $\mathcal{O}_q^{\{\mu_1\cdots\mu_n\}} = \bar{q}\,\gamma^{\{\mu_1} iD^{\mu_2}\cdots iD^{\mu_n\}}\,q$ is a local twist-two operator, symmetrized in its indices and with traces removed. Local operators have no extent in time and can be computed on a Euclidean lattice, which is how lattice QCD has traditionally accessed PDFs [79, 80]. In practice only the first few moments are available. On a hypercubic lattice, the operators for higher moments mix with operators of lower dimension, with coefficients that diverge as powers of the inverse lattice spacing. A handful of moments does not determine the x dependence of a distribution.

2.6 Form factors, GPDs and the transverse structure of the nucleon

PDFs describe the longitudinal momentum of partons. Information about their transverse position comes from elastic form factors and their generalization to GPDs. Because the lensing picture of the Sivers effect (Section 1.3) is built on this spatial information, we summarize the relevant definitions here.

The matrix element of the electromagnetic current between nucleon states of momenta P and $P' = P + \Delta$ is parametrized by the Dirac and Pauli form factors,

$$\langle P'|\,J^\mu(0)\,|P\rangle = \bar{u}(P')\left[\gamma^\mu F_1(t) + \frac{i\sigma^{\mu\nu}\Delta_\nu}{2m_N}\,F_2(t)\right]u(P)\;, \qquad t = \Delta^2\;. \tag{2.29}$$

At $t = 0$, F_1 gives the electric charge and F_2 the anomalous magnetic moment κ. The Sachs combinations $G_E = F_1 + \frac{t}{4m_N^2}F_2$ and $G_M = F_1 + F_2$ are what elastic scattering experiments measure, and the slope of G_E at $t = 0$ defines the charge radius, $\langle r^2\rangle = 6\,dG_E/dt|_{t=0}$.

GPDs are off-forward generalizations of PDFs. For unpolarized quarks they are defined by [16, 18, 17, 19]

$$\frac{1}{2}\int \frac{dz^-}{2\pi}\, e^{ix\bar{P}^+z^-}\, \langle P'|\, \bar{q}\big(-\tfrac{z}{2}\big)\, \gamma^+\, \mathcal{U}\, q\big(\tfrac{z}{2}\big)\, |P\rangle\Big|_{z^+=0,\, \boldsymbol{z}_\mathrm{T}=0} = \frac{1}{2\bar{P}^+}\left[H^q\, \bar{u}(P')\gamma^+ u(P) + E^q\, \bar{u}(P')\frac{i\sigma^{+\nu}\Delta_\nu}{2m_N}u(P)\right], \tag{2.30}$$

where $\bar{P} = (P + P')/2$, $\mathcal{U}$ is the straight light-like Wilson line between the quark fields, and H^q and E^q depend on x, the skewness $\xi = -\Delta^+/(2\bar{P}^+)$ and t. GPDs connect several quantities discussed so far. In the forward limit $H^q(x,0,0) = f_1^q(x)$. Their first moments are the form factors of Eq. (2.29), $\int dx\, H^q = F_1^q(t)$ and $\int dx\, E^q = F_2^q(t)$. Their second moments give the total angular momentum carried by quarks through Ji's sum rule [18],

$$J^q = \frac{1}{2}\int_{-1}^{1} dx\; x\,[H^q(x,\xi,0) + E^q(x,\xi,0)]\,, \tag{2.31}$$

which has been one of the main routes to the spin decomposition of the nucleon [40].

At $\xi = 0$ the momentum transfer is purely transverse, and the Fourier transform of H^q with respect to $\boldsymbol{\Delta}_\mathrm{T}$ gives the density of quarks with momentum fraction x at transverse distance $\boldsymbol{b}_\mathrm{T}$ from the center of momentum of the nucleon [20],

$$q(x, \boldsymbol{b}_\mathrm{T}) = \int \frac{d^2\boldsymbol{\Delta}_\mathrm{T}}{(2\pi)^2}\; e^{-i\boldsymbol{b}_\mathrm{T}\cdot\boldsymbol{\Delta}_\mathrm{T}}\; H^q(x,0,-\boldsymbol{\Delta}_\mathrm{T}^2)\;. \tag{2.32}$$

In a transversely polarized nucleon the density acquires a dipole term proportional to the Fourier transform of E^q. The quark distribution is shifted perpendicular to the spin, and the average shift of flavor q is set by its contribution to the anomalous magnetic moment, $\kappa_q/(2m_N)$ [54]. From the measured magnetic moments of the proton and neutron one finds $\kappa_u \approx 1.67$ and $\kappa_d \approx -2.03$, so up and down quarks are shifted in opposite directions. Burkardt argued that the final-state interaction in SIDIS is on average attractive, so that it converts this spatial shift into a transverse momentum shift in the opposite direction. This "chromodynamic lensing" picture relates the sign of the Sivers function of each flavor to the sign of κ_q [54, 55]. It is a model-dependent argument, but it predicts opposite signs for the up and down quark Sivers functions, in agreement with the experimental extractions.

2.7 SIDIS, Drell-Yan and TMD factorization

Transverse momentum dependent distributions enter processes in which a transverse momentum much smaller than the hard scale is measured. The two standard examples are semi-inclusive DIS and the Drell-Yan process.

In semi-inclusive DIS, $\ell(l)+N(P,S) \to \ell(l')+h(P_h)+X$, one hadron h in the final state is detected in addition to the scattered lepton. Besides x_B, y and Q^2, the kinematics are described by the fraction $z = P{\cdot}P_h/P{\cdot}q$ of the virtual photon energy carried by the hadron and by the transverse momentum $\boldsymbol{P}_{hT}$ of the hadron in the frame where the nucleon and the virtual photon move along the z axis. The azimuthal angles ϕ_h of $\boldsymbol{P}_{hT}$ and ϕ_S of the target spin $\boldsymbol{S}_{\rm T}$ are measured around the photon direction, following the Trento conventions [45]. For $|\boldsymbol{P}_{hT}| \ll Q$, the cross section on a transversely polarized target has the structure [59]

$$d\sigma \propto F_{UU,T} + \ldots + |\boldsymbol{S}_{\rm T}| \, \sin(\phi_h - \phi_S) \, F_{UT,T}^{\sin(\phi_h-\phi_S)} + \ldots \ , \tag{2.33}$$

where the dots denote other azimuthal modulations and polarization states. At leading power, the structure functions are convolutions of a TMD with a transverse momentum dependent fragmentation function D_1,

$$F_{UU,T} = \mathcal{C}\left[f_1 D_1\right] \ , \qquad\qquad F_{UT,T}^{\sin(\phi_h-\phi_S)} = \mathcal{C}\left[-\frac{\hat{\boldsymbol{h}}{\cdot}\boldsymbol{k}_{\rm T}}{m_N} \, f_{1T}^{\perp} D_1\right], \tag{2.34}$$

with $\hat{\boldsymbol{h}} = \boldsymbol{P}_{hT}/|\boldsymbol{P}_{hT}|$ and

$$\mathcal{C}[w\, f\, D] = x_B \sum_q e_q^2 \int d^2\boldsymbol{k}_{\rm T} \, d^2\boldsymbol{p}_{\rm T} \, \delta^{(2)}\left(\boldsymbol{k}_{\rm T} - \boldsymbol{p}_{\rm T} - \frac{\boldsymbol{P}_{hT}}{z}\right) w(\boldsymbol{k}_{\rm T}, \boldsymbol{p}_{\rm T}) \ f^q(x_B, \boldsymbol{k}_{\rm T}^2) \ D^q(z, \boldsymbol{p}_{\rm T}^2) \ . \tag{2.35}$$

Here $\boldsymbol{k}_{\rm T}$ is the transverse momentum of the quark in the nucleon and $\boldsymbol{p}_{\rm T}$ the transverse momentum of the fragmenting quark relative to the observed hadron. The ratio $A_{UT}^{\sin(\phi_h-\phi_S)} = F_{UT,T}^{\sin(\phi_h-\phi_S)}/F_{UU,T}$ is the Sivers asymmetry measured by HERMES, COMPASS and Jefferson Lab (Section 1.4). The transverse momentum of the observed hadron receives contributions from the intrinsic motion of the quark in the nucleon and from the fragmentation process, and the convolution cannot be undone without additional input. This is one reason why independent information on TMDs, for example from lattice QCD, is useful.

In the Drell-Yan process, $h_a(P_a) + h_b(P_b) \to \ell^+\ell^-(q) + X$, a quark from one hadron annihilates with an antiquark from the other into a virtual photon or Z boson, which decays into a lepton pair of invariant mass Q and transverse momentum $\boldsymbol{q}_{\rm T}$ [15]. No fragmentation is involved, and for $|\boldsymbol{q}_{\rm T}| \ll Q$ the cross section is

determined by two TMDs. The TMD factorization theorem for this process takes the form [26, 23, 75]

$$\frac{d\sigma}{dQ^2\,dy\,d^2\boldsymbol{q}_{\rm T}} = \sigma_0 \sum_q e_q^2\, H(Q,\mu) \int \frac{d^2\boldsymbol{b}_{\rm T}}{(2\pi)^2}\; e^{-i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{q}_{\rm T}}\; \tilde{f}_1^q(x_a,\boldsymbol{b}_{\rm T};\mu,\zeta_a)\; \tilde{f}_1^{\bar{q}}(x_b,\boldsymbol{b}_{\rm T};\mu,\zeta_b) + (q\leftrightarrow\bar{q}) + \mathcal{O}\!\left(\frac{|\boldsymbol{q}_{\rm T}|}{Q}\right), \tag{2.36}$$

where σ_0 is the Born cross section, H a perturbative hard factor, y the rapidity of the lepton pair, and $\tilde{f}_1(x,\boldsymbol{b}_{\rm T})$ the Fourier transform of the TMD in the transverse momentum, Eq. (4.8). The scales ζ_a and ζ_b, with $\zeta_a\zeta_b = Q^4$, arise from the rapidity divergences discussed in Section 2.9. Transverse position space is the natural setting for TMD factorization: the convolution in $\boldsymbol{k}_{\rm T}$ becomes a product, and evolution is multiplicative. Similar factorization theorems hold for SIDIS and for e^+e^- annihilation into two hadrons [30, 23]. A single-spin asymmetry in Drell-Yan with a transversely polarized hadron measures the Sivers function of that hadron in the Drell-Yan process, which makes the sign change of Eq. (1.2) testable.

2.8 The TMD correlator and gauge links

TMDs are defined through the quark-quark correlator

$$\Phi^{[\Gamma]}(x,\boldsymbol{k}_{\rm T};P,S;\mathcal{C}) = \int \frac{db^-\,d^2\boldsymbol{b}_{\rm T}}{(2\pi)^3}\; e^{ixP^+b^- - i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\; \frac{1}{2}\langle P,S|\,\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\,q(b)\,|P,S\rangle\Big|_{b^+=0}\,, \tag{2.37}$$

which generalizes Eq. (2.26) by allowing a transverse separation $\boldsymbol{b}_{\rm T}$ between the quark fields. Integrating over $\boldsymbol{k}_{\rm T}$ sets $\boldsymbol{b}_{\rm T}=0$ and returns the collinear correlator. Eq. (2.37) is schematic: it omits the soft factor and the regularization of rapidity divergences, which are included in the full definition, Eq. (4.1). It is enough to discuss the gauge link and the classification of TMDs.

2.8.1 Process-dependent gauge links

For collinear PDFs the fields sit on a light-like line, and the straight Wilson line along that line is the natural gauge-invariant choice. When the fields are separated transversely, no path is singled out by gauge invariance. The path is determined instead by the physics of the process [48, 50, 51]. In SIDIS, the quark struck by the virtual photon moves out of the nucleon with a large momentum along the light-cone direction n. Before it hadronizes, it can exchange gluons with the colored remnant of the target. In the eikonal approximation each such exchange with a gluon of momentum l contributes a factor proportional to $n\cdot A/(n\cdot l + i\epsilon)$, and summing all exchanges builds a Wilson line from the position of the quark field to infinity along n. Connecting the two

quark fields, together with a transverse link at infinity that is needed for gauge invariance in light-cone gauge [50], gives the staple-shaped gauge link

$$\mathcal{U}[\mathcal{C}_b]_{\rm SIDIS} = \mathcal{U}[0,\ +\infty n,\ +\infty n + b,\ b]\ . \tag{2.38}$$

In Drell-Yan the active quark interacts with the other hadron before it annihilates. The gluon exchanges are in the initial state, the eikonal propagators become $1/(n{\cdot}l - i\epsilon)$, and the link runs to past infinity,

$$\mathcal{U}[\mathcal{C}_b]_{\rm DY} = \mathcal{U}[0,\ -\infty n,\ -\infty n + b,\ b]\ . \tag{2.39}$$

Both paths are illustrated in Figure 4. The two gauge links differ only in the sign of the $i\epsilon$ in the eikonal propagators. Writing $1/(n{\cdot}l \pm i\epsilon) = \mathrm{PV}\,1/(n{\cdot}l) \mp i\pi\,\delta(n{\cdot}l)$ shows that the imaginary part, the phase that a single-spin asymmetry requires, is opposite in the two processes. This is the diagrammatic origin of the sign change found by Brodsky, Hwang and Schmidt in their model calculation [47, 49].



(a) SIDIS: future-pointing (b) Drell-Yan: past-pointing

Figure 4: Staple-shaped gauge links connecting the quark fields at 0 and b in the TMD correlator, Eq. (2.37). (a) In SIDIS the link runs to $+\infty$ along the light-cone direction n of the outgoing quark, Eq. (2.38). (b) In Drell-Yan it runs to $-\infty$, Eq. (2.39). The dashed segment is the transverse link at infinity. In the lattice calculation the legs have finite length $\eta|v|$ along a space-like direction v, and the two cases correspond to $\eta v{\cdot}P \to +\infty$ and $\eta v{\cdot}P \to -\infty$ [1].

2.8.2 Leading-twist TMDs

At leading twist, the correlator is projected with the Dirac structures $\Gamma = \gamma^+$, $\gamma^+\gamma_5$ and $i\sigma^{i+}\gamma_5$, which select the "good" light-cone components of the quark field. Parity, hermiticity and Lorentz covariance then allow eight functions of x and $\boldsymbol{k}_{\rm T}^2$ [27, 29, 22, 53, 102]:

$$\Phi^{[\gamma^+]} = f_1 - \left[\frac{\epsilon_{ij}\,k_{T,i}\,S_{T,j}}{m_N}\,f_{1T}^{\perp}\right]_{\rm odd}\ , \tag{2.40}$$

$$\Phi^{[\gamma^+\gamma_5]} = \Lambda\,g_1 + \frac{\boldsymbol{k}_{\rm T}{\cdot}\boldsymbol{S}_{\rm T}}{m_N}\,g_{1T}\ , \tag{2.41}$$

$$\Phi^{[i\sigma^{i+}\gamma_5]} = S_{T,i}\,h_1 + \frac{\Lambda\,k_{T,i}}{m_N}\,h_{1L}^{\perp} - \frac{\left(k_{T,i}k_{T,j} - \frac{1}{2}\boldsymbol{k}_{\rm T}^2\delta_{ij}\right)S_{T,j}}{m_N^2}\,h_{1T}^{\perp} + \left[\frac{\epsilon_{ij}\,k_{T,j}}{m_N}\,h_1^{\perp}\right]_{\rm odd}\ . \tag{2.42}$$

The eight functions are summarized in Table 1. Three of them survive integration over $\boldsymbol{k}_{\mathrm{T}}$ and have collinear counterparts: the unpolarized distribution f_1, the helicity distribution g_1 and the transversity distribution h_1. The other five exist only because of the transverse momentum. The worm-gear functions g_{1T} and $h_{1L}^{\perp}$ and the pretzelosity $h_{1T}^{\perp}$ correlate the quark transverse momentum with the quark and nucleon polarizations. The Sivers function $f_{1T}^{\perp}$ and the Boer-Mulders function $h_1^{\perp}$, marked by the brackets $[\ ]_{\mathrm{odd}}$, are the naively T-odd functions.

nucleon \ quark	U	L	T
U	f_1		$\big[h_1^{\perp}\big]_{\mathrm{odd}}$
L		g_1	$h_{1L}^{\perp}$
T	$\big[f_{1T}^{\perp}\big]_{\mathrm{odd}}$	g_{1T}	$h_1,\ h_{1T}^{\perp}$

Table 1: Leading-twist quark TMDs, arranged by the polarization of the nucleon (rows) and of the quark (columns): unpolarized (U), longitudinally polarized (L) and transversely polarized (T). The Sivers function $f_{1T}^{\perp}$ and the Boer-Mulders function $h_1^{\perp}$ are naively T-odd.

The γ^+ projection has a density interpretation. In light-cone gauge, $\Phi^{[\gamma^+]}(x, \boldsymbol{k}_{\mathrm{T}}; P, S)$ counts quarks with momentum fraction x and transverse momentum $\boldsymbol{k}_{\mathrm{T}}$ in a nucleon with spin S, summed over quark helicities [23, 103]. Eq. (2.40) is therefore the density of Eq. (1.1). Positivity of this density implies the bound $(|\boldsymbol{k}_{\mathrm{T}}|/m_N)\,|f_{1T}^{\perp}(x, \boldsymbol{k}_{\mathrm{T}}^2)| \le f_1(x, \boldsymbol{k}_{\mathrm{T}}^2)$ [104].

2.8.3 Time reversal and the sign change

Time reversal is a symmetry of QCD, but it also acts on the gauge link. Combined with parity, it maps the correlator with a future-pointing staple onto the correlator with a past-pointing staple, $n \to -n$ [48, 51]. Under this map the T-even functions are unchanged, whereas the terms in brackets in Eqs. (2.40) and (2.42) change sign. Combined with the link directions of Eqs. (2.38) and (2.39), this gives

$$f_{1T}^{\perp}\big|_{\mathrm{SIDIS}} = -f_{1T}^{\perp}\big|_{\mathrm{DY}}\ , \qquad h_1^{\perp}\big|_{\mathrm{SIDIS}} = -h_1^{\perp}\big|_{\mathrm{DY}}\ , \qquad f^{\text{T-even}}\big|_{\mathrm{SIDIS}} = f^{\text{T-even}}\big|_{\mathrm{DY}}\ . \tag{2.43}$$

Collins's 1993 argument [46] used a straight gauge link, which is invariant under time reversal. With such a link the T-odd functions do vanish. The Sivers function is nonzero only because the physical gauge link extends to infinity in a direction set by the process.

The same structure appears in the lattice calculation. The staple of Eq. (4.11) has finite legs of length $\eta|v|$ along a direction v, and the SIDIS and DY cases correspond to $\eta v{\cdot}P \to +\infty$ and $\eta v{\cdot}P \to -\infty$ [1]. T-even quantities are even functions of η, T-odd quantities are odd, and T-odd quantities vanish for a straight link, $\eta = 0$. The symmetry constraints on the correlator that lead to these statements are worked out in Chapter 4.

2.9 Rapidity divergences, the soft factor and evolution

The staple-shaped links of Eqs. (2.38) and (2.39) cause a problem that does not arise for collinear PDFs. Wilson lines along an exactly light-like direction produce divergences from gluons with large rapidity, moving parallel to the Wilson line, which are not regulated by dimensional regularization [103, 105, 23]. These rapidity divergences cancel in the cross section, between the TMDs and a soft factor, but they have to be regulated in the definition of each TMD.

Collins and Soper introduced the regulator used in this work [25, 21]: the Wilson lines are tilted away from the light cone into a space-like direction v. The approach to the light cone is measured by the Lorentz-invariant parameter

$$\hat{\zeta} = \frac{v{\cdot}P}{\sqrt{|v^2|}\,\sqrt{P^2}} = \sinh(y_P - y_v) \;, \tag{2.44}$$

Eq. (4.12), where y_P and y_v are the rapidities of P and v. The light-cone limit is $\hat{\zeta} \to \infty$. A space-like v has an important advantage for the lattice: one can always boost to a frame in which v has no time component, so that the entire operator lies at a single time [1, 31].

The soft factor $\tilde{\mathcal{S}}$ is a vacuum expectation value of Wilson lines, essentially a Wilson loop made from two staples. Dividing the unsubtracted correlator by an appropriate combination of soft factors, Eq. (4.1), removes the rapidity divergences and the dependence on the regulator in the limit where the Wilson lines approach the light cone [23, 31]. The Wilson lines also have a self-energy that diverges linearly with the ultraviolet cutoff, proportional to the length of the path. On the lattice this divergence appears as a factor $e^{-\delta m\, L}$ for a path of length L, with $\delta m \propto 1/a$, together with additional divergences from the corners of the path [106, 83]. For a fixed path, both the soft factor and these renormalization factors multiply the correlator independently of the Dirac structure Γ [1]. They therefore cancel in ratios of amplitudes taken at the same b, P and staple geometry.

The observables computed in this thesis are ratios of this type.

After the rapidity divergences are removed, the TMDs depend on two scales: the renormalization scale μ and the rapidity scale ζ. In the convention of Collins, the dependence on them is described by the Collins-Soper equations [25, 26, 23],

$$\frac{\partial \ln \tilde{f}(x, \boldsymbol{b}_{\mathrm{T}}; \mu, \zeta)}{\partial \ln \sqrt{\zeta}} = \tilde{K}(|\boldsymbol{b}_{\mathrm{T}}|; \mu) \ , \qquad \frac{d\tilde{K}(|\boldsymbol{b}_{\mathrm{T}}|; \mu)}{d \ln \mu} = -\gamma_K\big(\alpha_s(\mu)\big) \ , \tag{2.45}$$

together with a renormalization group equation in μ. The Collins-Soper kernel $\tilde{K}$ is independent of x and of the hadron, and γ_K is the cusp anomalous dimension. For $|\boldsymbol{b}_{\mathrm{T}}| \lesssim 1/\Lambda_{\mathrm{QCD}}$ the kernel can be computed in perturbation theory. At larger $|\boldsymbol{b}_{\mathrm{T}}|$ it is non-perturbative, and it is one of the quantities that recent lattice calculations have determined [90, 91, 92, 93]. The evolution of the Sivers function in these scales has been worked out in Refs. [107, 70]. In the lattice calculation the scale ζ is represented by $\hat{\zeta}$, which is finite at accessible nucleon momenta, and the light-cone limit has to be reached by extrapolation (Chapter 7).

2.10 Transverse position space and the Sivers shift

Lattice calculations produce matrix elements at fixed separations b, so it is natural to work with TMDs in transverse position space. The Fourier transform $\tilde{f}(x, \boldsymbol{b}_{\mathrm{T}}^2)$ and its derivatives $\tilde{f}^{(n)}(x, \boldsymbol{b}_{\mathrm{T}}^2)$ with respect to $\boldsymbol{b}_{\mathrm{T}}^2$ are defined in Eqs. (4.8) and (4.9). As $|\boldsymbol{b}_{\mathrm{T}}| \to 0$ they reduce to the transverse momentum moments

$$f^{(n)}(x) = \int d^2\boldsymbol{k}_{\mathrm{T}} \left(\frac{\boldsymbol{k}_{\mathrm{T}}^2}{2m_N^2}\right)^n f(x, \boldsymbol{k}_{\mathrm{T}}^2) \ , \tag{2.46}$$

Eq. (4.10). For the Sivers function, the first moment $f_{1T}^{\perp(1)}(x)$ is the quantity that enters the transverse momentum weighted Sivers asymmetry. It is also ultraviolet divergent in QCD, because the perturbative large-$\boldsymbol{k}_{\mathrm{T}}$ tail of the Sivers function falls off too slowly [108]. A finite $|\boldsymbol{b}_{\mathrm{T}}|$ acts as a regulator, and we keep $|\boldsymbol{b}_{\mathrm{T}}|$ finite throughout [1, 109]. We also use the notation $f^{[m]}$ for x-moments, $f^{[1]} = \int_{-1}^{1} dx\, f(x)$.

In $\boldsymbol{b}_{\mathrm{T}}$ space, the γ^+ projection of the correlator contains two Lorentz-invariant amplitudes [1],

$$\frac{1}{2P^+}\, \tilde{\Phi}^{[\gamma^+]}_{\text{unsubtr.}} = \widetilde{A}_{2B} + im_N\, \epsilon_{ij}\, b_i\, S_j\ \widetilde{A}_{12B} \ , \tag{2.47}$$

where $\widetilde{A}_{2B}$ gives the unpolarized TMD and $\widetilde{A}_{12B}$ the Sivers function after the Fourier transform in $b{\cdot}P$ and division by the soft factor. Their definitions in terms of the full set of invariant amplitudes are given in

Eq. (4.34). At $b{\cdot}P = 0$ the Fourier transform in $b{\cdot}P$ reduces to an integral over x, and one finds [1]

$$\tilde{f}_1^{[1](0)}(\boldsymbol{b}_{\rm T}^2) = \frac{2\widetilde{A}_{2B}}{\tilde{\mathcal{S}}} \ , \qquad \tilde{f}_{1T}^{\perp[1](1)}(\boldsymbol{b}_{\rm T}^2) = -\frac{2\widetilde{A}_{12B}}{\tilde{\mathcal{S}}} \ , \tag{2.48}$$

with both amplitudes evaluated at $b^2 = -\boldsymbol{b}_{\rm T}^2$ and $b{\cdot}P = 0$.

The Sivers function describes the average transverse momentum of unpolarized quarks in the direction perpendicular to the nucleon spin. For a nucleon polarized along $\hat{\boldsymbol{x}}$, $\boldsymbol{S}_{\rm T} = (1,0)$, Eq. (2.40) gives $\Phi^{[\gamma^+]} = f_1 + (k_y/m_N)\, f_{1T}^{\perp}$, and the average k_y is $m_N f_{1T}^{\perp(1)}/f_1^{(0)}$. The generalized Sivers shift extends this quantity to finite $|\boldsymbol{b}_{\rm T}|$ [1, 109],

$$\langle k_y\rangle_{TU}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\eta v{\cdot}P) \equiv m_N\, \frac{\tilde{f}_{1T}^{\perp[1](1)}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\ldots,\eta v{\cdot}P)}{\tilde{f}_1^{[1](0)}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\ldots,\eta v{\cdot}P)} = -m_N\, \frac{\widetilde{A}_{12B}(-\boldsymbol{b}_{\rm T}^2,0,0,-1/(m_N\hat{\zeta})^2,\eta v{\cdot}P)}{\widetilde{A}_{2B}(-\boldsymbol{b}_{\rm T}^2,0,0,-1/(m_N\hat{\zeta})^2,\eta v{\cdot}P)} \ . \tag{2.49}$$

The soft factor and the multiplicative renormalization factors cancel in the ratio, so $\langle k_y\rangle_{TU}$ can be computed on the lattice without constructing either. In the limit $|\boldsymbol{b}_{\rm T}| \to 0$ it would become the average transverse momentum of the quarks, with the numerator summing quark and antiquark contributions and the denominator counting quarks minus antiquarks [75]. Because $b{\cdot}P = 0$, Eq. (2.49) is integrated over x.

The x dependence enters through the second argument of the amplitudes. Since x and $b{\cdot}P$ are Fourier conjugate variables (Chapter 4), keeping $b{\cdot}P$ as a variable and Fourier transforming the numerator and denominator separately gives an x-dependent generalized Sivers shift,

$$\langle k_y\rangle_{TU}(x,\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\eta v{\cdot}P) = -m_N\, \frac{\int d(b{\cdot}P)\ e^{ix(b\cdot P)}\ \widetilde{A}_{12B}(b^2,b{\cdot}P,\ldots)}{\int d(b{\cdot}P)\ e^{ix(b\cdot P)}\ \widetilde{A}_{2B}(b^2,b{\cdot}P,\ldots)} \ . \tag{2.50}$$

The soft factor depends only on $\boldsymbol{b}_{\rm T}$, not on $b{\cdot}P$, so it still cancels [75]. This is the quantity computed in this thesis. The full expression, including the kinematic constraint that relates $v{\cdot}b$ to $b{\cdot}P$, is Eq. (5.6) in Chapter 5.

2.11 From $b{\cdot}P$ to x

Because the x dependence is obtained through a Fourier transform, it is useful to state what the $b{\cdot}P$ dependence of an amplitude tells us about the corresponding distribution. We take the unpolarized case as the example. At fixed $b^2 = -\boldsymbol{b}_{\rm T}^2$, the relation $f_1 = 2\int_{\mathcal{F}} \widetilde{A}_{2B}$ of Eq. (4.36) can be inverted in the longitudinal variable. Since $\tilde{f}_1(x,\boldsymbol{b}_{\rm T}^2)$ vanishes outside $-1 \le x \le 1$,

$$\widetilde{A}_{2B}(-\boldsymbol{b}_{\rm T}^2,b{\cdot}P) = \frac{\tilde{\mathcal{S}}}{2}\int_{-1}^{1} dx\ e^{-ix(b\cdot P)}\ \tilde{f}_1(x,\boldsymbol{b}_{\rm T}^2) \ , \tag{2.51}$$

where we have suppressed the remaining arguments of the amplitude and of the soft factor. Three properties follow directly from this representation.

First, the Taylor coefficients of the amplitude in $b{\cdot}P$ are the Mellin moments of the distribution at fixed $\boldsymbol{b}_{\mathrm{T}}$,

$$\frac{\widetilde{A}_{2B}(-\boldsymbol{b}_{\mathrm{T}}^2, b{\cdot}P)}{\widetilde{A}_{2B}(-\boldsymbol{b}_{\mathrm{T}}^2, 0)} = \sum_{n=0}^{\infty} \frac{(-i\, b{\cdot}P)^n}{n!}\, \langle x^n \rangle_{\boldsymbol{b}_{\mathrm{T}}} \,, \qquad \langle x^n \rangle_{\boldsymbol{b}_{\mathrm{T}}} = \frac{\int dx\, x^n\, \tilde{f}_1(x, \boldsymbol{b}_{\mathrm{T}}^2)}{\int dx\, \tilde{f}_1(x, \boldsymbol{b}_{\mathrm{T}}^2)} \,. \tag{2.52}$$

The value at $b{\cdot}P = 0$ is the x-integrated quantity studied in earlier work, and the first few derivatives give low moments of the kind that local operators provide. Musch used this relation with straight links to estimate the lowest moments of the unpolarized distribution [106].

Second, the real and imaginary parts of the amplitude separate quark and antiquark contributions. Using $f_1^{\bar{q}}(x) = -f_1^{q}(-x)$ for $x > 0$, and taking the soft factor to be real,

$$\begin{aligned} \mathrm{Re}\, \widetilde{A}_{2B} &= \frac{\tilde{\mathcal{S}}}{2} \int_0^1 dx\ \cos\big(x\, b{\cdot}P\big) \Big[\tilde{f}_1^{q}(x) - \tilde{f}_1^{\bar{q}}(x)\Big] \,, \\ \mathrm{Im}\, \widetilde{A}_{2B} &= -\frac{\tilde{\mathcal{S}}}{2} \int_0^1 dx\ \sin\big(x\, b{\cdot}P\big) \Big[\tilde{f}_1^{q}(x) + \tilde{f}_1^{\bar{q}}(x)\Big] \,. \end{aligned} \tag{2.53}$$

The real part, which is even in $b{\cdot}P$, is determined by the quark-minus-antiquark (valence-like) combination, and the imaginary part, which is odd, by the quark-plus-antiquark combination. This is the parity structure used in Chapter 5. For the naively T-odd Sivers function the antiquark enters with the opposite relative sign [75], so the roles of the two combinations are interchanged for $\widetilde{A}_{12B}$.

Third, the support property constrains how the amplitude can behave at large $|b{\cdot}P|$. If $\tilde{f}_1$ is integrable, the Riemann-Lebesgue lemma implies $\widetilde{A}_{2B} \to 0$ as $|b{\cdot}P| \to \infty$, and since the support is bounded, $\widetilde{A}_{2B}$ extends to an entire function of $b{\cdot}P$ of exponential type no larger than one. A parametrization of lattice data that grows with $|b{\cdot}P|$ at fixed b^2 is therefore inconsistent with a physical distribution and has no Fourier transform in the ordinary sense. This criterion is used in Section 6.2 to reject the Gaussian ansatz. A parametrization that decays like a power of $b{\cdot}P$ fast enough to be absolutely integrable does have a Fourier transform. It is in general not an entire function, so the resulting distribution is not guaranteed to vanish outside $|x| \leq 1$, and the size of any such tail is one measure of the model dependence.

Lattice data cover only a finite range of $b{\cdot}P$, because $|b{\cdot}P| \leq |\boldsymbol{P}|\,|\boldsymbol{b}|$ in a frame where b has no time component (Section 2.12). The behavior of $\tilde{f}$ at small $|x|$ is governed by the amplitude at large $|b{\cdot}P|$, outside

the range of the data. The x dependence reconstructed from a finite range of $b{\cdot}P$ therefore depends on the functional form used to describe the data, and it depends on it most strongly at small x. This is why the choice and validation of fit models take up a large part of this thesis (Chapter 6).

2.12 TMDs from lattice QCD

Lattice QCD works in Euclidean space-time. A non-local operator can be evaluated directly only if all of its points lie at a single Euclidean time, which corresponds to purely space-like separations in Minkowski space. Operators that extend along a light-like direction, like those in Eq. (2.26) and Eq. (2.37), cannot be placed on the lattice as they stand. Three strategies have been developed to deal with this.

The first is the local operator approach of Section 2.5. Mellin moments are matrix elements of local operators and are computed directly, but only the lowest moments are accessible. For TMDs this approach does not apply at all, because the staple-shaped link cannot be expanded into local operators.

The second is the Lorentz-invariant amplitude approach of Musch, Hägler, Engelhardt, Negele and Schäfer [82, 83, 1], used in this thesis. The correlator with a finite, space-like staple is decomposed into amplitudes $\widetilde{A}_i$ and $\widetilde{B}_i$ that depend only on Lorentz invariants: b^2, $b{\cdot}P$, $\hat{\zeta}$ and the staple length through $\eta v{\cdot}P$ [102]. Because b and v are space-like, there is a frame in which both have vanishing time components, and in that frame the operator can be evaluated at a single Euclidean time. The amplitudes computed there are, by Lorentz invariance, the same amplitudes that define TMDs in the light-cone frame. The accessible kinematics are limited by the lattice. In a frame with $b^0 = v^0 = 0$, the invariants become $b{\cdot}P = -\boldsymbol{b}{\cdot}\boldsymbol{P}$ and $\hat{\zeta} = -\boldsymbol{v}{\cdot}\boldsymbol{P}/(|\boldsymbol{v}|\, m_N)$, so that

$$|b{\cdot}P| \leq |\boldsymbol{P}|\,|\boldsymbol{b}| \;, \qquad |\hat{\zeta}| \leq \frac{|\boldsymbol{P}|}{m_N} \;. \tag{2.54}$$

Nonzero $b{\cdot}P$ requires a component of $\boldsymbol{b}$ along the nucleon momentum, and large $\hat{\zeta}$ requires large nucleon momenta, where the statistical signal deteriorates. The light-cone limit $\hat{\zeta} \to \infty$ and the SIDIS and DY limits $\eta v{\cdot}P \to \pm\infty$ are reached by extrapolation. Soft factors and renormalization constants cancel in the ratios discussed in Section 2.10. With this approach the Sivers and Boer-Mulders shifts, the worm-gear shift and the generalized tensor charge have been computed for the nucleon and the pion [1, 84, 85, 86].

The third strategy is large-momentum effective theory (LaMET) [87, 89]. Instead of Lorentz invariance,

it uses the fact that an equal-time correlator in a hadron with large momentum P^z approaches the light-cone correlator, up to a perturbative matching and corrections suppressed by powers of $1/P^z$. For TMDs, the quasi-TMD is defined with a staple along the z direction and a Fourier transform in the longitudinal separation, and it is related to the light-cone TMD by a matching coefficient, a factor involving the Collins-Soper kernel, and a reduced soft function that can itself be computed on the lattice [88, 90, 110, 111]. Ratios of quasi-TMDs at different momenta give the Collins-Soper kernel [90, 91, 93]. A related approach uses pseudo-distributions, which are functions of the Lorentz invariants $b{\cdot}P$ and b^2 [112]. For infinitely long staples, the Lorentz-invariant and LaMET definitions of the beam function coincide, and both can be matched to the continuum TMDs [111]. Reviews of these methods can be found in Refs. [80, 81, 75].

The Lorentz-invariant approach is well suited to ratio observables such as the Sivers shift, because it works directly with the amplitudes in which soft factors cancel and it makes the dependence on $b{\cdot}P$ explicit. That dependence is what this thesis uses to reach the x dependence. The next chapter describes how the correlation functions are computed on the lattice.

3 LATTICE QCD METHODOLOGY

This chapter describes how the QCD correlation functions used in this thesis are defined and computed on a Euclidean spacetime lattice. We follow Lin and Meyer [2] & Gattringer and Lang [78]. Section 3.1 defines the lattice and the Euclidean Dirac matrices used throughout. Section 3.2 introduces the link variables, the plaquette and the Wilson gauge action, and derives its continuum limit. Section 3.3 discretizes the Dirac action, identifies the fermion doublers and removes them with the Wilson term, and adds the clover term. Section 3.4 integrates out the quark fields, introduces quark propagators and the Monte Carlo evaluation of the path integral. Section 3.5 derives the spectral form of the nucleon two-point function, and Section 3.6 does the same for three-point functions of local operators and shows how the ratio of three- to two-point functions isolates the matrix element. Section 3.7 extends the construction to the non-local staple-link operators that define TMDs.

3.1 The Euclidean Lattice

The Euclidean path integral of Section 2.2 is regulated by restricting spacetime to a four-dimensional hypercubic lattice [77, 75],

$$\Lambda_4 = \{n = (n_1, n_2, n_3, n_4) \,|\, n_i \in \{0, 1, \ldots, L_i - 1\}\} \ , \tag{3.1}$$

where L_i is the number of sites in direction i and a site n sits at the spacetime point $x = an$, with a the lattice spacing. We write $n + \hat{\mu}$ for the neighboring site in direction μ, which is the point $x + a\hat{\mu}$. Spatial boundary conditions are periodic for all fields; in time they are periodic for the gauge field and anti-periodic for the quark fields [75]. The lattice spacing acts as an ultraviolet regulator, with momenta restricted to the Brillouin zone $p_\mu \in (-\pi/a, \pi/a]$, and the finite volume acts as an infrared regulator. Physical results are recovered in the continuum limit $a \to 0$ and the infinite-volume limit [75].

3.1.1 Euclidean Dirac matrices.

The Euclidean Dirac matrices used in this thesis are related to the Minkowski ones by

$$\gamma_E^1 = +i\gamma_M^1 \ , \quad \gamma_E^2 = -i\gamma_M^2 \ , \quad \gamma_E^3 = +i\gamma_M^3 \ , \quad \gamma_E^4 = \gamma_M^0 \ , \tag{3.2}$$

This is the convention of the lattice code that produced the matrix elements analyzed in Section 5.6. These matrices are hermitian and satisfy $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$, and $\gamma_5 \equiv \gamma_1\gamma_2\gamma_3\gamma_4 = -\gamma_M^5$ is hermitian and anticommutes with every γ_μ. Products of Dirac matrices are labeled by a four-bit integer, $\Gamma(n) = \gamma_1^{n_1}\gamma_2^{n_2}\gamma_3^{n_3}\gamma_4^{n_4}$ with $n = n_1 + 2n_2 + 4n_3 + 8n_4$, so that $\Gamma(1) = \gamma_1$, $\Gamma(8) = \gamma_4$ and $\Gamma(15) = \gamma_5$. For γ_1 and γ_3, Eq. (3.2) agrees with the convention $\gamma_i^M = -i\gamma_i^E$ of Ref. [2]; for γ_2 the sign is opposite. For a Dirac spinor $u(p,s)$ of mass m the spin sum reads

$$\sum_s u(p,s)\,\bar{u}(p,s) = -i\,\not{p} + m\ , \qquad \not{p} \equiv \gamma_\mu p_\mu\ , \qquad p_\mu = (-p^1,\, p^2,\, -p^3,\, iE)\ , \tag{3.3}$$

where (E, p^1, p^2, p^3) are the Minkowski components. Equation (3.3) follows from the Minkowski relation $\sum_s u\bar{u} = \gamma_M^\mu p_\mu + m$ and Eq. (3.2), with $\bar{u}u = 2m$. It has the form of Eq. (3.78) of Ref. [2], with the components of p_μ fixed by Eq. (3.2). For momenta along the 1-axis, $p_\mu = (-p^1, 0, 0, iE)$.

3.2 Gauge Fields on the Lattice

On the lattice the gauge field is represented by link variables, $SU(3)$ matrices that live on the links between neighboring sites. In this chapter we use the lattice convention of Ref. [78], in which the coupling is absorbed into the gauge field,

$$U_\mu(n) = \exp\big(iaA_\mu(n)\big)\ , \tag{3.4}$$

with $A_\mu(n)$ evaluated at the starting point of the link, which is accurate to $\mathcal{O}(a)$ [78]. This convention is related to that of Eqs. (2.10)–(2.14) by $A_\mu \to -gA_\mu$ and does not affect gauge-invariant quantities. The link $U_\mu(x)$ is the lattice version of the Wilson line between x and $x + a\hat{\mu}$ [2]. Under a gauge transformation $\Lambda(x) \in SU(3)$, the quark fields and links transform as [2]

$$\psi(x) \to \Lambda(x)\psi(x)\ , \qquad \bar{\psi}(x) \to \bar{\psi}(x)\Lambda(x)^{-1}\ , \qquad U_\mu(x) \to \Lambda(x)\,U_\mu(x)\,\Lambda(x+a\hat{\mu})^{-1}\ , \tag{3.5}$$

so that $U_\mu(x)\psi(x+a\hat{\mu})$ transforms like $\psi(x)$. A product of links along a path transforms with Λ at its two end points only. The trace of the product around a closed loop is therefore gauge invariant [78], and so is a product of links along an open path with a quark field attached at each end, which is how the non-local operators of Section 3.7 are built.

The shortest non-trivial closed loop is the plaquette (Fig. 5),

$$P_{\mu\nu}(x) = U_\mu(x)\, U_\nu(x + a\hat{\mu})\, U_\mu(x + a\hat{\nu})^\dagger\, U_\nu(x)^\dagger \ . \tag{3.6}$$



Figure 5: Quark fields $\psi(x)$ and $\psi(x + a\hat{\mu})$ on the sites of a cubic spacetime lattice, the link variable $U_\mu(x)$ between them, and the plaquette $P_{\mu\nu}(y)$ of Eq. (3.6). The product $U_\mu(x)\psi(x + a\hat{\mu})$ transforms in the same way as $\psi(x)$. Reproduced from Ref. [2], Fig. 1.1.

Wilson's gauge action is a sum over all plaquettes, each counted with one orientation [77, 2],

$$S_G[U] = \frac{2}{g^2} \sum_{n \in \Lambda_4} \sum_{\mu<\nu} \mathrm{Re}\,\mathrm{tr}\big[\mathbb{1} - P_{\mu\nu}(n)\big] \ . \tag{3.7}$$

Written as $\beta \sum [1 - \frac{1}{3}\mathrm{Re}\,\mathrm{tr} P_{\mu\nu}]$, the prefactor is the inverse coupling $\beta = 6/g^2$ [78].

3.2.1 Continuum limit of the Wilson action.

Equation (2.17) quoted the small-a behavior of the plaquette; here we derive it. To show that Eq. (3.7) approaches the Yang-Mills action, we insert Eq. (3.4) into the plaquette and combine the four exponentials with the Baker-Campbell-Hausdorff formula [78],

$$e^A e^B = \exp\Big(A + B + \tfrac{1}{2}[A, B] + \cdots\Big) \ . \tag{3.8}$$

Applying Eq. (3.8) three times gives

$$P_{\mu\nu}(n) = \exp\Big(iaA_\mu(n) + iaA_\nu(n+\hat{\mu}) - iaA_\mu(n+\hat{\nu}) - iaA_\nu(n) \\ - \tfrac{a^2}{2}[A_\mu(n), A_\nu(n+\hat{\mu})] - \tfrac{a^2}{2}[A_\mu(n+\hat{\nu}), A_\nu(n)] + \tfrac{a^2}{2}[A_\nu(n+\hat{\mu}), A_\mu(n+\hat{\nu})] \\ + \tfrac{a^2}{2}[A_\mu(n), A_\nu(n)] + \tfrac{a^2}{2}[A_\mu(n), A_\mu(n+\hat{\nu})] + \tfrac{a^2}{2}[A_\nu(n+\hat{\mu}), A_\nu(n)] + \mathcal{O}(a^3)\Big) \,. \tag{3.9}$$

The fields at shifted points are Taylor expanded, $A_\nu(n+\hat{\mu}) = A_\nu(n) + a\,\partial_\mu A_\nu(n) + \mathcal{O}(a^2)$. The linear terms combine into $ia^2(\partial_\mu A_\nu - \partial_\nu A_\mu)$, the last two commutators vanish at this order, and the remaining four combine into $-a^2[A_\mu, A_\nu]$, so that [78]

$$P_{\mu\nu}(n) = \exp\Big(ia^2\big(\partial_\mu A_\nu(n) - \partial_\nu A_\mu(n) + i[A_\mu(n), A_\nu(n)]\big) + \mathcal{O}(a^3)\Big) = \exp\big(ia^2 F_{\mu\nu}(n) + \mathcal{O}(a^3)\big) \,. \tag{3.10}$$

Expanding the exponential, $\mathbb{1} - P_{\mu\nu} = -ia^2 F_{\mu\nu} + \frac{a^4}{2}F_{\mu\nu}^2 + \ldots$. The $\mathcal{O}(a^2)$ term $-ia^2F_{\mu\nu}$ drops out of $\mathrm{Re\,tr}$, because $F_{\mu\nu}$ is traceless (equivalently, $\mathrm{tr}[P_{\mu\nu}]^* = \mathrm{tr}[P_{\nu\mu}]$), and the $\mathcal{O}(a^3)$ terms cancel in a similar way [78]. Using $\sum_{\mu<\nu} = \frac{1}{2}\sum_{\mu,\nu}$ for the symmetric summand,

$$S_G[U] = \frac{a^4}{2g^2}\sum_{n\in\Lambda_4}\sum_{\mu,\nu}\mathrm{tr}\big[F_{\mu\nu}(n)^2\big] + \mathcal{O}(a^2) \;\xrightarrow{a\to 0}\; \frac{1}{2g^2}\int d^4x \sum_{\mu,\nu}\mathrm{tr}\big[F_{\mu\nu}(x)^2\big] \,, \tag{3.11}$$

since $a^4\sum_n$ becomes $\int d^4x$. This is the Euclidean Yang-Mills action [2], and the corrections of Eq. (3.7) start at $\mathcal{O}(a^2)$ [78].

The integration over gauge fields uses the product of Haar measures of all links, $\mathcal{D}[U] = \prod_{x,\mu} dU_\mu(x)$, which is invariant under the gauge transformation of Eq. (3.5) [2].

3.3 Fermions on the Lattice

3.3.1 Naive discretization and doublers.

Replacing the derivative in the Euclidean Dirac action by a symmetric, gauge-covariant difference gives the naive lattice action [78]

$$S_F[\psi,\bar{\psi},U] = a^4\sum_{n\in\Lambda_4}\bar{\psi}(n)\left(\sum_{\mu=1}^{4}\gamma_\mu\frac{U_\mu(n)\psi(n+\hat{\mu}) - U_{-\mu}(n)\psi(n-\hat{\mu})}{2a} + m\,\psi(n)\right), \tag{3.12}$$

with $U_{-\mu}(n) \equiv U_\mu(n-\hat{\mu})^\dagger$. The action is bilinear, $S_F = a^4\sum_{n,m}\bar{\psi}(n)D(n|m)\psi(m)$, with the naive Dirac operator

$$D(n|m)^{\alpha\beta}_{ab} = \sum_{\mu=1}^{4}(\gamma_\mu)_{\alpha\beta}\,\frac{U_\mu(n)_{ab}\,\delta_{n+\hat{\mu},m} - U_{-\mu}(n)_{ab}\,\delta_{n-\hat{\mu},m}}{2a} + m\,\delta_{\alpha\beta}\,\delta_{ab}\,\delta_{n,m} \,. \tag{3.13}$$

For the free field, $U_\mu(n) = \mathbb{1}$, the Fourier transform of Eq. (3.13) is diagonal in momentum, with [78]

$$\widetilde{D}(p) = m\,\mathbb{1} + \frac{i}{a}\sum_{\mu=1}^{4}\gamma_\mu \sin(p_\mu a)\ , \qquad \widetilde{D}(p)^{-1} = \frac{m\,\mathbb{1} - ia^{-1}\sum_\mu \gamma_\mu \sin(p_\mu a)}{m^2 + a^{-2}\sum_\mu \sin^2(p_\mu a)}\ . \tag{3.14}$$

For $m = 0$ the propagator has the correct continuum limit at fixed p, $\widetilde{D}(p)^{-1} \to -i\sum_\mu \gamma_\mu p_\mu/p^2$, with a pole at $p = (0,0,0,0)$. On the lattice, however, $\sin(p_\mu a)$ also vanishes at $p_\mu = \pi/a$. Every momentum with components equal to 0 or π/a is a pole [78],

$$p = (\pi/a,0,0,0)\,,\ (0,\pi/a,0,0)\,,\ \ldots\,,\ (\pi/a,\pi/a,\pi/a,\pi/a)\ , \tag{3.15}$$

which gives 15 unphysical poles, the doublers, in addition to the physical one. By the Nielsen-Ninomiya theorem [113, 2], a lattice Dirac operator with the correct continuum limit, no doublers and exact chiral symmetry in the form $\gamma_5 D + D\gamma_5 = 0$ cannot be constructed.

3.3.2 Wilson fermions.

Wilson removed the doublers by adding a term that vanishes for $p_\mu = 0$ [77, 78],

$$\widetilde{D}(p) = m\,\mathbb{1} + \frac{i}{a}\sum_{\mu=1}^{4}\gamma_\mu \sin(p_\mu a) + \mathbb{1}\,\frac{1}{a}\sum_{\mu=1}^{4}\big(1-\cos(p_\mu a)\big)\ . \tag{3.16}$$

Each component $p_\mu = \pi/a$ contributes $2/a$, so a doubler with ℓ such components has mass $m + 2\ell/a$. In the limit $a \to 0$ the doublers become infinitely heavy and decouple. In position space the Wilson term is a discretization of $-\frac{a}{2}\partial_\mu\partial_\mu$, and combining it with Eq. (3.13) gives the Wilson-Dirac operator [78]

$$D(n|m)^{\alpha\beta}_{ab} = \Big(m + \frac{4}{a}\Big)\delta_{\alpha\beta}\,\delta_{ab}\,\delta_{n,m} - \frac{1}{2a}\sum_{\mu=\pm1}^{\pm4}(\mathbb{1}-\gamma_\mu)_{\alpha\beta}\,U_\mu(n)_{ab}\,\delta_{n+\hat{\mu},m}\ , \qquad \gamma_{-\mu} = -\gamma_\mu\ . \tag{3.17}$$

Rescaling the quark fields by $\sqrt{m+4/a}$ brings it to the hopping form [78]

$$D = \mathbb{1} - \kappa H\ , \qquad \kappa = \frac{1}{2(am+4)}\ , \qquad H(n|m)^{\alpha\beta}_{ab} = \sum_{\mu=\pm1}^{\pm4}(\mathbb{1}-\gamma_\mu)_{\alpha\beta}\,U_\mu(n)_{ab}\,\delta_{n+\hat{\mu},m}\ , \tag{3.18}$$

where H collects the nearest-neighbor terms and κ is the hopping parameter. The Wilson-Dirac operator is γ_5-hermitian [2, 78],

$$D^\dagger = \gamma_5\, D\, \gamma_5\ , \tag{3.19}$$

which follows from $\gamma_5^\dagger = \gamma_5$, $\gamma_5\gamma_\mu\gamma_5 = -\gamma_\mu$ and $U_{-\mu}(n) = U_\mu(n-\hat{\mu})^\dagger$, and which holds in the basis of Eq. (3.2). The Wilson term breaks chiral symmetry explicitly, even for massless quarks, and introduces discretization errors of $\mathcal{O}(a)$ [78].

3.3.3 Clover improvement.

The $\mathcal{O}(a)$ errors of the Wilson action are removed by adding the Sheikholeslami-Wohlert or clover term [114, 78],

$$S_{\text{clover}} = S_{\text{Wilson}} + c_{\text{sw}}\, a^5 \sum_{n\in\Lambda_4} \sum_{\mu<\nu} \bar{\psi}(n)\, \frac{1}{2}\, \sigma_{\mu\nu}\, \widehat{F}_{\mu\nu}(n)\, \psi(n) \ , \qquad \sigma_{\mu\nu} = \frac{1}{2i}[\gamma_\mu, \gamma_\nu] \ , \tag{3.20}$$

where the lattice field strength is built from the four plaquettes around the site n,

$$\widehat{F}_{\mu\nu}(n) = \frac{-i}{8a^2}\big(Q_{\mu\nu}(n) - Q_{\nu\mu}(n)\big) \ , \qquad Q_{\mu\nu}(n) = U_{\mu,\nu}(n) + U_{\nu,-\mu}(n) + U_{-\mu,-\nu}(n) + U_{-\nu,\mu}(n) \ , \tag{3.21}$$

with $U_{\mu,\nu}(n)$ the plaquette of Eq. (3.6) in the (μ,ν) orientation. The four plaquettes form the clover leaf that gives the term its name, and the coefficient c_{sw} is tuned to cancel the $\mathcal{O}(a)$ terms [2, 78].

3.3.4 Other formulations.

Twisted-mass fermions remove the doublers with a Wilson term, while staggered (Kogut-Susskind) fermions reduce the sixteen naive species to four tastes and keep a remnant chiral symmetry. Overlap fermions satisfy the Ginsparg-Wilson relation, and domain-wall fermions, which introduce a fifth dimension, satisfy it approximately at finite fifth-dimension extent [75, 78]. The gauge ensemble used in this work was generated with $n_f = 2+1$ flavors of clover-improved Wilson fermions [86].

3.4 Path Integral, Wick Contractions and Quark Propagators

With gauge action S_G and fermion action $S_F = a^4 \sum_f \sum_{n,m} \bar{\psi}^{(f)}(n) D^{(f)}(n|m) \psi^{(f)}(m)$, the lattice expectation value of an observable is [78, 2]

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}[U]\, \mathcal{D}[\psi,\bar{\psi}]\, \mathcal{O}[\psi,\bar{\psi},U]\, e^{-S_F[\psi,\bar{\psi},U] - S_G[U]} \ , \qquad Z = \int \mathcal{D}[U]\, \mathcal{D}[\psi,\bar{\psi}]\, e^{-S_F - S_G} \ . \tag{3.22}$$

The quark fields are Grassmann variables (Eq. (2.8)). For a quadratic form, the Grassmann integral with sources is [2]

$$\int \prod_i d\eta_i\, d\bar{\eta}_i\, \exp\Big(- \sum_{ij} \bar{\eta}_i A_{ij} \eta_j + \sum_i (\bar{\eta}_i \zeta_i + \bar{\zeta}_i \eta_i) \Big) = c\, \det(A)\, \exp\Big(\sum_{ij} \bar{\zeta}_i (A^{-1})_{ij} \zeta_j \Big) \ , \tag{3.23}$$

so the quark integration produces the fermion determinant, and derivatives with respect to the sources turn every pair $\psi\bar{\psi}$ into an element of A^{-1}. With $A = a^4 D$ this is Wick's theorem: the quark fields in $\mathcal{O}$ are

contracted in all possible ways and replaced by quark propagators [78, 75],

$$\left\langle \psi(n)_{\alpha a}\, \bar{\psi}(m)_{\beta b} \right\rangle_F = a^{-4} D^{-1}(n|m)_{\substack{\alpha\beta \\ ab}} \equiv S(n,m)_{\substack{\alpha\beta \\ ab}} \ . \tag{3.24}$$

After integrating out the quarks,

$$\langle \mathcal{O} \rangle = \frac{1}{Z} \int \mathcal{D}[U]\, \widetilde{\mathcal{O}}[U] \prod_f \det\left[D^{(f)}[U]\right] e^{-S_G[U]} \ , \tag{3.25}$$

where $\widetilde{\mathcal{O}}[U]$ is the contracted observable, a function of the propagators in the gauge field U [75].

3.4.1 Monte Carlo evaluation.

For real and positive determinants, $\mathcal{P}[U] = Z^{-1} \prod_f \det[D^{(f)}[U]]\, e^{-S_G[U]}$ is a probability distribution. A Markov chain whose transition probabilities satisfy detailed balance produces gauge configurations U_i distributed according to $\mathcal{P}[U]$ [78], and the expectation value becomes an ensemble average [2, 75],

$$\langle \mathcal{O} \rangle = \frac{1}{N} \sum_{i=1}^{N} \widetilde{\mathcal{O}}[U_i] + \mathcal{O}\big(N^{-1/2}\big) \ . \tag{3.26}$$

With dynamical quarks the standard algorithm is hybrid Monte Carlo [115]. Statistical uncertainties in this thesis are estimated with the Jackknife method (Section 6.6).

3.4.2 Computing propagators.

Rather than computing the full matrix D^{-1}, one solves the linear system [78]

$$\sum_m D(n|m)\, G(m) = S_0(n) \ , \qquad S_0(m)_{\substack{\beta \\ b}} = \delta_{m,m_0}\, \delta_{\beta\beta_0}\, \delta_{bb_0} \ , \tag{3.27}$$

for a point source S_0 at m_0 with iterative Krylov-space solvers. The twelve solutions for $\beta_0 = 1, \ldots, 4$ and $b_0 = 1, 2, 3$ give the point-to-all propagator $D^{-1}(n|m_0)$ from m_0 to every site n. Propagators in the opposite direction follow from Eq. (3.19) without further inversions [78],

$$S(m_0, n) = \gamma_5\, S(n, m_0)^\dagger\, \gamma_5 \ . \tag{3.28}$$

3.4.3 Example: the pion two-point function.

As an example of Wick contraction [78, 2], consider $\chi_\pi = \bar{d}\,\gamma_5\, u$, the conjugate operator is $\bar{\chi}_\pi = (\bar{d}\gamma_5 u)^\dagger = -\bar{u}\gamma_5 d$, using $\bar{\psi} = \psi^\dagger \gamma_4$ and $\gamma_4\gamma_5\gamma_4 = -\gamma_5$. Wick's theorem contracts $u(x)\bar{u}(0)$ and $d(0)\bar{d}(x)$; moving $d(0)$ to

the front past three Grassmann fields gives a sign, and the minus sign of $\bar{\chi}_\pi$ cancels it:

$$\big\langle \chi_\pi(x)\,\bar{\chi}_\pi(0)\big\rangle_F = \mathrm{tr}\big[S_d(0,x)\,\gamma_5\,S_u(x,0)\,\gamma_5\big] = \mathrm{tr}\big[S_d(x,0)^\dagger\,S_u(x,0)\big]\ , \tag{3.29}$$

where the second step uses Eq. (3.28) and the trace runs over Dirac and color indices. For degenerate u and d quarks this is $\sum |S_u(x,0)|^2$, so a single point-to-all propagator per configuration determines the pion correlator. The nucleon correlators below are built in the same way.

3.5 The Nucleon Two-Point Function

3.5.1 Interpolating operator and contraction.

A standard interpolating operator with the quantum numbers of the proton is [75, 2]

$$\chi_\alpha(\boldsymbol{x},t) = \epsilon^{ijk}\,u^i_\alpha(\boldsymbol{x},t)\,u^j_\gamma(\boldsymbol{x},t)\,[C^{-1}\gamma_5]_{\gamma\delta}\,d^k_\delta(\boldsymbol{x},t)\ , \tag{3.30}$$

with C the charge-conjugation matrix. The momentum-projected two-point function is

$$C_{\alpha\beta}(t,\boldsymbol{p}) = a^3\sum_{\boldsymbol{x}} e^{-i\boldsymbol{p}\cdot\boldsymbol{x}}\,\big\langle \chi_\alpha(\boldsymbol{x},t)\,\bar{\chi}_\beta(\boldsymbol{0},0)\big\rangle\ , \tag{3.31}$$

and Wick's theorem expresses it through u and d propagators from the source [75],

$$C_{\alpha\beta}(t,\boldsymbol{p}) = -a^3\sum_{\boldsymbol{x}} e^{-i\boldsymbol{p}\cdot\boldsymbol{x}}\,\epsilon^{ijk}\epsilon^{i'j'k'}\,[C^{-1}\gamma_5]_{\alpha'\alpha''}\,[\gamma_5 C]_{\beta'\beta''}\,\Big\langle \big[S_d\big]^{ki'}_{\alpha''\beta'}\Big\{\big[S_u\big]^{jj'}_{\alpha'\beta''}\big[S_u\big]^{ik'}_{\alpha\beta} - \big[S_u\big]^{ij'}_{\alpha\beta''}\big[S_u\big]^{jk'}_{\alpha'\beta}\Big\}\Big\rangle\ , \tag{3.32}$$

with $S_f \equiv S_f(\boldsymbol{x},t;\boldsymbol{0},0)$. The two terms correspond to the two ways of contracting the two u quarks.

3.5.2 Spectral decomposition.

On a lattice with time extent T, the two-point function is a trace over states [2],

$$\big\langle \chi(\boldsymbol{x},t)\,\bar{\chi}(\boldsymbol{0},0)\big\rangle = \frac{\mathrm{Tr}\big[e^{-H(T-t)}\chi(\boldsymbol{x})\,e^{-Ht}\,\bar{\chi}(\boldsymbol{0})\big]}{\mathrm{Tr}\big[e^{-HT}\big]}\ . \tag{3.33}$$

For $T\to\infty$ only the vacuum survives in the outer trace, and inserting a complete set of states between the operators gives the spectral sum of Eq. (2.7). We use relativistically normalized states, $\langle \boldsymbol{p}'|\boldsymbol{p}\rangle = (2\pi)^3 2E_{\boldsymbol{p}}\,\delta^{(3)}(\boldsymbol{p}-\boldsymbol{p}')$, for which the momentum projection in Eq. (3.31) selects states of momentum $\boldsymbol{p}$ weighted by $1/(2E_n(\boldsymbol{p}))$ [2, 75]. For a nucleon state the overlap with the interpolating operator is proportional to a Dirac spinor,

$$\langle\Omega|\chi_\alpha(0)|N_n(\boldsymbol{p},s)\rangle = \sqrt{Z_n(\boldsymbol{p})}\,u_\alpha(n,\boldsymbol{p},s)\ ,\qquad \langle N_n(\boldsymbol{p},s)|\bar{\chi}_\beta(0)|\Omega\rangle = \sqrt{Z_n(\boldsymbol{p})}^{\,*}\,\bar{u}_\beta(n,\boldsymbol{p},s)\ , \tag{3.34}$$

where n labels the nucleon and its excitations. If source and sink operators differ, for example through smearing, Z_n stands for $\sqrt{Z_n^{\rm snk}\bar{Z}_n^{\rm src}}$ [2]. Then

$$C_{\alpha\beta}(t,\boldsymbol{p}) = \sum_n \frac{|Z_n(\boldsymbol{p})|}{2E_n(\boldsymbol{p})}\, e^{-E_n(\boldsymbol{p})t} \sum_s u_\alpha(n,\boldsymbol{p},s)\,\bar{u}_\beta(n,\boldsymbol{p},s) = \sum_n \frac{|Z_n(\boldsymbol{p})|}{2E_n(\boldsymbol{p})}\, e^{-E_n(\boldsymbol{p})t}\,\big(-i\not{p}_n + m_n\big)_{\alpha\beta}\ , \tag{3.35}$$

where we used Eq. (3.3), with p_n the Euclidean four-momentum of Eq. (3.3) for energy $E_n(\boldsymbol{p})$. The correlator is projected onto positive parity with $\Gamma^+ = \frac{1}{2}(1+\gamma_4)$ [75]. Using $\mathrm{tr}[\gamma_\mu] = 0$, $\mathrm{tr}[\gamma_4\gamma_4] = 4$ and $p_4 = iE$,

$$\mathrm{tr}\big[\Gamma^+\big(-i\not{p} + m\big)\big] = \frac{1}{2}\Big(-i\,\mathrm{tr}\big[\gamma_4\gamma_4\big]\,iE + 4m\Big) = 2(E+m)\ , \tag{3.36}$$

so that, for $0 \ll t \ll T/2$,

$$C_{\Gamma^+}(t,\boldsymbol{p}) \equiv \Gamma^+_{\beta\alpha} C_{\alpha\beta}(t,\boldsymbol{p}) = |Z(\boldsymbol{p})|\,\frac{E(\boldsymbol{p})+m}{E(\boldsymbol{p})}\, e^{-E(\boldsymbol{p})t}\,\big[1 + \mathcal{O}\big(e^{-\Delta E\, t}\big)\big]\ , \tag{3.37}$$

where E and m refer to the nucleon ground state and ΔE is the gap to the first excitation. This agrees with Eq. (3.79) of Ref. [2] and Eq. (3.46) of Ref. [106]. On a finite lattice, the backward-propagating contribution $\propto e^{-E(T-t)}$ and the negative-parity states are not negligible near $t \approx T/2$ [2]. Fits of Eq. (3.37) at large t determine the nucleon mass and dispersion relation.

3.6 Three-Point Functions of Local Operators

3.6.1 Definition and spectral decomposition.

To probe nucleon structure, an operator is inserted at time τ between the source at time 0 and the sink at time t. For a local quark bilinear,

$$\mathcal{O}_\Gamma(x) = \bar{q}(x)\,\Gamma\, q(x)\ , \tag{3.38}$$

where Γ is any product of Dirac matrices, the three-point function is [2]

$$G(t,\tau;\boldsymbol{p},\boldsymbol{p}') = a^6 \sum_{\boldsymbol{x}_2,\boldsymbol{x}_1} e^{-i\boldsymbol{p}'\cdot(\boldsymbol{x}_2-\boldsymbol{x}_1)}\, e^{-i\boldsymbol{p}\cdot\boldsymbol{x}_1}\, \Gamma^{\rm 3pt}_{\beta\alpha}\, \big\langle\Omega\big|\, T\big\{\chi_\alpha(\boldsymbol{x}_2,t)\,\mathcal{O}(\boldsymbol{x}_1,\tau)\,\bar{\chi}_\beta(\boldsymbol{0},0)\big\}\big|\Omega\big\rangle\ , \tag{3.39}$$

with spin projector $\Gamma^{\rm 3pt}$ and momentum transfer $\boldsymbol{q} = \boldsymbol{p}' - \boldsymbol{p}$; the factor a^6 cancels in the ratio below. Inserting complete sets of states before and after $\mathcal{O}$ and using the translation $\chi(\boldsymbol{x},t) = e^{Ht}e^{-i\boldsymbol{P}\cdot\boldsymbol{x}}\chi(0)e^{i\boldsymbol{P}\cdot\boldsymbol{x}}e^{-Ht}$ [2],

$$G(t,\tau;\boldsymbol{p},\boldsymbol{p}') = \sum_{n,n'}\sum_{s,s'} \frac{e^{-E_{n'}(\boldsymbol{p}')(t-\tau)}\, e^{-E_n(\boldsymbol{p})\tau}}{2E_{n'}(\boldsymbol{p}')\,2E_n(\boldsymbol{p})}\, \Gamma^{\rm 3pt}_{\beta\alpha}\, \langle\Omega|\chi_\alpha|N_{n'}(\boldsymbol{p}',s')\rangle\,\langle N_{n'}(\boldsymbol{p}',s')|\mathcal{O}(\boldsymbol{q})|N_n(\boldsymbol{p},s)\rangle\,\langle N_n(\boldsymbol{p},s)|\bar{\chi}_\beta|\Omega\rangle\ . \tag{3.40}$$

For $0 \ll \tau \ll t$ the ground state dominates both sums [2]. The matrix element is written as a spinor sandwich,

$$\langle N(\boldsymbol{p}', s')|\mathcal{O}(\boldsymbol{q})|N(\boldsymbol{p}, s)\rangle = \bar{u}(\boldsymbol{p}', s')\, \mathcal{J}\, u(\boldsymbol{p}, s) \ , \tag{3.41}$$

where $\mathcal{J}$ is the combination of Dirac matrices and form factors allowed by the symmetries of $\mathcal{O}$ [2]. Inserting Eqs. (3.34) and (3.41), the spin sums close into Eq. (3.3),

$$\begin{aligned} G(t, \tau; \boldsymbol{p}, \boldsymbol{p}') &= \frac{\sqrt{Z(\boldsymbol{p}')}\sqrt{Z(\boldsymbol{p})}^*}{4E(\boldsymbol{p}')E(\boldsymbol{p})}\, e^{-E(\boldsymbol{p}')(t-\tau)}\, e^{-E(\boldsymbol{p})\tau}\, \Gamma^{\mathrm{3pt}}_{\beta\alpha} \sum_{s'} u_\alpha(\boldsymbol{p}', s')\bar{u}(\boldsymbol{p}', s')\, \mathcal{J} \sum_{s} u(\boldsymbol{p}, s)\bar{u}_\beta(\boldsymbol{p}, s) \\ &= \frac{\sqrt{Z(\boldsymbol{p}')}\sqrt{Z(\boldsymbol{p})}^*}{4E(\boldsymbol{p}')E(\boldsymbol{p})}\, e^{-E(\boldsymbol{p}')(t-\tau)}\, e^{-E(\boldsymbol{p})\tau}\, \mathrm{tr}\Big[\Gamma^{\mathrm{3pt}}\big(-i\not{p}' + m\big)\, \mathcal{J}\, \big(-i\not{p} + m\big)\Big] \ , \end{aligned} \tag{3.42}$$

(compare Eqs. (3.80)–(3.81) of Ref. [2] and Eq. (3.57) of Ref. [106]).

3.6.2 Quark-level contraction and sequential propagators.

After Wick contraction, the three-point function contains a connected part, in which the operator acts on a valence quark line, and a disconnected part, in which it closes a separate quark loop [2]. In the isospin limit $m_u = m_d$, the disconnected part cancels in the isovector combination $u - d$ [75, 86]. In the connected part, the propagator from the operator at $(\boldsymbol{x}_1, \tau)$ to the sink at $(\boldsymbol{x}_2, t)$ is summed over $\boldsymbol{x}_2$ and cannot be computed from point-to-all propagators. The sequential-source method solves this problem [2]. For a fixed sink time t, sink momentum $\boldsymbol{p}'$ and projector Γ^{3pt}, the forward propagators from the source are combined into a sequential source at the sink, and one further inversion gives the sequential propagator $\Sigma(\boldsymbol{0}, 0; \boldsymbol{x}_1, \tau; \boldsymbol{p}', t)$ from the sink to every point $(\boldsymbol{x}_1, \tau)$. The three-point function then becomes, with the phase convention of Eq. (3.39) [2],

$$G_\Gamma(t, \tau; \boldsymbol{p}, \boldsymbol{p}') = \sum_{\boldsymbol{x}_1} e^{+i\boldsymbol{q}\cdot\boldsymbol{x}_1} \left\langle \mathrm{tr}\big[\Sigma(\boldsymbol{0}, 0; \boldsymbol{x}_1, \tau; \boldsymbol{p}', t)\, \Gamma\, S(\boldsymbol{x}_1, \tau; \boldsymbol{0}, 0)\big] \right\rangle \ , \tag{3.43}$$

a contraction of the sequential propagator, the operator and an ordinary propagator. Because the sum over $\boldsymbol{x}_1$ is done last, any operator and any momentum transfer can be used with the same sequential propagator. A new inversion is needed for each sink momentum, sink time, projector and quark flavor [2, 106].

3.6.3 Ratio and zero momentum transfer.

The overlaps $Z(\boldsymbol{p})$ and the exponential time dependence are removed with the ratio [2]

$$R(t, \tau; \boldsymbol{p}', \boldsymbol{p}) = \frac{G_\Gamma(t, \tau; \boldsymbol{p}, \boldsymbol{p}')}{C_{\Gamma^+}(t, \boldsymbol{p}')} \left[\frac{C_{\Gamma^+}(\tau, \boldsymbol{p}')\, C_{\Gamma^+}(t, \boldsymbol{p}')\, C_{\Gamma^+}(t-\tau, \boldsymbol{p})}{C_{\Gamma^+}(\tau, \boldsymbol{p})\, C_{\Gamma^+}(t, \boldsymbol{p})\, C_{\Gamma^+}(t-\tau, \boldsymbol{p}')}\right]^{1/2} . \tag{3.44}$$

For the forward matrix elements that define TMDs, the source and sink momenta are equal, $\boldsymbol{p}' = \boldsymbol{p}$, so $\boldsymbol{q} = 0$. Every two-point function in the numerator of the bracket then appears in the denominator with the same arguments,

$$\left[\frac{C_{\Gamma^+}(\tau,\boldsymbol{p})\,C_{\Gamma^+}(t,\boldsymbol{p})\,C_{\Gamma^+}(t-\tau,\boldsymbol{p})}{C_{\Gamma^+}(\tau,\boldsymbol{p})\,C_{\Gamma^+}(t,\boldsymbol{p})\,C_{\Gamma^+}(t-\tau,\boldsymbol{p})}\right]^{1/2} = 1 \ , \tag{3.45}$$

and the ratio reduces to $R = G_\Gamma(t,\tau;\boldsymbol{p},\boldsymbol{p})/C_{\Gamma^+}(t,\boldsymbol{p})$. Inserting Eqs. (3.37) and (3.42) with $\boldsymbol{p}' = \boldsymbol{p}$, the overlaps and the factors $e^{-E(t-\tau)}e^{-E\tau} = e^{-Et}$ cancel,

$$R(t,\tau;\boldsymbol{p},\boldsymbol{p}) \xrightarrow{0\ll\tau\ll t} \frac{\mathrm{tr}\big[\Gamma^{3\mathrm{pt}}\big(-i\not{p}+m\big)\,\mathcal{J}\,\big(-i\not{p}+m\big)\big]}{4E(\boldsymbol{p})\big(E(\boldsymbol{p})+m\big)} = \frac{\mathrm{tr}\big[\Gamma^{3\mathrm{pt}}\big(-i\not{p}+m\big)\,\mathcal{J}\,\big(-i\not{p}+m\big)\big]}{2E(\boldsymbol{p})\,\mathrm{tr}\big[\Gamma^{+}\big(-i\not{p}+m\big)\big]} \ , \tag{3.46}$$

where the second form uses Eq. (3.36) and agrees with Eq. (3.61) of Ref. [106]. Equivalently, the kinematic prefactor $\sqrt{E_{p'}E_p/[(E_{p'}+m)(E_p+m)]}$ of Eq. (3.93) in Ref. [2] becomes $E/(E+m)$ at $\boldsymbol{p}' = \boldsymbol{p}$. The ratio is independent of τ and t in the region where the ground state dominates. Excited states are suppressed by increasing the source-sink separation, but the statistical noise grows with it, so the separation is a compromise between the two [106].

3.7 Non-Local Staple-Link Operators

A local operator such as Eq. (3.38) places the quark and antiquark fields at the same point. The TMD correlator of Eq. (4.1) instead contains the non-local operator $\bar{q}(0)\,\Gamma\,\mathcal{U}[0,\eta v,\eta v+b,b]\,q(b)$, in which the quark fields are separated by b and connected by the staple-shaped gauge link of Eq. (4.11) to make the operator gauge invariant. As discussed in Sections 2.8 and 4.3, the staple legs of the continuum definition extend to infinity; on the lattice v is space-like, η is finite, and the results are extrapolated to $\eta \to \infty$ [75, 86].

3.7.1 Lattice operator and three-point function.

The operator is evaluated in a Lorentz frame in which b and v are purely spatial, so that the whole operator lies in one time slice and can be computed with the Euclidean path integral [116, 86]. On the lattice the gauge link is a product of link variables along a step-wise path $\mathcal{C}^{\mathrm{lat}}$ that approximates the staple [106],

$$O^{\mathrm{lat}}_{\Gamma,q}(\mathcal{C}^{\mathrm{lat}};z) \equiv \bar{q}(z)\,\Gamma\,\mathcal{U}^{\mathrm{lat}}\big[\mathcal{C}^{\mathrm{lat}}+z\big]\,q\big(z+\ell[\mathcal{C}^{\mathrm{lat}}]\big) \ , \tag{3.47}$$

where $\ell[\mathcal{C}^{\mathrm{lat}}]$ is the displacement between the two ends of the path. We place $\bar{q}$ at the start of the path and q at its end, as in Eq. (4.1); Ref. [106] uses the opposite labeling. The paths used in this work for $\boldsymbol{b}$ and $\eta\boldsymbol{v}$ are given in Sections 5.4 and 5.5. The three-point function is [106]

$$C^{\mathrm{3pt}}_{\Gamma,q}(\tau,\boldsymbol{P};\mathcal{C}^{\mathrm{lat}}) = \frac{1}{\hat{L}^3}\sum_{z}\sum_{\beta\alpha}\Gamma^{\mathrm{3pt}}_{\beta\alpha}\left\langle \chi_\alpha(t_{\mathrm{snk}},\boldsymbol{P})\, O^{\mathrm{lat}}_{\Gamma,q}(\mathcal{C}^{\mathrm{lat}};z,\tau)\,\bar{\chi}_\beta(t_{\mathrm{src}},\boldsymbol{P})\right\rangle , \tag{3.48}$$

with $\hat{L}$ the spatial extent in lattice units. Here P denotes the nucleon momentum and $U(P,S)$ the corresponding spinor. Source and sink carry the same momentum $\boldsymbol{P}$: TMDs are defined by diagonal matrix elements, $P'=P$, whereas generalized TMDs also depend on the momentum transfer $\Delta = P'-P$ [75]. The spectral decomposition of Section 3.6 therefore applies with $\boldsymbol{p}'=\boldsymbol{p}=\boldsymbol{P}$, and Eq. (3.45) holds for every link path.

3.7.2 Contraction.

The connected contraction of Eq. (3.48) uses the forward propagator from the source to the quark field at one end of the link path and the sequential propagator from the sink to the antiquark field at the other end, joined by the link product [106]. The sequential propagator enters as a backward propagator, but it is stored in the format of a forward propagator, so Eq. (3.28) is applied before the contraction [106]. Schematically, for every site z in the operator time slice,

$$C^{\mathrm{3pt}}_{\Gamma}(z) = \mathrm{tr}\Big[\Gamma\,\mathcal{U}^{\mathrm{lat}}\big[\mathcal{C}^{\mathrm{lat}}+z\big]\; S_F\big(z+\ell\big)\;\gamma_5\, S_B(z)^\dagger\,\gamma_5\Big] , \tag{3.49}$$

where S_F is the forward propagator from the source, S_B the sequential propagator, and the trace runs over Dirac and color indices. In the lattice code used for this work, this contraction is implemented in Qlua as follows, with `gamma{n=15}` $=\Gamma(15)=\gamma_5$:

Listing 1: Contraction of the staple-link three-point function in Qlua.

```
local stap_fw_shifts_bw = (staple * fw_p:shift(b_dir, "from_forward")
                           * gamma{n=15} * bw_p:adjoin() * gamma{n=15})
for i_gn, gn in pairs(gamma_list) do
  -- bw_p is conjugated to do dot(bw, gamma{n=gn} * stap_fw_shifts)
  local bb = (gamma{n=gn} * stap_fw_shift_bw):trace()
  ...
end
```

The forward propagator `fw_p` is shifted along the separation direction `b_dir`, multiplied by the staple link product, and contracted with the γ_5-conjugated sequential propagator `bw_p`; the trace is then taken for every

Dirac structure $\Gamma(n)$ in `gamma_list`. Ref. [106] gives an equivalent implementation in Chroma. In the isospin limit the disconnected contribution again cancels in the isovector combination $u - d$ [75, 86].

3.7.3 Ratio and matrix element.

Since source and sink momenta are equal, the matrix element is extracted from the plateau of the forward ratio of Eq. (3.46) [106, 116],

$$R_{\Gamma,q}(\boldsymbol{P};\mathcal{C}^{\mathrm{lat}}) \equiv \mathrm{plateau}\left[\frac{C^{3\mathrm{pt}}_{\Gamma,q}(\tau,\boldsymbol{P};\mathcal{C}^{\mathrm{lat}})}{C^{2\mathrm{pt}}(t_{\mathrm{snk}}-t_{\mathrm{src}},\boldsymbol{P})}\right] = \sum_{S,S'} \frac{\bar{U}(P,S)\,\Gamma^{3\mathrm{pt}}\,U(P,S')}{2E_P\,\mathrm{tr}\big[\Gamma^{+}\big(-i\not{P}+m_N\big)\big]}\,\big\langle N(P,S')\big|O^{\mathrm{lat}}\big|N(P,S)\big\rangle\ . \tag{3.50}$$

The ratios $R_{\Gamma(n)}$ for $\Gamma(1),\Gamma(2),\Gamma(4),\Gamma(8)=\gamma_1,\gamma_2,\gamma_3,\gamma_4$ are the input from which the invariant amplitudes are solved in Section 5.6. As discussed in Section 2.9, the soft factor and the renormalization of the gauge link multiply the correlator independently of Γ, so they cancel in ratios of amplitudes taken at the same b, P and staple geometry, such as the Sivers shift [75, 86]. Whether this multiplicative structure carries over exactly to the lattice is not guaranteed and has been examined in Ref. [86]. The following chapters therefore work with such ratios. Section 4 gives the Lorentz-invariant parametrization of $\langle N|O^{\mathrm{lat}}|N\rangle$ that turns the ratios $R_{\Gamma(n)}$ into invariant amplitudes.

4 TMD CORRELATOR FOR LATTICE CALCULATIONS

This chapter defines the terms used in the rest of the thesis. Section 4.1 defines the TMD correlator as the Fourier transform of a bare non-local matrix element divided by the soft factor, and identifies the unpolarized distribution f_1 and the T-odd Sivers function $f_{1T}^{\perp}$ in its γ^+ projection. Section 4.2 introduces $\boldsymbol{b}_{\mathrm{T}}$-dependent TMDs and their derivatives, which the lattice evaluates directly, and keeps $|\boldsymbol{b}_{\mathrm{T}}|$ finite because $f_{1T}^{\perp(1)}(x)$ diverges without additional regularization. Section 4.3 introduces the staple-shaped gauge link and the Collins-Soper parameter $\hat{\zeta}$. Section 4.4 lists the Lorentz, parity, time-reversal, and Hermitian conjugation transformations, parametrizes the b-dependent correlator in terms of the Lorentz invariant amplitudes $\widetilde{A}_i$ and $\widetilde{B}_i$, and uses hermiticity to show that their real and imaginary parts are even and odd in b. Section 4.6 combines these amplitudes into $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$, from which f_1 and $f_{1T}^{\perp}$ follow by Fourier transformation.

4.1 Definition of TMDs

Consider a frame of reference where the nucleon has large momentum in the z-direction, i.e., $P^+ \gg m_N$, $\boldsymbol{P}_{\mathrm{T}} = 0$. In light cone coordinates, the components k^+, $\boldsymbol{k}_{\mathrm{T}}$, k^- of the quark momentum k scale as P^+/m_N, 1, m_N/P^+, respectively, under boosts along the z-axis. Thus, the quark longitudinal momentum fraction $x = k^+/P^+$ and its transverse momentum $\boldsymbol{k}_{\mathrm{T}}$ are invariant under boosts along the z-axis, while the k^- component is suppressed. This leads to transverse momentum-dependent parton distribution functions (TMDs), which depend on the longitudinal momentum fraction $x \equiv k^+/P^+$ and the quark transverse momentum $\boldsymbol{k}_{\mathrm{T}}$. The transverse momentum components $\boldsymbol{k}_{\mathrm{T}}$ describe an intrinsic motion of the quarks inside the proton that occurs independently of the momentum of the proton itself. TMDs are also an important ingredient for understanding the origin of the large angular and spin asymmetries observed in experiments such as semi-inclusive deep inelastic scattering (SIDIS) and the Drell-Yan process (DY).

We construct a correlator of the general form from the gauge-invariant bilocal operator $\bar{q}(0)\Gamma\mathcal{U}[\mathcal{C}_b]q(b)$ acting on the nucleon state $|P,S\rangle$, where $\bar{q}$ and q are local quark field operators, and the Wilson line $\mathcal{U}[\mathcal{C}_b]$ makes the bilocal operator gauge invariant. The Fourier transform of such a position space correlator $\langle P,S|\ \bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\ q(b)\ |P,S\rangle$ gives the momentum space correlator $\Phi^{[\Gamma]}(k,P,S;\ldots)$, or equivalently the TMDs. The Wilson line introduces

divergences that conventional dimensional regularization cannot treat [116], so we divide by the soft factor, $\widetilde{\mathcal{S}}(b^2;\ldots)$,

$$\Phi^{[\Gamma]}(k,P,S;\ldots) \;\equiv\; \int \frac{d^4b}{(2\pi)^4}\, e^{ik\cdot b}\, \frac{\overbrace{\frac{1}{2}\,\langle P,S|\;\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\;q(b)\;\;|P,S\rangle}^{\equiv\,\widetilde{\Phi}^{[\Gamma]}_{\text{unsubtr.}}(b,P,S;\ldots)}}{\widetilde{\mathcal{S}}(b^2;\ldots)}\,. \tag{4.1}$$

The numerator, $\widetilde{\Phi}^{[\Gamma]}_{\text{unsubtr.}}(b,P,S;\ldots)$, is the bare non-local matrix element in which a gauge link $\mathcal{U}[\mathcal{C}_b]$ connects the quark fields. The denominator is the soft factor, $\widetilde{\mathcal{S}}(b^2;\ldots)$, which regulates rapidity divergences. The ellipses ("…") denote additional dependence on parameters governing the Wilson line geometry and the soft factor, as well as an implicit dependence on the ultraviolet renormalization scale μ.

The soft factor, $\tilde{\mathcal{S}}$, stands symbolically for an expression that can, depending on the formalism, involve several vacuum expectation values. In the framework of Collins and Aybat [31, 23, 107], it takes the form

$$\widetilde{\mathcal{S}}(b^2;\ldots) = \sqrt{\frac{\tilde{S}_{(0)}(\boldsymbol{b}_{\rm T},+\infty,-\infty)\;\tilde{S}_{(0)}(\boldsymbol{b}_{\rm T},y_s,-\infty)}{\tilde{S}_{(0)}(\boldsymbol{b}_{\rm T},+\infty,y_s)}}\,, \tag{4.2}$$

here each of the objects $\tilde{S}_{(0)}(\boldsymbol{b}_{\rm T},\ldots)$ is a vacuum expectation value of Wilson line structures. In this framework, we start with space-like Wilson lines. Some Wilson lines remain tilted away from the light cone, leading to the dependence on the rapidity parameter y_s in the above expression. We need not specify $\tilde{\mathcal{S}}$ further, since it cancels in the observables we consider.

To project the correlator onto the relevant kinematics, we integrate over the suppressed momentum component k^-, which sets the position-space separation to $b^+=0$:

$$\begin{aligned}
\Phi^{[\Gamma]}(x,\boldsymbol{k}_{\rm T};P,S;\ldots) &\equiv \int dk^-\,\Phi^{[\Gamma]}(k,P,S;\ldots)\\
&= \int dk^- \int \frac{d^4b}{(2\pi)^4}\, e^{ik\cdot b}\,\frac{\frac{1}{2}\,\langle P,S|\;\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\;q(b)\;\;|P,S\rangle}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\ldots)} &(4.3)\\
&= \int dk^- \int \frac{db^+}{2\pi} e^{ik^-b^+}\int\frac{d^2\mathbf{b}_T}{(2\pi)^2}\int\frac{db^-}{2\pi}e^{ik^+b^- - i\mathbf{k}_T\cdot\mathbf{b}_T}\,\frac{\frac{1}{2}\,\langle P,S|\;\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\;q(b)\;\;|P,S\rangle}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\ldots)} &(4.4)\\
&= \int db^+\delta(b^+)\int\frac{d^2\mathbf{b}_T}{(2\pi)^2}\int\frac{db^-}{2\pi}e^{ik^+b^- - i\mathbf{k}_T\cdot\mathbf{b}_T}\,\frac{\frac{1}{2}\,\langle P,S|\;\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\;q(b)\;\;|P,S\rangle}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\ldots)} &(4.5)\\
&= \int\frac{d^2\boldsymbol{b}_{\rm T}}{(2\pi)^2}\int\frac{d(b\cdot P)}{(2\pi)P^+}\;e^{ix(b\cdot P)-i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\;\frac{\frac{1}{2}\,\langle P,S|\;\bar{q}(0)\,\Gamma\,\mathcal{U}[\mathcal{C}_b]\;q(b)\;\;|P,S\rangle}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\ldots)}\Bigg|_{b^+=0}\,. &(4.6)
\end{aligned}$$

As a consequence, $x\leftrightarrow(b\cdot P)$ and $\boldsymbol{k}_{\rm T}\leftrightarrow\boldsymbol{b}_{\rm T}$ are independent pairs of Fourier conjugate variables. With the Dirac matrix $\Gamma=\gamma^+$, which projects onto the leading-twist unpolarized quark distribution, the correlator becomes

[27, 29, 22, 102]:

$$\Phi^{[\gamma^+]}(x, \boldsymbol{k}_{\rm T}; P, S, \ldots) = f_1 - \left[\frac{\epsilon_{ij}\, \boldsymbol{k}_i\, \boldsymbol{S}_j}{m_N}\; f_{1T}^{\perp} \right]_{\rm odd} . \tag{4.7}$$

The term enclosed in $[\ ]_{\rm odd}$ isolates the naively time-reversal odd (T-odd) Sivers function $f_{1T}^{\perp}$ [44], which describes the distribution of unpolarized quarks inside a transversely polarized nucleon. The origin of this parametrization and the special role of T-odd TMDs become clear once we discuss the geometry of the gauge link path $\mathcal{C}_b$ and symmetry transformation properties.

4.2 TMDs in Fourier space

Our lattice method evaluates the b-dependent matrix elements $\widetilde{\Phi}^{[\Gamma]}_{\rm unsubtr.}(b, P, S; \ldots)$ introduced in Eq. (4.1). It is therefore more direct and natural to state our results in terms of Fourier-transformed, $\boldsymbol{b}_{\rm T}$-dependent TMDs and their $\boldsymbol{b}_{\rm T}$-derivatives. For a generic TMD f, we define the Fourier transform and its derivatives:

$$\tilde{f}(x, \boldsymbol{b}_{\rm T}^2; \ldots) \equiv \int d^2\boldsymbol{k}_{\rm T}\, e^{i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\; f(x, \boldsymbol{k}_{\rm T}^2; \ldots) = 2\pi \int d|\boldsymbol{k}_{\rm T}||\boldsymbol{k}_{\rm T}|\; J_0(|\boldsymbol{b}_{\rm T}||\boldsymbol{k}_{\rm T}|)\; f(x, \boldsymbol{k}_{\rm T}^2; \ldots) , \tag{4.8}$$

$$\tilde{f}^{(n)}(x, \boldsymbol{b}_{\rm T}^2 \ldots) \equiv n! \left(-\frac{2}{m_N^2} \partial_{\boldsymbol{b}_{\rm T}^2} \right)^n\; \tilde{f}(x, \boldsymbol{b}_{\rm T}^2; \ldots) = \frac{2\pi\; n!}{(m_N^2)^n} \int d|\boldsymbol{k}_{\rm T}||\boldsymbol{k}_{\rm T}| \left(\frac{|\boldsymbol{k}_{\rm T}|}{|\boldsymbol{b}_{\rm T}|} \right)^n J_n(|\boldsymbol{b}_{\rm T}||\boldsymbol{k}_{\rm T}|)\; f(x, \boldsymbol{k}_{\rm T}^2; \ldots) , \tag{4.9}$$

where J_n are the Bessel functions of the first kind, as shown in Ref. [117]; these Bessel-weighted quantities can be used to structure evolution equations.

In the formal limit $|\boldsymbol{b}_{\rm T}| \to 0$, they reduce to the conventional $\boldsymbol{k}_{\rm T}$-moments:

$$\tilde{f}^{(n)}(x, 0; \ldots) = \int d^2\boldsymbol{k}_{\rm T} \left(\frac{\boldsymbol{k}_{\rm T}^2}{2m_N^2} \right)^n f(x, \boldsymbol{k}_{\rm T}^2; \ldots) \equiv f^{(n)}(x) . \tag{4.10}$$

However, the first transverse moment of the Sivers function, $f_{1T}^{\perp(1)}(x)$, diverges without additional regularization, because the perturbatively calculable high-$\boldsymbol{k}_{\rm T}$ tails do not decay fast enough [108]. We therefore do not extrapolate to $\boldsymbol{b}_{\rm T} = 0$. At finite $|\boldsymbol{b}_{\rm T}|$, the integrals converge, and we access the x-dependence of the Sivers TMD through the conjugate variable $b{\cdot}P$ [82, 83].

4.3 Link geometry

The geometry of the gauge link path $\mathcal{C}_b$ affects both the number of allowed TMD structures and the numerical result.

The simplest link geometry is a straight line connecting the quark fields at 0 and b. Refs. [82, 83] have studied TMDs with straight gauge links on the lattice. These "process-independent" TMDs are theoretically interesting in their own right, but it is not yet known how to relate them quantitatively to the TMDs that appear in scattering experiments. The operator with straight gauge links has the greatest possible symmetry; as a result, T-odd TMDs vanish for straight gauge links.

To accommodate the symmetries required to isolate the Sivers effect and its x dependence, we use a staple-shaped gauge link; see Fig. 6:

$$\mathcal{U}[\mathcal{C}_b^{(\eta v)}] = \mathcal{U}[0, \eta v, \eta v + b, b] \ . \tag{4.11}$$



Figure 6: Staple-shaped gauge link $\mathcal{U}[0, \eta v, \eta v + b, b]$. The staple points along the direction v, in general $v = (0, v_1, v_2, v_3)$, and the limit $\eta \to \infty$ extends it.

Here v and b are both purely spatial, $v = (0, v_1, v_2, v_3)$ and $b = (0, b_1, b_2, b_3)$. The vector v sets the direction of the staple and η sets its extent. We need this off-axis staple because the x dependence requires a wide scan of $P \cdot b$ space, or equivalently b space. Later sections discuss this constraint in detail.

The Lorentz-invariant Collins-Soper parameter, $\zeta \equiv 2v{\cdot}P/\sqrt{|v^2|}$, parameterizes the approach toward the physical light-like limit ($v = n$). As v becomes light-like, $\zeta \to \infty$; the parameter acts as an artificial scale that regulates the rapidity divergences [25, 118, 107]. Expressing the vectors P and v via rapidities y_P and y_v (we use $P^\pm = m_N e^{\pm y_P}/\sqrt{2}$ and $v^+/v^- = -e^{2y_v}$), we define the dimensionless variable:

$$\hat{\zeta} \equiv \zeta/2m_N = \frac{v \cdot P}{\sqrt{|v^2|}\sqrt{P^2}} = \frac{v^+P^- + v^-P^+}{\sqrt{|v^2|}m_N} = \sinh(y_P - y_v) \ , \tag{4.12}$$

So $\hat{\zeta}$ measures the rapidity difference between the nucleon momentum and the Wilson line direction. This allows us to boost to a frame where v is purely spatial ($v^0 = 0, y_v = 0$).

4.4 Symmetry Constraints

Discrete symmetries constrain the correlation function. Consider the unsubtracted correlator [117]:

$$\Phi^{[\Gamma]}_{\text{unsub}}(k,P,S;v,\mu) = \int \frac{d^4b}{(2\pi)^4}\, e^{ik\cdot b} \underbrace{\frac{1}{2}\,\langle P,S|\ \bar{q}(0)\ \overbrace{\mathcal{U}[0,\infty v]\,\mathcal{U}[\infty v,b]}^{\mathcal{U}[\mathcal{C}_b]}\Gamma\ q(b)\ |P,S\rangle}_{\widetilde{\Phi}^{[\Gamma]}_{\text{unsub}}(b,P,S;v,\mu)} \ . \tag{4.13}$$

For unpolarized quarks ($\Gamma = \gamma^\mu$), its decomposition into real-valued Lorentz-invariant amplitudes reads

$$\begin{aligned}\frac{1}{2}\Phi^{[\gamma^\mu]}_{\text{unsub}} &= P^\mu\, A_2^{(+)} + k^\mu\, A_3^{(+)} + \frac{1}{m_N}\epsilon^{\mu\nu\alpha\beta}P_\nu k_\alpha S_\beta\, A_{12}^{(+)} + \frac{m_N^2}{(v\cdot P)}v^\mu\, B_1^{(+)} \\ &+ \frac{m_N}{v\cdot P}\epsilon^{\mu\nu\alpha\beta}P_\nu v_\alpha S_\beta\, B_7^{(+)} + \frac{m_N}{v\cdot P}\epsilon^{\mu\nu\alpha\beta}k_\nu v_\alpha S_\beta\, B_8^{(+)} \\ &+ \frac{1}{m_N(v\cdot P)}(k\cdot S)\epsilon^{\mu\nu\alpha\beta}P_\nu k_\alpha v_\beta\, B_9^{(+)} + \frac{m_N}{(v\cdot P)^2}(v\cdot S)\epsilon^{\mu\nu\alpha\beta}P_\nu k_\alpha v_\beta B_{10}^{(+)}\ .\end{aligned} \tag{4.14}$$

On applying Lorentz transformations (L), parity transformation (P), time-reversal (T), and Hermitian conjugation ($\dagger$) to the matrix elements, we find that the correlator fulfills [116, 117]:

$$(L): \qquad \Phi^{[\Gamma]}_{\text{unsub}}(k,P,S;v,\mu) = \Phi^{[\Lambda_{1/2}^{-1}\Gamma\Lambda_{1/2}]}_{\text{unsub}}(\Lambda k,\Lambda P,\Lambda S;\Lambda v,\mu)\ , \tag{4.15}$$

$$(P): \qquad \Phi^{[\Gamma]}_{\text{unsub}}(k,P,S;v,\mu) = \Phi^{[\gamma^0\Gamma\gamma^0]}_{\text{unsub}}(\overline{k},\overline{P},-\overline{S};\overline{v},\mu)\ , \tag{4.16}$$

$$(T): \qquad \Big[\Phi^{[\Gamma]}_{\text{unsub}}(k,P,S;v,\mu)\Big]^* = \Phi^{[\gamma^1\gamma^3\Gamma^*\gamma^3\gamma^1]}_{\text{unsub}}(\overline{k},\overline{P},\overline{S};-\overline{v},\mu)\ , \tag{4.17}$$

$$(\dagger): \qquad \Big[\Phi^{[\Gamma]}_{\text{unsub}}(k,P,S;v,\mu)\Big]^* = \Phi^{[\gamma^0\Gamma^\dagger\gamma^0]}_{\text{unsub}}(k,P,S;v,\mu)\ . \tag{4.18}$$

where, for a vector c, $\overline{c} \equiv (c^0,-c^1,-c^2,-c^3)$ denotes the vector with its spatial components reversed in sign. Hermiticity ($\dagger$) implies that the $A_i^{(+)}$ and $B_i^{(+)}$ in Eq. (4.14) are real valued. Time reversal (T) does not constrain the number of allowed structures, because it changes the sign of $(v\cdot P)$. Instead, time reversal (T) relates the SIDIS amplitudes $A_i^{(+)}$, $B_i^{(+)}$ to the Drell-Yan amplitudes $A_i^{(-)}$, $B_i^{(-)}$.

For any of the transformations $\mathcal{T} \in \{L,P,T,\dagger\}$, the Eqs. (4.15)-(4.18) are of the general form

$$\mathcal{T}_\Phi\left(\Phi(k,w)\right) = \Phi\left(\mathcal{T}_k(k),\mathcal{T}_w(w)\right) \tag{4.19}$$

where we omit the subscript "unsub" and the renormalization scale μ, and the symbol w collects all dependence on Γ, P, S and v. Here $\mathcal{T}_\Phi$ is either the identity function or complex conjugation. The transformation rule $\mathcal{T}_k(k)$

maps onto Λk, k or $\overline{k}$ and thus fulfills $a{\cdot}b = \mathcal{T}_k(a){\cdot}\mathcal{T}_k(b)$ for any two vectors a and b. The Fourier-transformed correlator

$$\widetilde{\Phi}(b,w) = \int d^4k\; e^{-ik\cdot b}\Phi(k,w) \tag{4.20}$$

transforms according to

$$\begin{aligned}
\mathcal{T}_\Phi\left(\widetilde{\Phi}(b,w)\right) &= \int d^4k\; e^{\mathcal{T}_\Phi(-i)\,k\cdot b}\;\mathcal{T}_\Phi\left(\Phi(k,w)\right) \\
&= \int d^4q\; e^{\mathcal{T}_\Phi(-i)\,\mathcal{T}_k^{-1}(q)\cdot b}\;\Phi\left(q,\mathcal{T}_w(w)\right) \text{.....(change of Variables: } k\to q=\mathcal{T}_k(k) \text{ or } k=\mathcal{T}_k^{-1}(q)) \\
&= \int d^4q\; e^{\mathcal{T}_\Phi(-i)\,q\cdot\mathcal{T}_k(b)}\;\Phi\left(q,\mathcal{T}_w(w)\right) \text{.....(}\because q{\cdot}b=\mathcal{T}_k(q){\cdot}\mathcal{T}_k(b)) \\
&= \int d^4q\; e^{-iq\left(\frac{\mathcal{T}_\Phi(i)}{i}\cdot\mathcal{T}_k(b)\right)}\;\Phi\left(q,\mathcal{T}_w(w)\right) \\
&= \widetilde{\Phi}\left(\frac{\mathcal{T}_\Phi(i)}{i}\mathcal{T}_k(b),\mathcal{T}_w(w)\right)\,.
\end{aligned} \tag{4.21}$$

In particular, under Hermitian conjugation the position-space correlator transforms as

$$(\dagger): \qquad \left[\widetilde{\Phi}^{[\Gamma]}_{\text{unsub}}(b,P,S;v)\right]^* = \widetilde{\Phi}^{[\gamma^0\Gamma^\dagger\gamma^0]}_{\text{unsub}}(-b,P,S;v)\;. \tag{4.22}$$

Let $f(k,w)$ be any of the structures preceding the invariant amplitudes in the parameterization of Φ. The structure $f(k,w)$ is a homogeneous function of some degree n in k, i.e., $f(\alpha k,w)=\alpha^n f(k,w)$ for any number α. For example, the structure $f(k,w)=\frac{1}{m_N(v\cdot P)}(k\cdot S)\epsilon^{\mu\nu\alpha\beta}P_\nu k_\alpha v_\beta$ preceding $B_9^{(+)}$ in Eq. (4.14) has degree $n=2$. If we define $\tilde{f}(b,w)\equiv f(-im_N^2 b,w)$, then

$$\mathcal{T}_\Phi\left(\tilde{f}(b,w)\right) = \mathcal{T}_\Phi(-im_N^2)^n\,\mathcal{T}_\Phi\left(f(b,w)\right) = f\left(\mathcal{T}_\Phi(-im_N^2)\mathcal{T}_k(b),\mathcal{T}_w(w)\right) = \tilde{f}\left(\frac{\mathcal{T}_\Phi(i)}{i}b,w\right)\,. \tag{4.23}$$

This shows that $\tilde{f}$ transforms like $\widetilde{\Phi}$ in Eq. (4.21). The parameterization of $\widetilde{\Phi}$ therefore follows from the substitution $k\to -im_N^2 b$ in the structures parameterizing Φ, which gives

$$\begin{aligned}
\frac{1}{2}\widetilde{\Phi}^{[\gamma^\mu]}_{\text{unsub}} =& P^\mu\,\widetilde{A}_2 - im_N^2 b^\mu\,\widetilde{A}_3 - im_N\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha S_\beta\,\widetilde{A}_{12} + \frac{m_N^2}{(v\cdot P)}v^\mu\,\widetilde{B}_1 \\
&+ \frac{m_N}{v\cdot P}\epsilon^{\mu\nu\alpha\beta}P_\nu v_\alpha S_\beta\,\widetilde{B}_7 - \frac{im_N^3}{v\cdot P}\epsilon^{\mu\nu\alpha\beta}b_\nu v_\alpha S_\beta\,\widetilde{B}_8 \\
&- \frac{m_N^3}{v\cdot P}(b\cdot S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha v_\beta\,\widetilde{B}_9 - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha v_\beta\widetilde{B}_{10}\;.
\end{aligned} \tag{4.24}$$

The amplitudes $\widetilde{A}_i$ and $\widetilde{B}_i$ introduced this way are no longer constrained to be real-valued functions. Instead,

Hermitian conjugation, Eq. (4.22), gives the relation

$$\left[\widetilde{A}_i(b^2, b{\cdot}P, v{\cdot}b/(v{\cdot}P), \zeta^{-2}, \mu^2)\right]^* = \widetilde{A}_i(b^2, -b{\cdot}P, -v{\cdot}b/(v{\cdot}P), \zeta^{-2}, \mu^2)\,. \tag{4.25}$$

In general, the amplitudes are complex,

$$\widetilde{A}_i = \widetilde{A}_i^{\mathrm{Re}} + i\widetilde{A}_i^{\mathrm{Im}} \tag{4.26}$$

$$\widetilde{B}_i = \widetilde{B}_i^{\mathrm{Re}} + i\widetilde{B}_i^{\mathrm{Im}} \tag{4.27}$$

Given the amplitudes at both $\pm b$, Eq. (4.25) implies

$$\widetilde{A}_i^{\mathrm{Re}} = \widetilde{A}_i^{(\text{b-even})} \tag{4.28a}$$

$$\widetilde{A}_i^{\mathrm{Im}} = \widetilde{A}_i^{(\text{b-odd})} \tag{4.28b}$$

$$\widetilde{B}_i^{\mathrm{Re}} = \widetilde{B}_i^{(\text{b-even})} \tag{4.28c}$$

$$\widetilde{B}_i^{\mathrm{Im}} = \widetilde{B}_i^{(\text{b-odd})} \tag{4.28d}$$

4.5 Correlator Parametrization

To translate Euclidean space lattice calculations into TMDs, which are defined and interpreted in light cone coordinates, we need the parametrization of the correlator $\tilde{\Phi}_{\text{unsubtr.}}$ in terms of manifestly Lorentz-invariant amplitudes. A parametrization of the correlator $\Phi_{\text{unsubtr.}}(k, P, S; \infty v, \mu)$ for link paths that extend to infinity in a direction v has been worked out in Ref. [102] and involves 32 independent amplitudes A_i and B_i that depend on the Lorentz-invariant quantities k^2, $k \cdot P$, $k{\cdot}v/v{\cdot}P$ and $\hat{\zeta}$. Section 4.4 shows that a parametrization of the corresponding b-dependent correlator $\tilde{\Phi}_{\text{unsubtr.}}$ is of the same form as the parametrization of $\Phi_{\text{unsubtr.}}$ if we substitute $k \to -im_N^2 b$, which is Eq. (4.24) (for $\Gamma = \gamma^\mu$):

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^\mu]}_{\text{unsub}} =& P^\mu\, \widetilde{A}_2 - im_N^2 b^\mu\, \widetilde{A}_3 - im_N \epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha S_\beta\, \widetilde{A}_{12} + \frac{m_N^2}{(v{\cdot}P)} v^\mu\, \widetilde{B}_1 \\ &+ \frac{m_N}{v{\cdot}P}\epsilon^{\mu\nu\alpha\beta} P_\nu v_\alpha S_\beta\, \widetilde{B}_7 - \frac{im_N^3}{v{\cdot}P}\epsilon^{\mu\nu\alpha\beta} b_\nu v_\alpha S_\beta\, \widetilde{B}_8 \\ &- \frac{m_N^3}{v{\cdot}P}(b{\cdot}S)\epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha v_\beta\, \widetilde{B}_9 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)\epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha v_\beta \widetilde{B}_{10}\,.\end{aligned} \tag{4.29}$$

For the lattice calculations, it is advantageous to start with an equivalent parametrization in which the structures explicitly depend on the staple extent η and which is still well-defined for $v{\cdot}P = 0$. Such a parametrization can

be obtained from the parametrization above by replacing $v \to \eta v$ and by leaving out the factors $(v{\cdot}P)^{-n}$.

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^\mu]}_{\text{unsub}} =& P^\mu\,\tilde{a}_2 - im_N^2 b^\mu\,\tilde{a}_3 - im_N\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha S_\beta\,\tilde{a}_{12}\\ &+ m_N^2\eta v^\mu\,\tilde{b}_1 + m_N\epsilon^{\mu\nu\alpha\beta}P_\nu\eta v_\alpha S_\beta\,\tilde{b}_7 - im_N^3\epsilon^{\mu\nu\alpha\beta}b_\nu\eta v_\alpha S_\beta\,\tilde{b}_8\\ &- m_N^3(b{\cdot}S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha\eta v_\beta\,\tilde{b}_9 - im_N^3(v{\cdot}S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha\eta^2 v_\beta\tilde{b}_{10}\end{aligned} \tag{4.30}$$

Here we have used lowercase amplitudes to distinguish the two parametrizations. The relation to the uppercase amplitudes is given by

$$\begin{aligned}\widetilde{A}_i\left(b^2, b{\cdot}P, \frac{v{\cdot}b}{v{\cdot}P}, \frac{v^2}{(v{\cdot}P)^2}, \eta v{\cdot}P\right) &= \tilde{a}_i(b^2, b{\cdot}P, \eta v{\cdot}b, (\eta v)^2, \eta v{\cdot}P)\,,\\ \widetilde{B}_i\left(b^2, b{\cdot}P, \frac{v{\cdot}b}{v{\cdot}P}, \frac{v^2}{(v{\cdot}P)^2}, \eta v{\cdot}P\right) &= (\eta v{\cdot}P)^n\;\tilde{b}_i(b^2, b{\cdot}P, \eta v{\cdot}b, (\eta v)^2, \eta v{\cdot}P)\,.\end{aligned} \tag{4.31}$$

where n is the power with which $v{\cdot}P$ appears in the denominator in front of the corresponding amplitude $\widetilde{B}_i$ in the parametrization. Notice that the $\tilde{a}_i$ and $\tilde{b}_i$ are functions of all the Lorentz-invariant products of b, P and ηv. For the uppercase amplitudes, however, we choose to represent the dependence on these invariants in the third and fourth arguments by η-independent expressions, to facilitate taking the limit $\eta \to \pm\infty$. The dependence on the Collins-Soper parameter $\hat{\zeta}$ is given by the fourth argument, $v^2/(v{\cdot}P)^2 = -1/(m_N\hat{\zeta})^2$, while the fifth argument, $\eta v{\cdot}P$, characterizes the length of the gauge link and distinguishes between future- and past-pointing Wilson lines. In our frame ($b^+ = 0$, $\boldsymbol{v}_\text{T} = \boldsymbol{P}_\text{T} = 0$), the kinematics fixes the relation between the invariants [106, 119]:

$$\frac{v{\cdot}b}{v{\cdot}P} = b{\cdot}P\frac{R(\hat{\zeta}^2)}{m_N^2}\,, \quad \text{where} \quad R(\hat{\zeta}^2) \equiv 1 - \sqrt{1+\hat{\zeta}^{-2}} = \frac{m_N^2}{v{\cdot}P}\frac{v^+}{P^+}\,. \tag{4.32}$$

The relation in Eq. (4.32) shows that the third argument of the $\widetilde{A}_i$ and $\widetilde{B}_i$ is not independent of the others in the context of TMDs. Moreover, in our lattice calculations, we have to choose the link directions b and v such that Eq. (4.32) is fulfilled. As a side remark, the parameter corresponding to Eq. (4.32) in momentum space is $v{\cdot}k/v{\cdot}P \approx x$, i.e., the amplitudes A_i and B_i acquire an explicit x-dependence, which has already been pointed out in Refs. [106, 119].

For the $\Gamma = \gamma^+$ structure at leading twist, the correlator can be written in the form

$$\frac{1}{2P^+}\widetilde{\Phi}^{[\gamma^+]}_{\text{unsubtr.}} = \widetilde{A}_{2B} + im_N\epsilon_{ij}\boldsymbol{b}_i\boldsymbol{S}_j\,\widetilde{A}_{12B}\,, \tag{4.33}$$

where the indices i, j correspond to transverse directions, $i, j \in \{1, 2\}$ (we use the convention $\epsilon^{0123} = 1$ for the totally antisymmetric Levi-Civita symbol, and introduce $\epsilon_{ij} \equiv \epsilon^{-+ij}$ such that $\epsilon_{12} = 1$), and where we have introduced the following abbreviations for combinations of amplitudes:

$$\begin{aligned} \widetilde{A}_{2B} &\equiv \widetilde{A}_2 + R(\hat{\zeta}^2)\widetilde{B}_1 \\ \widetilde{A}_{12B} &\equiv \widetilde{A}_{12} - R(\hat{\zeta}^2)\widetilde{B}_8 \,. \end{aligned} \tag{4.34}$$

The unpolarized distribution f_1 and the Sivers distribution $f_{1T}^\perp$ follow from the Fourier integration operator $\int_{\mathcal{F}}$:

$$\begin{aligned} \int_{\mathcal{F}} \widetilde{A}_i &\equiv \int \frac{d^2\boldsymbol{b}_\mathrm{T}}{(2\pi)^2}\, e^{-i\boldsymbol{b}_\mathrm{T}\cdot\boldsymbol{k}_\mathrm{T}}\, \frac{1}{\widetilde{\mathcal{S}}(b^2;\ldots)} \int \frac{d(b{\cdot}P)}{(2\pi)}\, e^{ix(b\cdot P)}\, \widetilde{A}_i(-\boldsymbol{b}_\mathrm{T}^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P) \\ &= \int_0^\infty \frac{d(-b^2)}{2(2\pi)}\, \frac{J_0(\sqrt{-b^2}\,|\boldsymbol{k}_\mathrm{T}|)}{\widetilde{\mathcal{S}}(b^2;\ldots)} \int \frac{d(b{\cdot}P)}{(2\pi)}\, e^{ix(b\cdot P)}\, \widetilde{A}_i(b^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P) \end{aligned} \tag{4.35}$$

Applied to the combined amplitudes, it gives the leading-twist distributions:

$$\begin{aligned} f_1(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, \eta v{\cdot}P) &= 2\int_{\mathcal{F}} \widetilde{A}_{2B}\,, \\ f_{1T}^\perp(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, \eta v{\cdot}P) &= 4m_N^2 \partial_{\boldsymbol{k}_\mathrm{T}^2} \int_{\mathcal{F}} \widetilde{A}_{12B}\,. \end{aligned} \tag{4.36}$$

Here, the dots "…" indicate further parameters that specify the geometry of the soft factor. The f_1 is a T-even distribution which fulfills

$$f^{\text{T-even}}(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, \eta v{\cdot}P) = f^{\text{T-even}}(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, -\eta v{\cdot}P) \tag{4.37}$$

while the Sivers function $f_{1T}^\perp$ is a T-odd distribution which fulfills

$$f^{\text{T-odd}}(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, \eta v{\cdot}P) = -f^{\text{T-odd}}(x, \boldsymbol{k}_\mathrm{T}^2; \hat{\zeta}, \ldots, -\eta v{\cdot}P) \tag{4.38}$$

As a result, T-odd distributions must vanish for $\eta = 0$, which corresponds to straight gauge links. TMDs for SIDIS and DY are obtained for $\eta v{\cdot}P \to \infty$ and $\eta v{\cdot}P \to -\infty$, respectively. In the following, we choose $v{\cdot}P \geq 0$, such that the SIDIS and DY limits for space-like v can also be written as $\eta|v| \to \infty$ and $\eta|v| \to -\infty$, respectively.

Equation (4.36) shows that the x-dependent TMDs in Fourier space, $\tilde{f}(x, \boldsymbol{b}_\mathrm{T}^2)$ as defined in Eqs. (4.8) and (4.9), are given directly by the Fourier transform of the amplitudes $\widetilde{A}_{iB}$ with respect to $b{\cdot}P$. For brevity, define

the $x-(b{\cdot}P)$-Fourier-transformed, soft-factor-divided amplitudes

$$\widetilde{A}_i^{[x]}(x,\boldsymbol{b}_{\rm T}^2) \equiv \frac{1}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\dots)} \int_{-\infty}^{\infty} \frac{d(b{\cdot}P)}{2\pi}\, e^{ix(b\cdot P)}\, \widetilde{A}_i\big(-\boldsymbol{b}_{\rm T}^2, b{\cdot}P, \dots\big)\,, \tag{4.39}$$

so that, by Eq. (4.35), $\int_{\mathcal{F}} \widetilde{A}_i = \int \frac{d^2\boldsymbol{b}'_{\rm T}}{(2\pi)^2}\, e^{-i\boldsymbol{b}'_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\, \widetilde{A}_i^{[x]}(x,\boldsymbol{b}'^2_{\rm T})$. Inserting $f_1 = 2\int_{\mathcal{F}} \widetilde{A}_{2B}$ into Eq. (4.8), we have Unpolarized TMD,

$$\begin{aligned}\tilde{f}_1(x,\boldsymbol{b}_{\rm T}^2) &= 2\int d^2\boldsymbol{k}_{\rm T}\, e^{i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}} \int \frac{d^2\boldsymbol{b}'_{\rm T}}{(2\pi)^2}\, e^{-i\boldsymbol{b}'_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\, \widetilde{A}_{2B}^{[x]}(x,\boldsymbol{b}'^2_{\rm T}) = 2\int d^2\boldsymbol{b}'_{\rm T}\, \delta^{(2)}(\boldsymbol{b}_{\rm T}-\boldsymbol{b}'_{\rm T})\, \widetilde{A}_{2B}^{[x]}(x,\boldsymbol{b}'^2_{\rm T}) \\ &= 2\,\widetilde{A}_{2B}^{[x]}(x,\boldsymbol{b}_{\rm T}^2)\,,\end{aligned} \tag{4.40}$$

For Sivers TMD, on starting from $f_{1T}^{\perp} = 4m_N^2\,\partial_{\boldsymbol{k}_{\rm T}^2}\int_{\mathcal{F}}\widetilde{A}_{12B}$ and using $\boldsymbol{k}_j\,\partial_{\boldsymbol{k}_{\rm T}^2} = \frac{1}{2}\,\partial_{\boldsymbol{k}_j}$, differentiate Eq. (4.8) with respect to $\boldsymbol{b}_j$, we have

$$\begin{aligned}\partial_{\boldsymbol{b}_j}\tilde{f}_{1T}^{\perp} &= \int d^2\boldsymbol{k}_{\rm T}\,(i\boldsymbol{k}_j)\, e^{i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\, 4m_N^2\,\partial_{\boldsymbol{k}_{\rm T}^2}\int_{\mathcal{F}}\widetilde{A}_{12B} = 2im_N^2\int d^2\boldsymbol{k}_{\rm T}\, e^{i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\,\partial_{\boldsymbol{k}_j}\int_{\mathcal{F}}\widetilde{A}_{12B} \\ &= -2im_N^2\,(i\boldsymbol{b}_j)\int d^2\boldsymbol{k}_{\rm T}\, e^{i\boldsymbol{b}_{\rm T}\cdot\boldsymbol{k}_{\rm T}}\int_{\mathcal{F}}\widetilde{A}_{12B} = 2m_N^2\,\boldsymbol{b}_j\,\widetilde{A}_{12B}^{[x]}(x,\boldsymbol{b}_{\rm T}^2)\,,\end{aligned} \tag{4.41}$$

where we integrated by parts in $\boldsymbol{k}_j$ (the surface term vanishes for $|\boldsymbol{b}_{\rm T}|>0$) and used the same δ-function step as for f_1. Since $\partial_{\boldsymbol{b}_j}\tilde{f} = 2\boldsymbol{b}_j\,\partial_{\boldsymbol{b}_{\rm T}^2}\tilde{f}$, this gives $\partial_{\boldsymbol{b}_{\rm T}^2}\tilde{f}_{1T}^{\perp} = m_N^2\,\widetilde{A}_{12B}^{[x]}$. Eq. (4.9) with $n=1$ then yields

$$\tilde{f}_{1T}^{\perp(1)}(x,\boldsymbol{b}_{\rm T}^2) = -\frac{2}{m_N^2}\,\partial_{\boldsymbol{b}_{\rm T}^2}\tilde{f}_{1T}^{\perp}(x,\boldsymbol{b}_{\rm T}^2) = -2\,\widetilde{A}_{12B}^{[x]}(x,\boldsymbol{b}_{\rm T}^2)\,. \tag{4.42}$$

In summary, we have,

$$\begin{aligned}\tilde{f}_1^{(0)}(x,\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\dots,\eta v{\cdot}P) &= 2\int_{-\infty}^{\infty}\frac{d(b{\cdot}P)}{2\pi}\, e^{ix(b\cdot P)}\,\frac{\widetilde{A}_{2B}\big(-\boldsymbol{b}_{\rm T}^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P\big)}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\dots)}\,, \\ &= 2\,\widetilde{A}_{2B}^{[x]}(x,\boldsymbol{b}_{\rm T}^2)\,.\end{aligned} \tag{4.43}$$

$$\begin{aligned}\tilde{f}_{1T}^{\perp(1)}(x,\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\dots,\eta v{\cdot}P) &= -2\int_{-\infty}^{\infty}\frac{d(b{\cdot}P)}{2\pi}\, e^{ix(b\cdot P)}\,\frac{\widetilde{A}_{12B}\big(-\boldsymbol{b}_{\rm T}^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P\big)}{\widetilde{\mathcal{S}}(-\boldsymbol{b}_{\rm T}^2;\dots)}\,, \\ &= -2\,\widetilde{A}_{12B}^{[x]}(x,\boldsymbol{b}_{\rm T}^2)\,.\end{aligned} \tag{4.44}$$

This framework lays the groundwork for extracting its x-dependence. In subsequent chapters, for brevity, we may drop the soft-factor term from the above definitions, as soft factors cancel in the ratios we form.

4.6 Generalized Sivers shift from amplitudes

Eqs. (4.43)-(4.44) identifies x-dependent derivatives of Fourier-transformed TMDs with simple linear combinations of amplitudes $\widetilde{A}_i$ and $\widetilde{B}_i$ evaluated at the same values of $\boldsymbol{b}_{\rm T}^2$, $b\cdot P$, $\hat{\zeta}$ and $\eta v{\cdot}P$. Forming ratios of these

objects thus amounts to taking ratios of linear combinations of the fundamental correlators $\tilde{\Phi}^{[\Gamma]}_{\text{unsubtr.}}$ evaluated at the same point, i.e., with the same values of b, P, and ηv. For a discussion of the renormalization properties of ratios of the objects in (4.43)-(4.44), it is thus sufficient to understand the renormalization properties of (ratios formed from) the correlators $\widetilde{\Phi}^{[\Gamma]}_{\text{unsubtr.}} = \frac{1}{2}\ \langle P,S|\ \bar{q}(0)\,\Gamma\,\mathcal{U}\ q(b)\ \ |P,S\rangle$.

As discussed in Refs. [116, 86], analytical studies of the operator $\bar{q}(0)\,\Gamma\,\mathcal{U}q(b)$ in the continuum suggest that for $b^2 \neq 0$ the renormalization factors are multiplicative and Γ-independent. The basic reason is that the quark field operators are at different locations and undergo wave function renormalization separately. We will assume here that our lattice representation of $\bar{q}(0)\,\Gamma\,\mathcal{U}\,q(b)$ is renormalized multiplicatively independent of Γ as long as we keep $\boldsymbol{b}_{\rm T}^2$ larger than a few lattice spacings. A more detailed discussion and numerical studies of this operator's renormalization properties appear in Ref. [83]. Under the assumption of multiplicative renormalization, the generalized Sivers shift can only depend on $\boldsymbol{b}_{\rm T}^2$, $P\cdot b$, $\hat{\zeta}$, and the staple extent $\eta v{\cdot}P$. All other renormalization and soft-factor-related dependence cancels out in the ratio.

From phenomenology, the Sivers shift is the ratio defined as

$$\langle \boldsymbol{k}_y\rangle_{TU} \equiv m_N \frac{f_{1T}^{\perp(1)}(x)}{f_1^{(0)}(x)} = \left.\frac{\int d^2\boldsymbol{k}_{\rm T}\,\boldsymbol{k}_y\ \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}{\int d^2\boldsymbol{k}_{\rm T}\quad\ \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}\right|_{\boldsymbol{S}_{\rm T}=(1,0)}, \tag{4.45}$$

where γ^+ projects onto the leading twist. In the context of the density interpretation of TMDs mentioned in section 4.1, the ratio above yields the average transverse momentum in the y-direction, for quarks with a given longitudinal momentum fraction x inside a proton polarized in the x-direction. As mentioned before, understanding $f_{1T}^{\perp(1)}$ simply as a $\boldsymbol{k}_{\rm T}$-weighted TMD is problematic, since the $\boldsymbol{k}_{\rm T}$-integral is expected to be UV divergent. A natural way of circumventing this divergence is to generalize the Sivers shift to an expression in terms of the Fourier-transformed TMDs:

$$\begin{aligned}
\langle \boldsymbol{k}_y\rangle_{TU}(\boldsymbol{b}_{\rm T}^2,x,\hat{\zeta},\eta v{\cdot}P) \ &\equiv\ m_N \frac{\tilde{f}_{1T}^{\perp(1)}(\boldsymbol{b}_{\rm T}^2,x;\hat{\zeta},\ldots,\eta v{\cdot}P)}{\tilde{f}_1^{(0)}(\boldsymbol{b}_{\rm T}^2,x;\hat{\zeta},\ldots,\eta v{\cdot}P)} && (4.46)\\
&= -m_N \frac{\int d(b\cdot P)e^{ix(b\cdot P)}\widetilde{A}_{12B}(b^2,b{\cdot}P,(b{\cdot}P)R(\hat{\zeta}^2)/m_N^2,-1/(m_N\hat{\zeta})^2,\eta v{\cdot}P)}{\int d(b\cdot P)e^{ix(b\cdot P)}\widetilde{A}_{2B}(b^2,b{\cdot}P,(b{\cdot}P)R(\hat{\zeta}^2)/m_N^2,-1/(m_N\hat{\zeta})^2,\eta v{\cdot}P)} && (4.47)\\
&\xrightarrow{\boldsymbol{b}_{\rm T}^2=0}\ \left.\frac{\int d^2\boldsymbol{k}_{\rm T}\,\boldsymbol{k}_y\ \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}{\int d^2\boldsymbol{k}_{\rm T}\quad\ \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}\right|_{\boldsymbol{S}_{\rm T}=(1,0)} && (4.48)
\end{aligned}$$

In the limit $\boldsymbol{b}_{\rm T}^2 = 0$ we recover the Sivers shift (4.45), because the Fourier transformed TMDs $\tilde{f}_{1T}^{\perp(1)}$ and $\tilde{f}_1^{(0)}$ coincide with the moments $f_{1T}^{\perp(1)}$ and $f_1^{(0)}$, respectively. We are, however, interested in the generalized Sivers

shift for non-zero $\boldsymbol{b}_{\mathrm{T}}^2$, where the said UV-divergence disappears. The variable $\boldsymbol{b}_{\mathrm{T}}^2$ effectively acts as a regulator. Moreover, the $\boldsymbol{b}_{\mathrm{T}}$-dependence allows us to study differences in the widths of distributions on a qualitative level.

Furthermore, the generalized Sivers shift is T-odd, i.e., we expect to obtain results of opposite sign in the SIDIS and DY limits $\eta v{\cdot}P \to \infty$ and $\eta v{\cdot}P \to -\infty$, respectively, as demonstrated in Ref. [116, 84, 86]

In summary, the generalized Sivers shift describes a feature of the transverse momentum distribution of (unpolarized) quarks in a transversely polarized proton. In the formal limit $\boldsymbol{b}_{\mathrm{T}}^2 = 0$, it measures the dipole moment of that distribution orthogonal to the polarization of the proton.

In the next chapter, we explore the lattice QCD calculations of the generalized Sivers shift and extract its dependence on the longitudinal momentum fraction x.

5 LATTICE DATA & EXTRACTING THE $x-$DEPENDENCE OF SIVERS SHIFT

This Chapter connects the lattice correlators to the x-dependent Sivers shift. Section 5.1 writes the Sivers shift as a ratio of Fourier transforms over $b \cdot P$ of the amplitudes $\widetilde{A}_{12B}$ and $\widetilde{A}_{2B}$, and shows that the real parts enter with the cosine and the imaginary parts with the sine. Section 5.2 explains why extrapolating the Collins-Soper parameter $\hat{\zeta} \to \infty$ is needed. Sections 5.3 and 5.4 fix the staple direction v for each nucleon momentum so that we scan the whole $b \cdot P$, and describe how the correlator at the off-axis target $\eta\boldsymbol{v}/a$ is obtained by path averaging and quadratic interpolation. Section 5.5 describes the quark separation vectors $\boldsymbol{b}$, and the gauge link paths that correspond to them.

Section 5.6 projects the correlator parametrization onto the lattice kinematics and solves for the real and imaginary amplitudes from the symmetric and antisymmetric combinations of the longitudinal b data. Section 5.7 combines them into $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$, and Section 5.8 maps these from (b_L, b_T) to $(P \cdot b, b^2)$. Section 5.9 checks the parity of the extracted amplitudes, shows them at the four simulated momenta, and discards $\widetilde{A}^{\mathrm{Im}}_{12B}$, which is statistically compatible with zero.

5.1 The Sivers Shift as a Function of Momentum Fraction x

To connect our lattice observables to the physical, x-dependent Transverse Momentum Dependent Parton Distribution Functions (TMDs), we must isolate the relevant amplitudes from the correlation function. Focusing on the leading-twist unpolarized quark distribution, the correlator is parameterized by the Lorentz-invariant amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$, which we construct from the primitive correlator amplitudes in Section 5.7. Our primary goal is to extract the generalized Sivers shift, $\langle \boldsymbol{k}_y \rangle_{TU}$, as a continuous function of the longitudinal momentum fraction $x = k^+/P^+$. The Sivers shift is defined phenomenologically as the ratio of the first transverse momentum moment of the Sivers function to the unpolarized distribution as discussed in Section 4.6.

Because x and the longitudinal spatial separation $b \cdot P$ are Fourier conjugate variables, mapping the x-dependence requires a Fourier transform of these amplitudes with respect to $b \cdot P$. Analytically, the x-dependent

Sivers shift takes the form:

$$\langle \boldsymbol{k}_y \rangle_{TU}(\boldsymbol{b}_{\rm T}^2, x, \hat{\zeta}, \eta v{\cdot}P) \;\equiv\; m_N \frac{\tilde{f}_{1T}^{\perp(1)}(\boldsymbol{b}_{\rm T}^2, x; \hat{\zeta}, \ldots, \eta v{\cdot}P)}{\tilde{f}_1^{(0)}(\boldsymbol{b}_{\rm T}^2, x; \hat{\zeta}, \ldots, \eta v{\cdot}P)} \tag{5.1}$$

$$= -m_N \frac{\int d(b\cdot P) e^{ix(b\cdot P)} \widetilde{A}_{12B}(b^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P)}{\int d(b\cdot P) e^{ix(b\cdot P)} \widetilde{A}_{2B}(b^2, b{\cdot}P, (b{\cdot}P)R(\hat{\zeta}^2)/m_N^2, -1/(m_N\hat{\zeta})^2, \eta v{\cdot}P)} \tag{5.2}$$

$$= -m_N \frac{\int d(b\cdot P) e^{ix(b\cdot P)} \left[\widetilde{A}_{12B}^{\rm Re}(\cdots) + i\widetilde{A}_{12B}^{\rm Im}(\cdots)\right]}{\int d(b\cdot P) e^{ix(b\cdot P)} \left[\widetilde{A}_{2B}^{\rm Re}(\cdots) + i\widetilde{A}_{2B}^{\rm Im}(\cdots)\right]} \tag{5.3}$$

$$= -m_N \frac{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right) + i\sin\left(x(b\cdot P)\right)\right] \left[\widetilde{A}_{12B}^{\rm Re}(\cdots) + i\widetilde{A}_{12B}^{\rm Im}(\cdots)\right]}{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right) + i\sin\left(x(b\cdot P)\right)\right] \left[\widetilde{A}_{2B}^{\rm Re}(\cdots) + i\widetilde{A}_{2B}^{\rm Im}(\cdots)\right]} \tag{5.4}$$

$$= -m_N \frac{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right)\widetilde{A}_{12B}^{\rm Re}(\cdots) - \sin\left(x(b\cdot P)\right)\widetilde{A}_{12B}^{\rm Im}(\cdots)\right]}{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right)\widetilde{A}_{2B}^{\rm Re}(\cdots) - \sin\left(x(b\cdot P)\right)\widetilde{A}_{2B}^{\rm Im}(\cdots)\right]} \tag{5.5}$$

$$= -m_N \frac{\left[\widetilde{A}_{12B}^{[x]\rm Re}(b^2, x, \hat{\zeta}, \eta v{\cdot}P) - \widetilde{A}_{12B}^{[x]\rm Im}(b^2, x, \hat{\zeta}, \eta v{\cdot}P)\right]}{\left[\widetilde{A}_{2B}^{[x]\rm Re}(b^2, x, \hat{\zeta}, \eta v{\cdot}P) - \widetilde{A}_{2B}^{[x]\rm Im}(b^2, x, \hat{\zeta}, \eta v{\cdot}P)\right]} \tag{5.6}$$

Expanding the Fourier phase factor into its trigonometric components isolates the real and imaginary contributions of the amplitudes to the Sivers shift. Because the integration over $d(b\cdot P)$ is performed over a symmetric domain, the non-vanishing terms of the integral are dictated by the parity of the integrands with respect to $b\cdot P$. The trigonometric weights $\cos\left(x(b\cdot P)\right)$ and $\sin\left(x(b\cdot P)\right)$ are even and odd, respectively.

This harmonic symmetry filters the invariant amplitudes based on their behavior under Hermitian conjugation. By Eqs. (4.25) and (4.28), the real components $\widetilde{A}_{2B}^{\rm Re}$ and $\widetilde{A}_{12B}^{\rm Re}$ are even functions of $b\cdot P$, while the imaginary components $\widetilde{A}_{2B}^{\rm Im}$ and $\widetilde{A}_{12B}^{\rm Im}$ are odd. (Parity in b is parity in $b\cdot P$ here, because b^2 is unchanged under $b \to -b$, $v\cdot b$ is tied to $b\cdot P$ by Eq. (4.32), and $R(\hat{\zeta}^2)$ is real.) Only the symmetric products therefore survive the integration, i.e., the even real amplitudes coupled with the cosine term and the odd imaginary amplitudes coupled with the sine term. Here, $\widetilde{A}_{iB}^{[x]\rm Re}$ and $\widetilde{A}_{iB}^{[x]\rm Im}$ are the Fourier-transformed amplitudes. Note that, since $\widetilde{A}_{iB}^{\rm Im}$ is odd in $(P\cdot b)$, its Fourier transform gives an imaginary solution, i. e. $\int \frac{d(b\cdot P)}{2\pi} e^{ix(b\cdot P)} \widetilde{A}_{iB}^{\rm Im} = i\widetilde{A}_{iB}^{[x]\rm Im}$, this results in

$$\int \frac{d(b\cdot P)}{2\pi} e^{ix(b\cdot P)} \left[\widetilde{A}_{iB}^{\rm Re} + i\widetilde{A}_{iB}^{\rm Im}\right] = \left[\widetilde{A}_{iB}^{[x]\rm Re} + i^2\widetilde{A}_{iB}^{[x]\rm Im}\right] \tag{5.7}$$

$$= \left[\widetilde{A}_{iB}^{[x]\rm Re} - \widetilde{A}_{iB}^{[x]\rm Im}\right] \tag{5.8}$$

for both unpolarized and Sivers TMD amplitudes.

Next, we extract these parity components from the discrete lattice data; Sections 5.5 and 5.6 describe how we build correlation functions to separate them.

5.2 The Collins-Soper Parameter ($\hat{\zeta}$)

We defined the Collins-Soper parameter $\hat{\zeta}$ in the previous Chapter. 4 (Eq. (4.12)):

$$\hat{\zeta} \equiv \zeta/2m_N = \frac{v \cdot P}{\sqrt{|v^2|}\sqrt{P^2}}, \tag{5.9}$$

On the lattice, the staple direction v must be space-like ($v^2 < 0$), so the Collins-Soper parameter $\hat{\zeta}$ and the physical TMD limit are recovered only by extrapolating $\hat{\zeta} \to \infty$. In practice, since v is a purely spatial vector on the lattice ($v_0 = 0$), increasing $\hat{\zeta}$ requires simulating the nucleon at progressively higher spatial momenta $\boldsymbol{P}$. Tracking the evolution of the Sivers shift across multiple $\hat{\zeta}$ values is therefore necessary for phenomenological predictions.

5.3 Kinematic Constraints to scan full $b \cdot P$

To perform the Fourier transform over $b \cdot P$, we require a lattice geometry with both non-zero $b \cdot P$ and zero $b \cdot P$. However, the Lorentz-invariant arguments of the amplitudes are not independent. Our chosen frame ($b^+ = 0, \boldsymbol{P}_{\mathrm{T}} = 0$) enforces the constraint of Eq. (4.32). Expanding the dot products in a frame where the momentum is directed along the 1-axis ($P = (E, P_1, 0, 0)$), we find:

$$b_1|v| \left[\frac{P_1}{m_N} \left(\hat{\zeta} - \sqrt{\hat{\zeta}^2 + 1} \right) + \hat{\zeta} \frac{m_N}{P_1} \right] = \vec{v}_T \cdot \vec{b}_T \ . \tag{5.10}$$

To satisfy this equation over a continuous range of $b \cdot P$ without coupling the transverse separation $\vec{b}_T$ directly to the longitudinal staple geometry, we orient the transverse separation vectors orthogonally. By choosing $\vec{v}_T = (0, v_3)$ and $\vec{b}_T = (b_2, 0)$, the right-hand side identically vanishes ($\vec{v}_T \cdot \vec{b}_T = 0$).

5.3.1 Explicit Calculation for Hadron Momentum $P_1 = -1$

Setting the right-hand side of Eq. (5.10) to zero implies that the bracketed term $\left[\frac{P_1}{m_N} \left(\hat{\zeta} - \sqrt{\hat{\zeta}^2 + 1} \right) + \hat{\zeta} \frac{m_N}{P_1} \right]$ must vanish. This fixes the required Collins-Soper parameter, up to a sign, for a given lattice momentum. Using the lattice ensemble parameters from our lattice size of $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit of $a = 0.11403$ fm,

the nucleon mass and the lowest momentum mode are:

$$m_N = \frac{0.6228(28)}{0.11403 \text{ fm}} = \frac{0.6228(28) \times 197.32698 \text{ MeV}}{0.11403} = 1077.8(48) \text{ MeV} , \tag{5.11}$$

$$P_1 = -\frac{2\pi}{L} = -\frac{2\pi}{32 \times 0.11403 \text{ fm}} = -339.8 \text{ MeV} . \tag{5.12}$$

Substituting these values into the constraint equation yields:

$$\begin{aligned} \implies \left[\frac{P_1}{m_N}\left(\hat{\zeta} - \sqrt{\hat{\zeta}^2 + 1}\right) + \hat{\zeta}\frac{m_N}{P_1}\right] = 0 \\ \implies \hat{\zeta} = \sqrt{\frac{P_1^2}{m_N^2\left(\frac{m_N^2}{P_1^2} + 2\right)}} \\ \hat{\zeta} = \pm 0.09077(75) \\ \boxed{\hat{\zeta} = \pm 0.090772} . \end{aligned} \tag{5.13}$$

With $\hat{\zeta}$ determined, we can derive the specific orientation of the space-like staple vector v. From the definition of $\hat{\zeta}$:

$$\hat{\zeta} = \frac{-v_1 P_1}{|v| m_N} = \frac{-v_1 P_1}{\left(\sqrt{v_1^2 + 0^2 + v_3^2}\right) m_N} . \tag{5.14}$$

Solving for the longitudinal staple component v_1 in terms of the transverse component v_3:

$$\implies \boxed{v_1 = \pm 0.301 \; v_3} . \tag{5.15}$$

5.4 Lattice Parameter Space and Off-Axis Links

This geometric calculation is repeated over a range of nucleon momenta to increase $\hat{\zeta}$ systematically. The resulting constraint parameters are summarized in Table 2.

$\boldsymbol{P} \cdot a\hat{L}/(2\pi)$	$\hat{\zeta}$	v_1
$(-1, 0, 0)$	± 0.090772	$v_1 = \pm 0.301 \; v_3$
$(-2, 0, 0)$	± 0.294367	$v_1 = \pm 0.531 \; v_3$
$(-3, 0, 0)$	± 0.539684	$v_1 = \pm 0.689 \; v_3$
$(-4, 0, 0)$	± 0.785756	$v_1 = \pm 0.786 \; v_3$

Table 2: Resulting Collins-Soper parameters ($\hat{\zeta}$) and the necessary geometric constraint on the staple vector components to ensure scan full $b \cdot P$.

$\boldsymbol{b}/a$	$\eta\boldsymbol{v}/a$	$\boldsymbol{P}\cdot a\hat{L}/(2\pi)$
$(b_1, b_2, 0)$	$\eta\cdot(v_1, 0, v_3)$, $\eta = 0, \cdots, 10$	$(n, 0, 0)$, $n = -1, -2, -3, -4$

Table 3: Configuration parameters for the Lattice QCD non-local matrix element extraction. Our lattice size is of $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is of $a = 0.11403$ fm

A consequence of Table 2 is that the staple vector v cannot align cleanly with the Cartesian axes of the Euclidean grid. Instead, it must point in an off-axis direction within the $v_1 - v_3$ plane. For example, to achieve $v_1 = 0.301\ v_3$ at $|P| = 1$, the Wilson line must trace an oblique, step-wise path across the discrete lattice sites (e.g., traversing roughly three lattice spacings in the 1-direction for every ten in the 3-direction). Generating these off-axis Wilson lines without injecting excessive discretization noise guides our strategies for link smearing and gauge path selection. Figure 7 plots the resulting staple directions in the $v_1 - v_3$ plane for all four Collins-Soper values in Table 2.



Figure 7: Directions of the spatial staple vectors $\boldsymbol{v}$ mapped in the $v_1 - v_3$ plane. Each vector corresponds to the requisite geometry to satisfy the $b \cdot P \neq 0$ constraint for the four sequential Collins-Soper parameters $\hat{\zeta}$.

Section 5.4.1 identifies the integer lattice sites that bound each off-axis target, and Section 5.4.2 averages over degenerate paths and interpolates to the target.

5.4.1 Off-Axis v and Spatial Interpolation

To evaluate the correlation function as a function of the staple extent, we scale the direction vector $\boldsymbol{v}$ by an increment η, and $\boldsymbol{v}$ is a unit vector. The physical length of the staple is thus governed by the coordinate $\eta\boldsymbol{v}/a$. Because $\boldsymbol{v}$ points in a continuous, off-axis direction, scaling it by integer values of η generally yields target coordinates that fall somewhere between the discrete Cartesian lattice sites.

To access the continuous off-axis coordinate $\eta\boldsymbol{v}/a$, we employ a spatial bounding and interpolation strategy. Rather than constructing a single, potentially biased step-wise path (a "staircase") that roughly approximates the vector, we compute the observable at multiple distinct integer lattice sites surrounding the exact ideal target coordinate. We reach these bounding coordinates through specific permutations of unit links. A path sequence denoted $\{1, 3, 1\}$ is a single link in the $\hat{x_1}$ direction, followed by one in the $\hat{x_3}$ direction and a final link in the $\hat{x_1}$ direction, which lands at the integer coordinate $(2, 0, 1)$. By varying these step sequences, we gather data at three to four adjacent lattice sites (e.g., on either side of the target along the x^3 slice).

Once these bounding matrix elements are computed, we perform a spatial interpolation inside the Jackknife resampling loop to extract the value at the continuous target coordinate $\eta\boldsymbol{v}/a$. The interpolation reduces discretization artifacts and the staircase errors of off-axis Wilson lines; the next section describes the procedure. Figures 8 through 15 show the ideal continuous trajectories for each nucleon momentum, together with the corresponding discrete bounding paths and link sequences.



Figure 8: Ideal continuous trajectory of the off-axis staple vector (solid blue) for $P_1 = -1$. Circular markers indicate target continuous coordinates for sequential values of η. Dashed lines illustrate the discrete, step-wise permutations of gauge links used to reach the surrounding bounding integer lattice sites.

$\eta\, \boldsymbol{v}/a$	Gauge-link paths for interpolation ($\hat{\zeta} = 0.090772$)
0·(0.301,0,1)	{{}}
1·(0.301,0,1)	{{3}, {1, 3}, {3, 1}, {1, 3, 1}}
2·(0.301,0,1)	{{3, 3}, {3, 1, 3}, {1, 3, 1, 3}, {3, 1, 3, 1}}
3·(0.301,0,1)	{{3, 3, 3}, {3, 1, 3, 3}, {3, 3, 1, 3}, {3, 1, 3, 1, 3}}
4·(0.301,0,1)	{{3, 3, 3, 3}, {3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3}}
5·(0.301,0,1)	{{3, 3, 1, 3, 3, 3}, {3, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 1, 3, 3}, {3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3}}
6·(0.301,0,1)	{{3, 3, 3, 1, 3, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 3, 1, 3}}
7·(0.301,0,1)	{{3, 3, 3, 1, 3, 3, 3, 3}, {3, 3, 3, 3, 1, 3, 3, 3}, {3, 3, 1, 3, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 1, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 1, 3, 3}}
8·(0.301,0,1)	{{3, 3, 3, 3, 1, 3, 3, 3, 3}, {3, 3, 3, 1, 3, 3, 1, 3, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3}}
9·(0.301,0,1)	{{3, 3, 3, 1, 3, 3, 3, 1, 3, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 3, 3, 1, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 3}}
10·(0.301,0,1)	{{3, 3, 3, 1, 3, 3, 3, 3, 1, 3, 3, 3}, {3, 3, 1, 3, 3, 3, 1, 3, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3}}

Figure 9: Discrete gauge link configurations defining the interpolation bounds for $P_1 = -1$. Empty brackets {} denote a zero-length origin link, while numerical arrays (e.g., $\{1, 3, 1\}$) give the sequence of directional steps.



Figure 10: Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -2$.

$\eta \cdot \nu / a$	Gauge-link paths for interpolation ($\hat{\zeta} = 0.294367$)
0·(0.531,0,1)	{{}}
1·(0.531,0,1)	{{3}, {1, 3}, {3, 1}, {1, 3, 1}}
2·(0.531,0,1)	{{3, 3}, {3, 1, 3}, {1, 3, 1, 3}, {3, 1, 3, 1}}
3·(0.531,0,1)	{{3, 1, 3, 3}, {3, 3, 1, 3}, {3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1}}
4·(0.531,0,1)	{{3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3}}
5·(0.531,0,1)	{{3, 3, 1, 3, 1, 3, 3}, {3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3}}
6·(0.531,0,1)	{{3, 3, 1, 3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3}}
7·(0.531,0,1)	{{3, 3, 1, 3, 1, 3, 3, 1, 3, 3}, {3, 3, 1, 3, 3, 1, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}}
8·(0.531,0,1)	{{3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3}}
9·(0.531,0,1)	{{3, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3}, {3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}}
10·(0.531,0,1)	{{3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3}, {3, 1, 3, 3, 1, 3, 3, 1, 3, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}}

Figure 11: Discrete gauge link configurations defining the interpolation bounds for $P_1 = -2$.



Figure 12: Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -3$.

$\eta\, \boldsymbol{\nu} / a$	Gauge-link paths for interpolation ($\hat{\zeta} = 0.539684$)
0·(0.689,0,1)	{{}}
1·(0.689,0,1)	{{3}, {1, 3}, {3, 1}, {1, 3, 1}}
2·(0.689,0,1)	{{3, 3}, {3, 1, 3}, {1, 3, 1, 3}, {3, 1, 3, 1}}
3·(0.689,0,1)	{{3, 1, 3, 3}, {3, 3, 1, 3}, {3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1}}
4·(0.689,0,1)	{{3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1}}
5·(0.689,0,1)	{{3, 3, 1, 3, 1, 3, 3}, {3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3}}
6·(0.689,0,1)	{{3, 1, 3, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
7·(0.689,0,1)	{{3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
8·(0.689,0,1)	{{3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
9·(0.689,0,1)	{{3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3}, {3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
10·(0.689,0,1)	{{3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 1, 3}}

Figure 13: Discrete gauge link configurations defining the interpolation bounds for $P_1 = -3$.



Figure 14: Ideal continuous trajectory and corresponding discrete bounding paths for the off-axis staple vector at $P_1 = -4$.

$\eta \cdot \mathbf{v} / a$	Gauge-link paths for interpolation ($\hat{\zeta} = 0.785756$)
0·(0.786,0,1)	{{}}
1·(0.786,0,1)	{{3}, {1, 3}, {3, 1}, {1, 3, 1}}
2·(0.786,0,1)	{{3, 1, 3}, {1, 3, 1, 3}, {3, 1, 3, 1}, {1, 3, 1, 3, 1}}
3·(0.786,0,1)	{{3, 1, 3, 3}, {3, 3, 1, 3}, {3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1}}
4·(0.786,0,1)	{{3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1}}
5·(0.786,0,1)	{{3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1}}
6·(0.786,0,1)	{{3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1}}
7·(0.786,0,1)	{{3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1}}
8·(0.786,0,1)	{{3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
9·(0.786,0,1)	{{3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}}
10·(0.786,0,1)	{{3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 3, 1, 3, 1, 3, 1, 3, 1, 3}, {3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3, 1, 3}}

Figure 15: Discrete gauge link configurations defining the interpolation bounds for $P_1 = -4$.

5.4.2 Spatial Interpolation of Off-Axis Coordinates

Before interpolating to the off-axis target $\eta \boldsymbol{v}/a$, we must account for path degeneracy. Because the lattice gauge links are restricted to the Cartesian axes, there are multiple step-wise permutations (different side-link paths) that terminate at the same physical lattice coordinate. While these paths are geometrically equivalent at their endpoints, they correspond to different local gauge field configurations and thus exhibit distinct discretization artifacts and gauge noise. As illustrated in Figure 16, to improve the symmetry, we average the correlation functions evaluated over all degenerate paths that share the same endpoint. Path averaging significantly improves the signal-to-noise ratio before any interpolation.



Figure 16: Illustration of path degeneracy. Two distinct step-wise gauge link permutations (e.g., solid and dashed lines) terminate at the identical discrete lattice coordinate. We average the correlators from these equivalent paths to suppress local gauge noise before interpolation.

Once we obtain the symmetry-improved (path-averaged) correlators at the discrete bounding sites, we perform a 1-dimensional interpolation. For a fixed transverse staple extent v_3, the kinematic constraint dictates an ideal continuous longitudinal component, $v_1^{\text{constraint}}$. The surrounding discrete lattice sites provide correlators at actual simulated coordinates, v_1^{lattice}. We define the fractional deviation from the ideal path as:

$$\Delta v_1 = \frac{v_1^{\text{constraint}} - v_1^{\text{lattice}}}{a} \, . \tag{5.16}$$

By calculating the observable at three or more bounding lattice sites, we generate a set of data points as a

function of this deviation Δv_1. Within each Jackknife resampling bin (to preserve the covariance and statistical error), we perform a quadratic interpolation over these points. We then extract the physical amplitude on the continuous constraint trajectory by evaluating the interpolating polynomial exactly at $\Delta v_1 = 0$.

The core Mathematica logic executed inside the data analysis pipeline to perform this quadratic interpolation is shown in the snippet below:

Listing 2: Core interpolation logic for evaluating the constrained correlator.

```
(* Pair fractional deviation (x) with path-averaged data (y) *)
plotforinterpolation = Transpose[{v1dataforInterpolation,
                                  dataforInterpolation}];

(* Construct quadratic interpolation over bounding points *)
fninterpol = Interpolation[plotforinterpolation,
                           InterpolationOrder -> 2];

(* Evaluate exactly on the constraint trajectory (deviation = 0) *)
interpol0 = {{0, fninterpol[0]}};
```

A representative example of this interpolation procedure is shown in Figure 17 for a specific staple extent ($\eta = 7$) and nucleon momentum ($P_1 = -1$). The red data points represent the path-averaged lattice evaluations at the discrete bounding sites, plotted against their deviation from the ideal trajectory. The solid gray curve denotes the quadratic interpolation, and the blue point highlights the final extracted physical value at $\Delta v_1 = 0$. We apply this procedure to all Dirac structures (Γ), separately for the real ($b \cdot P$-even) and imaginary ($b \cdot P$-odd) parts of the amplitudes.



Figure 17: Quadratic interpolation of the real and imaginary parts of the correlation function for $\boldsymbol{b} = \{1, 2, 0\}$, $\eta = 7$, and $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = (-1, 0, 0)$. Red markers indicate the path-averaged lattice data points as a function of their deviation from the ideal constraint trajectory, $\Delta v_1 = (v_1^{\text{constraint}} - v_1^{\text{lattice}})/a$. The gray band indicates the quadratic fit ('InterpolationOrder $\rightarrow$ 2'), and the central blue marker represents the final extracted value at $\Delta v_1 = 0$, lying exactly on the required off-axis spatial trajectory.

5.5 Quark Separation Vectors (b) and Amplitude Parity

In addition to the staple vector $\boldsymbol{v}$, the correlation function depends on the spatial separation $\boldsymbol{b}$ between the initial and final quark fields. Recall that the real and imaginary parts of the invariant amplitudes ($\widetilde{A}_{iB}^{\mathrm{Re}}$ and $\widetilde{A}_{iB}^{\mathrm{Im}}$) are even and odd functions of the invariant dot product $b \cdot P$, respectively.

Because the nucleon momentum in our lattice setup is fixed along the $\hat{x_1}$-axis ($P = P_1\hat{x_1}$), the longitudinal separation is directly proportional to the $\hat{x_1}$-component of the separation vector: $b \cdot P = -P_1 b_1$. Therefore, to isolate the real and imaginary parts of the amplitudes, we must evaluate the correlator at both positive and negative values of b_1. By taking the symmetric (even) and antisymmetric (odd) linear combinations of the correlators evaluated at $+b_1$ and $-b_1$, we project out the real and imaginary amplitudes required for the x-dependent Fourier transform. The numerical Fourier transform over $b \cdot P$ requires a wide and dense grid of the separation vector $\boldsymbol{b}$. Figure 20 shows the parameter space of our calculations. The left panel shows the discrete Cartesian grid in the $b_1 - b_2$ plane. The right panel shows how this grid maps onto the phenomenologically relevant invariant Variable Space spanned by $(b \cdot P)\hat{L}/(2\pi)$ and the absolute separation $|b|/a$; the mapping shifts with the target Collins-Soper parameter $\hat{\zeta}$. Where, $\hat{L}$ is the spatial lattice length $\hat{L} = 32$, and $a = 0.11403$ fm is lattice unit.



Figure 18: The sampled operator separation vectors mapped in the spatial $b_1 - b_2$ plane (where $b_1 \equiv b_L$ and $b_2 \equiv b_T$).



Figure 19: Mapping of the spatial lattice vectors into the physical invariant Variable Space $(b \cdot P/(2\pi), |b|/a)$. The trajectories diverge based on the selected target $\hat{\zeta}$.

Figure 20: Coverage of the quark separation vectors across the simulated lattice volume and their corresponding physical Variable Space representation. Our lattice size is of $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is of $a = 0.11403$ fm

Section 5.5.1 lists the gauge link paths used to reach each $\boldsymbol{b}$, and Section 5.5.2 averages over degenerate paths.

5.5.1 Gauge Link Construction for $\boldsymbol{b}$

Constructing the direct Wilson line connecting the quark fields at 0 and $\boldsymbol{b}$ requires generating step-wise paths across the lattice. We categorize these paths into purely transverse separations ($b_1 = 0$) and off-axis separations with a longitudinal component ($b_1 \neq 0$). For purely transverse separations, where $b \cdot P = 0$, the gauge links lie entirely within the transverse plane. Figure 21 lists the link sequences used to reach these target coordinates. An empty bracket {} denotes the local origin, while arrays such as $\{2, 2\}$ indicate a sequence of two consecutive links in the $\hat{x_2}$-direction.

$\{+b_1, b_3, b_3\} / a$	Gauge-link paths	$\{-b_1, b_3, b_3\} / a$	Gauge-link paths
{0, 0, 0}	{{}}	{0, 0, 0}	{{}}
{0, 1, 0}	{{2}}	{0, 1, 0}	{{2}}
{0, 2, 0}	{{2, 2}}	{0, 2, 0}	{{2, 2}}
{0, 3, 0}	{{2, 2, 2}}	{0, 3, 0}	{{2, 2, 2}}
{0, 4, 0}	{{2, 2, 2, 2}}	{0, 4, 0}	{{2, 2, 2, 2}}
{0, 5, 0}	{{2, 2, 2, 2, 2}}	{0, 5, 0}	{{2, 2, 2, 2, 2}}
{0, 6, 0}	{{2, 2, 2, 2, 2, 2}}	{0, 6, 0}	{{2, 2, 2, 2, 2, 2}}
{0, 7, 0}	{{2, 2, 2, 2, 2, 2, 2}}	{0, 7, 0}	{{2, 2, 2, 2, 2, 2, 2}}
{0, -1, 0}	{{-2}}	{0, -1, 0}	{{-2}}
{0, -2, 0}	{{-2, -2}}	{0, -2, 0}	{{-2, -2}}
{0, -3, 0}	{{-2, -2, -2}}	{0, -3, 0}	{{-2, -2, -2}}
{0, -4, 0}	{{-2, -2, -2, -2}}	{0, -4, 0}	{{-2, -2, -2, -2}}
{0, -5, 0}	{{-2, -2, -2, -2, -2}}	{0, -5, 0}	{{-2, -2, -2, -2, -2}}
{0, -6, 0}	{{-2, -2, -2, -2, -2, -2}}	{0, -6, 0}	{{-2, -2, -2, -2, -2, -2}}
{0, -7, 0}	{{-2, -2, -2, -2, -2, -2}}	{0, -7, 0}	{{-2, -2, -2, -2, -2, -2}}

Figure 21: Discrete gauge link configurations for purely transverse quark separations ($\boldsymbol{b} = \{0, b_2, 0\}$). The numerical arrays show the sequential lattice steps (e.g., '2' represents a shift in the x_2 direction). Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm.

To access finite $b \cdot P$, we construct off-axis paths with simultaneous steps in the longitudinal ($\hat{x_1}$) and transverse ($\hat{x_2}$) directions. To preserve the symmetry required for isolating parity states, every path that reaches a $+b_1$ coordinate has an exact mirror path reaching $-b_1$. Figures 22-23 show these mirrored sequences side by side. Sequences such as $\{2, 1\}$ indicate an alternating "staircase" step: one unit in the $\hat{x_2}$-direction followed by one unit in the $\hat{x_1}$-direction.

$\{+b_1,\ b_3,\ b_3\}\ /\ a$	Gauge-link paths	$\{-b_1,\ b_3,\ b_3\}\ /\ a$	Gauge-link paths
{1, 1, 0}	{{2, 1}, {1, 2}}	{-1, 1, 0}	{{2, -1}, {-1, 2}}
{2, 2, 0}	{{2, 1, 2, 1}, {1, 2, 1, 2}}	{-2, 2, 0}	{{2, -1, 2, -1}, {-1, 2, -1, 2}}
{3, 3, 0}	{{2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2}}	{-3, 3, 0}	{{2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2}}
{4, 4, 0}	{{2, 1, 2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2, 1, 2}}	{-4, 4, 0}	{{2, -1, 2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2, -1, 2}}
{5, 5, 0}	{{2, 1, 2, 1, 2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2, 1, 2, 1, 2}}	{-5, 5, 0}	{{2, -1, 2, -1, 2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2, -1, 2, -1, 2}}
{1, 2, 0}	{{2, 1, 2}}	{-1, 2, 0}	{{2, -1, 2}}
{2, 4, 0}	{{2, 1, 2, 2, 1, 2}}	{-2, 4, 0}	{{2, -1, 2, 2, -1, 2}}
{3, 6, 0}	{{2, 1, 2, 2, 1, 2, 2, 1, 2}}	{-3, 6, 0}	{{2, -1, 2, 2, -1, 2, 2, -1, 2}}
{2, 1, 0}	{{1, 2, 1}}	{-2, 1, 0}	{{-1, 2, -1}}
{4, 2, 0}	{{1, 2, 1, 1, 2, 1}}	{-4, 2, 0}	{{-1, 2, -1, -1, 2, -1}}
{6, 3, 0}	{{1, 2, 1, 1, 2, 1, 1, 2, 1}}	{-6, 3, 0}	{{-1, 2, -1, -1, 2, -1, -1, 2, -1}}
{4, 1, 0}	{{1, 1, 2, 1, 1}}	{-4, 1, 0}	{{-1, -1, 2, -1, -1}}
{8, 2, 0}	{{1, 1, 2, 1, 1, 1, 1, 2, 1, 1}}	{-8, 2, 0}	{{-1, -1, 2, -1, -1, -1, -1, 2, -1, -1}}
{1, -1, 0}	{{-2, 1}, {1, -2}}	{-1, -1, 0}	{{-2, -1}, {-1, -2}}
{2, -2, 0}	{{-2, 1, -2, 1}, {1, -2, 1, -2}}	{-2, -2, 0}	{{-2, -1, -2, -1}, {-1, -2, -1, -2}}
{3, -3, 0}	{{-2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2}}	{-3, -3, 0}	{{-2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2}}
{4, -4, 0}	{{-2, 1, -2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2, 1, -2}}	{-4, -4, 0}	{{-2, -1, -2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2, -1, -2}}
{5, -5, 0}	{{-2, 1, -2, 1, -2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2, 1, -2, 1, -2}}	{-5, -5, 0}	{{-2, -1, -2, -1, -2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2, -1, -2, -1, -2}}
{1, -2, 0}	{{-2, 1, -2}}	{-1, -2, 0}	{{-2, -1, -2}}
{2, -4, 0}	{{-2, 1, -2, -2, 1, -2}}	{-2, -4, 0}	{{-2, -1, -2, -2, -1, -2}}
{3, -6, 0}	{{-2, 1, -2, -2, 1, -2, -2, 1, -2}}	{-3, -6, 0}	{{-2, -1, -2, -2, -1, -2, -2, -1, -2}}
{2, -1, 0}	{{1, -2, 1}}	{-2, -1, 0}	{{-1, -2, -1}}
{4, -2, 0}	{{1, -2, 1, 1, -2, 1}}	{-4, -2, 0}	{{-1, -2, -1, -1, -2, -1}}
{6, -3, 0}	{{1, -2, 1, 1, -2, 1, 1, -2, 1}}	{-6, -3, 0}	{{-1, -2, -1, -1, -2, -1, -1, -2, -1}}
{4, -1, 0}	{{1, 1, -2, 1, 1}}	{-4, -1, 0}	{{-1, -1, -2, -1, -1}}
{8, -2, 0}	{{1, 1, -2, 1, 1, 1, 1, -2, 1, 1}}	{-8, -2, 0}	{{-1, -1, -2, -1, -1, -1, -1, -2, -1, -1}}

Figure 22: Gauge link sequences for $+b_1$ (positive longitudinal separation). Symmetric sets of step-wise gauge link paths required to access finite $\pm b \cdot P$ and isolate the real and imaginary components of the invariant amplitudes. Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm.

$\{+b_1, b_3, b_3\}/a$	Gauge-link paths	$\{-b_1, b_3, b_3\}/a$	Gauge-link paths
{-1, 1, 0}	{{2, -1}, {-1, 2}}	{1, 1, 0}	{{2, 1}, {1, 2}}
{-2, 2, 0}	{{2, -1, 2, -1}, {-1, 2, -1, 2}}	{2, 2, 0}	{{2, 1, 2, 1}, {1, 2, 1, 2}}
{-3, 3, 0}	{{2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2}}	{3, 3, 0}	{{2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2}}
{-4, 4, 0}	{{2, -1, 2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2, -1, 2}}	{4, 4, 0}	{{2, 1, 2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2, 1, 2}}
{-5, 5, 0}	{{2, -1, 2, -1, 2, -1, 2, -1, 2, -1}, {-1, 2, -1, 2, -1, 2, -1, 2, -1, 2}}	{5, 5, 0}	{{2, 1, 2, 1, 2, 1, 2, 1, 2, 1}, {1, 2, 1, 2, 1, 2, 1, 2, 1, 2}}
{-1, 2, 0}	{{2, -1, 2}}	{1, 2, 0}	{{2, 1, 2}}
{-2, 4, 0}	{{2, -1, 2, 2, -1, 2}}	{2, 4, 0}	{{2, 1, 2, 2, 1, 2}}
{-3, 6, 0}	{{2, -1, 2, 2, -1, 2, 2, -1, 2}}	{3, 6, 0}	{{2, 1, 2, 2, 1, 2, 2, 1, 2}}
{-2, 1, 0}	{{-1, 2, -1}}	{2, 1, 0}	{{1, 2, 1}}
{-4, 2, 0}	{{-1, 2, -1, -1, 2, -1}}	{4, 2, 0}	{{1, 2, 1, 1, 2, 1}}
{-6, 3, 0}	{{-1, 2, -1, -1, 2, -1, -1, 2, -1}}	{6, 3, 0}	{{1, 2, 1, 1, 2, 1, 1, 2, 1}}
{-4, 1, 0}	{{-1, -1, 2, -1, -1}}	{4, 1, 0}	{{1, 1, 2, 1, 1}}
{-8, 2, 0}	{{-1, -1, 2, -1, -1, -1, -1, 2, -1, -1}}	{8, 2, 0}	{{1, 1, 2, 1, 1, 1, 1, 2, 1, 1}}
{-1, -1, 0}	{{-2, -1}, {-1, -2}}	{1, -1, 0}	{{-2, 1}, {1, -2}}
{-2, -2, 0}	{{-2, -1, -2, -1}, {-1, -2, -1, -2}}	{2, -2, 0}	{{-2, 1, -2, 1}, {1, -2, 1, -2}}
{-3, -3, 0}	{{-2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2}}	{3, -3, 0}	{{-2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2}}
{-4, -4, 0}	{{-2, -1, -2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2, -1, -2}}	{4, -4, 0}	{{-2, 1, -2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2, 1, -2}}
{-5, -5, 0}	{{-2, -1, -2, -1, -2, -1, -2, -1, -2, -1}, {-1, -2, -1, -2, -1, -2, -1, -2, -1, -2}}	{5, -5, 0}	{{-2, 1, -2, 1, -2, 1, -2, 1, -2, 1}, {1, -2, 1, -2, 1, -2, 1, -2, 1, -2}}
{-1, -2, 0}	{{-2, -1, -2}}	{1, -2, 0}	{{-2, 1, -2}}
{-2, -4, 0}	{{-2, -1, -2, -2, -1, -2}}	{2, -4, 0}	{{-2, 1, -2, -2, 1, -2}}
{-3, -6, 0}	{{-2, -1, -2, -2, -1, -2, -2, -1, -2}}	{3, -6, 0}	{{-2, 1, -2, -2, 1, -2, -2, 1, -2}}
{-2, -1, 0}	{{-1, -2, -1}}	{2, -1, 0}	{{1, -2, 1}}
{-4, -2, 0}	{{-1, -2, -1, -1, -2, -1}}	{4, -2, 0}	{{1, -2, 1, 1, -2, 1}}
{-6, -3, 0}	{{-1, -2, -1, -1, -2, -1, -1, -2, -1}}	{6, -3, 0}	{{1, -2, 1, 1, -2, 1, 1, -2, 1}}
{-4, -1, 0}	{{-1, -1, -2, -1, -1}}	{4, -1, 0}	{{1, 1, -2, 1, 1}}
{-8, -2, 0}	{{-1, -1, -2, -1, -1, -1, -1, -2, -1, -1}}	{8, -2, 0}	{{1, 1, -2, 1, 1, 1, 1, -2, 1, 1}}

Figure 23: Mirrored gauge link sequences for $-b_1$ (negative longitudinal separation).Symmetric sets of step-wise gauge link paths required to access finite $\pm b \cdot P$ and isolate the real and imaginary components of the invariant amplitudes. Where the lattice size is $\{\hat{L}^3 \times \hat{T}\} = \{32^3 \times 96\}$ and lattice unit is $a = 0.11403$ fm.

5.5.2 Symmetry Improving Operation (Path Averaging) for $\boldsymbol{b}$

Several quark separation vectors $\boldsymbol{b}$ have the same path degeneracy as the staple direction vector $\boldsymbol{v}$ (Section 5.4.2). We treat it the same way: we average the correlation function over all equivalent step-wise paths that share the same origin and endpoint, as illustrated for a single coordinate in Figure 24.



Figure 24: Illustration of path degeneracy for the quark separation vector $\boldsymbol{b}$. Distinct step-wise permutations (solid and dashed lines) terminating at the same Cartesian coordinate are averaged to suppress local gauge noise.

Applying this symmetry-improving operation (path-averaging) strategy across the entire target grid yields the full network of smoothed off-axis trajectories, visualized in Figure 25.



Figure 25: The full network of path-averaged $\boldsymbol{b}$ vectors in the $b_1/a - b_2/a$ plane. Solid red lines represent the ideal vectors, while dashed lines illustrate the underlying degenerate staircase paths calculated and averaged on the discrete lattice.

From this path-averaged dataset, we obtain the parity-separated amplitudes used in the final Fourier transform.

5.6 Solving for Real and Imaginary Amplitudes $\widetilde{A}_{iB}^{\mathbf{Re}}$ and $\widetilde{A}_{iB}^{\mathbf{Im}}$ from correlator $\widetilde{\Phi}^{[\gamma^\mu]}_{\mathbf{unsubtr.}}$

To extract the physical Sivers shift from the lattice, we must isolate the specific invariant amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ from the raw correlator data. We start from the parametrization of the unsubtracted correlator in Eq. (4.24) [1]. While this parametrization describes a continuous gauge link extending to infinity, we adopt an equivalent representation for lattice calculations in which the structures depend explicitly on the finite staple extent η. This form remains well defined even in the $v \cdot P = 0$ limit. We obtain it by making the substitution $v \to \eta v$ and extracting the scaling factors $(v{\cdot}P)^{-n}$ as discussed in detail in Chapter 4:

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^\mu]}_{\text{unsubtr.}} = \frac{1}{2}R_{\gamma^\mu}E_N = P^\mu\,\tilde{a}_2 - im_N^2 b^\mu\,\tilde{a}_3 - im_N\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha S_\beta\,\tilde{a}_{12} \\ + m_N^2\eta v^\mu\,\tilde{b}_1 + m_N\epsilon^{\mu\nu\alpha\beta}P_\nu\eta v_\alpha S_\beta\,\tilde{b}_7 - im_N^3\epsilon^{\mu\nu\alpha\beta}b_\nu\eta v_\alpha S_\beta\,\tilde{b}_8 \\ - m_N^3(b{\cdot}S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha\eta v_\beta\,\tilde{b}_9 - im_N^3(v{\cdot}S)\epsilon^{\mu\nu\alpha\beta}P_\nu b_\alpha\eta^2 v_\beta\tilde{b}_{10}\end{aligned} \tag{5.17}$$

To evaluate Eq. (4.24), we project it onto the kinematics chosen for our lattice setup. We choose a frame where the nucleon moves along the negative $\hat{1}$-axis, the spin is polarized along the $\hat{3}$-axis, and the separation and staple vectors lie in the transverse/longitudinal planes. We set $P^\mu = (E_N, -|\vec{P}^1|, 0, 0)$, $b^\mu = (0, b_1, b_2, 0)$, $v^\mu = \pm n'(0, v_1, 0, v_3)$, and $S^\mu = (0, 0, 0, 1)$. Expanding the sum over Lorentz indices for the temporal component ($\mu = 0$) and canceling the terms that vanish (such as $b^0 = 0$, $v^0 = 0$, and $b{\cdot}S = 0$), we find for each γ^μ:

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^0]}_{\text{unsubtr.}} &= P^0\,\widetilde{A}_2 \cancelto{0}{- im_N^2 b^0\,\widetilde{A}_3} + im_N\underbrace{\epsilon^{0123}}_{1}|\vec{P}^1|b_2S_3\,\widetilde{A}_{12} \\ &+ \cancelto{0}{\frac{m_N^2}{(v{\cdot}P)}v^0\,\widetilde{B}_1} + \cancelto{0}{\frac{m_N}{v{\cdot}P}\epsilon^{0113}P_1v_1S_3\,\widetilde{B}_7} + \cancelto{0}{\frac{m_N}{v{\cdot}P}\epsilon^{0133}P_1v_3S_3\,\widetilde{B}_7} \\ &- \frac{im_N^3}{v{\cdot}P}\underbrace{\epsilon^{0213}}_{-1}b_2v_1S_3\,\widetilde{B}_8 - \cancelto{0}{\frac{im_N^3}{v{\cdot}P}\epsilon^{0113}b_1v_1S_3\,\widetilde{B}_8} - \cancelto{0}{\frac{im_N^3}{v{\cdot}P}\epsilon^{0133}b_1v_3S_3\,\widetilde{B}_8} - \cancelto{0}{\frac{im_N^3}{v{\cdot}P}\epsilon^{0233}b_2v_3S_3\,\widetilde{B}_8} \\ &+ \cancelto{0}{\frac{m_N^3}{v{\cdot}P}(b{\cdot}S)\underbrace{\epsilon^{0123}}_{1}}|\vec{P}_1|b_2v_3\,\widetilde{B}_9 - \cancelto{0}{\frac{m_N^3}{v{\cdot}P}(b{\cdot}S)\epsilon^{0121}P_1b_2v_1\,\widetilde{B}_9} - \cancelto{0}{\frac{m_N^3}{v{\cdot}P}(b{\cdot}S)\epsilon^{0113}P_1b_1v_3\,\widetilde{B}_9} \\ &- \cancelto{0}{\frac{m_N^3}{v{\cdot}P}(b{\cdot}S)\epsilon^{0111}P_1b_1v_1\,\widetilde{B}_9} + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)\underbrace{\epsilon^{0123}}_{1}|\vec{P}_1|b_2v_3\widetilde{B}_{10} - \cancelto{0}{\frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)\epsilon^{0121}P_1b_2v_1\widetilde{B}_{10}} \\ &- \cancelto{0}{\frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)\epsilon^{0111}P_1b_1v_1\widetilde{B}_{10}} - \cancelto{0}{\frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)\epsilon^{0113}P_1b_1v_3\widetilde{B}_{10}}\end{aligned} \tag{5.18}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^0]}_{\text{unsubtr.}} = P^0\,\widetilde{A}_2 + im_N|\vec{P}^1|b_2S_3\,\widetilde{A}_{12} + \frac{im_N^3}{v{\cdot}P}b_2v_1S_3\,\widetilde{B}_8 + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)|P_1|b_2v_3\widetilde{B}_{10}} \tag{5.19}$$

Repeating this tensor contraction for the longitudinal spatial component ($\mu = 1$), taking care to track the signs of the Levi-Civita symbol, yields:

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^1]}_{\text{unsubtr.}} &= -|\vec{P}^1|\,\widetilde{A}_2 - im_N^2 b^1\,\widetilde{A}_3 - im_N \underbrace{\epsilon^{1023}}_{-1} P_0 b_2 S_3\,\widetilde{A}_{12} \\ &\quad + \frac{m_N^2}{(v\cdot P)} v^1\,\widetilde{B}_1 + \cancelto{0}{\frac{m_N}{v\cdot P}\epsilon^{1013} P_0 v_1 S_3\,\widetilde{B}_7} + \cancelto{0}{\frac{m_N}{v\cdot P}\epsilon^{1033} P_0 v_3 S_3\,\widetilde{B}_7} \\ &\quad - \cancelto{0}{\frac{im_N^3}{v\cdot P}\epsilon^{1213} b_2 v_1 S_3\,\widetilde{B}_8} - \cancelto{0}{\frac{im_N^3}{v\cdot P}\epsilon^{1233} b_2 v_3 S_3\,\widetilde{B}_8} - \cancelto{0}{\frac{im_N^3}{v\cdot P}\epsilon^{1133} b_1 v_3 S_3\,\widetilde{B}_8} - \cancelto{0}{\frac{im_N^3}{v\cdot P}\epsilon^{1113} b_1 v_1 S_3\,\widetilde{B}_8} \\ &\quad - \cancelto{0}{\frac{m_N^3}{v\cdot P}(b\cdot S)\underbrace{\epsilon^{1023}}_{1}} P_0 b_2 v_3\,\widetilde{B}_9 - \cancelto{0}{\frac{m_N^3}{v\cdot P}(b\cdot S)\epsilon^{1021} P_0 b_2 v_1}\,\widetilde{B}_9 - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S)\underbrace{\epsilon^{1023}}_{-1} P_0 b_2 v_3 \widetilde{B}_{10} \\ &\quad - \cancelto{0}{\frac{im_N^3}{(v\cdot P)^2}(v\cdot S)\epsilon^{1021} P_0 b_2 v_1}\widetilde{B}_{10} \end{aligned} \tag{5.20}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^1]}_{\text{unsubtr.}} = -|\vec{P}^1|\,\widetilde{A}_2 + \frac{m_N^2}{(v\cdot P)} v^1\,\widetilde{B}_1 + im_N P_0 b_2 S_3\,\widetilde{A}_{12} + \frac{im_N^3}{(v\cdot P)^2}(v\cdot S) P_0 b_2 v_3 \widetilde{B}_{10} - im_N^2 b^1\,\widetilde{A}_3} \tag{5.21}$$

Evaluating the transverse component ($\mu = 2$) isolates terms that depend on P_0 and the longitudinal separation b_1:

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^2]}_{\text{unsubtr.}} &= -im_N^2 b^2\,\widetilde{A}_3 - im_N \underbrace{\epsilon^{2013}}_{1} P_0 b_1 S_3\,\widetilde{A}_{12} \\ &\quad + \frac{m_N}{v\cdot P}\underbrace{\epsilon^{2013}}_{1} P_0 v_1 S_3\,\widetilde{B}_7 \\ &\quad - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S)\underbrace{\epsilon^{2013}}_{1} P_0 b_1 v_3 \widetilde{B}_{10} \end{aligned} \tag{5.22}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^2]}_{\text{unsubtr.}} = -im_N^2 b^2\,\widetilde{A}_3 - im_N P_0 b_1 S_3\,\widetilde{A}_{12} + \frac{m_N}{v\cdot P} P_0 v_1 S_3\,\widetilde{B}_7 - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S) P_0 b_1 v_3 \widetilde{B}_{10}} \tag{5.23}$$

Finally, the orthogonal transverse component ($\mu = 3$) simplifies because of the alignment of the spin vector:

$$\begin{aligned}\frac{1}{2}\widetilde{\Phi}^{[\gamma^3]}_{\text{unsubtr.}} &= \frac{m_N^2}{(v\cdot P)} v^3\,\widetilde{B}_1 \\ &\quad - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S)\underbrace{\epsilon^{3021}}_{1} P_0 b_2 v_1 \widetilde{B}_{10} \end{aligned} \tag{5.24}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^3]}_{\text{unsubtr.}} = \frac{m_N^2}{(v\cdot P)} v^3\,\widetilde{B}_1 - \frac{im_N^3}{(v\cdot P)^2}(v\cdot S) P_0 b_2 v_1 \widetilde{B}_{10}} \tag{5.25}$$

Collecting these boxed results gives the full system of equations mapping the infinite-link continuous amplitudes to the projected correlator components:

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^0]}_{\text{unsubtr.}} = P^0\,\widetilde{A}_2 + im_N|\vec{P}^1|b_2S_3\,\widetilde{A}_{12} + \frac{im_N^3}{v{\cdot}P}b_2v_1S_3\,\widetilde{B}_8 + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)|P_1|b_2v_3\widetilde{B}_{10}} \tag{5.26}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^1]}_{\text{unsubtr.}} = -|\vec{P}^1|\,\widetilde{A}_2 - im_N^2b^1\,\widetilde{A}_3 + \frac{m_N^2}{(v{\cdot}P)}v^1\,\widetilde{B}_1 + im_NP_0b_2S_3\,\widetilde{A}_{12} + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_2v_3\widetilde{B}_{10}} \tag{5.27}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^2]}_{\text{unsubtr.}} = -im_N^2b^2\,\widetilde{A}_3 - im_NP_0b_1S_3\,\widetilde{A}_{12} + \frac{m_N}{v{\cdot}P}P_0v_1S_3\,\widetilde{B}_7 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_1v_3\widetilde{B}_{10}} \tag{5.28}$$

$$\boxed{\frac{1}{2}\widetilde{\Phi}^{[\gamma^3]}_{\text{unsubtr.}} = \frac{m_N^2}{(v{\cdot}P)}v^3\,\widetilde{B}_1 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_2v_1\widetilde{B}_{10}} \tag{5.29}$$

We structure this system as a column vector to streamline the upcoming algebraic inversions:

$$\begin{pmatrix}\widetilde{\Phi}^{[\gamma^0]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^1]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^2]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^3]}_{\text{unsubtr.}}\end{pmatrix} = 2\begin{pmatrix} P^0\,\widetilde{A}_2 + im_N|\vec{P}^1|b_2S_3\,\widetilde{A}_{12} + \frac{im_N^3}{v{\cdot}P}b_2v_1S_3\,\widetilde{B}_8 + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)|P_1|b_2v_3\widetilde{B}_{10}\\ -|\vec{P}^1|\,\widetilde{A}_2 - im_N^2b^1\,\widetilde{A}_3 + \frac{m_N^2}{(v{\cdot}P)}v^1\,\widetilde{B}_1 + im_NP_0b_2S_3\,\widetilde{A}_{12} + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_2v_3\widetilde{B}_{10}\\ -im_N^2b^2\,\widetilde{A}_3 - im_NP_0b_1S_3\,\widetilde{A}_{12} + \frac{m_N}{v{\cdot}P}P_0v_1S_3\,\widetilde{B}_7 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_1v_3\widetilde{B}_{10}\\ \frac{m_N^2}{(v{\cdot}P)}v^3\,\widetilde{B}_1 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P_0b_2v_1\widetilde{B}_{10}\end{pmatrix} \tag{5.30}$$

To match standard contravariant Lorentz notation, we raise the spatial indices using the Minkowski metric $g^{\mu\nu} = \text{diag}(1,-1,-1,-1)$. This introduces a relative minus sign for the spatial components (e.g., $P_1 = -P^1$, $b_2 = -b^2$):

$$\begin{pmatrix}\widetilde{\Phi}^{[\gamma^0]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^1]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^2]}_{\text{unsubtr.}}\\ \widetilde{\Phi}^{[\gamma^3]}_{\text{unsubtr.}}\end{pmatrix} = 2\begin{pmatrix} P^0\,\widetilde{A}_2 - im_N|\vec{P}^1|b^2S^3\,\widetilde{A}_{12} - \frac{im_N^3}{v{\cdot}P}b^2v^1S^3\,\widetilde{B}_8 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)|\vec{P}^1|b^2v^3\widetilde{B}_{10}\\ -|\vec{P}^1|\,\widetilde{A}_2 - im_N^2b^1\,\widetilde{A}_3 + \frac{m_N^2}{(v{\cdot}P)}v^1\,\widetilde{B}_1 + im_NP^0b^2S^3\,\widetilde{A}_{12} + \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P^0b^2v^3\widetilde{B}_{10}\\ -im_N^2b^2\,\widetilde{A}_3 - im_NP^0b^1S^3\,\widetilde{A}_{12} + \frac{m_N}{v{\cdot}P}P^0v^1S^3\,\widetilde{B}_7 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P^0b^1v^3\widetilde{B}_{10}\\ \frac{m_N^2}{(v{\cdot}P)}v^3\,\widetilde{B}_1 - \frac{im_N^3}{(v{\cdot}P)^2}(v{\cdot}S)P^0b^2v^1\widetilde{B}_{10}\end{pmatrix} \tag{5.31}$$

To connect these continuous amplitudes to the discretized finite-link observables simulated on the lattice, we use the mapping of Eq. (4.31) between the infinite-link amplitudes ($\widetilde{A}_i$, $\widetilde{B}_i$) and the finite-link amplitudes ($\tilde{a}_i$, $\tilde{b}_i$). The finite-link amplitudes are complex-valued functions, which we decompose into their real and imaginary components:

$$\tilde{a}_i = \tilde{a}_i^{\text{Re}} + i\tilde{a}_i^{\text{Im}} \tag{5.32}$$

$$\tilde{b}_i = \tilde{b}_i^{\text{Re}} + i\tilde{b}_i^{\text{Im}} \tag{5.33}$$

The lattice evaluation also requires converting the continuous Minkowski gamma matrices (γ_M) into the Euclidean conventions (γ_E) used by the lattice simulation software:

- Correct Euclidean $\longleftrightarrow$ Minkowski γ-matrices notation:

$$\boxed{\gamma_E^1 = +i\gamma_M^1} \tag{5.34}$$

$$\boxed{\gamma_E^2 = -i\gamma_M^2} \tag{5.35}$$

$$\boxed{\gamma_E^3 = +i\gamma_M^3} \tag{5.36}$$

$$\boxed{\gamma_E^4 = \gamma_M^4} \tag{5.37}$$

$$\boxed{\gamma_E^5} = \gamma_E^1\gamma_E^2\gamma_E^3\gamma_E^4 = \boxed{-\gamma_M^5} = -i\gamma_M^4\gamma_M^1\gamma_M^2\gamma_M^3 \tag{5.38}$$

Substituting the Euclidean gamma structures and mapping $P^0 \to E_N$, we rewrite the column vector entirely in terms of the lattice ratio observables $R_{\Gamma(i)}$ and the finite-link amplitudes. The index $i \in \{1, 2, 4, 8\}$ labels each Dirac structure by its bit-encoded position (CHROMA & QLUA convention discussed in Appendix A.) in our operator basis, matching the correspondence $\Gamma(1) = [+i\gamma^1]$, $\Gamma(2) = [-i\gamma^2]$, $\Gamma(4) = [+i\gamma^3]$, and $\Gamma(8) = [\gamma^0]$ used below:

$$\begin{pmatrix} R_{\Gamma(1)} \\ R_{\Gamma(2)} \\ R_{\Gamma(4)} \\ R_{\Gamma(8)} \end{pmatrix} = \begin{pmatrix} R_{[+i\gamma^1]} \\ R_{[-i\gamma^2]} \\ R_{[+i\gamma^3]} \\ R_{[\gamma^0]} \end{pmatrix} = 2 \begin{pmatrix} -i\frac{|\vec{P}^1|}{E_N}\,\tilde{a}_2 + \frac{m_N^2}{E_N} b^1\,\tilde{a}_3 + i\frac{m_N^2}{E_N}\eta v^1\,\tilde{b}_1 - m_N b^2\,\tilde{a}_{12} + m_N^3 b^2\eta^2 (v^3)^2 \tilde{b}_{10} \\ -\frac{m_N^2}{E_N} b^2\,\tilde{a}_3 - m_N b^1 \tilde{a}_{12} - i m_N \eta v^1 \tilde{b}_7 + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10} \\ i\frac{m_N^2}{E_N}\eta v^3\,\tilde{b}_1 - m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10} \\ \tilde{a}_2 - i m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12} - \frac{i m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8 + \frac{i m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10} \end{pmatrix} \tag{5.39}$$

To separate the real and imaginary amplitude contributions analytically, we split this matrix equation into two sets by flipping the longitudinal separation component b^1. For a strictly positive longitudinal separation ($+b^1$), writing out the real and imaginary parts of the correlator ratios gives:

$$\begin{pmatrix} \mathrm{Re}\{R_{\Gamma(1)}\}^{[+b^1]} \\ \mathrm{Im}\{R_{\Gamma(1)}\}^{[+b^1]} \\ \mathrm{Re}\{R_{\Gamma(2)}\}^{[+b^1]} \\ \mathrm{Im}\{R_{\Gamma(2)}\}^{[+b^1]} \\ \mathrm{Re}\{R_{\Gamma(4)}\}^{[+b^1]} \\ \mathrm{Im}\{R_{\Gamma(4)}\}^{[+b^1]} \\ \mathrm{Re}\{R_{\Gamma(8)}\}^{[+b^1]} \\ \mathrm{Im}\{R_{\Gamma(8)}\}^{[+b^1]} \end{pmatrix} = 2 \begin{pmatrix} \frac{m_N^2}{E_N} b^1\,\tilde{a}_3^{\mathrm{Re}} - m_N b^2\,\tilde{a}_{12}^{\mathrm{Re}} + m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Re}} + \frac{|\vec{P}^1|}{E_N}\,\tilde{a}_2^{\mathrm{Im}} - \frac{m_N^2}{E_N}\eta v^1\,\tilde{b}_1^{\mathrm{Im}} \\ -\frac{|\vec{P}^1|}{E_N}\,\tilde{a}_2^{\mathrm{Re}} + \frac{m_N^2}{E_N}\eta v^1\,\tilde{b}_1^{\mathrm{Re}} + \frac{m_N^2}{E_N} b^1\,\tilde{a}_3^{\mathrm{Im}} - m_N b^2\,\tilde{a}_{12}^{\mathrm{Im}} + m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Im}} \\ -\frac{m_N^2}{E_N} b^2\,\tilde{a}_3^{\mathrm{Re}} - m_N b^1 \tilde{a}_{12}^{\mathrm{Re}} + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Re}} + m_N \eta v^1 \tilde{b}_7^{\mathrm{Im}} \\ -m_N \eta v^1 \tilde{b}_7^{\mathrm{Re}} - \frac{m_N^2}{E_N} b^2\,\tilde{a}_3^{\mathrm{Im}} - m_N b^1 \tilde{a}_{12}^{\mathrm{Im}} + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Im}} \\ -m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\mathrm{Re}} - \frac{m_N^2}{E_N}\eta v^3\,\tilde{b}_1^{\mathrm{Im}} \\ \frac{m_N^2}{E_N}\eta v^3\,\tilde{b}_1^{\mathrm{Re}} - m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\mathrm{Im}} \\ \tilde{a}_2^{\mathrm{Re}} + m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\mathrm{Im}} + \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\mathrm{Im}} - \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Im}} \\ -m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\mathrm{Re}} - \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\mathrm{Re}} + \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\mathrm{Re}} + \tilde{a}_2^{\mathrm{Im}} \end{pmatrix} \tag{5.40}$$

Evaluating the system for the negated separation ($-b^1$) flips the signs of selected terms according to the parity constraints of Eq. (4.28). By construction, the imaginary parts of the amplitudes are odd functions of b^1

($\widetilde{A}_i^{\rm Im} = \widetilde{A}_i^{(\text{b-odd})} = \widetilde{A}_i^{(b^1\text{-odd})}$ and $\widetilde{B}_i^{\rm Im} = \widetilde{B}_i^{(\text{b-odd})} = \widetilde{B}_i^{(b^1\text{-odd})}$). Applying these parity reversals yields the conjugate matrix:

$$\begin{pmatrix} \mathrm{Re}\{R_{\Gamma(1)}\}^{[-b^1]} \\ \mathrm{Im}\{R_{\Gamma(1)}\}^{[-b^1]} \\ \mathrm{Re}\{R_{\Gamma(2)}\}^{[-b^1]} \\ \mathrm{Im}\{R_{\Gamma(2)}\}^{[-b^1]} \\ \mathrm{Re}\{R_{\Gamma(4)}\}^{[-b^1]} \\ \mathrm{Im}\{R_{\Gamma(4)}\}^{[-b^1]} \\ \mathrm{Re}\{R_{\Gamma(8)}\}^{[-b^1]} \\ \mathrm{Im}\{R_{\Gamma(8)}\}^{[-b^1]} \end{pmatrix} = 2 \begin{pmatrix} -\frac{m_N^2}{E_N} b^1\, \tilde{a}_3^{\rm Re} - m_N b^2\, \tilde{a}_{12}^{\rm Re} + m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} - \frac{|\vec{P}^1|}{E_N}\, \tilde{a}_2^{\rm Im} + \frac{m_N^2}{E_N} \eta v^1\, \tilde{b}_1^{\rm Im} \\ -\frac{|\vec{P}^1|}{E_N}\, \tilde{a}_2^{\rm Re} + \frac{m_N^2}{E_N} \eta v^1\, \tilde{b}_1^{\rm Re} + \frac{m_N^2}{E_N} b^1\, \tilde{a}_3^{\rm Im} + m_N b^2\, \tilde{a}_{12}^{\rm Im} - m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ -\frac{m_N^2}{E_N} b^2\, \tilde{a}_3^{\rm Re} + m_N b^1 \tilde{a}_{12}^{\rm Re} - m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} - m_N \eta v^1 \tilde{b}_7^{\rm Im} \\ -m_N \eta v^1 \tilde{b}_7^{\rm Re} + \frac{m_N^2}{E_N} b^2\, \tilde{a}_3^{\rm Im} - m_N b^1 \tilde{a}_{12}^{\rm Im} + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ -m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\rm Re} + \frac{m_N^2}{E_N} \eta v^3\, \tilde{b}_1^{\rm Im} \\ \frac{m_N^2}{E_N} \eta v^3\, \tilde{b}_1^{\rm Re} + m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\rm Im} \\ \tilde{a}_2^{\rm Re} - m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\rm Im} - \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\rm Im} + \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ -m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\rm Re} - \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\rm Re} + \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} - \tilde{a}_2^{\rm Im} \end{pmatrix} \tag{5.41}$$

By taking the symmetric average of the two matrices, $\frac{1}{2}\left(R^{[+b]} + R^{[-b]}\right)$, we project out the b-even combination, isolating the real parts of the requisite amplitudes:

$$\begin{pmatrix} \mathrm{Re}\{R_{\Gamma(1)}\}^{[b^1\text{-even}]} \\ \mathrm{Im}\{R_{\Gamma(1)}\}^{[b^1\text{-even}]} \\ \mathrm{Re}\{R_{\Gamma(2)}\}^{[b^1\text{-even}]} \\ \mathrm{Im}\{R_{\Gamma(2)}\}^{[b^1\text{-even}]} \\ \mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]} \\ \mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]} \\ \mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]} \\ \mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]} \end{pmatrix} = 2 \begin{pmatrix} -m_N b^2\, \tilde{a}_{12}^{\rm Re} + m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} \\ -\frac{|\vec{P}^1|}{E_N}\, \tilde{a}_2^{\rm Re} + \frac{m_N^2}{E_N} \eta v^1\, \tilde{b}_1^{\rm Re} + \frac{m_N^2}{E_N} b^1\, \tilde{a}_3^{\rm Im} \\ -\frac{m_N^2}{E_N} b^2\, \tilde{a}_3^{\rm Re} \\ -m_N \eta v^1 \tilde{b}_7^{\rm Re} - m_N b^1 \tilde{a}_{12}^{\rm Im} + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ -m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\rm Re} \\ \frac{m_N^2}{E_N} \eta v^3\, \tilde{b}_1^{\rm Re} \\ \tilde{a}_2^{\rm Re} \\ -m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\rm Re} - \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\rm Re} + \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} \end{pmatrix} \tag{5.42}$$

Conversely, taking the antisymmetric half-difference of the two matrices, $\frac{1}{2}\left(R^{[+b]} - R^{[-b]}\right)$, projects out the b-odd combination, isolating the imaginary parts:

$$\begin{pmatrix} \mathrm{Re}\{R_{\Gamma(1)}\}^{[b^1\text{-odd}]} \\ \mathrm{Im}\{R_{\Gamma(1)}\}^{[b^1\text{-odd}]} \\ \mathrm{Re}\{R_{\Gamma(2)}\}^{[b^1\text{-odd}]} \\ \mathrm{Im}\{R_{\Gamma(2)}\}^{[b^1\text{-odd}]} \\ \mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]} \\ \mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]} \\ \mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]} \\ \mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]} \end{pmatrix} = 2 \begin{pmatrix} \frac{m_N^2}{E_N} b^1\, \tilde{a}_3^{\rm Re} + \frac{|\vec{P}^1|}{E_N}\, \tilde{a}_2^{\rm Im} - \frac{m_N^2}{E_N} \eta v^1\, \tilde{b}_1^{\rm Im} \\ -m_N b^2\, \tilde{a}_{12}^{\rm Im} + m_N^3 b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ -m_N b^1 \tilde{a}_{12}^{\rm Re} + m_N^3 b^1 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Re} + m_N \eta v^1 \tilde{b}_7^{\rm Im} \\ -\frac{m_N^2}{E_N} b^2\, \tilde{a}_3^{\rm Im} \\ -\frac{m_N^2}{E_N} \eta v^3\, \tilde{b}_1^{\rm Im} \\ -m_N^3 b^2 \eta^2 v^1 v^3 \tilde{b}_{10}^{\rm Im} \\ m_N \frac{|\vec{P}^1|}{E_N} b^2 \tilde{a}_{12}^{\rm Im} + \frac{m_N^3}{E_N} b^2 \eta v^1 \tilde{b}_8^{\rm Im} - \frac{m_N^3}{E_N} |\vec{P}^1| b^2 \eta^2 (v^3)^2 \tilde{b}_{10}^{\rm Im} \\ \tilde{a}_2^{\rm Im} \end{pmatrix} \tag{5.43}$$

To analytically reconstruct the Sivers shift, we solve these projected systems for the amplitudes that drive the Sivers effect: $\tilde{a}_2$, $\tilde{b}_1$, $\tilde{a}_{12}$, and $\tilde{b}_8$. Reading directly from the b-even system, we isolate the real components:

$$\tilde{a}_2^{\mathrm{Re}} = \frac{1}{2}\mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]} \tag{5.44}$$

$$\tilde{b}_1^{\mathrm{Re}} = \frac{E_N}{2m_N^2\eta v^3}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]} \tag{5.45}$$

Substituting the appropriate matrix rows, we decouple $\tilde{a}_{12}$ and $\tilde{b}_8$:

$$\tilde{a}_{12}^{\mathrm{Re}} = \frac{-1}{2m_N b^2}\left[\mathrm{Re}\{R_{\Gamma(1)}\}^{[b^1\text{-even}]} + \frac{v^3}{v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]}\right] \tag{5.46}$$

$$\tilde{b}_8^{\mathrm{Re}} = \frac{-E_N}{2m_N^3 b^2\eta v^1}\left[\mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]} + \frac{|\vec{P}^1|v^3}{E_N v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]} + 2m_N\frac{|\vec{P}^1|}{E_N}b^2\tilde{a}_{12}^{\mathrm{Re}}\right] \tag{5.47}$$

Applying identical algebraic decoupling to the b-odd system, we isolate the corresponding imaginary components:

$$\tilde{a}_2^{\mathrm{Im}} = \frac{1}{2}\mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]} \tag{5.48}$$

$$\tilde{b}_1^{\mathrm{Im}} = \frac{-E_N}{2m_N^2\eta v^3}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]} \tag{5.49}$$

and the remaining transverse amplitudes:

$$\tilde{a}_{12}^{\mathrm{Im}} = \frac{-1}{2m_N b^2}\left[\mathrm{Im}\{R_{\Gamma(1)}\}^{[b^1\text{-odd}]} + \frac{v^3}{v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]}\right] \tag{5.50}$$

$$\tilde{b}_8^{\mathrm{Im}} = \frac{E_N}{2m_N^3 b^2\eta v^1}\left[\mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]} - \frac{|\vec{P}^1|v^3}{E_N v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]} - 2m_N\frac{|\vec{P}^1|}{E_N}b^2\tilde{a}_{12}^{\mathrm{Im}}\right] \tag{5.51}$$

Finally, we convert these finite-link evaluations $(\tilde{a}_i, \tilde{b}_i)$ back into the standard continuum-invariant amplitudes $(\widetilde{A}_i, \widetilde{B}_i)$ using the proportionality dictionary:

$$\widetilde{A}_i = \tilde{a}_i \tag{5.52}$$

$$\widetilde{B}_i = (\eta v\cdot P)^n\ \tilde{b}_i \tag{5.53}$$

Recalling that $v{\cdot}P = v_1 P^1 = v^1|P^1|$ in our chosen frame, each amplitude structure has its own conversion factor. Applying this mapping to the real components yields the final accessible amplitudes on the lattice:

$$\boxed{\widetilde{A}_2^{\mathrm{Re}} = \frac{1}{2}\mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]}} \tag{5.54}$$

$$\boxed{\widetilde{B}_1^{\mathrm{Re}} = \frac{E_N|P^1|v^1}{2m_N^2 v^3}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]}} \tag{5.55}$$

$$\boxed{\widetilde{A}_{12}^{\mathrm{Re}} = \frac{-1}{2m_N b^2}\left[\mathrm{Re}\{R_{\Gamma(1)}\}^{[b^1\text{-even}]} + \frac{v^3}{v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]}\right]} \tag{5.56}$$

$$\boxed{\widetilde{B}_8^{\mathrm{Re}} = \frac{-E_N|P^1|}{2m_N^3 b^2}\left[\mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-even}]} + \frac{|\vec{P}^1|v^3}{E_N v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-even}]} + 2m_N\frac{|\vec{P}^1|}{E_N}b^2\widetilde{A}_{12}^{\mathrm{Re}}\right]} \tag{5.57}$$

Applying the same scaling factors to the imaginary counterparts provides the complementary T-odd contributions:

$$\boxed{\widetilde{A}_2^{\mathrm{Im}} = \frac{1}{2}\mathrm{Im}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]}} \tag{5.58}$$

$$\boxed{\widetilde{B}_1^{\mathrm{Im}} = \frac{-E_N|P^1|v^1}{2m_N^2\eta v^3}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]}} \tag{5.59}$$

$$\boxed{\widetilde{A}_{12}^{\mathrm{Im}} = \frac{-1}{2m_N b^2}\left[\mathrm{Im}\{R_{\Gamma(1)}\}^{[b^1\text{-odd}]} + \frac{v^3}{v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]}\right]} \tag{5.60}$$

$$\boxed{\widetilde{B}_8^{\mathrm{Im}} = \frac{E_N|P^1|}{2m_N^3 b^2}\left[\mathrm{Re}\{R_{\Gamma(8)}\}^{[b^1\text{-odd}]} - \frac{|\vec{P}^1|v^3}{E_N v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b^1\text{-odd}]} - 2m_N\frac{|\vec{P}^1|}{E_N}b^2\widetilde{A}_{12}^{\mathrm{Im}}\right]} \tag{5.61}$$

5.7 Constructing the unpolarized $\widetilde{A}_{2B}$ and Sivers $\widetilde{A}_{12B}$ Amplitudes

With the invariant amplitudes isolated by parity projection, we now combine them into the amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ of Eq. (4.34), with $R(\hat{\zeta}^2)$ from Eq. (4.32). To apply these expressions in our simulated lattice geometry, we map the contravariant spatial indices to our physical grid parameters. We define the longitudinal spatial separation as $b_L \equiv b^1$ and the transverse spatial separation as $b_T \equiv b^2$. The squared spatial separation then expands to $b^2 = b_L^2 + b_T^2$, and we identify the longitudinal nucleon momentum as $P_L \equiv P^1$.

We begin with the defining linear combination for the unpolarized amplitude, $\widetilde{A}_{2B}$. Splitting this amplitude into its real and imaginary components yields:

$$\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \hat{\zeta}) = \widetilde{A}_2^{\mathrm{Re}} + R(\hat{\zeta}^2)\widetilde{B}_1^{\mathrm{Re}} \tag{5.62}$$

$$\widetilde{A}_{2B}^{\mathrm{Im}}(b_L, b_T, \hat{\zeta}) = \widetilde{A}_2^{\mathrm{Im}} + R(\hat{\zeta}^2)\widetilde{B}_1^{\mathrm{Im}} \tag{5.63}$$

Substituting the parity-projected lattice ratio observables derived in Section 5.6 (and applying our b_L, P_L geometric mapping), we obtain the real part of $\widetilde{A}_{2B}$ as an expression computable directly from our grid coordinates:

$$\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \hat{\zeta}) = \frac{1}{2}\mathrm{Re}\{R_{\Gamma(8)}\}^{[b_L\text{-even}]} + R(\hat{\zeta}^2)\frac{E_N|P_L|v^1}{2m_N^2 v^3}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b_L\text{-even}]} \tag{5.64}$$

The same substitution for the imaginary component, now with the b_L-odd projected data sets so that time-reversal parity is preserved, gives:

$$\widetilde{A}_{2B}^{\mathrm{Im}}(b_L, b_T, \hat{\zeta}) = \frac{1}{2}\mathrm{Im}\{R_{\Gamma(8)}\}^{[b_L\text{-odd}]} - R(\hat{\zeta}^2)\frac{E_N|P_L|v^1}{2m_N^2 \eta v^3}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b_L\text{-odd}]} \tag{5.65}$$

The amplitude that drives the Sivers shift, $\widetilde{A}_{12B}$, combines the structures $\widetilde{A}_{12}$ and $\widetilde{B}_8$. Its real and imaginary decompositions are defined as:

$$\widetilde{A}_{12B}^{\mathrm{Re}}(b_L, b_T, \hat{\zeta}) = \widetilde{A}_{12}^{\mathrm{Re}} - R(\hat{\zeta}^2)\widetilde{B}_8^{\mathrm{Re}} \tag{5.66}$$

$$\widetilde{A}_{12B}^{\mathrm{Im}}(b_L, b_T, \hat{\zeta}) = \widetilde{A}_{12}^{\mathrm{Im}} - R(\hat{\zeta}^2)\widetilde{B}_8^{\mathrm{Im}} \tag{5.67}$$

Because the terms within $\widetilde{A}_{12B}$ are more complex, we first write out the inner constituent $\widetilde{A}_{12}^{\mathrm{Re}}$ in terms of the grid parameters b_L and b_T, using the spatial kinematic constraint $b^2 = b_L^2 + b_T^2$ from above. The real part of the leading-twist term becomes:

$$\widetilde{A}_{12}^{\mathrm{Re}} = \frac{-1}{2m_N(b_L^2 + b_T^2)}\left[\mathrm{Re}\{R_{\Gamma(1)}\}^{[b_L\text{-even}]} + \frac{v^3}{v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b_L\text{-even}]}\right] \tag{5.68}$$

Substituting this internal structure alongside the expansion for $\widetilde{B}_8^{\mathrm{Re}}$ gives the full equation for the real part entering the Sivers integration:

$$\begin{aligned}\widetilde{A}_{12B}^{\mathrm{Re}}(b_L, b_T, \hat{\zeta}) = \widetilde{A}_{12}^{\mathrm{Re}} - R(\hat{\zeta}^2)\Bigg(&\frac{-E_N|P_L|}{2m_N^3(b_L^2 + b_T^2)}\Bigg[\mathrm{Im}\{R_{\Gamma(8)}\}^{[b_L\text{-even}]} + \frac{|P_L|v^3}{E_N v^1}\mathrm{Re}\{R_{\Gamma(4)}\}^{[b_L\text{-even}]} \\ &+ 2m_N\frac{|P_L|}{E_N}(b_L^2 + b_T^2)\widetilde{A}_{12}^{\mathrm{Re}}\Bigg]\Bigg)\end{aligned} \tag{5.69}$$

Following the same procedure for the imaginary (T-odd) components, we first define the internal b_L-odd constituent $\widetilde{A}_{12}^{\mathrm{Im}}$:

$$\widetilde{A}_{12}^{\mathrm{Im}} = \frac{-1}{2m_N(b_L^2 + b_T^2)}\left[\mathrm{Im}\{R_{\Gamma(1)}\}^{[b_L\text{-odd}]} + \frac{v^3}{v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b_L\text{-odd}]}\right] \tag{5.70}$$

We then insert it to obtain the full equation for the imaginary part of the Sivers amplitude:

$$\widetilde{A}^{\mathrm{Im}}_{12B}(b_L, b_T, \hat{\zeta}) = \widetilde{A}^{\mathrm{Im}}_{12} - R(\hat{\zeta}^2)\Bigg(\frac{E_N|P_L|}{2m_N^3(b_L^2+b_T^2)}\Bigg[\mathrm{Re}\{R_{\Gamma(8)}\}^{[b_L\text{-odd}]} - \frac{|P_L|v^3}{E_N v^1}\mathrm{Im}\{R_{\Gamma(4)}\}^{[b_L\text{-odd}]} - 2m_N\frac{|P_L|}{E_N}(b_L^2+b_T^2)\widetilde{A}^{\mathrm{Im}}_{12}\Bigg]\Bigg) \tag{5.71}$$

5.8 Transformation to the Lorentz-Invariant Variable Space

At this stage, the amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ are discrete functions of the Cartesian lattice grid parameters b_L and b_T. Extracting the dependence on the longitudinal momentum fraction x, however, requires the Fourier transform over the Lorentz-invariant inner product $P \cdot b$. To perform this integration, we first show that our simulated Cartesian lattice coordinates map one-to-one to the invariant Variable Space.

In our selected kinematic frame, the 4-vectors for the nucleon momentum and spatial separation are defined as:

$$P^\mu = (E_N, P_L, 0, 0) \tag{5.72}$$

$$b^\mu = (0, b_L, b_T, 0) \tag{5.73}$$

Evaluating the inner product using the Minkowski metric signature $(+,-,-,-)$, the temporal and transverse orthogonal components vanish, leaving:

$$\begin{aligned} P \cdot b &= P_0 b^0 - P_1 b^1 - P_2 b^2 - P_3 b^3 \\ &= -P_L b_L \end{aligned} \tag{5.74}$$

Inverting this identity expresses the longitudinal Cartesian lattice coordinate b_L as a continuous function of the invariant scalar $P \cdot b$ and the fixed lattice momentum P_L:

$$b_L = \frac{P \cdot b}{-P_L} \tag{5.75}$$

To complete the geometric mapping, we must also ensure the squared separation term b^2 present in the denominators of our amplitudes transforms invariantly. Substituting our Lorentz expression for b_L into $b^2 = b_L^2 + b_T^2$ yields:

$$b^2 = \left(\frac{P \cdot b}{-P_L}\right)^2 + b_T^2 \tag{5.76}$$

By replacing b_L and b^2 with these invariant expressions, we transform our discrete lattice amplitudes into continuous functions within the Lorentz-invariant frame:

$$\widetilde{A}_{iB}(b_L, b_T, \hat{\zeta}) \quad \xrightarrow{\text{Coordinate Transformation}} \quad \widetilde{A}_{iB}\left(P \cdot b, b^2, \hat{\zeta}\right) \tag{5.77}$$

With the amplitudes now expressed as functions of $P \cdot b$, the Fourier transform of Eq. (5.6) can be evaluated directly. This connects the Euclidean correlation ratios evaluated on the lattice to the physical, x-dependent phenomenological observables needed to study the Sivers effect.

5.9 Numerical Results and Parity Constraints of the Extracted Amplitudes

Before examining the lattice data across the different Collins-Soper scales ($\hat{\zeta}$), we check the parity constraints of Eq. (4.28). Since $P \cdot b = -P_L b_L$ (Eq. (5.74)), the real amplitudes $\widetilde{A}^{\mathrm{Re}}_{2B}$ and $\widetilde{A}^{\mathrm{Re}}_{12B}$ must be even in the Cartesian longitudinal separation b_L, and the imaginary amplitudes $\widetilde{A}^{\mathrm{Im}}_{2B}$ and $\widetilde{A}^{\mathrm{Im}}_{12B}$ must be odd. The lattice data in the figures below show these even and odd profiles along the b_L axis, which validates the parity projection.

However, the statistical quality of the extracted signals differs between amplitudes. While the unpolarized amplitudes ($\widetilde{A}^{\mathrm{Re}}_{2B}$ and $\widetilde{A}^{\mathrm{Im}}_{2B}$) and the real Sivers amplitude ($\widetilde{A}^{\mathrm{Re}}_{12B}$) show clear, statistically resolvable signals across the simulated data, the imaginary Sivers amplitude ($\widetilde{A}^{\mathrm{Im}}_{12B}$) does not. Across all evaluated nucleon momenta and staple extents, the extracted lattice data for $\widetilde{A}^{\mathrm{Im}}_{12B}$ remains completely dominated by gauge noise and is statistically compatible with zero. With no distinguishable signal, we cannot meaningfully constrain or fit a phenomenological model to this component. We therefore discard $\widetilde{A}^{\mathrm{Im}}_{12B}$ from further analysis: we drop the $\sin(x(b \cdot P))\widetilde{A}^{\mathrm{Im}}_{12B}$ term in Eq. (5.6) and evaluate the x-dependent Sivers shift using only the $\widetilde{A}^{\mathrm{Re}}_{12B}$ data.

Sections 5.9.1–5.9.4 show the amplitudes at $P_L = -1, -2, -3, -4$.

5.9.1 Amplitude Profiles at $\hat{\zeta} = 0.090772$ ($P_L = -1$)

Figures 26 through 28 show the extracted amplitudes at this momentum, first in the discrete $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)$ lattice grid and then mapped into the Lorentz-invariant $(P \cdot b, b^2)$ variable space of Section 5.8.



Figure 26: Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 27: Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 28: $\widetilde{A}^{\rm Re}_{2B}$, $\widetilde{A}^{\rm Im}_{2B}$, $\widetilde{A}^{\rm Re}_{12B}$ & $\widetilde{A}^{\rm Im}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space. Notice the lack of discernible signal in $\widetilde{A}^{\rm Im}_{12B}$.

5.9.2 Amplitude Profiles at $\hat{\zeta} = 0.294367$ ($P_L = -2$)

Figures 29 through 31 show the corresponding amplitudes at this larger Collins-Soper value.



Figure 29: Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 30: Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 31: $\widetilde{A}^{\rm Re}_{2B}$, $\widetilde{A}^{\rm Im}_{2B}$, $\widetilde{A}^{\rm Re}_{12B}$ & $\widetilde{A}^{\rm Im}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space.

5.9.3 Amplitude Profiles at $\hat{\zeta} = 0.539684$ ($P_L = -3$)

Figures 32 through 34 give the same amplitude decomposition at $P_L = -3$.



Figure 32: Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 33: Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 34: $\widetilde{A}^{\rm Re}_{2B}$, $\widetilde{A}^{\rm Im}_{2B}$, $\widetilde{A}^{\rm Re}_{12B}$ & $\widetilde{A}^{\rm Im}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space.

5.9.4 Amplitude Profiles at $\hat{\zeta} = 0.785756$ ($P_L = -4$)

Figures 35 through 37 complete the set at our largest simulated momentum, $P_L = -4$.



Figure 35: Real and imaginary parts of the $\widetilde{A}_{2B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 36: Real and imaginary parts of the $\widetilde{A}_{12B}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space.



Figure 37: $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$, $\widetilde{A}^{\mathrm{Re}}_{12B}$ & $\widetilde{A}^{\mathrm{Im}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4,0,0\}$ at fixed $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space.

With the real and imaginary parts of $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ verified across all four simulated Collins-Soper momenta, these path-averaged, interpolated amplitude sets are the data fitted in Section 6; the fitted models then enter the Fourier-transform integral of Eq. (5.6), from which the x-dependent Sivers shift is extracted.

6 FITTING METHODOLOGIES AND STABILITY ANALYSIS

As discussed in the previous section (Sec. 5), the lattice data show varying statistical precision among the invariant amplitudes. The real part of the unpolarized amplitude, $\widetilde{A}^{\mathrm{Re}}_{2B}$, has small statistical uncertainties across the simulated position space, whereas the imaginary unpolarized amplitude $\widetilde{A}^{\mathrm{Im}}_{2B}$ and the real Sivers amplitude $\widetilde{A}^{\mathrm{Re}}_{12B}$ have larger gauge noise. We already discarded the imaginary Sivers amplitude $\widetilde{A}^{\mathrm{Im}}_{12B}$ because it is consistent with zero. We first establish a stable global fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$ as the baseline for the remaining numerical analysis, as it corresponds to the leading-twist unpolarized TMD and has the highest data quality.

We parameterize the amplitudes in the Cartesian space (b_L, b_T) rather than the Lorentz-invariant variable space $(P \cdot b, b^2)$. Fitting in (b_L, b_T) provides a larger, more uniformly distributed set of discrete data points and resolves kinematic degeneracies. Because the squared separation distance $b^2 = b_L^2 + b_T^2$ is non-negative, multiple distinct combinations of longitudinal and transverse separations (such as a large b_L with a small b_T, or a small b_L with a large b_T) map to the same invariant b^2. Fitting in the $(P \cdot b, b^2)$ basis would therefore compress these distinct orthogonal spatial dependencies into a single coordinate. Fitting over the (b_L, b_T) grid allows the model to separate the longitudinal decay profile from the transverse decay profile.

Section 6.1 removes the data with $|\boldsymbol{b}| < 3a$, which contain discretization artifacts. Section 6.2 shows that a Gaussian model gives fails for $\widetilde{A}^{\mathrm{Re}}_{2B}$, which makes the Fourier integral over $P \cdot b$ diverge. Section 6.3 searches for a functional form by symbolic regression (PySR) with an asymptotic penalty and finds a power-law tail $(1 + b^2)^{-j}$ multiplied by $e^{-f\eta}$. Section 6.4 compares Gaussian and power-law fits over $6 \leq \eta \leq 10$ and fits individual η slices to determine the η-dependence of fit parameters (c, d), and Section 6.5 selects the model for $c(\eta)$ and gives the models for the three amplitudes.

Section 6.6 fits the three amplitudes simultaneously with the full Jackknife covariance matrix, cancels the shared $e^{-f\eta}$ factor, and evaluates the x-dependent Sivers shift at each of the four momenta. Section 6.7 tests whether the amplitudes factorize into separate $(P \cdot b)$ and b^2 profiles, first with ratios of the lattice data and then with explicit factorized fits.

6.1 Discretization Effects and Short-Distance Data Exclusion

Before fitting the lattice amplitudes, we account for the limits of the discretized spacetime. As discussed in Chapter 3, Lattice QCD defines the theory on a discrete grid with a finite lattice spacing a. Observables evaluated at length scales comparable to a contain discretization artifacts. At short quark-separation distances ($|\boldsymbol{b}| \sim 1a$ or $2a$), the continuous $O(4)$ rotational symmetry of Euclidean QCD breaks down to the discrete hypercubic group of the lattice. The step-wise Wilson lines connecting these closely spaced fields also introduce $\mathcal{O}(a)$ and $\mathcal{O}(a^2)$ finite-spacing errors [78, 2]. Fitting continuum models to these short-distance points would parameterize ultraviolet lattice artifacts rather than the nucleon's non-perturbative structure.

To separate our parameterizations from these short-distance effects, we apply a spatial cut to our dataset before fitting. We exclude data points where the magnitude of the quark spatial separation is less than three lattice units:

$$|\boldsymbol{b}| = \sqrt{b_L^2 + b_T^2} \geq 3a \ . \tag{6.1}$$

With the dataset restricted to $b \geq 3a$, the non-local operator spans enough lattice sites for the step-wise gauge link to approach a continuous trajectory. The fits and subsequent Fourier transforms then depend on long-range dynamics rather than the ultraviolet regulator.

From the simulation data itself, we note that the short-distance data ($|\boldsymbol{b}| < 3a$) curve in the opposite direction compared to the long-distance data ($|\boldsymbol{b}| \geq 3a$) for $\widetilde{A}_{2B}^{\mathrm{Re}}$ and $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitudes, as shown in Figures 38 and 39.



Figure 38: The $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude at fixed momentum ($P = -1$) and fixed η (here $\eta = 8$). The short-distance data ($|\boldsymbol{b}| < 3a$) curve in the opposite direction to the long-distance data ($|\boldsymbol{b}| \geq 3a$).



Figure 39: The $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitude at fixed momentum ($P = -1$) and fixed η (here $\eta = 8$). The short-distance data ($|\boldsymbol{b}| < 3a$) curve in the opposite direction to the long-distance data ($|\boldsymbol{b}| \geq 3a$).

All later stability analyses, global fits, and extracted phenomenological parameters in this chapter use only this artifact-suppressed ($|\boldsymbol{b}| \geq 3a$) dataset.

6.2 Failure of Simple Gaussian Model

For the real parts of the amplitudes ($\widetilde{A}^{\mathrm{Re}}_{iB}$), Hermitian conjugation requires the function to be even in the longitudinal separation b_L (Eq. (4.28)). We model the real amplitude with a standard 2D Gaussian centered at the origin

$$\widetilde{A}^{\mathrm{Re}}_{iB} = \gamma \exp\left[-\alpha b_L^2 - \beta b_T^2\right] \tag{6.2}$$

To transform this into the invariant variable space, we substitute $b_T^2 = b^2 - b_L^2$ and apply the Lorentz-invariant projection $b_L = (P \cdot b)/(-P_L)$ derived in Section 5.8. Squaring this substitution gives

$$\widetilde{A}^{\mathrm{Re}}_{iB} = \gamma \exp\left[-\frac{(\alpha - \beta)}{P_L^2}(P \cdot b)^2 - \beta b^2\right] \tag{6.3}$$

For the Fourier integral $\int d(P \cdot b) e^{ix(P \cdot b)} \widetilde{A}_{iB}$ to converge, the argument of the exponential must remain negative as $(P \cdot b) \to \pm\infty$. This requires $(\alpha - \beta) > 0$. If $\alpha < \beta$, the amplitude grows exponentially at large longitudinal separations and the integral diverges.

Fitting this Gaussian model to our lattice data yields $\alpha < \beta$ for the unpolarized real amplitude, $\widetilde{A}^{\mathrm{Re}}_{2B}$.



Figure 40: Failure of Gaussian model fit for $\widetilde{A}_{2B}^{\mathrm{Re}}$ (with $\alpha - \beta = -0.0427(27)$ and $\chi^2/\mathrm{dof} \approx 235$) and $\widetilde{A}_{12B}^{\mathrm{Re}}$ (with $\alpha - \beta = -0.022(25)$ and $\chi^2/\mathrm{dof} \approx 1.4$) amplitudes.

This means the amplitude decays faster in the transverse direction than in the longitudinal direction, so the expression in the $(P \cdot b, b^2)$ space is not integrable. Figure 40 also shows that the fit quality is poor. A symmetric 2D Gaussian decay profile is too rigid to describe the non-local operator.

Because of this divergence and poor fit quality, we cannot use the simple Gaussian model. Instead, we use multi-parameter functions that accommodate distinct decay rates and ensure convergence in the Fourier domain.

6.3 Physics-Informed Symbolic Regression via PySR

Choosing a multi-dimensional functional form by hand introduces human bias, so we use machine learning (ML) methods, specifically Symbolic Regression (SR), instead.

Unlike traditional regression, which optimizes the parameters of a pre-determined function, symbolic regression searches the space of mathematical expressions and their parameters simultaneously. We use `PySR` [94], a framework that uses Genetic Programming to discover interpretable equations from data by balancing accuracy and simplicity.

Section 6.3.1 describes the search configuration, Section 6.3.2 the custom loss function, and Section 6.3.3 the resulting Pareto front.

6.3.1 Expression Search and Configuration

`PySR` [94, 120, 121] uses a genetic algorithm to iteratively evolve mathematical expressions represented as binary trees. The algorithm evaluates a population of expressions against the lattice data using a custom loss function over thousands of generations.

Table 4 lists the `PySR` search configuration: the permitted mathematical operators, the nesting constraints, and the main hyperparameters. The operator library includes a custom Lorentzian-type binary operator, $\mathrm{lor}(x, y) = 1/(1 + x^2 + y^2)^2$, alongside standard arithmetic and exponential functions. Nesting constraints prevent the algorithm from generating physically meaningless double-exponential or double-decay structures.

Table 4: Hyperparameter and operator configuration for the Physics-Informed `PySR` Symbolic Regression search.

Configuration Parameter	**Specification / Value**
Binary Operators	$+$, $-$, $\times$, $/$, $\mathrm{pow}(x, y) = x^y$, $\mathrm{lor}(x, y) = \frac{1}{(1+x^2+y^2)^2}$
Unary Operators	$\mathrm{square}(x) = x^2$, $\exp(x)$, $\mathrm{decay}(x) = \exp(-x)$
Operator Constraints	$\mathrm{pow} : (-1, 1)$, $\mathrm{lor} : (10, 10)$, $\mathrm{decay} : 10$
Nesting Constraints	$\exp \to \exp : 0$, $\mathrm{decay} \to \mathrm{decay} : 0$, $\exp \to \mathrm{decay} : 0$
Populations	30
Evolution Generations	100
Maximum Tree Complexity	28
Loss Function	Physics-Informed Weighted $\chi^2/\mathrm{dof} + \epsilon\, \mathcal{P}_{\mathrm{asymptotic}}$ ($\epsilon = 0.1$)
Input Features	n (Staple length η), b_L (Longitudinal separation), b_T (Transverse separation)
Target Observable	$\widetilde{A}_{2B}^{\mathrm{Re}}(\eta, b_L, b_T)$ (Lattice momentum $P_1 = 1$, $b \geq 3a$, $\eta \in [6, 10]$)

6.3.2 The Physics-Informed Custom Loss Function

Standard symbolic regression optimizes metrics like Mean Squared Error (MSE). However, a genetic algorithm might discover a polynomial or diverging exponential function that fits the finite (b_L, b_T) lattice grid well, but diverges as $b_L \to \infty$. As shown in Section 6.2, this would cause the subsequent Fourier transform over $P \cdot b$ to diverge.

We implemented a custom loss function within `PySR` to enforce the required asymptotic decay. The reduced chi-squared statistic determines the base accuracy of the model:

$$\mathcal{L}_{\mathrm{base}} = \frac{\chi^2}{\mathrm{dof}} = \frac{1}{N-k} \sum_{i=1}^{N} \left(\frac{\widetilde{A}_{2B}^{\mathrm{Re,\ lattice}}(i) - \widetilde{A}_{2B}^{\mathrm{Re,\ model}}(i)}{\sigma_i} \right)^2 \tag{6.4}$$

Here, N is the number of lattice data points, k is the number of degrees of freedom in the generated expression, and σ_i is the statistical error of the i-th lattice point.

To enforce the asymptotic decay required for Fourier integrability, we introduce a penalty term. We evaluate the proposed model equation at large values of the longitudinal separation, b_L^{large}, while keeping the transverse separations b_T fixed to physical slices. We compute the sum of the absolute model predictions in this asymptotic regime:

$$\mathcal{P}_{\text{asymptotic}} = \sum_{b_T \in \{b_T^{\text{fixed}}\}} \sum_{b_L \in \{b_L^{\text{large}}\}} \left| \widetilde{A}_{2B}^{\text{Re, model}}(b_L, b_T) \right| \tag{6.5}$$

If a generated equation diverges or grows at large b_L, the penalty term $\mathcal{P}_{\text{asymptotic}}$ increases, lowering the equation's fitness score. If the equation decays toward zero, the penalty vanishes.

We combine these terms into the total objective function for `PySR`, with the penalty scaled by a small regularization parameter ϵ:

$$\mathcal{L}_{\text{custom}} = \mathcal{L}_{\text{base}} + \epsilon \, \mathcal{P}_{\text{asymptotic}} \tag{6.6}$$

$$\mathcal{L}_{\text{custom}} = \frac{\chi^2}{\text{dof}} + \epsilon \sum_{b_T \in \{b_T^{\text{fixed}}\}} \sum_{b_L \in \{b_L^{\text{large}}\}} \left| \widetilde{A}_{2B}^{\text{Re, model}}(b_L, b_T) \right| \tag{6.7}$$

By evaluating fitness with $\mathcal{L}_{\text{custom}}$, `PySR` restricts the search to mathematical expressions that fit the finite-lattice data and satisfy the asymptotic decay constraints. The Julia implementation of this objective function is shown below. The asymptotic penalty evaluates each candidate expression at five test coordinates in the regime $b_L \gg 1$, spanning staple extents $\eta \in \{6, 8, 10\}$ and longitudinal separations up to $b_L = 80a$. We set the regularization weight to $\epsilon = 0.1$.

```
using Symbolics

function eval_loss(tree, dataset::Dataset{T,L}, options)::L where {T,L}
    prediction, flag = eval_tree_array(tree, dataset.X, options)
    if !flag
        return L(Inf)
    end

    # 1. Base Chi-Squared per degree of freedom
    wmse = sum(((prediction .- dataset.y) .^ 2) .* dataset.weights) / (dataset.n - 3)

    # 2. Physics boundary condition: penalty for growth at large b
    points = [
        [10.0, 50.0, 1.0],
        [10.0, 80.0, 1.0],
        [10.0, 10.0, 1.0],
        [8.0,  20.0, 1.0],
        [6.0,  30.0, 2.0]
    ]

```

```
    penalty = 0.0
    for p in points
        Xp = reshape(p, :, 1)
        y_pred, ok = eval_tree_array(tree, Xp, options)
        if ok
            penalty += (y_pred[1])^2
        else
            penalty += 1e6
        end
    end

    return L(wmse + 0.1 * penalty)
end
```

6.3.3 Pareto Front of Discovered Models

Symbolic regression outputs a Pareto front of candidate models: a ranked list of equations that trade off mathematical simplicity (measured by expression tree complexity C) and predictive accuracy (measured by the custom loss $\mathcal{L}_{\text{custom}}$). Table 5 lists the Pareto front for the real unpolarized amplitude $\widetilde{A}_{2B}^{\text{Re}}$.

Table 5: Pareto front of analytical candidate models discovered by `PySR` for the real amplitude $\widetilde{A}_{2B}^{\text{Re}}(\eta, b_L, b_T)$, ranked by mathematical complexity and loss. The optimal physics-informed models ($C = 14$ and $C = 16$) are highlighted.

Complexity	**Loss**	**Discovered Analytical Expression** $\widetilde{A}_{2B}^{\text{Re}}(\eta, b_L, b_T)$	**Physical Interpretation**
1	276.43	0.00504	Flat constant baseline
3	171.56	$(0.4988)^{\eta} \approx e^{-0.695\eta}$	Pure exponential in η only
4	167.57	$\exp\left(-\eta^{0.808}\right)$	Stretched exponential in η
7	68.93	$\frac{11.58-\eta}{(1+b_L^2+b_T^2)^2}$	Linear in η $\times$ standard Lorentzian
9	30.81	$\frac{10.16-\eta}{(1+b_L^2+b_T^2)^{1.78}}$	Linear in η $\times$ generalized Lorentzian
11	5.16	$\frac{2473.5}{\eta^4(1+b_L^2+b_T^2)^{1.44}}$	Power-law in η (η^{-4}) $\times$ Power-law in b
12	4.76	$\left(1-\frac{14.49}{\eta}\right)^2 \frac{1}{(1+b_L^2+b_T^2)^{1.46}}$	Rational in η $\times$ Power-law in b
14	**3.93**	$\mathbf{\frac{39.71}{(1+b_L^2+b_T^2)^{1.39}}\, e^{-0.534\eta}}$	**Optimal factorized:** $e^{-0.534\eta} \times (1+b^2)^{-1.39}$
16	**2.03**	$\mathbf{\frac{245.5}{(1+b_L^2+b_T^2)^{1.26}}\, e^{-0.468\eta} - 0.00127}$	**Best-fit factorized with offset (χ^2/dof = 2.03)**
17	1.99	$\frac{1}{\eta}\left(\frac{10.46}{\eta}\right)^{2.34} \frac{1.68}{(1+b_L^2+b_T^2)^{1.21}} - 0.00166$	High-order power-law $\eta^{-3.34}\times$ Power-law in b
18	1.91	$\frac{0.28}{\eta^2}(17.98-\eta)^2 \frac{1}{(1+b_L^2+b_T^2)^{1.25}} - 0.00137$	Polynomial in η $\times$ Power-law in b
19	1.76	$\frac{1}{\eta}\left(\frac{10.38}{\eta}\right)^{2.37}\left(\frac{1.39}{(1+b^2)^{1.13}} - 0.00322\right) - 0.00149$	Asymptotic refinement
21	1.76	$\frac{1}{\eta}\left(\frac{10.30}{\eta}\right)^{2.43} \frac{1.40}{(1+b^2)^{1.13}} - \frac{0.00319}{\eta} - 0.00139$	Multi-scale refinement

At the lowest complexities ($C \leq 4$), `PySR` isolates the overall η-dependence of the amplitude as an expo-

nential decay $\sim (0.499)^{\eta} \approx e^{-0.695\eta}$, consistent with the area-law suppression of the extended Wilson line. At intermediate complexities ($C = 7$ to 12), the algorithm introduces spatial b-dependence through Lorentzian-type power-law denominators $(1 + b_L^2 + b_T^2)^{-j}$ alongside linear or power-law η-models. The algorithm does not select a Gaussian spatial profile e^{-cb^2} at any point along the Pareto front. Instead, it converges on the power-law tail structure, consistent with Section 6.2.

The models at complexities $C = 14$ and $C = 16$ best balance accuracy and structure. The $C = 14$ expression has a factorized form:

$$\widetilde{A}_{2B}^{\mathrm{Re}} = \frac{39.71}{(1 + b_L^2 + b_T^2)^{1.39}}\, e^{-0.534\eta} \ , \tag{6.8}$$

which is a product of an exponential decay in η and a generalized Lorentzian power-law decay in $b^2 = b_L^2 + b_T^2$. The $C = 16$ model introduces a small additive constant offset of -0.00127, reducing the loss to $\mathcal{L}_{\mathrm{custom}} = 2.03$:

$$\widetilde{A}_{2B}^{\mathrm{Re}} = \frac{245.5}{(1 + b_L^2 + b_T^2)^{1.26}}\, e^{-0.468\eta} - 0.00127 \ . \tag{6.9}$$

Beyond $C = 16$, added complexity yields only marginal improvements in loss while reducing interpretability.

The power-law tail $(1 + b^2)^{-j}$ decays more slowly than a Gaussian but, for $j > 1/2$, is absolutely integrable over the real line, so unlike the Gaussian model of Section 6.2 it guarantees convergent Fourier integrals. The factorized structure $e^{-f\eta} \times g(b^2)$ also allows the shared η-dependent exponential to cancel analytically in the Sivers shift ratio, which we use in Section 6.6.

As shown in Figure 41, the factorized $C = 14$ and $C = 16$ expressions track the lattice data throughout the accessible (b_L, b_T) grid across all staple extents. The pure power-law-η models ($C = 11, 17$) show slight systematic tension at the boundary η values, where their η^{-n} dependence departs from the exponential decay.

The straight-line diagnostic in Figure 42 confirms this: the lattice data curve upward at large b^2 rather than following the Gaussian's expected straight line, consistent with the generalized Lorentzian power-law tail discovered by the genetic algorithm.

We therefore adopt the Lorentzian power-law structure as the spatial profile for $\widetilde{A}_{2B}^{\mathrm{Re}}$. The following sections extend this functional form to the remaining amplitudes for the global simultaneous fit.



Figure 41: The top PySR analytical models ($C = 11, 14, 16, 17$) compared with the lattice data for $\widetilde{A}_{2B}^{\mathrm{Re}}$ at fixed staple extents $\eta = 6, 7, 8, 10$ (with $P_1 = -1$ and $|\boldsymbol{b}| \geq 3a$). Each panel shows the amplitude as a function of the longitudinal separation b_L/a at fixed transverse separations b_T, with the four Pareto-front models overlaid. The two exponential-η models ($C = 14$, solid; $C = 16$, dashed) give the most balanced description of the data across all η values, while the pure power-law-η models ($C = 11$, dotted; $C = 17$, dot-dashed) deviate slightly and systematically at the ends of the η range.



Figure 42: Diagnostic straight-line test for the Gaussian hypothesis. If the amplitude followed a pure Gaussian profile $\widetilde{A}_{2B}^{\mathrm{Re}} \propto e^{-c\,b^2}$, the data would appear as a straight line on a logarithmic ordinate versus b^2 (red dashed line, best Gaussian fit with $c = 0.058$). The lattice data (black markers) curve upward away from the Gaussian line at large b^2, indicating a heavier-than-Gaussian tail, which the `PySR` power-law form $(1 + b^2)^{-1.39}$ (blue solid line) accurately captures. The curvature is direct, model-independent evidence that the spatial decay of the non-local quark bilinear follows an algebraic power law rather than a Gaussian exponential.

6.4 Systematic Exploration of the Power-Law Model

Guided by the PySR result, we now determine the η-dependence of the power-law model. We fit $\widetilde{A}_{2B}^{\mathrm{Re}}$ at momentum $P_L = -1$, with $6 \le \eta \le 10$ and $|\boldsymbol{b}| \ge 3a$. For comparison, we first refit a family of Gaussian ansätze with η-dependent widths (Figure 43); for each model, the figure lists the model expression, the fit parameters, their Jackknife-averaged values, and the resulting χ^2/DoF.

(b_L, b_T, η) Fits for Re $\widetilde{A}_{2B}$ for $P_L=-1$
with $b \geq 3$ and $6 \geq \eta \geq 10$

Model Expression	Params List	Fitted Values	χ^2/DoF
$a\, e^{-f\eta}\, e^{-c\,(b_L P_L)^2 - d\,(b_L^2+b_T^2)}$	{a, c, d, f}	{⟨2.04(13)⟩$_{2902\,J}$, ⟨0.0005 ± 0.0018⟩$_{2902\,J}$, ⟨0.0578(16)⟩$_{2902\,J}$, ⟨0.5367(86)⟩$_{2902\,J}$}	5.19673
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}\big)\,(b_L P_L)^2 - d\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, d, f}	{⟨1.48(11)⟩$_{2902\,J}$, ⟨0.0397(89)⟩$_{2902\,J}$, ⟨6.45(29)⟩$_{2902\,J}$, ⟨0.0579(16)⟩$_{2902\,J}$, ⟨0.489(11)⟩$_{2902\,J}$}	4.64804
$a\, e^{-f\eta} \exp\Big(-c\,\big(1+\frac{k_1}{\eta}-\frac{k_2}{\sqrt{\eta}}\big)\,(b_L P_L)^2 - d\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, f}	{⟨1.43(11)⟩$_{2902\,J}$, ⟨0.34(11)⟩$_{2902\,J}$, ⟨5.45(38)⟩$_{2902\,J}$, ⟨4.69(16)⟩$_{2902\,J}$, ⟨0.0579(16)⟩$_{2902\,J}$, ⟨0.483(12)⟩$_{2902\,J}$}	4.58804
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\big)\,(b_L P_L)^2 - d\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, f}	{⟨1.43(11)⟩$_{2902\,J}$, ⟨0.122(33)⟩$_{2902\,J}$, ⟨11.61(48)⟩$_{2902\,J}$, ⟨-32.8(29)⟩$_{2902\,J}$, ⟨0.0579(16)⟩$_{2902\,J}$, ⟨0.483(12)⟩$_{2902\,J}$}	4.58644
$a\, e^{-f\eta} \exp\Big(-c\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, d, kd1, f}	{⟨0.742(72)⟩$_{2902\,J}$, ⟨0.0004 ± 0.0018⟩$_{2902\,J}$, ⟨0.1216(90)⟩$_{2902\,J}$, ⟨3.41(18)⟩$_{2902\,J}$, ⟨0.384(17)⟩$_{2902\,J}$}	4.08215
$a\, e^{-f\eta} \exp\Big(-c\,(b_L P_L)^2 - d\,\big(1+\frac{l_1}{\eta}-\frac{l_2}{\sqrt{\eta}}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, d, kd1, kd2, f}	{⟨0.458(65)⟩$_{2902\,J}$, ⟨0.0002 ± 0.0017⟩$_{2902\,J}$, ⟨0.661(97)⟩$_{2902\,J}$, ⟨4.60(19)⟩$_{2902\,J}$, ⟨4.14(10)⟩$_{2902\,J}$, ⟨0.311(24)⟩$_{2902\,J}$}	3.67038
$a\, e^{-f\eta} \exp\Big(-c\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}-\frac{l_2}{\eta^2}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, d, kd1, kd2, f}	{⟨0.459(65)⟩$_{2902\,J}$, ⟨0.0002 ± 0.0017⟩$_{2902\,J}$, ⟨0.287(34)⟩$_{2902\,J}$, ⟨8.91(33)⟩$_{2902\,J}$, ⟨-23.9(13)⟩$_{2902\,J}$, ⟨0.311(24)⟩$_{2902\,J}$}	3.66774
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, d, kd1, f}	{⟨0.737(70)⟩$_{2902\,J}$, ⟨0.015(10)⟩$_{2902\,J}$, ⟨6.03(69)⟩$_{2902\,J}$, ⟨0.1147(96)⟩$_{2902\,J}$, ⟨3.23(22)⟩$_{2902\,J}$, ⟨0.383(17)⟩$_{2902\,J}$}	4.04672
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}\big)\,(b_L P_L)^2 - d\,\big(1+\frac{l_1}{\eta}-\frac{l_2}{\sqrt{\eta}}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, d, kd1, kd2, f}	{⟨0.462(63)⟩$_{2902\,J}$, ⟨0.015(11)⟩$_{2902\,J}$, ⟨6.06(74)⟩$_{2902\,J}$, ⟨0.647(95)⟩$_{2902\,J}$, ⟨4.71(20)⟩$_{2902\,J}$, ⟨4.18(11)⟩$_{2902\,J}$, ⟨0.313(23)⟩$_{2902\,J}$}	3.67204
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}-\frac{l_2}{\eta^2}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, d, kd1, kd2, f}	{⟨0.463(63)⟩$_{2902\,J}$, ⟨0.015(11)⟩$_{2902\,J}$, ⟨6.06(74)⟩$_{2902\,J}$, ⟨0.277(34)⟩$_{2902\,J}$, ⟨8.95(33)⟩$_{2902\,J}$, ⟨-24.4(14)⟩$_{2902\,J}$, ⟨0.313(23)⟩$_{2902\,J}$}	3.66931
$a\, e^{-f\eta} \exp\Big(-c\,\big(1+\frac{k_1}{\eta}-\frac{k_2}{\sqrt{\eta}}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, f}	{⟨0.568(61)⟩$_{2902\,J}$, ⟨0.66(12)⟩$_{2902\,J}$, ⟨6.79(19)⟩$_{2902\,J}$, ⟨5.222(77)⟩$_{2902\,J}$, ⟨0.130(11)⟩$_{2902\,J}$, ⟨3.61(19)⟩$_{2902\,J}$, ⟨0.344(19)⟩$_{2902\,J}$}	3.80049
$a\, e^{-f\eta} \exp\Big(-c\,\big(1+\frac{k_1}{\eta}-\frac{k_2}{\sqrt{\eta}}\big)\,(b_L P_L)^2 - d\,\big(1+\frac{l_1}{\eta}-\frac{l_2}{\sqrt{\eta}}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, kd2, f}	{⟨0.464(64)⟩$_{2902\,J}$, ⟨0.06 ± 0.15⟩$_{2902\,J}$, ⟨26.8(87)⟩$_{2902\,J}$, ⟨13.1(35)⟩$_{2902\,J}$, ⟨0.62(13)⟩$_{2902\,J}$, ⟨4.67(32)⟩$_{2902\,J}$, ⟨4.16(17)⟩$_{2902\,J}$, ⟨0.313(24)⟩$_{2902\,J}$}	3.68333
$a\, e^{-f\eta} \exp\Big(-c\,\big(1+\frac{k_1}{\eta}-\frac{k_2}{\sqrt{\eta}}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}-\frac{l_2}{\eta^2}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, kd2, f}	{⟨0.464(64)⟩$_{2902\,J}$, ⟨0.05 ± 0.15⟩$_{2902\,J}$, ⟨27.2(89)⟩$_{2902\,J}$, ⟨13.2(35)⟩$_{2902\,J}$, ⟨0.271(43)⟩$_{2902\,J}$, ⟨8.88(52)⟩$_{2902\,J}$, ⟨-24.0(22)⟩$_{2902\,J}$, ⟨0.313(23)⟩$_{2902\,J}$}	3.68087
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, f}	{⟨0.568(61)⟩$_{2902\,J}$, ⟨0.188(35)⟩$_{2902\,J}$, ⟨13.64(42)⟩$_{2902\,J}$, ⟨-45.8(24)⟩$_{2902\,J}$, ⟨0.130(11)⟩$_{2902\,J}$, ⟨3.61(19)⟩$_{2902\,J}$, ⟨0.344(19)⟩$_{2902\,J}$}	3.79855
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\big)\,(b_L P_L)^2 - d\,\big(1+\frac{l_1}{\eta}-\frac{l_2}{\sqrt{\eta}}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, kd2, f}	{⟨0.464(64)⟩$_{2902\,J}$, ⟨0.028(45)⟩$_{2902\,J}$, ⟨17.5(51)⟩$_{2902\,J}$, ⟨-71(32)⟩$_{2902\,J}$, ⟨0.62(13)⟩$_{2902\,J}$, ⟨4.65(32)⟩$_{2902\,J}$, ⟨4.15(17)⟩$_{2902\,J}$, ⟨0.313(24)⟩$_{2902\,J}$}	3.68295
$a\, e^{-f\eta} \exp\Big(-c\,\big(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\big)\,(b_L P_L)^2 - d\,\big(1-\frac{l_1}{\eta}-\frac{l_2}{\eta^2}\big)\,(b_L^2+b_T^2)\Big)$	{a, c, kc1, kc2, d, kd1, kd2, f}	{⟨0.464(64)⟩$_{2902\,J}$, ⟨0.026(45)⟩$_{2902\,J}$, ⟨19.3(61)⟩$_{2902\,J}$, ⟨-83(38)⟩$_{2902\,J}$, ⟨0.271(43)⟩$_{2902\,J}$, ⟨8.88(53)⟩$_{2902\,J}$, ⟨-24.0(22)⟩$_{2902\,J}$, ⟨0.313(23)⟩$_{2902\,J}$}	3.68043

Figure 43: 3D jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of 16 Gaussian ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta} \exp\big[-c\,(b_L P_L)^2 - d\,(b_L^2 + b_T^2)\big]$, in which the coefficients c and d are each independently allowed to be either constant or one of three η-dependent forms, $c \to c(\eta)$ and/or $d \to d(\eta)$.

The coefficient c of $(P\cdot b)^2$ must be positive and stable for the Fourier transform to x to exist. In Figure 43, most of the Gaussian fits give an unstable or negative c and a higher χ^2/DoF, which confirms the conclusion of Section 6.2. The power-law ansätze in Figure 44 reduce χ^2/DoF to ≈ 1.5. We added two more columns for AIC (Akaike information criterion) and Fourier transform of the ansätze. Making d depend on η worsens the fit, while making c depend on η improves it. This raises a new ambiguity: several η-dependences give almost

the same χ^2/DoF. For each power-law model, the figure lists the model expression, the fit parameters, their Jackknife-averaged values, χ^2/DoF, AIC, and the resulting Fourier transform in x.

Gaussian-type fit for Re $\widetilde{A}_{2B}$ for P_L=-1
$b \geq 3$ are SELECTED for fit!
$6 \geq \eta \geq 10$ are SELECTED for fit!

Model Expression	Params List	Fitted Values	χ^2/DoF	AIC	FT Plot
$a\, e^{-f\eta}\left(1+(c+d)\, b_L^2+d\, b_T^2\right)^{-j}$	a c d f	⟨2.19(16)⟩$_{2902\,J}$ ⟨0.0218(44)⟩$_{2902\,J}$ ⟨0.0218(44)⟩$_{2902\,J}$ ⟨0.5382(88)⟩$_{2902\,J}$ ⟨2.62(33)⟩$_{2902\,J}$	24.778	5461.17	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k1}{\eta}\right)\right) b_L^2+d\, b_T^2\right)^{-j}$	a c d f k1 j	⟨2.85(30)⟩$_{2902\,J}$ ⟨0.078(14)⟩$_{2902\,J}$ ⟨0.078(14)⟩$_{2902\,J}$ ⟨0.4625(89)⟩$_{2902\,J}$ ⟨6.60(15)⟩$_{2902\,J}$ ⟨2.16(15)⟩$_{2902\,J}$	2.113	474.746	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k2}{\eta^2}-\frac{k1}{\eta}\right)\right) b_L^2+d\, b_T^2\right)^{-j}$	a c d f k1 k2	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.146(51)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨12.27(50)⟩$_{2902\,J}$ ⟨-36.2(28)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.56788	355.799	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k3}{\eta^3}-\frac{k2}{\eta^2}-\frac{k1}{\eta}\right)\right) b_L^2+d\, b_T^2\right)^{-j}$	a c d f k1 k2 k3 j	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.093(18)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨6.7(52)⟩$_{2902\,J}$ ⟨36(63)⟩$_{2902\,J}$ ⟨-220(200)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.57467	357.703	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k1\,\mathrm{Log}[\eta]}{\eta}\right)\right) b_L^2+d\, b_T^2\right)^{-j}$	a c d f k1 j	⟨3.90(43)⟩$_{2902\,J}$ ⟨0.092(17)⟩$_{2902\,J}$ ⟨0.092(17)⟩$_{2902\,J}$ ⟨0.4929(86)⟩$_{2902\,J}$ ⟨3.616(83)⟩$_{2902\,J}$ ⟨2.07(14)⟩$_{2902\,J}$	1.66147	375.863	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-e^{-n\eta}\, k2-\frac{k1}{\eta}\right)\right) b_L^2+d\, b_T^2\right)^{-j}$	a c d f k1 k2 n j	⟨3.98(58)⟩$_{2902\,J}$ ⟨0.093(18)⟩$_{2902\,J}$ ⟨0.093(18)⟩$_{2902\,J}$ ⟨0.495(12)⟩$_{2902\,J}$ ⟨7.77(98)⟩$_{2902\,J}$ ⟨82(76)⟩$_{2902\,J}$ ⟨0.64(45)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.57582	357.952	

Figure 44: 3D Jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of power-law ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta}\,(1+(c+d)b_L^2+db_T^2)^{-j}$, in which the coefficient c is either constant or one of several η-dependent forms, $c \to c(\eta)$.

For completeness, we explored more η-dependent forms $c \to c(\eta)$, which yield wider Fourier transforms in x space.

Gaussian-type fit for Re $\tilde{A}_{2B}$ for P_L=-1
$b \geq 3$ are SELECTED for fit!
$6 \geq \eta \geq 10$ are SELECTED for fit!

Model Expression	Params List	Fitted Values	χ^2/DoF	AIC	FT Plot ($\eta \to \infty$)
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k1}{\eta}-\frac{k2\,\mathrm{Log}[\eta]}{\eta}\right)\right)b_L^2+d\,b_T^2\right)^{-j}$	a c d f k1 k2 j	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.240(89)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨-4.50(52)⟩$_{2902\,J}$ ⟨5.94(29)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.56865	355.966	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k1}{\eta}+\frac{k2}{\sqrt{\eta}}\right)\right)b_L^2+d\,b_T^2\right)^{-j}$	a c d f k1 k2 j	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.43(17)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨-5.81(33)⟩$_{2902\,J}$ ⟨-4.86(14)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.56911	356.065	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1-\frac{k1}{\eta}+\frac{k2\,\mathrm{Log}[\eta]}{\sqrt{\eta}}\right)\right)b_L^2+d\,b_T^2\right)^{-j}$	a c d f k1 k2 j	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.81(32)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨0.409(88)⟩$_{2902\,J}$ ⟨-1.284(19)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.56961	356.175	
$a\, e^{-f\eta}\left(1+\left(d+c\left(1+\frac{k1\sqrt{\eta}}{1+k2\,\eta}\right)\right)b_L^2+d\,b_T^2\right)^{-j}$	a c d f k1 k2 j	⟨4.01(58)⟩$_{2902\,J}$ ⟨0.59(27)⟩$_{2902\,J}$ ⟨0.094(18)⟩$_{2902\,J}$ ⟨0.496(12)⟩$_{2902\,J}$ ⟨-0.855(43)⟩$_{2902\,J}$ ⟨0.179(17)⟩$_{2902\,J}$ ⟨2.06(14)⟩$_{2902\,J}$	1.57002	356.264	

Figure 45: 3D jackknife fits of $\widetilde{A}_{2B}^{\mathrm{Re}}(b_L, b_T, \eta)$ to a family of power-law ansätze, all built on the common baseline $\widetilde{A}_{2B}^{\mathrm{Re}} = a\, e^{-f\eta}\,(1+(c+d)b_L^2+db_T^2)^{-j}$, in which the coefficient c takes further η-dependent forms, $c \to c(\eta)$, that yield wider Fourier transforms in x space.

All viable asymptotic functions in η have a common power-law profile with a power $j \approx 2$ and an amplitude scale $a \approx 4$, damped by the area-law-like suppression factor $e^{-0.5\eta}$. To isolate the η-dependence of the width parameters c and d, we fix these structural parameters ($j = 2$, $a = 4$) and fit individual η slices sequentially (Figure 46).

Fit for Re $\tilde{A}_{2B}$ at Fixed η ($6 \le \eta \le 10$) & $P_L = -1$

Spatial Cut: $|\vec{b}| \ge 3a$ | *Model:* $(4\,e^{-0.5\,\eta})/(1 + (c + d)b_L^2 + db_T^2)^2$

η	Parameters {c, d}	χ^2/DoF
6	{-0.0052(18), 0.0964(13)}	0.925
7	{-0.0021(27), 0.0967(17)}	0.634
8	{0.0011(44), 0.1001(25)}	1.846
9	{0.0091(66), 0.0987(36)}	2.353
10	{0.0121(105), 0.1016(57)}	2.384

Figure 46: Sequential Lorentzian fits for $\widetilde{A}_{2B}^{\mathrm{Re}}$ at individual η slices, holding the global power $j = 2$ and amplitude $a = 4$ constant.

From these fits, we extract the parameters c and d. The fit parameter c is the coefficient of $b_L^2 = (P\cdot b)^2/P_L^2$, while d is a coefficient of the invariant squared separation b^2. We plot the extracted values of c and d against our candidate asymptotic models to analyze their η-dependence:



Figure 47: The extracted parameter $c(\eta)$ overlaid with the candidate asymptotic models $c_1(\eta), \ldots, c_{10}(\eta)$. The extracted values approach an asymptote at large η.

For the unpolarized real amplitude $\widetilde{A}_{2B}^{\mathrm{Re}}$, the transverse width parameter d remains constant across all η, but the longitudinal width parameter $c(\eta)$ requires an asymptotic model. We select the model $c_2(\eta) = c\left(1 - \frac{k_1}{\eta} - \frac{k_2}{\eta^2}\right)$ of Figure 47, for the reasons below.

For the real Sivers amplitude $\widetilde{A}_{12B}^{\mathrm{Re}}$, we repeated the sequential Lorentzian fitting at fixed η slices. As shown in Figure 49 and Table 6, the longitudinal parameter c and transverse parameter d are independent of η or weakly dependent on η. We therefore model c and d as constants for $\widetilde{A}_{12B}^{\mathrm{Re}}$.



Figure 48: The extracted parameter $d(\eta)$, which shows no visible dependence on the staple extent η.

Fit for Re $\tilde{A}_{12\,B}$ at Fixed η ($6 \le \eta \le 10$) & $P_L = -1$

Spatial Cut: $|\vec{b}| \ge 3a$ | *Model:* $(-0.658\, e^{-0.4702\,\eta})/(1 + (c + d)b_L^2 + d b_T^2)^{2.0054}$

η	Parameters {c, d}	χ^2/DoF
6	{0.0236(169), 0.0730(56)}	0.604
7	{0.0388(288), 0.0728(77)}	0.860
8	{0.0174(424), 0.0799(127)}	0.831
9	{−0.0105(575), 0.0925(214)}	0.730
10	{−0.0723(871), 0.1230(479)}	0.468

Table 6: Sequential Lorentzian fits for $\widetilde{A}_{12B}^{\mathrm{Re}}$ at individual η slices showing consistent χ^2/dof and stable parameters.



Figure 49: Lattice extracted parameters c (left) and d (right) for $\widetilde{A}_{12B}^{\mathrm{Re}}$ as a function of η. Both parameters show no visible dependence on η.

For the imaginary unpolarized amplitude $\widetilde{A}_{2B}^{\mathrm{Im}}$, the lattice data has more statistical noise, but the sequential fitting yields a similar trend. Mirroring the structural stability of the real amplitudes, we model c and d as constant in η. Figure 50 and Table 7 show the resulting fits.

Fit for Im $\tilde{A}_{2B}$ at Fixed η ($6 \le \eta \le 10$) & $P_L = -1$

Spatial Cut: $|\vec{b}| \ge 3a$ | *Model:* $(0.267\, b_L\, e^{-0.4702\,\eta})/(1 + (c + d)b_L^2 + db_T^2)^{1.108}$

η	Parameters {c, d}	χ^2/DoF
6	{0.0129(407), 0.3277(358)}	2.033
7	{0.0521(585), 0.3059(470)}	1.938
8	{0.0671(1104), 0.3346(840)}	1.252
9	{0.2724(1536), 0.2536(987)}	1.040
10	{0.6084(2810), 0.1409(1084)}	0.645

Table 7: Sequential Lorentzian fits for $\widetilde{A}_{2B}^{\mathrm{Im}}$ at individual η slices showing consistent χ^2/dof.



Figure 50: Lattice extracted parameters c (left) and d (right) for $\widetilde{A}_{2B}^{\mathrm{Im}}$ as a function of η. The data are noisier than for the real amplitudes, but consistent with the constant-parameter ansatz.

6.5 Candidate Selection

The χ^2/dof minimizations and the sequential fits at individual η slices determine the structural parameterizations for the three surviving invariant amplitudes ($\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, and $\widetilde{A}_{12B}^{\mathrm{Re}}$; $\widetilde{A}_{12B}^{\mathrm{Im}}$ remains excluded, as noted above).

In selecting the model for $c(\eta)$ for the $\widetilde{A}_{2B}^{\mathrm{Re}}$ amplitude, goodness of fit must be paired with physical boundary conditions on the light-cone momentum fraction $x \in [-1, 1]$:

- **Quark versus Anti-quark contribution for unpolarized TMD**: In light-cone field theory, positive momentum fractions ($x > 0$) correspond to quark distributions $q(x) = f_1(x)$, while negative momentum fractions ($x < 0$) govern anti-quark distributions through $\bar{q}(x) = -f_1(-x)$ [116]. In the physical limit, at larger momentum, the correct candidate model should give a near-zero contribution for anti-quarks, i.e.,

for unpolarized TMDs' momentum fraction $x \lesssim -0.2$.

$$\left[\tilde{f}_1^{(0)}(x \lesssim -0.2) = 2\widetilde{A}_{2B}^{[x]}(x \lesssim -0.2)\right] \longrightarrow 0 \tag{6.10}$$

where, $\widetilde{A}_{2B}^{[x]}(x) = \int_{-\infty}^{+\infty} d(b\cdot P) e^{ix(b\cdot P)} \left[\widetilde{A}_{2B}^{\rm Re}(\cdots) + i\widetilde{A}_{2B}^{\rm Im}(\cdots)\right]$. This crucial condition involves both $\widetilde{A}_{2B}^{\rm Re}$ and $\widetilde{A}_{2B}^{\rm Im}$.

- **Mellin moment (x-integrated) calculations**: The previous work on lattice TMDs [116, 85, 86] calculated the first Mellin moment in x (x-integrated TMD: $f^{[1]}(\boldsymbol{k}_{\rm T}^2;\ldots) \equiv \int_{-1}^{1} dx\ f(x,\boldsymbol{k}_{\rm T}^2;\ldots)$) of the Sivers shift, which is accessible by evaluating amplitudes at $(b\cdot P)=0$:

$$\begin{aligned}\langle \boldsymbol{k}_y\rangle_{TU}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\eta v\cdot P) \ &\equiv\ m_N \frac{\tilde{f}_{1T}^{\perp[1](1)}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\ldots,\eta v\cdot P)}{\tilde{f}_1^{[1](0)}(\boldsymbol{b}_{\rm T}^2;\hat{\zeta},\ldots,\eta v\cdot P)} = -m_N \frac{\widetilde{A}_{12B}(-\boldsymbol{b}_{\rm T}^2,0,0,-1/(m_N\hat{\zeta})^2,\eta v\cdot P)}{\widetilde{A}_{2B}(-\boldsymbol{b}_{\rm T}^2,0,0,-1/(m_N\hat{\zeta})^2,\eta v\cdot P)} \\ &\xrightarrow{\boldsymbol{b}_{\rm T}^2=0} \left.\frac{\int dx \int d^2\boldsymbol{k}_{\rm T}\, \boldsymbol{k}_y\ \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}{\int dx \int d^2\boldsymbol{k}_{\rm T}\quad \Phi^{[\gamma^+]}(x,\boldsymbol{k}_{\rm T},P,S;\ldots)}\right|_{\boldsymbol{S}_{\rm T}=(1,0)}\end{aligned} \tag{6.11}$$

  The correct candidate model must reproduce the previous results. This means we calculate

$$\widetilde{A}_{2B}^{[x]}(x) = \int_{-\infty}^{+\infty} d(b\cdot P) e^{ix(b\cdot P)} \left[\widetilde{A}_{2B}^{\rm Re}(\cdots) + i\widetilde{A}_{2B}^{\rm Im}(\cdots)\right] \tag{6.12}$$

$$\widetilde{A}_{12B}^{[x]}(x) = \int_{-\infty}^{+\infty} d(b\cdot P) e^{ix(b\cdot P)} \left[\widetilde{A}_{12B}^{\rm Re}(\cdots)\right] \tag{6.13}$$

  from candidate model for $\widetilde{A}_{2B}^{\rm Re}$, $\widetilde{A}_{2B}^{\rm Im}$, and $\widetilde{A}_{12B}^{\rm Re}$, then we calculate

$$\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\rm Re} - \widetilde{A}_{2B}^{[x]\rm Im}) \tag{6.14}$$

$$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\rm Re} \tag{6.15}$$

  to calculate $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$. This condition involves all $\widetilde{A}_{2B}^{\rm Re}$, $\widetilde{A}_{2B}^{\rm Im}$ and $\widetilde{A}_{12B}^{\rm Re}$ amplitudes.

The model $c_2(\eta)$ (of Figure 47) for real unpolarized amplitude $\widetilde{A}_{2B}^{\rm Re} = \left[\frac{ae^{-f\eta}}{\left[1+\left(d+c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\right)b_L^2+db_T^2\right]^j}\right]$ combined with the imaginary unpolarized amplitude model $\widetilde{A}_{2B}^{\rm Im} = \left[\frac{a''b_Le^{-f\eta}}{\left[1+(c''+d'')b_L^2+d''b_T^2\right]^{j''}}\right]$ suppresses anti-quark contributions in above mentioned regime and has the lowest AIC (and χ^2/DoF). The row 1 and column 3 of Figures 55, 61, 67, 73 plot the $\tilde{f}_1^{(0)}(x) = 2\widetilde{A}_{2B}^{[x]}(x)$, which shows near-zero contribution for anti-quark ($x \lesssim -0.2$) for higher momentum.

The selected model for $\widetilde{A}_{2B}^{\rm Re}$, $\widetilde{A}_{2B}^{\rm Im}$ and along with the model for $\widetilde{A}_{12B}^{\rm Re} = \left[\frac{-a'e^{-f\eta}}{\left[1+(c'+d')b_L^2+d'b_T^2\right]^{j'}}\right]$, produce Mellin moment (x-integrated) calculations of Sivers shift TMD $\langle \boldsymbol{k}_y\rangle_{TU} = m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ which is consistent previously

published results. Tables 9, 11, 13, 15, calculate the $\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and finally the Mellin moment of Sivers shift TMD $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ for integration limit of $[-\infty, \infty]$, $[-1, 1]$, $[-1, 0]$, and $[0, 1]$ which are consistent with previously published results.

In summary the final candidate analytical models in Cartesian (b_L, b_T) space for the three amplitudes are:

$$\widetilde{A}_{2B}^{\mathrm{Re}} = \left[\frac{ae^{-f\eta}}{\left[1 + \left(d + c\left(1 - \frac{k_1}{\eta} - \frac{k_2}{\eta^2}\right)\right) b_L^2 + db_T^2\right]^j}\right] \tag{6.16}$$

$$\widetilde{A}_{12B}^{\mathrm{Re}} = \left[\frac{-a'e^{-f\eta}}{\left[1 + (c' + d')b_L^2 + d'b_T^2\right]^{j'}}\right] \tag{6.17}$$

$$\widetilde{A}_{2B}^{\mathrm{Im}} = \left[\frac{a''b_L e^{-f\eta}}{\left[1 + (c'' + d'')b_L^2 + d''b_T^2\right]^{j''}}\right] \tag{6.18}$$

6.6 Simultaneous Fitting Strategy

The structural models in Eq. (6.16)-(6.18) show that all three surviving amplitudes share an η-dependence damped by a common exponential decay factor, $e^{-f\eta}$. Because the Sivers shift is a ratio, this shared factor cancels in the x-dependent result.

We evaluate the Sivers shift of Eq. (5.6). With $\widetilde{A}_{12B}^{\mathrm{Im}}$ dropped (Section 5.9) and only the even-parity combinations surviving the symmetric integration, it reduces to

$$\langle \boldsymbol{k}_y \rangle_{TU} = -m_N \frac{\int d(b\cdot P) \cos\left(x(b\cdot P)\right) \widetilde{A}_{12B}^{\mathrm{Re}}(\cdots)}{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right) \widetilde{A}_{2B}^{\mathrm{Re}}(\cdots) - \sin\left(x(b\cdot P)\right) \widetilde{A}_{2B}^{\mathrm{Im}}(\cdots)\right]} \,. \tag{6.19}$$

Substituting the analytical models from Eq. (6.16)-(6.18), the shared $e^{-f\eta}$ term is independent of the integration variable $b\cdot P$ and cancels between the numerator and denominator integrals. Applying the mapping of Section 5.8 ($b_L = P\cdot b/(-P_L)$, $b_T^2 = b^2 - b_L^2$), we arrive at the final integrands in the invariant frame:

$$\begin{aligned}
\langle \boldsymbol{k}_y \rangle_{TU} &= m_N \frac{\int d(b\cdot P) \cos\left(x(b\cdot P)\right) \frac{a'}{D_{12B}^{\mathrm{Re}}}}{\int d(b\cdot P) \left[\cos\left(x(b\cdot P)\right) \frac{a}{D_{2B}^{\mathrm{Re}}} - \sin\left(x(b\cdot P)\right) \frac{a''(P\cdot b)/(-P_L)}{D_{2B}^{\mathrm{Im}}}\right]} \,, \\
D_{12B}^{\mathrm{Re}} &= \left[1 + (c')\frac{(P\cdot b)^2}{P_L^2} + d'b^2\right]^{j'} \,, \\
D_{2B}^{\mathrm{Re}} &= \left[1 + \left(c\left(1 - \frac{k_1}{\eta} - \frac{k_2}{\eta^2}\right)\right)\frac{(P\cdot b)^2}{P_L^2} + db^2\right]^{j} \,, \\
D_{2B}^{\mathrm{Im}} &= \left[1 + (c'')\frac{(P\cdot b)^2}{P_L^2} + d''b^2\right]^{j''} \,.
\end{aligned} \tag{6.20}$$

We enforce this shared η-dependence with a simultaneous global fit in which the parameter f is the same in all three amplitude models ($\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, and $\widetilde{A}_{12B}^{\mathrm{Re}}$). This keeps the three models theoretically consistent and

stabilizes the parameter extraction.

Section 6.6.1 describes the correlated fit used at each momentum.

6.6.1 Simultaneous Fit with Full Covariance Matrix

Because we extract the amplitudes from the same gauge field configurations and evaluate them across nearby spatial coordinates (b_L, b_T), their statistical fluctuations are strongly correlated. An uncorrelated χ^2 minimization weights these data points incorrectly, which underestimates the uncertainties and biases the central values.

We therefore perform a correlated χ^2 minimization with the full covariance matrix, which accounts for the cross-correlations in the lattice data, both across different quark spatial separations and across the real and imaginary amplitude components.

We estimate these correlations using Jackknife resampling. Let $y_i^{(k)}$ represent the lattice evaluation of the i-th data point (spanning all spatial coordinates across $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, and $\widetilde{A}_{12B}^{\mathrm{Re}}$) in the k-th Jackknife bin. The full Jackknife covariance matrix C_{ij} is defined as [122]:

$$C_{ij} = \frac{N-1}{N} \sum_{k=1}^{N} \left(y_i^{(k)} - \bar{y}_i \right) \left(y_j^{(k)} - \bar{y}_j \right) \ , \tag{6.21}$$

where N is the total number of Jackknife samples and $\bar{y}_i = \frac{1}{N} \sum_k y_i^{(k)}$ is the sample mean.

The global simultaneous fit minimizes the correlated χ^2 objective function with respect to the model parameters $\mathbf{p}$:

$$\chi^2(\mathbf{p}) = \sum_{i,j} \left(\bar{y}_i - F_i(\mathbf{p}) \right) C_{ij}^{-1} \left(\bar{y}_j - F_j(\mathbf{p}) \right) \ , \tag{6.22}$$

where $F_i(\mathbf{p})$ represents the analytical models evaluated at the corresponding kinematics.

We use the `lsqfit` [123] non-linear least squares fitting package and the `gvar` library for Gaussian error propagation. We construct the full Jackknife covariance matrix that captures the cross-correlations between $\widetilde{A}_{2B}^{\mathrm{Re}}$, $\widetilde{A}_{2B}^{\mathrm{Im}}$, and $\widetilde{A}_{12B}^{\mathrm{Re}}$. The `gvar` framework binds the combined lattice data array to the inverse covariance matrix.

The following subsections give the results of these correlated simultaneous fits at each of the discrete lattice momenta.

6.6.2 Simultaneous Fit Results for $\hat{\zeta} = 0.090772$ $(\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-1, 0, 0\})$

The extracted parameters from the simultaneous fit for the lowest simulated nucleon momentum, $P_1 = -1$ ($\hat{\zeta} = 0.090772$), are in Table 8.

Table 8: Simultaneous Fit with common f: for $P_L = -1$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global $\chi^2/\text{DoF} = 1.85(10)$

Amplitude	Model Expression	Params List	Fitted Values
$\widetilde{A}_{2B}^{\text{Re}}$	$\frac{ae^{-f\eta}}{\left[1+\left(d+c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\right)b_L^2+db_T^2\right]^j}$	$\{a, c, d, k_1, k_2, j, f\}$	$[3.863(11), 0.336(40), 0.1630(74), 12.288(23), -36.1965(45), 1.627(25), 0.4702(46)]$
$\widetilde{A}_{2B}^{\text{Im}}$	$\frac{ab_Le^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.267(13), 0.2253(69), 0.2205(66), 1.108(21), 0.4702(46)]$
$\widetilde{A}_{12B}^{\text{Re}}$	$\frac{-ae^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.658(22), 0.0773(46), 0.0747(43), 2.0054(90), 0.4702(46)]$

Figures 51, 52, 53 show the 3D fit plots on the raw lattice data in both $\{b_L, b_T, \eta\}$-space and $\{(b \cdot P), b^2, \eta\}$-space. Because we exclude the short-distance data ($|\boldsymbol{b}| < 3a$), the fit describes the long-range dynamics, and ultraviolet lattice artifacts do not influence the results. Figure 54 shows the fit plots on the raw lattice data in both $\{b_L, b_T, \eta = 8\}$-space and $\{(b \cdot P), b^2, \eta = 8\}$-space at a specific η slice ($\eta = 8$).



Figure 51: Fit for $\widetilde{A}_{2B}^{\text{Re}}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-1, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 52: Fit for $\widetilde{A}_{2B}^{\mathrm{Im}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 53: Fit for $\widetilde{A}_{12B}^{\mathrm{Re}}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 54: Fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ & $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-1,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit.

From the continuous parametrization of the amplitudes, we evaluate the Fourier integrals over the invariant scalar $P \cdot b$,

$$\widetilde{A}_{2B}^{[x]\mathrm{Re}}(x, b^2; \ldots) = \int_{-\infty}^{+\infty} \frac{d(b\cdot P)}{2\pi} e^{ix(b\cdot P)}\, \widetilde{A}_{2B}^{\mathrm{Re}}((b\cdot P), b^2; \ldots)/\widetilde{\mathcal{S}}(b^2; \ldots) \tag{6.23}$$

$$i\widetilde{A}_{2B}^{[x]\mathrm{Im}}(x, b^2; \ldots) = \int_{-\infty}^{+\infty} \frac{d(b\cdot P)}{2\pi} e^{ix(b\cdot P)}\, \widetilde{A}_{2B}^{\mathrm{Im}}((b\cdot P), b^2; \ldots)/\widetilde{\mathcal{S}}(b^2; \ldots) \tag{6.24}$$

$$\widetilde{A}_{12B}^{[x]\mathrm{Re}}(x, b^2; \ldots) = \int_{-\infty}^{+\infty} \frac{d(b\cdot P)}{2\pi} e^{ix(b\cdot P)}\, \widetilde{A}_{12B}^{\mathrm{Re}}((b\cdot P), b^2; \ldots)/\widetilde{\mathcal{S}}(b^2; \ldots) \tag{6.25}$$

Note that, since $\widetilde{A}_{12B}^{\mathrm{Im}}$ is odd in $(b \cdot P)$, its Fourier transform gives an imaginary solution (as discussed in Eq. (5.8)). We can calculate the Fourier-transformed TMDs using Eqs. (4.43)-(4.44):

$$\tilde{f}_1^{(0)} = 2\left[\widetilde{A}_{2B}^{[x]\mathrm{Re}} + i(i\widetilde{A}_{2B}^{[x]\mathrm{Im}})\right] = 2\left[\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}}\right], \tag{6.26}$$

$$\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}. \tag{6.27}$$

As we detailed in Section 6.5, Tables 9, calculate the $\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and finally the Mellin moment of Sivers shift TMD $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ for integration limit of $[-\infty, \infty]$, $[-1, 1]$, $[-1, 0]$, and $[0, 1]$; the results are consistent with previous studies of x-moment lattice calculations [116, 86].

Table 9: Numerical Integration of Fourier Transforms: $P_L = -1, b_{min} \geq 3$

Integral	$[-\infty, \infty]$	$[-1, 1]$	$[-1, 0]$	$[0, 1]$
$\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$	0.889(24)	0.780(29)	0.390(14)	0.390(14)
$\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$	-1(108)e-17	2(282)e-17	0.1223(58)	-0.1223(58)
$\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$	1.778(47)	1.560(58)	0.535(25)	1.024(36)
$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$	0.469(23)	0.454(23)	0.227(12)	0.227(12)
$m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$	0.284(14)	0.313(18)	0.456(29)	0.238(13)

Figs. 55 and 56 show the x-dependence obtained from the Fourier transforms of the fitted amplitudes: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$, $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, the Sivers shift $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$ ([momentum fraction] $\times$ [Sivers shift]). Figure 55 is evaluated at fixed $b^2 = 9a^2$, and Figure 56 shows the same quantities as functions of x and b^2. The bands are the Jackknife uncertainties propagated from the fit parameters.



Figure 55: x-dependence at $b^2 = 9a^2$ for $P_L = -1$. Top row: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, and $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$. Bottom row: $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$. Bands show the Jackknife uncertainty.



Figure 56: The quantities of Figure 55 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -1$.

6.6.3 Simultaneous Fit Results for $\hat{\zeta} = 0.294367$ $(\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-2, 0, 0\})$

The extracted parameters from the simultaneous fit for the nucleon momentum, $P_1 = -2$ ($\hat{\zeta} = 0.294367$), are in Table 10.

Table 10: Simultaneous Fit with common f: for $P_L = -2$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global $\chi^2/\text{DoF} = 1.197(85)$

Amplitude	Model Expression	Params List	Fitted Values
$\widetilde{A}_{2B}^{\text{Re}}$	$\frac{ae^{-f\eta}}{\left[1+\left(d+c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\right)b_L^2+db_T^2\right]^j}$	$\{a, c, d, k_1, k_2, j, f\}$	$[3.9833(80), 0.166(35), 0.1020(46), 12.305(14), -36.1919(27), 2.061(20), 0.5357(66)]$
$\widetilde{A}_{2B}^{\text{Im}}$	$\frac{ab_Le^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.368(12), 0.1346(69), 0.1305(67), 1.141(20), 0.5357(66)]$
$\widetilde{A}_{12B}^{\text{Re}}$	$\frac{-ae^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.632(19), 0.0769(54), 0.0741(52), 2.0141(69), 0.5357(66)]$

Figures 57, 58, 59 and 60 show the fit on lattice data, and Figures 61 and 62 show the resulting x-dependence.



Figure 57: Fit for $\widetilde{A}_{2B}^{\text{Re}}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-2, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 58: Fit for $\widetilde{A}^{\mathrm{Im}}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 59: Fit for $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 60: Fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ & $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-2,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit.

From the continuous parametrization of the amplitudes, we evaluate the Fourier integrals over the invariant scalar $P \cdot b$. As we detailed in Section 6.5, Tables 11, calculate the $\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and finally the Mellin moment of Sivers shift TMD $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ for integration limit of $[-\infty, \infty]$, $[-1, 1]$, $[-1, 0]$, and $[0, 1]$; the results are consistent with previous studies of x-moment lattice calculations [116, 86].

Table 11: Numerical Integration of Fourier Transforms: $P_L = -2, b_{min} \geq 3$

Integral	$[-\infty, \infty]$	$[-1, 1]$	$[-1, 0]$	$[0, 1]$
$\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$	1.041(41)	1.035(43)	0.518(22)	0.518(22)
$\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$	-6(176)e-17	7(444)e-17	0.257(13)	-0.257(13)
$\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$	2.081(82)	2.070(86)	0.520(34)	1.550(63)
$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$	0.451(27)	0.451(27)	0.225(14)	0.225(14)
$m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$	0.233(14)	0.234(14)	0.466(36)	0.1565(95)

Figs. 61 and 62 show the x-dependence obtained from the Fourier transforms of the fitted amplitudes: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$, $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, the Sivers shift $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$ ([momentum fraction] $\times$ [Sivers shift]). Figure 61 is evaluated at fixed $b^2 = 9a^2$, and Figure 62 shows the same quantities as functions of x and b^2. The bands are the Jackknife uncertainties propagated from the fit parameters.



Figure 61: x-dependence at $b^2 = 9a^2$ for $P_L = -2$. Top row: $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$, $i\widetilde{A}^{[x]\mathrm{Im}}_{2B}$, and $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$. Bottom row: $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$, $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$. Bands show the Jackknife uncertainty.



Figure 62: The quantities of Figure 61 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -2$.

6.6.4 Simultaneous Fit Results for $\hat{\zeta} = 0.539684$ $(\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-3, 0, 0\})$

The extracted parameters from the simultaneous fit for the nucleon momentum, $P_1 = -3$ ($\hat{\zeta} = 0.539684$), are in Table 12.

Table 12: Simultaneous Fit with common f: for $P_L = -3$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global $\chi^2/\text{DoF} = 0.712(66)$

Amplitude	Model Expression	Params List	Fitted Values
$\widetilde{A}_{2B}^{\text{Re}}$	$\frac{ae^{-f\eta}}{\left[1+\left(d+c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\right)b_L^2+db_T^2\right]^j}$	$\{a, c, d, k_1, k_2, j, f\}$	$[4.0061(31), 0.45(12), 0.136(11), 12.284(11), -36.1973(19), 2.085(10), 0.540(15)]$
$\widetilde{A}_{2B}^{\text{Im}}$	$\frac{ab_Le^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.328(11), 0.163(15), 0.147(13), 1.057(15), 0.540(15)]$
$\widetilde{A}_{12B}^{\text{Re}}$	$\frac{-ae^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.569(13), 0.100(17), 0.094(16), 1.9969(49), 0.540(15)]$

Figures 63, 64, 65 and 66 show the fit on lattice data, and Figures 67 and 68 show the resulting x-dependence.



Figure 63: Fit for $\widetilde{A}_{2B}^{\text{Re}}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-3, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 64: Fit for $\widetilde{A}^{\mathrm{Im}}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 65: Fit for $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 66: Fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ & $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-3,0,0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit.

From the continuous parametrization of the amplitudes, we evaluate the Fourier integrals over the invariant scalar $P \cdot b$. As we detailed in Section 6.5, Tables 13, calculate the $\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and finally the Mellin moment of Sivers shift TMD $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ for integration limit of $[-\infty, \infty]$, $[-1, 1]$, $[-1, 0]$, and $[0, 1]$; the results are consistent with previous studies of x-moment lattice calculations [116, 86].

Table 13: Numerical Integration of Fourier Transforms: $P_L = -3, b_{min} \geq 3$

Integral	$[-\infty, \infty]$	$[-1, 1]$	$[-1, 0]$	$[0, 1]$
$\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$	0.759(69)	0.754(70)	0.377(35)	0.377(35)
$\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$	-1(320)e-17	2(33)e-16	0.234(23)	-0.235(23)
$\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$	1.52(14)	1.51(14)	0.286(55)	1.22(11)
$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$	0.333(51)	0.333(51)	0.167(25)	0.167(25)
$m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$	0.236(32)	0.238(32)	0.63(13)	0.147(20)

Figs. 67 and 68 show the x-dependence obtained from the Fourier transforms of the fitted amplitudes: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$, $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, the Sivers shift $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$ ([momentum fraction] $\times$ [Sivers shift]). Figure 67 is evaluated at fixed $b^2 = 9a^2$, and Figure 68 shows the same quantities as functions of x and b^2. The bands are the Jackknife uncertainties propagated from the fit parameters.



Figure 67: x-dependence at $b^2 = 9a^2$ for $P_L = -2$. Top row: $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$, $i\widetilde{A}^{[x]\mathrm{Im}}_{2B}$, and $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$. Bottom row: $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$, $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$. Bands show the Jackknife uncertainty.



Figure 68: The quantities of Figure 67 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -2$.

6.6.5 Simultaneous Fit Results for $\hat{\zeta} = 0.785756$ $(\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-4, 0, 0\})$

The extracted parameters from the simultaneous fit for the nucleon momentum, $P_1 = -4$ ($\hat{\zeta} = 0.785756$), are in Table 14.

Table 14: Simultaneous Fit with common f: for $P_L = -4$, $b_{min} \geq 3$, $\eta \in [6, 10]$. Global $\chi^2/\text{DoF} = 0.434(51)$

Amplitude	Model Expression	Params List	Fitted Values
$\widetilde{A}_{2B}^{\text{Re}}$	$\frac{ae^{-f\eta}}{\left[1+\left(d+c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\right)b_L^2+db_T^2\right]^j}$	$\{a, c, d, k_1, k_2, j, f\}$	$[4.0086(15), 0.27(21), 0.122(13), 12.2751(58), -36.19925(89), 2.0654(56), 0.577(19)]$
$\widetilde{A}_{2B}^{\text{Im}}$	$\frac{ab_Le^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.269(13), 0.264(47), 0.196(14), 1.0084(79), 0.577(19)]$
$\widetilde{A}_{12B}^{\text{Re}}$	$\frac{-ae^{-f\eta}}{\left[1+(c+d)b_L^2+db_T^2\right]^j}$	$\{a, c, d, j, f\}$	$[0.5824(55), 0.116(19), 0.102(18), 2.0059(34), 0.577(19)]$

Figures 69, 70, 71 and 72 show the fit on lattice data, and Figures 73 and 74 show the resulting x-dependence.



Figure 69: Fit for $\widetilde{A}_{2B}^{\text{Re}}$ amplitude for $\boldsymbol{P} \cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P} \cdot \boldsymbol{b}) \cdot \hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 70: Fit for $\widetilde{A}^{\mathrm{Im}}_{2B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 71: Fit for $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitude for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4,0,0\}$. The top row displays the $(b_L/a, b_T/a, \eta|\boldsymbol{v}|/a)-$space, while the bottom row displays the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2, \eta|\boldsymbol{v}|/a)-$space with the global simultaneous fit overlaid.



Figure 72: Fit for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ & $\widetilde{A}^{\mathrm{Re}}_{12B}$ amplitudes for $\boldsymbol{P}\cdot a\hat{L}/(2\pi) = \{-4, 0, 0\}$ at a fixed staple extent $\eta|\boldsymbol{v}|/a = 8$. The top two rows display the $(b_L/a, b_T/a)-$space, while the bottom two rows display the $((\boldsymbol{P}\cdot\boldsymbol{b})\cdot\hat{L}/(2\pi), b^2/a^2)-$space with the fit.

From the continuous parametrization of the amplitudes, we evaluate the Fourier integrals over the invariant scalar $P \cdot b$. As we detailed in Section 6.5, Tables 15, calculate the $\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$, $\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and finally the Mellin moment of Sivers shift TMD $m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$ for integration limit of $[-\infty, \infty]$, $[-1, 1]$, $[-1, 0]$, and $[0, 1]$; the results are consistent with previous studies of x-moment lattice calculations [116, 86].

Table 15: Numerical Integration of Fourier Transforms: $P_L = -4, b_{min} \geq 3$

Integral	$[-\infty, \infty]$	$[-1, 1]$	$[-1, 0]$	$[0, 1]$
$\int dx\, \widetilde{A}_{2B}^{[x]\mathrm{Re}}$	0.87(10)	0.87(10)	0.434(52)	0.434(52)
$\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Im}})$	3(207)e-17	2(21)e-16	0.153(26)	-0.154(26)
$\int dx\, \tilde{f}_1^{(0)} = 2\int dx\, (\widetilde{A}_{2B}^{[x]\mathrm{Re}} - \widetilde{A}_{2B}^{[x]\mathrm{Im}})$	1.74(21)	1.74(21)	0.56(12)	1.18(12)
$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2\int dx\, \widetilde{A}_{12B}^{[x]\mathrm{Re}}$	0.315(55)	0.315(55)	0.157(28)	0.157(28)
$m_N \frac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$	0.195(37)	0.195(37)	0.301(73)	0.144(27)

Figs. 73 and 74 show the x-dependence obtained from the Fourier transforms of the fitted amplitudes: $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$, $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, the Sivers shift $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$ ([momentum fraction] $\times$ [Sivers shift]). Figure 73 is evaluated at fixed $b^2 = 9a^2$, and Figure 74 shows the same quantities as functions of x and b^2. The bands are the Jackknife uncertainties propagated from the fit parameters.



Figure 73: x-dependence at $b^2 = 9a^2$ for $P_L = -2$. Top row: $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$, $i\widetilde{A}^{[x]\mathrm{Im}}_{2B}$, and $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$. Bottom row: $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$, $\langle k_y \rangle_{TU}$, and $x\langle k_y \rangle_{TU}$. Bands show the Jackknife uncertainty.



Figure 74: The quantities of Figure 73 as functions of x and $b^2/a^2 \in [9, 68]$ for $P_L = -2$.

6.7 Test for Factorization in $(P \cdot b)$ and b^2

The selected candidate model for amplitudes, and all other explored ansatze, were not factorized functions of $(P \cdot b)$ and b^2. In Section 6.2, we discussed the failure of a simple Gaussian model, which is a factorized function of $(P \cdot b)$ and b^2. In Section 6.3, we concluded that symbolic regression (PySR) did not find any factorized function. Figure 43 in Section 6.4 explored numerous factorized Gaussian models with different η dependencies in fit parameters. To provide more detail on non-factorization in lattice TMD amplitudes, we explore ratios of lattice data as a function of $(P \cdot b)$ and b^2 in Section 6.7.1 to test factorization. Finally, Section 6.7.2 explores more explicit factorized fits for TMD amplitudes and shows why it fails.

6.7.1 Ratio Test on the Lattice Data

We first check directly on the lattice data whether an amplitude can be written as a product of longitudinal and transverse profiles, $\widetilde{A}(P \cdot b, b^2) = F(P \cdot b)\, G(b^2)$. If it can, any ratio taken at a common b^2 between two values of $P \cdot b$ is a constant, and any ratio taken at a common $P \cdot b$ between two values of b^2 does not depend on $P \cdot b$. We form

$$T(P \cdot b, b^2) = \frac{\widetilde{A}(P \cdot b, b^2)}{\widetilde{A}(P \cdot b = 4, b^2)} \stackrel{?}{=} \frac{F(P \cdot b)}{F(4)} \,, \tag{6.28}$$

$$B(P \cdot b, b^2) = \frac{\widetilde{A}(P \cdot b, b^2)}{\widetilde{A}(P \cdot b, b^2 = 18)} \stackrel{?}{=} \frac{G(b^2)}{G(18)} \,, \tag{6.29}$$

where the second equality in each line holds only under factorization. T should be flat in b^2 and B should be flat in $P \cdot b$. Both ratios are computed on each Jackknife sample, and the error comes from the Jackknife spread of the ratio.

The limitation is in how the data are sampled. At fixed η and P_L, the (b_L, b_T) points computed on the lattice map onto an irregular set of $(P \cdot b, b^2)$ pairs, listed in Table 16 for $\eta|v|/a = 8$ and $P_L = -1$. There are 34 pairs, but the rows at $\pm P \cdot b$ are symmetric copies of each other (identical for the real amplitudes and opposite in sign for $\widetilde{A}^{\mathrm{Im}}_{2B}$), so only $P \cdot b \geq 0$ carries independent information. At each $(P \cdot b, b^2)$, the two points with $\pm b_T$ are independent measurements, and we average them sample by sample. After the cut $|\boldsymbol{b}| \geq 3a$, that is $b^2 \geq 9$, only three values of $P \cdot b$ have more than one b^2: $P \cdot b = 0$ with $b^2 = 9, 16, 25, 36, 49$; $P \cdot b = 3$ with $b^2 = 18, 45$; and $P \cdot b = 4$ with $b^2 = 17, 20, 32$. No two of these rows share two b^2 values, so the ratios in

Eqs. (6.28) and (6.29) cannot be built from data alone.

$P \cdot b$	b^2/a^2
0	0, 1, 4, **9, 16, 25, 36, 49**
±1	2, 5
±2	5, 8, 20
±3	**18, 45**
±4	**17, 20, 32**
±5	50
±6	45
±8	68

Table 16: All 34 $(P \cdot b, b^2)$ pairs available at $\eta|v|/a = 8$ and $P_L = -1$, where $P \cdot b = -P_L\, b_L$ and $b^2 = b_L^2 + b_T^2$. Grey entries fail the cut $|\boldsymbol{b}| \geq 3a$. Bold entries are the data used in the factorization test of Figure 75.

To address this, we place the three rows on a common grid $b^2 = 18, 20, 25, 32$ that spans all three rows. On the $P \cdot b = 0$ and $P \cdot b = 4$ rows, we use quadratic interpolation in b^2: the value at a grid point comes from the parabola through the three data points on the same row that lie closest to it. The $P \cdot b = 3$ row has only two data points, $b^2 = 18$ and 45, so there we use linear interpolation between them. There is no extrapolation. As a check, Figure 75 also shows the ratios obtained with linear interpolation on all three rows, between the nearest data points above and below each grid value. Table 17 lists which grid values are data and which are interpolated: 4 of the 12 are data, and the other 8 are interpolated. For $\widetilde{A}_{2B}^{\mathrm{Im}}$, which is odd in $P \cdot b$ and vanishes at $P \cdot b = 0$, only the $P \cdot b = 3$ and 4 rows are used.

$P \cdot b$	$b^2 = 18$	$b^2 = 20$	$b^2 = 25$	$b^2 = 32$
0	lin. $[16, 25]$ quad. $(9, 16, 25)$	lin. $[16, 25]$ quad. $(9, 16, 25)$	**data**	lin. $[25, 36]$ quad. $(16, 25, 36)$
3	**data**	lin. $[18, 45]$	lin. $[18, 45]$	lin. $[18, 45]$
4	lin. $[17, 20]$ quad. $(17, 20, 32)$	**data**	lin. $[20, 32]$ quad. $(17, 20, 32)$	**data**

Table 17: Origin of each grid value used in Figure 75. "Quad. (b_1^2, b_2^2, b_3^2)" means the parabola in b^2 through the data at those three points on the same $P \cdot b$ row; "lin. $[b_a^2, b_b^2]$" means linear interpolation in b^2 between the data at b_a^2 and b_b^2. Where a cell has two entries, the upper one gives the linear points (circles) and the lower one the quadratic points (triangles) of Figure 75. The pattern is the same for all three amplitudes.

From Figure 75, in certain region $P \cdot b = 0$, the ratio $T(P \cdot b, b^2) = \frac{\widetilde{A}(P \cdot b, b^2)}{\widetilde{A}(P \cdot b = 4, b^2)}$ is flat but for $P \cdot b = 3$, the ratio T is increasing monotonically. The same behavior is captured in both linear and quadratic interpolation. The other ratio $B(P \cdot b, b^2) = \frac{\widetilde{A}(P \cdot b, b^2)}{\widetilde{A}(P \cdot b, b^2 = 18)}$ is not flat, and there exists a cut in continuity at $P \cdot b = 3$ which



Figure 75: Factorization ratio test at $\eta|v|/a = 8$, $P_L = -1$ ($\hat{\zeta} = 0.090772$) for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$ and $\widetilde{A}^{\mathrm{Re}}_{12B}$. Top row: $T(P \cdot b, b^2)$ of Eq. (6.28), the ratio to the $P \cdot b = 4$ row, plotted against b^2. Bottom row: $B(P \cdot b, b^2)$ of Eq. (6.29), the ratio to $b^2 = 18$, plotted against $P \cdot b$. Under factorization, every set of points is flat. Filled circles use linear interpolation in b^2 on all three rows. Open triangles use quadratic interpolation on the $P \cdot b = 0$ and 4 rows and linear interpolation on the $P \cdot b = 3$ row (Table 17). A triangle is omitted where both schemes give the same value, which happens for every point that involves only the $P \cdot b = 3$ row and data points. Error bars are Jackknife errors and do not include any uncertainty from the interpolation.

could also arise from systematic uncertainty from interpolation and limitation of data.

The ratio test is thus only a qualitative check. Where the interpolated values rest on closely spaced data, the $P \cdot b = 0$ and 4 rows are compatible with factorization, but the sampling is too sparse to settle the question. We therefore test it below with explicit factorized fits, using the global simultaneous fit of Section 6.6 as the benchmark.

6.7.2 Rejection of Separable Longitudinal and Transverse Factorization

Because the ratio test cannot settle the question, we fit explicitly factorized models and compare them with the selected power-law fit. We constructed candidate models that multiply an independent longitudinal ($P \cdot b$) profile by various transverse b^2 profiles, including combinations of Gaussian, Lorentzian power-law, and exponential-decay forms, and evaluated each with full Jackknife fits.

The fit results are summarized in Figure 76 (see its caption for the poor χ^2/dof when exponential decay in $(P \cdot b)$ is paired with a Gaussian transverse profile). Even the better combinations give $\chi^2/\text{dof} \approx 2$, and the coefficient c governing $(P \cdot b)$ comes out negative and consistent with zero within Jackknife uncertainties. Such a coefficient is unstable and makes the Fourier transform diverge.

(b_L, b_T, η) Fits for Re $\tilde{A}_{2B}$ for P_L=−1
with $b \geq 3$ and $6 \geq \eta \geq 10$

Model Expression	Params List	Fitted Values	χ^2/DoF
$a\, e^{-f\eta}\, e^{-c\,\lvert b_L P_L\rvert}\, e^{-d\sqrt{b_L^2+b_T^2}}$	{a, c, d, f}	{⟨8.02(56)⟩$_{2902\,J}$, ⟨−0.0021(76)⟩$_{2902\,J}$, ⟨0.579(13)⟩$_{2902\,J}$, ⟨0.5348(84)⟩$_{2902\,J}$}	2.15678
$a\, e^{-f\eta}\, e^{-c\,\lvert b_L P_L\rvert}\, e^{-d\left(b_L^2+b_T^2\right)}$	{a, c, d, f}	{⟨2.08(13)⟩$_{2902\,J}$, ⟨0.0199(73)⟩$_{2902\,J}$, ⟨0.0564(15)⟩$_{2902\,J}$, ⟨0.5364(86)⟩$_{2902\,J}$}	4.98308
$\frac{a\, e^{-f\eta}\, e^{-c\,\lvert b_L P_L\rvert}}{\left(1+d\left(b_L^2+b_T^2\right)\right)^{j2}}$	{a, c, d, f, j2}	{⟨5.28(60)⟩$_{2902\,J}$, ⟨−0.0089(84)⟩$_{2902\,J}$, ⟨0.094(16)⟩$_{2902\,J}$, ⟨0.5344(84)⟩$_{2902\,J}$, ⟨2.07(13)⟩$_{2902\,J}$}	1.90543
$a\, e^{-f\eta}\, e^{-c\,(b_L P_L)^2}\, e^{-d\sqrt{b_L^2+b_T^2}}$	{a, c, d, f}	{⟨8.18(58)⟩$_{2902\,J}$, ⟨−0.0018(17)⟩$_{2902\,J}$, ⟨0.587(14)⟩$_{2902\,J}$, ⟨0.5348(85)⟩$_{2902\,J}$}	2.12445
$\frac{a\, e^{-f\eta}\, e^{-c\,(b_L P_L)^2}}{\left(1+d\left(b_L^2+b_T^2\right)\right)^{j2}}$	{a, c, d, f, j2}	{⟨4.99(50)⟩$_{2902\,J}$, ⟨−0.0015(16)⟩$_{2902\,J}$, ⟨0.085(13)⟩$_{2902\,J}$, ⟨0.5343(84)⟩$_{2902\,J}$, ⟨2.14(14)⟩$_{2902\,J}$}	1.90851

Figure 76: Fits testing the separability hypothesis. Pairing an exponential decay in $(P \cdot b)$ with a Gaussian transverse profile gives a poor fit ($\chi^2/\text{dof} \approx 4.98$). In models with a lower χ^2/dof, the coefficient c consistently fits to zero within uncertainties and is negative in each Jackknife set, so the data reject the separability hypothesis.

We also tested the separability hypothesis with an η-dependent c, written $c(\eta)$, as the coefficient governing $(P \cdot b)$; this forces c to be positive in the asymptotic limit. The asymptotic $c(\eta)$ lowers the χ^2/dof to ≈ 1.35, with c positive in the asymptotic limit (Figure 77). These fits show that the lattice data prefer a power-law in $(P \cdot b)$ rather than an exponential decay in $|(P \cdot b)|$.

(b_L, b_T, η) Fits for Re $\widetilde{A}_{2B}$ for P_L=-1
with $b \geq 3$ and $6 \geq \eta \geq 10$

Model Expression	Params List	Fitted Values	χ^2/DoF
$a\, e^{-f\eta}\, e^{-c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\lvert b_L P_L\rvert}\, e^{-d\sqrt{b_L^2+b_T^2}}$	{a, c, k1, k2, d, f}	{⟨5.28(42)⟩2902 J, ⟨0.51(12)⟩2902 J, ⟨11.60(44)⟩2902 J, ⟨-32.4(27)⟩2902 J, ⟨0.580(13)⟩2902 J, ⟨0.473(11)⟩2902 J}	1.57561
$a\, e^{-f\eta}\, e^{-c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\lvert b_L P_L\rvert}\, e^{-d\,(b_L^2+b_T^2)}$	{a, c, k1, k2, d, f}	{⟨1.36(10)⟩2902 J, ⟨0.53(13)⟩2902 J, ⟨11.09(49)⟩2902 J, ⟨-30.8(28)⟩2902 J, ⟨0.0564(15)⟩2902 J, ⟨0.473(12)⟩2902 J}	4.38385
$\frac{a\, e^{-f\eta}\, e^{-c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)\lvert b_L P_L\rvert}}{(1+d\,(b_L^2+b_T^2))^{j_2}}$	{a, c, k1, k2, d, f, j2}	{⟨3.53(47)⟩2902 J, ⟨0.49(12)⟩2902 J, ⟨11.79(45)⟩2902 J, ⟨-33.0(26)⟩2902 J, ⟨0.096(16)⟩2902 J, ⟨0.473(11)⟩2902 J, ⟨2.06(12)⟩2902 J}	1.3183
$a\, e^{-f\eta}\, e^{-c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)(b_L P_L)^2}\, e^{-d\sqrt{b_L^2+b_T^2}}$	{a, c, k1, k2, d, f}	{⟨5.94(48)⟩2902 J, ⟨0.114(30)⟩2902 J, ⟨11.96(44)⟩2902 J, ⟨-34.3(26)⟩2902 J, ⟨0.588(14)⟩2902 J, ⟨0.487(10)⟩2902 J}	1.54574
$\frac{a\, e^{-f\eta}\, e^{-c\left(1-\frac{k_1}{\eta}-\frac{k_2}{\eta^2}\right)(b_L P_L)^2}}{(1+d\,(b_L^2+b_T^2))^{j_2}}$	{a, c, k1, k2, d, f, j2}	{⟨3.68(38)⟩2902 J, ⟨0.110(29)⟩2902 J, ⟨12.02(44)⟩2902 J, ⟨-34.8(26)⟩2902 J, ⟨0.084(13)⟩2902 J, ⟨0.4896(98)⟩2902 J, ⟨2.16(14)⟩2902 J}	1.35274

Figure 77: Fits testing the separability hypothesis with η-dependent c, written $c(\eta)$. In models with a lower χ^2/dof, the coefficient c is now positive.

We studied four factorized ansatze with lower χ^2/dof and applied the same ansatze to all three invariant amplitudes ($\widetilde{A}^{\rm Re}_{2B}$, $\widetilde{A}^{\rm Im}_{2B}$, and $\widetilde{A}^{\rm Re}_{12B}$). These models combine either an exponential ($e^{-c|P\cdot b|}$) or Gaussian ($e^{-c(P\cdot b)^2}$) longitudinal profile with either an exponential ($e^{-d|b|}$) or power-law ($(1+db^2)^{-j}$) transverse envelope:

1. `ExpPb_Expb`:

$$\widetilde{A}^{\rm Re}_{2B} = a_{\rm re}\, e^{-c_{\rm re}|P\cdot b|-d_{\rm re}|b|}\, e^{-f\eta}\ , \tag{6.30}$$

$$\widetilde{A}^{\rm Im}_{2B} = a_{\rm im}\, (P\cdot b)\, e^{-c_{\rm im}|P\cdot b|-d_{\rm im}|b|}\, e^{-f\eta}\ , \tag{6.31}$$

$$\widetilde{A}^{\rm Re}_{12B} = -a_{\rm reA12B}\, e^{-c_{\rm reA12B}|P\cdot b|-d_{\rm reA12B}|b|}\, e^{-f\eta}\ . \tag{6.32}$$

2. `ExpPb_PowerLawb`:

$$\widetilde{A}^{\rm Re}_{2B} = \frac{a_{\rm re}\, e^{-c_{\rm re}|P\cdot b|-f\eta}}{(1+d_{\rm re}b^2)^{j_{\rm re}}}\ , \tag{6.33}$$

$$\widetilde{A}^{\rm Im}_{2B} = \frac{a_{\rm im}\, (P\cdot b)\, e^{-c_{\rm im}|P\cdot b|-f\eta}}{(1+d_{\rm im}b^2)^{j_{\rm im}}}\ , \tag{6.34}$$

$$\widetilde{A}^{\rm Re}_{12B} = \frac{-a_{\rm reA12B}\, e^{-c_{\rm reA12B}|P\cdot b|-f\eta}}{(1+d_{\rm reA12B}b^2)^{j_{\rm reA12B}}}\ . \tag{6.35}$$

3. `GaussianPb_Expb`:

$$\widetilde{A}^{\rm Re}_{2B} = a_{\rm re}\, e^{-c_{\rm re}(P\cdot b)^2-d_{\rm re}|b|}\, e^{-f\eta}\ , \tag{6.36}$$

$$\widetilde{A}^{\rm Im}_{2B} = a_{\rm im}\, (P\cdot b)\, e^{-c_{\rm im}(P\cdot b)^2-d_{\rm im}|b|}\, e^{-f\eta}\ , \tag{6.37}$$

$$\widetilde{A}^{\rm Re}_{12B} = -a_{\rm reA12B}\, e^{-c_{\rm reA12B}(P\cdot b)^2-d_{\rm reA12B}|b|}\, e^{-f\eta}\ . \tag{6.38}$$

4. `GaussianPb_PowerLawb`:

$$\widetilde{A}^{\mathrm{Re}}_{2B} = \frac{a_{\mathrm{re}}\, e^{-c_{\mathrm{re}}(P\cdot b)^2 - f\eta}}{(1 + d_{\mathrm{re}} b^2)^{j_{\mathrm{re}}}} \ , \tag{6.39}$$

$$\widetilde{A}^{\mathrm{Im}}_{2B} = \frac{a_{\mathrm{im}}\, (P\cdot b)\, e^{-c_{\mathrm{im}}(P\cdot b)^2 - f\eta}}{(1 + d_{\mathrm{im}} b^2)^{j_{\mathrm{im}}}} \ , \tag{6.40}$$

$$\widetilde{A}^{\mathrm{Re}}_{12B} = \frac{-a_{\mathrm{reA12B}}\, e^{-c_{\mathrm{reA12B}}(P\cdot b)^2 - f\eta}}{(1 + d_{\mathrm{reA12B}} b^2)^{j_{\mathrm{reA12B}}}} \ . \tag{6.41}$$

Each model has a closed-form Fourier transform, so the resulting x-space distributions are easy to inspect. We performed global simultaneous Jackknife fits on these four ansatze using the same kinematic cuts ($|\boldsymbol{b}| \geq 3a$, $\eta \in [6, 10]$) and shared area-law factor $e^{-f\eta}$. The fit quality and extracted parameters are summarized in Table 18.

Table 18: Comparison of the four factorized longitudinal-transverse models across nucleon momenta P_L, with extracted parameters shown at $P_L = -1$. While models with power-law transverse envelopes describe $\widetilde{A}^{\mathrm{Re}}_{2B}$ passably, all four factorized ansatze fail on $\widetilde{A}^{\mathrm{Im}}_{2B}$ and $\widetilde{A}^{\mathrm{Re}}_{12B}$.

Model Ansatz	**χ^2/dof Across Momenta**				**Key Fitted Parameters at $P_L = -1$**			**Primary Failure Mode**
	$P_L = -1$	$P_L = -2$	$P_L = -3$	$P_L = -4$	c_{im}	c_{reA12B}	a_{im}	
`ExpPb_Expb`	1.406(92)	0.956(75)	0.622(61)	0.455(53)	0.0010(262)	0.045(23)	−0.167(16)	$c_{\mathrm{im}} \to 0$; unphysical sign on a_{im}.
`ExpPb_PowerLawb`	1.286(89)	0.938(75)	0.623(61)	0.447(53)	0.0045(248)	0.028(24)	−0.167(31)	Fits $\widetilde{A}^{\mathrm{Re}}_{2B}$ well, but c_{im} and c_{reA12B} vanish within error; flips sign of a_{im}.
`GaussianPb_Expb`	2.389(121)	2.005(111)	0.945(76)	0.453(52)	0.568(171)	0.0038(69)	+0.050(121)	Poor χ^2/dof; c_{reA12B} pinned near zero; Gaussian over-damping at large x.
`GaussianPb_PowerLawb`	2.316(120)	1.981(111)	0.948(76)	0.448(52)	0.602(48)	0.0016(61)	+0.168(92)	Poor global fit at low momenta; $c_{\mathrm{reA12B}} \approx 0$; unphysical truncation of valence tails.
Power-law in $((P\cdot b), b^2)$	1.85(10)	1.197(85)	0.712(66)	0.434(51)	**0.2253(69)**	**0.0773(46)**	**+0.267(13)**	**All parameters stable, positive, and well-constrained across all amplitudes.**

The numerical results show several ways in which the factorized models fail. In `ExpPb_PowerLawb`, the real unpolarized amplitude fits with $\chi^2/\mathrm{dof} \approx 1.286$. Because $\widetilde{A}^{\mathrm{Re}}_{2B}$ dominates the statistical weight, the fit quality alone can make the model look adequate. The same parameterization fails for the imaginary amplitude $\widetilde{A}^{\mathrm{Im}}_{2B}$, however: the longitudinal decay width $c_{\mathrm{im}} = 0.0045 \pm 0.0248$ is zero within errors. The model then loses its longitudinal decay profile and gives an essentially flat distribution in $P \cdot b$. The overall amplitude also flips to an unphysical negative sign ($a_{\mathrm{im}} = -0.167$).

For the real Sivers amplitude $\widetilde{A}^{\mathrm{Re}}_{12B}$, the longitudinal parameter c_{reA12B} is similarly suppressed across all factorized models (0.028 ± 0.024 for `ExpPb_PowerLawb` and 0.0016 ± 0.0061 for `GaussianPb_PowerLawb`). The factorized models cannot capture the spin-dependent correlator's distinct longitudinal curvature without forcing the decay parameter to its lower boundary. Factoring out an exponential $|P \cdot b|$ dependence introduces a discontinuous first derivative (a cusp) at $P \cdot b = 0$. For an odd-parity amplitude like $\widetilde{A}^{\mathrm{Im}}_{2B}$, which must vanish linearly through the origin ($\widetilde{A}^{\mathrm{Im}}_{2B} \propto b_L$), the non-analytic factor $e^{-c|P\cdot b|}$ conflicts with the smooth behavior of the Euclidean correlation function. The fit responds by driving $c_{\mathrm{im}} \to 0$ to minimize the cusp penalty.

In the Lorentz-invariant variable space $(P \cdot b, b^2)$, the longitudinal separation $b_L = (P \cdot b)/(-P_L)$ and transverse separation $b_T = \sqrt{b^2 - b_L^2}$ are geometrically coupled. Factorized models artificially decouple $P \cdot b$ and b^2. In contrast, coupling $(P \cdot b)^2$ and b^2 within a single denominator,

$$D = \left[1 + c(\eta)\frac{(P \cdot b)^2}{P_L^2} + d\, b^2\right]^j \ , \tag{6.42}$$

preserves smooth analyticity at $P \cdot b = 0$, satisfies power-law integrability, and yields stable, positive parameters across all three amplitudes, matching the values already obtained from the global fit above (e.g., $c_{\mathrm{im}} = 0.2253(69)$, $c_{\mathrm{reA12B}} = 0.0773(46)$, $a_{\mathrm{im}} = +0.267(13)$). This confirms that the coupled power-law structure adopted for the global fit is preferred over the four explicit factorizations of $(P \cdot b)$ and b^2 tested above.

In the next section, we extrapolate the lattice calculation modeled by the adopted power-law ansatz to $\hat{\zeta} \to \infty$.

7 EXTRAPOLATIONS TO THE PHYSICAL LIMIT

Section 7.1 shows that the amplitudes, the TMD combinations, and the Sivers shift change systematically with $1/\hat{\zeta}$ across the four simulated momenta. Section 7.2 extrapolates the fit parameters to $1/\hat{\zeta} \to 0$ using $P_L = -2, -3, -4$, reconstructs the amplitudes and the Sivers shift in this limit, and lists the numerical Fourier-transform integrals over x at $|b_T| = 3a$. Section 7.3 tests whether the extrapolated Sivers shift and amplitudes factorize in x and $|b_T|$.

7.1 Motivation for the $\hat{\zeta} \to \infty$ Extrapolation

Section 6 established stable global simultaneous fits to the invariant amplitudes $\widetilde{A}_{2B}$ and $\widetilde{A}_{12B}$ at each of our four simulated nucleon momenta, corresponding to the finite Collins-Soper parameters $\hat{\zeta} = 0.090772$, 0.294367, 0.539684, 0.785756 for $P_L = -1, -2, -3, -4$ respectively (Table 2). As discussed in Section 5.2, these finite-$\hat{\zeta}$ results follow from the space-like staple direction that our Euclidean lattice geometry requires; the physical light-cone TMDs are recovered only in the limit $\hat{\zeta} \to \infty$. To compare our lattice results with phenomenological extractions of the Sivers function, we must therefore extrapolate each finite-momentum result to this physical limit.

Figure 78 shows why we need this extrapolation. At fixed transverse quark separation $|b_T| = 3a \approx 0.34$ fm, we evaluate the reconstructed amplitudes $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$ and $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, the corresponding unpolarized and Sivers TMD combinations $\tilde{f}_1^{(0)} = 2\widetilde{A}_{2B}^{[x]}$ and $\tilde{f}_{1T}^{\perp(1)} = -2\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, and the resulting Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$, as continuous functions of x at each of our four simulated momenta. Plotted against $1/\hat{\zeta}$, all six quantities vary systematically with the momentum. None of our finite lattice momenta has reached the physical limit, so the results can be interpreted physically only after the extrapolation to $1/\hat{\zeta} \to 0$ carried out in the rest of this chapter.



Figure 78: Discrete $\hat{\zeta}$-slices of the reconstructed amplitudes $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$ and $\widetilde{A}^{[x]\mathrm{Im}}_{2B}$ and the unpolarized and Sivers TMD combinations $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$ and $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$, plotted against both x and $1/\hat{\zeta}$ at fixed transverse separation $|b_T| = 3a \approx 0.34$ fm. Each ribbon corresponds to one of the four simulated momenta $P_L = -1, -2, -3, -4$ (equivalently $1/\hat{\zeta} = 11.02,\ 3.40,\ 1.85,\ 1.27$). The systematic drift of every panel with $1/\hat{\zeta}$ motivates the extrapolation to the physical light-cone limit $\hat{\zeta} \to \infty$ performed below.



Figure 79: Sivers shift $\langle k_y \rangle_{TU}$ and $x\langle k_y \rangle_{TU}$ for the same four momenta and the same $|b_T|$ as Fig. 78, restricted to the quark region $x \geq 0$.

7.2 Extrapolation at fit-parameters level

The lowest simulated momentum, $P_L = -1$ ($\hat{\zeta} = 0.090772$), sits too close to the origin in $1/\hat{\zeta}$ to resolve the curvature required to anchor an extrapolation; including it risks biasing the fit away from the true asymptotic ($\hat{\zeta} \to \infty$) behavior rather than constraining it. We therefore exclude $P_L = -1$ and extrapolate using only the three higher, better-separated momenta $P_L = -2, -3, -4$.

Rather than extrapolate the amplitudes point by point in x, we extrapolate each of the twelve parameters of the global simultaneous-fit models directly: the four parameters $\{a, c, d, j\}$ appearing in each of the power-law models for $\widetilde{A}^{\mathrm{Re}}_{2B}$, $\widetilde{A}^{\mathrm{Im}}_{2B}$, and $\widetilde{A}^{\mathrm{Re}}_{12B}$ (Eq. (6.16)-(6.18)). In these models c multiplies $b_L^2 = (P \cdot b)^2/P_L^2$ and a_{im} multiplies $b_L = (P \cdot b)/(-P_L)$, so the parameters c_{re}, c_{im}, and c_{reA12B} are first rescaled by $1/P_L^2$ and a_{im} by $1/P_L$. After rescaling, they multiply powers of the invariant $P \cdot b$ directly (up to an overall sign for a_{im}), so all three momenta can be fit on a common footing. For the resulting $\hat{\zeta}$-dependence of each rescaled parameter, we tested two functional forms,

$$p(\hat{\zeta}) = c + \frac{d}{\hat{\zeta}} \qquad \text{and} \qquad p(\hat{\zeta}) = c + \frac{d}{\hat{\zeta}^2}, \tag{7.1}$$

using correlated Jackknife fits to the three data points. The quadratic ansatz consistently gave a lower χ^2/dof than the linear form across the twelve parameters, and we adopt it for every extrapolation shown below. The extrapolated, physical-limit value of each parameter is its fitted intercept c at $1/\hat{\zeta}^2 = 0$.

Reconstructing the amplitudes from these twelve extrapolated parameters gives the unpolarized and Sivers TMD combinations directly at the physical limit, shown in Fig. 81 as continuous functions of x at fixed $|b_T| = 3a \approx 0.34$ fm.

Table 19 lists the corresponding numerical Fourier-transform integrals at $|b_T| = 3a \approx 0.34$ fm, evaluated over the full range $x \in \{-\infty, \infty\}$ as well as the restricted ranges $\{-1, 1\}$, $\{-1, 0\}$, and $\{0, 1\}$ used to isolate, respectively, the quark-plus-antiquark, antiquark, and quark contributions to the x-integrals $\int dx\, \tilde{f}_1^{(0)}$ and $\int dx\, \tilde{f}_{1T}^{\perp(1)}$.

Restricting the integration to $x \in \{-1, 0\}$ or $\{0, 1\}$ isolates the antiquark and quark contributions respectively; because $\widetilde{A}^{[x]\mathrm{Im}}_{2B}$ is an odd function of x while $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$ and $\widetilde{A}^{[x]\mathrm{Re}}_{12B}$ are even, the antiquark and quark ranges



Figure 80: Extrapolation of the twelve simultaneous-fit parameters to the physical limit $\hat{\zeta} \to \infty$, using only the three higher momenta $P_L = -2, -3, -4$ (black points, plotted against $1/\hat{\zeta}$). The dashed red curve and shaded band show the fitted ansatz $c + d/\hat{\zeta}^2$ of Eq. 7.1, with each panel's Jackknife χ^2/dof noted in its title; the blue square at $1/\hat{\zeta} = 0$ marks the extrapolated physical-limit value. The parameters $c_{\rm re}$, $c_{\rm im}$, and $c_{\rm reA12B}$ are shown rescaled by $1/P_L^2$, and $a_{\rm im}$ by $1/P_L$, as described in the text.

Table 19: Numerical integration of the Fourier-transform integrals at $|b_T| = 3a \approx 0.34$ fm, evaluated using the extrapolated ($\hat{\zeta} \to \infty$) amplitudes of Fig. 81, over the full range and three restricted x-ranges isolating the quark, antiquark, and combined contributions.

Integral	$\{-\infty, \infty\}$	$\{-1, 1\}$	$\{-1, 0\}$	$\{0, 1\}$
$\int dx\, \widetilde{A}_{2B}^{[x]\rm Re}$	0.751(79)	0.751(80)	0.375(40)	0.375(40)
$\int dx\, (\widetilde{A}_{2B}^{[x]\rm Im})$	$6(96) \times 10^{-17}$	$3(24) \times 10^{-16}$	0.165(27)	−0.169(29)
$\int dx\, \tilde{f}_1^{(0)} = 2 \int dx\, (\widetilde{A}_{2B}^{[x]\rm Re} - \widetilde{A}_{2B}^{[x]\rm Im})$	1.50(16)	1.50(16)	0.421(81)	1.09(11)
$\int dx\, \tilde{f}_{1T}^{\perp(1)} = -2 \int dx\, \widetilde{A}_{12B}^{[x]\rm Re}$	0.302(53)	0.302(53)	0.151(27)	0.151(27)
$m_N \dfrac{\int dx\, \tilde{f}_{1T}^{\perp(1)}}{\int dx\, \tilde{f}_1^{(0)}}$	0.217(38)	0.217(38)	0.387(89)	0.150(26)

give equal $\int dx\, \widetilde{A}_{2B}^{[x]\rm Re}$ and $\int dx\, \widetilde{A}_{12B}^{[x]\rm Re}$ but opposite-sign $\int dx\, (i\widetilde{A}_{2B}^{[x]\rm Im})$, so the combined $\{-1, 1\}$ integral of the imaginary term is consistent with zero within uncertainty. The integrals over the full range $\{-\infty, \infty\}$ agree



Figure 81: Unpolarized and Sivers TMD amplitudes reconstructed from the extrapolated ($\hat{\zeta} \to \infty$) fit parameters of Fig. 80, evaluated as continuous functions of x at fixed transverse separation $|b_T| = 3a \approx 0.34$ fm: $\widetilde{A}^{[x]\mathrm{Re}}_{2B}$, $\widetilde{A}^{[x]\mathrm{Im}}_{2B}$, the full unpolarized combination $\tilde{f}^{(0)}_1 = 2\widetilde{A}^{[x]}_{2B}$, and the Sivers combination $\tilde{f}^{\perp(1)}_{1T} = -2\widetilde{A}^{[x]\mathrm{Re}}_{12B}$. Shaded bands are the Jackknife uncertainty propagated from the twelve extrapolated parameters.



Figure 82: Sivers shift $\langle k_y\rangle_{TU}$ and $x\langle k_y\rangle_{TU}$ built from the extrapolated amplitudes of Fig. 81, restricted to the quark region $x \geq 0$. Shaded bands are the Jackknife uncertainty.

with those over the physically motivated range $\{-1, 1\}$ well within uncertainties. The extrapolated amplitudes therefore decay fast enough in x that truncating at the light-cone-momentum-fraction cutoff is negligible.

Figure 83 extends this reconstruction to the full dependence on both x and transverse separation, while Fig. 85 compares the extrapolated result with each of the three finite momenta used to obtain it.



Figure 83: The four extrapolated ($\hat{\zeta}\to\infty$) amplitude quantities of Fig. 81, shown as surfaces in x and $|b_T|$ (from $3a \approx 0.34$ fm to $\sqrt{68}\,a \approx 0.94$ fm). The joint x-$|b_T|$ dependence shows how the reconstructed amplitudes change with transverse quark separation at the physical limit.



Figure 84: The extrapolated Sivers shift $\langle k_y\rangle_{TU}$ and $x\langle k_y\rangle_{TU}$ of Fig. 82, shown as surfaces in $x \geq 0$ and $|b_T|$ over the same range.



Figure 85: Comparison, at fixed $|b_T| = 3a \approx 0.34$ fm, between the four amplitude quantities evaluated at each of the four simulated momenta ($P_L = -1, -2, -3, -4$, colored curves, each using its own unextrapolated fit parameters) and the extrapolated $\hat{\zeta} \to \infty$ result (black dashed curve), which uses only $P_L = -2, -3, -4$. The extrapolated curve differs from all three finite-momentum curves used in the extrapolation, including the one at the largest simulated momentum, so the fit-parameter extrapolation of Fig. 80 captures real momentum dependence and does not just reproduce the $P_L = -4$ result.



Figure 86: The same comparison as Fig. 85 for the Sivers shift $\langle k_y\rangle_{TU}$ and $x\langle k_y\rangle_{TU}$ ($x \geq 0$).

7.3 Testing Factorization in x and $|b_T|$ at the Physical Limit

Section 6.7 tested factorization of the position-space amplitudes in $(P \cdot b)$ and b^2. Here we ask a different question: whether the extrapolated distributions factorize in the momentum fraction x and the transverse separation $|b_T|$.

Figures 83 and 85 show that the extrapolated Sivers shift and amplitudes depend on both x and $|b_T|$ jointly. We ask whether this joint dependence is separable, that is, whether the extrapolated Sivers shift factorizes as

$$\langle k_y \rangle_{TU}(x, |b_T|) \stackrel{?}{=} F(x)\, G(|b_T|) \ , \tag{7.2}$$

with all of the x-shape carried by $F(x)$ and all of the $|b_T|$-shape by $G(|b_T|)$. Such a factorization is not automatic in our fit ansatz. Each amplitude enters only through the combination $d_X(b^2) = 1 + d_X\, b^2$ inside a Bessel function that also carries the x-dependence,

$$\widetilde{A}_X^{[x]}(x, b^2) \ \propto \ N_X(b^2)\ |x|^{-1/2+j_X}\ K_{\nu_X}\Big(|x|\sqrt{d_X(b^2)/c_X}\Big) \ , \tag{7.3}$$

so $|b_T|$ rescales the argument of the same Bessel function that carries the x-dependence, rather than multiplying it by an overall x-independent prefactor. The Sivers shift is a ratio of three such amplitudes ($\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, $\widetilde{A}_{12B}^{[x]\mathrm{Re}}$), each with its own independently fitted (j_X, d_X); an exact factorization would require these parameters to coincide across the three amplitudes so that the x- and $|b_T|$-dependence separates. Whether the extrapolated fit comes close enough to that coincidence is a numerical question, which we answer below.

Section 7.3.1 defines the relative distance from a factorized form, and Sections 7.3.2 and 7.3.3 apply it to the Sivers shift and to the individual amplitudes.

7.3.1 Quantifying the Distance from a Factorized Form

Given a function of two variables $f(x, |b_T|)$, we quantify how far it is from a factorized form $g(x)h(|b_T|)$ by finding the closest such pair in the least-squares sense and reporting the residual. Explicitly, we minimize

$$J = \int dx\, d|b_T|\, \big(f(x, |b_T|) - g(x)h(|b_T|)\big)^2 \tag{7.4}$$

over all functions g and h. Setting the functional derivatives with respect to $g(x)$ and $h(|b_T|)$ to zero gives

$$0 = \frac{\delta J}{\delta g(x)} = \int d|b_T|\, f(x,|b_T|)\, h(|b_T|) - g(x) \int d|b_T|\, h^2(|b_T|) \ , \tag{7.5}$$

$$0 = \frac{\delta J}{\delta h(|b_T|)} = \int dx\, f(x,|b_T|)\, g(x) - h(|b_T|) \int dx\, g^2(x) \ , \tag{7.6}$$

which can be solved for h in terms of g,

$$h(|b_T|) = \frac{\int dx\, f(x,|b_T|)\, g(x)}{\int dx\, g^2(x)} \ , \tag{7.7}$$

and reinserted to eliminate h, giving a closed, self-consistent equation for g alone:

$$g(x) = \left[\int d|b_T| \left[\frac{\int dx'\, f(x',|b_T|)\, g(x')}{\int dx'\, g^2(x')} \right]^2 \right]^{-1} \int d|b_T|\, f(x,|b_T|)\, \frac{\int dx'\, f(x',|b_T|)\, g(x')}{\int dx'\, g^2(x')} \ . \tag{7.8}$$

We solve Eq. (7.8) iteratively. Starting from an initial guess $g = 1$, we evaluate the right-hand side to obtain a refined g, reinsert it, and repeat until g (and the resulting relative distance below) stops changing within a fixed tolerance. For the grids used here, convergence is rapid in practice.

We then compare the residual $\sqrt{J}$ of the optimal separable pair $g(x)h(|b_T|)$ with the overall size of f, $N = \int dx\, d|b_T|\, f^2(x,|b_T|)$, which gives the relative distance from a factorized form:

$$\text{relative distance} = \sqrt{J/N} \ . \tag{7.9}$$

A relative distance of zero indicates exact factorization; the closer to zero, the better $F(x)G(|b_T|)$ approximates the true joint $(x,|b_T|)$ dependence. In practice, we replace the continuum integrals with sums over a discrete $(x,|b_T|)$ grid, apply the iteration of Eq. (7.8) independently to each Jackknife sample of the extrapolated amplitudes, and reduce the resulting distribution of relative distances to a Jackknife mean and error. We repeat the full test in four x-windows, $[0,1]$, $[0.1,0.9]$, $[0.2,0.8]$, and $[0.3,0.8]$, to check that the conclusion is not an artifact of the amplitudes' behavior near $x = 0$.

7.3.2 Factorization Test for the Sivers Shift

The extrapolated fit parameters already suggest, before any distance test, that exact factorization is unlikely. As noted in Section 7.3, factorization requires the Bessel-function order j_X and width parameter d_X to coincide across the three input amplitudes; the extrapolated values are

$$j_{\rm re} = 2.0667(69), \quad j_{\rm im} = 0.9921(92), \quad j_{\rm reA12B} = 2.0026(40) \ , \tag{7.10}$$

$$d_{\rm re} = 0.139(12), \quad d_{\rm im} = 0.183(14), \quad d_{\rm reA12B} = 0.105(18)\ . \tag{7.11}$$

The two j parameters entering the real amplitudes, $j_{\rm re}$ and $j_{\rm reA12B}$, agree with each other, but $j_{\rm im} \approx 1$ differs clearly from both. The three d parameters lie closer together but are still resolved from one another well outside their Jackknife uncertainties. This mismatch in the parameters, especially in j, gives a separate reason to expect a nonzero residual from exact factorization.

Table 20 lists the resulting relative distance of the extrapolated Sivers shift $\langle k_y \rangle_{TU}(x, |b_T|)$ from its best separable approximation, over $|b_T| \in [0.3421, 0.9403]$ fm (60 points, fixed across all four x-windows).

Table 20: Relative distance of the extrapolated Sivers shift $\langle k_y \rangle_{TU}(x, |b_T|)$ from its optimal factorized approximation $F(x)G(|b_T|)$, Eq. (7.9), over $|b_T| \in [0.3421, 0.9403]$ fm for four choices of x-window.

x-window	Relative distance
$[0.0, 1.0]$	0.0568(78)
$[0.1, 0.9]$	0.0714(95)
$[0.2, 0.8]$	0.0758(75)
$[0.3, 0.8]$	0.0620(88)

In every window the relative distance is at the level of 5.7-7.6% and nonzero well outside its Jackknife uncertainty. The Sivers shift is therefore close to, but not exactly, factorized. The residual stays stable as the x-window narrows away from $x = 0$, so behavior at the endpoint does not drive it. Figure 87 shows the optimal separable factors $g(x)$ and $h(|b_T|)$ for all four windows and compares the $g(x)h(|b_T|)$ reconstruction with the actual data at the two extreme $|b_T|$ values in the fit range.

Ratios of the Sivers shift at fixed x or fixed $|b_T|$ show the same non-factorization more directly. If $\langle k_y \rangle_{TU}(x, |b_T|) = F(x)G(|b_T|)$ exactly, then $R_b(x) \equiv \langle k_y \rangle_{TU}(x, |b_T|)/\langle k_y \rangle_{TU}(x, |b_T|_{\rm ref}) = G(|b_T|)/G(|b_T|_{\rm ref})$ must be flat in x for every choice of $|b_T|$ and reference $|b_T|_{\rm ref}$, and likewise $R_x(|b_T|) \equiv \langle k_y \rangle_{TU}(x, |b_T|)/\langle k_y \rangle_{TU}(x_{\rm ref}, |b_T|) = F(x)/F(x_{\rm ref})$ must be flat in $|b_T|$ for every x and $x_{\rm ref}$. Requiring this for every pair of references is both necessary and sufficient for factorization, so the ratio test is a complete test rather than a heuristic proxy. Figures 88 and 89 show that neither ratio is flat.

Finally, to check that the $\hat{\zeta} \to \infty$ extrapolation itself does not introduce this non-factorization, we repeat the $R_b(x)$ and $R_x(|b_T|)$ ratio tests on the raw, unextrapolated fit parameters at each finite momentum. Figures 90



Figure 87: Left and center: the optimal separable factors $g(x)$ and $h(|b_T|)$ (Eqs. (7.7)-(7.8), normalized so mean$[g^2] = 1$) for the extrapolated Sivers shift, shown for all four x-windows of Table 20. Right: the reconstructed $g(x)h(|b_T|)$ (dashed) compared to the actual extrapolated Sivers shift (solid) at the smallest and largest fitted $|b_T|$, using the $x \in [0, 1]$ fit; the visible gap between the solid and dashed curves is the residual quantified in Table 20.



Figure 88: $R_b(x) = \langle k_y\rangle_{TU}(x, |b_T|)/\langle k_y\rangle_{TU}(x, 0.34\text{ fm})$ for four values of $|b_T|$, as a function of x. Under exact factorization each curve would be flat. Instead, every curve rises from its $x \to 0$ value to a peak near $x \approx 0.1$ before decaying, so the $|b_T|$-dependence of the Sivers shift itself depends on x.

and 91 show that the same qualitative pattern, non-flat ratios in both x and $|b_T|$, is already present at every simulated momentum. If anything, the pattern becomes more pronounced at the higher momenta.

7.3.3 Factorization Test at the Amplitude Level

The Sivers-shift test above probes factorization of the full ratio $\langle k_y\rangle_{TU} = -m_N \widetilde{A}_{12B}^{[x]\mathrm{Re}}/(\widetilde{A}_{2B}^{[x]\mathrm{Re}} - i\widetilde{A}_{2B}^{[x]\mathrm{Im}})$, in which some of the non-factorized behavior of the individual amplitudes could in principle cancel. We therefore repeat the same distance test of Section 7.3.1 on each extrapolated amplitude separately: $\widetilde{A}_{12B}^{[x]\mathrm{Re}}$, $\widetilde{A}_{2B}^{[x]\mathrm{Re}}$, $i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$, and



Figure 89: $R_x(|b_T|) = \langle k_y \rangle_{TU}(x, |b_T|)/\langle k_y \rangle_{TU}(0.3, |b_T|)$ for four values of x, as a function of $|b_T|$. The $x = 0.2$ curve changes by more than a factor of two across the fitted $|b_T|$ range instead of staying flat, so the x-dependence of the Sivers shift itself depends on $|b_T|$, consistent with Fig. 88.

the combination $\widetilde{A}_{2B}^{[x]\mathrm{Re}} - i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$ (proportional to $\tilde{f}_1^{(0)}$) that appears in the Sivers-shift denominator. Table 21 lists the resulting relative distances, using the same $|b_T| \in [0.3421, 0.9403]$ fm grid and four x-windows as the Sivers-shift test.

Table 21: Relative distance of each extrapolated amplitude (and their combination $\widetilde{A}_{2B}^{[x]\mathrm{Re}} - i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$) from its optimal factorized approximation, over the same $|b_T| \in [0.3421, 0.9403]$ fm grid and x-windows as Table 20.

Amplitude	$x \in [0.0, 1.0]$	$x \in [0.1, 0.9]$	$x \in [0.2, 0.8]$	$x \in [0.3, 0.8]$
$\widetilde{A}_{12B}^{[x]\mathrm{Re}}$	0.0804(18)	0.0573(26)	0.0397(24)	0.0303(21)
$\widetilde{A}_{2B}^{[x]\mathrm{Re}}$	0.08138(65)	0.0718(36)	0.0581(55)	0.0478(56)
$i\widetilde{A}_{2B}^{[x]\mathrm{Im}}$	0.107(18)	0.0868(30)	0.0578(28)	0.0425(23)
$\widetilde{A}_{2B}^{[x]}$	0.1943(63)	0.0926(59)	0.0594(40)	0.0470(47)

Every amplitude has a nonzero relative distance from factorized form in every window. For the widest window, $x \in [0, 1]$, the amplitude-level residuals (8-19%) exceed the corresponding Sivers-ratio residual (5.7%, Table 20), so the ratio structure of the Sivers shift partially, but not completely, cancels the non-factorized behavior of the individual amplitudes. All four amplitude-level residuals shrink as the x-window narrows away from $x = 0$. This is consistent with the Bessel-function singularity structure of each amplitude near the origin,



Figure 90: The same ratio test as Fig. 88, $R_b(x)$, repeated at each finite momentum individually using its own raw (unextrapolated) fit parameters. None of the four panels shows a flat curve, so the non-factorization is a feature of the fitted amplitudes and not an artifact of the $\hat{\zeta} \to \infty$ extrapolation.

which differs between amplitudes because of the parameter mismatch identified in Section 7.3.2.

The Sivers-shift and amplitude-level tests agree. Neither the individual amplitudes nor the Sivers-shift ratio built from them is exactly factorized in x and $|b_T|$, but the deviation from a factorized form is a modest, well-resolved residual and not a qualitative failure. The Sivers shift itself is factorized to within roughly 5.7-7.6% in the relative L_2 sense. This is smaller than the 8-19% residuals of the amplitudes that enter it, which points to a partial cancellation of non-factorized behavior in the ratio. The residual traces back directly to the extrapolated fit parameters: the Bessel-function order $j_{\mathrm{im}} \approx 1$ governing $\widetilde{A}_{2B}^{[x]\mathrm{Im}}$ differs from $j_{\mathrm{re}}, j_{\mathrm{reA12B}} \approx 2$ governing the two real amplitudes, and the three width parameters d_X are likewise resolved from one another well outside their Jackknife uncertainties. We conclude that an exactly factorized ansatz $F(x)G(|b_T|)$ would be a good, but not exact, approximation to the extracted Sivers shift.



Figure 91: The same ratio test as Fig. 89, $R_x(|b_T|)$, repeated at each finite momentum individually. The departure from flatness grows with $|P_L|$, most visibly in the $x = 0.2$ curve, which is consistent with the parameter mismatch identified in the extrapolated j_X and d_X values above.



Figure 92: Optimal separable factor $g(x)$ for each of the four amplitude combinations of Table 21, for all four x-windows (normalized so mean$[g^2] = 1$). Each amplitude's $g(x)$ is dominated by a sharp rise as $x \to 0$, which steepens as the window narrows and reflects the Bessel-function behavior at small x.



Figure 93: Optimal separable factor $h(|b_T|)$ for each of the four amplitude combinations of Table 21, for all four x-windows. For all four amplitudes, $h(|b_T|)$ falls smoothly with increasing $|b_T|$, but its magnitude and slope depend on the x-window used to define the fit, which itself signals imperfect factorization.

8 CONCLUSION AND OUTLOOK



Figure 94: Reconstructed $g(x)h(|b_T|)$ (dashed) against the actual extrapolated amplitude data (solid) at the smallest and largest fitted $|b_T|$, using the $x \in [0, 1]$ fit, for each of the four amplitude combinations of Table 21. The largest reconstruction gap appears near $x \to 0$ for every amplitude, consistent with the $x \in [0, 1]$ column of Table 21 having the largest relative distance in every row.

A QLUA: Gamma $\Gamma(n)$

The binary notation for 16 basis matrices:

$$\Gamma(n) = \gamma_1^{n_0}\gamma_2^{n_1}\gamma_3^{n_2}\gamma_4^{n_3} \tag{A.1}$$

where, the binary representation of $[n]_{10}$ is $[n_3n_2n_1n_0]_2$

$$[n]_{10} = n_3 2^3 + n_2 2^2 + n_1 2^1 + n_0 2^0 \tag{A.2}$$

The below table contains all 16 gamma matrices with binary notations.

$[n]_{10}$	$[n_3 n_2 n_1 n_0]_2$	$\Gamma(n)$
0	0000	$\mathbb{1}$
1	0001	γ_1
2	0010	γ_2
3	0011	$\gamma_1\gamma_2$
4	0100	γ_3
5	0101	$\gamma_1\gamma_3$
6	0110	$\gamma_2\gamma_3$
7	0111	$\gamma_1\gamma_2\gamma_3 = \gamma_5\gamma_4$
8	1000	γ_4
9	1001	$\gamma_1\gamma_4$
10	1010	$\gamma_2\gamma_4$
11	1011	$\gamma_1\gamma_2\gamma_4 = -\gamma_5\gamma_3$
12	1100	$\gamma_3\gamma_4$
13	1101	$\gamma_1\gamma_3\gamma_4 = \gamma_5\gamma_2$
14	1110	$\gamma_2\gamma_3\gamma_4 = -\gamma_5\gamma_1$
15	1111	$\gamma_1\gamma_2\gamma_3\gamma_4 = \gamma_5$

References

[1] B. U. Musch, P. Hagler, M. Engelhardt, J. W. Negele, and A. Schafer, Phys. Rev. D **85**, 094510 (2012), arXiv:1111.4249 [hep-lat] .

[2] H.-W. Lin and H. B. Meyer, eds., Lattice QCD for Nuclear Physics, Lect. Notes Phys., Vol. 889 (Springer, 2015).

[3] R. Hofstadter, Rev. Mod. Phys. **28**, 214 (1956).

[4] E. D. Bloom et al., Phys. Rev. Lett. **23**, 930 (1969).

[5] M. Breidenbach, J. I. Friedman, H. W. Kendall, E. D. Bloom, D. H. Coward, H. C. DeStaebler, J. Drees, L. W. Mo, and R. E. Taylor, Phys. Rev. Lett. **23**, 935 (1969).

[6] J. D. Bjorken, Phys. Rev. **179**, 1547 (1969).

[7] R. P. Feynman, Phys. Rev. Lett. **23**, 1415 (1969).

[8] C. G. Callan, Jr. and D. J. Gross, Phys. Rev. Lett. **22**, 156 (1969).

[9] M. Gell-Mann, Phys. Lett. **8**, 214 (1964).

[10] G. Zweig, "An SU(3) model for strong interaction symmetry and its breaking. Version 2," in DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. VOL. 1. 1964 - 1978, edited by D. B. Lichtenberg and S. P. Rosen (1964) pp. 22–101.

[11] C.-N. Yang and R. L. Mills, Phys. Rev. **96**, 191 (1954).

[12] H. Fritzsch, M. Gell-Mann, and H. Leutwyler, Phys. Lett. B **47**, 365 (1973).

[13] D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30**, 1343 (1973).

[14] H. D. Politzer, Phys. Rev. Lett. **30**, 1346 (1973).

[15] S. D. Drell and T.-M. Yan, Phys. Rev. Lett. **25**, 316 (1970), [Erratum: Phys.Rev.Lett. 25, 902 (1970)].

[16] D. Müller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Hořejši, Fortsch. Phys. **42**, 101 (1994), arXiv:hep-ph/9812448 .

[17] A. V. Radyushkin, Phys. Rev. D **56**, 5524 (1997), arXiv:hep-ph/9704207 .

[18] X.-D. Ji, Phys. Rev. Lett. **78**, 610 (1997), arXiv:hep-ph/9603249 .

[19] M. Diehl, Phys. Rept. **388**, 41 (2003), arXiv:hep-ph/0307382 .

[20] M. Burkardt, Phys. Rev. D **62**, 071503 (2000), [Erratum: Phys.Rev.D 66, 119903 (2002)], arXiv:hep-ph/0005108 .

[21] J. C. Collins and D. E. Soper, Nucl. Phys. B **194**, 445 (1982).

[22] P. J. Mulders and R. D. Tangerman, Nucl. Phys. B **461**, 197 (1996), [Erratum: Nucl.Phys.B 484, 538–540 (1997)], arXiv:hep-ph/9510301 .

[23] J. Collins, Foundations of perturbative QCD, Vol. 32 (Cambridge University Press, 2011).

[24] R. N. Cahn, Phys. Lett. B **78**, 269 (1978).

[25] J. C. Collins and D. E. Soper, Nucl. Phys. B **193**, 381 (1981), [Erratum: Nucl.Phys.B 213, 545 (1983)].

[26] J. C. Collins, D. E. Soper, and G. F. Sterman, Nucl. Phys. B **250**, 199 (1985).

[27] J. P. Ralston and D. E. Soper, Nucl. Phys. B **152**, 109 (1979).

[28] A. Kotzinian, Nucl. Phys. B **441**, 234 (1995), arXiv:hep-ph/9412283 .

[29] R. D. Tangerman and P. J. Mulders, Phys. Rev. D **51**, 3357 (1995), arXiv:hep-ph/9403227 .

[30] X.-D. Ji, J.-P. Ma, and F. Yuan, Phys. Rev. D **71**, 034005 (2005), arXiv:hep-ph/0404183 .

[31] S. M. Aybat and T. C. Rogers, Phys. Rev. D **83**, 114042 (2011), arXiv:1101.5057 [hep-ph] .

[32] A. V. Belitsky, X.-d. Ji, and F. Yuan, Phys. Rev. D **69**, 074014 (2004), arXiv:hep-ph/0307383 .

[33] S. Meissner, A. Metz, and M. Schlegel, JHEP **08**, 056 (2009), arXiv:0906.5323 [hep-ph] .

[34] C. Lorce and B. Pasquini, Phys. Rev. D **84**, 014015 (2011), arXiv:1106.0139 [hep-ph] .

[35] D. Boer et al., (2011), arXiv:1108.1713 [nucl-th] .

[36] A. Accardi et al., Eur. Phys. J. A **52**, 268 (2016), arXiv:1212.1701 [nucl-ex] .

[37] R. Abdul Khalek et al., Nucl. Phys. A **1026**, 122447 (2022), arXiv:2103.05419 [physics.ins-det] .

[38] J. Ashman et al. (European Muon), Phys. Lett. B **206**, 364 (1988).

[39] R. L. Jaffe and A. Manohar, Nucl. Phys. B **337**, 509 (1990).

[40] E. Leader and C. Lorcé, Phys. Rept. **541**, 163 (2014), arXiv:1309.4235 [hep-ph] .

[41] G. Bunce et al., Phys. Rev. Lett. **36**, 1113 (1976).

[42] D. L. Adams et al. (FNAL-E704), Phys. Lett. B **264**, 462 (1991).

[43] G. L. Kane, J. Pumplin, and W. Repko, Phys. Rev. Lett. **41**, 1689 (1978).

[44] D. W. Sivers, Phys. Rev. D **41**, 83 (1990).

[45] A. Bacchetta, U. D'Alesio, M. Diehl, and C. A. Miller, Phys. Rev. D **70**, 117504 (2004), arXiv:hep-ph/0410050 .

[46] J. C. Collins, Nucl. Phys. B **396**, 161 (1993), arXiv:hep-ph/9208213 .

[47] S. J. Brodsky, D. S. Hwang, and I. Schmidt, Phys. Lett. B **530**, 99 (2002), arXiv:hep-ph/0201296 .

[48] J. C. Collins, Phys. Lett. B **536**, 43 (2002), arXiv:hep-ph/0204004 .

[49] S. J. Brodsky, D. S. Hwang, and I. Schmidt, Nucl. Phys. B **642**, 344 (2002), arXiv:hep-ph/0206259 .

[50] A. V. Belitsky, X. Ji, and F. Yuan, Nucl. Phys. B **656**, 165 (2003), arXiv:hep-ph/0208038 .

[51] D. Boer, P. J. Mulders, and F. Pijlman, Nucl. Phys. B **667**, 201 (2003), arXiv:hep-ph/0303034 .

[52] Z.-B. Kang and J.-W. Qiu, Phys. Rev. Lett. **103**, 172001 (2009), arXiv:0903.3629 [hep-ph] .

[53] D. Boer and P. J. Mulders, Phys. Rev. D **57**, 5780 (1998), arXiv:hep-ph/9711485 .

[54] M. Burkardt, Phys. Rev. D **66**, 114005 (2002), arXiv:hep-ph/0209179 .

[55] M. Burkardt, Nucl. Phys. A **735**, 185 (2004), arXiv:hep-ph/0302144 .

[56] M. Burkardt, Phys. Rev. D **69**, 091501 (2004), arXiv:hep-ph/0402014 .

[57] J.-w. Qiu and G. F. Sterman, Phys. Rev. Lett. **67**, 2264 (1991).

[58] X. Ji, J.-W. Qiu, W. Vogelsang, and F. Yuan, Phys. Rev. Lett. **97**, 082002 (2006), arXiv:hep-ph/0602239 .

[59] A. Bacchetta, M. Diehl, K. Goeke, A. Metz, P. J. Mulders, and M. Schlegel, JHEP **02**, 093 (2007), arXiv:hep-ph/0611265 .

[60] A. Airapetian et al. (HERMES), Phys. Rev. Lett. **103**, 152002 (2009), arXiv:0906.3918 [hep-ex] .

[61] A. Airapetian et al. (HERMES), JHEP **12**, 010 (2020), arXiv:2007.07755 [hep-ex] .

[62] M. Alekseev et al. (COMPASS), Phys. Lett. B **673**, 127 (2009), arXiv:0802.2160 [hep-ex] .

[63] C. Adolph et al. (COMPASS), Phys. Lett. B **744**, 250 (2015), arXiv:1408.4405 [hep-ex] .

[64] X. Qian et al. (Jefferson Lab Hall A), Phys. Rev. Lett. **107**, 072003 (2011), arXiv:1106.0363 [nucl-ex] .

[65] L. Adamczyk et al. (STAR), Phys. Rev. Lett. **116**, 132301 (2016), arXiv:1511.06003 [nucl-ex] .

[66] M. Aghasyan et al. (COMPASS), Phys. Rev. Lett. **119**, 112002 (2017), arXiv:1704.00488 [hep-ex] .

[67] C. Adolph et al. (COMPASS), Phys. Lett. B **770**, 138 (2017), arXiv:1609.07374 [hep-ex] .

[68] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, and A. Prokudin, Phys. Rev. D **72**, 094007 (2005), [Erratum: Phys.Rev.D 72, 099903 (2005)], arXiv:hep-ph/0507181 .

[69] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, S. Melis, F. Murgia, A. Prokudin, and C. Turk, Eur. Phys. J. A **39**, 89 (2009), arXiv:0805.2677 [hep-ph] .

[70] M. G. Echevarria, A. Idilbi, Z.-B. Kang, and I. Vitev, Phys. Rev. D **89**, 074013 (2014), arXiv:1401.5078 [hep-ph] .

[71] A. Bacchetta, F. Delcarro, C. Pisano, and M. Radici, Phys. Lett. B **827**, 136961 (2022), arXiv:2004.14278 [hep-ph] .

[72] M. Bury, A. Prokudin, and A. Vladimirov, Phys. Rev. Lett. **126**, 112002 (2021), arXiv:2012.05135 [hep-ph] .

[73] M. Bury, A. Prokudin, and A. Vladimirov, JHEP **05**, 151 (2021), arXiv:2103.03270 [hep-ph] .

[74] J. Cammarota, L. Gamberg, Z.-B. Kang, J. A. Miller, D. Pitonyak, A. Prokudin, T. C. Rogers, and N. Sato (Jefferson Lab Angular Momentum), Phys. Rev. D **102**, 054002 (2020), arXiv:2002.08384 [hep-ph] .

[75] R. Boussarie et al., (2023), arXiv:2304.03302 [hep-ph] .

[76] J. Dudek et al., Eur. Phys. J. A **48**, 187 (2012), arXiv:1208.1244 [hep-ex] .

[77] K. G. Wilson, Phys. Rev. D **10**, 2445 (1974).

[78] C. Gattringer and C. B. Lang, Quantum chromodynamics on the lattice, Lect. Notes Phys., Vol. 788 (Springer, 2010).

[79] M. Gockeler, R. Horsley, E.-M. Ilgenfritz, H. Perlt, P. E. L. Rakow, G. Schierholz, and A. Schiller, Phys. Rev. D **54**, 5705 (1996), arXiv:hep-lat/9602029 .

[80] H.-W. Lin et al., Prog. Part. Nucl. Phys. **100**, 107 (2018), arXiv:1711.07916 [hep-ph] .

[81] M. Constantinou et al., Prog. Part. Nucl. Phys. **121**, 103908 (2021), arXiv:2006.08636 [hep-ph] .

[82] P. Hagler, B. U. Musch, J. W. Negele, and A. Schafer, EPL **88**, 61001 (2009), arXiv:0908.1283 [hep-lat] .

[83] B. U. Musch, P. Hagler, J. W. Negele, and A. Schafer, Phys. Rev. D **83**, 094507 (2011), arXiv:1011.1213 [hep-lat] .

[84] M. Engelhardt, P. Hägler, B. Musch, J. Negele, and A. Schäfer, Phys. Rev. D **93**, 054501 (2016), arXiv:1506.07826 [hep-lat] .

[85] B. Yoon, T. Bhattacharya, M. Engelhardt, J. Green, R. Gupta, P. Hägler, B. Musch, J. Negele, A. Pochinsky, and S. Syritsyn, in 33rd International Symposium on Lattice Field Theory (SISSA, 2015) arXiv:1601.05717 [hep-lat] .

[86] B. Yoon, M. Engelhardt, R. Gupta, T. Bhattacharya, J. R. Green, B. U. Musch, J. W. Negele, A. V. Pochinsky, A. Schäfer, and S. N. Syritsyn, Phys. Rev. D **96**, 094508 (2017), arXiv:1706.03406 [hep-lat] .

[87] X. Ji, Phys. Rev. Lett. **110**, 262002 (2013), arXiv:1305.1539 [hep-ph] .

[88] X. Ji, P. Sun, X. Xiong, and F. Yuan, Phys. Rev. D **91**, 074009 (2015), arXiv:1405.7640 [hep-ph] .

[89] X. Ji, Y.-S. Liu, Y. Liu, J.-H. Zhang, and Y. Zhao, Rev. Mod. Phys. **93**, 035005 (2021), arXiv:2004.03543 [hep-ph] .

[90] M. A. Ebert, I. W. Stewart, and Y. Zhao, Phys. Rev. D **99**, 034505 (2019), arXiv:1811.00026 [hep-ph] .

[91] P. Shanahan, M. Wagman, and Y. Zhao, Phys. Rev. D **104**, 114502 (2021), arXiv:2107.11930 [hep-lat] .

[92] M. Schlemmer, A. Vladimirov, C. Zimmermann, M. Engelhardt, and A. Schäfer, JHEP **08**, 004 (2021), arXiv:2103.16991 [hep-lat] .

[93] M.-H. Chu et al. (Lattice Parton (LPC)), Phys. Rev. D **106**, 034509 (2022), arXiv:2204.00200 [hep-lat] .

[94] M. Cranmer, arXiv preprint arXiv:2305.01582 (2023), arXiv:2305.01582 [astro-ph.IM] .

[95] M. E. Peskin and D. V. Schroeder, An Introduction to quantum field theory (Addison-Wesley, Reading, USA, 1995).

[96] S. Navas et al. (Particle Data Group), Phys. Rev. D **110**, 030001 (2024).

[97] M. Creutz, Phys. Rev. D **21**, 2308 (1980).

[98] S. J. Brodsky, H.-C. Pauli, and S. S. Pinsky, Phys. Rept. **301**, 299 (1998), arXiv:hep-ph/9705477 .

[99] V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**, 438 (1972).

[100] G. Altarelli and G. Parisi, Nucl. Phys. B **126**, 298 (1977).

[101] Y. L. Dokshitzer, Sov. Phys. JETP **46**, 641 (1977).

[102] K. Goeke, A. Metz, and M. Schlegel, Phys. Lett. B **618**, 90 (2005), arXiv:hep-ph/0504130 .

[103] J. C. Collins, Acta Phys. Polon. B **34**, 3103 (2003), arXiv:hep-ph/0304122 .

[104] A. Bacchetta, M. Boglione, A. Henneman, and P. J. Mulders, Phys. Rev. Lett. **85**, 712 (2000), arXiv:hep-ph/9912490 .

[105] J. C. Collins, T. C. Rogers, and A. M. Stasto, Phys. Rev. D **77**, 085009 (2008), arXiv:0708.2833 [hep-ph] .

[106] B. U. Musch, Transverse momentum distributions inside the nucleon from lattice QCD, Ph.D. thesis, TU Munich (2009), arXiv:0907.2381 [hep-lat] .

[107] S. M. Aybat, J. C. Collins, J.-W. Qiu, and T. C. Rogers, Phys. Rev. D **85**, 034043 (2012), arXiv:1110.6428 [hep-ph] .

[108] A. Bacchetta, D. Boer, M. Diehl, and P. J. Mulders, JHEP **08**, 023 (2008), arXiv:0803.0227 [hep-ph] .

[109] D. Boer, L. Gamberg, B. Musch, and A. Prokudin, JHEP **10**, 021 (2011), arXiv:1107.5294 [hep-ph] .

[110] X. Ji, Y. Liu, and Y.-S. Liu, Nucl. Phys. B **955**, 115054 (2020), arXiv:1910.11415 [hep-ph] .

[111] M. A. Ebert, S. T. Schindler, I. W. Stewart, and Y. Zhao, JHEP **04**, 178 (2022), arXiv:2201.08401 [hep-ph] .

[112] A. V. Radyushkin, Phys. Rev. D **96**, 034025 (2017), arXiv:1705.01488 [hep-ph] .

[113] H. B. Nielsen and M. Ninomiya, Nucl. Phys. B **185**, 20 (1981), [Erratum: Nucl.Phys.B 195, 541 (1982)].

[114] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B **259**, 572 (1985).

[115] S. Duane, A. D. Kennedy, B. J. Pendleton, and D. Roweth, Phys. Lett. B **195**, 216 (1987).

[116] B. U. Musch, P. Hägler, M. Engelhardt, J. W. Negele, and A. Schäfer, Phys. Rev. D **85**, 094510 (2012).

[117] D. Boer, L. Gamberg, B. U. Musch, and A. Prokudin, Journal of High Energy Physics **2011**, 21 (2011).

[118] A. Idilbi, X.-D. Ji, J.-P. Ma, and F. Yuan, Phys. Rev. D **70**, 074021 (2004), arXiv:hep-ph/0406302 .

[119] A. Accardi, A. Bacchetta, W. Melnitchouk, and M. Schlegel, JHEP **11**, 093 (2009), arXiv:0907.2942 [hep-ph] .

[120] J. R. Koza, Genetic Programming: On the Programming of Computers by Means of Natural Selection (MIT Press, Cambridge, MA, USA, 1992).

[121] M. Schmidt and H. Lipson, Science **324**, 81 (2009).

[122] B. Yoon, Y.-C. Jang, C. Jung, and W. Lee, "Covariance fitting of highly correlated data in lattice qcd," (2012), arXiv:1101.2248 [hep-lat] .

[123] G. P. Lepage, B. Clark, C. T. H. Davies, K. Hornbostel, P. B. Mackenzie, C. Morningstar, and H. Trottier, Nucl. Phys. B Proc. Suppl. **106**, 12 (2002), arXiv:hep-lat/0110175 .