

(b_L, b_T, η) Fits for $\text{Re } \tilde{A}_{2B}$ for $P_L=-1$
 with $b \geq 3$ and $6 \geq \eta \geq 10$

Model Expression	Params List	Fitted Values	χ^2/DoF
$a e^{-f \eta} e^{-c b_L P_L } e^{-d \sqrt{b_L^2+b_T^2}}$	{a, c, d, f}	{ $\langle 8.02(56) \rangle_{2902 \text{ J}}$, $\langle -0.0021(76) \rangle_{2902 \text{ J}}$, $\langle 0.579(13) \rangle_{2902 \text{ J}}$, $\langle 0.5348(84) \rangle_{2902 \text{ J}}$ }	2.15678
$a e^{-f \eta} e^{-c b_L P_L } e^{-d (b_L^2+b_T^2)}$	{a, c, d, f}	{ $\langle 2.08(13) \rangle_{2902 \text{ J}}$, $\langle 0.0199(73) \rangle_{2902 \text{ J}}$, $\langle 0.0564(15) \rangle_{2902 \text{ J}}$, $\langle 0.5364(86) \rangle_{2902 \text{ J}}$ }	4.98308
$\frac{a e^{-f \eta} e^{-c b_L P_L }}{(1+d (b_L^2+b_T^2))^{j_2}}$	{a, c, d, f, j2}	{ $\langle 5.28(60) \rangle_{2902 \text{ J}}$, $\langle -0.0089(84) \rangle_{2902 \text{ J}}$, $\langle 0.094(16) \rangle_{2902 \text{ J}}$, $\langle 0.5344(84) \rangle_{2902 \text{ J}}$, $\langle 2.07(13) \rangle_{2902 \text{ J}}$ }	1.90543
$a e^{-f \eta} e^{-c (b_L P_L)^2} e^{-d \sqrt{b_L^2+b_T^2}}$	{a, c, d, f}	{ $\langle 8.18(58) \rangle_{2902 \text{ J}}$, $\langle -0.0018(17) \rangle_{2902 \text{ J}}$, $\langle 0.587(14) \rangle_{2902 \text{ J}}$, $\langle 0.5348(85) \rangle_{2902 \text{ J}}$ }	2.12445
$\frac{a e^{-f \eta} e^{-c (b_L P_L)^2}}{(1+d (b_L^2+b_T^2))^{j_2}}$	{a, c, d, f, j2}	{ $\langle 4.99(50) \rangle_{2902 \text{ J}}$, $\langle -0.0015(16) \rangle_{2902 \text{ J}}$, $\langle 0.085(13) \rangle_{2902 \text{ J}}$, $\langle 0.5343(84) \rangle_{2902 \text{ J}}$, $\langle 2.14(14) \rangle_{2902 \text{ J}}$ }	1.90851