

Gaussian-type fit for $\text{Re } \tilde{A}_{2B}$ for $P_L=-1$

$b \geq 3$ are SELECTED for fit!

$6 \geq \eta \geq 10$ are SELECTED for fit!

Model Expression	Params List	Fitted Values		χ^2/DoF	AIC	FT Plot
$a e^{-f \eta} \left(1 + (c + d) b_L^2 + d b_T^2 \right)^{-j}$	a c d f	$\langle 2.19 (16) \rangle_{2902 \text{ J}}$ $\langle 0.0218 (44) \rangle_{2902 \text{ J}}$ $\langle 2.62 (33) \rangle_{2902 \text{ J}}$	$\langle 0.0218 (44) \rangle_{2902 \text{ J}}$ $\langle 0.5382 (88) \rangle_{2902 \text{ J}}$	24.778	5461.17	
$a e^{-f \eta} \left(1 + \left(d + c \left(1 - \frac{k_1}{\eta} \right) \right) b_L^2 + d b_T^2 \right)^{-j}$	a c d f k1 j	$\langle 2.85 (30) \rangle_{2902 \text{ J}}$ $\langle 0.078 (14) \rangle_{2902 \text{ J}}$ $\langle 6.60 (15) \rangle_{2902 \text{ J}}$	$\langle 0.078 (14) \rangle_{2902 \text{ J}}$ $\langle 0.4625 (89) \rangle_{2902 \text{ J}}$ $\langle 2.16 (15) \rangle_{2902 \text{ J}}$	2.113	474.746	
$a e^{-f \eta} \left(1 + \left(d + c \left(1 - \frac{k_2}{\eta^2} - \frac{k_1}{\eta} \right) \right) b_L^2 + d b_T^2 \right)^{-j}$	a c d f k1 k2	$\langle 4.01 (58) \rangle_{2902 \text{ J}}$ $\langle 0.094 (18) \rangle_{2902 \text{ J}}$ $\langle 12.27 (50) \rangle_{2902 \text{ J}}$ $\langle 2.06 (14) \rangle_{2902 \text{ J}}$	$\langle 0.146 (51) \rangle_{2902 \text{ J}}$ $\langle 0.496 (12) \rangle_{2902 \text{ J}}$ $\langle -36.2 (28) \rangle_{2902 \text{ J}}$	1.56788	355.799	
$a e^{-f \eta} \left(1 + \left(d + c \left(1 - \frac{k_3}{\eta^3} - \frac{k_2}{\eta^2} - \frac{k_1}{\eta} \right) \right) b_L^2 + d b_T^2 \right)^{-j}$	a c d f k1 k2 k3 j	$\langle 4.01 (58) \rangle_{2902 \text{ J}}$ $\langle 0.094 (18) \rangle_{2902 \text{ J}}$ $\langle 6.7 (52) \rangle_{2902 \text{ J}}$ $\langle -220 (200) \rangle_{2902 \text{ J}}$	$\langle 0.093 (18) \rangle_{2902 \text{ J}}$ $\langle 0.496 (12) \rangle_{2902 \text{ J}}$ $\langle 36 (63) \rangle_{2902 \text{ J}}$ $\langle 2.06 (14) \rangle_{2902 \text{ J}}$	1.57467	357.703	
$a e^{-f \eta} \left(1 + \left(d + c \left(1 - \frac{k_1 \text{Log}[\eta]}{\eta} \right) \right) b_L^2 + d b_T^2 \right)^{-j}$	a c d f k1 j	$\langle 3.90 (43) \rangle_{2902 \text{ J}}$ $\langle 0.092 (17) \rangle_{2902 \text{ J}}$ $\langle 3.616 (83) \rangle_{2902 \text{ J}}$	$\langle 0.092 (17) \rangle_{2902 \text{ J}}$ $\langle 0.4929 (86) \rangle_{2902 \text{ J}}$ $\langle 2.07 (14) \rangle_{2902 \text{ J}}$	1.66147	375.863	
$a e^{-f \eta} \left(1 + \left(d + c \left(1 - e^{-n \eta} k_2 - \frac{k_1}{\eta} \right) \right) b_L^2 + d b_T^2 \right)^{-j}$	a c d f k1 k2 n j	$\langle 3.98 (58) \rangle_{2902 \text{ J}}$ $\langle 0.093 (18) \rangle_{2902 \text{ J}}$ $\langle 7.77 (98) \rangle_{2902 \text{ J}}$ $\langle 0.64 (45) \rangle_{2902 \text{ J}}$	$\langle 0.093 (18) \rangle_{2902 \text{ J}}$ $\langle 0.495 (12) \rangle_{2902 \text{ J}}$ $\langle 82 (76) \rangle_{2902 \text{ J}}$ $\langle 2.06 (14) \rangle_{2902 \text{ J}}$	1.57582	357.952	