



$$\tilde{A}_{2B}^{Re}(\eta|v|/a, (P \cdot b) \cdot \hat{L}/(2\pi), b^2/a^2, \hat{\zeta} = 0.090772)$$



- $b^2/a^2 = 0$
- $b^2/a^2 = 1$
- $b^2/a^2 = 2$
- $b^2/a^2 = 4$
- $b^2/a^2 = 5$
- $b^2/a^2 = 8$
- $b^2/a^2 = 9$
- $b^2/a^2 = 16$
- $b^2/a^2 = 17$
- $b^2/a^2 = 18$
- $b^2/a^2 = 20$
- $b^2/a^2 = 25$
- $b^2/a^2 = 32$
- $b^2/a^2 = 36$
- $b^2/a^2 = 45$
- $b^2/a^2 = 49$
- $b^2/a^2 = 50$
- $b^2/a^2 = 68$

$$\tilde{A}_{2B}^{Re}(\eta|v|/a, (P \cdot b) \cdot \hat{L}/(2\pi), b^2/a^2, \hat{\zeta} = 0.090772)$$



- $(P \cdot b) \cdot \hat{L}/(2\pi) = -8$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -6$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -5$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -4$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -3$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -2$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -1$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = -0$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 1$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 2$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 3$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 4$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 5$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 6$
- $(P \cdot b) \cdot \hat{L}/(2\pi) = 8$