

Simultaneous 4D fits for Re $\tilde{A}_{2B}$ for $P_L=-1$ with $b \geq 3$ and $7 \geq \eta \geq 10$				
Model Expression	Params List	Fitted Values	χ^2/DoF	
Invariant: $a \, e^{-f \, \eta} \, e^{-c \, P \cdot b } \, e^{-d \, b^2}$ Cartesian: $a \, e^{-f \, \eta} \, e^{-c \, b_L \, P_L } \, e^{-d \, (b_L^2+b_T^2)}$	{a, c, d, f}	{ $\langle 2.63(26) \rangle_{2902 \, \text{J}}$, $\langle 0.029(11) \rangle_{2902 \, \text{J}}$, $\langle 0.0649(26) \rangle_{2902 \, \text{J}}$, $\langle 0.543(12) \rangle_{2902 \, \text{J}}$ }	3.33953	
Invariant: $\frac{a \, e^{-f \, \eta} \, e^{-c \, P \cdot b }}{(1+d \, b^2)^j}$ Cartesian: $\frac{a \, e^{-f \, \eta} \, e^{-c \, b_L \, P_L }}{(1+d \, (b_L^2+b_T^2))^j}$	{a, c, d, f, j}	{ $\langle 7.2(14) \rangle_{2902 \, \text{J}}$, $\langle -0.007 \pm 0.012 \rangle_{2902 \, \text{J}}$, $\langle 0.102(26) \rangle_{2902 \, \text{J}}$, $\langle 0.541(12) \rangle_{2902 \, \text{J}}$, $\langle 2.19(19) \rangle_{2902 \, \text{J}}$ }	1.27672	
Invariant: $\frac{a \, e^{-f \, \eta} \, e^{-c \, P \cdot b }}{(1+d \, b^2)^2}$ Cartesian: $\frac{a \, e^{-f \, \eta} \, e^{-c \, b_L \, P_L }}{(1+d \, (b_L^2+b_T^2))^2}$	{a, c, d, f}	{ $\langle 8.9(14) \rangle_{2902 \, \text{J}}$, $\langle -0.009 \pm 0.012 \rangle_{2902 \, \text{J}}$, $\langle 0.144(15) \rangle_{2902 \, \text{J}}$, $\langle 0.541(12) \rangle_{2902 \, \text{J}}$ }	1.30998	
Invariant: $\frac{a \, e^{-f \, \eta} \, e^{-c \, P \cdot b } \, e^{-d_1 \, b^2}}{(1+d_2 \, b^2)^j}$ Cartesian: $\frac{a \, e^{-f \, \eta} \, e^{-c \, b_L \, P_L } \, e^{-d_1 \, (b_L^2+b_T^2)}}{(1+d_2 \, (b_L^2+b_T^2))^j}$	{a, c, d1, d2, f, j}	{ $\langle 5.7(29) \rangle_{2902 \, \text{J}}$, $\langle -0.007 \pm 0.012 \rangle_{2902 \, \text{J}}$, $\langle 0.007 \pm 0.019 \rangle_{2902 \, \text{J}}$, $\langle 0.03 \pm 0.12 \rangle_{2902 \, \text{J}}$, $\langle 0.541(12) \rangle_{2902 \, \text{J}}$, $\langle 1.6(13) \rangle_{2902 \, \text{J}}$ }	1.2761	