
Hierarchical Clustering of Alzheimer's Diagnostic Groups
Unsigned mean Fisher-z connectivity strength (FC)

cells are the ROI-level P-score = -log10(FDR-adjusted q) for each diagnostic group vs the pooled rest (BH over 246 x 6 = 1476 tests per biomarker)
groups clustered by correlation distance + average linkage over the 246-D P-SCORE profile
ROI rows ORDERED by euclidean + average on the same profiles (tree not drawn)



columns are ordered by the dendrogram above, NOT by disease stage

clinical progression order: CN → SMC → EMCI → MCI → LMCI → AD

bootstrap stability of the branches (1000 resamples of subjects within groups): {CN,MCI} 72%; {SMC,EMCI} 45%; {LMCI,AD} 38%

group P-score = -log10(q) from the MULTIVARIATE stage-vs-rest permutation test (24-test family); max ROI P = largest ROI-level P-score, DESCRIPTIVE only

d = merge DISTANCE (geometric, not a p-value). *q = permutation q for that cluster contrast, SELECTIVE (clusters came from these data) -- anti-conservative, descriptive only.

COLOUR SHOWS RELATIVE VARIATION WITHIN THIS BIOMARKER ONLY, NOT PROXIMITY TO SIGNIFICANCE. Figures are NOT comparable to one another.

Descriptive only. Branch heights are distances, not p-values. No group reaches q < 0.05 and no ROI x stage test in this project does either.