

Hierarchical Clustering of Alzheimer's Diagnostic Groups
Regional homogeneity (mReHo)

cells are the ROI-level P-score = -log10(FDR-adjusted q) for each diagnostic group vs the pooled rest (BH over 246 x 6 = 1476 tests per biomarker)
groups clustered by correlation distance + average linkage over the 246-D P-SCORE profile
ROI rows ORDERED by euclidean + average on the same profiles (tree not drawn)



	CN	LMCI	EMCI	AD	SMC	MCI
group P-score	**0.339**	**0.119**	**0.856**	**0.738**	**0.119**	**0.559**
max ROI P	0.524	0.353	0.928	0.928	0.524	0.498



clinical progression order: CN → SMC → EMCI → MCI → LMCI → AD

bootstrap stability of the branches (1000 resamples of subjects within groups): {CN,AD} 36%; {SMC,LMCI} 16%; {SMC,EMCI,MCI,LMCI} 12%

group P-score = -log10(q) from the MULTIVARIATE stage-vs-rest permutation test (24-test family); max ROI P = largest ROI-level P-score, DESCRIPTIVE only

d = merge DISTANCE (geometric, not a p-value). *q = permutation q for that cluster contrast, SELECTIVE (clusters came from these data) -- anti-conservative, descriptive only.

COLOUR SHOWS RELATIVE VARIATION WITHIN THIS BIOMARKER ONLY, NOT PROXIMITY TO SIGNIFICANCE. Figures are NOT comparable to one another.

Descriptive only. Branch heights are distances, not p-values. No group reaches q < 0.05 and no ROI x stage test in this project does either.