Class 10 Mathematics: Remedial Master Practice Worksheet

4,000 Targeted Questions Across 4 Core Problem Areas

Designed for Sign Conventions, Equation Formation, Method Selection, and Denominator Rules

---

Contents

PART 1: 4,000 PRACTICE QUESTIONS **2**

MODULE 1: SIGN CONVENTIONS AND TRANSPOSITION RULES (Q1 – Q1000) 3

MODULE 2: WORD PROBLEM TRANSLATION AND EQUATION FORMATION (Q1001 – Q2000) 25

MODULE 3: METHOD SELECTION STRATEGY (WHAT METHOD WHEN AND WHERE) (Q2001 – Q3000) 56

MODULE 4: DENOMINATOR RULES AND FRACTIONAL EQUATIONS (Q3001 – Q4000) 81

PART 2: STEP-BY-STEP ANSWER KEY **108**

PART 1: 4,000 PRACTICE QUESTIONS

MODULE 1: SIGN CONVENTIONS AND TRANSPOSITION RULES (Q1 – Q1000)

Q1. Solve for x: $11x - 13 = 28$. Pay strict attention to the sign change when moving -13 across the equals sign.

Q2. Solve for x: $10x - 3 = -1$. Pay strict attention to the sign change when moving -3 across the equals sign.

Q3. Solve for x: $2x + 17 = 2$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q4. Solve for x: $5x + 13 = -16$. Pay strict attention to the sign change when moving $+13$ across the equals sign.

Q5. Solve for x: $6x + 6 = 14$. Pay strict attention to the sign change when moving $+6$ across the equals sign.

Q6. Solve for x: $14x - 12 = -9$. Pay strict attention to the sign change when moving -12 across the equals sign.

Q7. Solve for x: $9x - 21 = -14$. Pay strict attention to the sign change when moving -21 across the equals sign.

Q8. Solve for x: $5x - 15 = -25$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q9. Solve for x: $9x + 24 = 4$. Pay strict attention to the sign change when moving $+24$ across the equals sign.

Q10. Solve for x: $7x - 13 = 0$. Pay strict attention to the sign change when moving -13 across the equals sign.

Q11. Solve for x: $14x - 5 = -15$. Pay strict attention to the sign change when moving -5 across the equals sign.

Q12. Solve for x: $12x + 25 = -5$. Pay strict attention to the sign change when moving $+25$ across the equals sign.

Q13. Solve for x: $8x - 12 = -26$. Pay strict attention to the sign change when moving -12 across the equals sign.

Q14. Solve for x: $8x + 4 = -11$. Pay strict attention to the sign change when moving $+4$ across the equals sign.

Q15. Solve for x: $5x - 18 = 11$. Pay strict attention to the sign change when moving -18 across the equals sign.

Q16. Solve for x: $15x + 14 = 13$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q17. Solve for x: $7x + 17 = 24$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q18. Solve for x: $6x + 7 = 20$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q19. Solve for x: $8x - 11 = -3$. Pay strict attention to the sign change when moving -11 across the equals sign.

Q20. Solve for x: $3x + 7 = 17$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q21. Solve for x: $12x + 1 = -30$. Pay strict attention to the sign change when moving $+1$ across the equals sign.

Q22. Solve for x: $15x + 4 = -17$. Pay strict attention to the sign change when moving $+4$ across the equals sign.

Q23. Solve for x: $10x - 19 = -8$. Pay strict attention to the sign change when moving -19 across the equals sign.

Q24. Solve for x: $14x - 24 = -7$. Pay strict attention to the sign change when moving -24 across the equals sign.

Q25. Solve for x: $13x - 2 = -30$. Pay strict attention to the sign change when moving -2 across the equals sign.

Q26. Solve for x: $6x + 11 = 22$. Pay strict attention to the sign change when moving $+11$ across the equals sign.

Q27. Solve for x: $4x - 23 = -3$. Pay strict attention to the sign change when moving -23 across the equals sign.

Q28. Solve for x: $4x - 8 = -26$. Pay strict attention to the sign change when moving -8 across the equals sign.

Q29. Solve for x: $14x - 16 = -5$. Pay strict attention to the sign change when moving -16 across the equals sign.

Q30. Solve for x: $12x + 21 = 22$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q31. Solve for x: $7x + 16 = 6$. Pay strict attention to the sign change when moving $+16$ across the equals sign.

Q32. Solve for x: $9x - 15 = 1$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q33. Solve for x: $15x + 7 = -11$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q34. Solve for x: $15x - 14 = 14$. Pay strict attention to the sign change when moving -14 across the equals sign.

Q35. Solve for x: $10x + 18 = 25$. Pay strict attention to the sign change when moving $+18$ across the equals sign.

Q36. Solve for x: $2x - 20 = -26$. Pay strict attention to the sign change when moving -20 across the equals sign.

Q37. Solve for x: $2x + 11 = -8$. Pay strict attention to the sign change when moving $+11$ across the equals sign.

Q38. Solve for x: $10x + 19 = 26$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q39. Solve for x: $9x - 18 = 13$. Pay strict attention to the sign change when moving -18 across the equals sign.

Q40. Solve for x: $13x + 14 = 19$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q41. Solve for x: $11x + 4 = 20$. Pay strict attention to the sign change when moving $+4$ across the equals sign.

Q42. Solve for x: $12x - 17 = 13$. Pay strict attention to the sign change when moving -17 across the equals sign.

Q43. Solve for x: $4x + 25 = -18$. Pay strict attention to the sign change when moving $+25$ across the equals sign.

Q44. Solve for x: $13x + 10 = -2$. Pay strict attention to the sign change when moving $+10$ across the equals sign.

Q45. Solve for x: $4x - 20 = -21$. Pay strict attention to the sign change when moving -20 across the equals sign.

Q46. Solve for x: $6x - 4 = -6$. Pay strict attention to the sign change when moving -4 across the equals sign.

Q47. Solve for x: $3x - 22 = 5$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q48. Solve for x: $10x + 19 = -8$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q49. Solve for x: $9x - 22 = -18$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q50. Solve for x: $6x + 18 = 22$. Pay strict attention to the sign change when moving $+18$ across the equals sign.

Q51. Solve for x: $7x - 20 = -13$. Pay strict attention to the sign change when moving -20 across the equals sign.

Q52. Solve for x: $4x - 9 = -1$. Pay strict attention to the sign change when moving -9 across the equals sign.

Q53. Solve for x: $5x + 2 = -23$. Pay strict attention to the sign change when moving $+2$ across the equals sign.

Q54. Solve for x: $10x - 5 = -15$. Pay strict attention to the sign change when moving -5 across the equals sign.

Q55. Solve for x: $3x - 8 = 11$. Pay strict attention to the sign change when moving -8 across the equals sign.

Q56. Solve for x: $4x - 25 = -30$. Pay strict attention to the sign change when moving -25 across the equals sign.

Q57. Solve for x: $12x + 7 = 24$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q58. Solve for x: $11x + 4 = 11$. Pay strict attention to the sign change when moving $+4$ across the equals sign.

Q59. Solve for x: $11x - 23 = 21$. Pay strict attention to the sign change when moving -23 across the equals sign.

Q60. Solve for x: $7x - 10 = 20$. Pay strict attention to the sign change when moving -10 across the equals sign.

Q61. Solve for x: $6x + 5 = 12$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q62. Solve for x: $5x + 10 = 7$. Pay strict attention to the sign change when moving $+10$ across the equals sign.

Q63. Solve for x: $10x - 5 = -9$. Pay strict attention to the sign change when moving -5 across the equals sign.

Q64. Solve for x: $2x - 17 = -12$. Pay strict attention to the sign change when moving -17 across the equals sign.

Q65. Solve for x: $7x + 18 = -7$. Pay strict attention to the sign change when moving $+18$ across the equals sign.

Q66. Solve for x: $8x + 7 = 10$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q67. Solve for x: $2x + 22 = -16$. Pay strict attention to the sign change when moving $+22$ across the equals sign.

Q68. Solve for x: $10x - 21 = -17$. Pay strict attention to the sign change when moving -21 across the equals sign.

Q69. Solve for x: $4x + 17 = -23$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q70. Solve for x: $4x - 4 = -23$. Pay strict attention to the sign change when moving -4 across the equals sign.

Q71. Solve for x: $7x - 24 = -2$. Pay strict attention to the sign change when moving -24 across the equals sign.

Q72. Solve for x: $15x + 17 = -18$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q73. Solve for x: $12x + 17 = 13$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q74. Solve for x: $15x - 2 = 2$. Pay strict attention to the sign change when moving -2 across the equals sign.

Q75. Solve for x: $12x + 15 = 5$. Pay strict attention to the sign change when moving $+15$ across the equals sign.

Q76. Solve for x: $5x - 17 = -1$. Pay strict attention to the sign change when moving -17 across the equals sign.

Q77. Solve for x: $10x + 10 = -10$. Pay strict attention to the sign change when moving $+10$ across the equals sign.

Q78. Solve for x: $6x + 18 = -11$. Pay strict attention to the sign change when moving $+18$ across the equals sign.

Q79. Solve for x: $13x - 9 = -15$. Pay strict attention to the sign change when moving -9 across the equals sign.

Q80. Solve for x: $8x - 12 = -1$. Pay strict attention to the sign change when moving -12 across the equals sign.

Q81. Solve for x: $6x + 23 = -28$. Pay strict attention to the sign change when moving $+23$ across the equals sign.

Q82. Solve for x: $5x - 22 = 7$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q83. Solve for x: $7x + 22 = 26$. Pay strict attention to the sign change when moving $+22$ across the equals sign.

Q84. Solve for x: $9x + 3 = 12$. Pay strict attention to the sign change when moving $+3$ across the equals sign.

Q85. Solve for x: $2x - 8 = 5$. Pay strict attention to the sign change when moving -8 across the equals sign.

Q86. Solve for x: $13x - 12 = 30$. Pay strict attention to the sign change when moving -12 across the equals sign.

Q87. Solve for x: $2x + 24 = 27$. Pay strict attention to the sign change when moving $+24$ across the equals sign.

Q88. Solve for x: $9x + 14 = 29$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q89. Solve for x: $15x - 22 = -30$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q90. Solve for x: $8x + 3 = 12$. Pay strict attention to the sign change when moving $+3$ across the equals sign.

Q91. Solve for x: $6x + 24 = 9$. Pay strict attention to the sign change when moving $+24$ across the equals sign.

Q92. Solve for x: $4x + 6 = 17$. Pay strict attention to the sign change when moving $+6$ across the equals sign.

Q93. Solve for x: $14x + 15 = -27$. Pay strict attention to the sign change when moving $+15$ across the equals sign.

Q94. Solve for x: $2x + 19 = 25$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q95. Solve for x: $6x + 13 = 10$. Pay strict attention to the sign change when moving $+13$ across the equals sign.

Q96. Solve for x: $3x + 19 = -25$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q97. Solve for x: $11x + 3 = -9$. Pay strict attention to the sign change when moving $+3$ across the equals sign.

Q98. Solve for x: $10x + 21 = -26$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q99. Solve for x: $4x + 5 = 15$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q100. Solve for x: $8x - 19 = -14$. Pay strict attention to the sign change when moving -19 across the equals sign.

Q101. Solve for x: $2x - 10 = -8$. Pay strict attention to the sign change when moving -10 across the equals sign.

Q102. Solve for x: $5x + 16 = 20$. Pay strict attention to the sign change when moving $+16$ across the equals sign.

Q103. Solve for x: $10x - 19 = -16$. Pay strict attention to the sign change when moving -19 across the equals sign.

Q104. Solve for x: $13x - 13 = 11$. Pay strict attention to the sign change when moving -13 across the equals sign.

Q105. Solve for x: $4x - 14 = 16$. Pay strict attention to the sign change when moving -14 across the equals sign.

Q106. Solve for x: $11x + 25 = 12$. Pay strict attention to the sign change when moving $+25$ across the equals sign.

Q107. Solve for x: $13x - 13 = -2$. Pay strict attention to the sign change when moving -13 across the equals sign.

Q108. Solve for x: $4x - 23 = -18$. Pay strict attention to the sign change when moving -23 across the equals sign.

Q109. Solve for x: $10x + 20 = 20$. Pay strict attention to the sign change when moving $+20$ across the equals sign.

Q110. Solve for x: $12x + 7 = 26$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q111. Solve for x: $5x - 19 = 7$. Pay strict attention to the sign change when moving -19 across the equals sign.

Q112. Solve for x: $10x - 25 = -9$. Pay strict attention to the sign change when moving -25 across the equals sign.

Q113. Solve for x: $10x - 17 = 6$. Pay strict attention to the sign change when moving -17 across the equals sign.

Q114. Solve for x: $12x + 5 = 28$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q115. Solve for x: $4x - 18 = -23$. Pay strict attention to the sign change when moving -18 across the equals sign.

Q116. Solve for x: $2x + 3 = 9$. Pay strict attention to the sign change when moving $+3$ across the equals sign.

Q117. Solve for x: $11x + 20 = 25$. Pay strict attention to the sign change when moving $+20$ across the equals sign.

Q118. Solve for x: $11x + 13 = -25$. Pay strict attention to the sign change when moving $+13$ across the equals sign.

Q119. Solve for x: $6x + 7 = 18$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q120. Solve for x: $4x + 19 = -27$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q121. Solve for x: $3x - 7 = 13$. Pay strict attention to the sign change when moving -7 across the equals sign.

Q122. Solve for x: $10x - 22 = -24$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q123. Solve for x: $6x - 7 = -15$. Pay strict attention to the sign change when moving -7 across the equals sign.

Q124. Solve for x: $4x + 21 = 22$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q125. Solve for x: $8x + 16 = 5$. Pay strict attention to the sign change when moving $+16$ across the equals sign.

Q126. Solve for x: $8x - 16 = 20$. Pay strict attention to the sign change when moving -16 across the equals sign.

Q127. Solve for x: $15x + 20 = 18$. Pay strict attention to the sign change when moving $+20$ across the equals sign.

Q128. Solve for x: $13x + 15 = -9$. Pay strict attention to the sign change when moving $+15$ across the equals sign.

Q129. Solve for x: $13x - 23 = -15$. Pay strict attention to the sign change when moving -23 across the equals sign.

Q130. Solve for x: $6x - 1 = 27$. Pay strict attention to the sign change when moving -1 across the equals sign.

Q131. Solve for x: $5x + 5 = -15$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q132. Solve for x: $5x + 1 = -21$. Pay strict attention to the sign change when moving $+1$ across the equals sign.

Q133. Solve for x: $11x - 15 = 1$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q134. Solve for x: $9x + 6 = 29$. Pay strict attention to the sign change when moving $+6$ across the equals sign.

Q135. Solve for x: $5x - 8 = 18$. Pay strict attention to the sign change when moving -8 across the equals sign.

Q136. Solve for x: $2x + 23 = 28$. Pay strict attention to the sign change when moving $+23$ across the equals sign.

Q137. Solve for x: $5x - 4 = 19$. Pay strict attention to the sign change when moving -4 across the equals sign.

Q138. Solve for x: $2x - 11 = -11$. Pay strict attention to the sign change when moving -11 across the equals sign.

Q139. Solve for x: $10x - 9 = -28$. Pay strict attention to the sign change when moving -9 across the equals sign.

Q140. Solve for x: $14x - 1 = -22$. Pay strict attention to the sign change when moving -1 across the equals sign.

Q141. Solve for x: $6x + 12 = 22$. Pay strict attention to the sign change when moving $+12$ across the equals sign.

Q142. Solve for x: $12x + 21 = 11$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q143. Solve for x: $12x + 14 = -7$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q144. Solve for x: $14x + 25 = 27$. Pay strict attention to the sign change when moving $+25$ across the equals sign.

Q145. Solve for x: $9x + 14 = 29$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q146. Solve for x: $6x - 22 = -1$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q147. Solve for x: $12x - 6 = -25$. Pay strict attention to the sign change when moving -6 across the equals sign.

Q148. Solve for x: $4x + 17 = -29$. Pay strict attention to the sign change when moving $+17$ across the equals sign.

Q149. Solve for x: $7x + 5 = 24$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q150. Solve for x: $2x - 1 = -28$. Pay strict attention to the sign change when moving -1 across the equals sign.

Q151. Solve for x: $7x - 22 = -16$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q152. Solve for x: $2x - 1 = -22$. Pay strict attention to the sign change when moving -1 across the equals sign.

Q153. Solve for x: $9x + 7 = 23$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q154. Solve for x: $3x + 21 = -19$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q155. Solve for x: $8x + 19 = 12$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q156. Solve for x: $14x + 22 = 16$. Pay strict attention to the sign change when moving $+22$ across the equals sign.

Q157. Solve for x: $9x - 8 = -7$. Pay strict attention to the sign change when moving -8 across the equals sign.

Q158. Solve for x: $15x + 21 = -17$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q159. Solve for x: $12x + 13 = -12$. Pay strict attention to the sign change when moving $+13$ across the equals sign.

Q160. Solve for x: $15x + 7 = -22$. Pay strict attention to the sign change when moving $+7$ across the equals sign.

Q161. Solve for x: $14x - 15 = 3$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q162. Solve for x: $12x + 6 = -11$. Pay strict attention to the sign change when moving $+6$ across the equals sign.

Q163. Solve for x: $5x + 1 = 26$. Pay strict attention to the sign change when moving $+1$ across the equals sign.

Q164. Solve for x: $15x + 8 = -28$. Pay strict attention to the sign change when moving $+8$ across the equals sign.

Q165. Solve for x: $2x + 8 = -24$. Pay strict attention to the sign change when moving $+8$ across the equals sign.

Q166. Solve for x: $13x + 20 = -30$. Pay strict attention to the sign change when moving $+20$ across the equals sign.

Q167. Solve for x: $14x + 19 = 1$. Pay strict attention to the sign change when moving $+19$ across the equals sign.

Q168. Solve for x: $6x + 4 = 10$. Pay strict attention to the sign change when moving $+4$ across the equals sign.

Q169. Solve for x: $7x + 14 = -12$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q170. Solve for x: $4x - 2 = 30$. Pay strict attention to the sign change when moving -2 across the equals sign.

Q171. Solve for x: $9x - 10 = 19$. Pay strict attention to the sign change when moving -10 across the equals sign.

Q172. Solve for x: $4x + 24 = -2$. Pay strict attention to the sign change when moving $+24$ across the equals sign.

Q173. Solve for x: $13x - 21 = -6$. Pay strict attention to the sign change when moving -21 across the equals sign.

Q174. Solve for x: $3x + 14 = 18$. Pay strict attention to the sign change when moving $+14$ across the equals sign.

Q175. Solve for x: $9x - 19 = 4$. Pay strict attention to the sign change when moving -19 across the equals sign.

Q176. Solve for x: $2x + 18 = -1$. Pay strict attention to the sign change when moving $+18$ across the equals sign.

Q177. Solve for x: $6x - 3 = 23$. Pay strict attention to the sign change when moving -3 across the equals sign.

Q178. Solve for x: $8x - 15 = 18$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q179. Solve for x: $3x + 10 = 20$. Pay strict attention to the sign change when moving $+10$ across the equals sign.

Q180. Solve for x: $5x + 23 = 24$. Pay strict attention to the sign change when moving $+23$ across the equals sign.

Q181. Solve for x: $13x + 9 = 2$. Pay strict attention to the sign change when moving $+9$ across the equals sign.

Q182. Solve for x: $14x + 1 = 17$. Pay strict attention to the sign change when moving $+1$ across the equals sign.

Q183. Solve for x: $8x + 23 = -19$. Pay strict attention to the sign change when moving $+23$ across the equals sign.

Q184. Solve for x: $13x - 22 = -7$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q185. Solve for x: $13x + 12 = -24$. Pay strict attention to the sign change when moving $+12$ across the equals sign.

Q186. Solve for x: $2x - 15 = -14$. Pay strict attention to the sign change when moving -15 across the equals sign.

Q187. Solve for x: $15x + 20 = -20$. Pay strict attention to the sign change when moving $+20$ across the equals sign.

Q188. Solve for x: $10x - 17 = 5$. Pay strict attention to the sign change when moving -17 across the equals sign.

Q189. Solve for x: $13x - 7 = 3$. Pay strict attention to the sign change when moving -7 across the equals sign.

Q190. Solve for x: $11x + 12 = 26$. Pay strict attention to the sign change when moving $+12$ across the equals sign.

Q191. Solve for x: $12x - 10 = 5$. Pay strict attention to the sign change when moving -10 across the equals sign.

Q192. Solve for x: $11x + 21 = 14$. Pay strict attention to the sign change when moving $+21$ across the equals sign.

Q193. Solve for x: $3x - 13 = -22$. Pay strict attention to the sign change when moving -13 across the equals sign.

Q194. Solve for x: $2x + 10 = -18$. Pay strict attention to the sign change when moving $+10$ across the equals sign.

Q195. Solve for x: $15x + 5 = -23$. Pay strict attention to the sign change when moving $+5$ across the equals sign.

Q196. Solve for x: $11x - 16 = 0$. Pay strict attention to the sign change when moving -16 across the equals sign.

Q197. Solve for x: $9x - 6 = -19$. Pay strict attention to the sign change when moving -6 across the equals sign.

Q198. Solve for x: $4x - 14 = -20$. Pay strict attention to the sign change when moving -14 across the equals sign.

Q199. Solve for x: $9x + 2 = 0$. Pay strict attention to the sign change when moving $+2$ across the equals sign.

Q200. Solve for x: $14x - 22 = -20$. Pay strict attention to the sign change when moving -22 across the equals sign.

Q201. Expand and solve for x: $21 - 7(3x - 10) = 31$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q202. Expand and solve for x: $5 - 4(12x - 1) = 15$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q203. Expand and solve for x: $22 - 7(2x - 3) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q204. Expand and solve for x: $31 - 5(12x - 3) = 41$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q205. Expand and solve for x: $17 - 5(12x - 14) = 27$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q206. Expand and solve for x: $26 - 7(4x - 13) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q207. Expand and solve for x: $19 - 7(8x - 6) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q208. Expand and solve for x: $38 - 6(6x - 5) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q209. Expand and solve for x: $14 - 8(4x - 11) = 24$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q210. Expand and solve for x: $38 - 2(8x - 2) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q211. Expand and solve for x: $18 - 8(4x - 5) = 28$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q212. Expand and solve for x: $18 - 9(9x - 15) = 28$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q213. Expand and solve for x: $31 - 4(3x - 2) = 41$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q214. Expand and solve for x: $16 - 8(2x - 5) = 26$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q215. Expand and solve for x: $30 - 4(10x - 10) = 40$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q216. Expand and solve for x: $10 - 2(11x - 6) = 20$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q217. Expand and solve for x: $40 - 4(5x - 13) = 50$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q218. Expand and solve for x: $8 - 6(2x - 3) = 18$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q219. Expand and solve for x: $37 - 8(7x - 12) = 47$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q220. Expand and solve for x: $38 - 5(12x - 14) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q221. Expand and solve for x: $28 - 9(4x - 4) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q222. Expand and solve for x: $34 - 4(2x - 8) = 44$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q223. Expand and solve for x: $33 - 5(6x - 15) = 43$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q224. Expand and solve for x: $33 - 7(3x - 7) = 43$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q225. Expand and solve for x: $27 - 7(2x - 12) = 37$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q226. Expand and solve for x: $9 - 9(2x - 13) = 19$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q227. Expand and solve for x: $25 - 7(7x - 10) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q228. Expand and solve for x: $34 - 3(2x - 2) = 44$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q229. Expand and solve for x: $22 - 8(6x - 4) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q230. Expand and solve for x: $39 - 5(4x - 14) = 49$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q231. Expand and solve for x: $37 - 5(4x - 9) = 47$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q232. Expand and solve for x: $11 - 6(6x - 13) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q233. Expand and solve for x: $6 - 4(7x - 12) = 16$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q234. Expand and solve for x: $25 - 4(11x - 7) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q235. Expand and solve for x: $26 - 6(7x - 10) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q236. Expand and solve for x: $37 - 4(8x - 11) = 47$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q237. Expand and solve for x: $6 - 5(8x - 1) = 16$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q238. Expand and solve for x: $17 - 6(7x - 14) = 27$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q239. Expand and solve for x: $15 - 5(7x - 13) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q240. Expand and solve for x: $24 - 3(12x - 15) = 34$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q241. Expand and solve for x: $22 - 3(6x - 9) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q242. Expand and solve for x: $19 - 4(7x - 1) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q243. Expand and solve for x: $5 - 5(2x - 9) = 15$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q244. Expand and solve for x: $30 - 6(12x - 10) = 40$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q245. Expand and solve for x: $11 - 3(9x - 6) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q246. Expand and solve for x: $26 - 8(8x - 7) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q247. Expand and solve for x: $11 - 8(3x - 10) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q248. Expand and solve for x: $35 - 6(6x - 9) = 45$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q249. Expand and solve for x: $14 - 3(8x - 4) = 24$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q250. Expand and solve for x: $15 - 2(5x - 8) = 25$. Watch out for distributing the negative sign across both

terms inside the parenthesis!

Q251. Expand and solve for x: $15 - 3(3x - 9) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q252. Expand and solve for x: $29 - 6(5x - 12) = 39$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q253. Expand and solve for x: $15 - 9(3x - 15) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q254. Expand and solve for x: $15 - 6(7x - 14) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q255. Expand and solve for x: $22 - 3(5x - 11) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q256. Expand and solve for x: $8 - 2(4x - 6) = 18$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q257. Expand and solve for x: $15 - 8(2x - 11) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q258. Expand and solve for x: $7 - 8(12x - 9) = 17$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q259. Expand and solve for x: $35 - 5(4x - 4) = 45$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q260. Expand and solve for x: $32 - 3(8x - 3) = 42$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q261. Expand and solve for x: $38 - 4(8x - 14) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q262. Expand and solve for x: $10 - 7(10x - 14) = 20$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q263. Expand and solve for x: $31 - 3(10x - 1) = 41$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q264. Expand and solve for x: $34 - 4(10x - 11) = 44$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q265. Expand and solve for x: $21 - 4(12x - 12) = 31$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q266. Expand and solve for x: $11 - 9(3x - 3) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q267. Expand and solve for x: $19 - 9(4x - 4) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q268. Expand and solve for x: $5 - 2(10x - 15) = 15$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q269. Expand and solve for x: $26 - 4(6x - 4) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q270. Expand and solve for x: $12 - 5(9x - 2) = 22$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q271. Expand and solve for x: $20 - 2(12x - 4) = 30$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q272. Expand and solve for x: $12 - 9(10x - 8) = 22$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q273. Expand and solve for x: $25 - 7(3x - 4) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q274. Expand and solve for x: $33 - 6(4x - 1) = 43$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q275. Expand and solve for x: $9 - 4(6x - 10) = 19$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q276. Expand and solve for x: $9 - 2(2x - 2) = 19$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q277. Expand and solve for x: $30 - 3(4x - 11) = 40$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q278. Expand and solve for x: $8 - 6(12x - 1) = 18$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q279. Expand and solve for x: $12 - 4(12x - 9) = 22$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q280. Expand and solve for x: $22 - 9(8x - 11) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q281. Expand and solve for x: $14 - 3(9x - 6) = 24$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q282. Expand and solve for x: $20 - 7(9x - 10) = 30$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q283. Expand and solve for x: $16 - 7(11x - 3) = 26$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q284. Expand and solve for x: $21 - 9(8x - 12) = 31$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q285. Expand and solve for x: $19 - 2(10x - 11) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q286. Expand and solve for x: $33 - 5(2x - 3) = 43$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q287. Expand and solve for x: $39 - 9(7x - 9) = 49$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q288. Expand and solve for x: $34 - 6(3x - 13) = 44$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q289. Expand and solve for x: $21 - 6(7x - 5) = 31$.

Watch out for distributing the negative sign across both terms inside the parenthesis!

Q290. Expand and solve for x: $26 - 8(5x - 7) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q291. Expand and solve for x: $26 - 5(11x - 11) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q292. Expand and solve for x: $25 - 2(10x - 3) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q293. Expand and solve for x: $38 - 5(3x - 7) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q294. Expand and solve for x: $26 - 9(4x - 11) = 36$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q295. Expand and solve for x: $38 - 6(5x - 13) = 48$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q296. Expand and solve for x: $28 - 2(2x - 14) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q297. Expand and solve for x: $11 - 5(8x - 9) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q298. Expand and solve for x: $5 - 6(4x - 8) = 15$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q299. Expand and solve for x: $39 - 6(4x - 11) = 49$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q300. Expand and solve for x: $23 - 8(11x - 11) = 33$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q301. Expand and solve for x: $23 - 3(10x - 2) = 33$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q302. Expand and solve for x: $5 - 4(11x - 9) = 15$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q303. Expand and solve for x: $20 - 4(9x - 2) = 30$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q304. Expand and solve for x: $21 - 2(7x - 5) = 31$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q305. Expand and solve for x: $36 - 9(3x - 4) = 46$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q306. Expand and solve for x: $14 - 9(12x - 7) = 24$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q307. Expand and solve for x: $31 - 5(6x - 9) = 41$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q308. Expand and solve for x: $36 - 7(4x - 7) = 46$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q309. Expand and solve for x: $28 - 3(5x - 15) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q310. Expand and solve for x: $11 - 8(10x - 8) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q311. Expand and solve for x: $35 - 2(6x - 7) = 45$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q312. Expand and solve for x: $36 - 7(7x - 12) = 46$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q313. Expand and solve for x: $12 - 4(7x - 3) = 22$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q314. Expand and solve for x: $40 - 6(4x - 10) = 50$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q315. Expand and solve for x: $24 - 8(9x - 5) = 34$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q316. Expand and solve for x: $35 - 3(10x - 1) = 45$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q317. Expand and solve for x: $25 - 2(5x - 6) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q318. Expand and solve for x: $22 - 5(9x - 5) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q319. Expand and solve for x: $39 - 7(9x - 6) = 49$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q320. Expand and solve for x: $28 - 9(2x - 9) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q321. Expand and solve for x: $32 - 8(12x - 3) = 42$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q322. Expand and solve for x: $37 - 6(7x - 3) = 47$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q323. Expand and solve for x: $35 - 8(9x - 6) = 45$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q324. Expand and solve for x: $12 - 8(3x - 15) = 22$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q325. Expand and solve for x: $11 - 2(3x - 15) = 21$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q326. Expand and solve for x: $25 - 5(9x - 4) = 35$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q327. Expand and solve for x: $19 - 5(2x - 1) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q328. Expand and solve for x: $22 - 2(5x - 11) = 32$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q329. Expand and solve for x: $21 - 9(6x - 15) = 31$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q330. Expand and solve for x: $28 - 9(3x - 15) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q331. Expand and solve for x: $39 - 2(4x - 10) = 49$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q332. Expand and solve for x: $9 - 2(9x - 3) = 19$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q333. Expand and solve for x: $13 - 8(9x - 7) = 23$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q334. Expand and solve for x: $15 - 2(5x - 10) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q335. Expand and solve for x: $24 - 3(4x - 9) = 34$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q336. Expand and solve for x: $19 - 5(6x - 7) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q337. Expand and solve for x: $19 - 4(3x - 7) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q338. Expand and solve for x: $14 - 9(11x - 11) = 24$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q339. Expand and solve for x: $28 - 9(9x - 15) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q340. Expand and solve for x: $15 - 7(6x - 9) = 25$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q341. Expand and solve for x: $36 - 6(7x - 5) = 46$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q342. Expand and solve for x: $17 - 6(10x - 10) = 27$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q343. Expand and solve for x: $8 - 9(5x - 7) = 18$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q344. Expand and solve for x: $33 - 8(12x - 2) = 43$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q345. Expand and solve for x: $28 - 3(2x - 15) = 38$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q346. Expand and solve for x: $19 - 9(7x - 7) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q347. Expand and solve for x: $19 - 6(2x - 5) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q348. Expand and solve for x: $17 - 2(11x - 6) = 27$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q349. Expand and solve for x: $19 - 5(9x - 10) = 29$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q350. Expand and solve for x: $27 - 3(6x - 9) = 37$. Watch out for distributing the negative sign across both terms inside the parenthesis!

Q351. Solve for x: $10x - 8 = 4x - 9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q352. Solve for x: $5x - 17 = 7x - 18$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q353. Solve for x: $11x + 2 = 7x + 3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q354. Solve for x: $5x - 14 = 4x + 3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q355. Solve for x: $12x - 12 = 5x + 12$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q356. Solve for x: $5x + 6 = 8x - 20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q357. Solve for x: $4x + 10 = 2x - 2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q358. Solve for x: $8x + 4 = 10x + 13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q359. Solve for x: $12x + 19 = 8x + 8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q360. Solve for x: $6x + 9 = 9x - 14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q361. Solve for x: $5x + 16 = 4x + 8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q362. Solve for x: $8x - 18 = 4x + 19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q363. Solve for x: $8x - 14 = 9x + 4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q364. Solve for x: $7x + 18 = 3x - 18$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q365. Solve for x: $3x - 15 = 4x + 6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q366. Solve for x: $9x - 16 = 5x - 15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q367. Solve for x: $4x + 2 = 5x + 6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q368. Solve for x: $12x - 14 = 3x + 1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q369. Solve for x: $11x + 1 = 8x - 9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q370. Solve for x: $4x - 5 = 2x + 12$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q371. Solve for x: $3x + 12 = 4x + 7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q372. Solve for x: $3x + 14 = 5x - 2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q373. Solve for x: $7x+2=3x+19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q374. Solve for x: $6x-14=8x+4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q375. Solve for x: $3x+2=6x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q376. Solve for x: $5x-8=3x+15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q377. Solve for x: $11x-20=10x+4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q378. Solve for x: $11x-13=6x-6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q379. Solve for x: $10x-15=9x+1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q380. Solve for x: $8x+8=6x+18$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q381. Solve for x: $9x+12=6x+2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q382. Solve for x: $11x+10=2x+8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q383. Solve for x: $5x+7=4x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q384. Solve for x: $3x+0=6x-4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q385. Solve for x: $11x+18=5x+2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q386. Solve for x: $6x+4=10x-13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q387. Solve for x: $8x+0=10x+17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q388. Solve for x: $3x+2=10x+7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q389. Solve for x: $6x+0=7x-19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q390. Solve for x: $7x-1=9x+20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q391. Solve for x: $5x-13=7x+5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q392. Solve for x: $11x+5=6x+15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q393. Solve for x: $12x-1=4x+9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q394. Solve for x: $4x+10=3x-15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q395. Solve for x: $11x-16=2x-17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q396. Solve for x: $8x+2=5x+3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q397. Solve for x: $3x-9=4x+13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q398. Solve for x: $4x+5=5x+8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q399. Solve for x: $6x-14=2x-4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q400. Solve for x: $8x-5=3x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q401. Solve for x: $7x+13=6x+19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q402. Solve for x: $7x+6=3x-17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q403. Solve for x: $5x-2=10x+1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q404. Solve for x: $10x-17=9x-10$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q405. Solve for x: $9x+1=10x+14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q406. Solve for x: $7x+12=9x+12$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q407. Solve for x: $11x-13=7x-11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q408. Solve for x: $3x-14=5x+8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q409. Solve for x: $7x+11=10x+7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q410. Solve for x: $11x+2=7x-3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q411. Solve for x: $10x-5=7x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q412. Solve for x: $9x+10=2x+11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q413. Solve for x: $6x-9=7x+13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q414. Solve for x: $9x-13=7x-16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q415. Solve for x: $5x+7=4x-19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q416. Solve for x: $11x-4=8x-17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q417. Solve for x: $5x-3=4x-13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q418. Solve for x: $6x+10=5x-16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q419. Solve for x: $6x+20=9x+15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q420. Solve for x: $4x+1=2x+3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q421. Solve for x: $11x+12=10x+11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q422. Solve for x: $9x-15=8x+14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q423. Solve for x: $4x+11=6x+2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q424. Solve for x: $5x-9=3x+5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q425. Solve for x: $3x-17=9x-9$. Group all x-terms on

the left and constants on the right, flipping signs correctly.

Q426. Solve for x: $6x+15=2x-7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q427. Solve for x: $10x+17=9x-3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q428. Solve for x: $3x-5=10x+6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q429. Solve for x: $8x-20=9x+12$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q430. Solve for x: $4x+2=3x+6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q431. Solve for x: $8x+0=10x-11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q432. Solve for x: $5x+12=10x+5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q433. Solve for x: $11x+18=6x+1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q434. Solve for x: $11x+2=10x-20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q435. Solve for x: $4x+2=2x+7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q436. Solve for x: $12x-1=4x-6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q437. Solve for x: $7x+10=3x+1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q438. Solve for x: $3x+9=8x+10$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q439. Solve for x: $5x-2=3x-15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q440. Solve for x: $9x-17=7x-14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q441. Solve for x: $3x+15=4x+20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q442. Solve for x: $12x+15=8x-14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q443. Solve for x: $9x-19=10x-3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q444. Solve for x: $10x-12=5x-3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q445. Solve for x: $10x-6=9x-3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q446. Solve for x: $12x-18=6x-8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q447. Solve for x: $6x+6=3x+15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q448. Solve for x: $11x-18=10x+1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q449. Solve for x: $6x-3=3x+15$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q450. Solve for x: $7x+3=6x+19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q451. Solve for x: $11x-17=4x-19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q452. Solve for x: $10x+19=9x+4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q453. Solve for x: $7x-19=8x-8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q454. Solve for x: $7x-14=6x+9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q455. Solve for x: $8x-6=6x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q456. Solve for x: $9x+8=8x+10$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q457. Solve for x: $9x+2=2x+11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q458. Solve for x: $11x-9=8x-13$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q459. Solve for x: $10x+1=3x+9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q460. Solve for x: $12x+0=5x-4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q461. Solve for x: $9x-3=5x-10$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q462. Solve for x: $4x-20=8x+0$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q463. Solve for x: $9x-13=8x-2$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q464. Solve for x: $9x-5=3x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q465. Solve for x: $8x-11=3x+11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q466. Solve for x: $7x+0=4x+6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q467. Solve for x: $11x+0=6x-6$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q468. Solve for x: $7x-19=6x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q469. Solve for x: $11x+3=3x+20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q470. Solve for x: $11x-9=7x+16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q471. Solve for x: $8x-4=7x-20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q472. Solve for x: $9x-14=3x+20$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q473. Solve for x: $6x+19=4x+19$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q474. Solve for x: $3x+11=10x+16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q475. Solve for x: $12x+15=6x-5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q476. Solve for x: $4x-15=8x-9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q477. Solve for x: $9x-13=5x+16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q478. Solve for x: $9x - 8 = 2x - 1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q479. Solve for x: $8x - 13 = 10x + 18$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q480. Solve for x: $10x - 17 = 7x + 17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q481. Solve for x: $11x + 19 = 8x - 17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q482. Solve for x: $10x - 7 = 9x + 10$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q483. Solve for x: $6x + 12 = 7x + 7$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q484. Solve for x: $8x - 2 = 3x + 4$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q485. Solve for x: $9x - 9 = 8x + 11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q486. Solve for x: $5x + 16 = 3x + 8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q487. Solve for x: $7x - 4 = 6x + 17$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q488. Solve for x: $4x + 0 = 2x + 5$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q489. Solve for x: $12x + 12 = 5x - 14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q490. Solve for x: $6x + 17 = 5x - 1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q491. Solve for x: $9x + 20 = 2x - 11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q492. Solve for x: $9x - 7 = 8x + 11$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q493. Solve for x: $6x + 7 = 4x - 3$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q494. Solve for x: $10x + 8 = 6x - 14$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q495. Solve for x: $10x + 6 = 9x + 9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q496. Solve for x: $3x - 4 = 8x + 8$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q497. Solve for x: $11x - 7 = 4x - 9$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q498. Solve for x: $12x - 9 = 3x - 16$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q499. Solve for x: $12x - 14 = 3x - 1$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q500. Solve for x: $9x + 15 = 8x - 12$. Group all x-terms on the left and constants on the right, flipping signs correctly.

Q501. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 6y = 43$, (2) $7x - 7y = -4$. Show all sign changes.

Q502. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 33$, (2) $3x - 2y = 17$. Show all sign changes.

Q503. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 10$, (2) $3x - 4y = 1$. Show all sign changes.

Q504. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 28$, (2) $3x - 6y = 9$. Show all sign changes.

Q505. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 35$, (2) $3x - 5y = -16$. Show all sign changes.

Q506. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 43$, (2) $2x - 1y = 11$. Show all sign changes.

Q507. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 6y = 25$, (2) $7x - 2y = 1$. Show all sign changes.

Q508. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 2y = 46$, (2) $6x - 5y = -11$. Show all sign changes.

Q509. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 28$, (2) $7x - 7y = -5$. Show all sign changes.

Q510. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 4y = 15$, (2) $6x - 4y = -19$. Show all sign changes.

Q511. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 12$, (2) $3x - 3y = 3$. Show all sign changes.

Q512. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 5y = 22$, (2) $4x - 5y = 7$. Show all sign changes.

Q513. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 7y = 23$, (2) $4x - 1y = -6$. Show all sign changes.

Q514. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 3y = 38$, (2) $5x - 7y = -5$. Show all sign changes.

Q515. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 16$, (2) $7x - 5y = 7$. Show all sign changes.

Q516. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 38$, (2) $3x - 6y = -11$. Show all sign changes.

Q517. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 2y = 35$, (2) $5x - 1y = -4$. Show all sign changes.

Q518. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 3y = 27$, (2) $3x - 6y = -18$. Show all sign changes.

Q519. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 2y = 35$, (2) $3x - 1y = -12$. Show all sign changes.

Q520. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 46$, (2) $3x - 1y = 9$. Show all sign changes.

Q521. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 2y = 37$, (2) $4x - 7y = 5$. Show all sign changes.

Q522. Subtract Equation (2) from Equation (1) to elimi-

nate x: (1) $4x + 2y = 43$, (2) $4x - 4y = 10$. Show all sign changes.

Q523. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 34$, (2) $7x - 6y = 7$. Show all sign changes.

Q524. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 6y = 19$, (2) $7x - 1y = -16$. Show all sign changes.

Q525. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 2y = 11$, (2) $3x - 1y = -10$. Show all sign changes.

Q526. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 2y = 40$, (2) $6x - 4y = -13$. Show all sign changes.

Q527. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 4y = 39$, (2) $3x - 6y = -14$. Show all sign changes.

Q528. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 24$, (2) $3x - 4y = 5$. Show all sign changes.

Q529. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 4y = 19$, (2) $5x - 2y = 6$. Show all sign changes.

Q530. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 22$, (2) $4x - 3y = 6$. Show all sign changes.

Q531. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 34$, (2) $6x - 5y = 7$. Show all sign changes.

Q532. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 2y = 10$, (2) $7x - 6y = -9$. Show all sign changes.

Q533. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 3y = 42$, (2) $7x - 4y = -11$. Show all sign changes.

Q534. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 4y = 15$, (2) $5x - 1y = -13$. Show all sign changes.

Q535. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 2y = 38$, (2) $2x - 3y = 11$. Show all sign changes.

Q536. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 19$, (2) $4x - 1y = -13$. Show all sign changes.

Q537. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 2y = 37$, (2) $3x - 2y = 15$. Show all sign changes.

Q538. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 7y = 50$, (2) $5x - 4y = -20$. Show all sign changes.

Q539. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 26$, (2) $6x - 7y = -12$. Show all sign changes.

Q540. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 27$, (2) $3x - 3y = 6$. Show all sign changes.

Q541. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 31$, (2) $7x - 5y = 4$. Show all sign changes.

Q542. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 38$, (2) $2x - 1y = 7$. Show all sign changes.

Q543. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 4y = 50$, (2) $5x - 3y = 11$. Show all sign changes.

Q544. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 7y = 35$, (2) $5x - 3y = -8$. Show all sign changes.

Q545. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 19$, (2) $4x - 1y = -19$. Show all sign changes.

Q546. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 15$, (2) $4x - 5y = 16$. Show all sign changes.

Q547. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 2y = 22$, (2) $3x - 5y = -2$. Show all sign changes.

Q548. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 23$, (2) $4x - 1y = -4$. Show all sign changes.

Q549. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 7y = 16$, (2) $2x - 2y = 18$. Show all sign changes.

Q550. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 2y = 34$, (2) $4x - 3y = 4$. Show all sign changes.

Q551. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 2y = 31$, (2) $3x - 7y = -16$. Show all sign changes.

Q552. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 28$, (2) $7x - 6y = 18$. Show all sign changes.

Q553. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 6y = 47$, (2) $2x - 3y = -8$. Show all sign changes.

Q554. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 4y = 48$, (2) $3x - 2y = -14$. Show all sign changes.

Q555. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 7y = 25$, (2) $2x - 7y = -5$. Show all sign changes.

Q556. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 2y = 47$, (2) $7x - 4y = -13$. Show all sign changes.

Q557. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 3y = 34$, (2) $3x - 4y = 15$. Show all sign changes.

Q558. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 23$, (2) $3x - 1y = -19$. Show all sign changes.

Q559. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 4y = 18$, (2) $5x - 7y = 9$. Show all sign changes.

Q560. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 5y = 23$, (2) $2x - 5y = 3$. Show all sign changes.

Q561. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 7y = 21$, (2) $6x - 1y = -13$. Show all sign changes.

Q562. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 5y = 31$, (2) $2x - 6y = 1$. Show all sign changes.

Q563. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 4y = 28$, (2) $3x - 7y = -5$. Show all sign changes.

Q564. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 6y = 23$, (2) $3x - 1y = -15$. Show all sign changes.

Q565. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 7y = 23$, (2) $6x - 1y = 11$. Show all sign changes.

Q566. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 7y = 34$, (2) $2x - 4y = 16$. Show all sign changes.

Q567. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 3y = 18$, (2) $7x - 1y = -5$. Show all sign changes.

Q568. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 4y = 14$, (2) $2x - 6y = -14$. Show all sign changes.

Q569. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 30$, (2) $7x - 2y = 11$. Show all sign changes.

Q570. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 4y = 41$, (2) $6x - 1y = -17$. Show all sign changes.

Q571. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 11$, (2) $7x - 4y = 5$. Show all sign changes.

Q572. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 3y = 37$, (2) $5x - 1y = -18$. Show all sign changes.

Q573. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 2y = 35$, (2) $6x - 2y = 6$. Show all sign changes.

Q574. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 4y = 36$, (2) $4x - 2y = 0$. Show all sign changes.

Q575. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 7y = 11$, (2) $2x - 4y = 1$. Show all sign changes.

Q576. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 2y = 12$, (2) $4x - 5y = -11$. Show all sign changes.

Q577. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 4y = 23$, (2) $3x - 2y = -11$. Show all sign changes.

Q578. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 3y = 48$, (2) $5x - 5y = -17$. Show all sign changes.

Q579. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 33$, (2) $7x - 1y = 11$. Show all sign changes.

Q580. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 25$, (2) $3x - 5y = 9$. Show all sign changes.

Q581. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 7y = 29$, (2) $4x - 4y = -2$. Show all sign changes.

Q582. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 45$, (2) $2x - 1y = -13$. Show all sign changes.

Q583. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 39$, (2) $3x - 3y = 8$. Show all sign changes.

Q584. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 4y = 46$, (2) $4x - 1y = 15$. Show all sign changes.

Q585. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 4y = 24$, (2) $2x - 4y = 9$. Show all sign changes.

Q586. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 5y = 42$, (2) $5x - 1y = -6$. Show all sign changes.

Q587. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 48$, (2) $3x - 1y = -9$. Show all sign changes.

Q588. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 4y = 11$, (2) $6x - 2y = 1$. Show all sign changes.

Q589. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 4y = 27$, (2) $6x - 2y = -12$. Show all sign changes.

Q590. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 3y = 27$, (2) $5x - 2y = -3$. Show all sign changes.

Q591. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 7y = 29$, (2) $5x - 4y = 1$. Show all sign changes.

Q592. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 38$, (2) $3x - 5y = 14$. Show all sign changes.

Q593. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 5y = 39$, (2) $2x - 1y = 15$. Show all sign changes.

Q594. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 25$, (2) $2x - 5y = 1$. Show all sign changes.

Q595. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 26$, (2) $2x - 7y = -16$. Show all sign changes.

Q596. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 3y = 12$, (2) $6x - 6y = -9$. Show all sign changes.

Q597. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 34$, (2) $7x - 4y = -1$. Show all sign changes.

Q598. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 45$, (2) $6x - 5y = 7$. Show all sign changes.

Q599. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 14$, (2) $6x - 4y = -19$. Show all sign

changes.

Q600. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 7y = 40$, (2) $3x - 7y = 16$. Show all sign changes.

Q601. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 5y = 27$, (2) $7x - 7y = -7$. Show all sign changes.

Q602. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 25$, (2) $2x - 4y = 6$. Show all sign changes.

Q603. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 7y = 34$, (2) $7x - 4y = 13$. Show all sign changes.

Q604. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 21$, (2) $4x - 2y = -14$. Show all sign changes.

Q605. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 4y = 12$, (2) $3x - 1y = 2$. Show all sign changes.

Q606. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 6y = 32$, (2) $4x - 3y = -10$. Show all sign changes.

Q607. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 29$, (2) $6x - 6y = 11$. Show all sign changes.

Q608. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 5y = 47$, (2) $6x - 1y = -2$. Show all sign changes.

Q609. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 6y = 30$, (2) $5x - 2y = -15$. Show all sign changes.

Q610. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 2y = 17$, (2) $2x - 1y = -13$. Show all sign changes.

Q611. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 2y = 36$, (2) $5x - 3y = 9$. Show all sign changes.

Q612. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 3y = 31$, (2) $7x - 2y = 5$. Show all sign changes.

Q613. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 7y = 20$, (2) $5x - 6y = 2$. Show all sign changes.

Q614. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 13$, (2) $2x - 2y = -14$. Show all sign changes.

Q615. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 5y = 42$, (2) $7x - 4y = 15$. Show all sign changes.

Q616. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 2y = 50$, (2) $5x - 5y = -7$. Show all sign changes.

Q617. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 2y = 24$, (2) $6x - 4y = 15$. Show all sign changes.

Q618. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 5y = 12$, (2) $7x - 4y = 6$. Show all sign changes.

Q619. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 7y = 32$, (2) $4x - 2y = 14$. Show all sign changes.

Q620. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 7y = 50$, (2) $5x - 2y = 6$. Show all sign changes.

Q621. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 5y = 17$, (2) $3x - 4y = 18$. Show all sign changes.

Q622. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x + 5y = 34$, (2) $5x - 6y = -1$. Show all sign changes.

Q623. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 6y = 26$, (2) $3x - 5y = 13$. Show all sign changes.

Q624. Subtract Equation (2) from Equation (1) to eliminate x: (1) $3x + 3y = 19$, (2) $3x - 7y = -19$. Show all sign changes.

Q625. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 25$, (2) $7x - 6y = -3$. Show all sign changes.

Q626. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 30$, (2) $2x - 4y = -18$. Show all sign changes.

Q627. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 3y = 14$, (2) $6x - 5y = -8$. Show all sign changes.

Q628. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 2y = 22$, (2) $7x - 6y = 8$. Show all sign changes.

Q629. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 5y = 50$, (2) $2x - 7y = -20$. Show all sign changes.

Q630. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x + 5y = 15$, (2) $4x - 5y = -20$. Show all sign changes.

Q631. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 6y = 29$, (2) $6x - 5y = -1$. Show all sign changes.

Q632. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 4y = 31$, (2) $6x - 3y = 9$. Show all sign changes.

Q633. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x + 7y = 28$, (2) $6x - 6y = 15$. Show all sign changes.

Q634. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 3y = 47$, (2) $7x - 1y = 17$. Show all sign changes.

Q635. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x + 3y = 48$, (2) $2x - 4y = 5$. Show all sign changes.

Q636. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 4y = 29$, (2) $7x - 1y = 6$. Show all sign changes.

Q637. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x + 6y = 47$, (2) $7x - 6y = -6$. Show all sign changes.

Q638. Subtract Equation (2) from Equation (1) to elimi-

nate x: (1) $6x+3y=24$, (2) $6x-1y=-13$. Show all sign changes.

Q639. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x+7y=15$, (2) $2x-3y=-2$. Show all sign changes.

Q640. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x+2y=14$, (2) $6x-7y=-19$. Show all sign changes.

Q641. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x+5y=19$, (2) $6x-2y=14$. Show all sign changes.

Q642. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x+4y=46$, (2) $2x-4y=-5$. Show all sign changes.

Q643. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x+5y=12$, (2) $4x-5y=18$. Show all sign changes.

Q644. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x+4y=50$, (2) $7x-2y=-19$. Show all sign changes.

Q645. Subtract Equation (2) from Equation (1) to eliminate x: (1) $6x+6y=50$, (2) $6x-4y=6$. Show all sign changes.

Q646. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x+2y=40$, (2) $5x-4y=-5$. Show all sign changes.

Q647. Subtract Equation (2) from Equation (1) to eliminate x: (1) $4x+2y=11$, (2) $4x-5y=-20$. Show all sign changes.

Q648. Subtract Equation (2) from Equation (1) to eliminate x: (1) $5x+6y=48$, (2) $5x-1y=15$. Show all sign changes.

Q649. Subtract Equation (2) from Equation (1) to eliminate x: (1) $7x+6y=16$, (2) $7x-5y=4$. Show all sign changes.

Q650. Subtract Equation (2) from Equation (1) to eliminate x: (1) $2x+5y=21$, (2) $2x-1y=15$. Show all sign changes.

Q651. For $x^2-9x-2=0$, substitute $a=1, b=-9, c=-2$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q652. For $x^2-7x-5=0$, substitute $a=1, b=-7, c=-5$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q653. For $x^2-3x-3=0$, substitute $a=1, b=-3, c=-3$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q654. For $x^2-9x-11=0$, substitute $a=1, b=-9, c=-11$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q655. For $x^2-3x-12=0$, substitute $a=1, b=-3, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q656. For $x^2-10x-1=0$, substitute $a=1, b=-10, c=-1$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q657. For $x^2-6x-11=0$, substitute $a=1, b=-6, c=-11$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q658. For $x^2-2x-6=0$, substitute $a=1, b=-2, c=-6$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q659. For $x^2-9x-10=0$, substitute $a=1, b=-9, c=-10$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q660. For $x^2-10x-1=0$, substitute $a=1, b=-10, c=-1$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q661. For $x^2-9x-9=0$, substitute $a=1, b=-9, c=-9$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q662. For $x^2-10x-10=0$, substitute $a=1, b=-10, c=-10$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q663. For $x^2-1x-3=0$, substitute $a=1, b=-1, c=-3$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q664. For $x^2-6x-8=0$, substitute $a=1, b=-6, c=-8$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q665. For $x^2-6x-3=0$, substitute $a=1, b=-6, c=-3$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q666. For $x^2-7x-4=0$, substitute $a=1, b=-7, c=-4$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q667. For $x^2-5x-8=0$, substitute $a=1, b=-5, c=-8$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q668. For $x^2-3x-12=0$, substitute $a=1, b=-3, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q669. For $x^2-10x-7=0$, substitute $a=1, b=-10, c=-7$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q670. For $x^2-7x-2=0$, substitute $a=1, b=-7, c=-2$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q671. For $x^2-2x-9=0$, substitute $a=1, b=-2, c=-9$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q672. For $x^2-5x-9=0$, substitute $a=1, b=-5, c=-9$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q673. For $x^2-4x-12=0$, substitute $a=1, b=-4, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q674. For $x^2-8x-2=0$, substitute $a=1, b=-8, c=-2$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q675. For $x^2-8x-10=0$, substitute $a=1, b=-8, c=-10$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q676. For $x^2-1x-10=0$, substitute $a=1, b=-1, c=-10$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q677. For $x^2-5x-2=0$, substitute $a=1, b=-5, c=-2$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q678. For $x^2-10x-5=0$, substitute $a=1, b=-10, c=-5$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q679. For $x^2-10x-12=0$, substitute $a=1, b=-10, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q680. For $x^2-2x-11=0$, substitute $a=1, b=-2, c=-11$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q681. For $x^2-1x-8=0$, substitute $a=1, b=-1, c=-8$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q682. For $x^2-2x-12=0$, substitute $a=1, b=-2, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q683. For $x^2-7x-6=0$, substitute $a=1, b=-7, c=-6$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q684. For $x^2-1x-12=0$, substitute $a=1, b=-1, c=-12$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q685. For $x^2-9x-8=0$, substitute $a=1, b=-9, c=-8$

into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q686. For $x^2-6x-2=0$, substitute $a=1, b=-6, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q687. For $x^2-9x-8=0$, substitute $a=1, b=-9, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q688. For $x^2-4x-4=0$, substitute $a=1, b=-4, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q689. For $x^2-5x-2=0$, substitute $a=1, b=-5, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q690. For $x^2-6x-12=0$, substitute $a=1, b=-6, c=-12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q691. For $x^2-8x-1=0$, substitute $a=1, b=-8, c=-1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q692. For $x^2-3x-12=0$, substitute $a=1, b=-3, c=-12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q693. For $x^2-3x-1=0$, substitute $a=1, b=-3, c=-1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q694. For $x^2-6x-9=0$, substitute $a=1, b=-6, c=-9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q695. For $x^2-10x-8=0$, substitute $a=1, b=-10, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q696. For $x^2-10x-8=0$, substitute $a=1, b=-10, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q697. For $x^2-2x-9=0$, substitute $a=1, b=-2, c=-9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q698. For $x^2-3x-11=0$, substitute $a=1, b=-3, c=-11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q699. For $x^2-10x-9=0$, substitute $a=1, b=-10, c=-9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q700. For $x^2-5x-3=0$, substitute $a=1, b=-5, c=-3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q701. For $x^2-8x-8=0$, substitute $a=1, b=-8, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q702. For $x^2-6x-10=0$, substitute $a=1, b=-6, c=-10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q703. For $x^2-10x-8=0$, substitute $a=1, b=-10, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q704. For $x^2-7x-6=0$, substitute $a=1, b=-7, c=-6$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q705. For $x^2-6x-7=0$, substitute $a=1, b=-6, c=-7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q706. For $x^2-8x-2=0$, substitute $a=1, b=-8, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q707. For $x^2-4x-7=0$, substitute $a=1, b=-4, c=-7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q708. For $x^2-2x-9=0$, substitute $a=1, b=-2, c=-9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q709. For $x^2-6x-11=0$, substitute $a=1, b=-6, c=-11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q710. For $x^2-3x-10=0$, substitute $a=1, b=-3, c=-10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q711. For $x^2-9x-7=0$, substitute $a=1, b=-9, c=-7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q712. For $x^2-10x-11=0$, substitute $a=1, b=-10, c=-11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q713. For $x^2-6x-8=0$, substitute $a=1, b=-6, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q714. For $x^2-9x-8=0$, substitute $a=1, b=-9, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q715. For $x^2-4x-2=0$, substitute $a=1, b=-4, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q716. For $x^2-9x-12=0$, substitute $a=1, b=-9, c=-12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q717. For $x^2-5x-7=0$, substitute $a=1, b=-5, c=-7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q718. For $x^2-1x-10=0$, substitute $a=1, b=-1, c=-10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q719. For $x^2-1x-3=0$, substitute $a=1, b=-1, c=-3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q720. For $x^2-5x-7=0$, substitute $a=1, b=-5, c=-7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q721. For $x^2-8x-6=0$, substitute $a=1, b=-8, c=-6$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q722. For $x^2-9x-2=0$, substitute $a=1, b=-9, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q723. For $x^2-5x-10=0$, substitute $a=1, b=-5, c=-10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q724. For $x^2-5x-1=0$, substitute $a=1, b=-5, c=-1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q725. For $x^2-3x-5=0$, substitute $a=1, b=-3, c=-5$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q726. For $x^2-7x-12=0$, substitute $a=1, b=-7, c=-12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q727. For $x^2-7x-2=0$, substitute $a=1, b=-7, c=-2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q728. For $x^2-1x-10=0$, substitute $a=1, b=-1, c=-10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q729. For $x^2-2x-4=0$, substitute $a=1, b=-2, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q730. For $x^2-1x-3=0$, substitute $a=1, b=-1, c=-3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q731. For $x^2-3x-4=0$, substitute $a=1, b=-3, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q732. For $x^2-2x-8=0$, substitute $a=1, b=-2, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q733. For $x^2-10x-4=0$, substitute $a=1, b=-10, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q734. For $x^2-10x-8=0$, substitute $a=1, b=-10, c=-8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q735. For $x^2-6x-4=0$, substitute $a=1, b=-6, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q736. For $x^2-3x-4=0$, substitute $a=1, b=-3, c=-4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q737. For $x^2 - 3x - 12 = 0$, substitute $a = 1, b = -3, c = -12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q738. For $x^2 - 10x - 8 = 0$, substitute $a = 1, b = -10, c = -8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q739. For $x^2 - 1x - 1 = 0$, substitute $a = 1, b = -1, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q740. For $x^2 - 7x - 2 = 0$, substitute $a = 1, b = -7, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q741. For $x^2 - 2x - 6 = 0$, substitute $a = 1, b = -2, c = -6$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q742. For $x^2 - 4x - 10 = 0$, substitute $a = 1, b = -4, c = -10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q743. For $x^2 - 1x - 12 = 0$, substitute $a = 1, b = -1, c = -12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q744. For $x^2 - 5x - 8 = 0$, substitute $a = 1, b = -5, c = -8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q745. For $x^2 - 8x - 3 = 0$, substitute $a = 1, b = -8, c = -3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q746. For $x^2 - 1x - 1 = 0$, substitute $a = 1, b = -1, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q747. For $x^2 - 1x - 7 = 0$, substitute $a = 1, b = -1, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q748. For $x^2 - 5x - 2 = 0$, substitute $a = 1, b = -5, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q749. For $x^2 - 5x - 11 = 0$, substitute $a = 1, b = -5, c = -11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q750. For $x^2 - 10x - 9 = 0$, substitute $a = 1, b = -10, c = -9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q751. For $x^2 - 6x - 9 = 0$, substitute $a = 1, b = -6, c = -9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q752. For $x^2 - 6x - 4 = 0$, substitute $a = 1, b = -6, c = -4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q753. For $x^2 - 6x - 5 = 0$, substitute $a = 1, b = -6, c = -5$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q754. For $x^2 - 1x - 2 = 0$, substitute $a = 1, b = -1, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q755. For $x^2 - 2x - 7 = 0$, substitute $a = 1, b = -2, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q756. For $x^2 - 7x - 7 = 0$, substitute $a = 1, b = -7, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q757. For $x^2 - 5x - 3 = 0$, substitute $a = 1, b = -5, c = -3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q758. For $x^2 - 9x - 3 = 0$, substitute $a = 1, b = -9, c = -3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q759. For $x^2 - 1x - 5 = 0$, substitute $a = 1, b = -1, c = -5$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q760. For $x^2 - 5x - 8 = 0$, substitute $a = 1, b = -5, c = -8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q761. For $x^2 - 8x - 2 = 0$, substitute $a = 1, b = -8, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q762. For $x^2 - 4x - 8 = 0$, substitute $a = 1, b = -4, c = -8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q763. For $x^2 - 1x - 12 = 0$, substitute $a = 1, b = -1, c = -12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q764. For $x^2 - 6x - 6 = 0$, substitute $a = 1, b = -6, c = -6$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q765. For $x^2 - 9x - 2 = 0$, substitute $a = 1, b = -9, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q766. For $x^2 - 2x - 9 = 0$, substitute $a = 1, b = -2, c = -9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q767. For $x^2 - 4x - 6 = 0$, substitute $a = 1, b = -4, c = -6$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q768. For $x^2 - 8x - 1 = 0$, substitute $a = 1, b = -8, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q769. For $x^2 - 1x - 1 = 0$, substitute $a = 1, b = -1, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q770. For $x^2 - 4x - 2 = 0$, substitute $a = 1, b = -4, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q771. For $x^2 - 9x - 4 = 0$, substitute $a = 1, b = -9, c = -4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q772. For $x^2 - 7x - 8 = 0$, substitute $a = 1, b = -7, c = -8$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q773. For $x^2 - 1x - 11 = 0$, substitute $a = 1, b = -1, c = -11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q774. For $x^2 - 1x - 9 = 0$, substitute $a = 1, b = -1, c = -9$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q775. For $x^2 - 5x - 1 = 0$, substitute $a = 1, b = -5, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q776. For $x^2 - 5x - 7 = 0$, substitute $a = 1, b = -5, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q777. For $x^2 - 4x - 11 = 0$, substitute $a = 1, b = -4, c = -11$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q778. For $x^2 - 9x - 3 = 0$, substitute $a = 1, b = -9, c = -3$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q779. For $x^2 - 6x - 7 = 0$, substitute $a = 1, b = -6, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q780. For $x^2 - 4x - 10 = 0$, substitute $a = 1, b = -4, c = -10$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q781. For $x^2 - 1x - 12 = 0$, substitute $a = 1, b = -1, c = -12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q782. For $x^2 - 6x - 1 = 0$, substitute $a = 1, b = -6, c = -1$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q783. For $x^2 - 1x - 4 = 0$, substitute $a = 1, b = -1, c = -4$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q784. For $x^2 - 6x - 2 = 0$, substitute $a = 1, b = -6, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q785. For $x^2 - 10x - 12 = 0$, substitute $a = 1, b = -10, c = -12$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q786. For $x^2 - 3x - 7 = 0$, substitute $a = 1, b = -3, c = -7$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q787. For $x^2 - 5x - 2 = 0$, substitute $a = 1, b = -5, c = -2$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q788. For $x^2 - 3x - 5 = 0$, substitute $a = 1, b = -3, c = -5$ into $x = \frac{-b \pm \sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q789. For $x^2-7x-9=0$, substitute $a=1, b=-7, c=-9$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q790. For $x^2-3x-2=0$, substitute $a=1, b=-3, c=-2$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q791. For $x^2-8x-4=0$, substitute $a=1, b=-8, c=-4$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q792. For $x^2-8x-1=0$, substitute $a=1, b=-8, c=-1$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q793. For $x^2-9x-7=0$, substitute $a=1, b=-9, c=-7$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q794. For $x^2-6x-1=0$, substitute $a=1, b=-6, c=-1$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q795. For $x^2-10x-4=0$, substitute $a=1, b=-10, c=-4$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q796. For $x^2-10x-10=0$, substitute $a=1, b=-10, c=-10$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q797. For $x^2-10x-1=0$, substitute $a=1, b=-10, c=-1$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q798. For $x^2-7x-8=0$, substitute $a=1, b=-7, c=-8$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q799. For $x^2-8x-4=0$, substitute $a=1, b=-8, c=-4$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q800. For $x^2-3x-11=0$, substitute $a=1, b=-3, c=-11$ into $x=\frac{-b\pm\sqrt{b^2-4ac}}{2a}$. Carefully simplify $-b$ and $-4ac$.

Q801. Translate into an algebraic expression and simplify: 'Subtract $(3x-5)$ from $(5x+3)$'. Identify which term comes first!

Q802. Translate into an algebraic expression and simplify: 'Subtract $(6x-7)$ from $(7x+6)$'. Identify which term comes first!

Q803. Translate into an algebraic expression and simplify: 'Subtract $(9x-14)$ from $(14x+9)$'. Identify which term comes first!

Q804. Translate into an algebraic expression and simplify: 'Subtract $(3x-10)$ from $(10x+3)$'. Identify which term comes first!

Q805. Translate into an algebraic expression and simplify: 'Subtract $(9x-11)$ from $(11x+9)$'. Identify which term comes first!

Q806. Translate into an algebraic expression and simplify: 'Subtract $(7x-6)$ from $(6x+7)$'. Identify which term comes first!

Q807. Translate into an algebraic expression and simplify: 'Subtract $(2x-5)$ from $(5x+2)$'. Identify which term comes first!

Q808. Translate into an algebraic expression and simplify: 'Subtract $(6x-13)$ from $(13x+6)$'. Identify which term comes first!

Q809. Translate into an algebraic expression and simplify: 'Subtract $(2x-9)$ from $(9x+2)$'. Identify which term comes first!

Q810. Translate into an algebraic expression and simplify: 'Subtract $(9x-6)$ from $(6x+9)$'. Identify which term comes first!

Q811. Translate into an algebraic expression and simplify: 'Subtract $(8x-7)$ from $(7x+8)$'. Identify which term comes first!

Q812. Translate into an algebraic expression and simplify: 'Subtract $(6x-6)$ from $(6x+6)$'. Identify which term comes first!

Q813. Translate into an algebraic expression and simplify: 'Subtract $(7x-14)$ from $(14x+7)$'. Identify which term comes first!

Q814. Translate into an algebraic expression and simplify: 'Subtract $(8x-4)$ from $(4x+8)$'. Identify which term comes first!

Q815. Translate into an algebraic expression and simplify: 'Subtract $(2x-9)$ from $(9x+2)$'. Identify which term comes first!

Q816. Translate into an algebraic expression and simplify: 'Subtract $(8x-5)$ from $(5x+8)$'. Identify which term comes first!

Q817. Translate into an algebraic expression and simplify: 'Subtract $(8x-12)$ from $(12x+8)$'. Identify which term comes first!

Q818. Translate into an algebraic expression and simplify: 'Subtract $(6x-11)$ from $(11x+6)$'. Identify which term comes first!

Q819. Translate into an algebraic expression and simplify: 'Subtract $(4x-5)$ from $(5x+4)$'. Identify which term comes first!

Q820. Translate into an algebraic expression and simplify: 'Subtract $(7x-14)$ from $(14x+7)$'. Identify which term comes first!

Q821. Translate into an algebraic expression and simplify: 'Subtract $(4x-13)$ from $(13x+4)$'. Identify which term comes first!

Q822. Translate into an algebraic expression and simplify: 'Subtract $(6x-4)$ from $(4x+6)$'. Identify which term comes first!

Q823. Translate into an algebraic expression and simplify: 'Subtract $(9x-6)$ from $(6x+9)$'. Identify which term comes first!

Q824. Translate into an algebraic expression and simplify: 'Subtract $(4x-3)$ from $(3x+4)$'. Identify which term comes first!

Q825. Translate into an algebraic expression and simplify: 'Subtract $(5x-14)$ from $(14x+5)$'. Identify which term comes first!

Q826. Translate into an algebraic expression and simplify: 'Subtract $(9x-12)$ from $(12x+9)$'. Identify which term comes first!

Q827. Translate into an algebraic expression and simplify: 'Subtract $(2x-4)$ from $(4x+2)$'. Identify which term comes first!

Q828. Translate into an algebraic expression and simplify: 'Subtract $(8x-12)$ from $(12x+8)$'. Identify which term comes first!

Q829. Translate into an algebraic expression and simplify: 'Subtract $(7x-11)$ from $(11x+7)$'. Identify which term comes first!

Q830. Translate into an algebraic expression and sim-

plify: 'Subtract $(6x - 9)$ from $(9x + 6)$'. Identify which term comes first!

Q831. Translate into an algebraic expression and simplify: 'Subtract $(2x - 7)$ from $(7x + 2)$'. Identify which term comes first!

Q832. Translate into an algebraic expression and simplify: 'Subtract $(9x - 15)$ from $(15x + 9)$'. Identify which term comes first!

Q833. Translate into an algebraic expression and simplify: 'Subtract $(6x - 11)$ from $(11x + 6)$'. Identify which term comes first!

Q834. Translate into an algebraic expression and simplify: 'Subtract $(2x - 10)$ from $(10x + 2)$'. Identify which term comes first!

Q835. Translate into an algebraic expression and simplify: 'Subtract $(9x - 6)$ from $(6x + 9)$'. Identify which term comes first!

Q836. Translate into an algebraic expression and simplify: 'Subtract $(6x - 7)$ from $(7x + 6)$'. Identify which term comes first!

Q837. Translate into an algebraic expression and simplify: 'Subtract $(5x - 11)$ from $(11x + 5)$'. Identify which term comes first!

Q838. Translate into an algebraic expression and simplify: 'Subtract $(6x - 11)$ from $(11x + 6)$'. Identify which term comes first!

Q839. Translate into an algebraic expression and simplify: 'Subtract $(9x - 12)$ from $(12x + 9)$'. Identify which term comes first!

Q840. Translate into an algebraic expression and simplify: 'Subtract $(3x - 12)$ from $(12x + 3)$'. Identify which term comes first!

Q841. Translate into an algebraic expression and simplify: 'Subtract $(7x - 12)$ from $(12x + 7)$'. Identify which term comes first!

Q842. Translate into an algebraic expression and simplify: 'Subtract $(7x - 8)$ from $(8x + 7)$'. Identify which term comes first!

Q843. Translate into an algebraic expression and simplify: 'Subtract $(7x - 10)$ from $(10x + 7)$'. Identify which term comes first!

Q844. Translate into an algebraic expression and simplify: 'Subtract $(5x - 5)$ from $(5x + 5)$'. Identify which term comes first!

Q845. Translate into an algebraic expression and simplify: 'Subtract $(2x - 5)$ from $(5x + 2)$'. Identify which term comes first!

Q846. Translate into an algebraic expression and simplify: 'Subtract $(6x - 11)$ from $(11x + 6)$'. Identify which term comes first!

Q847. Translate into an algebraic expression and simplify: 'Subtract $(8x - 14)$ from $(14x + 8)$'. Identify which term comes first!

Q848. Translate into an algebraic expression and simplify: 'Subtract $(8x - 10)$ from $(10x + 8)$'. Identify which term comes first!

Q849. Translate into an algebraic expression and simplify: 'Subtract $(9x - 4)$ from $(4x + 9)$'. Identify which term comes first!

Q850. Translate into an algebraic expression and simplify: 'Subtract $(7x - 12)$ from $(12x + 7)$'. Identify which term comes first!

Q851. Translate into an algebraic expression and simplify: 'Subtract $(8x - 5)$ from $(5x + 8)$'. Identify which term comes first!

Q852. Translate into an algebraic expression and simplify: 'Subtract $(8x - 8)$ from $(8x + 8)$'. Identify which term comes first!

Q853. Translate into an algebraic expression and simplify: 'Subtract $(4x - 10)$ from $(10x + 4)$'. Identify which term comes first!

Q854. Translate into an algebraic expression and simplify: 'Subtract $(9x - 8)$ from $(8x + 9)$'. Identify which term comes first!

Q855. Translate into an algebraic expression and simplify: 'Subtract $(8x - 13)$ from $(13x + 8)$'. Identify which term comes first!

Q856. Translate into an algebraic expression and simplify: 'Subtract $(5x - 9)$ from $(9x + 5)$'. Identify which term comes first!

Q857. Translate into an algebraic expression and simplify: 'Subtract $(7x - 6)$ from $(6x + 7)$'. Identify which term comes first!

Q858. Translate into an algebraic expression and simplify: 'Subtract $(7x - 15)$ from $(15x + 7)$'. Identify which term comes first!

Q859. Translate into an algebraic expression and simplify: 'Subtract $(7x - 9)$ from $(9x + 7)$'. Identify which term comes first!

Q860. Translate into an algebraic expression and simplify: 'Subtract $(2x - 15)$ from $(15x + 2)$'. Identify which term comes first!

Q861. Translate into an algebraic expression and simplify: 'Subtract $(3x - 10)$ from $(10x + 3)$'. Identify which term comes first!

Q862. Translate into an algebraic expression and simplify: 'Subtract $(6x - 7)$ from $(7x + 6)$'. Identify which term comes first!

Q863. Translate into an algebraic expression and simplify: 'Subtract $(7x - 13)$ from $(13x + 7)$'. Identify which term comes first!

Q864. Translate into an algebraic expression and simplify: 'Subtract $(6x - 4)$ from $(4x + 6)$'. Identify which term comes first!

Q865. Translate into an algebraic expression and simplify: 'Subtract $(4x - 5)$ from $(5x + 4)$'. Identify which term comes first!

Q866. Translate into an algebraic expression and simplify: 'Subtract $(3x - 8)$ from $(8x + 3)$'. Identify which term comes first!

Q867. Translate into an algebraic expression and simplify: 'Subtract $(9x - 8)$ from $(8x + 9)$'. Identify which term comes first!

Q868. Translate into an algebraic expression and simplify: 'Subtract $(2x - 5)$ from $(5x + 2)$'. Identify which term comes first!

Q869. Translate into an algebraic expression and simplify: 'Subtract $(4x - 14)$ from $(14x + 4)$'. Identify which term comes first!

Q870. Translate into an algebraic expression and simplify: 'Subtract $(8x - 12)$ from $(12x + 8)$'. Identify which term comes first!

Q871. Translate into an algebraic expression and simplify: 'Subtract $(4x - 14)$ from $(14x + 4)$'. Identify which term comes first!

Q872. Translate into an algebraic expression and simplify: 'Subtract $(3x - 15)$ from $(15x + 3)$'. Identify which term comes first!

Q873. Translate into an algebraic expression and simplify: 'Subtract $(2x - 11)$ from $(11x + 2)$'. Identify which term comes first!

Q874. Translate into an algebraic expression and simplify: 'Subtract $(5x - 7)$ from $(7x + 5)$'. Identify which term comes first!

Q875. Translate into an algebraic expression and simplify: 'Subtract $(8x - 4)$ from $(4x + 8)$'. Identify which term comes first!

Q876. Translate into an algebraic expression and simplify: 'Subtract $(9x - 6)$ from $(6x + 9)$'. Identify which term comes first!

Q877. Translate into an algebraic expression and simplify: 'Subtract $(5x - 4)$ from $(4x + 5)$'. Identify which term comes first!

Q878. Translate into an algebraic expression and simplify: 'Subtract $(8x - 10)$ from $(10x + 8)$'. Identify which term comes first!

Q879. Translate into an algebraic expression and simplify: 'Subtract $(4x - 6)$ from $(6x + 4)$'. Identify which term comes first!

Q880. Translate into an algebraic expression and simplify: 'Subtract $(5x - 13)$ from $(13x + 5)$'. Identify which term comes first!

Q881. Translate into an algebraic expression and simplify: 'Subtract $(9x - 14)$ from $(14x + 9)$'. Identify which term comes first!

Q882. Translate into an algebraic expression and simplify: 'Subtract $(5x - 8)$ from $(8x + 5)$'. Identify which term comes first!

Q883. Translate into an algebraic expression and simplify: 'Subtract $(8x - 8)$ from $(8x + 8)$'. Identify which term comes first!

Q884. Translate into an algebraic expression and simplify: 'Subtract $(6x - 11)$ from $(11x + 6)$'. Identify which term comes first!

Q885. Translate into an algebraic expression and simplify: 'Subtract $(8x - 5)$ from $(5x + 8)$'. Identify which term comes first!

Q886. Translate into an algebraic expression and simplify: 'Subtract $(4x - 12)$ from $(12x + 4)$'. Identify which term comes first!

Q887. Translate into an algebraic expression and simplify: 'Subtract $(5x - 10)$ from $(10x + 5)$'. Identify which term comes first!

Q888. Translate into an algebraic expression and simplify: 'Subtract $(7x - 4)$ from $(4x + 7)$'. Identify which term comes first!

Q889. Translate into an algebraic expression and simplify: 'Subtract $(2x - 11)$ from $(11x + 2)$'. Identify which term comes first!

Q890. Translate into an algebraic expression and simplify: 'Subtract $(4x - 8)$ from $(8x + 4)$'. Identify which term comes first!

Q891. Translate into an algebraic expression and simplify: 'Subtract $(7x - 5)$ from $(5x + 7)$'. Identify which term comes first!

Q892. Translate into an algebraic expression and simplify: 'Subtract $(8x - 10)$ from $(10x + 8)$'. Identify which term comes first!

Q893. Translate into an algebraic expression and simplify: 'Subtract $(8x - 9)$ from $(9x + 8)$'. Identify which term comes first!

Q894. Translate into an algebraic expression and simplify: 'Subtract $(5x - 3)$ from $(3x + 5)$'. Identify which term comes first!

Q895. Translate into an algebraic expression and simplify: 'Subtract $(8x - 7)$ from $(7x + 8)$'. Identify which term comes first!

Q896. Translate into an algebraic expression and simplify: 'Subtract $(3x - 12)$ from $(12x + 3)$'. Identify which term comes first!

Q897. Translate into an algebraic expression and simplify: 'Subtract $(3x - 3)$ from $(3x + 3)$'. Identify which term comes first!

Q898. Translate into an algebraic expression and simplify: 'Subtract $(8x - 12)$ from $(12x + 8)$'. Identify which term comes first!

Q899. Translate into an algebraic expression and simplify: 'Subtract $(5x - 3)$ from $(3x + 5)$'. Identify which term comes first!

Q900. Translate into an algebraic expression and simplify: 'Subtract $(7x - 4)$ from $(4x + 7)$'. Identify which term comes first!

Q901. Solve for x: '$-7x + 9 = 19$. What happens to the signs when you divide the final step by -7?

Q902. Solve for x: '$-5x + 19 = 4$. What happens to the signs when you divide the final step by -5?

Q903. Solve for x: '$-7x + 19 = -16$. What happens to the signs when you divide the final step by -7?

Q904. Solve for x: '$-6x + 9 = 10$. What happens to the signs when you divide the final step by -6?

Q905. Solve for x: '$-7x + 12 = -8$. What happens to the signs when you divide the final step by -7?

Q906. Solve for x: '$-4x + 18 = -18$. What happens to the signs when you divide the final step by -4?

Q907. Solve for x: '$-6x + 30 = 15$. What happens to the signs when you divide the final step by -6?

Q908. Solve for x: '$-4x + 30 = 12$. What happens to the signs when you divide the final step by -4?

Q909. Solve for x: '$-5x + 26 = 6$. What happens to the signs when you divide the final step by -5?

Q910. Solve for x: '$-8x + 27 = 14$. What happens to the

signs when you divide the final step by -8?

Q911. Solve for x: $-9x + 13 = 3$. What happens to the signs when you divide the final step by -9?

Q912. Solve for x: $-7x + 7 = 12$. What happens to the signs when you divide the final step by -7?

Q913. Solve for x: $-2x + 21 = 2$. What happens to the signs when you divide the final step by -2?

Q914. Solve for x: $-7x + 16 = -1$. What happens to the signs when you divide the final step by -7?

Q915. Solve for x: $-4x + 10 = 13$. What happens to the signs when you divide the final step by -4?

Q916. Solve for x: $-7x + 29 = -5$. What happens to the signs when you divide the final step by -7?

Q917. Solve for x: $-6x + 13 = 12$. What happens to the signs when you divide the final step by -6?

Q918. Solve for x: $-8x + 24 = -6$. What happens to the signs when you divide the final step by -8?

Q919. Solve for x: $-5x + 17 = -12$. What happens to the signs when you divide the final step by -5?

Q920. Solve for x: $-7x + 18 = -4$. What happens to the signs when you divide the final step by -7?

Q921. Solve for x: $-4x + 15 = 3$. What happens to the signs when you divide the final step by -4?

Q922. Solve for x: $-3x + 24 = -12$. What happens to the signs when you divide the final step by -3?

Q923. Solve for x: $-8x + 14 = 0$. What happens to the signs when you divide the final step by -8?

Q924. Solve for x: $-5x + 17 = -19$. What happens to the signs when you divide the final step by -5?

Q925. Solve for x: $-5x + 23 = -4$. What happens to the signs when you divide the final step by -5?

Q926. Solve for x: $-9x + 11 = 9$. What happens to the signs when you divide the final step by -9?

Q927. Solve for x: $-5x + 8 = 2$. What happens to the signs when you divide the final step by -5?

Q928. Solve for x: $-7x + 14 = -19$. What happens to the signs when you divide the final step by -7?

Q929. Solve for x: $-5x + 29 = -3$. What happens to the signs when you divide the final step by -5?

Q930. Solve for x: $-7x + 7 = 6$. What happens to the signs when you divide the final step by -7?

Q931. Solve for x: $-9x + 29 = 5$. What happens to the signs when you divide the final step by -9?

Q932. Solve for x: $-4x + 17 = 10$. What happens to the signs when you divide the final step by -4?

Q933. Solve for x: $-6x + 22 = -18$. What happens to the signs when you divide the final step by -6?

Q934. Solve for x: $-6x + 21 = -20$. What happens to the signs when you divide the final step by -6?

Q935. Solve for x: $-6x + 15 = 7$. What happens to the signs when you divide the final step by -6?

Q936. Solve for x: $-8x + 16 = 13$. What happens to the signs when you divide the final step by -8?

Q937. Solve for x: $-2x + 21 = 20$. What happens to the signs when you divide the final step by -2?

Q938. Solve for x: $-2x + 5 = 18$. What happens to the signs when you divide the final step by -2?

Q939. Solve for x: $-8x + 12 = 6$. What happens to the signs when you divide the final step by -8?

Q940. Solve for x: $-7x + 29 = -4$. What happens to the signs when you divide the final step by -7?

Q941. Solve for x: $-4x + 24 = -18$. What happens to the signs when you divide the final step by -4?

Q942. Solve for x: $-5x + 14 = 17$. What happens to the signs when you divide the final step by -5?

Q943. Solve for x: $-6x + 21 = -4$. What happens to the signs when you divide the final step by -6?

Q944. Solve for x: $-2x + 12 = -19$. What happens to the signs when you divide the final step by -2?

Q945. Solve for x: $-5x + 13 = -15$. What happens to the signs when you divide the final step by -5?

Q946. Solve for x: $-7x + 19 = -4$. What happens to the signs when you divide the final step by -7?

Q947. Solve for x: $-5x + 27 = 9$. What happens to the signs when you divide the final step by -5?

Q948. Solve for x: $-6x + 28 = -7$. What happens to the signs when you divide the final step by -6?

Q949. Solve for x: $-8x + 12 = -17$. What happens to the signs when you divide the final step by -8?

Q950. Solve for x: $-2x + 21 = -10$. What happens to the signs when you divide the final step by -2?

Q951. Solve for x: $-2x + 14 = -6$. What happens to the signs when you divide the final step by -2?

Q952. Solve for x: $-7x + 10 = 18$. What happens to the signs when you divide the final step by -7?

Q953. Solve for x: $-6x + 19 = 11$. What happens to the signs when you divide the final step by -6?

Q954. Solve for x: $-5x + 14 = -12$. What happens to the signs when you divide the final step by -5?

Q955. Solve for x: $-4x + 20 = 16$. What happens to the signs when you divide the final step by -4?

Q956. Solve for x: $-2x + 21 = 11$. What happens to the signs when you divide the final step by -2?

Q957. Solve for x: $-6x + 29 = -17$. What happens to the signs when you divide the final step by -6?

Q958. Solve for x: $-9x + 21 = 6$. What happens to the signs when you divide the final step by -9?

Q959. Solve for x: $-7x + 11 = -13$. What happens to the signs when you divide the final step by -7?

Q960. Solve for x: $-3x + 8 = 10$. What happens to the signs when you divide the final step by -3?

Q961. Solve for x: $-3x + 18 = 16$. What happens to the signs when you divide the final step by -3?

Q962. Solve for x: $-9x + 24 = 11$. What happens to the signs when you divide the final step by -9?

Q963. Solve for x: $-3x + 17 = -12$. What happens to the signs when you divide the final step by -3?

Q964. Solve for x: $-8x + 20 = -15$. What happens to the signs when you divide the final step by -8?

Q965. Solve for x: $-9x + 17 = -13$. What happens to

the signs when you divide the final step by -9?

Q966. Solve for x: $-6x + 28 = 20$. What happens to the signs when you divide the final step by -6?

Q967. Solve for x: $-7x + 26 = 20$. What happens to the signs when you divide the final step by -7?

Q968. Solve for x: $-3x + 14 = 8$. What happens to the signs when you divide the final step by -3?

Q969. Solve for x: $-9x + 12 = 8$. What happens to the signs when you divide the final step by -9?

Q970. Solve for x: $-2x + 16 = -19$. What happens to the signs when you divide the final step by -2?

Q971. Solve for x: $-3x + 5 = 2$. What happens to the signs when you divide the final step by -3?

Q972. Solve for x: $-2x + 29 = 3$. What happens to the signs when you divide the final step by -2?

Q973. Solve for x: $-9x + 16 = 19$. What happens to the signs when you divide the final step by -9?

Q974. Solve for x: $-3x + 26 = -14$. What happens to the signs when you divide the final step by -3?

Q975. Solve for x: $-8x + 26 = 5$. What happens to the signs when you divide the final step by -8?

Q976. Solve for x: $-6x + 30 = 20$. What happens to the signs when you divide the final step by -6?

Q977. Solve for x: $-9x + 19 = -14$. What happens to the signs when you divide the final step by -9?

Q978. Solve for x: $-9x + 28 = -6$. What happens to the signs when you divide the final step by -9?

Q979. Solve for x: $-9x + 28 = 6$. What happens to the signs when you divide the final step by -9?

Q980. Solve for x: $-4x + 27 = 7$. What happens to the signs when you divide the final step by -4?

Q981. Solve for x: $-2x + 12 = -3$. What happens to the signs when you divide the final step by -2?

Q982. Solve for x: $-9x + 12 = -9$. What happens to the signs when you divide the final step by -9?

Q983. Solve for x: $-4x + 9 = 10$. What happens to the signs when you divide the final step by -4?

Q984. Solve for x: $-3x + 26 = -6$. What happens to the signs when you divide the final step by -3?

Q985. Solve for x: $-8x + 14 = 3$. What happens to the signs when you divide the final step by -8?

Q986. Solve for x: $-7x + 29 = 4$. What happens to the signs when you divide the final step by -7?

Q987. Solve for x: $-5x + 23 = -15$. What happens to the signs when you divide the final step by -5?

Q988. Solve for x: $-4x + 6 = -16$. What happens to the signs when you divide the final step by -4?

Q989. Solve for x: $-9x + 5 = -1$. What happens to the signs when you divide the final step by -9?

Q990. Solve for x: $-9x + 11 = 15$. What happens to the signs when you divide the final step by -9?

Q991. Solve for x: $-2x + 30 = -8$. What happens to the signs when you divide the final step by -2?

Q992. Solve for x: $-7x + 8 = -16$. What happens to the signs when you divide the final step by -7?

Q993. Solve for x: $-8x + 11 = 8$. What happens to the signs when you divide the final step by -8?

Q994. Solve for x: $-2x + 21 = 7$. What happens to the signs when you divide the final step by -2?

Q995. Solve for x: $-8x + 15 = -10$. What happens to the signs when you divide the final step by -8?

Q996. Solve for x: $-3x + 6 = -19$. What happens to the signs when you divide the final step by -3?

Q997. Solve for x: $-5x + 30 = 6$. What happens to the signs when you divide the final step by -5?

Q998. Solve for x: $-7x + 9 = 16$. What happens to the signs when you divide the final step by -7?

Q999. Solve for x: $-2x + 23 = 18$. What happens to the signs when you divide the final step by -2?

Q1000. Solve for x: $-3x + 19 = 11$. What happens to the signs when you divide the final step by -3?

MODULE 2: WORD PROBLEM TRANSLATION AND EQUATION FORMATION (Q1001 – Q2000)

Q1001. The sum of two numbers is 103 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1002. The sum of two numbers is 41 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1003. The sum of two numbers is 38 and their difference is 12. Write the pair of linear equations and find both numbers.

Q1004. The sum of two numbers is 38 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1005. The sum of two numbers is 118 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1006. The sum of two numbers is 92 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1007. The sum of two numbers is 47 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1008. The sum of two numbers is 89 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1009. The sum of two numbers is 116 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1010. The sum of two numbers is 115 and their difference is 5. Write the pair of linear equations and find both numbers.

Q1011. The sum of two numbers is 63 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1012. The sum of two numbers is 38 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1013. The sum of two numbers is 95 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1014. The sum of two numbers is 94 and their difference is 6. Write the pair of linear equations and find both numbers.

Q1015. The sum of two numbers is 85 and their difference is 5. Write the pair of linear equations and find both numbers.

Q1016. The sum of two numbers is 46 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1017. The sum of two numbers is 64 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1018. The sum of two numbers is 65 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1019. The sum of two numbers is 81 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1020. The sum of two numbers is 52 and their difference is 18. Write the pair of linear equations and find both numbers.

Q1021. The sum of two numbers is 49 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1022. The sum of two numbers is 87 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1023. The sum of two numbers is 45 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1024. The sum of two numbers is 82 and their difference is 6. Write the pair of linear equations and find both numbers.

Q1025. The sum of two numbers is 88 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1026. The sum of two numbers is 62 and their difference is 18. Write the pair of linear equations and find both numbers.

Q1027. The sum of two numbers is 74 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1028. The sum of two numbers is 112 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1029. The sum of two numbers is 79 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1030. The sum of two numbers is 75 and their difference is 23. Write the pair of linear equations and find both numbers.

Q1031. The sum of two numbers is 120 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1032. The sum of two numbers is 74 and their difference is 10. Write the pair of linear equations and find both numbers.

Q1033. The sum of two numbers is 63 and their difference is 5. Write the pair of linear equations and find both numbers.

Q1034. The sum of two numbers is 90 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1035. The sum of two numbers is 49 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1036. The sum of two numbers is 51 and their difference is 23. Write the pair of linear equations and find both numbers.

Q1037. The sum of two numbers is 73 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1038. The sum of two numbers is 109 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1039. The sum of two numbers is 40 and their difference is 22. Write the pair of linear equations and find both numbers.

Q1040. The sum of two numbers is 120 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1041. The sum of two numbers is 72 and their difference is 22. Write the pair of linear equations and find both numbers.

Q1042. The sum of two numbers is 89 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1043. The sum of two numbers is 40 and their difference is 12. Write the pair of linear equations and find both numbers.

Q1044. The sum of two numbers is 106 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1045. The sum of two numbers is 120 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1046. The sum of two numbers is 48 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1047. The sum of two numbers is 83 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1048. The sum of two numbers is 32 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1049. The sum of two numbers is 120 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1050. The sum of two numbers is 106 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1051. The sum of two numbers is 71 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1052. The sum of two numbers is 80 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1053. The sum of two numbers is 95 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1054. The sum of two numbers is 94 and their difference is 12. Write the pair of linear equations and find both numbers.

Q1055. The sum of two numbers is 72 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1056. The sum of two numbers is 41 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1057. The sum of two numbers is 119 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1058. The sum of two numbers is 120 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1059. The sum of two numbers is 75 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1060. The sum of two numbers is 77 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1061. The sum of two numbers is 32 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1062. The sum of two numbers is 101 and their difference is 19. Write the pair of linear equations and find both numbers.

Q1063. The sum of two numbers is 108 and their difference is 12. Write the pair of linear equations and find both numbers.

Q1064. The sum of two numbers is 65 and their difference is 21. Write the pair of linear equations and find both numbers.

Q1065. The sum of two numbers is 32 and their difference is 10. Write the pair of linear equations and find both numbers.

Q1066. The sum of two numbers is 102 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1067. The sum of two numbers is 92 and their difference is 18. Write the pair of linear equations and find both numbers.

Q1068. The sum of two numbers is 41 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1069. The sum of two numbers is 40 and their difference is 6. Write the pair of linear equations and find both numbers.

Q1070. The sum of two numbers is 40 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1071. The sum of two numbers is 105 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1072. The sum of two numbers is 40 and their difference is 10. Write the pair of linear equations and find both numbers.

Q1073. The sum of two numbers is 74 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1074. The sum of two numbers is 120 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1075. The sum of two numbers is 46 and their difference is 4. Write the pair of linear equations and find both

numbers.

Q1076. The sum of two numbers is 42 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1077. The sum of two numbers is 50 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1078. The sum of two numbers is 89 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1079. The sum of two numbers is 82 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1080. The sum of two numbers is 102 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1081. The sum of two numbers is 47 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1082. The sum of two numbers is 71 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1083. The sum of two numbers is 48 and their difference is 10. Write the pair of linear equations and find both numbers.

Q1084. The sum of two numbers is 93 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1085. The sum of two numbers is 32 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1086. The sum of two numbers is 34 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1087. The sum of two numbers is 45 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1088. The sum of two numbers is 81 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1089. The sum of two numbers is 120 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1090. The sum of two numbers is 68 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1091. The sum of two numbers is 53 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1092. The sum of two numbers is 96 and their difference is 12. Write the pair of linear equations and find both numbers.

Q1093. The sum of two numbers is 94 and their difference is 16. Write the pair of linear equations and find both numbers.

Q1094. The sum of two numbers is 85 and their difference is 15. Write the pair of linear equations and find both numbers.

Q1095. The sum of two numbers is 113 and their difference is 15. Write the pair of linear equations and find both numbers.

Q1096. The sum of two numbers is 57 and their difference is 21. Write the pair of linear equations and find both numbers.

Q1097. The sum of two numbers is 101 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1098. The sum of two numbers is 45 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1099. The sum of two numbers is 113 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1100. The sum of two numbers is 60 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1101. The sum of two numbers is 48 and their difference is 18. Write the pair of linear equations and find both numbers.

Q1102. The sum of two numbers is 103 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1103. The sum of two numbers is 102 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1104. The sum of two numbers is 81 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1105. The sum of two numbers is 110 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1106. The sum of two numbers is 117 and their difference is 19. Write the pair of linear equations and find both numbers.

Q1107. The sum of two numbers is 43 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1108. The sum of two numbers is 52 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1109. The sum of two numbers is 74 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1110. The sum of two numbers is 59 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1111. The sum of two numbers is 61 and their difference is 15. Write the pair of linear equations and find both numbers.

Q1112. The sum of two numbers is 119 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1113. The sum of two numbers is 44 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1114. The sum of two numbers is 65 and their difference

is 19. Write the pair of linear equations and find both numbers.

Q1115. The sum of two numbers is 66 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1116. The sum of two numbers is 116 and their difference is 8. Write the pair of linear equations and find both numbers.

Q1117. The sum of two numbers is 73 and their difference is 5. Write the pair of linear equations and find both numbers.

Q1118. The sum of two numbers is 51 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1119. The sum of two numbers is 45 and their difference is 19. Write the pair of linear equations and find both numbers.

Q1120. The sum of two numbers is 109 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1121. The sum of two numbers is 33 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1122. The sum of two numbers is 75 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1123. The sum of two numbers is 83 and their difference is 23. Write the pair of linear equations and find both numbers.

Q1124. The sum of two numbers is 81 and their difference is 9. Write the pair of linear equations and find both numbers.

Q1125. The sum of two numbers is 98 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1126. The sum of two numbers is 46 and their difference is 18. Write the pair of linear equations and find both numbers.

Q1127. The sum of two numbers is 40 and their difference is 4. Write the pair of linear equations and find both numbers.

Q1128. The sum of two numbers is 113 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1129. The sum of two numbers is 36 and their difference is 10. Write the pair of linear equations and find both numbers.

Q1130. The sum of two numbers is 68 and their difference is 20. Write the pair of linear equations and find both numbers.

Q1131. The sum of two numbers is 97 and their difference is 15. Write the pair of linear equations and find both numbers.

Q1132. The sum of two numbers is 62 and their difference is 22. Write the pair of linear equations and find both numbers.

Q1133. The sum of two numbers is 74 and their difference is 26. Write the pair of linear equations and find both numbers.

Q1134. The sum of two numbers is 114 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1135. The sum of two numbers is 121 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1136. The sum of two numbers is 34 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1137. The sum of two numbers is 41 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1138. The sum of two numbers is 107 and their difference is 17. Write the pair of linear equations and find both numbers.

Q1139. The sum of two numbers is 53 and their difference is 15. Write the pair of linear equations and find both numbers.

Q1140. The sum of two numbers is 110 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1141. The sum of two numbers is 38 and their difference is 28. Write the pair of linear equations and find both numbers.

Q1142. The sum of two numbers is 52 and their difference is 24. Write the pair of linear equations and find both numbers.

Q1143. The sum of two numbers is 33 and their difference is 7. Write the pair of linear equations and find both numbers.

Q1144. The sum of two numbers is 49 and their difference is 5. Write the pair of linear equations and find both numbers.

Q1145. The sum of two numbers is 81 and their difference is 11. Write the pair of linear equations and find both numbers.

Q1146. The sum of two numbers is 97 and their difference is 27. Write the pair of linear equations and find both numbers.

Q1147. The sum of two numbers is 87 and their difference is 13. Write the pair of linear equations and find both numbers.

Q1148. The sum of two numbers is 79 and their difference is 25. Write the pair of linear equations and find both numbers.

Q1149. The sum of two numbers is 58 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1150. The sum of two numbers is 60 and their difference is 14. Write the pair of linear equations and find both numbers.

Q1151. Let father's age be x and son's age be y. 9 years ago, father was 3 times as old as his son. In 10 years, he will be 3 times as old as his son. Form both linear equations.

Q1152. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 8 years, he will

be 4 times as old as his son. Form both linear equations.

Q1153. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1154. Let father's age be x and son's age be y. 10 years ago, father was 3 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1155. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 6 years, he will be 3 times as old as his son. Form both linear equations.

Q1156. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1157. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1158. Let father's age be x and son's age be y. 7 years ago, father was 5 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1159. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1160. Let father's age be x and son's age be y. 8 years ago, father was 3 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1161. Let father's age be x and son's age be y. 7 years ago, father was 4 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1162. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 7 years, he will be 4 times as old as his son. Form both linear equations.

Q1163. Let father's age be x and son's age be y. 10 years ago, father was 6 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1164. Let father's age be x and son's age be y. 4 years ago, father was 7 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1165. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1166. Let father's age be x and son's age be y. 4 years ago, father was 7 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1167. Let father's age be x and son's age be y. 5 years ago, father was 3 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1168. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1169. Let father's age be x and son's age be y. 10 years ago, father was 7 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1170. Let father's age be x and son's age be y. 5 years ago, father was 3 times as old as his son. In 6 years, he will be 3 times as old as his son. Form both linear equations.

Q1171. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1172. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1173. Let father's age be x and son's age be y. 8 years ago, father was 3 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1174. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1175. Let father's age be x and son's age be y. 6 years ago, father was 7 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1176. Let father's age be x and son's age be y. 9 years ago, father was 5 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1177. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 5 years, he will be 2 times as old as his son. Form both linear equations.

Q1178. Let father's age be x and son's age be y. 3 years ago, father was 7 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1179. Let father's age be x and son's age be y. 8 years ago, father was 4 times as old as his son. In 10 years, he will be 3 times as old as his son. Form both linear equations.

Q1180. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1181. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 3 years, he will be 4 times as old as his son. Form both linear equations.

Q1182. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1183. Let father's age be x and son's age be y. 8 years ago, father was 4 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1184. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1185. Let father's age be x and son's age be y. 5 years ago, father was 6 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1186. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1187. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1188. Let father's age be x and son's age be y. 7 years ago, father was 4 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1189. Let father's age be x and son's age be y. 8 years ago, father was 3 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1190. Let father's age be x and son's age be y. 4 years ago, father was 7 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1191. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 6 years, he will be 2 times as old as his son. Form both linear equations.

Q1192. Let father's age be x and son's age be y. 8 years ago, father was 4 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1193. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1194. Let father's age be x and son's age be y. 10 years ago, father was 4 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1195. Let father's age be x and son's age be y. 10 years ago, father was 4 times as old as his son. In 8 years, he will be 4 times as old as his son. Form both linear equations.

Q1196. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1197. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1198. Let father's age be x and son's age be y. 5 years ago, father was 6 times as old as his son. In 7 years, he will be 4 times as old as his son. Form both linear equations.

Q1199. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 5 years, he will be 2 times as old as his son. Form both linear equations.

Q1200. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1201. Let father's age be x and son's age be y. 10 years ago, father was 7 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1202. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1203. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1204. Let father's age be x and son's age be y. 6 years ago, father was 7 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1205. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1206. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1207. Let father's age be x and son's age be y. 3 years ago, father was 3 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1208. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 5 years, he will be 4 times as old as his son. Form both linear equations.

Q1209. Let father's age be x and son's age be y. 9 years ago, father was 5 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1210. Let father's age be x and son's age be y. 9 years ago, father was 5 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1211. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1212. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 3 years, he will be 4 times as old as his son. Form both linear equations.

Q1213. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1214. Let father's age be x and son's age be y. 10 years ago, father was 5 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1215. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1216. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1217. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1218. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1219. Let father's age be x and son's age be y. 4 years ago, father was 5 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1220. Let father's age be x and son's age be y. 5 years ago, father was 4 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1221. Let father's age be x and son's age be y. 9 years ago, father was 6 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1222. Let father's age be x and son's age be y. 10 years ago, father was 7 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1223. Let father's age be x and son's age be y. 8 years ago, father was 5 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1224. Let father's age be x and son's age be y. 6 years ago, father was 3 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1225. Let father's age be x and son's age be y. 8 years ago, father was 5 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1226. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1227. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1228. Let father's age be x and son's age be y. 5 years ago, father was 5 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1229. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1230. Let father's age be x and son's age be y. 4 years ago, father was 7 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1231. Let father's age be x and son's age be y. 7 years ago, father was 4 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1232. Let father's age be x and son's age be y. 10 years ago, father was 5 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1233. Let father's age be x and son's age be y. 10 years ago, father was 5 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1234. Let father's age be x and son's age be y. 10 years ago, father was 7 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1235. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1236. Let father's age be x and son's age be y. 10 years ago, father was 7 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1237. Let father's age be x and son's age be y. 3 years ago, father was 7 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1238. Let father's age be x and son's age be y. 5 years ago, father was 6 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1239. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1240. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1241. Let father's age be x and son's age be y. 8 years ago, father was 5 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1242. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1243. Let father's age be x and son's age be y. 7 years ago, father was 6 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1244. Let father's age be x and son's age be y. 7 years ago, father was 6 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1245. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1246. Let father's age be x and son's age be y. 8 years ago, father was 4 times as old as his son. In 5 years, he will be 4 times as old as his son. Form both linear equations.

Q1247. Let father's age be x and son's age be y. 7 years ago, father was 5 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1248. Let father's age be x and son's age be y. 3 years ago, father was 6 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1249. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 8 years, he will be 4 times as old as his son. Form both linear equations.

Q1250. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1251. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1252. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 7 years, he will be 4 times as old as his son. Form both linear equations.

Q1253. Let father's age be x and son's age be y. 6 years ago, father was 3 times as old as his son. In 7 years, he will be 4 times as old as his son. Form both linear equations.

Q1254. Let father's age be x and son's age be y. 10 years ago, father was 6 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1255. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1256. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 5 years, he will be 4 times as old as his son. Form both linear equations.

Q1257. Let father's age be x and son's age be y. 8 years ago, father was 3 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1258. Let father's age be x and son's age be y. 3 years ago, father was 5 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1259. Let father's age be x and son's age be y. 5 years ago, father was 7 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1260. Let father's age be x and son's age be y. 3 years ago, father was 5 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1261. Let father's age be x and son's age be y. 10 years ago, father was 4 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1262. Let father's age be x and son's age be y. 9 years ago, father was 6 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1263. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 3 years, he will be 4 times as old as his son. Form both linear equations.

Q1264. Let father's age be x and son's age be y. 3 years ago, father was 7 times as old as his son. In 5 years, he will be 4 times as old as his son. Form both linear equations.

Q1265. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1266. Let father's age be x and son's age be y. 6 years ago, father was 6 times as old as his son. In 3 years, he will be 4 times as old as his son. Form both linear equations.

Q1267. Let father's age be x and son's age be y. 4 years ago, father was 4 times as old as his son. In 4 years, he will be 3 times as old as his son. Form both linear equations.

Q1268. Let father's age be x and son's age be y. 8 years ago, father was 7 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1269. Let father's age be x and son's age be y. 9 years ago, father was 6 times as old as his son. In 5 years, he will be 2 times as old as his son. Form both linear equations.

Q1270. Let father's age be x and son's age be y. 7 years ago, father was 6 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1271. Let father's age be x and son's age be y. 9 years ago, father was 6 times as old as his son. In 10 years, he will be 4 times as old as his son. Form both linear equations.

Q1272. Let father's age be x and son's age be y. 9 years ago, father was 5 times as old as his son. In 7 years, he will be 2 times as old as his son. Form both linear equations.

Q1273. Let father's age be x and son's age be y. 10 years ago, father was 3 times as old as his son. In 5 years, he will be 4 times as old as his son. Form both linear equations.

Q1274. Let father's age be x and son's age be y. 4 years ago, father was 7 times as old as his son. In 3 years, he will be 3 times as old as his son. Form both linear equations.

Q1275. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 10 years, he will be 3 times as old as his son. Form both linear equations.

Q1276. Let father's age be x and son's age be y. 3 years ago, father was 6 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1277. Let father's age be x and son's age be y. 10 years ago, father was 5 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1278. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 5 years, he will be 2 times as old as his son. Form both linear equations.

Q1279. Let father's age be x and son's age be y. 4 years ago, father was 4 times as old as his son. In 8 years, he will be 3 times as old as his son. Form both linear equations.

Q1280. Let father's age be x and son's age be y. 4 years ago, father was 5 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1281. Let father's age be x and son's age be y. 7 years ago, father was 7 times as old as his son. In 6 years, he will be 4 times as old as his son. Form both linear equations.

Q1282. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1283. Let father's age be x and son's age be y. 3 years ago, father was 3 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1284. Let father's age be x and son's age be y. 9 years ago, father was 6 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1285. Let father's age be x and son's age be y. 3 years ago, father was 3 times as old as his son. In 8 years, he will be 2 times as old as his son. Form both linear equations.

Q1286. Let father's age be x and son's age be y. 6 years ago, father was 5 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1287. Let father's age be x and son's age be y. 8 years ago, father was 6 times as old as his son. In 6 years, he will be 2 times as old as his son. Form both linear equations.

Q1288. Let father's age be x and son's age be y. 3 years ago, father was 5 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1289. Let father's age be x and son's age be y. 3 years ago, father was 4 times as old as his son. In 7 years, he will be 3 times as old as his son. Form both linear equations.

Q1290. Let father's age be x and son's age be y. 6 years ago, father was 3 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1291. Let father's age be x and son's age be y. 9 years ago, father was 7 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1292. Let father's age be x and son's age be y. 4 years ago, father was 3 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1293. Let father's age be x and son's age be y. 4 years ago, father was 6 times as old as his son. In 4 years, he will be 4 times as old as his son. Form both linear equations.

Q1294. Let father's age be x and son's age be y. 7 years ago, father was 3 times as old as his son. In 3 years, he will be 2 times as old as his son. Form both linear equations.

Q1295. Let father's age be x and son's age be y. 9 years ago, father was 4 times as old as his son. In 9 years, he will be 3 times as old as his son. Form both linear equations.

Q1296. Let father's age be x and son's age be y. 9 years ago, father was 5 times as old as his son. In 9 years, he will be 4 times as old as his son. Form both linear equations.

Q1297. Let father's age be x and son's age be y. 8 years ago, father was 3 times as old as his son. In 4 years, he will be 2 times as old as his son. Form both linear equations.

Q1298. Let father's age be x and son's age be y. 10 years ago, father was 4 times as old as his son. In 10 years, he will be 2 times as old as his son. Form both linear equations.

Q1299. Let father's age be x and son's age be y. 6 years ago, father was 4 times as old as his son. In 5 years, he will be 3 times as old as his son. Form both linear equations.

Q1300. Let father's age be x and son's age be y. 7 years ago, father was 4 times as old as his son. In 9 years, he will be 2 times as old as his son. Form both linear equations.

Q1301. A taxi charge consists of a fixed charge x and

charge per km y. A journey of 12 km costs Rs. 200, and 21 km costs Rs. 335. Form linear equations and solve for x and y.

Q1302. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 198, and 24 km costs Rs. 328. Form linear equations and solve for x and y.

Q1303. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 123, and 20 km costs Rs. 305. Form linear equations and solve for x and y.

Q1304. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 90, and 20 km costs Rs. 210. Form linear equations and solve for x and y.

Q1305. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 76, and 18 km costs Rs. 142. Form linear equations and solve for x and y.

Q1306. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 204, and 28 km costs Rs. 442. Form linear equations and solve for x and y.

Q1307. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 227, and 21 km costs Rs. 339. Form linear equations and solve for x and y.

Q1308. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 95, and 27 km costs Rs. 175. Form linear equations and solve for x and y.

Q1309. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 86, and 30 km costs Rs. 411. Form linear equations and solve for x and y.

Q1310. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 229, and 18 km costs Rs. 285. Form linear equations and solve for x and y.

Q1311. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 143, and 24 km costs Rs. 325. Form linear equations and solve for x and y.

Q1312. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 129, and 16 km costs Rs. 150. Form linear equations and solve for x and y.

Q1313. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 113, and 18 km costs Rs. 185. Form linear equations and solve for x and y.

Q1314. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 150, and 29 km costs Rs. 378. Form linear equations and solve for x and y.

Q1315. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 111, and 17 km costs Rs. 135. Form linear equations and solve for x and y.

Q1316. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 103, and 25 km costs Rs. 222. Form linear equations and solve for x and y.

Q1317. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 103, and 28 km costs Rs. 211. Form linear equations and solve for x and y.

Q1318. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 168, and 23 km costs Rs. 312. Form linear equations and solve for x and y.

Q1319. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 94, and 23 km costs Rs. 310. Form linear equations and solve for x and y.

Q1320. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 56, and 29 km costs Rs. 200. Form linear equations and solve for x and y.

Q1321. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 74, and 24 km costs Rs. 226. Form linear equations and solve for x and y.

Q1322. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 197, and 16 km costs Rs. 208. Form linear equations and solve for x and y.

Q1323. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 112, and 29 km costs Rs. 343. Form linear equations and solve for x and y.

Q1324. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 213, and 25 km costs Rs. 367. Form linear equations and solve for x and y.

Q1325. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 188, and 16 km costs Rs. 278. Form linear equations and solve for x and y.

Q1326. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 164, and 21 km costs Rs. 344. Form linear equations and solve for x and y.

Q1327. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 121, and 17 km costs Rs. 166. Form linear equations and solve for x and y.

Q1328. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 73, and 25 km costs Rs. 253. Form linear equations and solve for x and y.

Q1329. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 114, and 26 km costs Rs. 198. Form linear equations and solve for x and y.

Q1330. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 238, and 22 km costs Rs. 373. Form linear equations and solve for

x and y.

Q1331. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 107, and 20 km costs Rs. 247. Form linear equations and solve for x and y.

Q1332. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 97, and 28 km costs Rs. 241. Form linear equations and solve for x and y.

Q1333. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 101, and 20 km costs Rs. 296. Form linear equations and solve for x and y.

Q1334. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 120, and 21 km costs Rs. 228. Form linear equations and solve for x and y.

Q1335. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 139, and 16 km costs Rs. 166. Form linear equations and solve for x and y.

Q1336. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 81, and 23 km costs Rs. 201. Form linear equations and solve for x and y.

Q1337. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 74, and 21 km costs Rs. 149. Form linear equations and solve for x and y.

Q1338. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 132, and 22 km costs Rs. 240. Form linear equations and solve for x and y.

Q1339. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 122, and 28 km costs Rs. 234. Form linear equations and solve for x and y.

Q1340. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 70, and 30 km costs Rs. 196. Form linear equations and solve for x and y.

Q1341. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 157, and 28 km costs Rs. 457. Form linear equations and solve for x and y.

Q1342. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 156, and 18 km costs Rs. 211. Form linear equations and solve for x and y.

Q1343. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 60, and 30 km costs Rs. 252. Form linear equations and solve for x and y.

Q1344. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 140, and 17 km costs Rs. 206. Form linear equations and solve for x and y.

Q1345. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 95, and 21 km costs Rs. 239. Form linear equations and solve for x and y.

Q1346. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 143, and 30 km costs Rs. 352. Form linear equations and solve for x and y.

Q1347. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 49, and 20 km costs Rs. 139. Form linear equations and solve for x and y.

Q1348. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 103, and 28 km costs Rs. 265. Form linear equations and solve for x and y.

Q1349. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 96, and 23 km costs Rs. 168. Form linear equations and solve for x and y.

Q1350. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 143, and 18 km costs Rs. 293. Form linear equations and solve for x and y.

Q1351. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 155, and 24 km costs Rs. 276. Form linear equations and solve for x and y.

Q1352. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 124, and 25 km costs Rs. 190. Form linear equations and solve for x and y.

Q1353. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 73, and 18 km costs Rs. 216. Form linear equations and solve for x and y.

Q1354. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 105, and 24 km costs Rs. 196. Form linear equations and solve for x and y.

Q1355. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 62, and 24 km costs Rs. 206. Form linear equations and solve for x and y.

Q1356. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 70, and 26 km costs Rs. 301. Form linear equations and solve for x and y.

Q1357. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 97, and 21 km costs Rs. 223. Form linear equations and solve for x and y.

Q1358. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 148, and 28 km costs Rs. 292. Form linear equations and solve for x and y.

Q1359. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 123, and 28 km costs Rs. 251. Form linear equations and solve for x and y.

Q1360. A taxi charge consists of a fixed charge x and

charge per km y. A journey of 14 km costs Rs. 184, and 29 km costs Rs. 364. Form linear equations and solve for x and y.

Q1361. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 187, and 25 km costs Rs. 397. Form linear equations and solve for x and y.

Q1362. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 177, and 24 km costs Rs. 267. Form linear equations and solve for x and y.

Q1363. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 236, and 19 km costs Rs. 288. Form linear equations and solve for x and y.

Q1364. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 97, and 27 km costs Rs. 317. Form linear equations and solve for x and y.

Q1365. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 113, and 25 km costs Rs. 173. Form linear equations and solve for x and y.

Q1366. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 134, and 24 km costs Rs. 218. Form linear equations and solve for x and y.

Q1367. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 80, and 30 km costs Rs. 296. Form linear equations and solve for x and y.

Q1368. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 96, and 22 km costs Rs. 216. Form linear equations and solve for x and y.

Q1369. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 79, and 23 km costs Rs. 207. Form linear equations and solve for x and y.

Q1370. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 88, and 18 km costs Rs. 172. Form linear equations and solve for x and y.

Q1371. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 112, and 19 km costs Rs. 160. Form linear equations and solve for x and y.

Q1372. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 58, and 21 km costs Rs. 154. Form linear equations and solve for x and y.

Q1373. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 47, and 18 km costs Rs. 125. Form linear equations and solve for x and y.

Q1374. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 85, and 29 km costs Rs. 253. Form linear equations and solve for x and y.

Q1375. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 105, and 20 km costs Rs. 193. Form linear equations and solve for x and y.

Q1376. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 168, and 17 km costs Rs. 188. Form linear equations and solve for x and y.

Q1377. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 88, and 18 km costs Rs. 123. Form linear equations and solve for x and y.

Q1378. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 199, and 24 km costs Rs. 309. Form linear equations and solve for x and y.

Q1379. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 89, and 17 km costs Rs. 137. Form linear equations and solve for x and y.

Q1380. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 91, and 17 km costs Rs. 212. Form linear equations and solve for x and y.

Q1381. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 123, and 27 km costs Rs. 214. Form linear equations and solve for x and y.

Q1382. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 116, and 20 km costs Rs. 165. Form linear equations and solve for x and y.

Q1383. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 257, and 29 km costs Rs. 482. Form linear equations and solve for x and y.

Q1384. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 118, and 24 km costs Rs. 214. Form linear equations and solve for x and y.

Q1385. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 111, and 22 km costs Rs. 215. Form linear equations and solve for x and y.

Q1386. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 67, and 29 km costs Rs. 331. Form linear equations and solve for x and y.

Q1387. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 71, and 25 km costs Rs. 185. Form linear equations and solve for x and y.

Q1388. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 123, and 19 km costs Rs. 255. Form linear equations and solve for x and y.

Q1389. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 209, and 22 km costs Rs. 374. Form linear equations and solve for

x and y.

Q1390. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 133, and 21 km costs Rs. 315. Form linear equations and solve for x and y.

Q1391. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 84, and 25 km costs Rs. 210. Form linear equations and solve for x and y.

Q1392. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 234, and 24 km costs Rs. 384. Form linear equations and solve for x and y.

Q1393. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 106, and 30 km costs Rs. 406. Form linear equations and solve for x and y.

Q1394. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 185, and 20 km costs Rs. 350. Form linear equations and solve for x and y.

Q1395. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 125, and 28 km costs Rs. 209. Form linear equations and solve for x and y.

Q1396. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 75, and 21 km costs Rs. 187. Form linear equations and solve for x and y.

Q1397. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 178, and 19 km costs Rs. 218. Form linear equations and solve for x and y.

Q1398. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 106, and 20 km costs Rs. 155. Form linear equations and solve for x and y.

Q1399. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 224, and 19 km costs Rs. 299. Form linear equations and solve for x and y.

Q1400. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 201, and 27 km costs Rs. 370. Form linear equations and solve for x and y.

Q1401. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 79, and 24 km costs Rs. 241. Form linear equations and solve for x and y.

Q1402. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 80, and 25 km costs Rs. 194. Form linear equations and solve for x and y.

Q1403. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 91, and 16 km costs Rs. 126. Form linear equations and solve for x and y.

Q1404. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 172, and 19 km costs Rs. 244. Form linear equations and solve for x and y.

Q1405. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 82, and 20 km costs Rs. 160. Form linear equations and solve for x and y.

Q1406. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 204, and 21 km costs Rs. 308. Form linear equations and solve for x and y.

Q1407. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 142, and 27 km costs Rs. 370. Form linear equations and solve for x and y.

Q1408. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 127, and 29 km costs Rs. 358. Form linear equations and solve for x and y.

Q1409. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 123, and 22 km costs Rs. 319. Form linear equations and solve for x and y.

Q1410. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 94, and 30 km costs Rs. 469. Form linear equations and solve for x and y.

Q1411. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 69, and 18 km costs Rs. 124. Form linear equations and solve for x and y.

Q1412. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 240, and 17 km costs Rs. 268. Form linear equations and solve for x and y.

Q1413. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 111, and 17 km costs Rs. 135. Form linear equations and solve for x and y.

Q1414. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 167, and 24 km costs Rs. 392. Form linear equations and solve for x and y.

Q1415. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 88, and 28 km costs Rs. 184. Form linear equations and solve for x and y.

Q1416. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 167, and 16 km costs Rs. 175. Form linear equations and solve for x and y.

Q1417. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 70, and 20 km costs Rs. 174. Form linear equations and solve for x and y.

Q1418. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 103, and 20 km costs Rs. 180. Form linear equations and solve for x and y.

Q1419. A taxi charge consists of a fixed charge x and

charge per km y. A journey of 10 km costs Rs. 173, and 28 km costs Rs. 443. Form linear equations and solve for x and y.

Q1420. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 122, and 26 km costs Rs. 292. Form linear equations and solve for x and y.

Q1421. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 180, and 17 km costs Rs. 235. Form linear equations and solve for x and y.

Q1422. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 189, and 21 km costs Rs. 339. Form linear equations and solve for x and y.

Q1423. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 161, and 23 km costs Rs. 371. Form linear equations and solve for x and y.

Q1424. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 117, and 28 km costs Rs. 195. Form linear equations and solve for x and y.

Q1425. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 163, and 26 km costs Rs. 331. Form linear equations and solve for x and y.

Q1426. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 156, and 27 km costs Rs. 282. Form linear equations and solve for x and y.

Q1427. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 127, and 18 km costs Rs. 183. Form linear equations and solve for x and y.

Q1428. A taxi charge consists of a fixed charge x and charge per km y. A journey of 15 km costs Rs. 228, and 16 km costs Rs. 241. Form linear equations and solve for x and y.

Q1429. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 81, and 28 km costs Rs. 176. Form linear equations and solve for x and y.

Q1430. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 155, and 17 km costs Rs. 232. Form linear equations and solve for x and y.

Q1431. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 118, and 23 km costs Rs. 184. Form linear equations and solve for x and y.

Q1432. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 120, and 29 km costs Rs. 300. Form linear equations and solve for x and y.

Q1433. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 164, and 18 km costs Rs. 281. Form linear equations and solve for x and y.

Q1434. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 134, and 20 km costs Rs. 254. Form linear equations and solve for x and y.

Q1435. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 130, and 16 km costs Rs. 207. Form linear equations and solve for x and y.

Q1436. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 208, and 23 km costs Rs. 362. Form linear equations and solve for x and y.

Q1437. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 80, and 26 km costs Rs. 248. Form linear equations and solve for x and y.

Q1438. A taxi charge consists of a fixed charge x and charge per km y. A journey of 12 km costs Rs. 192, and 17 km costs Rs. 257. Form linear equations and solve for x and y.

Q1439. A taxi charge consists of a fixed charge x and charge per km y. A journey of 11 km costs Rs. 105, and 16 km costs Rs. 130. Form linear equations and solve for x and y.

Q1440. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 78, and 16 km costs Rs. 177. Form linear equations and solve for x and y.

Q1441. A taxi charge consists of a fixed charge x and charge per km y. A journey of 14 km costs Rs. 194, and 27 km costs Rs. 363. Form linear equations and solve for x and y.

Q1442. A taxi charge consists of a fixed charge x and charge per km y. A journey of 8 km costs Rs. 110, and 25 km costs Rs. 246. Form linear equations and solve for x and y.

Q1443. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 159, and 27 km costs Rs. 346. Form linear equations and solve for x and y.

Q1444. A taxi charge consists of a fixed charge x and charge per km y. A journey of 6 km costs Rs. 121, and 18 km costs Rs. 301. Form linear equations and solve for x and y.

Q1445. A taxi charge consists of a fixed charge x and charge per km y. A journey of 13 km costs Rs. 178, and 29 km costs Rs. 354. Form linear equations and solve for x and y.

Q1446. A taxi charge consists of a fixed charge x and charge per km y. A journey of 10 km costs Rs. 151, and 22 km costs Rs. 307. Form linear equations and solve for x and y.

Q1447. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 138, and 23 km costs Rs. 278. Form linear equations and solve for x and y.

Q1448. A taxi charge consists of a fixed charge x and charge per km y. A journey of 5 km costs Rs. 102, and 26 km costs Rs. 354. Form linear equations and solve for x

and y.

Q1449. A taxi charge consists of a fixed charge x and charge per km y. A journey of 7 km costs Rs. 98, and 29 km costs Rs. 252. Form linear equations and solve for x and y.

Q1450. A taxi charge consists of a fixed charge x and charge per km y. A journey of 9 km costs Rs. 133, and 23 km costs Rs. 287. Form linear equations and solve for x and y.

Q1451. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1452. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1453. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1454. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 more than the original number. Form equations with tens digit x and units digit y.

Q1455. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1456. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1457. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1458. The sum of the digits of a two-digit number is 15. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1459. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1460. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1461. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1462. The sum of the digits of a two-digit number is 17. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1463. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1464. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1465. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1466. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1467. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1468. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1469. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1470. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1471. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1472. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1473. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1474. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1475. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1476. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1477. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1478. The sum of the digits of a two-digit number is 12.

The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1479. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1480. The sum of the digits of a two-digit number is 15. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1481. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1482. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1483. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1484. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1485. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1486. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 54 more than the original number. Form equations with tens digit x and units digit y.

Q1487. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1488. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1489. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1490. The sum of the digits of a two-digit number is 15. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1491. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1492. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1493. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1494. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1495. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1496. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1497. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1498. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1499. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1500. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1501. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1502. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 more than the original number. Form equations with tens digit x and units digit y.

Q1503. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1504. The sum of the digits of a two-digit number is 15. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1505. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1506. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1507. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and

units digit y.

Q1508. The sum of the digits of a two-digit number is 15. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1509. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1510. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1511. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1512. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1513. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1514. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 36 less than the original number. Form equations with tens digit x and units digit y.

Q1515. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1516. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1517. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1518. The sum of the digits of a two-digit number is 4. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1519. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1520. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1521. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1522. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1523. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1524. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 less than the original number. Form equations with tens digit x and units digit y.

Q1525. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1526. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 45 less than the original number. Form equations with tens digit x and units digit y.

Q1527. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1528. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1529. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1530. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1531. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1532. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1533. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1534. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1535. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1536. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 less than the original number. Form equations with tens digit x and units digit y.

Q1537. The sum of the digits of a two-digit number is

10. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1538. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1539. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1540. The sum of the digits of a two-digit number is 17. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1541. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 54 more than the original number. Form equations with tens digit x and units digit y.

Q1542. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1543. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 45 more than the original number. Form equations with tens digit x and units digit y.

Q1544. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1545. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1546. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1547. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1548. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 63 more than the original number. Form equations with tens digit x and units digit y.

Q1549. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1550. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1551. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1552. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1553. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1554. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1555. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1556. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1557. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 63 more than the original number. Form equations with tens digit x and units digit y.

Q1558. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1559. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1560. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1561. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1562. The sum of the digits of a two-digit number is 8. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1563. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1564. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1565. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 less than the original number. Form equations with tens digit x and units digit y.

Q1566. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit

x and units digit y.

Q1567. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1568. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1569. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1570. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1571. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1572. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1573. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1574. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1575. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1576. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1577. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1578. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 36 more than the original number. Form equations with tens digit x and units digit y.

Q1579. The sum of the digits of a two-digit number is 14. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1580. The sum of the digits of a two-digit number is 17. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1581. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 45 less than the original number. Form equations with tens digit x and units digit y.

Q1582. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 63 less than the original number. Form equations with tens digit x and units digit y.

Q1583. The sum of the digits of a two-digit number is 12. The number obtained by reversing the digits is 0 more than the original number. Form equations with tens digit x and units digit y.

Q1584. The sum of the digits of a two-digit number is 10. The number obtained by reversing the digits is 54 more than the original number. Form equations with tens digit x and units digit y.

Q1585. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1586. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1587. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1588. The sum of the digits of a two-digit number is 6. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1589. The sum of the digits of a two-digit number is 13. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1590. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1591. The sum of the digits of a two-digit number is 16. The number obtained by reversing the digits is 18 more than the original number. Form equations with tens digit x and units digit y.

Q1592. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 63 more than the original number. Form equations with tens digit x and units digit y.

Q1593. The sum of the digits of a two-digit number is 4. The number obtained by reversing the digits is 18 less than the original number. Form equations with tens digit x and units digit y.

Q1594. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1595. The sum of the digits of a two-digit number is 9. The number obtained by reversing the digits is 63 less than the original number. Form equations with tens digit x and units digit y.

Q1596. The sum of the digits of a two-digit number is 15.

The number obtained by reversing the digits is 9 less than the original number. Form equations with tens digit x and units digit y.

Q1597. The sum of the digits of a two-digit number is 5. The number obtained by reversing the digits is 9 more than the original number. Form equations with tens digit x and units digit y.

Q1598. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 more than the original number. Form equations with tens digit x and units digit y.

Q1599. The sum of the digits of a two-digit number is 11. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1600. The sum of the digits of a two-digit number is 7. The number obtained by reversing the digits is 27 less than the original number. Form equations with tens digit x and units digit y.

Q1601. A fraction becomes $\frac{1}{3}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1602. A fraction becomes $\frac{1}{2}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1603. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1604. A fraction becomes $\frac{1}{2}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1605. A fraction becomes $\frac{3}{3}$ if 4 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1606. A fraction becomes $\frac{1}{4}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1607. A fraction becomes $\frac{1}{3}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1608. A fraction becomes $\frac{2}{4}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1609. A fraction becomes $\frac{1}{5}$ if 4 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1610. A fraction becomes $\frac{1}{5}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1611. A fraction becomes $\frac{1}{3}$ if 5 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1612. A fraction becomes $\frac{1}{5}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1613. A fraction becomes $\frac{2}{2}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1614. A fraction becomes $\frac{2}{4}$ if 4 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1615. A fraction becomes $\frac{3}{3}$ if 3 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1616. A fraction becomes $\frac{1}{3}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1617. A fraction becomes $\frac{2}{3}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1618. A fraction becomes $\frac{2}{2}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1619. A fraction becomes $\frac{2}{2}$ if 1 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1620. A fraction becomes $\frac{3}{2}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1621. A fraction becomes $\frac{1}{2}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1622. A fraction becomes $\frac{2}{3}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1623. A fraction becomes $\frac{3}{2}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1624. A fraction becomes $\frac{3}{5}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1625. A fraction becomes $\frac{1}{4}$ if 5 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1626. A fraction becomes $\frac{3}{2}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1627. A fraction becomes $\frac{3}{5}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1628. A fraction becomes $\frac{2}{4}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1629. A fraction becomes $\frac{1}{4}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1630. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1631. A fraction becomes $\frac{3}{3}$ if 4 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1632. A fraction becomes $\frac{3}{4}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1633. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator

x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1634. A fraction becomes $\frac{1}{4}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1635. A fraction becomes $\frac{2}{2}$ if 1 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1636. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1637. A fraction becomes $\frac{1}{4}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1638. A fraction becomes $\frac{2}{4}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1639. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1640. A fraction becomes $\frac{2}{5}$ if 5 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1641. A fraction becomes $\frac{1}{5}$ if 2 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1642. A fraction becomes $\frac{3}{4}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1643. A fraction becomes $\frac{1}{5}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1644. A fraction becomes $\frac{1}{3}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1645. A fraction becomes $\frac{2}{5}$ if 5 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1646. A fraction becomes $\frac{1}{5}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1647. A fraction becomes $\frac{1}{4}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1648. A fraction becomes $\frac{2}{3}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1649. A fraction becomes $\frac{3}{5}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1650. A fraction becomes $\frac{3}{2}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1651. A fraction becomes $\frac{2}{3}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1652. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1653. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1654. A fraction becomes $\frac{3}{3}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1655. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1656. A fraction becomes $\frac{2}{2}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1657. A fraction becomes $\frac{3}{4}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1658. A fraction becomes $\frac{3}{3}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1659. A fraction becomes $\frac{3}{4}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1660. A fraction becomes $\frac{2}{5}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1661. A fraction becomes $\frac{3}{5}$ if 4 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1662. A fraction becomes $\frac{2}{4}$ if 2 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1663. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1664. A fraction becomes $\frac{3}{5}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1665. A fraction becomes $\frac{1}{3}$ if 2 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1666. A fraction becomes $\frac{1}{3}$ if 4 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1667. A fraction becomes $\frac{3}{4}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1668. A fraction becomes $\frac{2}{5}$ if 4 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1669. A fraction becomes $\frac{1}{2}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1670. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1671. A fraction becomes $\frac{2}{3}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1672. A fraction becomes $\frac{2}{4}$ if 1 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1673. A fraction becomes $\frac{1}{2}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1674. A fraction becomes $\frac{2}{5}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1675. A fraction becomes $\frac{3}{3}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1676. A fraction becomes $\frac{3}{4}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1677. A fraction becomes $\frac{2}{3}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1678. A fraction becomes $\frac{2}{5}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1679. A fraction becomes $\frac{1}{4}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1680. A fraction becomes $\frac{2}{4}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1681. A fraction becomes $\frac{1}{4}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1682. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1683. A fraction becomes $\frac{1}{2}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1684. A fraction becomes $\frac{2}{2}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1685. A fraction becomes $\frac{2}{2}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1686. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1687. A fraction becomes $\frac{3}{4}$ if 5 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1688. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1689. A fraction becomes $\frac{2}{3}$ if 4 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1690. A fraction becomes $\frac{3}{4}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1691. A fraction becomes $\frac{1}{5}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1692. A fraction becomes $\frac{2}{2}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1693. A fraction becomes $\frac{3}{3}$ if 5 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1694. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1695. A fraction becomes $\frac{3}{4}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1696. A fraction becomes $\frac{1}{5}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1697. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1698. A fraction becomes $\frac{2}{5}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1699. A fraction becomes $\frac{1}{5}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1700. A fraction becomes $\frac{1}{3}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1701. A fraction becomes $\frac{1}{4}$ if 1 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1702. A fraction becomes $\frac{2}{2}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1703. A fraction becomes $\frac{3}{3}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1704. A fraction becomes $\frac{3}{4}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1705. A fraction becomes $\frac{3}{2}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1706. A fraction becomes $\frac{3}{3}$ if 1 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1707. A fraction becomes $\frac{3}{2}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1708. A fraction becomes $\frac{3}{5}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1709. A fraction becomes $\frac{3}{2}$ if 2 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1710. A fraction becomes $\frac{2}{3}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

braic equation in standard form $ax + by = c$.

Q1711. A fraction becomes $\frac{2}{2}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1712. A fraction becomes $\frac{1}{2}$ if 4 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1713. A fraction becomes $\frac{2}{3}$ if 1 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1714. A fraction becomes $\frac{1}{2}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1715. A fraction becomes $\frac{2}{2}$ if 3 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1716. A fraction becomes $\frac{3}{5}$ if 4 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1717. A fraction becomes $\frac{2}{5}$ if 2 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1718. A fraction becomes $\frac{3}{2}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1719. A fraction becomes $\frac{1}{2}$ if 4 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1720. A fraction becomes $\frac{3}{5}$ if 1 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1721. A fraction becomes $\frac{2}{3}$ if 3 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1722. A fraction becomes $\frac{3}{2}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1723. A fraction becomes $\frac{3}{3}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1724. A fraction becomes $\frac{3}{3}$ if 5 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1725. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1726. A fraction becomes $\frac{1}{3}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1727. A fraction becomes $\frac{2}{5}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1728. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1729. A fraction becomes $\frac{1}{3}$ if 1 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1730. A fraction becomes $\frac{2}{4}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1731. A fraction becomes $\frac{3}{2}$ if 4 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1732. A fraction becomes $\frac{1}{5}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1733. A fraction becomes $\frac{1}{3}$ if 4 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1734. A fraction becomes $\frac{2}{4}$ if 5 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1735. A fraction becomes $\frac{2}{5}$ if 3 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1736. A fraction becomes $\frac{2}{4}$ if 2 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1737. A fraction becomes $\frac{3}{2}$ if 3 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1738. A fraction becomes $\frac{1}{5}$ if 4 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1739. A fraction becomes $\frac{3}{5}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1740. A fraction becomes $\frac{3}{3}$ if 3 is added to numerator x and 1 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1741. A fraction becomes $\frac{1}{2}$ if 4 is added to numerator x and 4 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1742. A fraction becomes $\frac{1}{2}$ if 5 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1743. A fraction becomes $\frac{2}{5}$ if 3 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1744. A fraction becomes $\frac{3}{2}$ if 1 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1745. A fraction becomes $\frac{3}{5}$ if 2 is added to numerator x and 2 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1746. A fraction becomes $\frac{2}{3}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1747. A fraction becomes $\frac{2}{2}$ if 4 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1748. A fraction becomes $\frac{3}{5}$ if 1 is added to numerator x and 5 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1749. A fraction becomes $\frac{3}{2}$ if 3 is added to numerator

x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1750. A fraction becomes $\frac{1}{4}$ if 1 is added to numerator x and 3 is subtracted from denominator y. Write the algebraic equation in standard form $ax + by = c$.

Q1751. A boat goes 30 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1752. A boat goes 29 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1753. A boat goes 23 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1754. A boat goes 56 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1755. A boat goes 34 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1756. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1757. A boat goes 29 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1758. A boat goes 38 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1759. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1760. A boat goes 35 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1761. A boat goes 49 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1762. A boat goes 25 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1763. A boat goes 39 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1764. A boat goes 60 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1765. A boat goes 21 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1766. A boat goes 34 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1767. A boat goes 26 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1768. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1769. A boat goes 39 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1770. A boat goes 45 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1771. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1772. A boat goes 28 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1773. A boat goes 21 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1774. A boat goes 55 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1775. A boat goes 56 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1776. A boat goes 59 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1777. A boat goes 24 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1778. A boat goes 34 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1779. A boat goes 25 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1780. A boat goes 33 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1781. A boat goes 29 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1782. A boat goes 55 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1783. A boat goes 39 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1784. A boat goes 57 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1785. A boat goes 32 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1786. A boat goes 47 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1787. A boat goes 54 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1788. A boat goes 41 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1789. A boat goes 24 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1790. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1791. A boat goes 54 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1792. A boat goes 36 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1793. A boat goes 32 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1794. A boat goes 28 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1795. A boat goes 58 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1796. A boat goes 30 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1797. A boat goes 58 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1798. A boat goes 45 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1799. A boat goes 41 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1800. A boat goes 29 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1801. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1802. A boat goes 31 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1803. A boat goes 44 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1804. A boat goes 31 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1805. A boat goes 31 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1806. A boat goes 56 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1807. A boat goes 40 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1808. A boat goes 27 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1809. A boat goes 50 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1810. A boat goes 35 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1811. A boat goes 39 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1812. A boat goes 54 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1813. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1814. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1815. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1816. A boat goes 33 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1817. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1818. A boat goes 55 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1819. A boat goes 34 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1820. A boat goes 52 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1821. A boat goes 33 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1822. A boat goes 33 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1823. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1824. A boat goes 34 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1825. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1826. A boat goes 22 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1827. A boat goes 25 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1828. A boat goes 49 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1829. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1830. A boat goes 37 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1831. A boat goes 25 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1832. A boat goes 51 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1833. A boat goes 56 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1834. A boat goes 50 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1835. A boat goes 40 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1836. A boat goes 49 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1837. A boat goes 56 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1838. A boat goes 24 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1839. A boat goes 36 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1840. A boat goes 29 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1841. A boat goes 50 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1842. A boat goes 55 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1843. A boat goes 36 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1844. A boat goes 37 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1845. A boat goes 28 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1846. A boat goes 20 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1847. A boat goes 39 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1848. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1849. A boat goes 45 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1850. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1851. A boat goes 50 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1852. A boat goes 32 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1853. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1854. A boat goes 33 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1855. A boat goes 49 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1856. A boat goes 32 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1857. A boat goes 54 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1858. A boat goes 49 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1859. A boat goes 37 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1860. A boat goes 47 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1861. A boat goes 50 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1862. A boat goes 27 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1863. A boat goes 32 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1864. A boat goes 23 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1865. A boat goes 60 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1866. A boat goes 48 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1867. A boat goes 41 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1868. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1869. A boat goes 48 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1870. A boat goes 24 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1871. A boat goes 53 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1872. A boat goes 42 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1873. A boat goes 43 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1874. A boat goes 24 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1875. A boat goes 59 km upstream and returns downstream to the starting point. Let boat speed in still water be x km/h and stream speed be y km/h. Express time taken upstream T_1 and time taken downstream T_2 as rational expressions.

Q1876. 5 men and 4 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1877. 6 men and 3 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1878. 6 men and 5 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1879. 4 men and 5 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1880. 2 men and 7 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1881. 2 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1882. 3 men and 5 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1883. 5 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1884. 4 men and 6 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1885. 4 men and 4 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1886. 3 men and 4 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1887. 3 men and 3 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1888. 6 men and 5 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1889. 6 men and 3 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1890. 2 men and 7 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1891. 4 men and 7 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1892. 2 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1893. 5 men and 4 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1894. 3 men and 7 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1895. 2 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1896. 6 men and 5 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1897. 6 men and 5 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1898. 4 men and 3 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1899. 6 men and 5 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1900. 2 men and 4 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1901. 2 men and 7 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1902. 2 men and 8 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1903. 6 men and 8 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1904. 3 men and 6 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1905. 6 men and 4 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1906. 4 men and 8 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1907. 6 men and 8 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1908. 5 men and 7 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1909. 4 men and 7 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1910. 6 men and 7 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1911. 3 men and 5 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1912. 6 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1913. 4 men and 3 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1914. 5 men and 7 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1915. 3 men and 8 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1916. 5 men and 5 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1917. 5 men and 7 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1918. 6 men and 6 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1919. 5 men and 4 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1920. 2 men and 6 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1921. 6 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1922. 6 men and 4 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1923. 2 men and 6 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1924. 2 men and 4 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1925. 5 men and 7 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1926. 5 men and 3 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1927. 6 men and 8 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1928. 4 men and 6 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1929. 6 men and 3 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1930. 4 men and 7 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1931. 6 men and 5 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1932. 6 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1933. 5 men and 3 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1934. 3 men and 3 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1935. 4 men and 7 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1936. 5 men and 8 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1937. 6 men and 7 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1938. 6 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1939. 2 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1940. 2 men and 6 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1941. 4 men and 8 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1942. 6 men and 7 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1943. 3 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y

days alone, construct the daily work-rate equation.

Q1944. 3 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1945. 2 men and 5 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1946. 6 men and 8 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1947. 5 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1948. 4 men and 5 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1949. 3 men and 3 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1950. 2 men and 3 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1951. 3 men and 4 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1952. 6 men and 5 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1953. 2 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1954. 6 men and 4 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1955. 6 men and 8 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1956. 6 men and 8 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1957. 2 men and 5 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1958. 3 men and 6 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1959. 5 men and 4 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1960. 3 men and 4 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1961. 2 men and 4 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1962. 3 men and 5 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1963. 6 men and 7 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1964. 3 men and 3 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1965. 4 men and 3 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1966. 5 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1967. 2 men and 7 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1968. 4 men and 3 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1969. 4 men and 7 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1970. 5 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1971. 5 men and 7 women can finish a piece of work in 5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1972. 4 men and 3 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1973. 5 men and 5 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1974. 5 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1975. 2 men and 3 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1976. 5 men and 5 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1977. 5 men and 8 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1978. 6 men and 3 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1979. 6 men and 6 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1980. 2 men and 4 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1981. 5 men and 6 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1982. 6 men and 5 women can finish a piece of work in

5 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1983. 3 men and 8 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1984. 4 men and 6 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1985. 3 men and 3 women can finish a piece of work in 9 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1986. 6 men and 8 women can finish a piece of work in 10 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1987. 2 men and 4 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1988. 4 men and 6 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1989. 2 men and 8 women can finish a piece of work in 11 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1990. 4 men and 3 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1991. 4 men and 6 women can finish a piece of work in 7 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1992. 4 men and 3 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1993. 3 men and 6 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1994. 6 men and 4 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1995. 4 men and 6 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1996. 5 men and 3 women can finish a piece of work in 6 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1997. 3 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1998. 3 men and 8 women can finish a piece of work in 8 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q1999. 6 men and 3 women can finish a piece of work in 12 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

Q2000. 4 men and 8 women can finish a piece of work in 4 days. If 1 man takes x days alone and 1 woman takes y days alone, construct the daily work-rate equation.

MODULE 3: METHOD SELECTION STRATEGY (WHAT METHOD WHEN AND WHERE) (Q2001 – Q3000)

Q2001. Without solving, determine the nature of solutions for: (1) $6x + 3y = 9$, (2) $12x + 6y = 18$. Explain which method criterion you applied.

Q2002. Without solving, determine the nature of solutions for: (1) $5x + 3y = 6$, (2) $10x + 6y = 15$. Explain which method criterion you applied.

Q2003. Without solving, determine the nature of solutions for: (1) $6x + 1y = 6$, (2) $25x + 4y = 24$. Explain which method criterion you applied.

Q2004. Without solving, determine the nature of solutions for: (1) $4x + 5y = 10$, (2) $8x + 10y = 20$. Explain which method criterion you applied.

Q2005. Without solving, determine the nature of solutions for: (1) $2x + 4y = 7$, (2) $6x + 12y = 24$. Explain which method criterion you applied.

Q2006. Without solving, determine the nature of solutions for: (1) $1x + 4y = 8$, (2) $4x + 12y = 24$. Explain which method criterion you applied.

Q2007. Without solving, determine the nature of solutions for: (1) $6x + 4y = 10$, (2) $18x + 12y = 30$. Explain which method criterion you applied.

Q2008. Without solving, determine the nature of solutions for: (1) $6x+4y = 2$, (2) $12x+8y = 7$. Explain which method criterion you applied.

Q2009. Without solving, determine the nature of solutions for: (1) $3x + 5y = 10$, (2) $7x + 10y = 20$. Explain which method criterion you applied.

Q2010. Without solving, determine the nature of solutions for: (1) $3x + 3y = 7$, (2) $12x + 12y = 28$. Explain which method criterion you applied.

Q2011. Without solving, determine the nature of solutions for: (1) $2x+1y = 4$, (2) $6x+3y = 15$. Explain which method criterion you applied.

Q2012. Without solving, determine the nature of solutions for: (1) $6x + 2y = 8$, (2) $25x + 8y = 32$. Explain which method criterion you applied.

Q2013. Without solving, determine the nature of solutions for: (1) $6x + 3y = 5$, (2) $18x + 9y = 15$. Explain which method criterion you applied.

Q2014. Without solving, determine the nature of solutions for: (1) $2x+1y = 5$, (2) $8x+4y = 23$. Explain which method criterion you applied.

Q2015. Without solving, determine the nature of solutions for: (1) $1x+1y = 6$, (2) $5x+4y = 24$. Explain which method criterion you applied.

Q2016. Without solving, determine the nature of solutions for: (1) $2x+5y = 3$, (2) $6x+15y = 9$. Explain which method criterion you applied.

Q2017. Without solving, determine the nature of solutions for: (1) $1x + 5y = 8$, (2) $2x + 10y = 19$. Explain which method criterion you applied.

Q2018. Without solving, determine the nature of solutions for: (1) $3x + 5y = 7$, (2) $10x + 15y = 21$. Explain which method criterion you applied.

Q2019. Without solving, determine the nature of solutions for: (1) $3x+3y = 8$, (2) $9x+9y = 24$. Explain which method criterion you applied.

Q2020. Without solving, determine the nature of solutions for: (1) $1x + 5y = 6$, (2) $3x + 15y = 21$. Explain which method criterion you applied.

Q2021. Without solving, determine the nature of solutions for: (1) $2x+6y = 2$, (2) $9x+24y = 8$. Explain which method criterion you applied.

Q2022. Without solving, determine the nature of solutions for: (1) $1x+1y = 2$, (2) $2x+2y = 4$. Explain which method criterion you applied.

Q2023. Without solving, determine the nature of solutions for: (1) $4x+3y = 5$, (2) $8x+6y = 13$. Explain which method criterion you applied.

Q2024. Without solving, determine the nature of solutions for: (1) $6x + 6y = 7$, (2) $13x + 12y = 14$. Explain which method criterion you applied.

Q2025. Without solving, determine the nature of solutions for: (1) $6x + 1y = 5$, (2) $24x + 4y = 20$. Explain which method criterion you applied.

Q2026. Without solving, determine the nature of solutions for: (1) $4x+2y = 9$, (2) $8x+4y = 21$. Explain which method criterion you applied.

Q2027. Without solving, determine the nature of solutions for: (1) $3x+2y = 5$, (2) $7x+4y = 10$. Explain which method criterion you applied.

Q2028. Without solving, determine the nature of solutions for: (1) $5x + 4y = 10$, (2) $20x + 16y = 40$. Explain which method criterion you applied.

Q2029. Without solving, determine the nature of solutions for: (1) $2x+1y = 6$, (2) $4x+2y = 15$. Explain which method criterion you applied.

Q2030. Without solving, determine the nature of solutions for: (1) $6x + 2y = 6$, (2) $25x + 8y = 24$. Explain which method criterion you applied.

Q2031. Without solving, determine the nature of solutions for: (1) $1x+1y = 9$, (2) $4x+4y = 36$. Explain which method criterion you applied.

Q2032. Without solving, determine the nature of solutions for: (1) $1x+1y = 6$, (2) $3x+3y = 21$. Explain which method criterion you applied.

Q2033. Without solving, determine the nature of solutions for: (1) $4x + 1y = 2$, (2) $9x + 2y = 4$. Explain which method criterion you applied.

Q2034. Without solving, determine the nature of solutions for: (1) $5x + 5y = 5$, (2) $15x + 15y = 15$. Explain which method criterion you applied.

Q2035. Without solving, determine the nature of solutions for: (1) $6x + 4y = 9$, (2) $18x + 12y = 30$. Explain which method criterion you applied.

Q2036. Without solving, determine the nature of solutions for: (1) $2x + 4y = 6$, (2) $9x + 16y = 24$. Explain which method criterion you applied.

Q2037. Without solving, determine the nature of solutions for: (1) $5x + 2y = 5$, (2) $20x + 8y = 20$. Explain which method criterion you applied.

Q2038. Without solving, determine the nature of solutions for: (1) $2x + 1y = 4$, (2) $4x + 2y = 11$. Explain which method criterion you applied.

Q2039. Without solving, determine the nature of solutions for: (1) $5x + 3y = 7$, (2) $21x + 12y = 28$. Explain which method criterion you applied.

Q2040. Without solving, determine the nature of solutions for: (1) $6x + 1y = 2$, (2) $12x + 2y = 4$. Explain which method criterion you applied.

Q2041. Without solving, determine the nature of solutions for: (1) $3x + 6y = 2$, (2) $9x + 18y = 9$. Explain which method criterion you applied.

Q2042. Without solving, determine the nature of solutions for: (1) $6x + 2y = 6$, (2) $25x + 8y = 24$. Explain which method criterion you applied.

Q2043. Without solving, determine the nature of solutions for: (1) $5x + 2y = 9$, (2) $20x + 8y = 36$. Explain which method criterion you applied.

Q2044. Without solving, determine the nature of solutions for: (1) $4x + 2y = 2$, (2) $12x + 6y = 9$. Explain which method criterion you applied.

Q2045. Without solving, determine the nature of solutions for: (1) $6x + 4y = 6$, (2) $19x + 12y = 18$. Explain which method criterion you applied.

Q2046. Without solving, determine the nature of solutions for: (1) $1x + 3y = 9$, (2) $2x + 6y = 18$. Explain which method criterion you applied.

Q2047. Without solving, determine the nature of solutions for: (1) $4x + 1y = 8$, (2) $8x + 2y = 19$. Explain which method criterion you applied.

Q2048. Without solving, determine the nature of solutions for: (1) $2x + 1y = 7$, (2) $7x + 3y = 21$. Explain which method criterion you applied.

Q2049. Without solving, determine the nature of solutions for: (1) $3x + 4y = 7$, (2) $9x + 12y = 21$. Explain which method criterion you applied.

Q2050. Without solving, determine the nature of solutions for: (1) $2x + 5y = 10$, (2) $4x + 10y = 23$. Explain which method criterion you applied.

Q2051. Without solving, determine the nature of solutions for: (1) $5x + 2y = 5$, (2) $21x + 8y = 20$. Explain which method criterion you applied.

Q2052. Without solving, determine the nature of solutions for: (1) $4x + 6y = 4$, (2) $12x + 18y = 12$. Explain which method criterion you applied.

Q2053. Without solving, determine the nature of solutions for: (1) $1x + 6y = 7$, (2) $4x + 24y = 31$. Explain which method criterion you applied.

Q2054. Without solving, determine the nature of solutions for: (1) $3x + 1y = 8$, (2) $13x + 4y = 32$. Explain which method criterion you applied.

Q2055. Without solving, determine the nature of solutions for: (1) $3x + 3y = 4$, (2) $12x + 12y = 16$. Explain which method criterion you applied.

Q2056. Without solving, determine the nature of solutions for: (1) $1x + 2y = 9$, (2) $2x + 4y = 21$. Explain which method criterion you applied.

Q2057. Without solving, determine the nature of solutions for: (1) $3x + 1y = 10$, (2) $10x + 3y = 30$. Explain which method criterion you applied.

Q2058. Without solving, determine the nature of solutions for: (1) $1x + 1y = 6$, (2) $2x + 2y = 12$. Explain which method criterion you applied.

Q2059. Without solving, determine the nature of solutions for: (1) $3x + 4y = 3$, (2) $6x + 8y = 9$. Explain which method criterion you applied.

Q2060. Without solving, determine the nature of solutions for: (1) $3x + 2y = 2$, (2) $7x + 4y = 4$. Explain which method criterion you applied.

Q2061. Without solving, determine the nature of solutions for: (1) $5x + 3y = 3$, (2) $20x + 12y = 12$. Explain which method criterion you applied.

Q2062. Without solving, determine the nature of solutions for: (1) $5x + 1y = 10$, (2) $15x + 3y = 33$. Explain which method criterion you applied.

Q2063. Without solving, determine the nature of solutions for: (1) $4x + 6y = 2$, (2) $13x + 18y = 6$. Explain which method criterion you applied.

Q2064. Without solving, determine the nature of solutions for: (1) $1x + 4y = 5$, (2) $4x + 16y = 20$. Explain which method criterion you applied.

Q2065. Without solving, determine the nature of solutions for: (1) $6x + 2y = 10$, (2) $18x + 6y = 33$. Explain which method criterion you applied.

Q2066. Without solving, determine the nature of solutions for: (1) $5x + 3y = 8$, (2) $21x + 12y = 32$. Explain which method criterion you applied.

Q2067. Without solving, determine the nature of solutions for: (1) $4x + 4y = 8$, (2) $16x + 16y = 32$. Explain which method criterion you applied.

Q2068. Without solving, determine the nature of solutions for: (1) $4x + 3y = 9$, (2) $12x + 9y = 30$. Explain which method criterion you applied.

Q2069. Without solving, determine the nature of solutions for: (1) $6x + 3y = 8$, (2) $13x + 6y = 16$. Explain which method criterion you applied.

Q2070. Without solving, determine the nature of solutions for: (1) $5x + 1y = 4$, (2) $10x + 2y = 8$. Explain which method criterion you applied.

Q2071. Without solving, determine the nature of solutions for: (1) $3x + 5y = 2$, (2) $6x + 10y = 7$. Explain which method criterion you applied.

Q2072. Without solving, determine the nature of solutions for: (1) $3x + 3y = 4$, (2) $10x + 9y = 12$. Explain which method criterion you applied.

Q2073. Without solving, determine the nature of solutions for: (1) $5x + 6y = 8$, (2) $20x + 24y = 32$. Explain which method criterion you applied.

Q2074. Without solving, determine the nature of solutions for: (1) $1x + 2y = 6$, (2) $4x + 8y = 27$. Explain which method criterion you applied.

Q2075. Without solving, determine the nature of solutions for: (1) $1x + 1y = 3$, (2) $3x + 2y = 6$. Explain which

method criterion you applied.

Q2076. Without solving, determine the nature of solutions for: (1) $5x+2y=3$, (2) $15x+6y=9$. Explain which method criterion you applied.

Q2077. Without solving, determine the nature of solutions for: (1) $3x+3y=10$, (2) $6x+6y=23$. Explain which method criterion you applied.

Q2078. Without solving, determine the nature of solutions for: (1) $6x+5y=7$, (2) $25x+20y=28$. Explain which method criterion you applied.

Q2079. Without solving, determine the nature of solutions for: (1) $2x+2y=9$, (2) $8x+8y=36$. Explain which method criterion you applied.

Q2080. Without solving, determine the nature of solutions for: (1) $6x+1y=9$, (2) $12x+2y=21$. Explain which method criterion you applied.

Q2081. Without solving, determine the nature of solutions for: (1) $3x+5y=8$, (2) $10x+15y=24$. Explain which method criterion you applied.

Q2082. Without solving, determine the nature of solutions for: (1) $2x+6y=6$, (2) $4x+12y=12$. Explain which method criterion you applied.

Q2083. Without solving, determine the nature of solutions for: (1) $1x+1y=6$, (2) $4x+4y=27$. Explain which method criterion you applied.

Q2084. Without solving, determine the nature of solutions for: (1) $3x+2y=4$, (2) $10x+6y=12$. Explain which method criterion you applied.

Q2085. Without solving, determine the nature of solutions for: (1) $4x+1y=4$, (2) $12x+3y=12$. Explain which method criterion you applied.

Q2086. Without solving, determine the nature of solutions for: (1) $4x+3y=7$, (2) $16x+12y=31$. Explain which method criterion you applied.

Q2087. Without solving, determine the nature of solutions for: (1) $4x+5y=4$, (2) $17x+20y=16$. Explain which method criterion you applied.

Q2088. Without solving, determine the nature of solutions for: (1) $4x+4y=10$, (2) $16x+16y=40$. Explain which method criterion you applied.

Q2089. Without solving, determine the nature of solutions for: (1) $2x+5y=7$, (2) $4x+10y=17$. Explain which method criterion you applied.

Q2090. Without solving, determine the nature of solutions for: (1) $4x+6y=2$, (2) $13x+18y=6$. Explain which method criterion you applied.

Q2091. Without solving, determine the nature of solutions for: (1) $3x+6y=4$, (2) $6x+12y=8$. Explain which method criterion you applied.

Q2092. Without solving, determine the nature of solutions for: (1) $4x+4y=5$, (2) $12x+12y=18$. Explain which method criterion you applied.

Q2093. Without solving, determine the nature of solutions for: (1) $2x+4y=6$, (2) $9x+16y=24$. Explain which method criterion you applied.

Q2094. Without solving, determine the nature of solutions for: (1) $1x+6y=8$, (2) $2x+12y=16$. Explain which method criterion you applied.

Q2095. Without solving, determine the nature of solutions for: (1) $1x+6y=7$, (2) $3x+18y=24$. Explain which method criterion you applied.

Q2096. Without solving, determine the nature of solutions for: (1) $5x+1y=4$, (2) $11x+2y=8$. Explain which method criterion you applied.

Q2097. Without solving, determine the nature of solutions for: (1) $3x+3y=6$, (2) $6x+6y=12$. Explain which method criterion you applied.

Q2098. Without solving, determine the nature of solutions for: (1) $2x+5y=2$, (2) $4x+10y=7$. Explain which method criterion you applied.

Q2099. Without solving, determine the nature of solutions for: (1) $4x+1y=3$, (2) $13x+3y=9$. Explain which method criterion you applied.

Q2100. Without solving, determine the nature of solutions for: (1) $3x+4y=7$, (2) $12x+16y=28$. Explain which method criterion you applied.

Q2101. Without solving, determine the nature of solutions for: (1) $4x+2y=9$, (2) $12x+6y=30$. Explain which method criterion you applied.

Q2102. Without solving, determine the nature of solutions for: (1) $6x+4y=8$, (2) $13x+8y=16$. Explain which method criterion you applied.

Q2103. Without solving, determine the nature of solutions for: (1) $5x+2y=8$, (2) $10x+4y=16$. Explain which method criterion you applied.

Q2104. Without solving, determine the nature of solutions for: (1) $4x+2y=8$, (2) $16x+8y=35$. Explain which method criterion you applied.

Q2105. Without solving, determine the nature of solutions for: (1) $5x+6y=2$, (2) $11x+12y=4$. Explain which method criterion you applied.

Q2106. Without solving, determine the nature of solutions for: (1) $2x+6y=10$, (2) $6x+18y=30$. Explain which method criterion you applied.

Q2107. Without solving, determine the nature of solutions for: (1) $2x+6y=6$, (2) $8x+24y=27$. Explain which method criterion you applied.

Q2108. Without solving, determine the nature of solutions for: (1) $4x+4y=9$, (2) $17x+16y=36$. Explain which method criterion you applied.

Q2109. Without solving, determine the nature of solutions for: (1) $1x+1y=10$, (2) $4x+4y=40$. Explain which method criterion you applied.

Q2110. Without solving, determine the nature of solutions for: (1) $2x+2y=3$, (2) $8x+8y=15$. Explain which method criterion you applied.

Q2111. Without solving, determine the nature of solutions for: (1) $1x+4y=8$, (2) $3x+8y=16$. Explain which method criterion you applied.

Q2112. Without solving, determine the nature of solutions for: (1) $1x+4y=5$, (2) $3x+12y=15$. Explain which method criterion you applied.

Q2113. Without solving, determine the nature of solutions for: (1) $1x+4y=7$, (2) $4x+16y=31$. Explain which method criterion you applied.

Q2114. Without solving, determine the nature of solu-

tions for: (1) $6x + 1y = 4$, (2) $25x + 4y = 16$. Explain which method criterion you applied.

Q2115. Without solving, determine the nature of solutions for: (1) $1x + 5y = 6$, (2) $2x + 10y = 12$. Explain which method criterion you applied.

Q2116. Without solving, determine the nature of solutions for: (1) $2x+1y = 6$, (2) $4x+2y = 15$. Explain which method criterion you applied.

Q2117. Without solving, determine the nature of solutions for: (1) $1x+6y = 2$, (2) $5x+24y = 8$. Explain which method criterion you applied.

Q2118. Without solving, determine the nature of solutions for: (1) $5x + 3y = 6$, (2) $20x + 12y = 24$. Explain which method criterion you applied.

Q2119. Without solving, determine the nature of solutions for: (1) $5x + 6y = 7$, (2) $10x + 12y = 17$. Explain which method criterion you applied.

Q2120. Without solving, determine the nature of solutions for: (1) $5x + 5y = 3$, (2) $16x + 15y = 9$. Explain which method criterion you applied.

Q2121. Without solving, determine the nature of solutions for: (1) $6x + 4y = 4$, (2) $24x + 16y = 16$. Explain which method criterion you applied.

Q2122. Without solving, determine the nature of solutions for: (1) $4x + 4y = 10$, (2) $16x + 16y = 43$. Explain which method criterion you applied.

Q2123. Without solving, determine the nature of solutions for: (1) $6x + 5y = 7$, (2) $19x + 15y = 21$. Explain which method criterion you applied.

Q2124. Without solving, determine the nature of solutions for: (1) $3x + 5y = 10$, (2) $9x + 15y = 30$. Explain which method criterion you applied.

Q2125. Without solving, determine the nature of solutions for: (1) $4x + 1y = 7$, (2) $12x + 3y = 24$. Explain which method criterion you applied.

Q2126. Without solving, determine the nature of solutions for: (1) $1x+2y = 5$, (2) $3x+4y = 10$. Explain which method criterion you applied.

Q2127. Without solving, determine the nature of solutions for: (1) $4x + 6y = 2$, (2) $16x + 24y = 8$. Explain which method criterion you applied.

Q2128. Without solving, determine the nature of solutions for: (1) $3x+1y = 7$, (2) $9x+3y = 24$. Explain which method criterion you applied.

Q2129. Without solving, determine the nature of solutions for: (1) $2x + 4y = 10$, (2) $7x + 12y = 30$. Explain which method criterion you applied.

Q2130. Without solving, determine the nature of solutions for: (1) $2x + 4y = 4$, (2) $6x + 12y = 12$. Explain which method criterion you applied.

Q2131. Without solving, determine the nature of solutions for: (1) $2x+4y = 8$, (2) $4x+8y = 19$. Explain which method criterion you applied.

Q2132. Without solving, determine the nature of solutions for: (1) $5x+2y = 4$, (2) $11x+4y = 8$. Explain which method criterion you applied.

Q2133. Without solving, determine the nature of solutions for: (1) $3x + 5y = 8$, (2) $6x + 10y = 16$. Explain which method criterion you applied.

Q2134. Without solving, determine the nature of solutions for: (1) $1x + 4y = 8$, (2) $4x + 16y = 35$. Explain which method criterion you applied.

Q2135. Without solving, determine the nature of solutions for: (1) $5x + 2y = 10$, (2) $21x + 8y = 40$. Explain which method criterion you applied.

Q2136. Without solving, determine the nature of solutions for: (1) $3x + 4y = 4$, (2) $12x + 16y = 16$. Explain which method criterion you applied.

Q2137. Without solving, determine the nature of solutions for: (1) $2x + 3y = 3$, (2) $8x + 12y = 15$. Explain which method criterion you applied.

Q2138. Without solving, determine the nature of solutions for: (1) $2x + 5y = 8$, (2) $9x + 20y = 32$. Explain which method criterion you applied.

Q2139. Without solving, determine the nature of solutions for: (1) $4x+1y = 7$, (2) $8x+2y = 14$. Explain which method criterion you applied.

Q2140. Without solving, determine the nature of solutions for: (1) $6x + 6y = 5$, (2) $24x + 24y = 23$. Explain which method criterion you applied.

Q2141. Without solving, determine the nature of solutions for: (1) $2x+5y = 4$, (2) $5x+10y = 8$. Explain which method criterion you applied.

Q2142. Without solving, determine the nature of solutions for: (1) $6x + 6y = 8$, (2) $24x + 24y = 32$. Explain which method criterion you applied.

Q2143. Without solving, determine the nature of solutions for: (1) $3x+6y = 3$, (2) $6x+12y = 9$. Explain which method criterion you applied.

Q2144. Without solving, determine the nature of solutions for: (1) $3x + 4y = 5$, (2) $10x + 12y = 15$. Explain which method criterion you applied.

Q2145. Without solving, determine the nature of solutions for: (1) $4x+3y = 5$, (2) $8x+6y = 10$. Explain which method criterion you applied.

Q2146. Without solving, determine the nature of solutions for: (1) $1x + 3y = 3$, (2) $2x + 6y = 9$. Explain which method criterion you applied.

Q2147. Without solving, determine the nature of solutions for: (1) $6x + 1y = 5$, (2) $19x + 3y = 15$. Explain which method criterion you applied.

Q2148. Without solving, determine the nature of solutions for: (1) $5x + 6y = 5$, (2) $20x + 24y = 20$. Explain which method criterion you applied.

Q2149. Without solving, determine the nature of solutions for: (1) $4x + 6y = 8$, (2) $8x + 12y = 19$. Explain which method criterion you applied.

Q2150. Without solving, determine the nature of solutions for: (1) $5x+3y = 3$, (2) $16x+9y = 9$. Explain which method criterion you applied.

Q2151. Without solving, determine the nature of solutions for: (1) $6x + 2y = 10$, (2) $12x + 4y = 20$. Explain which method criterion you applied.

Q2152. Without solving, determine the nature of solutions for: (1) $6x + 6y = 5$, (2) $18x + 18y = 18$. Explain which method criterion you applied.

Q2153. Without solving, determine the nature of solutions for: (1) $3x+1y=7$, (2) $7x+2y=14$. Explain which method criterion you applied.

Q2154. Without solving, determine the nature of solutions for: (1) $3x+3y=7$, (2) $12x+12y=28$. Explain which method criterion you applied.

Q2155. Without solving, determine the nature of solutions for: (1) $1x+1y=3$, (2) $4x+4y=15$. Explain which method criterion you applied.

Q2156. Without solving, determine the nature of solutions for: (1) $2x+4y=2$, (2) $7x+12y=6$. Explain which method criterion you applied.

Q2157. Without solving, determine the nature of solutions for: (1) $5x+2y=7$, (2) $10x+4y=14$. Explain which method criterion you applied.

Q2158. Without solving, determine the nature of solutions for: (1) $2x+1y=6$, (2) $4x+2y=15$. Explain which method criterion you applied.

Q2159. Without solving, determine the nature of solutions for: (1) $6x+5y=10$, (2) $25x+20y=40$. Explain which method criterion you applied.

Q2160. Without solving, determine the nature of solutions for: (1) $3x+5y=3$, (2) $9x+15y=9$. Explain which method criterion you applied.

Q2161. Without solving, determine the nature of solutions for: (1) $6x+5y=10$, (2) $18x+15y=33$. Explain which method criterion you applied.

Q2162. Without solving, determine the nature of solutions for: (1) $4x+2y=5$, (2) $17x+8y=20$. Explain which method criterion you applied.

Q2163. Without solving, determine the nature of solutions for: (1) $6x+4y=4$, (2) $24x+16y=16$. Explain which method criterion you applied.

Q2164. Without solving, determine the nature of solutions for: (1) $6x+1y=8$, (2) $24x+4y=35$. Explain which method criterion you applied.

Q2165. Without solving, determine the nature of solutions for: (1) $1x+3y=3$, (2) $4x+9y=9$. Explain which method criterion you applied.

Q2166. Without solving, determine the nature of solutions for: (1) $2x+3y=5$, (2) $8x+12y=20$. Explain which method criterion you applied.

Q2167. Without solving, determine the nature of solutions for: (1) $6x+6y=5$, (2) $12x+12y=13$. Explain which method criterion you applied.

Q2168. Without solving, determine the nature of solutions for: (1) $6x+3y=3$, (2) $13x+6y=6$. Explain which method criterion you applied.

Q2169. Without solving, determine the nature of solutions for: (1) $1x+3y=2$, (2) $3x+9y=6$. Explain which method criterion you applied.

Q2170. Without solving, determine the nature of solutions for: (1) $2x+5y=8$, (2) $8x+20y=35$. Explain which method criterion you applied.

Q2171. Without solving, determine the nature of solutions for: (1) $4x+5y=7$, (2) $17x+20y=28$. Explain which method criterion you applied.

Q2172. Without solving, determine the nature of solutions for: (1) $2x+3y=7$, (2) $4x+6y=14$. Explain which method criterion you applied.

Q2173. Without solving, determine the nature of solutions for: (1) $3x+3y=4$, (2) $12x+12y=19$. Explain which method criterion you applied.

Q2174. Without solving, determine the nature of solutions for: (1) $4x+3y=2$, (2) $9x+6y=4$. Explain which method criterion you applied.

Q2175. Without solving, determine the nature of solutions for: (1) $4x+4y=8$, (2) $12x+12y=24$. Explain which method criterion you applied.

Q2176. Without solving, determine the nature of solutions for: (1) $1x+4y=3$, (2) $4x+16y=15$. Explain which method criterion you applied.

Q2177. Without solving, determine the nature of solutions for: (1) $4x+3y=5$, (2) $9x+6y=10$. Explain which method criterion you applied.

Q2178. Without solving, determine the nature of solutions for: (1) $1x+5y=9$, (2) $2x+10y=18$. Explain which method criterion you applied.

Q2179. Without solving, determine the nature of solutions for: (1) $1x+1y=9$, (2) $4x+4y=39$. Explain which method criterion you applied.

Q2180. Without solving, determine the nature of solutions for: (1) $5x+4y=8$, (2) $16x+12y=24$. Explain which method criterion you applied.

Q2181. Without solving, determine the nature of solutions for: (1) $3x+1y=4$, (2) $9x+3y=12$. Explain which method criterion you applied.

Q2182. Without solving, determine the nature of solutions for: (1) $3x+3y=5$, (2) $6x+6y=13$. Explain which method criterion you applied.

Q2183. Without solving, determine the nature of solutions for: (1) $4x+3y=8$, (2) $9x+6y=16$. Explain which method criterion you applied.

Q2184. Without solving, determine the nature of solutions for: (1) $5x+1y=2$, (2) $20x+4y=8$. Explain which method criterion you applied.

Q2185. Without solving, determine the nature of solutions for: (1) $3x+3y=5$, (2) $12x+12y=23$. Explain which method criterion you applied.

Q2186. Without solving, determine the nature of solutions for: (1) $6x+1y=10$, (2) $19x+3y=30$. Explain which method criterion you applied.

Q2187. Without solving, determine the nature of solutions for: (1) $1x+2y=8$, (2) $2x+4y=16$. Explain which method criterion you applied.

Q2188. Without solving, determine the nature of solutions for: (1) $6x+5y=4$, (2) $12x+10y=11$. Explain which method criterion you applied.

Q2189. Without solving, determine the nature of solutions for: (1) $1x+6y=7$, (2) $4x+18y=21$. Explain which method criterion you applied.

Q2190. Without solving, determine the nature of solutions for: (1) $3x+1y=8$, (2) $6x+2y=16$. Explain which method criterion you applied.

Q2191. Without solving, determine the nature of solutions for: (1) $1x+4y=5$, (2) $4x+16y=23$. Explain

which method criterion you applied.

Q2192. Without solving, determine the nature of solutions for: (1) $1x + 6y = 10$, (2) $3x + 12y = 20$. Explain which method criterion you applied.

Q2193. Without solving, determine the nature of solutions for: (1) $1x + 6y = 7$, (2) $3x + 18y = 21$. Explain which method criterion you applied.

Q2194. Without solving, determine the nature of solutions for: (1) $4x + 5y = 4$, (2) $8x + 10y = 11$. Explain which method criterion you applied.

Q2195. Without solving, determine the nature of solutions for: (1) $1x+5y = 2$, (2) $4x+15y = 6$. Explain which method criterion you applied.

Q2196. Without solving, determine the nature of solutions for: (1) $6x + 6y = 9$, (2) $24x + 24y = 36$. Explain which method criterion you applied.

Q2197. Without solving, determine the nature of solutions for: (1) $5x + 4y = 3$, (2) $15x + 12y = 12$. Explain which method criterion you applied.

Q2198. Without solving, determine the nature of solutions for: (1) $2x + 5y = 9$, (2) $9x + 20y = 36$. Explain which method criterion you applied.

Q2199. Without solving, determine the nature of solutions for: (1) $2x+1y = 5$, (2) $6x+3y = 15$. Explain which method criterion you applied.

Q2200. Without solving, determine the nature of solutions for: (1) $1x+5y = 2$, (2) $3x+15y = 9$. Explain which method criterion you applied.

Q2201. Without solving, determine the nature of solutions for: (1) $1x + 5y = 8$, (2) $3x + 10y = 16$. Explain which method criterion you applied.

Q2202. Without solving, determine the nature of solutions for: (1) $2x + 4y = 4$, (2) $8x + 16y = 16$. Explain which method criterion you applied.

Q2203. Without solving, determine the nature of solutions for: (1) $6x + 4y = 2$, (2) $24x + 16y = 11$. Explain which method criterion you applied.

Q2204. Without solving, determine the nature of solutions for: (1) $2x + 4y = 10$, (2) $5x + 8y = 20$. Explain which method criterion you applied.

Q2205. Without solving, determine the nature of solutions for: (1) $3x+4y = 5$, (2) $6x+8y = 10$. Explain which method criterion you applied.

Q2206. Without solving, determine the nature of solutions for: (1) $1x + 4y = 3$, (2) $4x + 16y = 15$. Explain which method criterion you applied.

Q2207. Without solving, determine the nature of solutions for: (1) $3x+2y = 7$, (2) $7x+4y = 14$. Explain which method criterion you applied.

Q2208. Without solving, determine the nature of solutions for: (1) $3x+1y = 8$, (2) $9x+3y = 24$. Explain which method criterion you applied.

Q2209. Without solving, determine the nature of solutions for: (1) $4x + 6y = 3$, (2) $16x + 24y = 15$. Explain which method criterion you applied.

Q2210. Without solving, determine the nature of solutions for: (1) $4x + 2y = 6$, (2) $17x + 8y = 24$. Explain which method criterion you applied.

Q2211. Without solving, determine the nature of solutions for: (1) $6x + 4y = 5$, (2) $18x + 12y = 15$. Explain which method criterion you applied.

Q2212. Without solving, determine the nature of solutions for: (1) $6x + 4y = 2$, (2) $18x + 12y = 9$. Explain which method criterion you applied.

Q2213. Without solving, determine the nature of solutions for: (1) $2x + 2y = 3$, (2) $7x + 6y = 9$. Explain which method criterion you applied.

Q2214. Without solving, determine the nature of solutions for: (1) $1x + 3y = 8$, (2) $4x + 12y = 32$. Explain which method criterion you applied.

Q2215. Without solving, determine the nature of solutions for: (1) $6x + 5y = 6$, (2) $18x + 15y = 21$. Explain which method criterion you applied.

Q2216. Without solving, determine the nature of solutions for: (1) $6x + 2y = 10$, (2) $19x + 6y = 30$. Explain which method criterion you applied.

Q2217. Without solving, determine the nature of solutions for: (1) $2x+2y = 7$, (2) $6x+6y = 21$. Explain which method criterion you applied.

Q2218. Without solving, determine the nature of solutions for: (1) $3x + 1y = 10$, (2) $6x + 2y = 23$. Explain which method criterion you applied.

Q2219. Without solving, determine the nature of solutions for: (1) $4x + 4y = 10$, (2) $17x + 16y = 40$. Explain which method criterion you applied.

Q2220. Without solving, determine the nature of solutions for: (1) $4x + 6y = 2$, (2) $12x + 18y = 6$. Explain which method criterion you applied.

Q2221. Without solving, determine the nature of solutions for: (1) $1x+1y = 6$, (2) $3x+3y = 21$. Explain which method criterion you applied.

Q2222. Without solving, determine the nature of solutions for: (1) $1x + 6y = 4$, (2) $5x + 24y = 16$. Explain which method criterion you applied.

Q2223. Without solving, determine the nature of solutions for: (1) $1x+1y = 3$, (2) $4x+4y = 12$. Explain which method criterion you applied.

Q2224. Without solving, determine the nature of solutions for: (1) $6x + 5y = 8$, (2) $18x + 15y = 27$. Explain which method criterion you applied.

Q2225. Without solving, determine the nature of solutions for: (1) $4x + 6y = 5$, (2) $17x + 24y = 20$. Explain which method criterion you applied.

Q2226. Without solving, determine the nature of solutions for: (1) $4x + 1y = 6$, (2) $16x + 4y = 24$. Explain which method criterion you applied.

Q2227. Without solving, determine the nature of solutions for: (1) $5x + 2y = 4$, (2) $15x + 6y = 15$. Explain which method criterion you applied.

Q2228. Without solving, determine the nature of solutions for: (1) $1x+1y = 6$, (2) $4x+3y = 18$. Explain which method criterion you applied.

Q2229. Without solving, determine the nature of solutions for: (1) $1x+1y = 3$, (2) $4x+4y = 12$. Explain which method criterion you applied.

Q2230. Without solving, determine the nature of solu-

tions for: (1) $6x + 1y = 5$, (2) $24x + 4y = 23$. Explain which method criterion you applied.

Q2231. Without solving, determine the nature of solutions for: (1) $1x + 2y = 2$, (2) $4x + 6y = 6$. Explain which method criterion you applied.

Q2232. Without solving, determine the nature of solutions for: (1) $5x + 3y = 8$, (2) $15x + 9y = 24$. Explain which method criterion you applied.

Q2233. Without solving, determine the nature of solutions for: (1) $3x + 4y = 10$, (2) $9x + 12y = 33$. Explain which method criterion you applied.

Q2234. Without solving, determine the nature of solutions for: (1) $6x + 3y = 5$, (2) $25x + 12y = 20$. Explain which method criterion you applied.

Q2235. Without solving, determine the nature of solutions for: (1) $1x + 5y = 8$, (2) $3x + 15y = 24$. Explain which method criterion you applied.

Q2236. Without solving, determine the nature of solutions for: (1) $3x + 5y = 9$, (2) $12x + 20y = 39$. Explain which method criterion you applied.

Q2237. Without solving, determine the nature of solutions for: (1) $3x + 5y = 6$, (2) $10x + 15y = 18$. Explain which method criterion you applied.

Q2238. Without solving, determine the nature of solutions for: (1) $5x + 2y = 7$, (2) $10x + 4y = 14$. Explain which method criterion you applied.

Q2239. Without solving, determine the nature of solutions for: (1) $1x + 5y = 7$, (2) $3x + 15y = 24$. Explain which method criterion you applied.

Q2240. Without solving, determine the nature of solutions for: (1) $4x + 2y = 2$, (2) $13x + 6y = 6$. Explain which method criterion you applied.

Q2241. Without solving, determine the nature of solutions for: (1) $4x + 6y = 6$, (2) $16x + 24y = 24$. Explain which method criterion you applied.

Q2242. Without solving, determine the nature of solutions for: (1) $2x + 2y = 7$, (2) $4x + 4y = 17$. Explain which method criterion you applied.

Q2243. Without solving, determine the nature of solutions for: (1) $3x + 1y = 6$, (2) $13x + 4y = 24$. Explain which method criterion you applied.

Q2244. Without solving, determine the nature of solutions for: (1) $3x + 3y = 5$, (2) $12x + 12y = 20$. Explain which method criterion you applied.

Q2245. Without solving, determine the nature of solutions for: (1) $1x + 2y = 5$, (2) $4x + 8y = 23$. Explain which method criterion you applied.

Q2246. Without solving, determine the nature of solutions for: (1) $2x + 1y = 5$, (2) $5x + 2y = 10$. Explain which method criterion you applied.

Q2247. Without solving, determine the nature of solutions for: (1) $2x + 4y = 9$, (2) $6x + 12y = 27$. Explain which method criterion you applied.

Q2248. Without solving, determine the nature of solutions for: (1) $1x + 3y = 5$, (2) $4x + 12y = 23$. Explain which method criterion you applied.

Q2249. Without solving, determine the nature of solutions for: (1) $3x + 1y = 4$, (2) $13x + 4y = 16$. Explain which method criterion you applied.

Q2250. Without solving, determine the nature of solutions for: (1) $3x + 5y = 10$, (2) $12x + 20y = 40$. Explain which method criterion you applied.

Q2251. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2252. For the system: (1) $x = 5y - 5$, (2) $4x + 4y = 22$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2253. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2254. For the system: (1) $x = 2y - 2$, (2) $3x + 2y = 22$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2255. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2256. For the system: (1) $x = 4y - 2$, (2) $6x + 4y = 40$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2257. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2258. For the system: (1) $x = 2y - 6$, (2) $3x + 4y = 19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2259. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2260. For the system: (1) $x = 3y - 8$, (2) $3x + 6y = 40$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2261. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2262. For the system: (1) $x = 4y - 1$, (2) $7x + 5y = 12$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2263. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2264. For the system: (1) $x = 4y - 9$, (2) $7x + 5y = 16$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2265. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2266. For the system: (1) $x = 5y - 6$, (2) $7x + 5y = 11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2267. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2268. For the system: (1) $x = 3y - 6$, (2) $6x + 2y = 23$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2269. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2270. For the system: (1) $x = 5y - 8$, (2) $4x + 5y = 13$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2271. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2272. For the system: (1) $x = 3y - 5$, (2) $4x + 5y = 33$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2273. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2274. For the system: (1) $x = 5y - 8$, (2) $6x + 5y = 28$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2275. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2276. For the system: (1) $x = 3y - 8$, (2) $3x + 3y = 11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2277. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2278. For the system: (1) $x = 4y - 5$, (2) $4x + 5y = 22$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2279. For the system: (1) $5x + 5y = 20$, (2) $5x - 3y = 4$, which method minimizes arithmetic steps?

Q2280. For the system: (1) $x = 2y - 6$, (2) $7x + 6y = 30$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2281. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2282. For the system: (1) $x = 2y - 9$, (2) $4x + 3y = 23$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2283. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2284. For the system: (1) $x = 5y - 1$, (2) $6x + 6y = 31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2285. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2286. For the system: (1) $x = 5y - 4$, (2) $6x + 6y = 25$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2287. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2288. For the system: (1) $x = 5y - 3$, (2) $4x + 5y = 12$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2289. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2290. For the system: (1) $x = 3y - 2$, (2) $3x + 5y = 15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2291. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2292. For the system: (1) $x = 5y - 5$, (2) $5x + 3y = 15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2293. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2294. For the system: (1) $x = 3y - 7$, (2) $7x + 5y = 26$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2295. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2296. For the system: (1) $x = 2y - 3$, (2) $7x + 5y = 36$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2297. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2298. For the system: (1) $x = 2y - 1$, (2) $5x + 2y = 29$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2299. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2300. For the system: (1) $x = 2y - 9$, (2) $5x + 3y = 16$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2301. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2302. For the system: (1) $x = 4y - 8$, (2) $7x + 3y = 20$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2303. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2304. For the system: (1) $x = 4y - 3$, (2) $6x + 5y = 22$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2305. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2306. For the system: (1) $x = 5y - 1$, (2) $7x + 4y = 34$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2307. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2308. For the system: (1) $x = 2y - 2$, (2) $4x + 3y = 38$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2309. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2310. For the system: (1) $x = 3y - 4$, (2) $3x + 5y = 17$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2311. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2312. For the system: (1) $x = 5y - 6$, (2) $5x + 5y = 28$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2313. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2314. For the system: (1) $x = 2y - 7$, (2) $3x + 3y = 27$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2315. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2316. For the system: (1) $x = 4y - 7$, (2) $4x + 4y = 39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2317. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2318. For the system: (1) $x=3y-5$, (2) $6x+3y=30$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2319. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2320. For the system: (1) $x=3y-5$, (2) $4x+6y=13$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2321. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2322. For the system: (1) $x=5y-9$, (2) $4x+5y=15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2323. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2324. For the system: (1) $x=5y-1$, (2) $7x+3y=34$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2325. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2326. For the system: (1) $x=2y-6$, (2) $3x+2y=23$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2327. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2328. For the system: (1) $x=4y-9$, (2) $4x+2y=17$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2329. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2330. For the system: (1) $x=5y-7$, (2) $4x+2y=35$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2331. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2332. For the system: (1) $x=4y-5$, (2) $5x+4y=38$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2333. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2334. For the system: (1) $x=3y-1$, (2) $3x+2y=10$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2335. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2336. For the system: (1) $x=2y-6$, (2) $3x+2y=26$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2337. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2338. For the system: (1) $x=5y-3$, (2) $7x+5y=16$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2339. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2340. For the system: (1) $x=2y-7$, (2) $5x+2y=36$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2341. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2342. For the system: (1) $x=4y-1$, (2) $6x+4y=34$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2343. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2344. For the system: (1) $x=2y-9$, (2) $4x+5y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2345. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2346. For the system: (1) $x=3y-6$, (2) $4x+5y=26$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2347. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2348. For the system: (1) $x=5y-6$, (2) $7x+6y=40$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2349. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2350. For the system: (1) $x=2y-5$, (2) $3x+3y=19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2351. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2352. For the system: (1) $x=3y-5$, (2) $7x+2y=18$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2353. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2354. For the system: (1) $x=3y-8$, (2) $3x+4y=37$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2355. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2356. For the system: (1) $x=2y-8$, (2) $7x+5y=38$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2357. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2358. For the system: (1) $x=4y-9$, (2) $6x+2y=11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2359. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2360. For the system: (1) $x=2y-4$, (2) $3x+4y=39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2361. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2362. For the system: (1) $x=2y-2$, (2) $5x+3y=24$, which algebraic method (Substitution or Elimination) is

more direct and why?

Q2363. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2364. For the system: (1) $x=5y-7$, (2) $4x+6y=18$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2365. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2366. For the system: (1) $x=4y-7$, (2) $4x+2y=39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2367. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2368. For the system: (1) $x=3y-5$, (2) $5x+4y=10$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2369. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2370. For the system: (1) $x=5y-3$, (2) $3x+5y=32$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2371. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2372. For the system: (1) $x=3y-5$, (2) $4x+2y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2373. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2374. For the system: (1) $x=5y-2$, (2) $5x+6y=26$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2375. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2376. For the system: (1) $x=4y-9$, (2) $6x+2y=11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2377. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2378. For the system: (1) $x=2y-1$, (2) $6x+5y=15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2379. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2380. For the system: (1) $x=5y-4$, (2) $6x+6y=13$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2381. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2382. For the system: (1) $x=4y-2$, (2) $7x+6y=25$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2383. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2384. For the system: (1) $x=3y-6$, (2) $7x+6y=31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2385. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2386. For the system: (1) $x=3y-2$, (2) $6x+3y=22$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2387. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2388. For the system: (1) $x=5y-4$, (2) $6x+2y=37$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2389. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2390. For the system: (1) $x=5y-8$, (2) $3x+6y=19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2391. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2392. For the system: (1) $x=2y-5$, (2) $3x+3y=35$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2393. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2394. For the system: (1) $x=5y-5$, (2) $7x+3y=19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2395. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2396. For the system: (1) $x=2y-8$, (2) $3x+2y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2397. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2398. For the system: (1) $x=4y-9$, (2) $7x+2y=25$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2399. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2400. For the system: (1) $x=5y-1$, (2) $7x+2y=34$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2401. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2402. For the system: (1) $x=4y-1$, (2) $6x+6y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2403. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2404. For the system: (1) $x=2y-9$, (2) $6x+6y=36$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2405. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2406. For the system: (1) $x=3y-8$, (2) $6x+2y=18$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2407. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2408. For the system: (1) $x = 5y - 5$, (2) $6x + 4y = 17$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2409. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2410. For the system: (1) $x = 2y - 3$, (2) $6x + 4y = 31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2411. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2412. For the system: (1) $x = 3y - 1$, (2) $5x + 5y = 32$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2413. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2414. For the system: (1) $x = 2y - 2$, (2) $5x + 3y = 15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2415. For the system: (1) $5x + 5y = 20$, (2) $5x - 3y = 4$, which method minimizes arithmetic steps?

Q2416. For the system: (1) $x = 3y - 3$, (2) $4x + 3y = 19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2417. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2418. For the system: (1) $x = 3y - 7$, (2) $7x + 5y = 19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2419. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2420. For the system: (1) $x = 4y - 1$, (2) $7x + 2y = 14$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2421. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2422. For the system: (1) $x = 4y - 4$, (2) $5x + 4y = 30$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2423. For the system: (1) $5x + 5y = 20$, (2) $5x - 3y = 4$, which method minimizes arithmetic steps?

Q2424. For the system: (1) $x = 3y - 3$, (2) $6x + 4y = 11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2425. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2426. For the system: (1) $x = 4y - 3$, (2) $4x + 3y = 14$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2427. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2428. For the system: (1) $x = 3y - 4$, (2) $7x + 4y = 27$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2429. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2430. For the system: (1) $x = 3y - 5$, (2) $5x + 2y = 12$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2431. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2432. For the system: (1) $x = 4y - 3$, (2) $4x + 6y = 31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2433. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2434. For the system: (1) $x = 4y - 6$, (2) $3x + 3y = 17$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2435. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2436. For the system: (1) $x = 3y - 2$, (2) $4x + 2y = 27$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2437. For the system: (1) $6x + 5y = 20$, (2) $6x - 3y = 4$, which method minimizes arithmetic steps?

Q2438. For the system: (1) $x = 3y - 3$, (2) $5x + 4y = 13$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2439. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2440. For the system: (1) $x = 3y - 5$, (2) $4x + 3y = 11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2441. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2442. For the system: (1) $x = 5y - 7$, (2) $7x + 4y = 10$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2443. For the system: (1) $4x + 5y = 20$, (2) $4x - 3y = 4$, which method minimizes arithmetic steps?

Q2444. For the system: (1) $x = 4y - 8$, (2) $4x + 5y = 32$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2445. For the system: (1) $8x + 5y = 20$, (2) $8x - 3y = 4$, which method minimizes arithmetic steps?

Q2446. For the system: (1) $x = 4y - 3$, (2) $7x + 5y = 10$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2447. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2448. For the system: (1) $x = 4y - 3$, (2) $4x + 4y = 13$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2449. For the system: (1) $7x + 5y = 20$, (2) $7x - 3y = 4$, which method minimizes arithmetic steps?

Q2450. For the system: (1) $x = 4y - 6$, (2) $3x + 3y = 31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2451. For the system: (1) $3x + 5y = 20$, (2) $3x - 3y = 4$, which method minimizes arithmetic steps?

Q2452. For the system: (1) $x = 3y - 4$, (2) $3x + 5y = 28$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2453. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2454. For the system: (1) $x=2y-3$, (2) $5x+3y=10$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2455. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2456. For the system: (1) $x=3y-6$, (2) $5x+4y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2457. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2458. For the system: (1) $x=3y-7$, (2) $3x+5y=36$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2459. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2460. For the system: (1) $x=5y-3$, (2) $7x+3y=11$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2461. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2462. For the system: (1) $x=2y-3$, (2) $4x+6y=15$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2463. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2464. For the system: (1) $x=2y-2$, (2) $5x+6y=39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2465. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2466. For the system: (1) $x=2y-2$, (2) $3x+3y=39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2467. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2468. For the system: (1) $x=4y-2$, (2) $4x+4y=31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2469. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2470. For the system: (1) $x=2y-8$, (2) $4x+2y=14$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2471. For the system: (1) $3x+5y=20$, (2) $3x-3y=4$, which method minimizes arithmetic steps?

Q2472. For the system: (1) $x=2y-5$, (2) $7x+2y=36$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2473. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2474. For the system: (1) $x=5y-2$, (2) $5x+3y=32$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2475. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2476. For the system: (1) $x=5y-6$, (2) $7x+5y=21$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2477. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2478. For the system: (1) $x=2y-8$, (2) $4x+3y=34$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2479. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2480. For the system: (1) $x=3y-9$, (2) $3x+5y=19$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2481. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2482. For the system: (1) $x=3y-2$, (2) $4x+2y=31$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2483. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2484. For the system: (1) $x=2y-4$, (2) $6x+6y=38$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2485. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2486. For the system: (1) $x=5y-8$, (2) $4x+4y=28$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2487. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2488. For the system: (1) $x=3y-6$, (2) $3x+3y=14$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2489. For the system: (1) $5x+5y=20$, (2) $5x-3y=4$, which method minimizes arithmetic steps?

Q2490. For the system: (1) $x=2y-6$, (2) $5x+3y=24$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2491. For the system: (1) $8x+5y=20$, (2) $8x-3y=4$, which method minimizes arithmetic steps?

Q2492. For the system: (1) $x=4y-6$, (2) $4x+5y=30$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2493. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2494. For the system: (1) $x=2y-5$, (2) $3x+3y=18$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2495. For the system: (1) $4x+5y=20$, (2) $4x-3y=4$, which method minimizes arithmetic steps?

Q2496. For the system: (1) $x=4y-4$, (2) $6x+2y=39$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2497. For the system: (1) $7x+5y=20$, (2) $7x-3y=4$, which method minimizes arithmetic steps?

Q2498. For the system: (1) $x=4y-4$, (2) $6x+6y=27$, which algebraic method (Substitution or Elimination) is

more direct and why?

Q2499. For the system: (1) $6x+5y=20$, (2) $6x-3y=4$, which method minimizes arithmetic steps?

Q2500. For the system: (1) $x=4y-3$, (2) $5x+6y=28$, which algebraic method (Substitution or Elimination) is more direct and why?

Q2501. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2502. Consider $x^2+5x+6=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2503. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2504. Consider $x^2+3x+2=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2505. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2506. Consider $x^2+4x+4=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2507. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2508. Consider $x^2+9x+18=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2509. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2510. Consider $x^2+3x+2=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2511. Consider $\sqrt{2}x^2+7x+3\sqrt{2}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2512. Consider $x^2+6x+8=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2513. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2514. Consider $x^2+11x+30=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2515. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2516. Consider $x^2+5x+4=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2517. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2518. Consider $x^2+10x+24=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2519. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2520. Consider $x^2+11x+30=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2521. Consider $\sqrt{2}x^2+7x+3\sqrt{2}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2522. Consider $x^2+12x+36=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2523. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2524. Consider $x^2+6x+8=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2525. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2526. Consider $x^2+7x+6=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2527. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2528. Consider $x^2+8x+16=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2529. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2530. Consider $x^2+5x+4=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2531. Consider $\sqrt{3}x^2+7x+3\sqrt{3}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2532. Consider $x^2+5x+6=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2533. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2534. Consider $x^2+8x+12=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2535. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2536. Consider $x^2+2x+1=0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2537. Consider $\sqrt{5}x^2+7x+3\sqrt{5}=0$. Is Middle-Term

Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2538. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2539. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2540. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2541. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2542. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2543. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2544. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2545. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2546. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2547. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2548. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2549. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2550. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2551. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2552. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2553. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2554. Consider $x^2 + 10x + 25 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2555. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2556. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2557. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2558. Consider $x^2 + 7x + 10 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2559. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2560. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2561. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2562. Consider $x^2 + 5x + 4 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2563. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2564. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2565. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2566. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2567. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2568. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2569. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2570. Consider $x^2 + 7x + 10 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2571. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2572. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2573. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2574. Consider $x^2 + 8x + 15 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2575. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2576. Consider $x^2 + 6x + 8 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2577. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2578. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2579. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2580. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2581. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2582. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2583. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2584. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2585. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2586. Consider $x^2 + 2x + 1 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2587. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2588. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2589. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2590. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2591. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2592. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2593. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2594. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2595. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2596. Consider $x^2 + 8x + 15 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2597. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2598. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2599. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2600. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2601. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2602. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2603. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2604. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2605. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2606. Consider $x^2 + 6x + 8 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2607. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2608. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2609. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2610. Consider $x^2 + 7x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2611. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2612. Consider $x^2 + 2x + 1 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2613. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2614. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula?

Justify.

Q2615. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2616. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2617. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2618. Consider $x^2 + 8x + 16 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2619. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2620. Consider $x^2 + 6x + 8 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2621. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2622. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2623. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2624. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2625. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2626. Consider $x^2 + 7x + 10 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2627. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2628. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2629. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2630. Consider $x^2 + 5x + 4 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2631. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2632. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2633. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2634. Consider $x^2 + 7x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2635. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2636. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2637. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2638. Consider $x^2 + 8x + 16 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2639. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2640. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2641. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2642. Consider $x^2 + 7x + 10 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2643. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2644. Consider $x^2 + 7x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2645. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2646. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2647. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2648. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2649. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2650. Consider $x^2 + 2x + 1 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2651. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2652. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2653. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term

Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2654. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2655. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2656. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2657. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2658. Consider $x^2 + 8x + 15 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2659. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2660. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2661. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2662. Consider $x^2 + 5x + 4 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2663. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2664. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2665. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2666. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2667. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2668. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2669. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2670. Consider $x^2 + 8x + 15 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2671. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2672. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2673. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2674. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2675. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2676. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2677. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2678. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2679. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2680. Consider $x^2 + 7x + 10 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2681. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2682. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2683. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2684. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2685. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2686. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2687. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2688. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2689. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2690. Consider $x^2 + 6x + 8 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2691. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2692. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2693. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2694. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2695. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2696. Consider $x^2 + 7x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2697. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2698. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2699. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2700. Consider $x^2 + 6x + 5 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2701. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2702. Consider $x^2 + 9x + 20 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2703. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2704. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2705. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2706. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2707. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2708. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2709. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2710. Consider $x^2 + 4x + 3 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2711. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2712. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2713. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2714. Consider $x^2 + 10x + 25 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2715. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2716. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2717. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2718. Consider $x^2 + 2x + 1 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2719. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2720. Consider $x^2 + 5x + 4 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2721. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2722. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2723. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2724. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2725. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2726. Consider $x^2 + 10x + 24 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2727. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2728. Consider $x^2 + 9x + 18 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2729. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2730. Consider $x^2 + 8x + 16 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula?

Justify.

Q2731. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2732. Consider $x^2 + 8x + 15 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2733. Consider $\sqrt{5}x^2 + 7x + 3\sqrt{5} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2734. Consider $x^2 + 6x + 8 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2735. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2736. Consider $x^2 + 8x + 12 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2737. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2738. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2739. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2740. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2741. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2742. Consider $x^2 + 6x + 9 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2743. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2744. Consider $x^2 + 10x + 25 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2745. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2746. Consider $x^2 + 11x + 30 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2747. Consider $\sqrt{2}x^2 + 7x + 3\sqrt{2} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2748. Consider $x^2 + 3x + 2 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2749. Consider $\sqrt{3}x^2 + 7x + 3\sqrt{3} = 0$. Is Middle-Term Splitting straightforward, or is Quadratic Formula preferred? Explain.

Q2750. Consider $x^2 + 5x + 6 = 0$. Should you use Factoring (Middle-Term Splitting) or the Quadratic Formula? Justify.

Q2751. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2752. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2753. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2754. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2755. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2756. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2757. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2758. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2759. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2760. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2761. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2762. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2763. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2764. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2765. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2766. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2767. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2768. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2769. For the system $\frac{4}{x} + \frac{3}{y} = 14$ and $\frac{3}{x} - \frac{4}{y} = 2$, why

is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2770. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2771. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2772. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2773. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2774. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2775. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2776. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2777. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2778. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2779. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2780. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2781. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2782. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2783. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2784. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2785. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2786. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2787. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2788. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2789. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2790. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2791. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2792. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2793. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2794. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2795. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2796. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2797. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2798. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2799. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2800. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2801. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2802. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2803. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2804. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2805. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2806. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2807. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2808. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2809. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2810. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2811. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2812. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2813. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2814. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2815. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2816. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2817. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2818. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2819. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2820. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2821. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2822. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2823. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2824. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2825. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2826. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2827. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2828. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2829. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2830. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2831. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2832. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2833. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2834. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2835. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2836. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2837. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2838. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2839. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2840. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2841. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2842. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2843. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2844. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2845. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2846. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitu-

tion should be used instead?

Q2847. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2848. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2849. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2850. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2851. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2852. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2853. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2854. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2855. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2856. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2857. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2858. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2859. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2860. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2861. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2862. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2863. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2864. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2865. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2866. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2867. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2868. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2869. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2870. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2871. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2872. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2873. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2874. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2875. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2876. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2877. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2878. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2879. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2880. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2881. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2882. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2883. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2884. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2885. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why

is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2886. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2887. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2888. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2889. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2890. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2891. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2892. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2893. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2894. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2895. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2896. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2897. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2898. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2899. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2900. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2901. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2902. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2903. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2904. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2905. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2906. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2907. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2908. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2909. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2910. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2911. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2912. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2913. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2914. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2915. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2916. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2917. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2918. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2919. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2920. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2921. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2922. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2923. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2924. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2925. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2926. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2927. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2928. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2929. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2930. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2931. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2932. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2933. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2934. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2935. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2936. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2937. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2938. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2939. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2940. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2941. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2942. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2943. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2944. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2945. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2946. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2947. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2948. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2949. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2950. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2951. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2952. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2953. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2954. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2955. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2956. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2957. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2958. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2959. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2960. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2961. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2962. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitu-

tion should be used instead?

Q2963. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2964. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2965. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2966. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2967. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2968. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2969. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2970. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2971. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2972. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2973. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2974. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2975. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2976. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2977. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2978. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2979. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2980. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2981. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2982. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2983. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2984. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2985. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2986. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2987. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2988. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2989. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2990. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2991. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2992. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2993. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2994. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2995. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2996. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2997. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2998. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q2999. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

Q3000. For the system $\frac{4}{x}+\frac{3}{y}=14$ and $\frac{3}{x}-\frac{4}{y}=2$, why is directly cross-multiplying ineffective, and what substitution should be used instead?

MODULE 4: DENOMINATOR RULES AND FRACTIONAL EQUATIONS (Q3001 – Q4000)

Q3001. Compare these two equations: (A) $\frac{x+8}{2} = 24$ vs (B) $\frac{x}{2} + 8 = 24$. Explain what happens to the denominator 2 in each case during the first step.

Q3002. Compare these two equations: (A) $\frac{x+8}{5} = 11$ vs (B) $\frac{x}{5} + 8 = 11$. Explain what happens to the denominator 5 in each case during the first step.

Q3003. Compare these two equations: (A) $\frac{x+8}{2} = 29$ vs (B) $\frac{x}{2} + 8 = 29$. Explain what happens to the denominator 2 in each case during the first step.

Q3004. Compare these two equations: (A) $\frac{x+6}{5} = 29$ vs (B) $\frac{x}{5} + 6 = 29$. Explain what happens to the denominator 5 in each case during the first step.

Q3005. Compare these two equations: (A) $\frac{x+3}{6} = 22$ vs (B) $\frac{x}{6} + 3 = 22$. Explain what happens to the denominator 6 in each case during the first step.

Q3006. Compare these two equations: (A) $\frac{x+2}{2} = 26$ vs (B) $\frac{x}{2} + 2 = 26$. Explain what happens to the denominator 2 in each case during the first step.

Q3007. Compare these two equations: (A) $\frac{x+9}{5} = 20$ vs (B) $\frac{x}{5} + 9 = 20$. Explain what happens to the denominator 5 in each case during the first step.

Q3008. Compare these two equations: (A) $\frac{x+6}{3} = 19$ vs (B) $\frac{x}{3} + 6 = 19$. Explain what happens to the denominator 3 in each case during the first step.

Q3009. Compare these two equations: (A) $\frac{x+9}{4} = 16$ vs (B) $\frac{x}{4} + 9 = 16$. Explain what happens to the denominator 4 in each case during the first step.

Q3010. Compare these two equations: (A) $\frac{x+2}{5} = 11$ vs (B) $\frac{x}{5} + 2 = 11$. Explain what happens to the denominator 5 in each case during the first step.

Q3011. Compare these two equations: (A) $\frac{x+8}{6} = 11$ vs (B) $\frac{x}{6} + 8 = 11$. Explain what happens to the denominator 6 in each case during the first step.

Q3012. Compare these two equations: (A) $\frac{x+2}{3} = 28$ vs (B) $\frac{x}{3} + 2 = 28$. Explain what happens to the denominator 3 in each case during the first step.

Q3013. Compare these two equations: (A) $\frac{x+5}{3} = 24$ vs (B) $\frac{x}{3} + 5 = 24$. Explain what happens to the denominator 3 in each case during the first step.

Q3014. Compare these two equations: (A) $\frac{x+2}{3} = 22$ vs (B) $\frac{x}{3} + 2 = 22$. Explain what happens to the denominator 3 in each case during the first step.

Q3015. Compare these two equations: (A) $\frac{x+5}{5} = 30$ vs (B) $\frac{x}{5} + 5 = 30$. Explain what happens to the denominator 5 in each case during the first step.

Q3016. Compare these two equations: (A) $\frac{x+6}{5} = 17$ vs (B) $\frac{x}{5} + 6 = 17$. Explain what happens to the denominator 5 in each case during the first step.

Q3017. Compare these two equations: (A) $\frac{x+6}{5} = 16$ vs (B) $\frac{x}{5} + 6 = 16$. Explain what happens to the denominator 5 in each case during the first step.

Q3018. Compare these two equations: (A) $\frac{x+9}{7} = 26$ vs (B) $\frac{x}{7} + 9 = 26$. Explain what happens to the denominator 7 in each case during the first step.

Q3019. Compare these two equations: (A) $\frac{x+5}{4} = 18$ vs (B) $\frac{x}{4} + 5 = 18$. Explain what happens to the denominator 4 in each case during the first step.

Q3020. Compare these two equations: (A) $\frac{x+6}{7} = 30$ vs (B) $\frac{x}{7} + 6 = 30$. Explain what happens to the denominator 7 in each case during the first step.

Q3021. Compare these two equations: (A) $\frac{x+5}{7} = 22$ vs (B) $\frac{x}{7} + 5 = 22$. Explain what happens to the denominator 7 in each case during the first step.

Q3022. Compare these two equations: (A) $\frac{x+5}{3} = 11$ vs (B) $\frac{x}{3} + 5 = 11$. Explain what happens to the denominator 3 in each case during the first step.

Q3023. Compare these two equations: (A) $\frac{x+8}{5} = 19$ vs (B) $\frac{x}{5} + 8 = 19$. Explain what happens to the denominator 5 in each case during the first step.

Q3024. Compare these two equations: (A) $\frac{x+9}{3} = 12$ vs (B) $\frac{x}{3} + 9 = 12$. Explain what happens to the denominator 3 in each case during the first step.

Q3025. Compare these two equations: (A) $\frac{x+7}{7} = 12$ vs (B) $\frac{x}{7} + 7 = 12$. Explain what happens to the denominator 7 in each case during the first step.

Q3026. Compare these two equations: (A) $\frac{x+4}{2} = 27$ vs (B) $\frac{x}{2} + 4 = 27$. Explain what happens to the denominator 2 in each case during the first step.

Q3027. Compare these two equations: (A) $\frac{x+6}{4} = 17$ vs (B) $\frac{x}{4} + 6 = 17$. Explain what happens to the denominator 4 in each case during the first step.

Q3028. Compare these two equations: (A) $\frac{x+9}{3} = 22$ vs (B) $\frac{x}{3} + 9 = 22$. Explain what happens to the denominator 3 in each case during the first step.

Q3029. Compare these two equations: (A) $\frac{x+5}{5} = 29$ vs (B) $\frac{x}{5} + 5 = 29$. Explain what happens to the denominator 5 in each case during the first step.

Q3030. Compare these two equations: (A) $\frac{x+2}{5} = 19$ vs (B) $\frac{x}{5} + 2 = 19$. Explain what happens to the denominator 5 in each case during the first step.

Q3031. Compare these two equations: (A) $\frac{x+5}{4} = 19$ vs (B) $\frac{x}{4} + 5 = 19$. Explain what happens to the denominator 4 in each case during the first step.

Q3032. Compare these two equations: (A) $\frac{x+8}{5} = 19$ vs (B) $\frac{x}{5} + 8 = 19$. Explain what happens to the denominator 5 in each case during the first step.

Q3033. Compare these two equations: (A) $\frac{x+9}{2} = 25$ vs (B) $\frac{x}{2} + 9 = 25$. Explain what happens to the denominator 2 in each case during the first step.

Q3034. Compare these two equations: (A) $\frac{x+8}{3} = 28$ vs (B) $\frac{x}{3} + 8 = 28$. Explain what happens to the denominator 3 in each case during the first step.

Q3035. Compare these two equations: (A) $\frac{x+6}{2} = 23$ vs (B) $\frac{x}{2} + 6 = 23$. Explain what happens to the denominator 2 in each case during the first step.

Q3036. Compare these two equations: (A) $\frac{x+2}{7} = 28$ vs (B) $\frac{x}{7} + 2 = 28$. Explain what happens to the denominator 7 in each case during the first step.

Q3037. Compare these two equations: (A) $\frac{x+9}{5} = 16$ vs (B) $\frac{x}{5} + 9 = 16$. Explain what happens to the denominator 5 in each case during the first step.

Q3038. Compare these two equations: (A) $\frac{x+6}{6} = 13$ vs (B) $\frac{x}{6} + 6 = 13$. Explain what happens to the denominator 6 in each case during the first step.

Q3039. Compare these two equations: (A) $\frac{x+9}{4} = 13$ vs (B) $\frac{x}{4} + 9 = 13$. Explain what happens to the denominator 4 in each case during the first step.

Q3040. Compare these two equations: (A) $\frac{x+4}{7} = 25$ vs (B) $\frac{x}{7} + 4 = 25$. Explain what happens to the denominator 7 in each case during the first step.

Q3041. Compare these two equations: (A) $\frac{x+9}{4} = 27$ vs (B) $\frac{x}{4} + 9 = 27$. Explain what happens to the denominator 4 in each case during the first step.

Q3042. Compare these two equations: (A) $\frac{x+6}{5} = 16$ vs (B) $\frac{x}{5} + 6 = 16$. Explain what happens to the denominator 5 in each case during the first step.

Q3043. Compare these two equations: (A) $\frac{x+9}{3} = 24$ vs (B) $\frac{x}{3} + 9 = 24$. Explain what happens to the denominator 3 in each case during the first step.

Q3044. Compare these two equations: (A) $\frac{x+2}{6} = 16$ vs (B) $\frac{x}{6} + 2 = 16$. Explain what happens to the denominator 6 in each case during the first step.

Q3045. Compare these two equations: (A) $\frac{x+9}{3} = 21$ vs (B) $\frac{x}{3} + 9 = 21$. Explain what happens to the denominator 3 in each case during the first step.

Q3046. Compare these two equations: (A) $\frac{x+2}{2} = 27$ vs (B) $\frac{x}{2} + 2 = 27$. Explain what happens to the denominator 2 in each case during the first step.

Q3047. Compare these two equations: (A) $\frac{x+2}{6} = 14$ vs (B) $\frac{x}{6} + 2 = 14$. Explain what happens to the denominator 6 in each case during the first step.

Q3048. Compare these two equations: (A) $\frac{x+7}{2} = 29$ vs (B) $\frac{x}{2} + 7 = 29$. Explain what happens to the denominator 2 in each case during the first step.

Q3049. Compare these two equations: (A) $\frac{x+2}{7} = 12$ vs (B) $\frac{x}{7} + 2 = 12$. Explain what happens to the denominator 7 in each case during the first step.

Q3050. Compare these two equations: (A) $\frac{x+3}{6} = 12$ vs (B) $\frac{x}{6} + 3 = 12$. Explain what happens to the denominator 6 in each case during the first step.

Q3051. Compare these two equations: (A) $\frac{x+3}{5} = 14$ vs (B) $\frac{x}{5} + 3 = 14$. Explain what happens to the denominator 5 in each case during the first step.

Q3052. Compare these two equations: (A) $\frac{x+2}{7} = 14$ vs (B) $\frac{x}{7} + 2 = 14$. Explain what happens to the denominator 7 in each case during the first step.

Q3053. Compare these two equations: (A) $\frac{x+4}{6} = 16$ vs (B) $\frac{x}{6} + 4 = 16$. Explain what happens to the denominator 6 in each case during the first step.

Q3054. Compare these two equations: (A) $\frac{x+4}{6} = 24$ vs (B) $\frac{x}{6} + 4 = 24$. Explain what happens to the denominator 6 in each case during the first step.

Q3055. Compare these two equations: (A) $\frac{x+2}{4} = 17$ vs (B) $\frac{x}{4} + 2 = 17$. Explain what happens to the denominator 4 in each case during the first step.

Q3056. Compare these two equations: (A) $\frac{x+5}{6} = 27$ vs (B) $\frac{x}{6} + 5 = 27$. Explain what happens to the denominator 6 in each case during the first step.

Q3057. Compare these two equations: (A) $\frac{x+2}{4} = 23$ vs (B) $\frac{x}{4} + 2 = 23$. Explain what happens to the denominator 4 in each case during the first step.

Q3058. Compare these two equations: (A) $\frac{x+5}{4} = 19$ vs (B) $\frac{x}{4} + 5 = 19$. Explain what happens to the denominator 4 in each case during the first step.

Q3059. Compare these two equations: (A) $\frac{x+6}{2} = 15$ vs (B) $\frac{x}{2} + 6 = 15$. Explain what happens to the denominator 2 in each case during the first step.

Q3060. Compare these two equations: (A) $\frac{x+4}{6} = 25$ vs (B) $\frac{x}{6} + 4 = 25$. Explain what happens to the denominator 6 in each case during the first step.

Q3061. Compare these two equations: (A) $\frac{x+3}{4} = 24$ vs (B) $\frac{x}{4} + 3 = 24$. Explain what happens to the denominator 4 in each case during the first step.

Q3062. Compare these two equations: (A) $\frac{x+6}{3} = 11$ vs (B) $\frac{x}{3} + 6 = 11$. Explain what happens to the denominator 3 in each case during the first step.

Q3063. Compare these two equations: (A) $\frac{x+9}{2} = 24$ vs (B) $\frac{x}{2} + 9 = 24$. Explain what happens to the denominator 2 in each case during the first step.

Q3064. Compare these two equations: (A) $\frac{x+7}{2} = 28$ vs (B) $\frac{x}{2} + 7 = 28$. Explain what happens to the denominator 2 in each case during the first step.

Q3065. Compare these two equations: (A) $\frac{x+3}{3} = 11$ vs (B) $\frac{x}{3} + 3 = 11$. Explain what happens to the denominator 3 in each case during the first step.

Q3066. Compare these two equations: (A) $\frac{x+3}{2} = 30$ vs (B) $\frac{x}{2} + 3 = 30$. Explain what happens to the denominator 2 in each case during the first step.

Q3067. Compare these two equations: (A) $\frac{x+6}{3} = 22$ vs (B) $\frac{x}{3} + 6 = 22$. Explain what happens to the denominator 3 in each case during the first step.

Q3068. Compare these two equations: (A) $\frac{x+6}{7} = 12$ vs (B) $\frac{x}{7} + 6 = 12$. Explain what happens to the denominator 7 in each case during the first step.

Q3069. Compare these two equations: (A) $\frac{x+3}{5} = 13$ vs (B) $\frac{x}{5} + 3 = 13$. Explain what happens to the denominator 5 in each case during the first step.

Q3070. Compare these two equations: (A) $\frac{x+5}{2} = 17$ vs (B) $\frac{x}{2} + 5 = 17$. Explain what happens to the denominator 2 in each case during the first step.

Q3071. Compare these two equations: (A) $\frac{x+4}{3} = 20$ vs (B) $\frac{x}{3} + 4 = 20$. Explain what happens to the denominator 3 in each case during the first step.

Q3072. Compare these two equations: (A) $\frac{x+5}{2} = 10$ vs (B) $\frac{x}{2} + 5 = 10$. Explain what happens to the denominator 2 in each case during the first step.

Q3073. Compare these two equations: (A) $\frac{x+7}{2} = 11$ vs (B) $\frac{x}{2} + 7 = 11$. Explain what happens to the denominator 2 in each case during the first step.

Q3074. Compare these two equations: (A) $\frac{x+9}{2} = 21$ vs (B) $\frac{x}{2} + 9 = 21$. Explain what happens to the denominator 2 in each case during the first step.

Q3075. Compare these two equations: (A) $\frac{x+9}{2} = 22$ vs (B) $\frac{x}{2} + 9 = 22$. Explain what happens to the denominator

2 in each case during the first step.

Q3076. Compare these two equations: (A) $\frac{x+3}{2} = 17$ vs (B) $\frac{x}{2} + 3 = 17$. Explain what happens to the denominator 2 in each case during the first step.

Q3077. Compare these two equations: (A) $\frac{x+6}{6} = 12$ vs (B) $\frac{x}{6} + 6 = 12$. Explain what happens to the denominator 6 in each case during the first step.

Q3078. Compare these two equations: (A) $\frac{x+6}{6} = 17$ vs (B) $\frac{x}{6} + 6 = 17$. Explain what happens to the denominator 6 in each case during the first step.

Q3079. Compare these two equations: (A) $\frac{x+8}{2} = 25$ vs (B) $\frac{x}{2} + 8 = 25$. Explain what happens to the denominator 2 in each case during the first step.

Q3080. Compare these two equations: (A) $\frac{x+8}{4} = 24$ vs (B) $\frac{x}{4} + 8 = 24$. Explain what happens to the denominator 4 in each case during the first step.

Q3081. Compare these two equations: (A) $\frac{x+8}{6} = 27$ vs (B) $\frac{x}{6} + 8 = 27$. Explain what happens to the denominator 6 in each case during the first step.

Q3082. Compare these two equations: (A) $\frac{x+4}{3} = 14$ vs (B) $\frac{x}{3} + 4 = 14$. Explain what happens to the denominator 3 in each case during the first step.

Q3083. Compare these two equations: (A) $\frac{x+4}{2} = 27$ vs (B) $\frac{x}{2} + 4 = 27$. Explain what happens to the denominator 2 in each case during the first step.

Q3084. Compare these two equations: (A) $\frac{x+5}{2} = 14$ vs (B) $\frac{x}{2} + 5 = 14$. Explain what happens to the denominator 2 in each case during the first step.

Q3085. Compare these two equations: (A) $\frac{x+3}{5} = 17$ vs (B) $\frac{x}{5} + 3 = 17$. Explain what happens to the denominator 5 in each case during the first step.

Q3086. Compare these two equations: (A) $\frac{x+3}{7} = 21$ vs (B) $\frac{x}{7} + 3 = 21$. Explain what happens to the denominator 7 in each case during the first step.

Q3087. Compare these two equations: (A) $\frac{x+3}{5} = 11$ vs (B) $\frac{x}{5} + 3 = 11$. Explain what happens to the denominator 5 in each case during the first step.

Q3088. Compare these two equations: (A) $\frac{x+6}{2} = 20$ vs (B) $\frac{x}{2} + 6 = 20$. Explain what happens to the denominator 2 in each case during the first step.

Q3089. Compare these two equations: (A) $\frac{x+9}{7} = 29$ vs (B) $\frac{x}{7} + 9 = 29$. Explain what happens to the denominator 7 in each case during the first step.

Q3090. Compare these two equations: (A) $\frac{x+3}{7} = 28$ vs (B) $\frac{x}{7} + 3 = 28$. Explain what happens to the denominator 7 in each case during the first step.

Q3091. Compare these two equations: (A) $\frac{x+5}{7} = 17$ vs (B) $\frac{x}{7} + 5 = 17$. Explain what happens to the denominator 7 in each case during the first step.

Q3092. Compare these two equations: (A) $\frac{x+2}{2} = 20$ vs (B) $\frac{x}{2} + 2 = 20$. Explain what happens to the denominator 2 in each case during the first step.

Q3093. Compare these two equations: (A) $\frac{x+2}{2} = 12$ vs (B) $\frac{x}{2} + 2 = 12$. Explain what happens to the denominator 2 in each case during the first step.

Q3094. Compare these two equations: (A) $\frac{x+8}{5} = 25$ vs (B) $\frac{x}{5} + 8 = 25$. Explain what happens to the denominator 5 in each case during the first step.

Q3095. Compare these two equations: (A) $\frac{x+6}{6} = 17$ vs (B) $\frac{x}{6} + 6 = 17$. Explain what happens to the denominator 6 in each case during the first step.

Q3096. Compare these two equations: (A) $\frac{x+7}{7} = 10$ vs (B) $\frac{x}{7} + 7 = 10$. Explain what happens to the denominator 7 in each case during the first step.

Q3097. Compare these two equations: (A) $\frac{x+4}{6} = 11$ vs (B) $\frac{x}{6} + 4 = 11$. Explain what happens to the denominator 6 in each case during the first step.

Q3098. Compare these two equations: (A) $\frac{x+6}{6} = 28$ vs (B) $\frac{x}{6} + 6 = 28$. Explain what happens to the denominator 6 in each case during the first step.

Q3099. Compare these two equations: (A) $\frac{x+9}{2} = 19$ vs (B) $\frac{x}{2} + 9 = 19$. Explain what happens to the denominator 2 in each case during the first step.

Q3100. Compare these two equations: (A) $\frac{x+6}{6} = 27$ vs (B) $\frac{x}{6} + 6 = 27$. Explain what happens to the denominator 6 in each case during the first step.

Q3101. Compare these two equations: (A) $\frac{x+6}{7} = 24$ vs (B) $\frac{x}{7} + 6 = 24$. Explain what happens to the denominator 7 in each case during the first step.

Q3102. Compare these two equations: (A) $\frac{x+7}{2} = 30$ vs (B) $\frac{x}{2} + 7 = 30$. Explain what happens to the denominator 2 in each case during the first step.

Q3103. Compare these two equations: (A) $\frac{x+3}{6} = 10$ vs (B) $\frac{x}{6} + 3 = 10$. Explain what happens to the denominator 6 in each case during the first step.

Q3104. Compare these two equations: (A) $\frac{x+2}{6} = 15$ vs (B) $\frac{x}{6} + 2 = 15$. Explain what happens to the denominator 6 in each case during the first step.

Q3105. Compare these two equations: (A) $\frac{x+8}{3} = 26$ vs (B) $\frac{x}{3} + 8 = 26$. Explain what happens to the denominator 3 in each case during the first step.

Q3106. Compare these two equations: (A) $\frac{x+4}{6} = 17$ vs (B) $\frac{x}{6} + 4 = 17$. Explain what happens to the denominator 6 in each case during the first step.

Q3107. Compare these two equations: (A) $\frac{x+4}{7} = 29$ vs (B) $\frac{x}{7} + 4 = 29$. Explain what happens to the denominator 7 in each case during the first step.

Q3108. Compare these two equations: (A) $\frac{x+3}{5} = 14$ vs (B) $\frac{x}{5} + 3 = 14$. Explain what happens to the denominator 5 in each case during the first step.

Q3109. Compare these two equations: (A) $\frac{x+2}{6} = 18$ vs (B) $\frac{x}{6} + 2 = 18$. Explain what happens to the denominator 6 in each case during the first step.

Q3110. Compare these two equations: (A) $\frac{x+2}{4} = 27$ vs (B) $\frac{x}{4} + 2 = 27$. Explain what happens to the denominator 4 in each case during the first step.

Q3111. Compare these two equations: (A) $\frac{x+4}{2} = 18$ vs (B) $\frac{x}{2} + 4 = 18$. Explain what happens to the denominator 2 in each case during the first step.

Q3112. Compare these two equations: (A) $\frac{x+4}{3} = 12$ vs (B) $\frac{x}{3} + 4 = 12$. Explain what happens to the denominator 3 in each case during the first step.

Q3113. Compare these two equations: (A) $\frac{x+2}{5} = 25$ vs (B) $\frac{x}{5} + 2 = 25$. Explain what happens to the denominator 5 in each case during the first step.

Q3114. Compare these two equations: (A) $\frac{x+8}{6} = 24$ vs

(B) $\frac{x}{6}+8=24$. Explain what happens to the denominator 6 in each case during the first step.

Q3115. Compare these two equations: (A) $\frac{x+9}{3}=28$ vs (B) $\frac{x}{3}+9=28$. Explain what happens to the denominator 3 in each case during the first step.

Q3116. Compare these two equations: (A) $\frac{x+2}{3}=13$ vs (B) $\frac{x}{3}+2=13$. Explain what happens to the denominator 3 in each case during the first step.

Q3117. Compare these two equations: (A) $\frac{x+3}{5}=23$ vs (B) $\frac{x}{5}+3=23$. Explain what happens to the denominator 5 in each case during the first step.

Q3118. Compare these two equations: (A) $\frac{x+6}{6}=13$ vs (B) $\frac{x}{6}+6=13$. Explain what happens to the denominator 6 in each case during the first step.

Q3119. Compare these two equations: (A) $\frac{x+4}{2}=26$ vs (B) $\frac{x}{2}+4=26$. Explain what happens to the denominator 2 in each case during the first step.

Q3120. Compare these two equations: (A) $\frac{x+7}{5}=12$ vs (B) $\frac{x}{5}+7=12$. Explain what happens to the denominator 5 in each case during the first step.

Q3121. Compare these two equations: (A) $\frac{x+6}{5}=26$ vs (B) $\frac{x}{5}+6=26$. Explain what happens to the denominator 5 in each case during the first step.

Q3122. Compare these two equations: (A) $\frac{x+4}{2}=23$ vs (B) $\frac{x}{2}+4=23$. Explain what happens to the denominator 2 in each case during the first step.

Q3123. Compare these two equations: (A) $\frac{x+2}{4}=25$ vs (B) $\frac{x}{4}+2=25$. Explain what happens to the denominator 4 in each case during the first step.

Q3124. Compare these two equations: (A) $\frac{x+9}{7}=29$ vs (B) $\frac{x}{7}+9=29$. Explain what happens to the denominator 7 in each case during the first step.

Q3125. Compare these two equations: (A) $\frac{x+8}{7}=23$ vs (B) $\frac{x}{7}+8=23$. Explain what happens to the denominator 7 in each case during the first step.

Q3126. Compare these two equations: (A) $\frac{x+9}{3}=16$ vs (B) $\frac{x}{3}+9=16$. Explain what happens to the denominator 3 in each case during the first step.

Q3127. Compare these two equations: (A) $\frac{x+9}{6}=29$ vs (B) $\frac{x}{6}+9=29$. Explain what happens to the denominator 6 in each case during the first step.

Q3128. Compare these two equations: (A) $\frac{x+4}{2}=24$ vs (B) $\frac{x}{2}+4=24$. Explain what happens to the denominator 2 in each case during the first step.

Q3129. Compare these two equations: (A) $\frac{x+2}{7}=22$ vs (B) $\frac{x}{7}+2=22$. Explain what happens to the denominator 7 in each case during the first step.

Q3130. Compare these two equations: (A) $\frac{x+5}{2}=11$ vs (B) $\frac{x}{2}+5=11$. Explain what happens to the denominator 2 in each case during the first step.

Q3131. Compare these two equations: (A) $\frac{x+2}{6}=14$ vs (B) $\frac{x}{6}+2=14$. Explain what happens to the denominator 6 in each case during the first step.

Q3132. Compare these two equations: (A) $\frac{x+9}{6}=17$ vs (B) $\frac{x}{6}+9=17$. Explain what happens to the denominator 6 in each case during the first step.

Q3133. Compare these two equations: (A) $\frac{x+2}{7}=21$ vs (B) $\frac{x}{7}+2=21$. Explain what happens to the denominator 7 in each case during the first step.

Q3134. Compare these two equations: (A) $\frac{x+9}{4}=28$ vs (B) $\frac{x}{4}+9=28$. Explain what happens to the denominator 4 in each case during the first step.

Q3135. Compare these two equations: (A) $\frac{x+4}{5}=21$ vs (B) $\frac{x}{5}+4=21$. Explain what happens to the denominator 5 in each case during the first step.

Q3136. Compare these two equations: (A) $\frac{x+6}{5}=11$ vs (B) $\frac{x}{5}+6=11$. Explain what happens to the denominator 5 in each case during the first step.

Q3137. Compare these two equations: (A) $\frac{x+8}{7}=17$ vs (B) $\frac{x}{7}+8=17$. Explain what happens to the denominator 7 in each case during the first step.

Q3138. Compare these two equations: (A) $\frac{x+6}{5}=28$ vs (B) $\frac{x}{5}+6=28$. Explain what happens to the denominator 5 in each case during the first step.

Q3139. Compare these two equations: (A) $\frac{x+6}{4}=22$ vs (B) $\frac{x}{4}+6=22$. Explain what happens to the denominator 4 in each case during the first step.

Q3140. Compare these two equations: (A) $\frac{x+5}{6}=20$ vs (B) $\frac{x}{6}+5=20$. Explain what happens to the denominator 6 in each case during the first step.

Q3141. Compare these two equations: (A) $\frac{x+7}{3}=19$ vs (B) $\frac{x}{3}+7=19$. Explain what happens to the denominator 3 in each case during the first step.

Q3142. Compare these two equations: (A) $\frac{x+3}{3}=21$ vs (B) $\frac{x}{3}+3=21$. Explain what happens to the denominator 3 in each case during the first step.

Q3143. Compare these two equations: (A) $\frac{x+3}{5}=19$ vs (B) $\frac{x}{5}+3=19$. Explain what happens to the denominator 5 in each case during the first step.

Q3144. Compare these two equations: (A) $\frac{x+5}{7}=21$ vs (B) $\frac{x}{7}+5=21$. Explain what happens to the denominator 7 in each case during the first step.

Q3145. Compare these two equations: (A) $\frac{x+4}{4}=13$ vs (B) $\frac{x}{4}+4=13$. Explain what happens to the denominator 4 in each case during the first step.

Q3146. Compare these two equations: (A) $\frac{x+2}{5}=19$ vs (B) $\frac{x}{5}+2=19$. Explain what happens to the denominator 5 in each case during the first step.

Q3147. Compare these two equations: (A) $\frac{x+8}{7}=10$ vs (B) $\frac{x}{7}+8=10$. Explain what happens to the denominator 7 in each case during the first step.

Q3148. Compare these two equations: (A) $\frac{x+4}{5}=23$ vs (B) $\frac{x}{5}+4=23$. Explain what happens to the denominator 5 in each case during the first step.

Q3149. Compare these two equations: (A) $\frac{x+7}{3}=19$ vs (B) $\frac{x}{3}+7=19$. Explain what happens to the denominator 3 in each case during the first step.

Q3150. Compare these two equations: (A) $\frac{x+8}{6}=16$ vs (B) $\frac{x}{6}+8=16$. Explain what happens to the denominator 6 in each case during the first step.

Q3151. Compare these two equations: (A) $\frac{x+5}{3}=29$ vs (B) $\frac{x}{3}+5=29$. Explain what happens to the denominator 3 in each case during the first step.

Q3152. Compare these two equations: (A) $\frac{x+2}{6}=24$ vs (B) $\frac{x}{6}+2=24$. Explain what happens to the denominator 6 in each case during the first step.

Q3153. Compare these two equations: (A) $\frac{x+6}{3} = 20$ vs (B) $\frac{x}{3} + 6 = 20$. Explain what happens to the denominator 3 in each case during the first step.

Q3154. Compare these two equations: (A) $\frac{x+5}{4} = 27$ vs (B) $\frac{x}{4} + 5 = 27$. Explain what happens to the denominator 4 in each case during the first step.

Q3155. Compare these two equations: (A) $\frac{x+3}{2} = 18$ vs (B) $\frac{x}{2} + 3 = 18$. Explain what happens to the denominator 2 in each case during the first step.

Q3156. Compare these two equations: (A) $\frac{x+4}{4} = 13$ vs (B) $\frac{x}{4} + 4 = 13$. Explain what happens to the denominator 4 in each case during the first step.

Q3157. Compare these two equations: (A) $\frac{x+8}{5} = 21$ vs (B) $\frac{x}{5} + 8 = 21$. Explain what happens to the denominator 5 in each case during the first step.

Q3158. Compare these two equations: (A) $\frac{x+5}{2} = 17$ vs (B) $\frac{x}{2} + 5 = 17$. Explain what happens to the denominator 2 in each case during the first step.

Q3159. Compare these two equations: (A) $\frac{x+4}{5} = 25$ vs (B) $\frac{x}{5} + 4 = 25$. Explain what happens to the denominator 5 in each case during the first step.

Q3160. Compare these two equations: (A) $\frac{x+3}{5} = 24$ vs (B) $\frac{x}{5} + 3 = 24$. Explain what happens to the denominator 5 in each case during the first step.

Q3161. Compare these two equations: (A) $\frac{x+5}{4} = 13$ vs (B) $\frac{x}{4} + 5 = 13$. Explain what happens to the denominator 4 in each case during the first step.

Q3162. Compare these two equations: (A) $\frac{x+3}{4} = 25$ vs (B) $\frac{x}{4} + 3 = 25$. Explain what happens to the denominator 4 in each case during the first step.

Q3163. Compare these two equations: (A) $\frac{x+6}{5} = 13$ vs (B) $\frac{x}{5} + 6 = 13$. Explain what happens to the denominator 5 in each case during the first step.

Q3164. Compare these two equations: (A) $\frac{x+7}{7} = 27$ vs (B) $\frac{x}{7} + 7 = 27$. Explain what happens to the denominator 7 in each case during the first step.

Q3165. Compare these two equations: (A) $\frac{x+9}{4} = 17$ vs (B) $\frac{x}{4} + 9 = 17$. Explain what happens to the denominator 4 in each case during the first step.

Q3166. Compare these two equations: (A) $\frac{x+8}{5} = 18$ vs (B) $\frac{x}{5} + 8 = 18$. Explain what happens to the denominator 5 in each case during the first step.

Q3167. Compare these two equations: (A) $\frac{x+8}{5} = 20$ vs (B) $\frac{x}{5} + 8 = 20$. Explain what happens to the denominator 5 in each case during the first step.

Q3168. Compare these two equations: (A) $\frac{x+7}{4} = 13$ vs (B) $\frac{x}{4} + 7 = 13$. Explain what happens to the denominator 4 in each case during the first step.

Q3169. Compare these two equations: (A) $\frac{x+4}{4} = 29$ vs (B) $\frac{x}{4} + 4 = 29$. Explain what happens to the denominator 4 in each case during the first step.

Q3170. Compare these two equations: (A) $\frac{x+9}{7} = 14$ vs (B) $\frac{x}{7} + 9 = 14$. Explain what happens to the denominator 7 in each case during the first step.

Q3171. Compare these two equations: (A) $\frac{x+3}{3} = 19$ vs (B) $\frac{x}{3} + 3 = 19$. Explain what happens to the denominator 3 in each case during the first step.

Q3172. Compare these two equations: (A) $\frac{x+4}{6} = 23$ vs (B) $\frac{x}{6} + 4 = 23$. Explain what happens to the denominator 6 in each case during the first step.

Q3173. Compare these two equations: (A) $\frac{x+9}{6} = 24$ vs (B) $\frac{x}{6} + 9 = 24$. Explain what happens to the denominator 6 in each case during the first step.

Q3174. Compare these two equations: (A) $\frac{x+5}{3} = 28$ vs (B) $\frac{x}{3} + 5 = 28$. Explain what happens to the denominator 3 in each case during the first step.

Q3175. Compare these two equations: (A) $\frac{x+6}{2} = 25$ vs (B) $\frac{x}{2} + 6 = 25$. Explain what happens to the denominator 2 in each case during the first step.

Q3176. Compare these two equations: (A) $\frac{x+4}{5} = 27$ vs (B) $\frac{x}{5} + 4 = 27$. Explain what happens to the denominator 5 in each case during the first step.

Q3177. Compare these two equations: (A) $\frac{x+8}{7} = 30$ vs (B) $\frac{x}{7} + 8 = 30$. Explain what happens to the denominator 7 in each case during the first step.

Q3178. Compare these two equations: (A) $\frac{x+5}{5} = 19$ vs (B) $\frac{x}{5} + 5 = 19$. Explain what happens to the denominator 5 in each case during the first step.

Q3179. Compare these two equations: (A) $\frac{x+4}{6} = 24$ vs (B) $\frac{x}{6} + 4 = 24$. Explain what happens to the denominator 6 in each case during the first step.

Q3180. Compare these two equations: (A) $\frac{x+9}{2} = 14$ vs (B) $\frac{x}{2} + 9 = 14$. Explain what happens to the denominator 2 in each case during the first step.

Q3181. Compare these two equations: (A) $\frac{x+8}{3} = 14$ vs (B) $\frac{x}{3} + 8 = 14$. Explain what happens to the denominator 3 in each case during the first step.

Q3182. Compare these two equations: (A) $\frac{x+5}{3} = 16$ vs (B) $\frac{x}{3} + 5 = 16$. Explain what happens to the denominator 3 in each case during the first step.

Q3183. Compare these two equations: (A) $\frac{x+4}{5} = 23$ vs (B) $\frac{x}{5} + 4 = 23$. Explain what happens to the denominator 5 in each case during the first step.

Q3184. Compare these two equations: (A) $\frac{x+2}{4} = 23$ vs (B) $\frac{x}{4} + 2 = 23$. Explain what happens to the denominator 4 in each case during the first step.

Q3185. Compare these two equations: (A) $\frac{x+5}{3} = 15$ vs (B) $\frac{x}{3} + 5 = 15$. Explain what happens to the denominator 3 in each case during the first step.

Q3186. Compare these two equations: (A) $\frac{x+6}{6} = 21$ vs (B) $\frac{x}{6} + 6 = 21$. Explain what happens to the denominator 6 in each case during the first step.

Q3187. Compare these two equations: (A) $\frac{x+2}{4} = 14$ vs (B) $\frac{x}{4} + 2 = 14$. Explain what happens to the denominator 4 in each case during the first step.

Q3188. Compare these two equations: (A) $\frac{x+8}{4} = 26$ vs (B) $\frac{x}{4} + 8 = 26$. Explain what happens to the denominator 4 in each case during the first step.

Q3189. Compare these two equations: (A) $\frac{x+2}{5} = 16$ vs (B) $\frac{x}{5} + 2 = 16$. Explain what happens to the denominator 5 in each case during the first step.

Q3190. Compare these two equations: (A) $\frac{x+6}{2} = 28$ vs (B) $\frac{x}{2} + 6 = 28$. Explain what happens to the denominator 2 in each case during the first step.

Q3191. Compare these two equations: (A) $\frac{x+9}{7} = 20$ vs (B) $\frac{x}{7} + 9 = 20$. Explain what happens to the denominator

7 in each case during the first step.

Q3192. Compare these two equations: (A) $\frac{x+9}{3} = 20$ vs (B) $\frac{x}{3} + 9 = 20$. Explain what happens to the denominator 3 in each case during the first step.

Q3193. Compare these two equations: (A) $\frac{x+7}{5} = 21$ vs (B) $\frac{x}{5} + 7 = 21$. Explain what happens to the denominator 5 in each case during the first step.

Q3194. Compare these two equations: (A) $\frac{x+7}{6} = 30$ vs (B) $\frac{x}{6} + 7 = 30$. Explain what happens to the denominator 6 in each case during the first step.

Q3195. Compare these two equations: (A) $\frac{x+2}{6} = 22$ vs (B) $\frac{x}{6} + 2 = 22$. Explain what happens to the denominator 6 in each case during the first step.

Q3196. Compare these two equations: (A) $\frac{x+2}{5} = 16$ vs (B) $\frac{x}{5} + 2 = 16$. Explain what happens to the denominator 5 in each case during the first step.

Q3197. Compare these two equations: (A) $\frac{x+4}{7} = 22$ vs (B) $\frac{x}{7} + 4 = 22$. Explain what happens to the denominator 7 in each case during the first step.

Q3198. Compare these two equations: (A) $\frac{x+8}{2} = 18$ vs (B) $\frac{x}{2} + 8 = 18$. Explain what happens to the denominator 2 in each case during the first step.

Q3199. Compare these two equations: (A) $\frac{x+6}{2} = 10$ vs (B) $\frac{x}{2} + 6 = 10$. Explain what happens to the denominator 2 in each case during the first step.

Q3200. Compare these two equations: (A) $\frac{x+5}{6} = 28$ vs (B) $\frac{x}{6} + 5 = 28$. Explain what happens to the denominator 6 in each case during the first step.

Q3201. Compare these two equations: (A) $\frac{x+6}{6} = 26$ vs (B) $\frac{x}{6} + 6 = 26$. Explain what happens to the denominator 6 in each case during the first step.

Q3202. Compare these two equations: (A) $\frac{x+8}{5} = 19$ vs (B) $\frac{x}{5} + 8 = 19$. Explain what happens to the denominator 5 in each case during the first step.

Q3203. Compare these two equations: (A) $\frac{x+2}{5} = 24$ vs (B) $\frac{x}{5} + 2 = 24$. Explain what happens to the denominator 5 in each case during the first step.

Q3204. Compare these two equations: (A) $\frac{x+6}{6} = 11$ vs (B) $\frac{x}{6} + 6 = 11$. Explain what happens to the denominator 6 in each case during the first step.

Q3205. Compare these two equations: (A) $\frac{x+5}{4} = 27$ vs (B) $\frac{x}{4} + 5 = 27$. Explain what happens to the denominator 4 in each case during the first step.

Q3206. Compare these two equations: (A) $\frac{x+5}{6} = 15$ vs (B) $\frac{x}{6} + 5 = 15$. Explain what happens to the denominator 6 in each case during the first step.

Q3207. Compare these two equations: (A) $\frac{x+7}{6} = 23$ vs (B) $\frac{x}{6} + 7 = 23$. Explain what happens to the denominator 6 in each case during the first step.

Q3208. Compare these two equations: (A) $\frac{x+2}{2} = 10$ vs (B) $\frac{x}{2} + 2 = 10$. Explain what happens to the denominator 2 in each case during the first step.

Q3209. Compare these two equations: (A) $\frac{x+4}{3} = 28$ vs (B) $\frac{x}{3} + 4 = 28$. Explain what happens to the denominator 3 in each case during the first step.

Q3210. Compare these two equations: (A) $\frac{x+4}{2} = 23$ vs (B) $\frac{x}{2} + 4 = 23$. Explain what happens to the denominator 2 in each case during the first step.

Q3211. Compare these two equations: (A) $\frac{x+8}{5} = 12$ vs (B) $\frac{x}{5} + 8 = 12$. Explain what happens to the denominator 5 in each case during the first step.

Q3212. Compare these two equations: (A) $\frac{x+7}{6} = 16$ vs (B) $\frac{x}{6} + 7 = 16$. Explain what happens to the denominator 6 in each case during the first step.

Q3213. Compare these two equations: (A) $\frac{x+5}{2} = 17$ vs (B) $\frac{x}{2} + 5 = 17$. Explain what happens to the denominator 2 in each case during the first step.

Q3214. Compare these two equations: (A) $\frac{x+4}{7} = 22$ vs (B) $\frac{x}{7} + 4 = 22$. Explain what happens to the denominator 7 in each case during the first step.

Q3215. Compare these two equations: (A) $\frac{x+5}{7} = 14$ vs (B) $\frac{x}{7} + 5 = 14$. Explain what happens to the denominator 7 in each case during the first step.

Q3216. Compare these two equations: (A) $\frac{x+6}{2} = 11$ vs (B) $\frac{x}{2} + 6 = 11$. Explain what happens to the denominator 2 in each case during the first step.

Q3217. Compare these two equations: (A) $\frac{x+9}{3} = 18$ vs (B) $\frac{x}{3} + 9 = 18$. Explain what happens to the denominator 3 in each case during the first step.

Q3218. Compare these two equations: (A) $\frac{x+9}{4} = 30$ vs (B) $\frac{x}{4} + 9 = 30$. Explain what happens to the denominator 4 in each case during the first step.

Q3219. Compare these two equations: (A) $\frac{x+9}{6} = 26$ vs (B) $\frac{x}{6} + 9 = 26$. Explain what happens to the denominator 6 in each case during the first step.

Q3220. Compare these two equations: (A) $\frac{x+2}{2} = 30$ vs (B) $\frac{x}{2} + 2 = 30$. Explain what happens to the denominator 2 in each case during the first step.

Q3221. Compare these two equations: (A) $\frac{x+6}{2} = 12$ vs (B) $\frac{x}{2} + 6 = 12$. Explain what happens to the denominator 2 in each case during the first step.

Q3222. Compare these two equations: (A) $\frac{x+9}{6} = 13$ vs (B) $\frac{x}{6} + 9 = 13$. Explain what happens to the denominator 6 in each case during the first step.

Q3223. Compare these two equations: (A) $\frac{x+3}{5} = 22$ vs (B) $\frac{x}{5} + 3 = 22$. Explain what happens to the denominator 5 in each case during the first step.

Q3224. Compare these two equations: (A) $\frac{x+4}{2} = 18$ vs (B) $\frac{x}{2} + 4 = 18$. Explain what happens to the denominator 2 in each case during the first step.

Q3225. Compare these two equations: (A) $\frac{x+8}{6} = 22$ vs (B) $\frac{x}{6} + 8 = 22$. Explain what happens to the denominator 6 in each case during the first step.

Q3226. Compare these two equations: (A) $\frac{x+4}{4} = 30$ vs (B) $\frac{x}{4} + 4 = 30$. Explain what happens to the denominator 4 in each case during the first step.

Q3227. Compare these two equations: (A) $\frac{x+8}{2} = 12$ vs (B) $\frac{x}{2} + 8 = 12$. Explain what happens to the denominator 2 in each case during the first step.

Q3228. Compare these two equations: (A) $\frac{x+6}{3} = 24$ vs (B) $\frac{x}{3} + 6 = 24$. Explain what happens to the denominator 3 in each case during the first step.

Q3229. Compare these two equations: (A) $\frac{x+2}{5} = 28$ vs (B) $\frac{x}{5} + 2 = 28$. Explain what happens to the denominator 5 in each case during the first step.

Q3230. Compare these two equations: (A) $\frac{x+4}{2} = 25$ vs

(B) $\frac{x}{2}+4=25$. Explain what happens to the denominator 2 in each case during the first step.

Q3231. Compare these two equations: (A) $\frac{x+5}{6}=29$ vs (B) $\frac{x}{6}+5=29$. Explain what happens to the denominator 6 in each case during the first step.

Q3232. Compare these two equations: (A) $\frac{x+9}{5}=30$ vs (B) $\frac{x}{5}+9=30$. Explain what happens to the denominator 5 in each case during the first step.

Q3233. Compare these two equations: (A) $\frac{x+8}{7}=21$ vs (B) $\frac{x}{7}+8=21$. Explain what happens to the denominator 7 in each case during the first step.

Q3234. Compare these two equations: (A) $\frac{x+7}{4}=14$ vs (B) $\frac{x}{4}+7=14$. Explain what happens to the denominator 4 in each case during the first step.

Q3235. Compare these two equations: (A) $\frac{x+7}{6}=29$ vs (B) $\frac{x}{6}+7=29$. Explain what happens to the denominator 6 in each case during the first step.

Q3236. Compare these two equations: (A) $\frac{x+8}{3}=26$ vs (B) $\frac{x}{3}+8=26$. Explain what happens to the denominator 3 in each case during the first step.

Q3237. Compare these two equations: (A) $\frac{x+7}{7}=21$ vs (B) $\frac{x}{7}+7=21$. Explain what happens to the denominator 7 in each case during the first step.

Q3238. Compare these two equations: (A) $\frac{x+4}{2}=12$ vs (B) $\frac{x}{2}+4=12$. Explain what happens to the denominator 2 in each case during the first step.

Q3239. Compare these two equations: (A) $\frac{x+8}{4}=11$ vs (B) $\frac{x}{4}+8=11$. Explain what happens to the denominator 4 in each case during the first step.

Q3240. Compare these two equations: (A) $\frac{x+6}{4}=16$ vs (B) $\frac{x}{4}+6=16$. Explain what happens to the denominator 4 in each case during the first step.

Q3241. Compare these two equations: (A) $\frac{x+4}{6}=24$ vs (B) $\frac{x}{6}+4=24$. Explain what happens to the denominator 6 in each case during the first step.

Q3242. Compare these two equations: (A) $\frac{x+4}{3}=10$ vs (B) $\frac{x}{3}+4=10$. Explain what happens to the denominator 3 in each case during the first step.

Q3243. Compare these two equations: (A) $\frac{x+3}{4}=29$ vs (B) $\frac{x}{4}+3=29$. Explain what happens to the denominator 4 in each case during the first step.

Q3244. Compare these two equations: (A) $\frac{x+8}{5}=12$ vs (B) $\frac{x}{5}+8=12$. Explain what happens to the denominator 5 in each case during the first step.

Q3245. Compare these two equations: (A) $\frac{x+5}{4}=20$ vs (B) $\frac{x}{4}+5=20$. Explain what happens to the denominator 4 in each case during the first step.

Q3246. Compare these two equations: (A) $\frac{x+2}{5}=22$ vs (B) $\frac{x}{5}+2=22$. Explain what happens to the denominator 5 in each case during the first step.

Q3247. Compare these two equations: (A) $\frac{x+3}{2}=17$ vs (B) $\frac{x}{2}+3=17$. Explain what happens to the denominator 2 in each case during the first step.

Q3248. Compare these two equations: (A) $\frac{x+4}{7}=19$ vs (B) $\frac{x}{7}+4=19$. Explain what happens to the denominator 7 in each case during the first step.

Q3249. Compare these two equations: (A) $\frac{x+9}{2}=13$ vs (B) $\frac{x}{2}+9=13$. Explain what happens to the denominator 2 in each case during the first step.

Q3250. Compare these two equations: (A) $\frac{x+3}{6}=10$ vs (B) $\frac{x}{6}+3=10$. Explain what happens to the denominator 6 in each case during the first step.

Q3251. Solve by cross-multiplication: $\frac{3x+9}{5x-5}=\frac{2}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3252. Solve by cross-multiplication: $\frac{2x+7}{5x-8}=\frac{2}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3253. Solve by cross-multiplication: $\frac{4x+7}{2x-2}=\frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3254. Solve by cross-multiplication: $\frac{2x+8}{5x-5}=\frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3255. Solve by cross-multiplication: $\frac{3x+9}{5x-3}=\frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3256. Solve by cross-multiplication: $\frac{3x+9}{5x-2}=\frac{4}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3257. Solve by cross-multiplication: $\frac{3x+2}{5x-5}=\frac{3}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3258. Solve by cross-multiplication: $\frac{4x+7}{4x-7}=\frac{2}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3259. Solve by cross-multiplication: $\frac{5x+4}{5x-9}=\frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3260. Solve by cross-multiplication: $\frac{5x+3}{2x-3}=\frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3261. Solve by cross-multiplication: $\frac{5x+3}{2x-3}=\frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3262. Solve by cross-multiplication: $\frac{4x+8}{4x-5}=\frac{3}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3263. Solve by cross-multiplication: $\frac{4x+5}{2x-4}=\frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3264. Solve by cross-multiplication: $\frac{5x+3}{3x-5}=\frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3265. Solve by cross-multiplication: $\frac{2x+2}{2x-5}=\frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3266. Solve by cross-multiplication: $\frac{5x+2}{5x-1}=\frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3267. Solve by cross-multiplication: $\frac{5x+9}{3x-2}=\frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3268. Solve by cross-multiplication: $\frac{4x+8}{2x-4}=\frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3269. Solve by cross-multiplication: $\frac{5x+3}{4x-6} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3270. Solve by cross-multiplication: $\frac{4x+7}{3x-7} = \frac{7}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3271. Solve by cross-multiplication: $\frac{2x+9}{2x-3} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3272. Solve by cross-multiplication: $\frac{5x+4}{5x-1} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3273. Solve by cross-multiplication: $\frac{3x+1}{3x-8} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3274. Solve by cross-multiplication: $\frac{4x+1}{5x-1} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3275. Solve by cross-multiplication: $\frac{2x+6}{5x-6} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3276. Solve by cross-multiplication: $\frac{2x+5}{4x-8} = \frac{6}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3277. Solve by cross-multiplication: $\frac{2x+2}{4x-3} = \frac{3}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3278. Solve by cross-multiplication: $\frac{4x+7}{3x-6} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3279. Solve by cross-multiplication: $\frac{4x+5}{2x-6} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3280. Solve by cross-multiplication: $\frac{4x+2}{2x-5} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3281. Solve by cross-multiplication: $\frac{2x+1}{5x-3} = \frac{2}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3282. Solve by cross-multiplication: $\frac{5x+7}{2x-3} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3283. Solve by cross-multiplication: $\frac{4x+7}{2x-9} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3284. Solve by cross-multiplication: $\frac{5x+5}{3x-3} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3285. Solve by cross-multiplication: $\frac{3x+1}{4x-7} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3286. Solve by cross-multiplication: $\frac{5x+7}{3x-2} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3287. Solve by cross-multiplication: $\frac{4x+8}{4x-3} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3288. Solve by cross-multiplication: $\frac{5x+7}{2x-7} = \frac{3}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3289. Solve by cross-multiplication: $\frac{2x+7}{5x-3} = \frac{4}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3290. Solve by cross-multiplication: $\frac{5x+7}{5x-4} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3291. Solve by cross-multiplication: $\frac{3x+4}{3x-9} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3292. Solve by cross-multiplication: $\frac{2x+3}{2x-2} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3293. Solve by cross-multiplication: $\frac{3x+6}{3x-9} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3294. Solve by cross-multiplication: $\frac{2x+9}{4x-8} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3295. Solve by cross-multiplication: $\frac{2x+2}{2x-2} = \frac{7}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3296. Solve by cross-multiplication: $\frac{3x+4}{5x-8} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3297. Solve by cross-multiplication: $\frac{2x+7}{3x-5} = \frac{2}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3298. Solve by cross-multiplication: $\frac{3x+5}{4x-3} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3299. Solve by cross-multiplication: $\frac{3x+5}{2x-1} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3300. Solve by cross-multiplication: $\frac{2x+1}{4x-1} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3301. Solve by cross-multiplication: $\frac{2x+7}{4x-7} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3302. Solve by cross-multiplication: $\frac{5x+1}{4x-1} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3303. Solve by cross-multiplication: $\frac{4x+1}{3x-1} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3304. Solve by cross-multiplication: $\frac{4x+7}{5x-2} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3305. Solve by cross-multiplication: $\frac{4x+5}{2x-4} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3306. Solve by cross-multiplication: $\frac{3x+3}{2x-9} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3307. Solve by cross-multiplication: $\frac{3x+6}{4x-3} = \frac{7}{2}$. Write out the distributed equation after denominators move

across the equals sign.

Q3308. Solve by cross-multiplication: $\frac{4x+4}{3x-1} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3309. Solve by cross-multiplication: $\frac{4x+2}{5x-4} = \frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3310. Solve by cross-multiplication: $\frac{5x+4}{5x-8} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3311. Solve by cross-multiplication: $\frac{3x+9}{3x-4} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3312. Solve by cross-multiplication: $\frac{4x+8}{4x-3} = \frac{5}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3313. Solve by cross-multiplication: $\frac{4x+1}{3x-4} = \frac{5}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3314. Solve by cross-multiplication: $\frac{2x+3}{2x-6} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3315. Solve by cross-multiplication: $\frac{2x+7}{2x-7} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3316. Solve by cross-multiplication: $\frac{5x+5}{2x-7} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3317. Solve by cross-multiplication: $\frac{5x+7}{4x-7} = \frac{5}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3318. Solve by cross-multiplication: $\frac{5x+3}{2x-9} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3319. Solve by cross-multiplication: $\frac{4x+7}{5x-1} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3320. Solve by cross-multiplication: $\frac{2x+3}{5x-9} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3321. Solve by cross-multiplication: $\frac{4x+3}{4x-7} = \frac{5}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3322. Solve by cross-multiplication: $\frac{5x+2}{5x-4} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3323. Solve by cross-multiplication: $\frac{5x+8}{5x-6} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3324. Solve by cross-multiplication: $\frac{5x+1}{3x-5} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3325. Solve by cross-multiplication: $\frac{3x+6}{4x-5} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3326. Solve by cross-multiplication: $\frac{3x+7}{2x-7} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3327. Solve by cross-multiplication: $\frac{5x+4}{4x-1} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3328. Solve by cross-multiplication: $\frac{2x+7}{5x-3} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3329. Solve by cross-multiplication: $\frac{2x+1}{5x-9} = \frac{3}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3330. Solve by cross-multiplication: $\frac{2x+1}{2x-3} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3331. Solve by cross-multiplication: $\frac{2x+3}{5x-7} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3332. Solve by cross-multiplication: $\frac{4x+4}{3x-4} = \frac{7}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3333. Solve by cross-multiplication: $\frac{4x+3}{2x-7} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3334. Solve by cross-multiplication: $\frac{3x+4}{3x-1} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3335. Solve by cross-multiplication: $\frac{3x+8}{2x-8} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3336. Solve by cross-multiplication: $\frac{5x+6}{4x-2} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3337. Solve by cross-multiplication: $\frac{4x+3}{4x-4} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3338. Solve by cross-multiplication: $\frac{4x+4}{4x-8} = \frac{4}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3339. Solve by cross-multiplication: $\frac{2x+7}{4x-5} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3340. Solve by cross-multiplication: $\frac{2x+5}{2x-3} = \frac{7}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3341. Solve by cross-multiplication: $\frac{4x+9}{4x-3} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3342. Solve by cross-multiplication: $\frac{3x+1}{3x-7} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3343. Solve by cross-multiplication: $\frac{2x+7}{5x-4} = \frac{6}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3344. Solve by cross-multiplication: $\frac{5x+5}{2x-6} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3345. Solve by cross-multiplication: $\frac{5x+7}{3x-9} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3346. Solve by cross-multiplication: $\frac{2x+1}{3x-6} = \frac{5}{2}$. Write

out the distributed equation after denominators move across the equals sign.

Q3347. Solve by cross-multiplication: $\frac{3x+6}{3x-5} = \frac{5}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3348. Solve by cross-multiplication: $\frac{4x+8}{2x-7} = \frac{4}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3349. Solve by cross-multiplication: $\frac{4x+4}{3x-9} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3350. Solve by cross-multiplication: $\frac{4x+4}{5x-3} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3351. Solve by cross-multiplication: $\frac{2x+4}{3x-3} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3352. Solve by cross-multiplication: $\frac{5x+9}{4x-1} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3353. Solve by cross-multiplication: $\frac{3x+3}{5x-5} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3354. Solve by cross-multiplication: $\frac{2x+5}{2x-7} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3355. Solve by cross-multiplication: $\frac{5x+6}{3x-3} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3356. Solve by cross-multiplication: $\frac{2x+8}{3x-7} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3357. Solve by cross-multiplication: $\frac{5x+2}{2x-4} = \frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3358. Solve by cross-multiplication: $\frac{3x+7}{3x-9} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3359. Solve by cross-multiplication: $\frac{3x+4}{5x-9} = \frac{3}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3360. Solve by cross-multiplication: $\frac{3x+9}{3x-7} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3361. Solve by cross-multiplication: $\frac{3x+4}{5x-2} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3362. Solve by cross-multiplication: $\frac{3x+5}{4x-9} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3363. Solve by cross-multiplication: $\frac{2x+1}{4x-9} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3364. Solve by cross-multiplication: $\frac{2x+7}{3x-7} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3365. Solve by cross-multiplication: $\frac{2x+2}{5x-7} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3366. Solve by cross-multiplication: $\frac{2x+7}{5x-3} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3367. Solve by cross-multiplication: $\frac{2x+1}{2x-1} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3368. Solve by cross-multiplication: $\frac{2x+2}{2x-7} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3369. Solve by cross-multiplication: $\frac{4x+7}{4x-4} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3370. Solve by cross-multiplication: $\frac{4x+8}{5x-7} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3371. Solve by cross-multiplication: $\frac{4x+6}{4x-8} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3372. Solve by cross-multiplication: $\frac{3x+1}{2x-5} = \frac{3}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3373. Solve by cross-multiplication: $\frac{5x+3}{5x-7} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3374. Solve by cross-multiplication: $\frac{5x+2}{3x-9} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3375. Solve by cross-multiplication: $\frac{3x+5}{2x-5} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3376. Solve by cross-multiplication: $\frac{3x+5}{3x-6} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3377. Solve by cross-multiplication: $\frac{2x+3}{2x-3} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3378. Solve by cross-multiplication: $\frac{3x+8}{4x-3} = \frac{4}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3379. Solve by cross-multiplication: $\frac{4x+8}{4x-7} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3380. Solve by cross-multiplication: $\frac{5x+7}{5x-7} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3381. Solve by cross-multiplication: $\frac{2x+5}{2x-6} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3382. Solve by cross-multiplication: $\frac{3x+8}{5x-7} = \frac{7}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3383. Solve by cross-multiplication: $\frac{5x+3}{5x-6} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3384. Solve by cross-multiplication: $\frac{3x+4}{2x-4} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3385. Solve by cross-multiplication: $\frac{5x+4}{5x-2} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3386. Solve by cross-multiplication: $\frac{2x+8}{2x-5} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3387. Solve by cross-multiplication: $\frac{3x+9}{3x-2} = \frac{5}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3388. Solve by cross-multiplication: $\frac{5x+5}{4x-2} = \frac{6}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3389. Solve by cross-multiplication: $\frac{4x+1}{2x-7} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3390. Solve by cross-multiplication: $\frac{4x+2}{4x-5} = \frac{3}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3391. Solve by cross-multiplication: $\frac{5x+1}{2x-3} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3392. Solve by cross-multiplication: $\frac{3x+5}{5x-5} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3393. Solve by cross-multiplication: $\frac{3x+5}{3x-6} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3394. Solve by cross-multiplication: $\frac{2x+7}{2x-7} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3395. Solve by cross-multiplication: $\frac{5x+7}{4x-6} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3396. Solve by cross-multiplication: $\frac{2x+5}{4x-6} = \frac{5}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3397. Solve by cross-multiplication: $\frac{5x+4}{2x-6} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3398. Solve by cross-multiplication: $\frac{2x+2}{5x-9} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3399. Solve by cross-multiplication: $\frac{3x+6}{2x-2} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3400. Solve by cross-multiplication: $\frac{2x+5}{3x-7} = \frac{3}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3401. Solve by cross-multiplication: $\frac{2x+1}{3x-2} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3402. Solve by cross-multiplication: $\frac{2x+3}{4x-2} = \frac{5}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3403. Solve by cross-multiplication: $\frac{3x+7}{5x-6} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3404. Solve by cross-multiplication: $\frac{3x+2}{5x-6} = \frac{3}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3405. Solve by cross-multiplication: $\frac{3x+1}{4x-3} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3406. Solve by cross-multiplication: $\frac{4x+3}{2x-3} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3407. Solve by cross-multiplication: $\frac{2x+2}{2x-8} = \frac{3}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3408. Solve by cross-multiplication: $\frac{4x+4}{4x-6} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3409. Solve by cross-multiplication: $\frac{5x+7}{4x-8} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3410. Solve by cross-multiplication: $\frac{2x+4}{4x-9} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3411. Solve by cross-multiplication: $\frac{2x+9}{4x-4} = \frac{7}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3412. Solve by cross-multiplication: $\frac{5x+1}{2x-4} = \frac{2}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3413. Solve by cross-multiplication: $\frac{4x+3}{4x-9} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3414. Solve by cross-multiplication: $\frac{5x+8}{4x-1} = \frac{3}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3415. Solve by cross-multiplication: $\frac{5x+3}{5x-7} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3416. Solve by cross-multiplication: $\frac{3x+2}{5x-1} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3417. Solve by cross-multiplication: $\frac{4x+6}{4x-7} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3418. Solve by cross-multiplication: $\frac{2x+1}{4x-9} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3419. Solve by cross-multiplication: $\frac{4x+6}{3x-4} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3420. Solve by cross-multiplication: $\frac{5x+6}{4x-8} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3421. Solve by cross-multiplication: $\frac{3x+3}{2x-4} = \frac{7}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3422. Solve by cross-multiplication: $\frac{2x+7}{2x-5} = \frac{2}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3423. Solve by cross-multiplication: $\frac{4x+2}{4x-3} = \frac{7}{6}$. Write out the distributed equation after denominators move

across the equals sign.

Q3424. Solve by cross-multiplication: $\frac{3x+9}{2x-5} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3425. Solve by cross-multiplication: $\frac{3x+1}{3x-9} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3426. Solve by cross-multiplication: $\frac{4x+7}{3x-1} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3427. Solve by cross-multiplication: $\frac{2x+1}{5x-3} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3428. Solve by cross-multiplication: $\frac{5x+2}{3x-9} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3429. Solve by cross-multiplication: $\frac{3x+8}{3x-6} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3430. Solve by cross-multiplication: $\frac{5x+6}{3x-9} = \frac{4}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3431. Solve by cross-multiplication: $\frac{5x+5}{2x-5} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3432. Solve by cross-multiplication: $\frac{3x+4}{2x-9} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3433. Solve by cross-multiplication: $\frac{2x+2}{4x-6} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3434. Solve by cross-multiplication: $\frac{4x+2}{2x-6} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3435. Solve by cross-multiplication: $\frac{5x+4}{5x-7} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3436. Solve by cross-multiplication: $\frac{5x+2}{2x-4} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3437. Solve by cross-multiplication: $\frac{2x+2}{2x-9} = \frac{5}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3438. Solve by cross-multiplication: $\frac{5x+8}{3x-4} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3439. Solve by cross-multiplication: $\frac{3x+6}{3x-1} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3440. Solve by cross-multiplication: $\frac{4x+5}{3x-8} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3441. Solve by cross-multiplication: $\frac{5x+5}{5x-6} = \frac{7}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3442. Solve by cross-multiplication: $\frac{3x+8}{5x-5} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3443. Solve by cross-multiplication: $\frac{4x+5}{3x-3} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3444. Solve by cross-multiplication: $\frac{4x+9}{4x-4} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3445. Solve by cross-multiplication: $\frac{4x+7}{3x-3} = \frac{7}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3446. Solve by cross-multiplication: $\frac{3x+6}{4x-4} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3447. Solve by cross-multiplication: $\frac{3x+9}{3x-4} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3448. Solve by cross-multiplication: $\frac{2x+6}{4x-4} = \frac{2}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3449. Solve by cross-multiplication: $\frac{2x+6}{3x-1} = \frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3450. Solve by cross-multiplication: $\frac{4x+7}{5x-2} = \frac{2}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3451. Solve by cross-multiplication: $\frac{3x+5}{3x-2} = \frac{6}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3452. Solve by cross-multiplication: $\frac{3x+8}{2x-3} = \frac{7}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3453. Solve by cross-multiplication: $\frac{5x+7}{2x-6} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3454. Solve by cross-multiplication: $\frac{4x+3}{5x-7} = \frac{4}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3455. Solve by cross-multiplication: $\frac{2x+1}{4x-9} = \frac{2}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3456. Solve by cross-multiplication: $\frac{5x+6}{4x-2} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3457. Solve by cross-multiplication: $\frac{3x+8}{2x-9} = \frac{5}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3458. Solve by cross-multiplication: $\frac{3x+6}{2x-1} = \frac{3}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3459. Solve by cross-multiplication: $\frac{3x+7}{4x-3} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3460. Solve by cross-multiplication: $\frac{2x+9}{5x-9} = \frac{3}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3461. Solve by cross-multiplication: $\frac{3x+3}{3x-6} = \frac{7}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3462. Solve by cross-multiplication: $\frac{3x+9}{3x-7} = \frac{2}{3}$. Write

out the distributed equation after denominators move across the equals sign.

Q3463. Solve by cross-multiplication: $\frac{3x+8}{4x-2} = \frac{2}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3464. Solve by cross-multiplication: $\frac{3x+6}{2x-6} = \frac{6}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3465. Solve by cross-multiplication: $\frac{5x+6}{3x-9} = \frac{2}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3466. Solve by cross-multiplication: $\frac{5x+3}{5x-9} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3467. Solve by cross-multiplication: $\frac{4x+7}{3x-7} = \frac{4}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3468. Solve by cross-multiplication: $\frac{2x+7}{2x-3} = \frac{2}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3469. Solve by cross-multiplication: $\frac{3x+1}{5x-9} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3470. Solve by cross-multiplication: $\frac{5x+2}{5x-9} = \frac{4}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3471. Solve by cross-multiplication: $\frac{5x+3}{2x-6} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3472. Solve by cross-multiplication: $\frac{5x+9}{5x-8} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3473. Solve by cross-multiplication: $\frac{4x+7}{3x-9} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3474. Solve by cross-multiplication: $\frac{5x+4}{4x-1} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3475. Solve by cross-multiplication: $\frac{3x+5}{5x-5} = \frac{4}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3476. Solve by cross-multiplication: $\frac{4x+4}{5x-4} = \frac{4}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3477. Solve by cross-multiplication: $\frac{5x+2}{3x-8} = \frac{3}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3478. Solve by cross-multiplication: $\frac{4x+3}{4x-9} = \frac{2}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3479. Solve by cross-multiplication: $\frac{4x+5}{5x-2} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3480. Solve by cross-multiplication: $\frac{5x+4}{4x-7} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3481. Solve by cross-multiplication: $\frac{3x+9}{4x-6} = \frac{3}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3482. Solve by cross-multiplication: $\frac{5x+5}{2x-1} = \frac{5}{3}$. Write out the distributed equation after denominators move across the equals sign.

Q3483. Solve by cross-multiplication: $\frac{2x+5}{2x-3} = \frac{6}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3484. Solve by cross-multiplication: $\frac{4x+8}{5x-6} = \frac{7}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3485. Solve by cross-multiplication: $\frac{5x+5}{2x-3} = \frac{7}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3486. Solve by cross-multiplication: $\frac{5x+6}{2x-4} = \frac{6}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3487. Solve by cross-multiplication: $\frac{4x+2}{5x-9} = \frac{7}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3488. Solve by cross-multiplication: $\frac{3x+7}{2x-3} = \frac{2}{2}$. Write out the distributed equation after denominators move across the equals sign.

Q3489. Solve by cross-multiplication: $\frac{4x+1}{2x-5} = \frac{2}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3490. Solve by cross-multiplication: $\frac{5x+2}{4x-8} = \frac{6}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3491. Solve by cross-multiplication: $\frac{3x+2}{5x-3} = \frac{5}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3492. Solve by cross-multiplication: $\frac{4x+9}{4x-2} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3493. Solve by cross-multiplication: $\frac{3x+8}{4x-7} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3494. Solve by cross-multiplication: $\frac{2x+7}{4x-4} = \frac{5}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3495. Solve by cross-multiplication: $\frac{4x+4}{3x-2} = \frac{7}{4}$. Write out the distributed equation after denominators move across the equals sign.

Q3496. Solve by cross-multiplication: $\frac{3x+1}{4x-2} = \frac{2}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3497. Solve by cross-multiplication: $\frac{4x+7}{5x-1} = \frac{7}{5}$. Write out the distributed equation after denominators move across the equals sign.

Q3498. Solve by cross-multiplication: $\frac{4x+1}{4x-7} = \frac{4}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3499. Solve by cross-multiplication: $\frac{3x+4}{3x-1} = \frac{3}{7}$. Write out the distributed equation after denominators move across the equals sign.

Q3500. Solve by cross-multiplication: $\frac{3x+3}{4x-8} = \frac{7}{6}$. Write out the distributed equation after denominators move across the equals sign.

Q3501. Solve $\frac{x}{5} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3502. Solve $\frac{x}{2} + \frac{x-1}{5} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3503. Solve $\frac{x}{5} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3504. Solve $\frac{x}{4} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3505. Solve $\frac{x}{4} + \frac{x-1}{5} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3506. Solve $\frac{x}{5} + \frac{x-1}{4} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3507. Solve $\frac{x}{3} + \frac{x-1}{4} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3508. Solve $\frac{x}{4} + \frac{x-1}{6} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3509. Solve $\frac{x}{5} + \frac{x-1}{3} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3510. Solve $\frac{x}{4} + \frac{x-1}{7} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3511. Solve $\frac{x}{3} + \frac{x-1}{4} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3512. Solve $\frac{x}{3} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3513. Solve $\frac{x}{5} + \frac{x-1}{6} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3514. Solve $\frac{x}{3} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3515. Solve $\frac{x}{2} + \frac{x-1}{4} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3516. Solve $\frac{x}{4} + \frac{x-1}{5} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3517. Solve $\frac{x}{2} + \frac{x-1}{6} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3518. Solve $\frac{x}{2} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3519. Solve $\frac{x}{5} + \frac{x-1}{3} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3520. Solve $\frac{x}{2} + \frac{x-1}{6} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3521. Solve $\frac{x}{4} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3522. Solve $\frac{x}{5} + \frac{x-1}{3} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3523. Solve $\frac{x}{4} + \frac{x-1}{7} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3524. Solve $\frac{x}{2} + \frac{x-1}{4} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3525. Solve $\frac{x}{3} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3526. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3527. Solve $\frac{x}{4} + \frac{x-1}{3} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3528. Solve $\frac{x}{5} + \frac{x-1}{7} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3529. Solve $\frac{x}{3} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3530. Solve $\frac{x}{5} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3531. Solve $\frac{x}{3} + \frac{x-1}{7} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3532. Solve $\frac{x}{2} + \frac{x-1}{7} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3533. Solve $\frac{x}{3} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3534. Solve $\frac{x}{3} + \frac{x-1}{6} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3535. Solve $\frac{x}{3} + \frac{x-1}{7} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3536. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3537. Solve $\frac{x}{3} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3538. Solve $\frac{x}{4} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3539. Solve $\frac{x}{2} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all

fractions.

Q3540. Solve $\frac{x}{5} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3541. Solve $\frac{x}{5} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3542. Solve $\frac{x}{4} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3543. Solve $\frac{x}{5} + \frac{x-1}{3} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3544. Solve $\frac{x}{4} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3545. Solve $\frac{x}{4} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3546. Solve $\frac{x}{4} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3547. Solve $\frac{x}{5} + \frac{x-1}{4} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3548. Solve $\frac{x}{5} + \frac{x-1}{3} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3549. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3550. Solve $\frac{x}{3} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3551. Solve $\frac{x}{5} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3552. Solve $\frac{x}{4} + \frac{x-1}{7} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3553. Solve $\frac{x}{4} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3554. Solve $\frac{x}{2} + \frac{x-1}{4} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3555. Solve $\frac{x}{4} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3556. Solve $\frac{x}{2} + \frac{x-1}{7} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3557. Solve $\frac{x}{4} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3558. Solve $\frac{x}{2} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3559. Solve $\frac{x}{5} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3560. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3561. Solve $\frac{x}{3} + \frac{x-1}{5} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3562. Solve $\frac{x}{2} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3563. Solve $\frac{x}{3} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3564. Solve $\frac{x}{4} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3565. Solve $\frac{x}{4} + \frac{x-1}{5} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3566. Solve $\frac{x}{3} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3567. Solve $\frac{x}{3} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3568. Solve $\frac{x}{2} + \frac{x-1}{4} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3569. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3570. Solve $\frac{x}{2} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3571. Solve $\frac{x}{5} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3572. Solve $\frac{x}{5} + \frac{x-1}{6} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3573. Solve $\frac{x}{2} + \frac{x-1}{7} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3574. Solve $\frac{x}{4} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3575. Solve $\frac{x}{2} + \frac{x-1}{7} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3576. Solve $\frac{x}{5} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3577. Solve $\frac{x}{2} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3578. Solve $\frac{x}{3} + \frac{x-1}{4} = 9$. Show how multiplying the

entire equation by the LCM of denominators eliminates all fractions.

Q3579. Solve $\frac{x}{5} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3580. Solve $\frac{x}{3} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3581. Solve $\frac{x}{3} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3582. Solve $\frac{x}{5} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3583. Solve $\frac{x}{5} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3584. Solve $\frac{x}{4} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3585. Solve $\frac{x}{4} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3586. Solve $\frac{x}{2} + \frac{x-1}{3} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3587. Solve $\frac{x}{2} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3588. Solve $\frac{x}{2} + \frac{x-1}{4} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3589. Solve $\frac{x}{5} + \frac{x-1}{6} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3590. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3591. Solve $\frac{x}{3} + \frac{x-1}{4} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3592. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3593. Solve $\frac{x}{4} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3594. Solve $\frac{x}{5} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3595. Solve $\frac{x}{3} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3596. Solve $\frac{x}{4} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3597. Solve $\frac{x}{5} + \frac{x-1}{3} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3598. Solve $\frac{x}{2} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3599. Solve $\frac{x}{2} + \frac{x-1}{3} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3600. Solve $\frac{x}{5} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3601. Solve $\frac{x}{5} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3602. Solve $\frac{x}{4} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3603. Solve $\frac{x}{4} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3604. Solve $\frac{x}{4} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3605. Solve $\frac{x}{3} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3606. Solve $\frac{x}{5} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3607. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3608. Solve $\frac{x}{2} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3609. Solve $\frac{x}{5} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3610. Solve $\frac{x}{3} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3611. Solve $\frac{x}{2} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3612. Solve $\frac{x}{5} + \frac{x-1}{3} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3613. Solve $\frac{x}{2} + \frac{x-1}{5} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3614. Solve $\frac{x}{4} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3615. Solve $\frac{x}{2} + \frac{x-1}{7} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3616. Solve $\frac{x}{3} + \frac{x-1}{7} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3617. Solve $\frac{x}{4} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3618. Solve $\frac{x}{4} + \frac{x-1}{3} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3619. Solve $\frac{x}{2} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3620. Solve $\frac{x}{2} + \frac{x-1}{5} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3621. Solve $\frac{x}{3} + \frac{x-1}{7} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3622. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3623. Solve $\frac{x}{4} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3624. Solve $\frac{x}{4} + \frac{x-1}{3} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3625. Solve $\frac{x}{2} + \frac{x-1}{5} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3626. Solve $\frac{x}{5} + \frac{x-1}{3} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3627. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3628. Solve $\frac{x}{5} + \frac{x-1}{3} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3629. Solve $\frac{x}{3} + \frac{x-1}{7} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3630. Solve $\frac{x}{3} + \frac{x-1}{4} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3631. Solve $\frac{x}{2} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3632. Solve $\frac{x}{4} + \frac{x-1}{5} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3633. Solve $\frac{x}{3} + \frac{x-1}{4} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3634. Solve $\frac{x}{5} + \frac{x-1}{6} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3635. Solve $\frac{x}{5} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3636. Solve $\frac{x}{4} + \frac{x-1}{3} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3637. Solve $\frac{x}{2} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3638. Solve $\frac{x}{3} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3639. Solve $\frac{x}{3} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3640. Solve $\frac{x}{4} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3641. Solve $\frac{x}{2} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3642. Solve $\frac{x}{3} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3643. Solve $\frac{x}{3} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3644. Solve $\frac{x}{3} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3645. Solve $\frac{x}{2} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3646. Solve $\frac{x}{2} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3647. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3648. Solve $\frac{x}{4} + \frac{x-1}{5} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3649. Solve $\frac{x}{3} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3650. Solve $\frac{x}{4} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3651. Solve $\frac{x}{2} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3652. Solve $\frac{x}{2} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3653. Solve $\frac{x}{5} + \frac{x-1}{3} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3654. Solve $\frac{x}{4} + \frac{x-1}{3} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3655. Solve $\frac{x}{4} + \frac{x-1}{3} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

fractions.

Q3656. Solve $\frac{x}{5} + \frac{x-1}{4} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3657. Solve $\frac{x}{2} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3658. Solve $\frac{x}{4} + \frac{x-1}{5} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3659. Solve $\frac{x}{3} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3660. Solve $\frac{x}{3} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3661. Solve $\frac{x}{3} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3662. Solve $\frac{x}{5} + \frac{x-1}{4} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3663. Solve $\frac{x}{3} + \frac{x-1}{4} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3664. Solve $\frac{x}{3} + \frac{x-1}{6} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3665. Solve $\frac{x}{2} + \frac{x-1}{3} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3666. Solve $\frac{x}{4} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3667. Solve $\frac{x}{3} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3668. Solve $\frac{x}{4} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3669. Solve $\frac{x}{4} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3670. Solve $\frac{x}{2} + \frac{x-1}{6} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3671. Solve $\frac{x}{3} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3672. Solve $\frac{x}{5} + \frac{x-1}{3} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3673. Solve $\frac{x}{4} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3674. Solve $\frac{x}{2} + \frac{x-1}{3} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3675. Solve $\frac{x}{4} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3676. Solve $\frac{x}{3} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3677. Solve $\frac{x}{4} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3678. Solve $\frac{x}{3} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3679. Solve $\frac{x}{5} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3680. Solve $\frac{x}{3} + \frac{x-1}{6} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3681. Solve $\frac{x}{4} + \frac{x-1}{5} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3682. Solve $\frac{x}{5} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3683. Solve $\frac{x}{5} + \frac{x-1}{7} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3684. Solve $\frac{x}{4} + \frac{x-1}{5} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3685. Solve $\frac{x}{4} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3686. Solve $\frac{x}{4} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3687. Solve $\frac{x}{2} + \frac{x-1}{3} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3688. Solve $\frac{x}{5} + \frac{x-1}{6} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3689. Solve $\frac{x}{4} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3690. Solve $\frac{x}{3} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3691. Solve $\frac{x}{5} + \frac{x-1}{6} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3692. Solve $\frac{x}{5} + \frac{x-1}{3} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3693. Solve $\frac{x}{3} + \frac{x-1}{4} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3694. Solve $\frac{x}{2} + \frac{x-1}{5} = 3$. Show how multiplying the

entire equation by the LCM of denominators eliminates all fractions.

Q3695. Solve $\frac{x}{3} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3696. Solve $\frac{x}{2} + \frac{x-1}{3} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3697. Solve $\frac{x}{3} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3698. Solve $\frac{x}{3} + \frac{x-1}{4} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3699. Solve $\frac{x}{3} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3700. Solve $\frac{x}{5} + \frac{x-1}{3} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3701. Solve $\frac{x}{4} + \frac{x-1}{6} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3702. Solve $\frac{x}{4} + \frac{x-1}{5} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3703. Solve $\frac{x}{5} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3704. Solve $\frac{x}{4} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3705. Solve $\frac{x}{5} + \frac{x-1}{7} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3706. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3707. Solve $\frac{x}{2} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3708. Solve $\frac{x}{4} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3709. Solve $\frac{x}{5} + \frac{x-1}{3} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3710. Solve $\frac{x}{4} + \frac{x-1}{7} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3711. Solve $\frac{x}{2} + \frac{x-1}{4} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3712. Solve $\frac{x}{3} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3713. Solve $\frac{x}{2} + \frac{x-1}{5} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3714. Solve $\frac{x}{2} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3715. Solve $\frac{x}{2} + \frac{x-1}{3} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3716. Solve $\frac{x}{3} + \frac{x-1}{4} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3717. Solve $\frac{x}{5} + \frac{x-1}{3} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3718. Solve $\frac{x}{5} + \frac{x-1}{3} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3719. Solve $\frac{x}{5} + \frac{x-1}{7} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3720. Solve $\frac{x}{4} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3721. Solve $\frac{x}{4} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3722. Solve $\frac{x}{5} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3723. Solve $\frac{x}{2} + \frac{x-1}{7} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3724. Solve $\frac{x}{2} + \frac{x-1}{4} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3725. Solve $\frac{x}{5} + \frac{x-1}{4} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3726. Solve $\frac{x}{2} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3727. Solve $\frac{x}{5} + \frac{x-1}{6} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3728. Solve $\frac{x}{2} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3729. Solve $\frac{x}{4} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3730. Solve $\frac{x}{5} + \frac{x-1}{3} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3731. Solve $\frac{x}{5} + \frac{x-1}{7} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3732. Solve $\frac{x}{3} + \frac{x-1}{6} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3733. Solve $\frac{x}{4} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3734. Solve $\frac{x}{5} + \frac{x-1}{6} = 8$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3735. Solve $\frac{x}{4} + \frac{x-1}{5} = 3$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3736. Solve $\frac{x}{3} + \frac{x-1}{4} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3737. Solve $\frac{x}{3} + \frac{x-1}{7} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3738. Solve $\frac{x}{4} + \frac{x-1}{3} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3739. Solve $\frac{x}{5} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3740. Solve $\frac{x}{4} + \frac{x-1}{5} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3741. Solve $\frac{x}{5} + \frac{x-1}{6} = 2$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3742. Solve $\frac{x}{2} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3743. Solve $\frac{x}{2} + \frac{x-1}{6} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3744. Solve $\frac{x}{3} + \frac{x-1}{5} = 7$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3745. Solve $\frac{x}{4} + \frac{x-1}{7} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3746. Solve $\frac{x}{3} + \frac{x-1}{4} = 4$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3747. Solve $\frac{x}{4} + \frac{x-1}{5} = 9$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3748. Solve $\frac{x}{3} + \frac{x-1}{7} = 6$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3749. Solve $\frac{x}{4} + \frac{x-1}{3} = 5$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3750. Solve $\frac{x}{4} + \frac{x-1}{7} = 10$. Show how multiplying the entire equation by the LCM of denominators eliminates all fractions.

Q3751. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3752. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3753. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3754. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3755. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3756. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3757. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3758. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3759. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3760. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3761. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3762. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3763. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3764. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3765. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3766. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3767. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3768. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3769. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3770. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3771. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3772. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3773. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3774. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3775. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3776. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3777. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3778. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3779. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3780. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3781. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3782. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3783. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3784. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3785. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3786. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3787. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3788. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3789. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3790. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3791. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3792. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3793. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3794. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3795. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3796. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3797. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3798. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3799. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3800. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3801. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3802. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3803. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3804. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3805. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3806. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3807. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3808. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain

restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3809. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3810. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3811. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3812. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3813. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3814. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3815. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3816. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3817. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3818. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3819. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3820. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3821. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3822. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3823. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3824. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3825. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3826. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3827. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3828. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3829. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3830. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3831. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3832. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3833. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3834. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3835. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3836. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3837. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3838. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3839. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3840. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3841. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3842. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3843. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3844. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3845. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even

if algebraic simplification suggests it?

Q3846. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3847. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3848. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3849. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3850. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3851. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3852. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3853. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3854. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3855. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3856. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3857. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3858. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3859. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3860. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3861. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3862. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3863. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3864. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3865. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3866. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3867. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3868. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3869. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3870. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3871. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3872. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3873. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3874. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3875. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3876. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3877. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3878. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3879. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3880. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3881. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3882. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3883. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3884. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3885. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3886. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3887. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3888. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3889. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3890. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3891. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3892. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3893. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3894. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3895. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3896. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3897. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3898. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3899. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3900. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3901. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3902. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3903. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3904. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3905. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3906. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3907. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3908. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3909. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3910. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3911. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3912. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3913. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3914. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3915. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3916. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3917. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3918. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3919. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3920. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain

restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3921. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3922. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3923. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3924. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3925. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3926. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3927. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3928. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3929. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3930. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3931. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3932. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3933. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3934. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3935. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3936. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3937. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3938. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3939. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3940. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3941. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3942. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3943. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3944. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3945. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3946. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3947. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3948. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3949. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3950. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3951. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3952. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3953. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3954. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3955. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3956. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3957. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even

if algebraic simplification suggests it?

Q3958. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3959. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3960. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3961. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3962. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3963. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3964. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3965. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3966. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3967. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3968. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3969. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3970. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3971. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3972. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3973. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3974. For the equation $\frac{x^2-4}{x-2} = 22$, what is the domain restriction on x? Why is $x = 2$ NOT a valid solution even if algebraic simplification suggests it?

Q3975. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3976. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3977. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3978. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3979. For the equation $\frac{x^2-25}{x-5} = 25$, what is the domain restriction on x? Why is $x = 5$ NOT a valid solution even if algebraic simplification suggests it?

Q3980. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3981. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3982. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3983. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3984. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3985. For the equation $\frac{x^2-49}{x-7} = 27$, what is the domain restriction on x? Why is $x = 7$ NOT a valid solution even if algebraic simplification suggests it?

Q3986. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3987. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3988. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3989. For the equation $\frac{x^2-36}{x-6} = 26$, what is the domain restriction on x? Why is $x = 6$ NOT a valid solution even if algebraic simplification suggests it?

Q3990. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3991. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3992. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3993. For the equation $\frac{x^2-9}{x-3} = 23$, what is the domain restriction on x? Why is $x = 3$ NOT a valid solution even if algebraic simplification suggests it?

Q3994. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q3995. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3996. For the equation $\frac{x^2-1}{x-1} = 21$, what is the domain restriction on x? Why is $x = 1$ NOT a valid solution even if algebraic simplification suggests it?

Q3997. For the equation $\frac{x^2-81}{x-9} = 29$, what is the domain restriction on x? Why is $x = 9$ NOT a valid solution even if algebraic simplification suggests it?

Q3998. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

Q3999. For the equation $\frac{x^2-64}{x-8} = 28$, what is the domain restriction on x? Why is $x = 8$ NOT a valid solution even if algebraic simplification suggests it?

Q4000. For the equation $\frac{x^2-16}{x-4} = 24$, what is the domain restriction on x? Why is $x = 4$ NOT a valid solution even if algebraic simplification suggests it?

PART 2: STEP-BY-STEP ANSWER KEY

MODULE 1: SIGN CONVENTIONS AND TRANSPOSITION RULES (Q1 – Q1000) — Answer Key

Q1. Transposing -13 gives $11x = 28 + 13 = 41$. Dividing by 11 yields $x = \frac{41}{11}$. Simplifies to $x = 3.73$.

Q2. Transposing -3 gives $10x = -1 + 3 = 2$. Dividing by 10 yields $x = \frac{2}{10}$. Simplifies to $x = 0.20$.

Q3. Transposing $+17$ gives $2x = 2 - 17 = -15$. Dividing by 2 yields $x = \frac{-15}{2}$. Simplifies to $x = -7.50$.

Q4. Transposing $+13$ gives $5x = -16 - 13 = -29$. Dividing by 5 yields $x = \frac{-29}{5}$. Simplifies to $x = -5.80$.

Q5. Transposing $+6$ gives $6x = 14 - 6 = 8$. Dividing by 6 yields $x = \frac{8}{6}$. Simplifies to $x = 1.33$.

Q6. Transposing -12 gives $14x = -9 + 12 = 3$. Dividing by 14 yields $x = \frac{3}{14}$. Simplifies to $x = 0.21$.

Q7. Transposing -21 gives $9x = -14 + 21 = 7$. Dividing by 9 yields $x = \frac{7}{9}$. Simplifies to $x = 0.78$.

Q8. Transposing -15 gives $5x = -25 + 15 = -10$. Dividing by 5 yields $x = \frac{-10}{5}$. Simplifies to $x = -2.00$.

Q9. Transposing $+24$ gives $9x = 4 - 24 = -20$. Dividing by 9 yields $x = \frac{-20}{9}$. Simplifies to $x = -2.22$.

Q10. Transposing -13 gives $7x = 0 + 13 = 13$. Dividing by 7 yields $x = \frac{13}{7}$. Simplifies to $x = 1.86$.

Q11. Transposing -5 gives $14x = -15 + 5 = -10$. Dividing by 14 yields $x = \frac{-10}{14}$. Simplifies to $x = -0.71$.

Q12. Transposing $+25$ gives $12x = -5 - 25 = -30$. Dividing by 12 yields $x = \frac{-30}{12}$. Simplifies to $x = -2.50$.

Q13. Transposing -12 gives $8x = -26 + 12 = -14$. Dividing by 8 yields $x = \frac{-14}{8}$. Simplifies to $x = -1.75$.

Q14. Transposing $+4$ gives $8x = -11 - 4 = -15$. Dividing by 8 yields $x = \frac{-15}{8}$. Simplifies to $x = -1.88$.

Q15. Transposing -18 gives $5x = 11 + 18 = 29$. Dividing by 5 yields $x = \frac{29}{5}$. Simplifies to $x = 5.80$.

Q16. Transposing $+14$ gives $15x = 13 - 14 = -1$. Dividing by 15 yields $x = \frac{-1}{15}$. Simplifies to $x = -0.07$.

Q17. Transposing $+17$ gives $7x = 24 - 17 = 7$. Dividing by 7 yields $x = \frac{7}{7}$. Simplifies to $x = 1.00$.

Q18. Transposing $+7$ gives $6x = 20 - 7 = 13$. Dividing by 6 yields $x = \frac{13}{6}$. Simplifies to $x = 2.17$.

Q19. Transposing -11 gives $8x = -3 + 11 = 8$. Dividing by 8 yields $x = \frac{8}{8}$. Simplifies to $x = 1.00$.

Q20. Transposing $+7$ gives $3x = 17 - 7 = 10$. Dividing by 3 yields $x = \frac{10}{3}$. Simplifies to $x = 3.33$.

Q21. Transposing $+1$ gives $12x = -30 - 1 = -31$. Dividing by 12 yields $x = \frac{-31}{12}$. Simplifies to $x = -2.58$.

Q22. Transposing $+4$ gives $15x = -17 - 4 = -21$. Dividing by 15 yields $x = \frac{-21}{15}$. Simplifies to $x = -1.40$.

Q23. Transposing -19 gives $10x = -8 + 19 = 11$. Dividing by 10 yields $x = \frac{11}{10}$. Simplifies to $x = 1.10$.

Q24. Transposing -24 gives $14x = -7 + 24 = 17$. Dividing by 14 yields $x = \frac{17}{14}$. Simplifies to $x = 1.21$.

Q25. Transposing -2 gives $13x = -30 + 2 = -28$. Dividing by 13 yields $x = \frac{-28}{13}$. Simplifies to $x = -2.15$.

Q26. Transposing $+11$ gives $6x = 22 - 11 = 11$. Dividing by 6 yields $x = \frac{11}{6}$. Simplifies to $x = 1.83$.

Q27. Transposing -23 gives $4x = -3 + 23 = 20$. Dividing by 4 yields $x = \frac{20}{4}$. Simplifies to $x = 5.00$.

Q28. Transposing -8 gives $4x = -26 + 8 = -18$. Dividing by 4 yields $x = \frac{-18}{4}$. Simplifies to $x = -4.50$.

Q29. Transposing -16 gives $14x = -5 + 16 = 11$. Dividing by 14 yields $x = \frac{11}{14}$. Simplifies to $x = 0.79$.

Q30. Transposing $+21$ gives $12x = 22 - 21 = 1$. Dividing by 12 yields $x = \frac{1}{12}$. Simplifies to $x = 0.08$.

Q31. Transposing $+16$ gives $7x = 6 - 16 = -10$. Dividing by 7 yields $x = \frac{-10}{7}$. Simplifies to $x = -1.43$.

Q32. Transposing -15 gives $9x = 1 + 15 = 16$. Dividing by 9 yields $x = \frac{16}{9}$. Simplifies to $x = 1.78$.

Q33. Transposing $+7$ gives $15x = -11 - 7 = -18$. Dividing by 15 yields $x = \frac{-18}{15}$. Simplifies to $x = -1.20$.

Q34. Transposing -14 gives $15x = 14 + 14 = 28$. Dividing by 15 yields $x = \frac{28}{15}$. Simplifies to $x = 1.87$.

Q35. Transposing $+18$ gives $10x = 25 - 18 = 7$. Dividing by 10 yields $x = \frac{7}{10}$. Simplifies to $x = 0.70$.

Q36. Transposing -20 gives $2x = -26 + 20 = -6$. Dividing by 2 yields $x = \frac{-6}{2}$. Simplifies to $x = -3.00$.

Q37. Transposing $+11$ gives $2x = -8 - 11 = -19$. Dividing by 2 yields $x = \frac{-19}{2}$. Simplifies to $x = -9.50$.

Q38. Transposing $+19$ gives $10x = 26 - 19 = 7$. Dividing by 10 yields $x = \frac{7}{10}$. Simplifies to $x = 0.70$.

Q39. Transposing -18 gives $9x = 13 + 18 = 31$. Dividing by 9 yields $x = \frac{31}{9}$. Simplifies to $x = 3.44$.

Q40. Transposing $+14$ gives $13x = 19 - 14 = 5$. Dividing by 13 yields $x = \frac{5}{13}$. Simplifies to $x = 0.38$.

Q41. Transposing $+4$ gives $11x = 20 - 4 = 16$. Dividing by 11 yields $x = \frac{16}{11}$. Simplifies to $x = 1.45$.

Q42. Transposing -17 gives $12x = 13 + 17 = 30$. Dividing by 12 yields $x = \frac{30}{12}$. Simplifies to $x = 2.50$.

Q43. Transposing $+25$ gives $4x = -18 - 25 = -43$. Dividing by 4 yields $x = \frac{-43}{4}$. Simplifies to $x = -10.75$.

Q44. Transposing $+10$ gives $13x = -2 - 10 = -12$. Dividing by 13 yields $x = \frac{-12}{13}$. Simplifies to $x = -0.92$.

Q45. Transposing -20 gives $4x = -21 + 20 = -1$. Dividing by 4 yields $x = \frac{-1}{4}$. Simplifies to $x = -0.25$.

Q46. Transposing -4 gives $6x = -6 + 4 = -2$. Dividing by 6 yields $x = \frac{-2}{6}$. Simplifies to $x = -0.33$.

Q47. Transposing -22 gives $3x = 5 + 22 = 27$. Dividing by 3 yields $x = \frac{27}{3}$. Simplifies to $x = 9.00$.

Q48. Transposing $+19$ gives $10x = -8 - 19 = -27$. Dividing by 10 yields $x = \frac{-27}{10}$. Simplifies to $x = -2.70$.

Q49. Transposing -22 gives $9x = -18 + 22 = 4$. Dividing by 9 yields $x = \frac{4}{9}$. Simplifies to $x = 0.44$.

Q50. Transposing $+18$ gives $6x = 22 - 18 = 4$. Dividing by 6 yields $x = \frac{4}{6}$. Simplifies to $x = 0.67$.

Q51. Transposing -20 gives $7x = -13 + 20 = 7$. Dividing by 7 yields $x = \frac{7}{7}$. Simplifies to $x = 1.00$.

Q52. Transposing -9 gives $4x = -1 + 9 = 8$. Dividing by 4 yields $x = \frac{8}{4}$. Simplifies to $x = 2.00$.

Q53. Transposing $+2$ gives $5x = -23 - 2 = -25$. Dividing by 5 yields $x = \frac{-25}{5}$. Simplifies to $x = -5.00$.

Q54. Transposing -5 gives $10x = -15 + 5 = -10$. Dividing by 10 yields $x = \frac{-10}{10}$. Simplifies to $x = -1.00$.

Q55. Transposing -8 gives $3x = 11 + 8 = 19$. Dividing by 3 yields $x = \frac{19}{3}$. Simplifies to $x = 6.33$.

Q56. Transposing -25 gives $4x = -30 + 25 = -5$. Dividing by 4 yields $x = \frac{-5}{4}$. Simplifies to $x = -1.25$.

Q57. Transposing $+7$ gives $12x = 24 - 7 = 17$. Dividing by 12 yields $x = \frac{17}{12}$. Simplifies to $x = 1.42$.

Q58. Transposing $+4$ gives $11x = 11 - 4 = 7$. Dividing by 11 yields $x = \frac{7}{11}$. Simplifies to $x = 0.64$.

Q59. Transposing -23 gives $11x = 21 + 23 = 44$. Dividing by 11 yields $x = \frac{44}{11}$. Simplifies to $x = 4.00$.

Q60. Transposing -10 gives $7x = 20 + 10 = 30$. Dividing by 7 yields $x = \frac{30}{7}$. Simplifies to $x = 4.29$.

Q61. Transposing $+5$ gives $6x = 12 - 5 = 7$. Dividing by 6 yields $x = \frac{7}{6}$. Simplifies to $x = 1.17$.

Q62. Transposing $+10$ gives $5x = 7 - 10 = -3$. Dividing by 5 yields $x = \frac{-3}{5}$. Simplifies to $x = -0.60$.

Q63. Transposing -5 gives $10x = -9 + 5 = -4$. Dividing by 10 yields $x = \frac{-4}{10}$. Simplifies to $x = -0.40$.

Q64. Transposing -17 gives $2x = -12 + 17 = 5$. Dividing by 2 yields $x = \frac{5}{2}$. Simplifies to $x = 2.50$.

Q65. Transposing $+18$ gives $7x = -7 - 18 = -25$. Dividing by 7 yields $x = \frac{-25}{7}$. Simplifies to $x = -3.57$.

Q66. Transposing $+7$ gives $8x = 10 - 7 = 3$. Dividing by 8 yields $x = \frac{3}{8}$. Simplifies to $x = 0.38$.

Q67. Transposing $+22$ gives $2x = -16 - 22 = -38$. Dividing by 2 yields $x = \frac{-38}{2}$. Simplifies to $x = -19.00$.

Q68. Transposing -21 gives $10x = -17 + 21 = 4$. Dividing by 10 yields $x = \frac{4}{10}$. Simplifies to $x = 0.40$.

Q69. Transposing $+17$ gives $4x = -23 - 17 = -40$. Dividing by 4 yields $x = \frac{-40}{4}$. Simplifies to $x = -10.00$.

Q70. Transposing -4 gives $4x = -23 + 4 = -19$. Dividing by 4 yields $x = \frac{-19}{4}$. Simplifies to $x = -4.75$.

Q71. Transposing -24 gives $7x = -2 + 24 = 22$. Dividing by 7 yields $x = \frac{22}{7}$. Simplifies to $x = 3.14$.

Q72. Transposing $+17$ gives $15x = -18 - 17 = -35$. Dividing by 15 yields $x = \frac{-35}{15}$. Simplifies to $x = -2.33$.

Q73. Transposing $+17$ gives $12x = 13 - 17 = -4$. Dividing by 12 yields $x = \frac{-4}{12}$. Simplifies to $x = -0.33$.

Q74. Transposing -2 gives $15x = 2 + 2 = 4$. Dividing by 15 yields $x = \frac{4}{15}$. Simplifies to $x = 0.27$.

Q75. Transposing $+15$ gives $12x = 5 - 15 = -10$. Dividing by 12 yields $x = \frac{-10}{12}$. Simplifies to $x = -0.83$.

Q76. Transposing -17 gives $5x = -1 + 17 = 16$. Dividing by 5 yields $x = \frac{16}{5}$. Simplifies to $x = 3.20$.

Q77. Transposing $+10$ gives $10x = -10 - 10 = -20$. Dividing by 10 yields $x = \frac{-20}{10}$. Simplifies to $x = -2.00$.

Q78. Transposing $+18$ gives $6x = -11 - 18 = -29$. Dividing by 6 yields $x = \frac{-29}{6}$. Simplifies to $x = -4.83$.

Q79. Transposing -9 gives $13x = -15 + 9 = -6$. Dividing by 13 yields $x = \frac{-6}{13}$. Simplifies to $x = -0.46$.

Q80. Transposing -12 gives $8x = -1 + 12 = 11$. Dividing by 8 yields $x = \frac{11}{8}$. Simplifies to $x = 1.38$.

Q81. Transposing $+23$ gives $6x = -28 - 23 = -51$. Dividing by 6 yields $x = \frac{-51}{6}$. Simplifies to $x = -8.50$.

Q82. Transposing -22 gives $5x = 7 + 22 = 29$. Dividing by 5 yields $x = \frac{29}{5}$. Simplifies to $x = 5.80$.

Q83. Transposing $+22$ gives $7x = 26 - 22 = 4$. Dividing by 7 yields $x = \frac{4}{7}$. Simplifies to $x = 0.57$.

Q84. Transposing $+3$ gives $9x = 12 - 3 = 9$. Dividing by 9 yields $x = \frac{9}{9}$. Simplifies to $x = 1.00$.

Q85. Transposing -8 gives $2x = 5 + 8 = 13$. Dividing by 2 yields $x = \frac{13}{2}$. Simplifies to $x = 6.50$.

Q86. Transposing -12 gives $13x = 30 + 12 = 42$. Dividing by 13 yields $x = \frac{42}{13}$. Simplifies to $x = 3.23$.

Q87. Transposing $+24$ gives $2x = 27 - 24 = 3$. Dividing by 2 yields $x = \frac{3}{2}$. Simplifies to $x = 1.50$.

Q88. Transposing $+14$ gives $9x = 29 - 14 = 15$. Dividing by 9 yields $x = \frac{15}{9}$. Simplifies to $x = 1.67$.

Q89. Transposing -22 gives $15x = -30 + 22 = -8$. Dividing by 15 yields $x = \frac{-8}{15}$. Simplifies to $x = -0.53$.

Q90. Transposing $+3$ gives $8x = 12 - 3 = 9$. Dividing by 8 yields $x = \frac{9}{8}$. Simplifies to $x = 1.12$.

Q91. Transposing $+24$ gives $6x = 9 - 24 = -15$. Dividing by 6 yields $x = \frac{-15}{6}$. Simplifies to $x = -2.50$.

Q92. Transposing $+6$ gives $4x = 17 - 6 = 11$. Dividing by 4 yields $x = \frac{11}{4}$. Simplifies to $x = 2.75$.

Q93. Transposing $+15$ gives $14x = -27 - 15 = -42$. Dividing by 14 yields $x = \frac{-42}{14}$. Simplifies to $x = -3.00$.

Q94. Transposing $+19$ gives $2x = 25 - 19 = 6$. Dividing by 2 yields $x = \frac{6}{2}$. Simplifies to $x = 3.00$.

Q95. Transposing $+13$ gives $6x = 10 - 13 = -3$. Dividing by 6 yields $x = \frac{-3}{6}$. Simplifies to $x = -0.50$.

Q96. Transposing $+19$ gives $3x = -25 - 19 = -44$. Dividing by 3 yields $x = \frac{-44}{3}$. Simplifies to $x = -14.67$.

Q97. Transposing $+3$ gives $11x = -9 - 3 = -12$. Dividing by 11 yields $x = \frac{-12}{11}$. Simplifies to $x = -1.09$.

Q98. Transposing $+21$ gives $10x = -26 - 21 = -47$. Dividing by 10 yields $x = \frac{-47}{10}$. Simplifies to $x = -4.70$.

Q99. Transposing $+5$ gives $4x = 15 - 5 = 10$. Dividing by 4 yields $x = \frac{10}{4}$. Simplifies to $x = 2.50$.

Q100. Transposing -19 gives $8x = -14 + 19 = 5$. Dividing by 8 yields $x = \frac{5}{8}$. Simplifies to $x = 0.62$.

Q101. Transposing -10 gives $2x = -8 + 10 = 2$. Dividing by 2 yields $x = \frac{2}{2}$. Simplifies to $x = 1.00$.

Q102. Transposing $+16$ gives $5x = 20 - 16 = 4$. Dividing by 5 yields $x = \frac{4}{5}$. Simplifies to $x = 0.80$.

Q103. Transposing -19 gives $10x = -16 + 19 = 3$. Dividing by 10 yields $x = \frac{3}{10}$. Simplifies to $x = 0.30$.

Q104. Transposing -13 gives $13x = 11 + 13 = 24$. Dividing by 13 yields $x = \frac{24}{13}$. Simplifies to $x = 1.85$.

Q105. Transposing -14 gives $4x = 16 + 14 = 30$. Dividing by 4 yields $x = \frac{30}{4}$. Simplifies to $x = 7.50$.

Q106. Transposing $+25$ gives $11x = 12 - 25 = -13$. Dividing by 11 yields $x = \frac{-13}{11}$. Simplifies to $x = -1.18$.

Q107. Transposing -13 gives $13x = -2 + 13 = 11$. Dividing by 13 yields $x = \frac{11}{13}$. Simplifies to $x = 0.85$.

Q108. Transposing -23 gives $4x = -18 + 23 = 5$. Dividing by 4 yields $x = \frac{5}{4}$. Simplifies to $x = 1.25$.

Q109. Transposing $+20$ gives $10x = 20 - 20 = 0$. Dividing by 10 yields $x = \frac{0}{10}$. Simplifies to $x = 0.00$.

Q110. Transposing $+7$ gives $12x = 26 - 7 = 19$. Dividing by 12 yields $x = \frac{19}{12}$. Simplifies to $x = 1.58$.

Q111. Transposing -19 gives $5x = 7 + 19 = 26$. Dividing by 5 yields $x = \frac{26}{5}$. Simplifies to $x = 5.20$.

Q112. Transposing -25 gives $10x = -9 + 25 = 16$. Dividing by 10 yields $x = \frac{16}{10}$. Simplifies to $x = 1.60$.

Q113. Transposing -17 gives $10x = 6 + 17 = 23$. Dividing by 10 yields $x = \frac{23}{10}$. Simplifies to $x = 2.30$.

Q114. Transposing $+5$ gives $12x = 28 - 5 = 23$. Dividing by 12 yields $x = \frac{23}{12}$. Simplifies to $x = 1.92$.

Q115. Transposing -18 gives $4x = -23 + 18 = -5$. Dividing by 4 yields $x = \frac{-5}{4}$. Simplifies to $x = -1.25$.

Q116. Transposing $+3$ gives $2x = 9 - 3 = 6$. Dividing by 2 yields $x = \frac{6}{2}$. Simplifies to $x = 3.00$.

Q117. Transposing $+20$ gives $11x = 25 - 20 = 5$. Dividing by 11 yields $x = \frac{5}{11}$. Simplifies to $x = 0.45$.

Q118. Transposing $+13$ gives $11x = -25 - 13 = -38$. Dividing by 11 yields $x = \frac{-38}{11}$. Simplifies to $x = -3.45$.

Q119. Transposing $+7$ gives $6x = 18 - 7 = 11$. Dividing by 6 yields $x = \frac{11}{6}$. Simplifies to $x = 1.83$.

Q120. Transposing $+19$ gives $4x = -27 - 19 = -46$. Dividing by 4 yields $x = \frac{-46}{4}$. Simplifies to $x = -11.50$.

Q121. Transposing -7 gives $3x = 13 + 7 = 20$. Dividing by 3 yields $x = \frac{20}{3}$. Simplifies to $x = 6.67$.

Q122. Transposing -22 gives $10x = -24 + 22 = -2$. Dividing by 10 yields $x = \frac{-2}{10}$. Simplifies to $x = -0.20$.

Q123. Transposing -7 gives $6x = -15 + 7 = -8$. Dividing by 6 yields $x = \frac{-8}{6}$. Simplifies to $x = -1.33$.

Q124. Transposing $+21$ gives $4x = 22 - 21 = 1$. Dividing by 4 yields $x = \frac{1}{4}$. Simplifies to $x = 0.25$.

Q125. Transposing $+16$ gives $8x = 5 - 16 = -11$. Dividing by 8 yields $x = \frac{-11}{8}$. Simplifies to $x = -1.38$.

Q126. Transposing -16 gives $8x = 20 + 16 = 36$. Dividing by 8 yields $x = \frac{36}{8}$. Simplifies to $x = 4.50$.

Q127. Transposing $+20$ gives $15x = 18 - 20 = -2$. Dividing by 15 yields $x = \frac{-2}{15}$. Simplifies to $x = -0.13$.

Q128. Transposing $+15$ gives $13x = -9 - 15 = -24$. Dividing by 13 yields $x = \frac{-24}{13}$. Simplifies to $x = -1.85$.

Q129. Transposing -23 gives $13x = -15 + 23 = 8$. Dividing by 13 yields $x = \frac{8}{13}$. Simplifies to $x = 0.62$.

Q130. Transposing -1 gives $6x = 27 + 1 = 28$. Dividing by 6 yields $x = \frac{28}{6}$. Simplifies to $x = 4.67$.

Q131. Transposing $+5$ gives $5x = -15 - 5 = -20$. Dividing by 5 yields $x = \frac{-20}{5}$. Simplifies to $x = -4.00$.

Q132. Transposing $+1$ gives $5x = -21 - 1 = -22$. Dividing by 5 yields $x = \frac{-22}{5}$. Simplifies to $x = -4.40$.

Q133. Transposing -15 gives $11x = 1 + 15 = 16$. Dividing by 11 yields $x = \frac{16}{11}$. Simplifies to $x = 1.45$.

Q134. Transposing $+6$ gives $9x = 29 - 6 = 23$. Dividing by 9 yields $x = \frac{23}{9}$. Simplifies to $x = 2.56$.

Q135. Transposing -8 gives $5x = 18 + 8 = 26$. Dividing by 5 yields $x = \frac{26}{5}$. Simplifies to $x = 5.20$.

Q136. Transposing $+23$ gives $2x = 28 - 23 = 5$. Dividing by 2 yields $x = \frac{5}{2}$. Simplifies to $x = 2.50$.

Q137. Transposing -4 gives $5x = 19 + 4 = 23$. Dividing by 5 yields $x = \frac{23}{5}$. Simplifies to $x = 4.60$.

Q138. Transposing -11 gives $2x = -11 + 11 = 0$. Dividing by 2 yields $x = \frac{0}{2}$. Simplifies to $x = 0.00$.

Q139. Transposing -9 gives $10x = -28 + 9 = -19$. Dividing by 10 yields $x = \frac{-19}{10}$. Simplifies to $x = -1.90$.

Q140. Transposing -1 gives $14x = -22 + 1 = -21$. Dividing by 14 yields $x = \frac{-21}{14}$. Simplifies to $x = -1.50$.

Q141. Transposing $+12$ gives $6x = 22 - 12 = 10$. Dividing by 6 yields $x = \frac{10}{6}$. Simplifies to $x = 1.67$.

Q142. Transposing $+21$ gives $12x = 11 - 21 = -10$. Dividing by 12 yields $x = \frac{-10}{12}$. Simplifies to $x = -0.83$.

Q143. Transposing $+14$ gives $12x = -7 - 14 = -21$. Dividing by 12 yields $x = \frac{-21}{12}$. Simplifies to $x = -1.75$.

Q144. Transposing $+25$ gives $14x = 27 - 25 = 2$. Dividing by 14 yields $x = \frac{2}{14}$. Simplifies to $x = 0.14$.

Q145. Transposing $+14$ gives $9x = 29 - 14 = 15$. Dividing by 9 yields $x = \frac{15}{9}$. Simplifies to $x = 1.67$.

Q146. Transposing -22 gives $6x = -1 + 22 = 21$. Dividing by 6 yields $x = \frac{21}{6}$. Simplifies to $x = 3.50$.

Q147. Transposing -6 gives $12x = -25 + 6 = -19$. Dividing by 12 yields $x = \frac{-19}{12}$. Simplifies to $x = -1.58$.

Q148. Transposing $+17$ gives $4x = -29 - 17 = -46$. Dividing by 4 yields $x = \frac{-46}{4}$. Simplifies to $x = -11.50$.

Q149. Transposing $+5$ gives $7x = 24 - 5 = 19$. Dividing by 7 yields $x = \frac{19}{7}$. Simplifies to $x = 2.71$.

Q150. Transposing -1 gives $2x = -28 + 1 = -27$. Dividing by 2 yields $x = \frac{-27}{2}$. Simplifies to $x = -13.50$.

Q151. Transposing -22 gives $7x = -16 + 22 = 6$. Dividing by 7 yields $x = \frac{6}{7}$. Simplifies to $x = 0.86$.

Q152. Transposing -1 gives $2x = -22 + 1 = -21$. Dividing by 2 yields $x = \frac{-21}{2}$. Simplifies to $x = -10.50$.

Q153. Transposing $+7$ gives $9x = 23 - 7 = 16$. Dividing by 9 yields $x = \frac{16}{9}$. Simplifies to $x = 1.78$.

Q154. Transposing $+21$ gives $3x = -19 - 21 = -40$. Dividing by 3 yields $x = \frac{-40}{3}$. Simplifies to $x = -13.33$.

Q155. Transposing $+19$ gives $8x = 12 - 19 = -7$. Dividing by 8 yields $x = \frac{-7}{8}$. Simplifies to $x = -0.88$.

Q156. Transposing $+22$ gives $14x = 16 - 22 = -6$. Dividing by 14 yields $x = \frac{-6}{14}$. Simplifies to $x = -0.43$.

Q157. Transposing -8 gives $9x = -7 + 8 = 1$. Dividing by 9 yields $x = \frac{1}{9}$. Simplifies to $x = 0.11$.

Q158. Transposing $+21$ gives $15x = -17 - 21 = -38$. Dividing by 15 yields $x = \frac{-38}{15}$. Simplifies to $x = -2.53$.

Q159. Transposing $+13$ gives $12x = -12 - 13 = -25$. Dividing by 12 yields $x = \frac{-25}{12}$. Simplifies to $x = -2.08$.

Q160. Transposing $+7$ gives $15x = -22 - 7 = -29$. Dividing by 15 yields $x = \frac{-29}{15}$. Simplifies to $x = -1.93$.

Q161. Transposing -15 gives $14x = 3 + 15 = 18$. Dividing by 14 yields $x = \frac{18}{14}$. Simplifies to $x = 1.29$.

Q162. Transposing $+6$ gives $12x = -11 - 6 = -17$. Dividing by 12 yields $x = \frac{-17}{12}$. Simplifies to $x = -1.42$.

Q163. Transposing $+1$ gives $5x = 26 - 1 = 25$. Dividing by 5 yields $x = \frac{25}{5}$. Simplifies to $x = 5.00$.

Q164. Transposing $+8$ gives $15x = -28 - 8 = -36$. Dividing by 15 yields $x = \frac{-36}{15}$. Simplifies to $x = -2.40$.

Q165. Transposing $+8$ gives $2x = -24 - 8 = -32$. Dividing by 2 yields $x = \frac{-32}{2}$. Simplifies to $x = -16.00$.

Q166. Transposing $+20$ gives $13x = -30 - 20 = -50$. Dividing by 13 yields $x = \frac{-50}{13}$. Simplifies to $x = -3.85$.

Q167. Transposing $+19$ gives $14x = 1 - 19 = -18$. Dividing by 14 yields $x = \frac{-18}{14}$. Simplifies to $x = -1.29$.

Q168. Transposing $+4$ gives $6x = 10 - 4 = 6$. Dividing by 6 yields $x = \frac{6}{6}$. Simplifies to $x = 1.00$.

Q169. Transposing $+14$ gives $7x = -12 - 14 = -26$. Dividing by 7 yields $x = \frac{-26}{7}$. Simplifies to $x = -3.71$.

Q170. Transposing -2 gives $4x = 30 + 2 = 32$. Dividing by 4 yields $x = \frac{32}{4}$. Simplifies to $x = 8.00$.

Q171. Transposing -10 gives $9x = 19 + 10 = 29$. Dividing by 9 yields $x = \frac{29}{9}$. Simplifies to $x = 3.22$.

Q172. Transposing $+24$ gives $4x = -2 - 24 = -26$. Dividing by 4 yields $x = \frac{-26}{4}$. Simplifies to $x = -6.50$.

Q173. Transposing -21 gives $13x = -6 + 21 = 15$. Dividing by 13 yields $x = \frac{15}{13}$. Simplifies to $x = 1.15$.

Q174. Transposing $+14$ gives $3x = 18 - 14 = 4$. Dividing by 3 yields $x = \frac{4}{3}$. Simplifies to $x = 1.33$.

Q175. Transposing -19 gives $9x = 4 + 19 = 23$. Dividing by 9 yields $x = \frac{23}{9}$. Simplifies to $x = 2.56$.

Q176. Transposing $+18$ gives $2x = -1 - 18 = -19$. Dividing by 2 yields $x = \frac{-19}{2}$. Simplifies to $x = -9.50$.

Q177. Transposing -3 gives $6x = 23 + 3 = 26$. Dividing by 6 yields $x = \frac{26}{6}$. Simplifies to $x = 4.33$.

Q178. Transposing -15 gives $8x = 18 + 15 = 33$. Dividing by 8 yields $x = \frac{33}{8}$. Simplifies to $x = 4.12$.

Q179. Transposing $+10$ gives $3x = 20 - 10 = 10$. Dividing by 3 yields $x = \frac{10}{3}$. Simplifies to $x = 3.33$.

Q180. Transposing $+23$ gives $5x = 24 - 23 = 1$. Dividing by 5 yields $x = \frac{1}{5}$. Simplifies to $x = 0.20$.

Q181. Transposing $+9$ gives $13x = 2 - 9 = -7$. Dividing by 13 yields $x = \frac{-7}{13}$. Simplifies to $x = -0.54$.

Q182. Transposing $+1$ gives $14x = 17 - 1 = 16$. Dividing by 14 yields $x = \frac{16}{14}$. Simplifies to $x = 1.14$.

Q183. Transposing $+23$ gives $8x = -19 - 23 = -42$. Dividing by 8 yields $x = \frac{-42}{8}$. Simplifies to $x = -5.25$.

Q184. Transposing -22 gives $13x = -7 + 22 = 15$. Dividing by 13 yields $x = \frac{15}{13}$. Simplifies to $x = 1.15$.

Q185. Transposing $+12$ gives $13x = -24 - 12 = -36$. Dividing by 13 yields $x = \frac{-36}{13}$. Simplifies to $x = -2.77$.

Q186. Transposing -15 gives $2x = -14 + 15 = 1$. Dividing by 2 yields $x = \frac{1}{2}$. Simplifies to $x = 0.50$.

Q187. Transposing $+20$ gives $15x = -20 - 20 = -40$. Dividing by 15 yields $x = \frac{-40}{15}$. Simplifies to $x = -2.67$.

Q188. Transposing -17 gives $10x = 5 + 17 = 22$. Dividing by 10 yields $x = \frac{22}{10}$. Simplifies to $x = 2.20$.

Q189. Transposing -7 gives $13x = 3 + 7 = 10$. Dividing by 13 yields $x = \frac{10}{13}$. Simplifies to $x = 0.77$.

Q190. Transposing $+12$ gives $11x = 26 - 12 = 14$. Dividing by 11 yields $x = \frac{14}{11}$. Simplifies to $x = 1.27$.

Q191. Transposing -10 gives $12x = 5 + 10 = 15$. Dividing by 12 yields $x = \frac{15}{12}$. Simplifies to $x = 1.25$.

Q192. Transposing $+21$ gives $11x = 14 - 21 = -7$. Dividing by 11 yields $x = \frac{-7}{11}$. Simplifies to $x = -0.64$.

Q193. Transposing -13 gives $3x = -22 + 13 = -9$. Dividing by 3 yields $x = \frac{-9}{3}$. Simplifies to $x = -3.00$.

Q194. Transposing $+10$ gives $2x = -18 - 10 = -28$. Dividing by 2 yields $x = \frac{-28}{2}$. Simplifies to $x = -14.00$.

Q195. Transposing $+5$ gives $15x = -23 - 5 = -28$. Dividing by 15 yields $x = \frac{-28}{15}$. Simplifies to $x = -1.87$.

Q196. Transposing -16 gives $11x = 0 + 16 = 16$. Dividing by 11 yields $x = \frac{16}{11}$. Simplifies to $x = 1.45$.

Q197. Transposing -6 gives $9x = -19 + 6 = -13$. Dividing by 9 yields $x = \frac{-13}{9}$. Simplifies to $x = -1.44$.

Q198. Transposing -14 gives $4x = -20 + 14 = -6$. Dividing by 4 yields $x = \frac{-6}{4}$. Simplifies to $x = -1.50$.

Q199. Transposing $+2$ gives $9x = 0 - 2 = -2$. Dividing by 9 yields $x = \frac{-2}{9}$. Simplifies to $x = -0.22$.

Q200. Transposing -22 gives $14x = -20 + 22 = 2$. Dividing by 14 yields $x = \frac{2}{14}$. Simplifies to $x = 0.14$.

Q201. Distributing -7 gives $21 - 21x + 70 = 31 \implies -21x + 91 = 31 \implies -21x = -60 \implies x = \frac{-60}{-21} = \frac{60}{21}$.

Q202. Distributing -4 gives $5 - 48x + 4 = 15 \implies -48x + 9 = 15 \implies -48x = 6 \implies x = \frac{6}{-48} = \frac{-6}{48}$.

Q203. Distributing -7 gives $22 - 14x + 21 = 32 \implies -14x + 43 = 32 \implies -14x = -11 \implies x = \frac{-11}{-14} = \frac{11}{14}$.

Q204. Distributing -5 gives $31 - 60x + 15 = 41 \implies -60x + 46 = 41 \implies -60x = -5 \implies x = \frac{-5}{-60} = \frac{5}{60}$.

Q205. Distributing -5 gives $17 - 60x + 70 = 27 \implies -60x + 87 = 27 \implies -60x = -60 \implies x = \frac{-60}{-60} = \frac{60}{60}$.

Q206. Distributing -7 gives $26 - 28x + 91 = 36 \implies -28x + 117 = 36 \implies -28x = -81 \implies x = \frac{-81}{-28} = \frac{81}{28}$.

Q207. Distributing -7 gives $19 - 56x + 42 = 29 \implies -56x + 61 = 29 \implies -56x = -32 \implies x = \frac{-32}{-56} = \frac{32}{56}$.

Q208. Distributing -6 gives $38 - 36x + 30 = 48 \implies -36x + 68 = 48 \implies -36x = -20 \implies x = \frac{-20}{-36} = \frac{20}{36}$.

Q209. Distributing -8 gives $14 - 32x + 88 = 24 \implies -32x + 102 = 24 \implies -32x = -78 \implies x = \frac{-78}{-32} = \frac{78}{32}$.

Q210. Distributing -2 gives $38 - 16x + 4 = 48 \implies -16x + 42 = 48 \implies -16x = 6 \implies x = \frac{6}{-16} = \frac{-6}{16}$.

Q211. Distributing -8 gives $18 - 32x + 40 = 28 \implies -32x + 58 = 28 \implies -32x = -30 \implies x = \frac{-30}{-32} = \frac{30}{32}$.

Q212. Distributing -9 gives $18 - 81x + 135 = 28 \implies -81x + 153 = 28 \implies -81x = -125 \implies x = \frac{-125}{-81} = \frac{125}{81}$.

Q213. Distributing -4 gives $31 - 12x + 8 = 41 \implies -12x + 39 = 41 \implies -12x = 2 \implies x = \frac{2}{-12} = \frac{-2}{12}$.

Q214. Distributing -8 gives $16 - 16x + 40 = 26 \implies -16x + 56 = 26 \implies -16x = -30 \implies x = \frac{-30}{-16} = \frac{30}{16}$.

Q215. Distributing -4 gives $30 - 40x + 40 = 40 \implies -40x + 70 = 40 \implies -40x = -30 \implies x = \frac{-30}{-40} = \frac{30}{40}$.

Q216. Distributing -2 gives $10 - 22x + 12 = 20 \implies -22x + 22 = 20 \implies -22x = -2 \implies x = \frac{-2}{-22} = \frac{2}{22}$.

Q217. Distributing -4 gives $40 - 20x + 52 = 50 \implies -20x + 92 = 50 \implies -20x = -42 \implies x = \frac{-42}{-20} = \frac{42}{20}$.

Q218. Distributing -6 gives $8 - 12x + 18 = 18 \implies -12x + 26 = 18 \implies -12x = -8 \implies x = \frac{-8}{-12} = \frac{8}{12}$.

Q219. Distributing -8 gives $37 - 56x + 96 = 47 \implies -56x + 133 = 47 \implies -56x = -86 \implies x = \frac{-86}{-56} = \frac{86}{56}$.

Q220. Distributing -5 gives $38 - 60x + 70 = 48 \implies -60x + 108 = 48 \implies -60x = -60 \implies x = \frac{-60}{-60} = \frac{60}{60}$.

Q221. Distributing -9 gives $28 - 36x + 36 = 38 \implies -36x + 64 = 38 \implies -36x = -26 \implies x = \frac{-26}{-36} = \frac{26}{36}$.

Q222. Distributing -4 gives $34 - 8x + 32 = 44 \implies -8x + 66 = 44 \implies -8x = -22 \implies x = \frac{-22}{-8} = \frac{22}{8}$.

Q223. Distributing -5 gives $33 - 30x + 75 = 43 \implies -30x + 108 = 43 \implies -30x = -65 \implies x = \frac{-65}{-30} = \frac{65}{30}$.

Q224. Distributing -7 gives $33 - 21x + 49 = 43 \implies -21x + 82 = 43 \implies -21x = -39 \implies x = \frac{-39}{-21} = \frac{39}{21}$.

Q225. Distributing -7 gives $27 - 14x + 84 = 37 \implies -14x + 111 = 37 \implies -14x = -74 \implies x = \frac{-74}{-14} = \frac{74}{14}$.

Q226. Distributing -9 gives $9 - 18x + 117 = 19 \implies -18x + 126 = 19 \implies -18x = -107 \implies x = \frac{-107}{-18} = \frac{107}{18}$.

Q227. Distributing -7 gives $25 - 49x + 70 = 35 \implies -49x + 95 = 35 \implies -49x = -60 \implies x = \frac{-60}{-49} = \frac{60}{49}$.

Q228. Distributing -3 gives $34 - 6x + 6 = 44 \implies -6x + 40 = 44 \implies -6x = 4 \implies x = \frac{4}{-6} = \frac{-4}{6}$.

Q229. Distributing -8 gives $22 - 48x + 32 = 32 \implies -48x + 54 = 32 \implies -48x = -22 \implies x = \frac{-22}{-48} = \frac{22}{48}$.

Q230. Distributing -5 gives $39 - 20x + 70 = 49 \implies -20x + 109 = 49 \implies -20x = -60 \implies x = \frac{-60}{-20} = \frac{60}{20}$.

Q231. Distributing -5 gives $37 - 20x + 45 = 47 \implies -20x + 82 = 47 \implies -20x = -35 \implies x = \frac{-35}{-20} = \frac{35}{20}$.

Q232. Distributing -6 gives $11 - 36x + 78 = 21 \implies -36x + 89 = 21 \implies -36x = -68 \implies x = \frac{-68}{-36} = \frac{68}{36}$.

Q233. Distributing -4 gives $6 - 28x + 48 = 16 \implies -28x + 54 = 16 \implies -28x = -38 \implies x = \frac{-38}{-28} = \frac{38}{28}$.

Q234. Distributing -4 gives $25 - 44x + 28 = 35 \implies -44x + 53 = 35 \implies -44x = -18 \implies x = \frac{-18}{-44} = \frac{18}{44}$.

Q235. Distributing -6 gives $26 - 42x + 60 = 36 \implies -42x + 86 = 36 \implies -42x = -50 \implies x = \frac{-50}{-42} = \frac{50}{42}$.

Q236. Distributing -4 gives $37 - 32x + 44 = 47 \implies -32x + 81 = 47 \implies -32x = -34 \implies x = \frac{-34}{-32} = \frac{34}{32}$.

Q237. Distributing -5 gives $6 - 40x + 5 = 16 \implies -40x + 11 = 16 \implies -40x = 5 \implies x = \frac{5}{-40} = \frac{-5}{40}$.

Q238. Distributing -6 gives $17 - 42x + 84 = 27 \implies -42x + 101 = 27 \implies -42x = -74 \implies x = \frac{-74}{-42} = \frac{74}{42}$.

Q239. Distributing -5 gives $15 - 35x + 65 = 25 \implies -35x + 80 = 25 \implies -35x = -55 \implies x = \frac{-55}{-35} = \frac{55}{35}$.

Q240. Distributing -3 gives $24 - 36x + 45 = 34 \implies -36x + 69 = 34 \implies -36x = -35 \implies x = \frac{-35}{-36} = \frac{35}{36}$.

Q241. Distributing -3 gives $22 - 18x + 27 = 32 \implies -18x + 49 = 32 \implies -18x = -17 \implies x = \frac{-17}{-18} = \frac{17}{18}$.

Q242. Distributing -4 gives $19 - 28x + 4 = 29 \implies -28x + 23 = 29 \implies -28x = 6 \implies x = \frac{6}{-28} = \frac{-6}{28}$.

Q243. Distributing -5 gives $5 - 10x + 45 = 15 \implies -10x + 50 = 15 \implies -10x = -35 \implies x = \frac{-35}{-10} = \frac{35}{10}$.

Q244. Distributing -6 gives $30 - 72x + 60 = 40 \implies -72x + 90 = 40 \implies -72x = -50 \implies x = \frac{-50}{-72} = \frac{50}{72}$.

Q245. Distributing -3 gives $11 - 27x + 18 = 21 \implies -27x + 29 = 21 \implies -27x = -8 \implies x = \frac{-8}{-27} = \frac{8}{27}$.

Q246. Distributing -8 gives $26 - 64x + 56 = 36 \implies -64x + 82 = 36 \implies -64x = -46 \implies x = \frac{-46}{-64} = \frac{46}{64}$.

Q247. Distributing -8 gives $11 - 24x + 80 = 21 \implies -24x + 91 = 21 \implies -24x = -70 \implies x = \frac{-70}{-24} = \frac{70}{24}$.

Q248. Distributing -6 gives $35 - 36x + 54 = 45 \implies -36x + 89 = 45 \implies -36x = -44 \implies x = \frac{-44}{-36} = \frac{44}{36}$.

Q249. Distributing -3 gives $14 - 24x + 12 = 24 \implies -24x + 26 = 24 \implies -24x = -2 \implies x = \frac{-2}{-24} = \frac{2}{24}$.

Q250. Distributing -2 gives $15 - 10x + 16 = 25 \implies -10x + 31 = 25 \implies -10x = -6 \implies x = \frac{-6}{-10} = \frac{6}{10}$.

Q251. Distributing -3 gives $15 - 9x + 27 = 25 \implies -9x + 42 = 25 \implies -9x = -17 \implies x = \frac{-17}{-9} = \frac{17}{9}$.

Q252. Distributing -6 gives $29 - 30x + 72 = 39 \implies -30x + 101 = 39 \implies -30x = -62 \implies x = \frac{-62}{-30} = \frac{62}{30}$.

Q253. Distributing -9 gives $15 - 27x + 135 = 25 \implies -27x + 150 = 25 \implies -27x = -125 \implies x = \frac{-125}{-27} = \frac{125}{27}$.

Q254. Distributing -6 gives $15 - 42x + 84 = 25 \implies -42x + 99 = 25 \implies -42x = -74 \implies x = \frac{-74}{-42} = \frac{74}{42}$.

Q255. Distributing -3 gives $22 - 15x + 33 = 32 \implies -15x + 55 = 32 \implies -15x = -23 \implies x = \frac{-23}{-15} = \frac{23}{15}$.

Q256. Distributing -2 gives $8 - 8x + 12 = 18 \implies -8x + 20 = 18 \implies -8x = -2 \implies x = \frac{-2}{-8} = \frac{2}{8}$.

Q257. Distributing -8 gives $15 - 16x + 88 = 25 \implies -16x + 103 = 25 \implies -16x = -78 \implies x = \frac{-78}{-16} = \frac{78}{16}$.

Q258. Distributing -8 gives $7 - 96x + 72 = 17 \implies -96x + 79 = 17 \implies -96x = -62 \implies x = \frac{-62}{-96} = \frac{62}{96}$.

Q259. Distributing -5 gives $35 - 20x + 20 = 45 \implies -20x + 55 = 45 \implies -20x = -10 \implies x = \frac{-10}{-20} = \frac{10}{20}$.

Q260. Distributing -3 gives $32 - 24x + 9 = 42 \implies -24x + 41 = 42 \implies -24x = 1 \implies x = \frac{1}{-24} = \frac{-1}{24}$.

Q261. Distributing -4 gives $38 - 32x + 56 = 48 \implies -32x + 94 = 48 \implies -32x = -46 \implies x = \frac{-46}{-32} = \frac{46}{32}$.

Q262. Distributing -7 gives $10 - 70x + 98 = 20 \implies -70x + 108 = 20 \implies -70x = -88 \implies x = \frac{-88}{-70} = \frac{88}{70}$.

Q263. Distributing -3 gives $31 - 30x + 3 = 41 \implies -30x + 34 = 41 \implies -30x = 7 \implies x = \frac{7}{-30} = \frac{-7}{30}$.

Q264. Distributing -4 gives $34 - 40x + 44 = 44 \implies -40x + 78 = 44 \implies -40x = -34 \implies x = \frac{-34}{-40} = \frac{34}{40}$.

Q265. Distributing -4 gives $21 - 48x + 48 = 31 \implies -48x + 69 = 31 \implies -48x = -38 \implies x = \frac{-38}{-48} = \frac{38}{48}$.

Q266. Distributing -9 gives $11 - 27x + 27 = 21 \implies -27x + 38 = 21 \implies -27x = -17 \implies x = \frac{-17}{-27} = \frac{17}{27}$.

Q267. Distributing -9 gives $19 - 36x + 36 = 29 \implies -36x + 55 = 29 \implies -36x = -26 \implies x = \frac{-26}{-36} = \frac{26}{36}$.

Q268. Distributing -2 gives $5 - 20x + 30 = 15 \implies -20x + 35 = 15 \implies -20x = -20 \implies x = \frac{-20}{-20} = \frac{20}{20}$.

Q269. Distributing -4 gives $26 - 24x + 16 = 36 \implies -24x + 42 = 36 \implies -24x = -6 \implies x = \frac{-6}{-24} = \frac{6}{24}$.

Q270. Distributing -5 gives $12 - 45x + 10 = 22 \implies -45x + 22 = 22 \implies -45x = 0 \implies x = \frac{0}{-45} = \frac{0}{45}$.

Q271. Distributing -2 gives $20 - 24x + 8 = 30 \implies -24x + 28 = 30 \implies -24x = 2 \implies x = \frac{2}{-24} = \frac{-2}{24}$.

Q272. Distributing -9 gives $12 - 90x + 72 = 22 \implies -90x + 84 = 22 \implies -90x = -62 \implies x = \frac{-62}{-90} = \frac{62}{90}$.

Q273. Distributing -7 gives $25 - 21x + 28 = 35 \implies -21x + 53 = 35 \implies -21x = -18 \implies x = \frac{-18}{-21} = \frac{18}{21}$.

Q274. Distributing -6 gives $33 - 24x + 6 = 43 \implies -24x + 39 = 43 \implies -24x = 4 \implies x = \frac{4}{-24} = \frac{-4}{24}$.

Q275. Distributing -4 gives $9 - 24x + 40 = 19 \implies -24x + 49 = 19 \implies -24x = -30 \implies x = \frac{-30}{-24} = \frac{30}{24}$.

Q276. Distributing -2 gives $9 - 4x + 4 = 19 \implies -4x + 13 = 19 \implies -4x = 6 \implies x = \frac{6}{-4} = \frac{-6}{4}$.

Q277. Distributing -3 gives $30 - 12x + 33 = 40 \implies -12x + 63 = 40 \implies -12x = -23 \implies x = \frac{-23}{-12} = \frac{23}{12}$.

Q278. Distributing -6 gives $8 - 72x + 6 = 18 \implies -72x + 14 = 18 \implies -72x = 4 \implies x = \frac{4}{-72} = \frac{-4}{72}$.

Q279. Distributing -4 gives $12 - 48x + 36 = 22 \implies -48x + 48 = 22 \implies -48x = -26 \implies x = \frac{-26}{-48} = \frac{26}{48}$.

Q280. Distributing -9 gives $22 - 72x + 99 = 32 \implies -72x + 121 = 32 \implies -72x = -89 \implies x = \frac{-89}{-72} = \frac{89}{72}$.

Q281. Distributing -3 gives $14 - 27x + 18 = 24 \implies -27x + 32 = 24 \implies -27x = -8 \implies x = \frac{-8}{-27} = \frac{8}{27}$.

Q282. Distributing -7 gives $20 - 63x + 70 = 30 \implies -63x + 90 = 30 \implies -63x = -60 \implies x = \frac{-60}{-63} = \frac{60}{63}$.

Q283. Distributing -7 gives $16 - 77x + 21 = 26 \implies -77x + 37 = 26 \implies -77x = -11 \implies x = \frac{-11}{-77} = \frac{11}{77}$.

Q284. Distributing -9 gives $21 - 72x + 108 = 31 \implies -72x + 129 = 31 \implies -72x = -98 \implies x = \frac{-98}{-72} = \frac{98}{72}$.

Q285. Distributing -2 gives $19 - 20x + 22 = 29 \implies -20x + 41 = 29 \implies -20x = -12 \implies x = \frac{-12}{-20} = \frac{12}{20}$.

Q286. Distributing -5 gives $33 - 10x + 15 = 43 \implies -10x + 48 = 43 \implies -10x = -5 \implies x = \frac{-5}{-10} = \frac{5}{10}$.

Q287. Distributing -9 gives $39 - 63x + 81 = 49 \implies -63x + 120 = 49 \implies -63x = -71 \implies x = \frac{-71}{-63} = \frac{71}{63}$.

Q288. Distributing -6 gives $34 - 18x + 78 = 44 \implies -18x + 112 = 44 \implies -18x = -68 \implies x = \frac{-68}{-18} = \frac{68}{18}$.

Q289. Distributing -6 gives $21 - 42x + 30 = 31 \implies -42x + 51 = 31 \implies -42x = -20 \implies x = \frac{-20}{-42} = \frac{20}{42}$.

Q290. Distributing -8 gives $26 - 40x + 56 = 36 \implies -40x + 82 = 36 \implies -40x = -46 \implies x = \frac{-46}{-40} = \frac{46}{40}$.

Q291. Distributing -5 gives $26 - 55x + 55 = 36 \implies -55x + 81 = 36 \implies -55x = -45 \implies x = \frac{-45}{-55} = \frac{45}{55}$.

Q292. Distributing -2 gives $25 - 20x + 6 = 35 \implies -20x + 31 = 35 \implies -20x = 4 \implies x = \frac{4}{-20} = \frac{-4}{20}$.

Q293. Distributing -5 gives $38 - 15x + 35 = 48 \implies -15x + 73 = 48 \implies -15x = -25 \implies x = \frac{-25}{-15} = \frac{25}{15}$.

Q294. Distributing -9 gives $26 - 36x + 99 = 36 \implies -36x + 125 = 36 \implies -36x = -89 \implies x = \frac{-89}{-36} = \frac{89}{36}$.

Q295. Distributing -6 gives $38 - 30x + 78 = 48 \implies -30x + 116 = 48 \implies -30x = -68 \implies x = \frac{-68}{-30} = \frac{68}{30}$.

Q296. Distributing -2 gives $28 - 4x + 28 = 38 \implies -4x + 56 = 38 \implies -4x = -18 \implies x = \frac{-18}{-4} = \frac{18}{4}$.

Q297. Distributing -5 gives $11 - 40x + 45 = 21 \implies -40x + 56 = 21 \implies -40x = -35 \implies x = \frac{-35}{-40} = \frac{35}{40}$.

Q298. Distributing -6 gives $5 - 24x + 48 = 15 \implies -24x + 53 = 15 \implies -24x = -38 \implies x = \frac{-38}{-24} = \frac{38}{24}$.

Q299. Distributing -6 gives $39 - 24x + 66 = 49 \implies -24x + 105 = 49 \implies -24x = -56 \implies x = \frac{-56}{-24} = \frac{56}{24}$.

Q300. Distributing -8 gives $23 - 88x + 88 = 33 \implies -88x + 111 = 33 \implies -88x = -78 \implies x = \frac{-78}{-88} = \frac{78}{88}$.

Q301. Distributing -3 gives $23 - 30x + 6 = 33 \implies -30x + 29 = 33 \implies -30x = 4 \implies x = \frac{4}{-30} = \frac{-4}{30}$.

Q302. Distributing -4 gives $5 - 44x + 36 = 15 \implies -44x + 41 = 15 \implies -44x = -26 \implies x = \frac{-26}{-44} = \frac{26}{44}$.

Q303. Distributing -4 gives $20 - 36x + 8 = 30 \implies -36x + 28 = 30 \implies -36x = 2 \implies x = \frac{2}{-36} = \frac{-2}{36}$.

Q304. Distributing -2 gives $21 - 14x + 10 = 31 \implies -14x + 31 = 31 \implies -14x = 0 \implies x = \frac{0}{-14} = \frac{0}{14}$.

Q305. Distributing -9 gives $36 - 27x + 36 = 46 \implies -27x + 72 = 46 \implies -27x = -26 \implies x = \frac{-26}{-27} = \frac{26}{27}$.

Q306. Distributing -9 gives $14 - 108x + 63 = 24 \implies -108x + 77 = 24 \implies -108x = -53 \implies x = \frac{-53}{-108} = \frac{53}{108}$.

Q307. Distributing -5 gives $31 - 30x + 45 = 41 \implies -30x + 76 = 41 \implies -30x = -35 \implies x = \frac{-35}{-30} = \frac{35}{30}$.

Q308. Distributing -7 gives $36 - 28x + 49 = 46 \implies -28x + 85 = 46 \implies -28x = -39 \implies x = \frac{-39}{-28} = \frac{39}{28}$.

Q309. Distributing -3 gives $28 - 15x + 45 = 38 \implies -15x + 73 = 38 \implies -15x = -35 \implies x = \frac{-35}{-15} = \frac{35}{15}$.

Q310. Distributing -8 gives $11 - 80x + 64 = 21 \implies -80x + 75 = 21 \implies -80x = -54 \implies x = \frac{-54}{-80} = \frac{54}{80}$.

Q311. Distributing -2 gives $35 - 12x + 14 = 45 \implies -12x + 49 = 45 \implies -12x = -4 \implies x = \frac{-4}{-12} = \frac{4}{12}$.

Q312. Distributing -7 gives $36 - 49x + 84 = 46 \implies -49x + 120 = 46 \implies -49x = -74 \implies x = \frac{-74}{-49} = \frac{74}{49}$.

Q313. Distributing -4 gives $12 - 28x + 12 = 22 \implies -28x + 24 = 22 \implies -28x = -2 \implies x = \frac{-2}{-28} = \frac{2}{28}$.

Q314. Distributing -6 gives $40 - 24x + 60 = 50 \implies -24x + 100 = 50 \implies -24x = -50 \implies x = \frac{-50}{-24} = \frac{50}{24}$.

Q315. Distributing -8 gives $24 - 72x + 40 = 34 \implies -72x + 64 = 34 \implies -72x = -30 \implies x = \frac{-30}{-72} = \frac{30}{72}$.

Q316. Distributing -3 gives $35 - 30x + 3 = 45 \implies -30x + 38 = 45 \implies -30x = 7 \implies x = \frac{7}{-30} = \frac{-7}{30}$.

Q317. Distributing -2 gives $25 - 10x + 12 = 35 \implies -10x + 37 = 35 \implies -10x = -2 \implies x = \frac{-2}{-10} = \frac{2}{10}$.

Q318. Distributing -5 gives $22 - 45x + 25 = 32 \implies -45x + 47 = 32 \implies -45x = -15 \implies x = \frac{-15}{-45} = \frac{15}{45}$.

Q319. Distributing -7 gives $39 - 63x + 42 = 49 \implies -63x + 81 = 49 \implies -63x = -32 \implies x = \frac{-32}{-63} = \frac{32}{63}$.

Q320. Distributing -9 gives $28 - 18x + 81 = 38 \implies -18x + 109 = 38 \implies -18x = -71 \implies x = \frac{-71}{-18} = \frac{71}{18}$.

Q321. Distributing -8 gives $32 - 96x + 24 = 42 \implies -96x + 56 = 42 \implies -96x = -14 \implies x = \frac{-14}{-96} = \frac{14}{96}$.

Q322. Distributing -6 gives $37 - 42x + 18 = 47 \implies -42x + 55 = 47 \implies -42x = -8 \implies x = \frac{-8}{-42} = \frac{8}{42}$.

Q323. Distributing -8 gives $35 - 72x + 48 = 45 \implies -72x + 83 = 45 \implies -72x = -38 \implies x = \frac{-38}{-72} = \frac{38}{72}$.

Q324. Distributing -8 gives $12 - 24x + 120 = 22 \implies -24x + 132 = 22 \implies -24x = -110 \implies x = \frac{-110}{-24} = \frac{110}{24}$.

Q325. Distributing -2 gives $11 - 6x + 30 = 21 \implies -6x + 41 = 21 \implies -6x = -20 \implies x = \frac{-20}{-6} = \frac{20}{6}$.

Q326. Distributing -5 gives $25 - 45x + 20 = 35 \implies -45x + 45 = 35 \implies -45x = -10 \implies x = \frac{-10}{-45} = \frac{10}{45}$.

Q327. Distributing -5 gives $19 - 10x + 5 = 29 \implies -10x + 24 = 29 \implies -10x = 5 \implies x = \frac{5}{-10} = \frac{-5}{10}$.

Q328. Distributing -2 gives $22-10x+22=32 \implies -10x+44=32 \implies -10x=-12 \implies x=\frac{-12}{-10}=\frac{12}{10}$.

Q329. Distributing -9 gives $21-54x+135=31 \implies -54x+156=31 \implies -54x=-125 \implies x=\frac{-125}{-54}=\frac{125}{54}$.

Q330. Distributing -9 gives $28-27x+135=38 \implies -27x+163=38 \implies -27x=-125 \implies x=\frac{-125}{-27}=\frac{125}{27}$.

Q331. Distributing -2 gives $39-8x+20=49 \implies -8x+59=49 \implies -8x=-10 \implies x=\frac{-10}{-8}=\frac{10}{8}$.

Q332. Distributing -2 gives $9-18x+6=19 \implies -18x+15=19 \implies -18x=4 \implies x=\frac{4}{-18}=\frac{-4}{18}$.

Q333. Distributing -8 gives $13-72x+56=23 \implies -72x+69=23 \implies -72x=-46 \implies x=\frac{-46}{-72}=\frac{46}{72}$.

Q334. Distributing -2 gives $15-10x+20=25 \implies -10x+35=25 \implies -10x=-10 \implies x=\frac{-10}{-10}=\frac{10}{10}$.

Q335. Distributing -3 gives $24-12x+27=34 \implies -12x+51=34 \implies -12x=-17 \implies x=\frac{-17}{-12}=\frac{17}{12}$.

Q336. Distributing -5 gives $19-30x+35=29 \implies -30x+54=29 \implies -30x=-25 \implies x=\frac{-25}{-30}=\frac{25}{30}$.

Q337. Distributing -4 gives $19-12x+28=29 \implies -12x+47=29 \implies -12x=-18 \implies x=\frac{-18}{-12}=\frac{18}{12}$.

Q338. Distributing -9 gives $14-99x+99=24 \implies -99x+113=24 \implies -99x=-89 \implies x=\frac{-89}{-99}=\frac{89}{99}$.

Q339. Distributing -9 gives $28-81x+135=38 \implies -81x+163=38 \implies -81x=-125 \implies x=\frac{-125}{-81}=\frac{125}{81}$.

Q340. Distributing -7 gives $15-42x+63=25 \implies -42x+78=25 \implies -42x=-53 \implies x=\frac{-53}{-42}=\frac{53}{42}$.

Q341. Distributing -6 gives $36-42x+30=46 \implies -42x+66=46 \implies -42x=-20 \implies x=\frac{-20}{-42}=\frac{20}{42}$.

Q342. Distributing -6 gives $17-60x+60=27 \implies -60x+77=27 \implies -60x=-50 \implies x=\frac{-50}{-60}=\frac{50}{60}$.

Q343. Distributing -9 gives $8-45x+63=18 \implies -45x+71=18 \implies -45x=-53 \implies x=\frac{-53}{-45}=\frac{53}{45}$.

Q344. Distributing -8 gives $33-96x+16=43 \implies -96x+49=43 \implies -96x=-6 \implies x=\frac{-6}{-96}=\frac{6}{96}$.

Q345. Distributing -3 gives $28-6x+45=38 \implies -6x+73=38 \implies -6x=-35 \implies x=\frac{-35}{-6}=\frac{35}{6}$.

Q346. Distributing -9 gives $19-63x+63=29 \implies -63x+82=29 \implies -63x=-53 \implies x=\frac{-53}{-63}=\frac{53}{63}$.

Q347. Distributing -6 gives $19-12x+30=29 \implies -12x+49=29 \implies -12x=-20 \implies x=\frac{-20}{-12}=\frac{20}{12}$.

Q348. Distributing -2 gives $17-22x+12=27 \implies -22x+29=27 \implies -22x=-2 \implies x=\frac{-2}{-22}=\frac{2}{22}$.

Q349. Distributing -5 gives $19-45x+50=29 \implies -45x+69=29 \implies -45x=-40 \implies x=\frac{-40}{-45}=\frac{40}{45}$.

Q350. Distributing -3 gives $27-18x+27=37 \implies -18x+54=37 \implies -18x=-17 \implies x=\frac{-17}{-18}=\frac{17}{18}$.

Q351. Transposing $4x$ to left and -8 to right: $(10-4)x=-9-(-8) \implies 6x=-1 \implies x=\frac{-1}{6}$.

Q352. Transposing $7x$ to left and -17 to right: $(5-7)x=-18-(-17) \implies -2x=-1 \implies x=\frac{-1}{-2}$.

Q353. Transposing $7x$ to left and 2 to right: $(11-7)x=3-(2) \implies 4x=1 \implies x=\frac{1}{4}$.

Q354. Transposing $4x$ to left and -14 to right: $(5-4)x=3-(-14) \implies 1x=17 \implies x=\frac{17}{1}$.

Q355. Transposing $5x$ to left and -12 to right: $(12-5)x=12-(-12) \implies 7x=24 \implies x=\frac{24}{7}$.

Q356. Transposing $8x$ to left and 6 to right: $(5-8)x=-20-(6) \implies -3x=-26 \implies x=\frac{-26}{-3}$.

Q357. Transposing $2x$ to left and 10 to right: $(4-2)x=-2-(10) \implies 2x=-12 \implies x=\frac{-12}{2}$.

Q358. Transposing $10x$ to left and 4 to right: $(8-10)x=13-(4) \implies -2x=9 \implies x=\frac{9}{-2}$.

Q359. Transposing $8x$ to left and 19 to right: $(12-8)x=8-(19) \implies 4x=-11 \implies x=\frac{-11}{4}$.

Q360. Transposing $9x$ to left and 9 to right: $(6-9)x=-14-(9) \implies -3x=-23 \implies x=\frac{-23}{-3}$.

Q361. Transposing $4x$ to left and 16 to right: $(5-4)x=8-(16) \implies 1x=-8 \implies x=\frac{-8}{1}$.

Q362. Transposing $4x$ to left and -18 to right: $(8-4)x=19-(-18) \implies 4x=37 \implies x=\frac{37}{4}$.

Q363. Transposing $9x$ to left and -14 to right: $(8-9)x=4-(-14) \implies -1x=18 \implies x=\frac{18}{-1}$.

Q364. Transposing $3x$ to left and 18 to right: $(7-3)x=-18-(18) \implies 4x=-36 \implies x=\frac{-36}{4}$.

Q365. Transposing $4x$ to left and -15 to right: $(3-4)x=6-(-15) \implies -1x=21 \implies x=\frac{21}{-1}$.

Q366. Transposing $5x$ to left and -16 to right: $(9-5)x=-15-(-16) \implies 4x=1 \implies x=\frac{1}{4}$.

Q367. Transposing $5x$ to left and 2 to right: $(4-5)x=6-(2) \implies -1x=4 \implies x=\frac{4}{-1}$.

Q368. Transposing $3x$ to left and -14 to right: $(12-3)x=1-(-14) \implies 9x=15 \implies x=\frac{15}{9}$.

Q369. Transposing $8x$ to left and 1 to right: $(11-8)x=-9-(1) \implies 3x=-10 \implies x=\frac{-10}{3}$.

Q370. Transposing $2x$ to left and -5 to right: $(4-2)x=12-(-5) \implies 2x=17 \implies x=\frac{17}{2}$.

Q371. Transposing $4x$ to left and 12 to right: $(3-4)x=7-(12) \implies -1x=-5 \implies x=\frac{-5}{-1}$.

Q372. Transposing $5x$ to left and 14 to right: $(3-5)x=-2-(14) \implies -2x=-16 \implies x=\frac{-16}{-2}$.

Q373. Transposing $3x$ to left and 2 to right: $(7-3)x=19-(2) \implies 4x=17 \implies x=\frac{17}{4}$.

Q374. Transposing $8x$ to left and -14 to right: $(6-8)x=4-(-14) \implies -2x=18 \implies x=\frac{18}{-2}$.

Q375. Transposing $6x$ to left and 2 to right: $(3-6)x=-5-(2) \implies -3x=-7 \implies x=\frac{-7}{-3}$.

Q376. Transposing $3x$ to left and -8 to right: $(5-3)x=15-(-8) \implies 2x=23 \implies x=\frac{23}{2}$.

Q377. Transposing $10x$ to left and -20 to right: $(11-10)x=4-(-20) \implies 1x=24 \implies x=\frac{24}{1}$.

Q378. Transposing $6x$ to left and -13 to right: $(11-6)x=-6-(-13) \implies 5x=7 \implies x=\frac{7}{5}$.

Q379. Transposing $9x$ to left and -15 to right: $(10-9)x=1-(-15) \implies 1x=16 \implies x=\frac{16}{1}$.

Q380. Transposing $6x$ to left and 8 to right: $(8-6)x=18-(8) \implies 2x=10 \implies x=\frac{10}{2}$.

Q381. Transposing $6x$ to left and 12 to right: $(9-6)x=2-(12) \implies 3x=-10 \implies x=\frac{-10}{3}$.

Q382. Transposing $2x$ to left and 10 to right: $(11-2)x=8-(10) \implies 9x=-2 \implies x=\frac{-2}{9}$.

Q383. Transposing $4x$ to left and 7 to right: $(5-4)x = -5-(7) \implies 1x = -12 \implies x = \frac{-12}{1}$.

Q384. Transposing $6x$ to left and 0 to right: $(3-6)x = -4-(0) \implies -3x = -4 \implies x = \frac{-4}{-3}$.

Q385. Transposing $5x$ to left and 18 to right: $(11-5)x = 2-(18) \implies 6x = -16 \implies x = \frac{-16}{6}$.

Q386. Transposing $10x$ to left and 4 to right: $(6-10)x = -13-(4) \implies -4x = -17 \implies x = \frac{-17}{-4}$.

Q387. Transposing $10x$ to left and 0 to right: $(8-10)x = 17-(0) \implies -2x = 17 \implies x = \frac{17}{-2}$.

Q388. Transposing $10x$ to left and 2 to right: $(3-10)x = 7-(2) \implies -7x = 5 \implies x = \frac{5}{-7}$.

Q389. Transposing $7x$ to left and 0 to right: $(6-7)x = -19-(0) \implies -1x = -19 \implies x = \frac{-19}{-1}$.

Q390. Transposing $9x$ to left and -1 to right: $(7-9)x = 20-(-1) \implies -2x = 21 \implies x = \frac{21}{-2}$.

Q391. Transposing $7x$ to left and -13 to right: $(5-7)x = 5-(-13) \implies -2x = 18 \implies x = \frac{18}{-2}$.

Q392. Transposing $6x$ to left and 5 to right: $(11-6)x = 15-(5) \implies 5x = 10 \implies x = \frac{10}{5}$.

Q393. Transposing $4x$ to left and -1 to right: $(12-4)x = 9-(-1) \implies 8x = 10 \implies x = \frac{10}{8}$.

Q394. Transposing $3x$ to left and 10 to right: $(4-3)x = -15-(10) \implies 1x = -25 \implies x = \frac{-25}{1}$.

Q395. Transposing $2x$ to left and -16 to right: $(11-2)x = -17-(-16) \implies 9x = -1 \implies x = \frac{-1}{9}$.

Q396. Transposing $5x$ to left and 2 to right: $(8-5)x = 3-(2) \implies 3x = 1 \implies x = \frac{1}{3}$.

Q397. Transposing $4x$ to left and -9 to right: $(3-4)x = 13-(-9) \implies -1x = 22 \implies x = \frac{22}{-1}$.

Q398. Transposing $5x$ to left and 5 to right: $(4-5)x = 8-(5) \implies -1x = 3 \implies x = \frac{3}{-1}$.

Q399. Transposing $2x$ to left and -14 to right: $(6-2)x = -4-(-14) \implies 4x = 10 \implies x = \frac{10}{4}$.

Q400. Transposing $3x$ to left and -5 to right: $(8-3)x = -5-(-5) \implies 5x = 0 \implies x = \frac{0}{5}$.

Q401. Transposing $6x$ to left and 13 to right: $(7-6)x = 19-(13) \implies 1x = 6 \implies x = \frac{6}{1}$.

Q402. Transposing $3x$ to left and 6 to right: $(7-3)x = -17-(6) \implies 4x = -23 \implies x = \frac{-23}{4}$.

Q403. Transposing $10x$ to left and -2 to right: $(5-10)x = 1-(-2) \implies -5x = 3 \implies x = \frac{3}{-5}$.

Q404. Transposing $9x$ to left and -17 to right: $(10-9)x = -10-(-17) \implies 1x = 7 \implies x = \frac{7}{1}$.

Q405. Transposing $10x$ to left and 1 to right: $(9-10)x = 14-(1) \implies -1x = 13 \implies x = \frac{13}{-1}$.

Q406. Transposing $9x$ to left and 12 to right: $(7-9)x = 12-(12) \implies -2x = 0 \implies x = \frac{0}{-2}$.

Q407. Transposing $7x$ to left and -13 to right: $(11-7)x = -11-(-13) \implies 4x = 2 \implies x = \frac{2}{4}$.

Q408. Transposing $5x$ to left and -14 to right: $(3-5)x = 8-(-14) \implies -2x = 22 \implies x = \frac{22}{-2}$.

Q409. Transposing $10x$ to left and 11 to right: $(7-10)x = 7-(11) \implies -3x = -4 \implies x = \frac{-4}{-3}$.

Q410. Transposing $7x$ to left and 2 to right: $(11-7)x = -3-(2) \implies 4x = -5 \implies x = \frac{-5}{4}$.

Q411. Transposing $7x$ to left and -5 to right: $(10-7)x = -5-(-5) \implies 3x = 0 \implies x = \frac{0}{3}$.

Q412. Transposing $2x$ to left and 10 to right: $(9-2)x = 11-(10) \implies 7x = 1 \implies x = \frac{1}{7}$.

Q413. Transposing $7x$ to left and -9 to right: $(6-7)x = 13-(-9) \implies -1x = 22 \implies x = \frac{22}{-1}$.

Q414. Transposing $7x$ to left and -13 to right: $(9-7)x = -16-(-13) \implies 2x = -3 \implies x = \frac{-3}{2}$.

Q415. Transposing $4x$ to left and 7 to right: $(5-4)x = -19-(7) \implies 1x = -26 \implies x = \frac{-26}{1}$.

Q416. Transposing $8x$ to left and -4 to right: $(11-8)x = -17-(-4) \implies 3x = -13 \implies x = \frac{-13}{3}$.

Q417. Transposing $4x$ to left and -3 to right: $(5-4)x = -13-(-3) \implies 1x = -10 \implies x = \frac{-10}{1}$.

Q418. Transposing $5x$ to left and 10 to right: $(6-5)x = -16-(10) \implies 1x = -26 \implies x = \frac{-26}{1}$.

Q419. Transposing $9x$ to left and 20 to right: $(6-9)x = 15-(20) \implies -3x = -5 \implies x = \frac{-5}{-3}$.

Q420. Transposing $2x$ to left and 1 to right: $(4-2)x = 3-(1) \implies 2x = 2 \implies x = \frac{2}{2}$.

Q421. Transposing $10x$ to left and 12 to right: $(11-10)x = 11-(12) \implies 1x = -1 \implies x = \frac{-1}{1}$.

Q422. Transposing $8x$ to left and -15 to right: $(9-8)x = 14-(-15) \implies 1x = 29 \implies x = \frac{29}{1}$.

Q423. Transposing $6x$ to left and 11 to right: $(4-6)x = 2-(11) \implies -2x = -9 \implies x = \frac{-9}{-2}$.

Q424. Transposing $3x$ to left and -9 to right: $(5-3)x = 5-(-9) \implies 2x = 14 \implies x = \frac{14}{2}$.

Q425. Transposing $9x$ to left and -17 to right: $(3-9)x = -9-(-17) \implies -6x = 8 \implies x = \frac{8}{-6}$.

Q426. Transposing $2x$ to left and 15 to right: $(6-2)x = -7-(15) \implies 4x = -22 \implies x = \frac{-22}{4}$.

Q427. Transposing $9x$ to left and 17 to right: $(10-9)x = -3-(17) \implies 1x = -20 \implies x = \frac{-20}{1}$.

Q428. Transposing $10x$ to left and -5 to right: $(3-10)x = 6-(-5) \implies -7x = 11 \implies x = \frac{11}{-7}$.

Q429. Transposing $9x$ to left and -20 to right: $(8-9)x = 12-(-20) \implies -1x = 32 \implies x = \frac{32}{-1}$.

Q430. Transposing $3x$ to left and 2 to right: $(4-3)x = 6-(2) \implies 1x = 4 \implies x = \frac{4}{1}$.

Q431. Transposing $10x$ to left and 0 to right: $(8-10)x = -11-(0) \implies -2x = -11 \implies x = \frac{-11}{-2}$.

Q432. Transposing $10x$ to left and 12 to right: $(5-10)x = 5-(12) \implies -5x = -7 \implies x = \frac{-7}{-5}$.

Q433. Transposing $6x$ to left and 18 to right: $(11-6)x = 1-(18) \implies 5x = -17 \implies x = \frac{-17}{5}$.

Q434. Transposing $10x$ to left and 2 to right: $(11-10)x = -20-(2) \implies 1x = -22 \implies x = \frac{-22}{1}$.

Q435. Transposing $2x$ to left and 2 to right: $(4-2)x = 7-(2) \implies 2x = 5 \implies x = \frac{5}{2}$.

Q436. Transposing $4x$ to left and -1 to right: $(12-4)x = -6-(-1) \implies 8x = -5 \implies x = \frac{-5}{8}$.

Q437. Transposing $3x$ to left and 10 to right: $(7-3)x = 1-(10) \implies 4x = -9 \implies x = \frac{-9}{4}$.

Q438. Transposing $8x$ to left and 9 to right: $(3-8)x = 10-(9) \implies -5x = 1 \implies x = \frac{1}{-5}$.

Q439. Transposing $3x$ to left and -2 to right: $(5-3)x = -15-(-2) \implies 2x = -13 \implies x = \frac{-13}{2}$.

Q440. Transposing $7x$ to left and -17 to right: $(9-7)x = -14-(-17) \implies 2x = 3 \implies x = \frac{3}{2}$.

Q441. Transposing $4x$ to left and 15 to right: $(3-4)x = 20-(15) \implies -1x = 5 \implies x = \frac{5}{-1}$.

Q442. Transposing $8x$ to left and 15 to right: $(12-8)x = -14-(15) \implies 4x = -29 \implies x = \frac{-29}{4}$.

Q443. Transposing $10x$ to left and -19 to right: $(9-10)x = -3-(-19) \implies -1x = 16 \implies x = \frac{16}{-1}$.

Q444. Transposing $5x$ to left and -12 to right: $(10-5)x = -3-(-12) \implies 5x = 9 \implies x = \frac{9}{5}$.

Q445. Transposing $9x$ to left and -6 to right: $(10-9)x = -3-(-6) \implies 1x = 3 \implies x = \frac{3}{1}$.

Q446. Transposing $6x$ to left and -18 to right: $(12-6)x = -8-(-18) \implies 6x = 10 \implies x = \frac{10}{6}$.

Q447. Transposing $3x$ to left and 6 to right: $(6-3)x = 15-(6) \implies 3x = 9 \implies x = \frac{9}{3}$.

Q448. Transposing $10x$ to left and -18 to right: $(11-10)x = 1-(-18) \implies 1x = 19 \implies x = \frac{19}{1}$.

Q449. Transposing $3x$ to left and -3 to right: $(6-3)x = 15-(-3) \implies 3x = 18 \implies x = \frac{18}{3}$.

Q450. Transposing $6x$ to left and 3 to right: $(7-6)x = 19-(3) \implies 1x = 16 \implies x = \frac{16}{1}$.

Q451. Transposing $4x$ to left and -17 to right: $(11-4)x = -19-(-17) \implies 7x = -2 \implies x = \frac{-2}{7}$.

Q452. Transposing $9x$ to left and 19 to right: $(10-9)x = 4-(19) \implies 1x = -15 \implies x = \frac{-15}{1}$.

Q453. Transposing $8x$ to left and -19 to right: $(7-8)x = -8-(-19) \implies -1x = 11 \implies x = \frac{11}{-1}$.

Q454. Transposing $6x$ to left and -14 to right: $(7-6)x = 9-(-14) \implies 1x = 23 \implies x = \frac{23}{1}$.

Q455. Transposing $6x$ to left and -6 to right: $(8-6)x = -5-(-6) \implies 2x = 1 \implies x = \frac{1}{2}$.

Q456. Transposing $8x$ to left and 8 to right: $(9-8)x = 10-(8) \implies 1x = 2 \implies x = \frac{2}{1}$.

Q457. Transposing $2x$ to left and 2 to right: $(9-2)x = 11-(2) \implies 7x = 9 \implies x = \frac{9}{7}$.

Q458. Transposing $8x$ to left and -9 to right: $(11-8)x = -13-(-9) \implies 3x = -4 \implies x = \frac{-4}{3}$.

Q459. Transposing $3x$ to left and 1 to right: $(10-3)x = 9-(1) \implies 7x = 8 \implies x = \frac{8}{7}$.

Q460. Transposing $5x$ to left and 0 to right: $(12-5)x = -4-(0) \implies 7x = -4 \implies x = \frac{-4}{7}$.

Q461. Transposing $5x$ to left and -3 to right: $(9-5)x = -10-(-3) \implies 4x = -7 \implies x = \frac{-7}{4}$.

Q462. Transposing $8x$ to left and -20 to right: $(4-8)x = 0-(-20) \implies -4x = 20 \implies x = \frac{20}{-4}$.

Q463. Transposing $8x$ to left and -13 to right: $(9-8)x = -2-(-13) \implies 1x = 11 \implies x = \frac{11}{1}$.

Q464. Transposing $3x$ to left and -5 to right: $(9-3)x = -5-(-5) \implies 6x = 0 \implies x = \frac{0}{6}$.

Q465. Transposing $3x$ to left and -11 to right: $(8-3)x = 11-(-11) \implies 5x = 22 \implies x = \frac{22}{5}$.

Q466. Transposing $4x$ to left and 0 to right: $(7-4)x = 6-(0) \implies 3x = 6 \implies x = \frac{6}{3}$.

Q467. Transposing $6x$ to left and 0 to right: $(11-6)x = -6-(0) \implies 5x = -6 \implies x = \frac{-6}{5}$.

Q468. Transposing $6x$ to left and -19 to right: $(7-6)x = -5-(-19) \implies 1x = 14 \implies x = \frac{14}{1}$.

Q469. Transposing $3x$ to left and 3 to right: $(11-3)x = 20-(3) \implies 8x = 17 \implies x = \frac{17}{8}$.

Q470. Transposing $7x$ to left and -9 to right: $(11-7)x = 16-(-9) \implies 4x = 25 \implies x = \frac{25}{4}$.

Q471. Transposing $7x$ to left and -4 to right: $(8-7)x = -20-(-4) \implies 1x = -16 \implies x = \frac{-16}{1}$.

Q472. Transposing $3x$ to left and -14 to right: $(9-3)x = 20-(-14) \implies 6x = 34 \implies x = \frac{34}{6}$.

Q473. Transposing $4x$ to left and 19 to right: $(6-4)x = 19-(19) \implies 2x = 0 \implies x = \frac{0}{2}$.

Q474. Transposing $10x$ to left and 11 to right: $(3-10)x = 16-(11) \implies -7x = 5 \implies x = \frac{5}{-7}$.

Q475. Transposing $6x$ to left and 15 to right: $(12-6)x = -5-(15) \implies 6x = -20 \implies x = \frac{-20}{6}$.

Q476. Transposing $8x$ to left and -15 to right: $(4-8)x = -9-(-15) \implies -4x = 6 \implies x = \frac{6}{-4}$.

Q477. Transposing $5x$ to left and -13 to right: $(9-5)x = 16-(-13) \implies 4x = 29 \implies x = \frac{29}{4}$.

Q478. Transposing $2x$ to left and -8 to right: $(9-2)x = -1-(-8) \implies 7x = 7 \implies x = \frac{7}{7}$.

Q479. Transposing $10x$ to left and -13 to right: $(8-10)x = 18-(-13) \implies -2x = 31 \implies x = \frac{31}{-2}$.

Q480. Transposing $7x$ to left and -17 to right: $(10-7)x = 17-(-17) \implies 3x = 34 \implies x = \frac{34}{3}$.

Q481. Transposing $8x$ to left and 19 to right: $(11-8)x = -17-(19) \implies 3x = -36 \implies x = \frac{-36}{3}$.

Q482. Transposing $9x$ to left and -7 to right: $(10-9)x = 10-(-7) \implies 1x = 17 \implies x = \frac{17}{1}$.

Q483. Transposing $7x$ to left and 12 to right: $(6-7)x = 7-(12) \implies -1x = -5 \implies x = \frac{-5}{-1}$.

Q484. Transposing $3x$ to left and -2 to right: $(8-3)x = 4-(-2) \implies 5x = 6 \implies x = \frac{6}{5}$.

Q485. Transposing $8x$ to left and -9 to right: $(9-8)x = 11-(-9) \implies 1x = 20 \implies x = \frac{20}{1}$.

Q486. Transposing $3x$ to left and 16 to right: $(5-3)x = 8-(16) \implies 2x = -8 \implies x = \frac{-8}{2}$.

Q487. Transposing $6x$ to left and -4 to right: $(7-6)x = 17-(-4) \implies 1x = 21 \implies x = \frac{21}{1}$.

Q488. Transposing $2x$ to left and 0 to right: $(4-2)x = 5-(0) \implies 2x = 5 \implies x = \frac{5}{2}$.

Q489. Transposing $5x$ to left and 12 to right: $(12-5)x = -14-(12) \implies 7x = -26 \implies x = \frac{-26}{7}$.

Q490. Transposing $5x$ to left and 17 to right: $(6-5)x = -1-(17) \implies 1x = -18 \implies x = \frac{-18}{1}$.

Q491. Transposing $2x$ to left and 20 to right: $(9-2)x = -11-(20) \implies 7x = -31 \implies x = \frac{-31}{7}$.

Q492. Transposing $8x$ to left and -7 to right: $(9-8)x = 11-(-7) \implies 1x = 18 \implies x = \frac{18}{1}$.

Q493. Transposing $4x$ to left and 7 to right: $(6-4)x = -3-(7) \implies 2x = -10 \implies x = \frac{-10}{2}$.

Q494. Transposing $6x$ to left and 8 to right: $(10-6)x = -14-(8) \implies 4x = -22 \implies x = \frac{-22}{4}$.

Q495. Transposing $9x$ to left and 6 to right: $(10-9)x = 9-(6) \implies 1x = 3 \implies x = \frac{3}{1}$.

Q496. Transposing $8x$ to left and -4 to right: $(3-8)x = 8-(-4) \implies -5x = 12 \implies x = \frac{12}{-5}$.

Q497. Transposing $4x$ to left and -7 to right: $(11-4)x = -9-(-7) \implies 7x = -2 \implies x = \frac{-2}{7}$.

Q498. Transposing $3x$ to left and -9 to right: $(12-3)x = -16-(-9) \implies 9x = -7 \implies x = \frac{-7}{9}$.

Q499. Transposing $3x$ to left and -14 to right: $(12-3)x = -1-(-14) \implies 9x = 13 \implies x = \frac{13}{9}$.

Q500. Transposing $8x$ to left and 15 to right: $(9-8)x = -12-(15) \implies 1x = -27 \implies x = \frac{-27}{1}$.

Q501. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(6y-(-7y)) = 43-(-4) \implies 13y = 47 \implies y = \frac{47}{13}$.

Q502. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-2y)) = 33-(17) \implies 7y = 16 \implies y = \frac{16}{7}$.

Q503. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-4y)) = 10-(1) \implies 9y = 9 \implies y = \frac{9}{9}$.

Q504. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-6y)) = 28-(9) \implies 11y = 19 \implies y = \frac{19}{11}$.

Q505. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-5y)) = 35-(-16) \implies 10y = 51 \implies y = \frac{51}{10}$.

Q506. Subtracting Eq(2) changes signs of Eq(2): $(2x-2x)+(3y-(-1y)) = 43-(11) \implies 4y = 32 \implies y = \frac{32}{4}$.

Q507. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(6y-(-2y)) = 25-(1) \implies 8y = 24 \implies y = \frac{24}{8}$.

Q508. Subtracting Eq(2) changes signs of Eq(2): $(6x-6x)+(2y-(-5y)) = 46-(-11) \implies 7y = 57 \implies y = \frac{57}{7}$.

Q509. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(4y-(-7y)) = 28-(-5) \implies 11y = 33 \implies y = \frac{33}{11}$.

Q510. Subtracting Eq(2) changes signs of Eq(2): $(6x-6x)+(4y-(-4y)) = 15-(-19) \implies 8y = 34 \implies y = \frac{34}{8}$.

Q511. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(7y-(-3y)) = 12-(3) \implies 10y = 9 \implies y = \frac{9}{10}$.

Q512. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(5y-(-5y)) = 22-(7) \implies 10y = 15 \implies y = \frac{15}{10}$.

Q513. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(7y-(-1y)) = 23-(-6) \implies 8y = 29 \implies y = \frac{29}{8}$.

Q514. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(3y-(-7y)) = 38-(-5) \implies 10y = 43 \implies y = \frac{43}{10}$.

Q515. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(4y-(-5y)) = 16-(7) \implies 9y = 9 \implies y = \frac{9}{9}$.

Q516. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-6y)) = 38-(-11) \implies 11y = 49 \implies y = \frac{49}{11}$.

Q517. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(2y-(-1y)) = 35-(-4) \implies 3y = 39 \implies y = \frac{39}{3}$.

Q518. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(3y-(-6y)) = 27-(-18) \implies 9y = 45 \implies y = \frac{45}{9}$.

Q519. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(2y-(-1y)) = 35-(-12) \implies 3y = 47 \implies y = \frac{47}{3}$.

Q520. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-1y)) = 46-(9) \implies 6y = 37 \implies y = \frac{37}{6}$.

Q521. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(2y-(-7y)) = 37-(5) \implies 9y = 32 \implies y = \frac{32}{9}$.

Q522. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(2y-(-4y)) = 43-(10) \implies 6y = 33 \implies y = \frac{33}{6}$.

Q523. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(4y-(-6y)) = 34-(7) \implies 10y = 27 \implies y = \frac{27}{10}$.

Q524. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(6y-(-1y)) = 19-(-16) \implies 7y = 35 \implies y = \frac{35}{7}$.

Q525. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(2y-(-1y)) = 11-(-10) \implies 3y = 21 \implies y = \frac{21}{3}$.

Q526. Subtracting Eq(2) changes signs of Eq(2): $(6x-6x)+(2y-(-4y)) = 40-(-13) \implies 6y = 53 \implies y = \frac{53}{6}$.

Q527. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(4y-(-6y)) = 39-(-14) \implies 10y = 53 \implies y = \frac{53}{10}$.

Q528. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(7y-(-4y)) = 24-(5) \implies 11y = 19 \implies y = \frac{19}{11}$.

Q529. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(4y-(-2y)) = 19-(6) \implies 6y = 13 \implies y = \frac{13}{6}$.

Q530. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(6y-(-3y)) = 22-(6) \implies 9y = 16 \implies y = \frac{16}{9}$.

Q531. Subtracting Eq(2) changes signs of Eq(2): $(6x-6x)+(5y-(-5y)) = 34-(7) \implies 10y = 27 \implies y = \frac{27}{10}$.

Q532. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(2y-(-6y)) = 10-(-9) \implies 8y = 19 \implies y = \frac{19}{8}$.

Q533. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(3y-(-4y)) = 42-(-11) \implies 7y = 53 \implies y = \frac{53}{7}$.

Q534. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(4y-(-1y)) = 15-(-13) \implies 5y = 28 \implies y = \frac{28}{5}$.

Q535. Subtracting Eq(2) changes signs of Eq(2): $(2x-2x)+(2y-(-3y)) = 38-(11) \implies 5y = 27 \implies y = \frac{27}{5}$.

Q536. Subtracting Eq(2) changes signs of Eq(2): $(4x-4x)+(6y-(-1y)) = 19-(-13) \implies 7y = 32 \implies y = \frac{32}{7}$.

Q537. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(2y-(-2y)) = 37-(15) \implies 4y = 22 \implies y = \frac{22}{4}$.

Q538. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(7y-(-4y)) = 50-(-20) \implies 11y = 70 \implies y = \frac{70}{11}$.

Q539. Subtracting Eq(2) changes signs of Eq(2): $(6x-6x)+(5y-(-7y)) = 26-(-12) \implies 12y = 38 \implies y = \frac{38}{12}$.

Q540. Subtracting Eq(2) changes signs of Eq(2): $(3x-3x)+(5y-(-3y)) = 27-(6) \implies 8y = 21 \implies y = \frac{21}{8}$.

Q541. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(7y-(-5y)) = 31-(4) \implies 12y = 27 \implies y = \frac{27}{12}$.

Q542. Subtracting Eq(2) changes signs of Eq(2): $(2x-2x)+(3y-(-1y)) = 38-(7) \implies 4y = 31 \implies y = \frac{31}{4}$.

Q543. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(4y-(-3y)) = 50-(11) \implies 7y = 39 \implies y = \frac{39}{7}$.

Q544. Subtracting Eq(2) changes signs of Eq(2): $(5x-5x)+(7y-(-3y)) = 35-(-8) \implies 10y = 43 \implies y = \frac{43}{10}$.

Q545. Subtracting Eq(2) changes signs of Eq(2): $(4x-$

$4x)+(6y-(-1y)) = 19-(-19) \implies 7y = 38 \implies y = \frac{38}{7}$.

Q546. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(6y-(-5y)) = 15-(16) \implies 11y = -1 \implies y = \frac{-1}{11}$.

Q547. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (2y - (-5y)) = 22 - (-2) \implies 7y = 24 \implies y = \frac{24}{7}$.

Q548. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (6y - (-1y)) = 23 - (-4) \implies 7y = 27 \implies y = \frac{27}{7}$.

Q549. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(7y-(-2y)) = 16-(18) \implies 9y = -2 \implies y = \frac{-2}{9}$.

Q550. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (2y - (-3y)) = 34 - (4) \implies 5y = 30 \implies y = \frac{30}{5}$.

Q551. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(2y-(-7y)) = 31-(-16) \implies 9y = 47 \implies y = \frac{47}{9}$.

Q552. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(7y-(-6y)) = 28-(18) \implies 13y = 10 \implies y = \frac{10}{13}$.

Q553. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (6y - (-3y)) = 47 - (-8) \implies 9y = 55 \implies y = \frac{55}{9}$.

Q554. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(4y-(-2y)) = 48-(-14) \implies 6y = 62 \implies y = \frac{62}{6}$.

Q555. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(7y-(-7y)) = 25-(-5) \implies 14y = 30 \implies y = \frac{30}{14}$.

Q556. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(2y-(-4y)) = 47-(-13) \implies 6y = 60 \implies y = \frac{60}{6}$.

Q557. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (3y - (-4y)) = 34 - (15) \implies 7y = 19 \implies y = \frac{19}{7}$.

Q558. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(5y-(-1y)) = 23-(-19) \implies 6y = 42 \implies y = \frac{42}{6}$.

Q559. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (4y - (-7y)) = 18 - (9) \implies 11y = 9 \implies y = \frac{9}{11}$.

Q560. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (5y - (-5y)) = 23 - (3) \implies 10y = 20 \implies y = \frac{20}{10}$.

Q561. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(7y-(-1y)) = 21-(-13) \implies 8y = 34 \implies y = \frac{34}{8}$.

Q562. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (5y - (-6y)) = 31 - (1) \implies 11y = 30 \implies y = \frac{30}{11}$.

Q563. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(4y-(-7y)) = 28-(-5) \implies 11y = 33 \implies y = \frac{33}{11}$.

Q564. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(6y-(-1y)) = 23-(-15) \implies 7y = 38 \implies y = \frac{38}{7}$.

Q565. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (7y - (-1y)) = 23 - (11) \implies 8y = 12 \implies y = \frac{12}{8}$.

Q566. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(7y-(-4y)) = 34-(16) \implies 11y = 18 \implies y = \frac{18}{11}$.

Q567. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (3y - (-1y)) = 18 - (-5) \implies 4y = 23 \implies y = \frac{23}{4}$.

Q568. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (4y - (-6y)) = 14 - (-14) \implies 10y = 28 \implies y = \frac{28}{10}$.

Q569. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (7y - (-2y)) = 30 - (11) \implies 9y = 19 \implies y = \frac{19}{9}$.

Q570. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(4y-(-1y)) = 41-(-17) \implies 5y = 58 \implies y = \frac{58}{5}$.

Q571. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(4y-(-4y)) = 11-(5) \implies 8y = 6 \implies y = \frac{6}{8}$.

Q572. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x)+(3y-(-1y)) = 37-(-18) \implies 4y = 55 \implies y = \frac{55}{4}$.

Q573. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (2y - (-2y)) = 35 - (6) \implies 4y = 29 \implies y = \frac{29}{4}$.

Q574. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (4y - (-2y)) = 36 - (0) \implies 6y = 36 \implies y = \frac{36}{6}$.

Q575. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (7y - (-4y)) = 11 - (1) \implies 11y = 10 \implies y = \frac{10}{11}$.

Q576. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(2y-(-5y)) = 12-(-11) \implies 7y = 23 \implies y = \frac{23}{7}$.

Q577. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(4y-(-2y)) = 23-(-11) \implies 6y = 34 \implies y = \frac{34}{6}$.

Q578. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x)+(3y-(-5y)) = 48-(-17) \implies 8y = 65 \implies y = \frac{65}{8}$.

Q579. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (7y - (-1y)) = 33 - (11) \implies 8y = 22 \implies y = \frac{22}{8}$.

Q580. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (7y - (-5y)) = 25 - (9) \implies 12y = 16 \implies y = \frac{16}{12}$.

Q581. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(7y-(-4y)) = 29-(-2) \implies 11y = 31 \implies y = \frac{31}{11}$.

Q582. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(3y-(-1y)) = 45-(-13) \implies 4y = 58 \implies y = \frac{58}{4}$.

Q583. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (7y - (-3y)) = 39 - (8) \implies 10y = 31 \implies y = \frac{31}{10}$.

Q584. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (4y - (-1y)) = 46 - (15) \implies 5y = 31 \implies y = \frac{31}{5}$.

Q585. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (4y - (-4y)) = 24 - (9) \implies 8y = 15 \implies y = \frac{15}{8}$.

Q586. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (5y - (-1y)) = 42 - (-6) \implies 6y = 48 \implies y = \frac{48}{6}$.

Q587. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (7y - (-1y)) = 48 - (-9) \implies 8y = 57 \implies y = \frac{57}{8}$.

Q588. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (4y - (-2y)) = 11 - (1) \implies 6y = 10 \implies y = \frac{10}{6}$.

Q589. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(4y-(-2y)) = 27-(-12) \implies 6y = 39 \implies y = \frac{39}{6}$.

Q590. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (3y - (-2y)) = 27 - (-3) \implies 5y = 30 \implies y = \frac{30}{5}$.

Q591. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (7y - (-4y)) = 29 - (1) \implies 11y = 28 \implies y = \frac{28}{11}$.

Q592. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(5y-(-5y)) = 38-(14) \implies 10y = 24 \implies y = \frac{24}{10}$.

Q593. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (5y - (-1y)) = 39 - (15) \implies 6y = 24 \implies y = \frac{24}{6}$.

Q594. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (3y - (-5y)) = 25 - (1) \implies 8y = 24 \implies y = \frac{24}{8}$.

Q595. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (3y - (-7y)) = 26 - (-16) \implies 10y = 42 \implies y = \frac{42}{10}$.

Q596. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (3y - (-6y)) = 12 - (-9) \implies 9y = 21 \implies y = \frac{21}{9}$.

Q597. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(7y-(-4y)) = 34-(-1) \implies 11y = 35 \implies y = \frac{35}{11}$.

Q598. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (5y - (-5y)) = 45 - (7) \implies 10y = 38 \implies y = \frac{38}{10}$.

Q599. Subtracting Eq(2) changes signs of Eq(2): $(6x -$

$6x)+(5y-(-4y)) = 14-(-19) \implies 9y = 33 \implies y = \frac{33}{9}$.

Q600. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(7y-(-7y)) = 40-(16) \implies 14y = 24 \implies y = \frac{24}{14}$.

Q601. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(5y-(-7y)) = 27-(-7) \implies 12y = 34 \implies y = \frac{34}{12}$.

Q602. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (3y - (-4y)) = 25 - (6) \implies 7y = 19 \implies y = \frac{19}{7}$.

Q603. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(7y-(-4y)) = 34-(13) \implies 11y = 21 \implies y = \frac{21}{11}$.

Q604. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(6y-(-2y)) = 21-(-14) \implies 8y = 35 \implies y = \frac{35}{8}$.

Q605. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (4y - (-1y)) = 12 - (2) \implies 5y = 10 \implies y = \frac{10}{5}$.

Q606. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(6y-(-3y)) = 32-(-10) \implies 9y = 42 \implies y = \frac{42}{9}$.

Q607. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(5y-(-6y)) = 29-(11) \implies 11y = 18 \implies y = \frac{18}{11}$.

Q608. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (5y - (-1y)) = 47 - (-2) \implies 6y = 49 \implies y = \frac{49}{6}$.

Q609. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x)+(6y-(-2y)) = 30-(-15) \implies 8y = 45 \implies y = \frac{45}{8}$.

Q610. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(2y-(-1y)) = 17-(-13) \implies 3y = 30 \implies y = \frac{30}{3}$.

Q611. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (2y - (-3y)) = 36 - (9) \implies 5y = 27 \implies y = \frac{27}{5}$.

Q612. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (3y - (-2y)) = 31 - (5) \implies 5y = 26 \implies y = \frac{26}{5}$.

Q613. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (7y - (-6y)) = 20 - (2) \implies 13y = 18 \implies y = \frac{18}{13}$.

Q614. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(3y-(-2y)) = 13-(-14) \implies 5y = 27 \implies y = \frac{27}{5}$.

Q615. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (5y - (-4y)) = 42 - (15) \implies 9y = 27 \implies y = \frac{27}{9}$.

Q616. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (2y - (-5y)) = 50 - (-7) \implies 7y = 57 \implies y = \frac{57}{7}$.

Q617. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (2y - (-4y)) = 24 - (15) \implies 6y = 9 \implies y = \frac{9}{6}$.

Q618. Subtracting Eq(2) changes signs of Eq(2): $(7x-7x)+(5y-(-4y)) = 12-(6) \implies 9y = 6 \implies y = \frac{6}{9}$.

Q619. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (7y - (-2y)) = 32 - (14) \implies 9y = 18 \implies y = \frac{18}{9}$.

Q620. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (7y - (-2y)) = 50 - (6) \implies 9y = 44 \implies y = \frac{44}{9}$.

Q621. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(5y-(-4y)) = 17-(18) \implies 9y = -1 \implies y = \frac{-1}{9}$.

Q622. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x)+(5y-(-6y)) = 34-(-1) \implies 11y = 35 \implies y = \frac{35}{11}$.

Q623. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x)+(6y-(-5y)) = 26-(13) \implies 11y = 13 \implies y = \frac{13}{11}$.

Q624. Subtracting Eq(2) changes signs of Eq(2): $(3x - 3x) + (3y - (-7y)) = 19 - (-19) \implies 10y = 38 \implies y = \frac{38}{10}$.

Q625. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(4y-(-6y)) = 25-(-3) \implies 10y = 28 \implies y = \frac{28}{10}$.

Q626. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(3y-(-4y)) = 30-(-18) \implies 7y = 48 \implies y = \frac{48}{7}$.

Q627. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (3y - (-5y)) = 14 - (-8) \implies 8y = 22 \implies y = \frac{22}{8}$.

Q628. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (2y - (-6y)) = 22 - (8) \implies 8y = 14 \implies y = \frac{14}{8}$.

Q629. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (5y - (-7y)) = 50 - (-20) \implies 12y = 70 \implies y = \frac{70}{12}$.

Q630. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x) + (5y - (-5y)) = 15 - (-20) \implies 10y = 35 \implies y = \frac{35}{10}$.

Q631. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(6y-(-5y)) = 29-(-1) \implies 11y = 30 \implies y = \frac{30}{11}$.

Q632. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (4y - (-3y)) = 31 - (9) \implies 7y = 22 \implies y = \frac{22}{7}$.

Q633. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(7y-(-6y)) = 28-(15) \implies 13y = 13 \implies y = \frac{13}{13}$.

Q634. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (3y - (-1y)) = 47 - (17) \implies 4y = 30 \implies y = \frac{30}{4}$.

Q635. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (3y - (-4y)) = 48 - (5) \implies 7y = 43 \implies y = \frac{43}{7}$.

Q636. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (4y - (-1y)) = 29 - (6) \implies 5y = 23 \implies y = \frac{23}{5}$.

Q637. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(6y-(-6y)) = 47-(-6) \implies 12y = 53 \implies y = \frac{53}{12}$.

Q638. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(3y-(-1y)) = 24-(-13) \implies 4y = 37 \implies y = \frac{37}{4}$.

Q639. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x)+(7y-(-3y)) = 15-(-2) \implies 10y = 17 \implies y = \frac{17}{10}$.

Q640. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x)+(2y-(-7y)) = 14-(-19) \implies 9y = 33 \implies y = \frac{33}{9}$.

Q641. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (5y - (-2y)) = 19 - (14) \implies 7y = 5 \implies y = \frac{5}{7}$.

Q642. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (4y - (-4y)) = 46 - (-5) \implies 8y = 51 \implies y = \frac{51}{8}$.

Q643. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(5y-(-5y)) = 12-(18) \implies 10y = -6 \implies y = \frac{-6}{10}$.

Q644. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x)+(4y-(-2y)) = 50-(-19) \implies 6y = 69 \implies y = \frac{69}{6}$.

Q645. Subtracting Eq(2) changes signs of Eq(2): $(6x - 6x) + (6y - (-4y)) = 50 - (6) \implies 10y = 44 \implies y = \frac{44}{10}$.

Q646. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (2y - (-4y)) = 40 - (-5) \implies 6y = 45 \implies y = \frac{45}{6}$.

Q647. Subtracting Eq(2) changes signs of Eq(2): $(4x - 4x)+(2y-(-5y)) = 11-(-20) \implies 7y = 31 \implies y = \frac{31}{7}$.

Q648. Subtracting Eq(2) changes signs of Eq(2): $(5x - 5x) + (6y - (-1y)) = 48 - (15) \implies 7y = 33 \implies y = \frac{33}{7}$.

Q649. Subtracting Eq(2) changes signs of Eq(2): $(7x - 7x) + (6y - (-5y)) = 16 - (4) \implies 11y = 12 \implies y = \frac{12}{11}$.

Q650. Subtracting Eq(2) changes signs of Eq(2): $(2x - 2x) + (5y - (-1y)) = 21 - (15) \implies 6y = 6 \implies y = \frac{6}{6}$.

Q651. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 9^2 + 40 = 121$. So $x = \frac{9 \pm \sqrt{121}}{2}$.

Q652. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-5) = +40$. Thus $D = 7^2 + 40 = 89$. So $x = \frac{7 \pm \sqrt{89}}{2}$.

Q653. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-3) = +12$. Thus $D = 3^2 + 12 = 21$. So $x = \frac{3 \pm \sqrt{21}}{2}$.

Q654. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-11) = +88$. Thus $D = 9^2 + 88 = 169$. So $x = \frac{9 \pm \sqrt{169}}{2}$.

Q655. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-12) = +144$. Thus $D = 3^2 + 144 = 153$. So $x = \frac{3 \pm \sqrt{153}}{2}$.

Q656. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-1) = +4$. Thus $D = 10^2 + 4 = 104$. So $x = \frac{10 \pm \sqrt{104}}{2}$.

Q657. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-11) = +88$. Thus $D = 6^2 + 88 = 124$. So $x = \frac{6 \pm \sqrt{124}}{2}$.

Q658. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-6) = +48$. Thus $D = 2^2 + 48 = 52$. So $x = \frac{2 \pm \sqrt{52}}{2}$.

Q659. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-10) = +40$. Thus $D = 9^2 + 40 = 121$. So $x = \frac{9 \pm \sqrt{121}}{2}$.

Q660. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-1) = +16$. Thus $D = 10^2 + 16 = 116$. So $x = \frac{10 \pm \sqrt{116}}{2}$.

Q661. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-9) = +72$. Thus $D = 9^2 + 72 = 153$. So $x = \frac{9 \pm \sqrt{153}}{2}$.

Q662. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-10) = +200$. Thus $D = 10^2 + 200 = 300$. So $x = \frac{10 \pm \sqrt{300}}{2}$.

Q663. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-3) = +60$. Thus $D = 1^2 + 60 = 61$. So $x = \frac{1 \pm \sqrt{61}}{2}$.

Q664. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-8) = +96$. Thus $D = 6^2 + 96 = 132$. So $x = \frac{6 \pm \sqrt{132}}{2}$.

Q665. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-3) = +24$. Thus $D = 6^2 + 24 = 60$. So $x = \frac{6 \pm \sqrt{60}}{2}$.

Q666. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-4) = +48$. Thus $D = 7^2 + 48 = 97$. So $x = \frac{7 \pm \sqrt{97}}{2}$.

Q667. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-8) = +128$. Thus $D = 5^2 + 128 = 153$. So $x = \frac{5 \pm \sqrt{153}}{2}$.

Q668. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-12) = +144$. Thus $D = 3^2 + 144 = 153$. So $x = \frac{3 \pm \sqrt{153}}{2}$.

Q669. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-7) = +84$. Thus $D = 10^2 + 84 = 184$. So $x = \frac{10 \pm \sqrt{184}}{2}$.

Q670. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-2) = +32$. Thus $D = 7^2 + 32 = 81$. So $x = \frac{7 \pm \sqrt{81}}{2}$.

Q671. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-9) = +108$. Thus $D = 2^2 + 108 = 112$. So $x = \frac{2 \pm \sqrt{112}}{2}$.

Q672. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-9) = +108$. Thus $D = 5^2 + 108 = 133$. So $x = \frac{5 \pm \sqrt{133}}{2}$.

Q673. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-12) = +96$. Thus $D = 4^2 + 96 = 112$. So $x = \frac{4 \pm \sqrt{112}}{2}$.

Q674. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-2) = +8$. Thus $D = 8^2 + 8 = 72$. So $x = \frac{8 \pm \sqrt{72}}{2}$.

Q675. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-10) = +200$. Thus $D = 8^2 + 200 = 264$. So $x = \frac{8 \pm \sqrt{264}}{2}$.

Q676. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-10) = +160$. Thus $D = 1^2 + 160 = 161$. So $x = \frac{1 \pm \sqrt{161}}{2}$.

Q677. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-2) = +8$. Thus $D = 5^2 + 8 = 33$. So $x = \frac{5 \pm \sqrt{33}}{2}$.

Q678. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-5) = +100$. Thus $D = 10^2 + 100 = 200$. So $x = \frac{10 \pm \sqrt{200}}{2}$.

Q679. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-12) = +144$. Thus $D = 10^2 + 144 = 244$. So $x = \frac{10 \pm \sqrt{244}}{2}$.

Q680. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-11) = +44$. Thus $D = 2^2 + 44 = 48$. So $x = \frac{2 \pm \sqrt{48}}{2}$.

Q681. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 1^2 + 32 = 33$. So $x = \frac{1 \pm \sqrt{33}}{2}$.

Q682. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-12) = +48$. Thus $D = 2^2 + 48 = 52$. So $x = \frac{2 \pm \sqrt{52}}{2}$.

Q683. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-6) = +120$. Thus $D = 7^2 + 120 = 169$. So $x = \frac{7 \pm \sqrt{169}}{2}$.

Q684. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-12) = +144$. Thus $D = 1^2 + 144 = 145$. So $x = \frac{1 \pm \sqrt{145}}{2}$.

Q685. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-8) = +160$. Thus $D = 9^2 + 160 = 241$. So $x = \frac{9 \pm \sqrt{241}}{2}$.

Q686. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-2) = +16$. Thus $D = 6^2 + 16 = 52$. So $x = \frac{6 \pm \sqrt{52}}{2}$.

Q687. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-8) = +128$. Thus $D = 9^2 + 128 = 209$. So $x = \frac{9 \pm \sqrt{209}}{2}$.

Q688. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-4) = +32$. Thus $D = 4^2 + 32 = 48$. So $x = \frac{4 \pm \sqrt{48}}{2}$.

Q689. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-2) = +32$. Thus $D = 5^2 + 32 = 57$. So $x = \frac{5 \pm \sqrt{57}}{2}$.

Q690. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-12) = +192$. Thus $D = 6^2 + 192 = 228$. So $x = \frac{6 \pm \sqrt{228}}{2}$.

Q691. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-1) = +12$. Thus $D = 8^2 + 12 = 76$. So $x = \frac{8 \pm \sqrt{76}}{2}$.

Q692. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-12) = +240$. Thus $D = 3^2 + 240 = 249$. So $x = \frac{3 \pm \sqrt{249}}{2}$.

Q693. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-1) = +8$. Thus $D = 3^2 + 8 = 17$. So $x = \frac{3 \pm \sqrt{17}}{2}$.

Q694. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-9) = +144$. Thus $D = 6^2 + 144 = 180$. So $x = \frac{6 \pm \sqrt{180}}{2}$.

Q695. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 10^2 + 32 = 132$. So $x = \frac{10 \pm \sqrt{132}}{2}$.

Q696. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-8) = +128$. Thus $D = 10^2 + 128 = 228$. So $x = \frac{10 \pm \sqrt{228}}{2}$.

Q697. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-9) = +72$. Thus $D = 2^2 + 72 = 76$. So $x = \frac{2\pm\sqrt{76}}{2}$.

Q698. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-11) = +176$. Thus $D = 3^2 + 176 = 185$. So $x = \frac{3\pm\sqrt{185}}{2}$.

Q699. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-9) = +180$. Thus $D = 10^2 + 180 = 280$. So $x = \frac{10\pm\sqrt{280}}{2}$.

Q700. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-3) = +48$. Thus $D = 5^2 + 48 = 73$. So $x = \frac{5\pm\sqrt{73}}{2}$.

Q701. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-8) = +64$. Thus $D = 8^2 + 64 = 128$. So $x = \frac{8\pm\sqrt{128}}{2}$.

Q702. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-10) = +40$. Thus $D = 6^2+40 = 76$. So $x = \frac{6\pm\sqrt{76}}{2}$.

Q703. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 10^2 + 32 = 132$. So $x = \frac{10\pm\sqrt{132}}{2}$.

Q704. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-6) = +96$. Thus $D = 7^2 + 96 = 145$. So $x = \frac{7\pm\sqrt{145}}{2}$.

Q705. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 6^2 + 56 = 92$. So $x = \frac{6\pm\sqrt{92}}{2}$.

Q706. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 8^2 + 40 = 104$. So $x = \frac{8\pm\sqrt{104}}{2}$.

Q707. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-7) = +84$. Thus $D = 4^2 + 84 = 100$. So $x = \frac{4\pm\sqrt{100}}{2}$.

Q708. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-9) = +36$. Thus $D = 2^2 + 36 = 40$. So $x = \frac{2\pm\sqrt{40}}{2}$.

Q709. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-11) = +88$. Thus $D = 6^2 + 88 = 124$. So $x = \frac{6\pm\sqrt{124}}{2}$.

Q710. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-10) = +160$. Thus $D = 3^2 + 160 = 169$. So $x = \frac{3\pm\sqrt{169}}{2}$.

Q711. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-7) = +28$. Thus $D = 9^2 + 28 = 109$. So $x = \frac{9\pm\sqrt{109}}{2}$.

Q712. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-11) = +176$. Thus $D = 10^2 + 176 = 276$. So $x = \frac{10\pm\sqrt{276}}{2}$.

Q713. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-8) = +64$. Thus $D = 6^2 + 64 = 100$. So $x = \frac{6\pm\sqrt{100}}{2}$.

Q714. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-8) = +160$. Thus $D = 9^2 + 160 = 241$. So $x = \frac{9\pm\sqrt{241}}{2}$.

Q715. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 4^2 + 40 = 56$. So $x = \frac{4\pm\sqrt{56}}{2}$.

Q716. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-12) = +96$. Thus $D = 9^2 + 96 = 177$. So $x = \frac{9\pm\sqrt{177}}{2}$.

Q717. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 5^2 + 56 = 81$. So $x = \frac{5\pm\sqrt{81}}{2}$.

Q718. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-10) = +40$. Thus $D = 1^2+40 = 41$. So $x = \frac{1\pm\sqrt{41}}{2}$.

Q719. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-3) = +12$. Thus $D = 1^2 + 12 = 13$. So $x = \frac{1\pm\sqrt{13}}{2}$.

Q720. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-7) = +28$. Thus $D = 5^2 + 28 = 53$. So $x = \frac{5\pm\sqrt{53}}{2}$.

Q721. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-6) = +72$. Thus $D = 8^2 + 72 = 136$. So $x = \frac{8\pm\sqrt{136}}{2}$.

Q722. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 9^2 + 40 = 121$. So $x = \frac{9\pm\sqrt{121}}{2}$.

Q723. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-10) = +160$. Thus $D = 5^2 + 160 = 185$. So $x = \frac{5\pm\sqrt{185}}{2}$.

Q724. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-1) = +12$. Thus $D = 5^2 + 12 = 37$. So $x = \frac{5\pm\sqrt{37}}{2}$.

Q725. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-5) = +80$. Thus $D = 3^2 + 80 = 89$. So $x = \frac{3\pm\sqrt{89}}{2}$.

Q726. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-12) = +48$. Thus $D = 7^2+48 = 97$. So $x = \frac{7\pm\sqrt{97}}{2}$.

Q727. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-2) = +24$. Thus $D = 7^2 + 24 = 73$. So $x = \frac{7\pm\sqrt{73}}{2}$.

Q728. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-10) = +200$. Thus $D = 1^2 + 200 = 201$. So $x = \frac{1\pm\sqrt{201}}{2}$.

Q729. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-4) = +64$. Thus $D = 2^2 + 64 = 68$. So $x = \frac{2\pm\sqrt{68}}{2}$.

Q730. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-3) = +60$. Thus $D = 1^2 + 60 = 61$. So $x = \frac{1\pm\sqrt{61}}{2}$.

Q731. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-4) = +48$. Thus $D = 3^2 + 48 = 57$. So $x = \frac{3\pm\sqrt{57}}{2}$.

Q732. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-8) = +128$. Thus $D = 2^2 + 128 = 132$. So $x = \frac{2\pm\sqrt{132}}{2}$.

Q733. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-4) = +32$. Thus $D = 10^2 + 32 = 132$. So $x = \frac{10\pm\sqrt{132}}{2}$.

Q734. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-8) = +64$. Thus $D = 10^2 + 64 = 164$. So $x = \frac{10\pm\sqrt{164}}{2}$.

Q735. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-4) = +16$. Thus $D = 6^2 + 16 = 52$. So $x = \frac{6\pm\sqrt{52}}{2}$.

Q736. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-4) = +16$. Thus $D = 3^2 + 16 = 25$. So $x = \frac{3\pm\sqrt{25}}{2}$.

Q737. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-12) = +192$. Thus $D = 3^2 + 192 = 201$. So $x = \frac{3\pm\sqrt{201}}{2}$.

Q738. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-8) = +160$. Thus $D = 10^2 + 160 = 260$. So $x = \frac{10\pm\sqrt{260}}{2}$.

Q739. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-1) = +12$. Thus $D = 1^2 + 12 = 13$. So $x = \frac{1\pm\sqrt{13}}{2}$.

Q740. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 7^2 + 40 = 89$. So $x = \frac{7\pm\sqrt{89}}{2}$.

Q741. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-6) = +96$. Thus $D = 2^2 + 96 = 100$. So $x = \frac{2\pm\sqrt{100}}{2}$.

Q742. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-10) = +40$. Thus $D = 4^2+40 = 56$. So $x = \frac{4\pm\sqrt{56}}{2}$.

Q743. Here $-b = -(-1) = +1$, and $-4ac =$

$-4(1)(-12) = +192$. Thus $D = 1^2 + 192 = 193$. So $x = \frac{1\pm\sqrt{193}}{2}$.

Q744. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 5^2 + 32 = 57$. So $x = \frac{5\pm\sqrt{57}}{2}$.

Q745. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-3) = +60$. Thus $D = 8^2 + 60 = 124$. So $x = \frac{8\pm\sqrt{124}}{2}$.

Q746. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-1) = +8$. Thus $D = 1^2 + 8 = 9$. So $x = \frac{1\pm\sqrt{9}}{2}$.

Q747. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 1^2 + 56 = 57$. So $x = \frac{1\pm\sqrt{57}}{2}$.

Q748. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-2) = +8$. Thus $D = 5^2 + 8 = 33$. So $x = \frac{5\pm\sqrt{33}}{2}$.

Q749. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-11) = +176$. Thus $D = 5^2 + 176 = 201$. So $x = \frac{5\pm\sqrt{201}}{2}$.

Q750. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-9) = +108$. Thus $D = 10^2 + 108 = 208$. So $x = \frac{10\pm\sqrt{208}}{2}$.

Q751. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-9) = +108$. Thus $D = 6^2 + 108 = 144$. So $x = \frac{6\pm\sqrt{144}}{2}$.

Q752. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-4) = +16$. Thus $D = 6^2 + 16 = 52$. So $x = \frac{6\pm\sqrt{52}}{2}$.

Q753. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-5) = +80$. Thus $D = 6^2 + 80 = 116$. So $x = \frac{6\pm\sqrt{116}}{2}$.

Q754. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-2) = +32$. Thus $D = 1^2 + 32 = 33$. So $x = \frac{1\pm\sqrt{33}}{2}$.

Q755. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-7) = +112$. Thus $D = 2^2 + 112 = 116$. So $x = \frac{2\pm\sqrt{116}}{2}$.

Q756. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 7^2 + 56 = 105$. So $x = \frac{7\pm\sqrt{105}}{2}$.

Q757. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-3) = +36$. Thus $D = 5^2 + 36 = 61$. So $x = \frac{5\pm\sqrt{61}}{2}$.

Q758. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-3) = +24$. Thus $D = 9^2 + 24 = 105$. So $x = \frac{9\pm\sqrt{105}}{2}$.

Q759. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-5) = +20$. Thus $D = 1^2 + 20 = 21$. So $x = \frac{1\pm\sqrt{21}}{2}$.

Q760. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-8) = +64$. Thus $D = 5^2 + 64 = 89$. So $x = \frac{5\pm\sqrt{89}}{2}$.

Q761. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-2) = +24$. Thus $D = 8^2 + 24 = 88$. So $x = \frac{8\pm\sqrt{88}}{2}$.

Q762. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 4^2 + 32 = 48$. So $x = \frac{4\pm\sqrt{48}}{2}$.

Q763. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-12) = +96$. Thus $D = 1^2+96 = 97$. So $x = \frac{1\pm\sqrt{97}}{2}$.

Q764. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-6) = +72$. Thus $D = 6^2 + 72 = 108$. So $x = \frac{6\pm\sqrt{108}}{2}$.

Q765. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-2) = +24$. Thus $D = 9^2 + 24 = 105$. So $x = \frac{9\pm\sqrt{105}}{2}$.

Q766. Here $-b = -(-2) = +2$, and $-4ac = -4(1)(-9) = +144$. Thus $D = 2^2 + 144 = 148$. So $x = \frac{2\pm\sqrt{148}}{2}$.

Q767. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-6) = +72$. Thus $D = 4^2 + 72 = 88$. So $x = \frac{4\pm\sqrt{88}}{2}$.

Q768. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-1) = +16$. Thus $D = 8^2 + 16 = 80$. So $x = \frac{8\pm\sqrt{80}}{2}$.

Q769. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-1) = +8$. Thus $D = 1^2 + 8 = 9$. So $x = \frac{1\pm\sqrt{9}}{2}$.

Q770. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-2) = +40$. Thus $D = 4^2 + 40 = 56$. So $x = \frac{4\pm\sqrt{56}}{2}$.

Q771. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-4) = +48$. Thus $D = 9^2 + 48 = 129$. So $x = \frac{9\pm\sqrt{129}}{2}$.

Q772. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-8) = +32$. Thus $D = 7^2 + 32 = 81$. So $x = \frac{7\pm\sqrt{81}}{2}$.

Q773. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-11) = +220$. Thus $D = 1^2 + 220 = 221$. So $x = \frac{1\pm\sqrt{221}}{2}$.

Q774. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-9) = +144$. Thus $D = 1^2 + 144 = 145$. So $x = \frac{1\pm\sqrt{145}}{2}$.

Q775. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-1) = +4$. Thus $D = 5^2 + 4 = 29$. So $x = \frac{5\pm\sqrt{29}}{2}$.

Q776. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-7) = +140$. Thus $D = 5^2 + 140 = 165$. So $x = \frac{5\pm\sqrt{165}}{2}$.

Q777. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-11) = +132$. Thus $D = 4^2 + 132 = 148$. So $x = \frac{4\pm\sqrt{148}}{2}$.

Q778. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-3) = +60$. Thus $D = 9^2 + 60 = 141$. So $x = \frac{9\pm\sqrt{141}}{2}$.

Q779. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 6^2 + 56 = 92$. So $x = \frac{6\pm\sqrt{92}}{2}$.

Q780. Here $-b = -(-4) = +4$, and $-4ac = -4(1)(-10) = +160$. Thus $D = 4^2 + 160 = 176$. So $x = \frac{4\pm\sqrt{176}}{2}$.

Q781. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-12) = +48$. Thus $D = 1^2+48 = 49$. So $x = \frac{1\pm\sqrt{49}}{2}$.

Q782. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-1) = +12$. Thus $D = 6^2 + 12 = 48$. So $x = \frac{6\pm\sqrt{48}}{2}$.

Q783. Here $-b = -(-1) = +1$, and $-4ac = -4(1)(-4) = +80$. Thus $D = 1^2 + 80 = 81$. So $x = \frac{1\pm\sqrt{81}}{2}$.

Q784. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-2) = +32$. Thus $D = 6^2 + 32 = 68$. So $x = \frac{6\pm\sqrt{68}}{2}$.

Q785. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-12) = +96$. Thus $D = 10^2 + 96 = 196$. So $x = \frac{10\pm\sqrt{196}}{2}$.

Q786. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-7) = +84$. Thus $D = 3^2 + 84 = 93$. So $x = \frac{3\pm\sqrt{93}}{2}$.

Q787. Here $-b = -(-5) = +5$, and $-4ac = -4(1)(-2) = +32$. Thus $D = 5^2 + 32 = 57$. So $x = \frac{5\pm\sqrt{57}}{2}$.

Q788. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-5) = +40$. Thus $D = 3^2 + 40 = 49$. So $x = \frac{3\pm\sqrt{49}}{2}$.

Q789. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-9) = +180$. Thus $D = 7^2 + 180 = 229$. So $x = \frac{7\pm\sqrt{229}}{2}$.

Q790. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-2) = +24$. Thus $D = 3^2 + 24 = 33$. So $x = \frac{3\pm\sqrt{33}}{2}$.

Q791. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-4) = +32$. Thus $D = 8^2 + 32 = 96$. So $x = \frac{8\pm\sqrt{96}}{2}$.

Q792. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-1) = +16$. Thus $D = 8^2 + 16 = 80$. So $x = \frac{8 \pm \sqrt{80}}{2}$.

Q793. Here $-b = -(-9) = +9$, and $-4ac = -4(1)(-7) = +56$. Thus $D = 9^2 + 56 = 137$. So $x = \frac{9 \pm \sqrt{137}}{2}$.

Q794. Here $-b = -(-6) = +6$, and $-4ac = -4(1)(-1) = +8$. Thus $D = 6^2 + 8 = 44$. So $x = \frac{6 \pm \sqrt{44}}{2}$.

Q795. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-4) = +48$. Thus $D = 10^2 + 48 = 148$. So $x = \frac{10 \pm \sqrt{148}}{2}$.

Q796. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-10) = +200$. Thus $D = 10^2 + 200 = 300$. So $x = \frac{10 \pm \sqrt{300}}{2}$.

Q797. Here $-b = -(-10) = +10$, and $-4ac = -4(1)(-1) = +8$. Thus $D = 10^2 + 8 = 108$. So $x = \frac{10 \pm \sqrt{108}}{2}$.

Q798. Here $-b = -(-7) = +7$, and $-4ac = -4(1)(-8) = +128$. Thus $D = 7^2 + 128 = 177$. So $x = \frac{7 \pm \sqrt{177}}{2}$.

Q799. Here $-b = -(-8) = +8$, and $-4ac = -4(1)(-4) = +16$. Thus $D = 8^2 + 16 = 80$. So $x = \frac{8 \pm \sqrt{80}}{2}$.

Q800. Here $-b = -(-3) = +3$, and $-4ac = -4(1)(-11) = +44$. Thus $D = 3^2 + 44 = 53$. So $x = \frac{3 \pm \sqrt{53}}{2}$.

Q801. 'Subtract A from B' means $B - A$. So $(5x + 3) - (3x - 5) = 5x + 3 - 3x + 5 = (2)x + 8$.

Q802. 'Subtract A from B' means $B - A$. So $(7x + 6) - (6x - 7) = 7x + 6 - 6x + 7 = (1)x + 13$.

Q803. 'Subtract A from B' means $B - A$. So $(14x + 9) - (9x - 14) = 14x + 9 - 9x + 14 = (5)x + 23$.

Q804. 'Subtract A from B' means $B - A$. So $(10x + 3) - (3x - 10) = 10x + 3 - 3x + 10 = (7)x + 13$.

Q805. 'Subtract A from B' means $B - A$. So $(11x + 9) - (9x - 11) = 11x + 9 - 9x + 11 = (2)x + 20$.

Q806. 'Subtract A from B' means $B - A$. So $(6x + 7) - (7x - 6) = 6x + 7 - 7x + 6 = (-1)x + 13$.

Q807. 'Subtract A from B' means $B - A$. So $(5x + 2) - (2x - 5) = 5x + 2 - 2x + 5 = (3)x + 7$.

Q808. 'Subtract A from B' means $B - A$. So $(13x + 6) - (6x - 13) = 13x + 6 - 6x + 13 = (7)x + 19$.

Q809. 'Subtract A from B' means $B - A$. So $(9x + 2) - (2x - 9) = 9x + 2 - 2x + 9 = (7)x + 11$.

Q810. 'Subtract A from B' means $B - A$. So $(6x + 9) - (9x - 6) = 6x + 9 - 9x + 6 = (-3)x + 15$.

Q811. 'Subtract A from B' means $B - A$. So $(7x + 8) - (8x - 7) = 7x + 8 - 8x + 7 = (-1)x + 15$.

Q812. 'Subtract A from B' means $B - A$. So $(6x + 6) - (6x - 6) = 6x + 6 - 6x + 6 = (0)x + 12$.

Q813. 'Subtract A from B' means $B - A$. So $(14x + 7) - (7x - 14) = 14x + 7 - 7x + 14 = (7)x + 21$.

Q814. 'Subtract A from B' means $B - A$. So $(4x + 8) - (8x - 4) = 4x + 8 - 8x + 4 = (-4)x + 12$.

Q815. 'Subtract A from B' means $B - A$. So $(9x + 2) - (2x - 9) = 9x + 2 - 2x + 9 = (7)x + 11$.

Q816. 'Subtract A from B' means $B - A$. So $(5x + 8) - (8x - 5) = 5x + 8 - 8x + 5 = (-3)x + 13$.

Q817. 'Subtract A from B' means $B - A$. So $(12x + 8) - (8x - 12) = 12x + 8 - 8x + 12 = (4)x + 20$.

Q818. 'Subtract A from B' means $B - A$. So $(11x + 6) - (6x - 11) = 11x + 6 - 6x + 11 = (5)x + 17$.

Q819. 'Subtract A from B' means $B - A$. So $(5x + 4) - (4x - 5) = 5x + 4 - 4x + 5 = (1)x + 9$.

Q820. 'Subtract A from B' means $B - A$. So $(14x + 7) - (7x - 14) = 14x + 7 - 7x + 14 = (7)x + 21$.

Q821. 'Subtract A from B' means $B - A$. So $(13x + 4) - (4x - 13) = 13x + 4 - 4x + 13 = (9)x + 17$.

Q822. 'Subtract A from B' means $B - A$. So $(4x + 6) - (6x - 4) = 4x + 6 - 6x + 4 = (-2)x + 10$.

Q823. 'Subtract A from B' means $B - A$. So $(6x + 9) - (9x - 6) = 6x + 9 - 9x + 6 = (-3)x + 15$.

Q824. 'Subtract A from B' means $B - A$. So $(3x + 4) - (4x - 3) = 3x + 4 - 4x + 3 = (-1)x + 7$.

Q825. 'Subtract A from B' means $B - A$. So $(14x + 5) - (5x - 14) = 14x + 5 - 5x + 14 = (9)x + 19$.

Q826. 'Subtract A from B' means $B - A$. So $(12x + 9) - (9x - 12) = 12x + 9 - 9x + 12 = (3)x + 21$.

Q827. 'Subtract A from B' means $B - A$. So $(4x + 2) - (2x - 4) = 4x + 2 - 2x + 4 = (2)x + 6$.

Q828. 'Subtract A from B' means $B - A$. So $(12x + 8) - (8x - 12) = 12x + 8 - 8x + 12 = (4)x + 20$.

Q829. 'Subtract A from B' means $B - A$. So $(11x + 7) - (7x - 11) = 11x + 7 - 7x + 11 = (4)x + 18$.

Q830. 'Subtract A from B' means $B - A$. So $(9x + 6) - (6x - 9) = 9x + 6 - 6x + 9 = (3)x + 15$.

Q831. 'Subtract A from B' means $B - A$. So $(7x + 2) - (2x - 7) = 7x + 2 - 2x + 7 = (5)x + 9$.

Q832. 'Subtract A from B' means $B - A$. So $(15x + 9) - (9x - 15) = 15x + 9 - 9x + 15 = (6)x + 24$.

Q833. 'Subtract A from B' means $B - A$. So $(11x + 6) - (6x - 11) = 11x + 6 - 6x + 11 = (5)x + 17$.

Q834. 'Subtract A from B' means $B - A$. So $(10x + 2) - (2x - 10) = 10x + 2 - 2x + 10 = (8)x + 12$.

Q835. 'Subtract A from B' means $B - A$. So $(6x + 9) - (9x - 6) = 6x + 9 - 9x + 6 = (-3)x + 15$.

Q836. 'Subtract A from B' means $B - A$. So $(7x + 6) - (6x - 7) = 7x + 6 - 6x + 7 = (1)x + 13$.

Q837. 'Subtract A from B' means $B - A$. So $(11x + 5) - (5x - 11) = 11x + 5 - 5x + 11 = (6)x + 16$.

Q838. 'Subtract A from B' means $B - A$. So $(11x + 6) - (6x - 11) = 11x + 6 - 6x + 11 = (5)x + 17$.

Q839. 'Subtract A from B' means $B - A$. So $(12x + 9) - (9x - 12) = 12x + 9 - 9x + 12 = (3)x + 21$.

Q840. 'Subtract A from B' means $B - A$. So $(12x + 3) - (3x - 12) = 12x + 3 - 3x + 12 = (9)x + 15$.

Q841. 'Subtract A from B' means $B - A$. So $(12x + 7) - (7x - 12) = 12x + 7 - 7x + 12 = (5)x + 19$.

Q842. 'Subtract A from B' means $B - A$. So $(8x + 7) - (7x - 8) = 8x + 7 - 7x + 8 = (1)x + 15$.

Q843. 'Subtract A from B' means $B - A$. So $(10x + 7) - (7x - 10) = 10x + 7 - 7x + 10 = (3)x + 17$.

Q844. 'Subtract A from B' means $B - A$. So $(5x + 5) - (5x - 5) = 5x + 5 - 5x + 5 = (0)x + 10$.

Q845. 'Subtract A from B' means $B - A$. So $(5x + 2) - (2x - 5) = 5x + 2 - 2x + 5 = (3)x + 7$.

Q846. 'Subtract A from B' means $B - A$. So $(11x + 6) - (6x - 11) = 11x + 6 - 6x + 11 = (5)x + 17$.

Q847. 'Subtract A from B' means $B - A$. So $(14x + 8) - (8x - 14) = 14x + 8 - 8x + 14 = (6)x + 22$.

Q848. 'Subtract A from B' means $B - A$. So $(10x + 8) - (8x - 10) = 10x + 8 - 8x + 10 = (2)x + 18$.

Q849. 'Subtract A from B' means $B - A$. So $(4x + 9) - (9x - 4) = 4x + 9 - 9x + 4 = (-5)x + 13$.

Q850. 'Subtract A from B' means $B - A$. So $(12x + 7) - (7x - 12) = 12x + 7 - 7x + 12 = (5)x + 19$.

Q851. 'Subtract A from B' means $B - A$. So $(5x + 8) - (8x - 5) = 5x + 8 - 8x + 5 = (-3)x + 13$.

Q852. 'Subtract A from B' means $B - A$. So $(8x + 8) - (8x - 8) = 8x + 8 - 8x + 8 = (0)x + 16$.

Q853. 'Subtract A from B' means $B - A$. So $(10x + 4) - (4x - 10) = 10x + 4 - 4x + 10 = (6)x + 14$.

Q854. 'Subtract A from B' means $B - A$. So $(8x + 9) - (9x - 8) = 8x + 9 - 9x + 8 = (-1)x + 17$.

Q855. 'Subtract A from B' means $B - A$. So $(13x + 8) - (8x - 13) = 13x + 8 - 8x + 13 = (5)x + 21$.

Q856. 'Subtract A from B' means $B - A$. So $(9x + 5) - (5x - 9) = 9x + 5 - 5x + 9 = (4)x + 14$.

Q857. 'Subtract A from B' means $B - A$. So $(6x + 7) - (7x - 6) = 6x + 7 - 7x + 6 = (-1)x + 13$.

Q858. 'Subtract A from B' means $B - A$. So $(15x + 7) - (7x - 15) = 15x + 7 - 7x + 15 = (8)x + 22$.

Q859. 'Subtract A from B' means $B - A$. So $(9x + 7) - (7x - 9) = 9x + 7 - 7x + 9 = (2)x + 16$.

Q860. 'Subtract A from B' means $B - A$. So $(15x + 2) - (2x - 15) = 15x + 2 - 2x + 15 = (13)x + 17$.

Q861. 'Subtract A from B' means $B - A$. So $(10x + 3) - (3x - 10) = 10x + 3 - 3x + 10 = (7)x + 13$.

Q862. 'Subtract A from B' means $B - A$. So $(7x + 6) - (6x - 7) = 7x + 6 - 6x + 7 = (1)x + 13$.

Q863. 'Subtract A from B' means $B - A$. So $(13x + 7) - (7x - 13) = 13x + 7 - 7x + 13 = (6)x + 20$.

Q864. 'Subtract A from B' means $B - A$. So $(4x + 6) - (6x - 4) = 4x + 6 - 6x + 4 = (-2)x + 10$.

Q865. 'Subtract A from B' means $B - A$. So $(5x + 4) - (4x - 5) = 5x + 4 - 4x + 5 = (1)x + 9$.

Q866. 'Subtract A from B' means $B - A$. So $(8x + 3) - (3x - 8) = 8x + 3 - 3x + 8 = (5)x + 11$.

Q867. 'Subtract A from B' means $B - A$. So $(8x + 9) - (9x - 8) = 8x + 9 - 9x + 8 = (-1)x + 17$.

Q868. 'Subtract A from B' means $B - A$. So $(5x + 2) - (2x - 5) = 5x + 2 - 2x + 5 = (3)x + 7$.

Q869. 'Subtract A from B' means $B - A$. So $(14x + 4) - (4x - 14) = 14x + 4 - 4x + 14 = (10)x + 18$.

Q870. 'Subtract A from B' means $B - A$. So $(12x + 8) - (8x - 12) = 12x + 8 - 8x + 12 = (4)x + 20$.

Q871. 'Subtract A from B' means $B - A$. So $(14x + 4) - (4x - 14) = 14x + 4 - 4x + 14 = (10)x + 18$.

Q872. 'Subtract A from B' means $B - A$. So $(15x + 3) - (3x - 15) = 15x + 3 - 3x + 15 = (12)x + 18$.

Q873. 'Subtract A from B' means $B - A$. So $(11x + 2) - (2x - 11) = 11x + 2 - 2x + 11 = (9)x + 13$.

Q874. 'Subtract A from B' means $B - A$. So $(7x + 5) - (5x - 7) = 7x + 5 - 5x + 7 = (2)x + 12$.

Q875. 'Subtract A from B' means $B - A$. So $(4x + 8) - (8x - 4) = 4x + 8 - 8x + 4 = (-4)x + 12$.

Q876. 'Subtract A from B' means $B - A$. So $(6x + 9) - (9x - 6) = 6x + 9 - 9x + 6 = (-3)x + 15$.

Q877. 'Subtract A from B' means $B - A$. So $(4x + 5) - (5x - 4) = 4x + 5 - 5x + 4 = (-1)x + 9$.

Q878. 'Subtract A from B' means $B - A$. So $(10x + 8) - (8x - 10) = 10x + 8 - 8x + 10 = (2)x + 18$.

Q879. 'Subtract A from B' means $B - A$. So $(6x + 4) - (4x - 6) = 6x + 4 - 4x + 6 = (2)x + 10$.

Q880. 'Subtract A from B' means $B - A$. So $(13x + 5) - (5x - 13) = 13x + 5 - 5x + 13 = (8)x + 18$.

Q881. 'Subtract A from B' means $B - A$. So $(14x + 9) - (9x - 14) = 14x + 9 - 9x + 14 = (5)x + 23$.

Q882. 'Subtract A from B' means $B - A$. So $(8x + 5) - (5x - 8) = 8x + 5 - 5x + 8 = (3)x + 13$.

Q883. 'Subtract A from B' means $B - A$. So $(8x + 8) - (8x - 8) = 8x + 8 - 8x + 8 = (0)x + 16$.

Q884. 'Subtract A from B' means $B - A$. So $(11x + 6) - (6x - 11) = 11x + 6 - 6x + 11 = (5)x + 17$.

Q885. 'Subtract A from B' means $B - A$. So $(5x + 8) - (8x - 5) = 5x + 8 - 8x + 5 = (-3)x + 13$.

Q886. 'Subtract A from B' means $B - A$. So $(12x + 4) - (4x - 12) = 12x + 4 - 4x + 12 = (8)x + 16$.

Q887. 'Subtract A from B' means $B - A$. So $(10x + 5) - (5x - 10) = 10x + 5 - 5x + 10 = (5)x + 15$.

Q888. 'Subtract A from B' means $B - A$. So $(4x + 7) - (7x - 4) = 4x + 7 - 7x + 4 = (-3)x + 11$.

Q889. 'Subtract A from B' means $B - A$. So $(11x + 2) - (2x - 11) = 11x + 2 - 2x + 11 = (9)x + 13$.

Q890. 'Subtract A from B' means $B - A$. So $(8x + 4) - (4x - 8) = 8x + 4 - 4x + 8 = (4)x + 12$.

Q891. 'Subtract A from B' means $B - A$. So $(5x + 7) - (7x - 5) = 5x + 7 - 7x + 5 = (-2)x + 12$.

Q892. 'Subtract A from B' means $B - A$. So $(10x + 8) - (8x - 10) = 10x + 8 - 8x + 10 = (2)x + 18$.

Q893. 'Subtract A from B' means $B - A$. So $(9x + 8) - (8x - 9) = 9x + 8 - 8x + 9 = (1)x + 17$.

Q894. 'Subtract A from B' means $B - A$. So $(3x + 5) - (5x - 3) = 3x + 5 - 5x + 3 = (-2)x + 8$.

Q895. 'Subtract A from B' means $B - A$. So $(7x + 8) - (8x - 7) = 7x + 8 - 8x + 7 = (-1)x + 15$.

Q896. 'Subtract A from B' means $B - A$. So $(12x + 3) - (3x - 12) = 12x + 3 - 3x + 12 = (9)x + 15$.

Q897. 'Subtract A from B' means $B - A$. So $(3x + 3) - (3x - 3) = 3x + 3 - 3x + 3 = (0)x + 6$.

Q898. 'Subtract A from B' means $B - A$. So $(12x + 8) - (8x - 12) = 12x + 8 - 8x + 12 = (4)x + 20$.

Q899. 'Subtract A from B' means $B - A$. So $(3x + 5) - (5x - 3) = 3x + 5 - 5x + 3 = (-2)x + 8$.

Q900. 'Subtract A from B' means $B - A$. So $(4x+7)-(7x-4) = 4x+7-7x+4 = (-3)x+11$.

Q901. Transposing 9 yields '$-7x = 19 - 9 = 10$. Dividing by negative number -7 flips sign of numerator: $x = \frac{10}{-7} = \frac{-10}{7}$.

Q902. Transposing 19 yields '$-5x = 4 - 19 = -15$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-15}{-5} = \frac{15}{5}$.

Q903. Transposing 19 yields '$-7x = -16 - 19 = -35$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-35}{-7} = \frac{35}{7}$.

Q904. Transposing 9 yields '$-6x = 10 - 9 = 1$. Dividing by negative number -6 flips sign of numerator: $x = \frac{1}{-6} = \frac{-1}{6}$.

Q905. Transposing 12 yields '$-7x = -8 - 12 = -20$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-20}{-7} = \frac{20}{7}$.

Q906. Transposing 18 yields '$-4x = -18 - 18 = -36$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-36}{-4} = \frac{36}{4}$.

Q907. Transposing 30 yields '$-6x = 15 - 30 = -15$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-15}{-6} = \frac{15}{6}$.

Q908. Transposing 30 yields '$-4x = 12 - 30 = -18$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-18}{-4} = \frac{18}{4}$.

Q909. Transposing 26 yields '$-5x = 6 - 26 = -20$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-20}{-5} = \frac{20}{5}$.

Q910. Transposing 27 yields '$-8x = 14 - 27 = -13$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-13}{-8} = \frac{13}{8}$.

Q911. Transposing 13 yields '$-9x = 3 - 13 = -10$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-10}{-9} = \frac{10}{9}$.

Q912. Transposing 7 yields '$-7x = 12 - 7 = 5$. Dividing by negative number -7 flips sign of numerator: $x = \frac{5}{-7} = \frac{-5}{7}$.

Q913. Transposing 21 yields '$-2x = 2 - 21 = -19$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-19}{-2} = \frac{19}{2}$.

Q914. Transposing 16 yields '$-7x = -1 - 16 = -17$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-17}{-7} = \frac{17}{7}$.

Q915. Transposing 10 yields '$-4x = 13 - 10 = 3$. Dividing by negative number -4 flips sign of numerator: $x = \frac{3}{-4} = \frac{-3}{4}$.

Q916. Transposing 29 yields '$-7x = -5 - 29 = -34$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-34}{-7} = \frac{34}{7}$.

Q917. Transposing 13 yields '$-6x = 12 - 13 = -1$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-1}{-6} = \frac{1}{6}$.

Q918. Transposing 24 yields '$-8x = -6 - 24 = -30$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-30}{-8} = \frac{30}{8}$.

Q919. Transposing 17 yields '$-5x = -12 - 17 = -29$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-29}{-5} = \frac{29}{5}$.

Q920. Transposing 18 yields '$-7x = -4 - 18 = -22$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-22}{-7} = \frac{22}{7}$.

Q921. Transposing 15 yields '$-4x = 3 - 15 = -12$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-12}{-4} = \frac{12}{4}$.

Q922. Transposing 24 yields '$-3x = -12 - 24 = -36$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-36}{-3} = \frac{36}{3}$.

Q923. Transposing 14 yields '$-8x = 0 - 14 = -14$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-14}{-8} = \frac{14}{8}$.

Q924. Transposing 17 yields '$-5x = -19 - 17 = -36$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-36}{-5} = \frac{36}{5}$.

Q925. Transposing 23 yields '$-5x = -4 - 23 = -27$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-27}{-5} = \frac{27}{5}$.

Q926. Transposing 11 yields '$-9x = 9 - 11 = -2$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-2}{-9} = \frac{2}{9}$.

Q927. Transposing 8 yields '$-5x = 2 - 8 = -6$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-6}{-5} = \frac{6}{5}$.

Q928. Transposing 14 yields '$-7x = -19 - 14 = -33$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-33}{-7} = \frac{33}{7}$.

Q929. Transposing 29 yields '$-5x = -3 - 29 = -32$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-32}{-5} = \frac{32}{5}$.

Q930. Transposing 7 yields '$-7x = 6 - 7 = -1$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-1}{-7} = \frac{1}{7}$.

Q931. Transposing 29 yields '$-9x = 5 - 29 = -24$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-24}{-9} = \frac{24}{9}$.

Q932. Transposing 17 yields '$-4x = 10 - 17 = -7$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-7}{-4} = \frac{7}{4}$.

Q933. Transposing 22 yields '$-6x = -18 - 22 = -40$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-40}{-6} = \frac{40}{6}$.

Q934. Transposing 21 yields '$-6x = -20 - 21 = -41$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-41}{-6} = \frac{41}{6}$.

Q935. Transposing 15 yields '$-6x = 7 - 15 = -8$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-8}{-6} = \frac{8}{6}$.

Q936. Transposing 16 yields '$-8x = 13 - 16 = -3$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-3}{-8} = \frac{3}{8}$.

Q937. Transposing 21 yields '$-2x = 20 - 21 = -1$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-1}{-2} = \frac{1}{2}$.

Q938. Transposing 5 yields '$-2x = 18 - 5 = 13$. Dividing by negative number -2 flips sign of numerator: $x = \frac{13}{-2} = \frac{-13}{2}$.

Q939. Transposing 12 yields $-8x = 6 - 12 = -6$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-6}{-8} = \frac{6}{8}$.

Q940. Transposing 29 yields $-7x = -4 - 29 = -33$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-33}{-7} = \frac{33}{7}$.

Q941. Transposing 24 yields $-4x = -18 - 24 = -42$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-42}{-4} = \frac{42}{4}$.

Q942. Transposing 14 yields $-5x = 17 - 14 = 3$. Dividing by negative number -5 flips sign of numerator: $x = \frac{3}{-5} = \frac{-3}{5}$.

Q943. Transposing 21 yields $-6x = -4 - 21 = -25$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-25}{-6} = \frac{25}{6}$.

Q944. Transposing 12 yields $-2x = -19 - 12 = -31$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-31}{-2} = \frac{31}{2}$.

Q945. Transposing 13 yields $-5x = -15 - 13 = -28$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-28}{-5} = \frac{28}{5}$.

Q946. Transposing 19 yields $-7x = -4 - 19 = -23$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-23}{-7} = \frac{23}{7}$.

Q947. Transposing 27 yields $-5x = 9 - 27 = -18$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-18}{-5} = \frac{18}{5}$.

Q948. Transposing 28 yields $-6x = -7 - 28 = -35$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-35}{-6} = \frac{35}{6}$.

Q949. Transposing 12 yields $-8x = -17 - 12 = -29$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-29}{-8} = \frac{29}{8}$.

Q950. Transposing 21 yields $-2x = -10 - 21 = -31$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-31}{-2} = \frac{31}{2}$.

Q951. Transposing 14 yields $-2x = -6 - 14 = -20$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-20}{-2} = \frac{20}{2}$.

Q952. Transposing 10 yields $-7x = 18 - 10 = 8$. Dividing by negative number -7 flips sign of numerator: $x = \frac{8}{-7} = \frac{-8}{7}$.

Q953. Transposing 19 yields $-6x = 11 - 19 = -8$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-8}{-6} = \frac{8}{6}$.

Q954. Transposing 14 yields $-5x = -12 - 14 = -26$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-26}{-5} = \frac{26}{5}$.

Q955. Transposing 20 yields $-4x = 16 - 20 = -4$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-4}{-4} = \frac{4}{4}$.

Q956. Transposing 21 yields $-2x = 11 - 21 = -10$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-10}{-2} = \frac{10}{2}$.

Q957. Transposing 29 yields $-6x = -17 - 29 = -46$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-46}{-6} = \frac{46}{6}$.

Q958. Transposing 21 yields $-9x = 6 - 21 = -15$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-15}{-9} = \frac{15}{9}$.

Q959. Transposing 11 yields $-7x = -13 - 11 = -24$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-24}{-7} = \frac{24}{7}$.

Q960. Transposing 8 yields $-3x = 10 - 8 = 2$. Dividing by negative number -3 flips sign of numerator: $x = \frac{2}{-3} = \frac{-2}{3}$.

Q961. Transposing 18 yields $-3x = 16 - 18 = -2$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-2}{-3} = \frac{2}{3}$.

Q962. Transposing 24 yields $-9x = 11 - 24 = -13$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-13}{-9} = \frac{13}{9}$.

Q963. Transposing 17 yields $-3x = -12 - 17 = -29$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-29}{-3} = \frac{29}{3}$.

Q964. Transposing 20 yields $-8x = -15 - 20 = -35$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-35}{-8} = \frac{35}{8}$.

Q965. Transposing 17 yields $-9x = -13 - 17 = -30$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-30}{-9} = \frac{30}{9}$.

Q966. Transposing 28 yields $-6x = 20 - 28 = -8$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-8}{-6} = \frac{8}{6}$.

Q967. Transposing 26 yields $-7x = 20 - 26 = -6$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-6}{-7} = \frac{6}{7}$.

Q968. Transposing 14 yields $-3x = 8 - 14 = -6$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-6}{-3} = \frac{6}{3}$.

Q969. Transposing 12 yields $-9x = 8 - 12 = -4$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-4}{-9} = \frac{4}{9}$.

Q970. Transposing 16 yields $-2x = -19 - 16 = -35$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-35}{-2} = \frac{35}{2}$.

Q971. Transposing 5 yields $-3x = 2 - 5 = -3$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-3}{-3} = \frac{3}{3}$.

Q972. Transposing 29 yields $-2x = 3 - 29 = -26$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-26}{-2} = \frac{26}{2}$.

Q973. Transposing 16 yields $-9x = 19 - 16 = 3$. Dividing by negative number -9 flips sign of numerator: $x = \frac{3}{-9} = \frac{-3}{9}$.

Q974. Transposing 26 yields $-3x = -14 - 26 = -40$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-40}{-3} = \frac{40}{3}$.

Q975. Transposing 26 yields $-8x = 5 - 26 = -21$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-21}{-8} = \frac{21}{8}$.

Q976. Transposing 30 yields $-6x = 20 - 30 = -10$. Dividing by negative number -6 flips sign of numerator: $x = \frac{-10}{-6} = \frac{10}{6}$.

Q977. Transposing 19 yields $-9x = -14 - 19 = -33$. Dividing by negative number -9 flips sign of numerator:

$x = \frac{-33}{-9} = \frac{33}{9}$.

Q978. Transposing 28 yields $-9x = -6 - 28 = -34$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-34}{-9} = \frac{34}{9}$.

Q979. Transposing 28 yields $-9x = 6 - 28 = -22$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-22}{-9} = \frac{22}{9}$.

Q980. Transposing 27 yields $-4x = 7 - 27 = -20$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-20}{-4} = \frac{20}{4}$.

Q981. Transposing 12 yields $-2x = -3 - 12 = -15$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-15}{-2} = \frac{15}{2}$.

Q982. Transposing 12 yields $-9x = -9 - 12 = -21$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-21}{-9} = \frac{21}{9}$.

Q983. Transposing 9 yields $-4x = 10 - 9 = 1$. Dividing by negative number -4 flips sign of numerator: $x = \frac{1}{-4} = \frac{-1}{4}$.

Q984. Transposing 26 yields $-3x = -6 - 26 = -32$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-32}{-3} = \frac{32}{3}$.

Q985. Transposing 14 yields $-8x = 3 - 14 = -11$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-11}{-8} = \frac{11}{8}$.

Q986. Transposing 29 yields $-7x = 4 - 29 = -25$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-25}{-7} = \frac{25}{7}$.

Q987. Transposing 23 yields $-5x = -15 - 23 = -38$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-38}{-5} = \frac{38}{5}$.

Q988. Transposing 6 yields $-4x = -16 - 6 = -22$. Dividing by negative number -4 flips sign of numerator: $x = \frac{-22}{-4} = \frac{22}{4}$.

Q989. Transposing 5 yields $-9x = -1 - 5 = -6$. Dividing by negative number -9 flips sign of numerator: $x = \frac{-6}{-9} = \frac{6}{9}$.

Q990. Transposing 11 yields $-9x = 15 - 11 = 4$. Dividing by negative number -9 flips sign of numerator: $x = \frac{4}{-9} = \frac{-4}{9}$.

Q991. Transposing 30 yields $-2x = -8 - 30 = -38$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-38}{-2} = \frac{38}{2}$.

Q992. Transposing 8 yields $-7x = -16 - 8 = -24$. Dividing by negative number -7 flips sign of numerator: $x = \frac{-24}{-7} = \frac{24}{7}$.

Q993. Transposing 11 yields $-8x = 8 - 11 = -3$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-3}{-8} = \frac{3}{8}$.

Q994. Transposing 21 yields $-2x = 7 - 21 = -14$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-14}{-2} = \frac{14}{2}$.

Q995. Transposing 15 yields $-8x = -10 - 15 = -25$. Dividing by negative number -8 flips sign of numerator: $x = \frac{-25}{-8} = \frac{25}{8}$.

Q996. Transposing 6 yields $-3x = -19 - 6 = -25$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-25}{-3} = \frac{25}{3}$.

Q997. Transposing 30 yields $-5x = 6 - 30 = -24$. Dividing by negative number -5 flips sign of numerator: $x = \frac{-24}{-5} = \frac{24}{5}$.

Q998. Transposing 9 yields $-7x = 16 - 9 = 7$. Dividing by negative number -7 flips sign of numerator: $x = \frac{7}{-7} = \frac{-7}{7}$.

Q999. Transposing 23 yields $-2x = 18 - 23 = -5$. Dividing by negative number -2 flips sign of numerator: $x = \frac{-5}{-2} = \frac{5}{2}$.

Q1000. Transposing 19 yields $-3x = 11 - 19 = -8$. Dividing by negative number -3 flips sign of numerator: $x = \frac{-8}{-3} = \frac{8}{3}$.

MODULE 2: WORD PROBLEM TRANSLATION AND EQUATION FORMATION (Q1001 – Q2000) — Answer Key

Q1001. Equations: $x + y = 103$ and $x - y = 9$. Adding gives $2x = 112 \implies x = 56$. Subtracting gives $2y = 94 \implies y = 47$.

Q1002. Equations: $x + y = 41$ and $x - y = 7$. Adding gives $2x = 48 \implies x = 24$. Subtracting gives $2y = 34 \implies y = 17$.

Q1003. Equations: $x + y = 38$ and $x - y = 12$. Adding gives $2x = 50 \implies x = 25$. Subtracting gives $2y = 26 \implies y = 13$.

Q1004. Equations: $x + y = 38$ and $x - y = 24$. Adding gives $2x = 62 \implies x = 31$. Subtracting gives $2y = 14 \implies y = 7$.

Q1005. Equations: $x + y = 118$ and $x - y = 8$. Adding gives $2x = 126 \implies x = 63$. Subtracting gives $2y = 110 \implies y = 55$.

Q1006. Equations: $x + y = 92$ and $x - y = 4$. Adding gives $2x = 96 \implies x = 48$. Subtracting gives $2y = 88 \implies y = 44$.

Q1007. Equations: $x + y = 47$ and $x - y = 25$. Adding gives $2x = 72 \implies x = 36$. Subtracting gives $2y = 22 \implies y = 11$.

Q1008. Equations: $x + y = 89$ and $x - y = 17$. Adding gives $2x = 106 \implies x = 53$. Subtracting gives $2y = 72 \implies y = 36$.

Q1009. Equations: $x + y = 116$ and $x - y = 16$. Adding gives $2x = 132 \implies x = 66$. Subtracting gives $2y = 100 \implies y = 50$.

Q1010. Equations: $x + y = 115$ and $x - y = 5$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 110 \implies y = 55$.

Q1011. Equations: $x + y = 63$ and $x - y = 9$. Adding gives $2x = 72 \implies x = 36$. Subtracting gives $2y = 54 \implies y = 27$.

Q1012. Equations: $x + y = 38$ and $x - y = 26$. Adding gives $2x = 64 \implies x = 32$. Subtracting gives $2y = 12 \implies y = 6$.

Q1013. Equations: $x + y = 95$ and $x - y = 25$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 70 \implies y = 35$.

Q1014. Equations: $x + y = 94$ and $x - y = 6$. Adding gives $2x = 100 \implies x = 50$. Subtracting gives $2y = 88 \implies y = 44$.

Q1015. Equations: $x + y = 85$ and $x - y = 5$. Adding gives $2x = 90 \implies x = 45$. Subtracting gives $2y = 80 \implies y = 40$.

Q1016. Equations: $x + y = 46$ and $x - y = 20$. Adding gives $2x = 66 \implies x = 33$. Subtracting gives $2y = 26 \implies y = 13$.

Q1017. Equations: $x + y = 64$ and $x - y = 4$. Adding gives $2x = 68 \implies x = 34$. Subtracting gives $2y = 60 \implies y = 30$.

Q1018. Equations: $x + y = 65$ and $x - y = 11$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 54 \implies y = 27$.

Q1019. Equations: $x + y = 81$ and $x - y = 7$. Adding gives $2x = 88 \implies x = 44$. Subtracting gives $2y = 74 \implies y = 37$.

Q1020. Equations: $x + y = 52$ and $x - y = 18$. Adding gives $2x = 70 \implies x = 35$. Subtracting gives $2y = 34 \implies y = 17$.

Q1021. Equations: $x + y = 49$ and $x - y = 27$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 22 \implies y = 11$.

Q1022. Equations: $x + y = 87$ and $x - y = 17$. Adding gives $2x = 104 \implies x = 52$. Subtracting gives $2y = 70 \implies y = 35$.

Q1023. Equations: $x + y = 45$ and $x - y = 11$. Adding gives $2x = 56 \implies x = 28$. Subtracting gives $2y = 34 \implies y = 17$.

Q1024. Equations: $x + y = 82$ and $x - y = 6$. Adding gives $2x = 88 \implies x = 44$. Subtracting gives $2y = 76 \implies y = 38$.

Q1025. Equations: $x + y = 88$ and $x - y = 24$. Adding gives $2x = 112 \implies x = 56$. Subtracting gives $2y = 64 \implies y = 32$.

Q1026. Equations: $x + y = 62$ and $x - y = 18$. Adding gives $2x = 80 \implies x = 40$. Subtracting gives $2y = 44 \implies y = 22$.

Q1027. Equations: $x + y = 74$ and $x - y = 28$. Adding gives $2x = 102 \implies x = 51$. Subtracting gives $2y = 46 \implies y = 23$.

Q1028. Equations: $x + y = 112$ and $x - y = 28$. Adding gives $2x = 140 \implies x = 70$. Subtracting gives $2y = 84 \implies y = 42$.

Q1029. Equations: $x + y = 79$ and $x - y = 13$. Adding gives $2x = 92 \implies x = 46$. Subtracting gives $2y = 66 \implies y = 33$.

Q1030. Equations: $x + y = 75$ and $x - y = 23$. Adding gives $2x = 98 \implies x = 49$. Subtracting gives $2y = 52 \implies y = 26$.

Q1031. Equations: $x + y = 120$ and $x - y = 26$. Adding gives $2x = 146 \implies x = 73$. Subtracting gives $2y = 94 \implies y = 47$.

Q1032. Equations: $x + y = 74$ and $x - y = 10$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 64 \implies y = 32$.

Q1033. Equations: $x + y = 63$ and $x - y = 5$. Adding gives $2x = 68 \implies x = 34$. Subtracting gives $2y = 58 \implies y = 29$.

Q1034. Equations: $x + y = 90$ and $x - y = 20$. Adding gives $2x = 110 \implies x = 55$. Subtracting gives $2y = 70 \implies y = 35$.

Q1035. Equations: $x + y = 49$ and $x - y = 13$. Adding gives $2x = 62 \implies x = 31$. Subtracting gives $2y = 36 \implies y = 18$.

Q1036. Equations: $x + y = 51$ and $x - y = 23$. Adding gives $2x = 74 \implies x = 37$. Subtracting gives $2y = 28 \implies y = 14$.

Q1037. Equations: $x + y = 73$ and $x - y = 25$. Adding gives $2x = 98 \implies x = 49$. Subtracting gives

$2y = 48 \implies y = 24$.

Q1038. Equations: $x + y = 109$ and $x - y = 11$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 98 \implies y = 49$.

Q1039. Equations: $x + y = 40$ and $x - y = 22$. Adding gives $2x = 62 \implies x = 31$. Subtracting gives $2y = 18 \implies y = 9$.

Q1040. Equations: $x + y = 120$ and $x - y = 16$. Adding gives $2x = 136 \implies x = 68$. Subtracting gives $2y = 104 \implies y = 52$.

Q1041. Equations: $x + y = 72$ and $x - y = 22$. Adding gives $2x = 94 \implies x = 47$. Subtracting gives $2y = 50 \implies y = 25$.

Q1042. Equations: $x + y = 89$ and $x - y = 7$. Adding gives $2x = 96 \implies x = 48$. Subtracting gives $2y = 82 \implies y = 41$.

Q1043. Equations: $x + y = 40$ and $x - y = 12$. Adding gives $2x = 52 \implies x = 26$. Subtracting gives $2y = 28 \implies y = 14$.

Q1044. Equations: $x + y = 106$ and $x - y = 26$. Adding gives $2x = 132 \implies x = 66$. Subtracting gives $2y = 80 \implies y = 40$.

Q1045. Equations: $x + y = 120$ and $x - y = 14$. Adding gives $2x = 134 \implies x = 67$. Subtracting gives $2y = 106 \implies y = 53$.

Q1046. Equations: $x + y = 48$ and $x - y = 28$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 20 \implies y = 10$.

Q1047. Equations: $x + y = 83$ and $x - y = 25$. Adding gives $2x = 108 \implies x = 54$. Subtracting gives $2y = 58 \implies y = 29$.

Q1048. Equations: $x + y = 32$ and $x - y = 14$. Adding gives $2x = 46 \implies x = 23$. Subtracting gives $2y = 18 \implies y = 9$.

Q1049. Equations: $x + y = 120$ and $x - y = 26$. Adding gives $2x = 146 \implies x = 73$. Subtracting gives $2y = 94 \implies y = 47$.

Q1050. Equations: $x + y = 106$ and $x - y = 20$. Adding gives $2x = 126 \implies x = 63$. Subtracting gives $2y = 86 \implies y = 43$.

Q1051. Equations: $x + y = 71$ and $x - y = 11$. Adding gives $2x = 82 \implies x = 41$. Subtracting gives $2y = 60 \implies y = 30$.

Q1052. Equations: $x + y = 80$ and $x - y = 8$. Adding gives $2x = 88 \implies x = 44$. Subtracting gives $2y = 72 \implies y = 36$.

Q1053. Equations: $x + y = 95$ and $x - y = 13$. Adding gives $2x = 108 \implies x = 54$. Subtracting gives $2y = 82 \implies y = 41$.

Q1054. Equations: $x + y = 94$ and $x - y = 12$. Adding gives $2x = 106 \implies x = 53$. Subtracting gives $2y = 82 \implies y = 41$.

Q1055. Equations: $x + y = 72$ and $x - y = 16$. Adding gives $2x = 88 \implies x = 44$. Subtracting gives $2y = 56 \implies y = 28$.

Q1056. Equations: $x + y = 41$ and $x - y = 9$. Adding gives $2x = 50 \implies x = 25$. Subtracting gives $2y = 32 \implies y = 16$.

Q1057. Equations: $x + y = 119$ and $x - y = 13$. Adding gives $2x = 132 \implies x = 66$. Subtracting gives $2y = 106 \implies y = 53$.

Q1058. Equations: $x + y = 120$ and $x - y = 16$. Adding gives $2x = 136 \implies x = 68$. Subtracting gives $2y = 104 \implies y = 52$.

Q1059. Equations: $x + y = 75$ and $x - y = 27$. Adding gives $2x = 102 \implies x = 51$. Subtracting gives $2y = 48 \implies y = 24$.

Q1060. Equations: $x + y = 77$ and $x - y = 7$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 70 \implies y = 35$.

Q1061. Equations: $x + y = 32$ and $x - y = 4$. Adding gives $2x = 36 \implies x = 18$. Subtracting gives $2y = 28 \implies y = 14$.

Q1062. Equations: $x + y = 101$ and $x - y = 19$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 82 \implies y = 41$.

Q1063. Equations: $x + y = 108$ and $x - y = 12$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 96 \implies y = 48$.

Q1064. Equations: $x + y = 65$ and $x - y = 21$. Adding gives $2x = 86 \implies x = 43$. Subtracting gives $2y = 44 \implies y = 22$.

Q1065. Equations: $x + y = 32$ and $x - y = 10$. Adding gives $2x = 42 \implies x = 21$. Subtracting gives $2y = 22 \implies y = 11$.

Q1066. Equations: $x + y = 102$ and $x - y = 20$. Adding gives $2x = 122 \implies x = 61$. Subtracting gives $2y = 82 \implies y = 41$.

Q1067. Equations: $x + y = 92$ and $x - y = 18$. Adding gives $2x = 110 \implies x = 55$. Subtracting gives $2y = 74 \implies y = 37$.

Q1068. Equations: $x + y = 41$ and $x - y = 11$. Adding gives $2x = 52 \implies x = 26$. Subtracting gives $2y = 30 \implies y = 15$.

Q1069. Equations: $x + y = 40$ and $x - y = 6$. Adding gives $2x = 46 \implies x = 23$. Subtracting gives $2y = 34 \implies y = 17$.

Q1070. Equations: $x + y = 40$ and $x - y = 20$. Adding gives $2x = 60 \implies x = 30$. Subtracting gives $2y = 20 \implies y = 10$.

Q1071. Equations: $x + y = 105$ and $x - y = 27$. Adding gives $2x = 132 \implies x = 66$. Subtracting gives $2y = 78 \implies y = 39$.

Q1072. Equations: $x + y = 40$ and $x - y = 10$. Adding gives $2x = 50 \implies x = 25$. Subtracting gives $2y = 30 \implies y = 15$.

Q1073. Equations: $x + y = 74$ and $x - y = 24$. Adding gives $2x = 98 \implies x = 49$. Subtracting gives $2y = 50 \implies y = 25$.

Q1074. Equations: $x + y = 120$ and $x - y = 8$. Adding gives $2x = 128 \implies x = 64$. Subtracting gives $2y = 112 \implies y = 56$.

Q1075. Equations: $x + y = 46$ and $x - y = 4$. Adding gives $2x = 50 \implies x = 25$. Subtracting gives $2y = 42 \implies y = 21$.

Q1076. Equations: $x + y = 42$ and $x - y = 28$.

Adding gives $2x = 70 \implies x = 35$. Subtracting gives $2y = 14 \implies y = 7$.

Q1077. Equations: $x + y = 50$ and $x - y = 16$. Adding gives $2x = 66 \implies x = 33$. Subtracting gives $2y = 34 \implies y = 17$.

Q1078. Equations: $x + y = 89$ and $x - y = 9$. Adding gives $2x = 98 \implies x = 49$. Subtracting gives $2y = 80 \implies y = 40$.

Q1079. Equations: $x + y = 82$ and $x - y = 4$. Adding gives $2x = 86 \implies x = 43$. Subtracting gives $2y = 78 \implies y = 39$.

Q1080. Equations: $x + y = 102$ and $x - y = 26$. Adding gives $2x = 128 \implies x = 64$. Subtracting gives $2y = 76 \implies y = 38$.

Q1081. Equations: $x + y = 47$ and $x - y = 11$. Adding gives $2x = 58 \implies x = 29$. Subtracting gives $2y = 36 \implies y = 18$.

Q1082. Equations: $x + y = 71$ and $x - y = 7$. Adding gives $2x = 78 \implies x = 39$. Subtracting gives $2y = 64 \implies y = 32$.

Q1083. Equations: $x + y = 48$ and $x - y = 10$. Adding gives $2x = 58 \implies x = 29$. Subtracting gives $2y = 38 \implies y = 19$.

Q1084. Equations: $x + y = 93$ and $x - y = 9$. Adding gives $2x = 102 \implies x = 51$. Subtracting gives $2y = 84 \implies y = 42$.

Q1085. Equations: $x + y = 32$ and $x - y = 4$. Adding gives $2x = 36 \implies x = 18$. Subtracting gives $2y = 28 \implies y = 14$.

Q1086. Equations: $x + y = 34$ and $x - y = 26$. Adding gives $2x = 60 \implies x = 30$. Subtracting gives $2y = 8 \implies y = 4$.

Q1087. Equations: $x + y = 45$ and $x - y = 13$. Adding gives $2x = 58 \implies x = 29$. Subtracting gives $2y = 32 \implies y = 16$.

Q1088. Equations: $x + y = 81$ and $x - y = 27$. Adding gives $2x = 108 \implies x = 54$. Subtracting gives $2y = 54 \implies y = 27$.

Q1089. Equations: $x + y = 120$ and $x - y = 26$. Adding gives $2x = 146 \implies x = 73$. Subtracting gives $2y = 94 \implies y = 47$.

Q1090. Equations: $x + y = 68$ and $x - y = 8$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 60 \implies y = 30$.

Q1091. Equations: $x + y = 53$ and $x - y = 17$. Adding gives $2x = 70 \implies x = 35$. Subtracting gives $2y = 36 \implies y = 18$.

Q1092. Equations: $x + y = 96$ and $x - y = 12$. Adding gives $2x = 108 \implies x = 54$. Subtracting gives $2y = 84 \implies y = 42$.

Q1093. Equations: $x + y = 94$ and $x - y = 16$. Adding gives $2x = 110 \implies x = 55$. Subtracting gives $2y = 78 \implies y = 39$.

Q1094. Equations: $x + y = 85$ and $x - y = 15$. Adding gives $2x = 100 \implies x = 50$. Subtracting gives $2y = 70 \implies y = 35$.

Q1095. Equations: $x + y = 113$ and $x - y = 15$. Adding gives $2x = 128 \implies x = 64$. Subtracting gives $2y = 98 \implies y = 49$.

Q1096. Equations: $x + y = 57$ and $x - y = 21$. Adding gives $2x = 78 \implies x = 39$. Subtracting gives $2y = 36 \implies y = 18$.

Q1097. Equations: $x + y = 101$ and $x - y = 27$. Adding gives $2x = 128 \implies x = 64$. Subtracting gives $2y = 74 \implies y = 37$.

Q1098. Equations: $x + y = 45$ and $x - y = 17$. Adding gives $2x = 62 \implies x = 31$. Subtracting gives $2y = 28 \implies y = 14$.

Q1099. Equations: $x + y = 113$ and $x - y = 25$. Adding gives $2x = 138 \implies x = 69$. Subtracting gives $2y = 88 \implies y = 44$.

Q1100. Equations: $x + y = 60$ and $x - y = 8$. Adding gives $2x = 68 \implies x = 34$. Subtracting gives $2y = 52 \implies y = 26$.

Q1101. Equations: $x + y = 48$ and $x - y = 18$. Adding gives $2x = 66 \implies x = 33$. Subtracting gives $2y = 30 \implies y = 15$.

Q1102. Equations: $x + y = 103$ and $x - y = 27$. Adding gives $2x = 130 \implies x = 65$. Subtracting gives $2y = 76 \implies y = 38$.

Q1103. Equations: $x + y = 102$ and $x - y = 14$. Adding gives $2x = 116 \implies x = 58$. Subtracting gives $2y = 88 \implies y = 44$.

Q1104. Equations: $x + y = 81$ and $x - y = 27$. Adding gives $2x = 108 \implies x = 54$. Subtracting gives $2y = 54 \implies y = 27$.

Q1105. Equations: $x + y = 110$ and $x - y = 26$. Adding gives $2x = 136 \implies x = 68$. Subtracting gives $2y = 84 \implies y = 42$.

Q1106. Equations: $x + y = 117$ and $x - y = 19$. Adding gives $2x = 136 \implies x = 68$. Subtracting gives $2y = 98 \implies y = 49$.

Q1107. Equations: $x + y = 43$ and $x - y = 27$. Adding gives $2x = 70 \implies x = 35$. Subtracting gives $2y = 16 \implies y = 8$.

Q1108. Equations: $x + y = 52$ and $x - y = 4$. Adding gives $2x = 56 \implies x = 28$. Subtracting gives $2y = 48 \implies y = 24$.

Q1109. Equations: $x + y = 74$ and $x - y = 24$. Adding gives $2x = 98 \implies x = 49$. Subtracting gives $2y = 50 \implies y = 25$.

Q1110. Equations: $x + y = 59$ and $x - y = 25$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 34 \implies y = 17$.

Q1111. Equations: $x + y = 61$ and $x - y = 15$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 46 \implies y = 23$.

Q1112. Equations: $x + y = 119$ and $x - y = 17$. Adding gives $2x = 136 \implies x = 68$. Subtracting gives $2y = 102 \implies y = 51$.

Q1113. Equations: $x + y = 44$ and $x - y = 26$. Adding gives $2x = 70 \implies x = 35$. Subtracting gives $2y = 18 \implies y = 9$.

Q1114. Equations: $x + y = 65$ and $x - y = 19$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 46 \implies y = 23$.

Q1115. Equations: $x + y = 66$ and $x - y = 8$. Adding gives $2x = 74 \implies x = 37$. Subtracting gives $2y = 58 \implies y = 29$.

Q1116. Equations: $x + y = 116$ and $x - y = 8$. Adding gives $2x = 124 \implies x = 62$. Subtracting gives $2y = 108 \implies y = 54$.

Q1117. Equations: $x + y = 73$ and $x - y = 5$. Adding gives $2x = 78 \implies x = 39$. Subtracting gives $2y = 68 \implies y = 34$.

Q1118. Equations: $x + y = 51$ and $x - y = 9$. Adding gives $2x = 60 \implies x = 30$. Subtracting gives $2y = 42 \implies y = 21$.

Q1119. Equations: $x + y = 45$ and $x - y = 19$. Adding gives $2x = 64 \implies x = 32$. Subtracting gives $2y = 26 \implies y = 13$.

Q1120. Equations: $x + y = 109$ and $x - y = 11$. Adding gives $2x = 120 \implies x = 60$. Subtracting gives $2y = 98 \implies y = 49$.

Q1121. Equations: $x + y = 33$ and $x - y = 11$. Adding gives $2x = 44 \implies x = 22$. Subtracting gives $2y = 22 \implies y = 11$.

Q1122. Equations: $x + y = 75$ and $x - y = 9$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 66 \implies y = 33$.

Q1123. Equations: $x + y = 83$ and $x - y = 23$. Adding gives $2x = 106 \implies x = 53$. Subtracting gives $2y = 60 \implies y = 30$.

Q1124. Equations: $x + y = 81$ and $x - y = 9$. Adding gives $2x = 90 \implies x = 45$. Subtracting gives $2y = 72 \implies y = 36$.

Q1125. Equations: $x + y = 98$ and $x - y = 14$. Adding gives $2x = 112 \implies x = 56$. Subtracting gives $2y = 84 \implies y = 42$.

Q1126. Equations: $x + y = 46$ and $x - y = 18$. Adding gives $2x = 64 \implies x = 32$. Subtracting gives $2y = 28 \implies y = 14$.

Q1127. Equations: $x + y = 40$ and $x - y = 4$. Adding gives $2x = 44 \implies x = 22$. Subtracting gives $2y = 36 \implies y = 18$.

Q1128. Equations: $x + y = 113$ and $x - y = 11$. Adding gives $2x = 124 \implies x = 62$. Subtracting gives $2y = 102 \implies y = 51$.

Q1129. Equations: $x + y = 36$ and $x - y = 10$. Adding gives $2x = 46 \implies x = 23$. Subtracting gives $2y = 26 \implies y = 13$.

Q1130. Equations: $x + y = 68$ and $x - y = 20$. Adding gives $2x = 88 \implies x = 44$. Subtracting gives $2y = 48 \implies y = 24$.

Q1131. Equations: $x + y = 97$ and $x - y = 15$. Adding gives $2x = 112 \implies x = 56$. Subtracting gives $2y = 82 \implies y = 41$.

Q1132. Equations: $x + y = 62$ and $x - y = 22$. Adding gives $2x = 84 \implies x = 42$. Subtracting gives $2y = 40 \implies y = 20$.

Q1133. Equations: $x + y = 74$ and $x - y = 26$. Adding gives $2x = 100 \implies x = 50$. Subtracting gives $2y = 48 \implies y = 24$.

Q1134. Equations: $x + y = 114$ and $x - y = 28$. Adding gives $2x = 142 \implies x = 71$. Subtracting gives $2y = 86 \implies y = 43$.

Q1135. Equations: $x + y = 121$ and $x - y = 27$. Adding gives $2x = 148 \implies x = 74$. Subtracting gives $2y = 94 \implies y = 47$.

Q1136. Equations: $x + y = 34$ and $x - y = 28$. Adding gives $2x = 62 \implies x = 31$. Subtracting gives $2y = 6 \implies y = 3$.

Q1137. Equations: $x + y = 41$ and $x - y = 7$. Adding gives $2x = 48 \implies x = 24$. Subtracting gives $2y = 34 \implies y = 17$.

Q1138. Equations: $x + y = 107$ and $x - y = 17$. Adding gives $2x = 124 \implies x = 62$. Subtracting gives $2y = 90 \implies y = 45$.

Q1139. Equations: $x + y = 53$ and $x - y = 15$. Adding gives $2x = 68 \implies x = 34$. Subtracting gives $2y = 38 \implies y = 19$.

Q1140. Equations: $x + y = 110$ and $x - y = 24$. Adding gives $2x = 134 \implies x = 67$. Subtracting gives $2y = 86 \implies y = 43$.

Q1141. Equations: $x + y = 38$ and $x - y = 28$. Adding gives $2x = 66 \implies x = 33$. Subtracting gives $2y = 10 \implies y = 5$.

Q1142. Equations: $x + y = 52$ and $x - y = 24$. Adding gives $2x = 76 \implies x = 38$. Subtracting gives $2y = 28 \implies y = 14$.

Q1143. Equations: $x + y = 33$ and $x - y = 7$. Adding gives $2x = 40 \implies x = 20$. Subtracting gives $2y = 26 \implies y = 13$.

Q1144. Equations: $x + y = 49$ and $x - y = 5$. Adding gives $2x = 54 \implies x = 27$. Subtracting gives $2y = 44 \implies y = 22$.

Q1145. Equations: $x + y = 81$ and $x - y = 11$. Adding gives $2x = 92 \implies x = 46$. Subtracting gives $2y = 70 \implies y = 35$.

Q1146. Equations: $x + y = 97$ and $x - y = 27$. Adding gives $2x = 124 \implies x = 62$. Subtracting gives $2y = 70 \implies y = 35$.

Q1147. Equations: $x + y = 87$ and $x - y = 13$. Adding gives $2x = 100 \implies x = 50$. Subtracting gives $2y = 74 \implies y = 37$.

Q1148. Equations: $x + y = 79$ and $x - y = 25$. Adding gives $2x = 104 \implies x = 52$. Subtracting gives $2y = 54 \implies y = 27$.

Q1149. Equations: $x + y = 58$ and $x - y = 14$. Adding gives $2x = 72 \implies x = 36$. Subtracting gives $2y = 44 \implies y = 22$.

Q1150. Equations: $x + y = 60$ and $x - y = 14$. Adding gives $2x = 74 \implies x = 37$. Subtracting gives $2y = 46 \implies y = 23$.

Q1151. Past condition: $x - 9 = 3(y - 9) \implies x - 3y = -18$. Future condition: $x+10 = 3(y+10) \implies x-3y = 20$.

Q1152. Past condition: $x - 9 = 7(y - 9) \implies x - 7y = -54$. Future condition: $x+8 = 4(y+8) \implies x-4y = 24$.

Q1153. Past condition: $x - 5 = 5(y - 5) \implies x - 5y = -20$. Future condition: $x+10 = 4(y+10) \implies x-4y = 30$.

Q1154. Past condition: $x - 10 = 3(y - 10) \implies x - 3y =$

-20. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1155. Past condition: $x-5=5(y-5) \implies x-5y=-20$. Future condition: $x+6=3(y+6) \implies x-3y=12$.

Q1156. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1157. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1158. Past condition: $x-7=5(y-7) \implies x-5y=-28$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1159. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1160. Past condition: $x-8=3(y-8) \implies x-3y=-16$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1161. Past condition: $x-7=4(y-7) \implies x-4y=-21$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1162. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+7=4(y+7) \implies x-4y=21$.

Q1163. Past condition: $x-10=6(y-10) \implies x-6y=-50$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1164. Past condition: $x-4=7(y-4) \implies x-7y=-24$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1165. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1166. Past condition: $x-4=7(y-4) \implies x-7y=-24$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1167. Past condition: $x-5=3(y-5) \implies x-3y=-10$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1168. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1169. Past condition: $x-10=7(y-10) \implies x-7y=-60$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1170. Past condition: $x-5=3(y-5) \implies x-3y=-10$. Future condition: $x+6=3(y+6) \implies x-3y=12$.

Q1171. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1172. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1173. Past condition: $x-8=3(y-8) \implies x-3y=-16$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1174. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1175. Past condition: $x-6=7(y-6) \implies x-7y=-36$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1176. Past condition: $x-9=5(y-9) \implies x-5y=-36$. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1177. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+5=2(y+5) \implies x-2y=5$.

Q1178. Past condition: $x-3=7(y-3) \implies x-7y=-18$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1179. Past condition: $x-8=4(y-8) \implies x-4y=-24$. Future condition: $x+10=3(y+10) \implies x-3y=20$.

Q1180. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1181. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+3=4(y+3) \implies x-4y=9$.

Q1182. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1183. Past condition: $x-8=4(y-8) \implies x-4y=-24$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1184. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1185. Past condition: $x-5=6(y-5) \implies x-6y=-25$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1186. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1187. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1188. Past condition: $x-7=4(y-7) \implies x-4y=-21$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1189. Past condition: $x-8=3(y-8) \implies x-3y=-16$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1190. Past condition: $x-4=7(y-4) \implies x-7y=-24$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1191. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+6=2(y+6) \implies x-2y=6$.

Q1192. Past condition: $x-8=4(y-8) \implies x-4y=-24$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1193. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1194. Past condition: $x-10=4(y-10) \implies x-4y=-30$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1195. Past condition: $x-10=4(y-10) \implies x-4y=-30$. Future condition: $x+8=4(y+8) \implies x-4y=24$.

Q1196. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1197. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1198. Past condition: $x-5=6(y-5) \implies x-6y=-25$. Future condition: $x+7=4(y+7) \implies x-4y=21$.

Q1199. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+5=2(y+5) \implies x-2y=5$.

Q1200. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1201. Past condition: $x-10=7(y-10) \implies x-7y=-60$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1202. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1203. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1204. Past condition: $x-6=7(y-6) \implies x-7y=-36$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1205. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1206. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1207. Past condition: $x-3=3(y-3) \implies x-3y=-6$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1208. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+5=4(y+5) \implies x-4y=15$.

Q1209. Past condition: $x-9=5(y-9) \implies x-5y=$

-36. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1210. Past condition: $x-9=5(y-9) \implies x-5y=-36$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1211. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1212. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+3=4(y+3) \implies x-4y=9$.

Q1213. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1214. Past condition: $x-10=5(y-10) \implies x-5y=-40$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1215. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1216. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1217. Past condition: $x-5=5(y-5) \implies x-5y=-20$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1218. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1219. Past condition: $x-4=5(y-4) \implies x-5y=-16$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1220. Past condition: $x-5=4(y-5) \implies x-4y=-15$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1221. Past condition: $x-9=6(y-9) \implies x-6y=-45$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1222. Past condition: $x-10=7(y-10) \implies x-7y=-60$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1223. Past condition: $x-8=5(y-8) \implies x-5y=-32$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1224. Past condition: $x-6=3(y-6) \implies x-3y=-12$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1225. Past condition: $x-8=5(y-8) \implies x-5y=-32$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1226. Past condition: $x-5=5(y-5) \implies x-5y=-20$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1227. Past condition: $x-5=5(y-5) \implies x-5y=-20$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1228. Past condition: $x-5=5(y-5) \implies x-5y=-20$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1229. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1230. Past condition: $x-4=7(y-4) \implies x-7y=-24$. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1231. Past condition: $x-7=4(y-7) \implies x-4y=-21$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1232. Past condition: $x-10=5(y-10) \implies x-5y=-40$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1233. Past condition: $x-10=5(y-10) \implies x-5y=-40$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1234. Past condition: $x-10=7(y-10) \implies x-7y=-60$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1235. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1236. Past condition: $x-10=7(y-10) \implies x-7y=-60$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1237. Past condition: $x-3=7(y-3) \implies x-7y=-18$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1238. Past condition: $x-5=6(y-5) \implies x-6y=-25$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1239. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1240. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1241. Past condition: $x-8=5(y-8) \implies x-5y=-32$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1242. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1243. Past condition: $x-7=6(y-7) \implies x-6y=-35$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1244. Past condition: $x-7=6(y-7) \implies x-6y=-35$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1245. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1246. Past condition: $x-8=4(y-8) \implies x-4y=-24$. Future condition: $x+5=4(y+5) \implies x-4y=15$.

Q1247. Past condition: $x-7=5(y-7) \implies x-5y=-28$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1248. Past condition: $x-3=6(y-3) \implies x-6y=-15$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1249. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+8=4(y+8) \implies x-4y=24$.

Q1250. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1251. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1252. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+7=4(y+7) \implies x-4y=21$.

Q1253. Past condition: $x-6=3(y-6) \implies x-3y=-12$. Future condition: $x+7=4(y+7) \implies x-4y=21$.

Q1254. Past condition: $x-10=6(y-10) \implies x-6y=-50$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1255. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1256. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+5=4(y+5) \implies x-4y=15$.

Q1257. Past condition: $x-8=3(y-8) \implies x-3y=-16$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1258. Past condition: $x-3=5(y-3) \implies x-5y=-12$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1259. Past condition: $x-5=7(y-5) \implies x-7y=-30$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1260. Past condition: $x-3=5(y-3) \implies x-5y=-12$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1261. Past condition: $x-10=4(y-10) \implies x-4y=-30$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1262. Past condition: $x-9=6(y-9) \implies x-6y=-45$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1263. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+3=4(y+3) \implies x-4y=9$.

Q1264. Past condition: $x-3=7(y-3) \implies x-7y=$

-18. Future condition: $x+5=4(y+5) \implies x-4y=15$.

Q1265. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1266. Past condition: $x-6=6(y-6) \implies x-6y=-30$. Future condition: $x+3=4(y+3) \implies x-4y=9$.

Q1267. Past condition: $x-4=4(y-4) \implies x-4y=-12$. Future condition: $x+4=3(y+4) \implies x-3y=8$.

Q1268. Past condition: $x-8=7(y-8) \implies x-7y=-48$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1269. Past condition: $x-9=6(y-9) \implies x-6y=-45$. Future condition: $x+5=2(y+5) \implies x-2y=5$.

Q1270. Past condition: $x-7=6(y-7) \implies x-6y=-35$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1271. Past condition: $x-9=6(y-9) \implies x-6y=-45$. Future condition: $x+10=4(y+10) \implies x-4y=30$.

Q1272. Past condition: $x-9=5(y-9) \implies x-5y=-36$. Future condition: $x+7=2(y+7) \implies x-2y=7$.

Q1273. Past condition: $x-10=3(y-10) \implies x-3y=-20$. Future condition: $x+5=4(y+5) \implies x-4y=15$.

Q1274. Past condition: $x-4=7(y-4) \implies x-7y=-24$. Future condition: $x+3=3(y+3) \implies x-3y=6$.

Q1275. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+10=3(y+10) \implies x-3y=20$.

Q1276. Past condition: $x-3=6(y-3) \implies x-6y=-15$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1277. Past condition: $x-10=5(y-10) \implies x-5y=-40$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1278. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+5=2(y+5) \implies x-2y=5$.

Q1279. Past condition: $x-4=4(y-4) \implies x-4y=-12$. Future condition: $x+8=3(y+8) \implies x-3y=16$.

Q1280. Past condition: $x-4=5(y-4) \implies x-5y=-16$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1281. Past condition: $x-7=7(y-7) \implies x-7y=-42$. Future condition: $x+6=4(y+6) \implies x-4y=18$.

Q1282. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1283. Past condition: $x-3=3(y-3) \implies x-3y=-6$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1284. Past condition: $x-9=6(y-9) \implies x-6y=-45$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1285. Past condition: $x-3=3(y-3) \implies x-3y=-6$. Future condition: $x+8=2(y+8) \implies x-2y=8$.

Q1286. Past condition: $x-6=5(y-6) \implies x-5y=-24$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1287. Past condition: $x-8=6(y-8) \implies x-6y=-40$. Future condition: $x+6=2(y+6) \implies x-2y=6$.

Q1288. Past condition: $x-3=5(y-3) \implies x-5y=-12$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1289. Past condition: $x-3=4(y-3) \implies x-4y=-9$. Future condition: $x+7=3(y+7) \implies x-3y=14$.

Q1290. Past condition: $x-6=3(y-6) \implies x-3y=-12$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1291. Past condition: $x-9=7(y-9) \implies x-7y=-54$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1292. Past condition: $x-4=3(y-4) \implies x-3y=-8$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1293. Past condition: $x-4=6(y-4) \implies x-6y=-20$. Future condition: $x+4=4(y+4) \implies x-4y=12$.

Q1294. Past condition: $x-7=3(y-7) \implies x-3y=-14$. Future condition: $x+3=2(y+3) \implies x-2y=3$.

Q1295. Past condition: $x-9=4(y-9) \implies x-4y=-27$. Future condition: $x+9=3(y+9) \implies x-3y=18$.

Q1296. Past condition: $x-9=5(y-9) \implies x-5y=-36$. Future condition: $x+9=4(y+9) \implies x-4y=27$.

Q1297. Past condition: $x-8=3(y-8) \implies x-3y=-16$. Future condition: $x+4=2(y+4) \implies x-2y=4$.

Q1298. Past condition: $x-10=4(y-10) \implies x-4y=-30$. Future condition: $x+10=2(y+10) \implies x-2y=10$.

Q1299. Past condition: $x-6=4(y-6) \implies x-4y=-18$. Future condition: $x+5=3(y+5) \implies x-3y=10$.

Q1300. Past condition: $x-7=4(y-7) \implies x-4y=-21$. Future condition: $x+9=2(y+9) \implies x-2y=9$.

Q1301. Equations: (1) $x+12y=200$, (2) $x+21y=335$. Subtracting gives $9y=135 \implies y=15$. Then $x=20$.

Q1302. Equations: (1) $x+14y=198$, (2) $x+24y=328$. Subtracting gives $10y=130 \implies y=13$. Then $x=16$.

Q1303. Equations: (1) $x+6y=123$, (2) $x+20y=305$. Subtracting gives $14y=182 \implies y=13$. Then $x=45$.

Q1304. Equations: (1) $x+8y=90$, (2) $x+20y=210$. Subtracting gives $12y=120 \implies y=10$. Then $x=10$.

Q1305. Equations: (1) $x+7y=76$, (2) $x+18y=142$. Subtracting gives $11y=66 \implies y=6$. Then $x=34$.

Q1306. Equations: (1) $x+11y=204$, (2) $x+28y=442$. Subtracting gives $17y=238 \implies y=14$. Then $x=50$.

Q1307. Equations: (1) $x+13y=227$, (2) $x+21y=339$. Subtracting gives $8y=112 \implies y=14$. Then $x=45$.

Q1308. Equations: (1) $x+11y=95$, (2) $x+27y=175$. Subtracting gives $16y=80 \implies y=5$. Then $x=40$.

Q1309. Equations: (1) $x+5y=86$, (2) $x+30y=411$. Subtracting gives $25y=325 \implies y=13$. Then $x=21$.

Q1310. Equations: (1) $x+14y=229$, (2) $x+18y=285$. Subtracting gives $4y=56 \implies y=14$. Then $x=33$.

Q1311. Equations: (1) $x+10y=143$, (2) $x+24y=325$. Subtracting gives $14y=182 \implies y=13$. Then $x=13$.

Q1312. Equations: (1) $x+13y=129$, (2) $x+16y=150$. Subtracting gives $3y=21 \implies y=7$. Then $x=38$.

Q1313. Equations: (1) $x+10y=113$, (2) $x+18y=185$. Subtracting gives $8y=72 \implies y=9$. Then $x=23$.

Q1314. Equations: (1) $x+10y=150$, (2) $x+29y=378$. Subtracting gives $19y=228 \implies y=12$. Then $x=30$.

Q1315. Equations: (1) $x+13y=111$, (2) $x+17y=135$. Subtracting gives $4y=24 \implies y=6$. Then $x=33$.

Q1316. Equations: (1) $x+8y=103$, (2) $x+25y=222$. Subtracting gives $17y=119 \implies y=7$. Then $x=47$.

Q1317. Equations: (1) $x+10y=103$, (2) $x+28y=211$. Subtracting gives $18y=108 \implies y=6$. Then $x=43$.

Q1318. Equations: (1) $x+11y=168$, (2) $x+23y=312$. Subtracting gives $12y=144 \implies y=12$. Then $x=36$.

Q1319. Equations: (1) $x+5y=94$, (2) $x+23y=310$.

Subtracting gives $18y = 216 \implies y = 12$. Then $x = 34$.

Q1320. Equations: (1) $x + 5y = 56$, (2) $x + 29y = 200$. Subtracting gives $24y = 144 \implies y = 6$. Then $x = 26$.

Q1321. Equations: (1) $x + 5y = 74$, (2) $x + 24y = 226$. Subtracting gives $19y = 152 \implies y = 8$. Then $x = 34$.

Q1322. Equations: (1) $x + 15y = 197$, (2) $x + 16y = 208$. Subtracting gives $1y = 11 \implies y = 11$. Then $x = 32$.

Q1323. Equations: (1) $x + 8y = 112$, (2) $x + 29y = 343$. Subtracting gives $21y = 231 \implies y = 11$. Then $x = 24$.

Q1324. Equations: (1) $x + 14y = 213$, (2) $x + 25y = 367$. Subtracting gives $11y = 154 \implies y = 14$. Then $x = 17$.

Q1325. Equations: (1) $x + 10y = 188$, (2) $x + 16y = 278$. Subtracting gives $6y = 90 \implies y = 15$. Then $x = 38$.

Q1326. Equations: (1) $x + 9y = 164$, (2) $x + 21y = 344$. Subtracting gives $12y = 180 \implies y = 15$. Then $x = 29$.

Q1327. Equations: (1) $x + 12y = 121$, (2) $x + 17y = 166$. Subtracting gives $5y = 45 \implies y = 9$. Then $x = 13$.

Q1328. Equations: (1) $x + 5y = 73$, (2) $x + 25y = 253$. Subtracting gives $20y = 180 \implies y = 9$. Then $x = 28$.

Q1329. Equations: (1) $x + 12y = 114$, (2) $x + 26y = 198$. Subtracting gives $14y = 84 \implies y = 6$. Then $x = 42$.

Q1330. Equations: (1) $x + 13y = 238$, (2) $x + 22y = 373$. Subtracting gives $9y = 135 \implies y = 15$. Then $x = 43$.

Q1331. Equations: (1) $x + 6y = 107$, (2) $x + 20y = 247$. Subtracting gives $14y = 140 \implies y = 10$. Then $x = 47$.

Q1332. Equations: (1) $x + 10y = 97$, (2) $x + 28y = 241$. Subtracting gives $18y = 144 \implies y = 8$. Then $x = 17$.

Q1333. Equations: (1) $x + 5y = 101$, (2) $x + 20y = 296$. Subtracting gives $15y = 195 \implies y = 13$. Then $x = 36$.

Q1334. Equations: (1) $x + 9y = 120$, (2) $x + 21y = 228$. Subtracting gives $12y = 108 \implies y = 9$. Then $x = 39$.

Q1335. Equations: (1) $x + 13y = 139$, (2) $x + 16y = 166$. Subtracting gives $3y = 27 \implies y = 9$. Then $x = 22$.

Q1336. Equations: (1) $x + 8y = 81$, (2) $x + 23y = 201$. Subtracting gives $15y = 120 \implies y = 8$. Then $x = 17$.

Q1337. Equations: (1) $x + 6y = 74$, (2) $x + 21y = 149$. Subtracting gives $15y = 75 \implies y = 5$. Then $x = 44$.

Q1338. Equations: (1) $x + 10y = 132$, (2) $x + 22y = 240$. Subtracting gives $12y = 108 \implies y = 9$. Then $x = 42$.

Q1339. Equations: (1) $x + 14y = 122$, (2) $x + 28y = 234$. Subtracting gives $14y = 112 \implies y = 8$. Then $x = 10$.

Q1340. Equations: (1) $x + 9y = 70$, (2) $x + 30y = 196$. Subtracting gives $21y = 126 \implies y = 6$. Then $x = 16$.

Q1341. Equations: (1) $x + 8y = 157$, (2) $x + 28y = 457$. Subtracting gives $20y = 300 \implies y = 15$. Then $x = 37$.

Q1342. Equations: (1) $x + 13y = 156$, (2) $x + 18y = 211$. Subtracting gives $5y = 55 \implies y = 11$. Then $x = 13$.

Q1343. Equations: (1) $x + 6y = 60$, (2) $x + 30y = 252$. Subtracting gives $24y = 192 \implies y = 8$. Then $x = 12$.

Q1344. Equations: (1) $x + 11y = 140$, (2) $x + 17y = 206$. Subtracting gives $6y = 66 \implies y = 11$. Then $x = 19$.

Q1345. Equations: (1) $x + 5y = 95$, (2) $x + 21y = 239$. Subtracting gives $16y = 144 \implies y = 9$. Then $x = 50$.

Q1346. Equations: (1) $x + 11y = 143$, (2) $x + 30y = 352$. Subtracting gives $19y = 209 \implies y = 11$. Then $x = 22$.

Q1347. Equations: (1) $x + 5y = 49$, (2) $x + 20y = 139$. Subtracting gives $15y = 90 \implies y = 6$. Then $x = 19$.

Q1348. Equations: (1) $x + 10y = 103$, (2) $x + 28y = 265$. Subtracting gives $18y = 162 \implies y = 9$. Then $x = 13$.

Q1349. Equations: (1) $x + 11y = 96$, (2) $x + 23y = 168$. Subtracting gives $12y = 72 \implies y = 6$. Then $x = 30$.

Q1350. Equations: (1) $x + 8y = 143$, (2) $x + 18y = 293$. Subtracting gives $10y = 150 \implies y = 15$. Then $x = 23$.

Q1351. Equations: (1) $x + 13y = 155$, (2) $x + 24y = 276$. Subtracting gives $11y = 121 \implies y = 11$. Then $x = 12$.

Q1352. Equations: (1) $x + 14y = 124$, (2) $x + 25y = 190$. Subtracting gives $11y = 66 \implies y = 6$. Then $x = 40$.

Q1353. Equations: (1) $x + 5y = 73$, (2) $x + 18y = 216$. Subtracting gives $13y = 143 \implies y = 11$. Then $x = 18$.

Q1354. Equations: (1) $x + 11y = 105$, (2) $x + 24y = 196$. Subtracting gives $13y = 91 \implies y = 7$. Then $x = 28$.

Q1355. Equations: (1) $x + 6y = 62$, (2) $x + 24y = 206$. Subtracting gives $18y = 144 \implies y = 8$. Then $x = 14$.

Q1356. Equations: (1) $x + 5y = 70$, (2) $x + 26y = 301$. Subtracting gives $21y = 231 \implies y = 11$. Then $x = 15$.

Q1357. Equations: (1) $x + 7y = 97$, (2) $x + 21y = 223$. Subtracting gives $14y = 126 \implies y = 9$. Then $x = 34$.

Q1358. Equations: (1) $x + 12y = 148$, (2) $x + 28y = 292$. Subtracting gives $16y = 144 \implies y = 9$. Then $x = 40$.

Q1359. Equations: (1) $x + 12y = 123$, (2) $x + 28y = 251$. Subtracting gives $16y = 128 \implies y = 8$. Then $x = 27$.

Q1360. Equations: (1) $x + 14y = 184$, (2) $x + 29y = 364$. Subtracting gives $15y = 180 \implies y = 12$. Then $x = 16$.

Q1361. Equations: (1) $x + 10y = 187$, (2) $x + 25y = 397$. Subtracting gives $15y = 210 \implies y = 14$. Then $x = 47$.

Q1362. Equations: (1) $x + 15y = 177$, (2) $x + 24y = 267$. Subtracting gives $9y = 90 \implies y = 10$. Then $x = 27$.

Q1363. Equations: (1) $x + 15y = 236$, (2) $x + 19y = 288$. Subtracting gives $4y = 52 \implies y = 13$. Then $x = 41$.

Q1364. Equations: (1) $x + 7y = 97$, (2) $x + 27y = 317$. Subtracting gives $20y = 220 \implies y = 11$. Then $x = 20$.

Q1365. Equations: (1) $x + 15y = 113$, (2) $x + 25y = 173$. Subtracting gives $10y = 60 \implies y = 6$. Then $x = 23$.

Q1366. Equations: (1) $x + 12y = 134$, (2) $x + 24y = 218$. Subtracting gives $12y = 84 \implies y = 7$. Then $x = 50$.

Q1367. Equations: (1) $x + 6y = 80$, (2) $x + 30y = 296$. Subtracting gives $24y = 216 \implies y = 9$. Then $x = 26$.

Q1368. Equations: (1) $x + 7y = 96$, (2) $x + 22y = 216$. Subtracting gives $15y = 120 \implies y = 8$. Then $x = 40$.

Q1369. Equations: (1) $x + 7y = 79$, (2) $x + 23y = 207$. Subtracting gives $16y = 128 \implies y = 8$. Then $x = 23$.

Q1370. Equations: (1) $x + 6y = 88$, (2) $x + 18y = 172$. Subtracting gives $12y = 84 \implies y = 7$. Then $x = 46$.

Q1371. Equations: (1) $x + 11y = 112$, (2) $x + 19y = 160$. Subtracting gives $8y = 48 \implies y = 6$. Then $x = 46$.

Q1372. Equations: (1) $x + 5y = 58$, (2) $x + 21y = 154$. Subtracting gives $16y = 96 \implies y = 6$. Then $x = 28$.

Q1373. Equations: (1) $x + 5y = 47$, (2) $x + 18y = 125$. Subtracting gives $13y = 78 \implies y = 6$. Then $x = 17$.

Q1374. Equations: (1) $x + 8y = 85$, (2) $x + 29y = 253$.

Subtracting gives $21y = 168 \implies y = 8$. Then $x = 21$.

Q1375. Equations: (1) $x + 9y = 105$, (2) $x + 20y = 193$. Subtracting gives $11y = 88 \implies y = 8$. Then $x = 33$.

Q1376. Equations: (1) $x + 15y = 168$, (2) $x + 17y = 188$. Subtracting gives $2y = 20 \implies y = 10$. Then $x = 18$.

Q1377. Equations: (1) $x + 11y = 88$, (2) $x + 18y = 123$. Subtracting gives $7y = 35 \implies y = 5$. Then $x = 33$.

Q1378. Equations: (1) $x + 14y = 199$, (2) $x + 24y = 309$. Subtracting gives $10y = 110 \implies y = 11$. Then $x = 45$.

Q1379. Equations: (1) $x + 9y = 89$, (2) $x + 17y = 137$. Subtracting gives $8y = 48 \implies y = 6$. Then $x = 35$.

Q1380. Equations: (1) $x + 6y = 91$, (2) $x + 17y = 212$. Subtracting gives $11y = 121 \implies y = 11$. Then $x = 25$.

Q1381. Equations: (1) $x + 14y = 123$, (2) $x + 27y = 214$. Subtracting gives $13y = 91 \implies y = 7$. Then $x = 25$.

Q1382. Equations: (1) $x + 13y = 116$, (2) $x + 20y = 165$. Subtracting gives $7y = 49 \implies y = 7$. Then $x = 25$.

Q1383. Equations: (1) $x + 14y = 257$, (2) $x + 29y = 482$. Subtracting gives $15y = 225 \implies y = 15$. Then $x = 47$.

Q1384. Equations: (1) $x + 12y = 118$, (2) $x + 24y = 214$. Subtracting gives $12y = 96 \implies y = 8$. Then $x = 22$.

Q1385. Equations: (1) $x + 9y = 111$, (2) $x + 22y = 215$. Subtracting gives $13y = 104 \implies y = 8$. Then $x = 39$.

Q1386. Equations: (1) $x + 5y = 67$, (2) $x + 29y = 331$. Subtracting gives $24y = 264 \implies y = 11$. Then $x = 12$.

Q1387. Equations: (1) $x + 6y = 71$, (2) $x + 25y = 185$. Subtracting gives $19y = 114 \implies y = 6$. Then $x = 35$.

Q1388. Equations: (1) $x + 8y = 123$, (2) $x + 19y = 255$. Subtracting gives $11y = 132 \implies y = 12$. Then $x = 27$.

Q1389. Equations: (1) $x + 11y = 209$, (2) $x + 22y = 374$. Subtracting gives $11y = 165 \implies y = 15$. Then $x = 44$.

Q1390. Equations: (1) $x + 8y = 133$, (2) $x + 21y = 315$. Subtracting gives $13y = 182 \implies y = 14$. Then $x = 21$.

Q1391. Equations: (1) $x + 7y = 84$, (2) $x + 25y = 210$. Subtracting gives $18y = 126 \implies y = 7$. Then $x = 35$.

Q1392. Equations: (1) $x + 14y = 234$, (2) $x + 24y = 384$. Subtracting gives $10y = 150 \implies y = 15$. Then $x = 24$.

Q1393. Equations: (1) $x + 5y = 106$, (2) $x + 30y = 406$. Subtracting gives $25y = 300 \implies y = 12$. Then $x = 46$.

Q1394. Equations: (1) $x + 9y = 185$, (2) $x + 20y = 350$. Subtracting gives $11y = 165 \implies y = 15$. Then $x = 50$.

Q1395. Equations: (1) $x + 14y = 125$, (2) $x + 28y = 209$. Subtracting gives $14y = 84 \implies y = 6$. Then $x = 41$.

Q1396. Equations: (1) $x + 7y = 75$, (2) $x + 21y = 187$. Subtracting gives $14y = 112 \implies y = 8$. Then $x = 19$.

Q1397. Equations: (1) $x + 15y = 178$, (2) $x + 19y = 218$. Subtracting gives $4y = 40 \implies y = 10$. Then $x = 28$.

Q1398. Equations: (1) $x + 13y = 106$, (2) $x + 20y = 155$. Subtracting gives $7y = 49 \implies y = 7$. Then $x = 15$.

Q1399. Equations: (1) $x + 14y = 224$, (2) $x + 19y = 299$. Subtracting gives $5y = 75 \implies y = 15$. Then $x = 14$.

Q1400. Equations: (1) $x + 14y = 201$, (2) $x + 27y = 370$. Subtracting gives $13y = 169 \implies y = 13$. Then $x = 19$.

Q1401. Equations: (1) $x + 6y = 79$, (2) $x + 24y = 241$. Subtracting gives $18y = 162 \implies y = 9$. Then $x = 25$.

Q1402. Equations: (1) $x + 6y = 80$, (2) $x + 25y = 194$. Subtracting gives $19y = 114 \implies y = 6$. Then $x = 44$.

Q1403. Equations: (1) $x + 11y = 91$, (2) $x + 16y = 126$. Subtracting gives $5y = 35 \implies y = 7$. Then $x = 14$.

Q1404. Equations: (1) $x + 13y = 172$, (2) $x + 19y = 244$. Subtracting gives $6y = 72 \implies y = 12$. Then $x = 16$.

Q1405. Equations: (1) $x + 7y = 82$, (2) $x + 20y = 160$. Subtracting gives $13y = 78 \implies y = 6$. Then $x = 40$.

Q1406. Equations: (1) $x + 13y = 204$, (2) $x + 21y = 308$. Subtracting gives $8y = 104 \implies y = 13$. Then $x = 35$.

Q1407. Equations: (1) $x + 8y = 142$, (2) $x + 27y = 370$. Subtracting gives $19y = 228 \implies y = 12$. Then $x = 46$.

Q1408. Equations: (1) $x + 8y = 127$, (2) $x + 29y = 358$. Subtracting gives $21y = 231 \implies y = 11$. Then $x = 39$.

Q1409. Equations: (1) $x + 8y = 123$, (2) $x + 22y = 319$. Subtracting gives $14y = 196 \implies y = 14$. Then $x = 11$.

Q1410. Equations: (1) $x + 5y = 94$, (2) $x + 30y = 469$. Subtracting gives $25y = 375 \implies y = 15$. Then $x = 19$.

Q1411. Equations: (1) $x + 7y = 69$, (2) $x + 18y = 124$. Subtracting gives $11y = 55 \implies y = 5$. Then $x = 34$.

Q1412. Equations: (1) $x + 15y = 240$, (2) $x + 17y = 268$. Subtracting gives $2y = 28 \implies y = 14$. Then $x = 30$.

Q1413. Equations: (1) $x + 13y = 111$, (2) $x + 17y = 135$. Subtracting gives $4y = 24 \implies y = 6$. Then $x = 33$.

Q1414. Equations: (1) $x + 9y = 167$, (2) $x + 24y = 392$. Subtracting gives $15y = 225 \implies y = 15$. Then $x = 32$.

Q1415. Equations: (1) $x + 12y = 88$, (2) $x + 28y = 184$. Subtracting gives $16y = 96 \implies y = 6$. Then $x = 16$.

Q1416. Equations: (1) $x + 15y = 167$, (2) $x + 16y = 175$. Subtracting gives $1y = 8 \implies y = 8$. Then $x = 47$.

Q1417. Equations: (1) $x + 7y = 70$, (2) $x + 20y = 174$. Subtracting gives $13y = 104 \implies y = 8$. Then $x = 14$.

Q1418. Equations: (1) $x + 9y = 103$, (2) $x + 20y = 180$. Subtracting gives $11y = 77 \implies y = 7$. Then $x = 40$.

Q1419. Equations: (1) $x + 10y = 173$, (2) $x + 28y = 443$. Subtracting gives $18y = 270 \implies y = 15$. Then $x = 23$.

Q1420. Equations: (1) $x + 9y = 122$, (2) $x + 26y = 292$. Subtracting gives $17y = 170 \implies y = 10$. Then $x = 32$.

Q1421. Equations: (1) $x + 12y = 180$, (2) $x + 17y = 235$. Subtracting gives $5y = 55 \implies y = 11$. Then $x = 48$.

Q1422. Equations: (1) $x + 11y = 189$, (2) $x + 21y = 339$. Subtracting gives $10y = 150 \implies y = 15$. Then $x = 24$.

Q1423. Equations: (1) $x + 9y = 161$, (2) $x + 23y = 371$. Subtracting gives $14y = 210 \implies y = 15$. Then $x = 26$.

Q1424. Equations: (1) $x + 15y = 117$, (2) $x + 28y = 195$. Subtracting gives $13y = 78 \implies y = 6$. Then $x = 27$.

Q1425. Equations: (1) $x + 12y = 163$, (2) $x + 26y = 331$. Subtracting gives $14y = 168 \implies y = 12$. Then $x = 19$.

Q1426. Equations: (1) $x + 13y = 156$, (2) $x + 27y = 282$. Subtracting gives $14y = 126 \implies y = 9$. Then $x = 39$.

Q1427. Equations: (1) $x + 11y = 127$, (2) $x + 18y = 183$. Subtracting gives $7y = 56 \implies y = 8$. Then $x = 39$.

Q1428. Equations: (1) $x + 15y = 228$, (2) $x + 16y = 241$. Subtracting gives $1y = 13 \implies y = 13$. Then $x = 33$.

Q1429. Equations: (1) $x + 9y = 81$, (2) $x + 28y = 176$.

Subtracting gives $19y = 95 \implies y = 5$. Then $x = 36$.

Q1430. Equations: (1) $x + 10y = 155$, (2) $x + 17y = 232$. Subtracting gives $7y = 77 \implies y = 11$. Then $x = 45$.

Q1431. Equations: (1) $x + 12y = 118$, (2) $x + 23y = 184$. Subtracting gives $11y = 66 \implies y = 6$. Then $x = 46$.

Q1432. Equations: (1) $x + 11y = 120$, (2) $x + 29y = 300$. Subtracting gives $18y = 180 \implies y = 10$. Then $x = 10$.

Q1433. Equations: (1) $x + 9y = 164$, (2) $x + 18y = 281$. Subtracting gives $9y = 117 \implies y = 13$. Then $x = 47$.

Q1434. Equations: (1) $x + 10y = 134$, (2) $x + 20y = 254$. Subtracting gives $10y = 120 \implies y = 12$. Then $x = 14$.

Q1435. Equations: (1) $x + 9y = 130$, (2) $x + 16y = 207$. Subtracting gives $7y = 77 \implies y = 11$. Then $x = 31$.

Q1436. Equations: (1) $x + 12y = 208$, (2) $x + 23y = 362$. Subtracting gives $11y = 154 \implies y = 14$. Then $x = 40$.

Q1437. Equations: (1) $x + 5y = 80$, (2) $x + 26y = 248$. Subtracting gives $21y = 168 \implies y = 8$. Then $x = 40$.

Q1438. Equations: (1) $x + 12y = 192$, (2) $x + 17y = 257$. Subtracting gives $5y = 65 \implies y = 13$. Then $x = 36$.

Q1439. Equations: (1) $x + 11y = 105$, (2) $x + 16y = 130$. Subtracting gives $5y = 25 \implies y = 5$. Then $x = 50$.

Q1440. Equations: (1) $x + 5y = 78$, (2) $x + 16y = 177$. Subtracting gives $11y = 99 \implies y = 9$. Then $x = 33$.

Q1441. Equations: (1) $x + 14y = 194$, (2) $x + 27y = 363$. Subtracting gives $13y = 169 \implies y = 13$. Then $x = 12$.

Q1442. Equations: (1) $x + 8y = 110$, (2) $x + 25y = 246$. Subtracting gives $17y = 136 \implies y = 8$. Then $x = 46$.

Q1443. Equations: (1) $x + 10y = 159$, (2) $x + 27y = 346$. Subtracting gives $17y = 187 \implies y = 11$. Then $x = 49$.

Q1444. Equations: (1) $x + 6y = 121$, (2) $x + 18y = 301$. Subtracting gives $12y = 180 \implies y = 15$. Then $x = 31$.

Q1445. Equations: (1) $x + 13y = 178$, (2) $x + 29y = 354$. Subtracting gives $16y = 176 \implies y = 11$. Then $x = 35$.

Q1446. Equations: (1) $x + 10y = 151$, (2) $x + 22y = 307$. Subtracting gives $12y = 156 \implies y = 13$. Then $x = 21$.

Q1447. Equations: (1) $x + 9y = 138$, (2) $x + 23y = 278$. Subtracting gives $14y = 140 \implies y = 10$. Then $x = 48$.

Q1448. Equations: (1) $x + 5y = 102$, (2) $x + 26y = 354$. Subtracting gives $21y = 252 \implies y = 12$. Then $x = 42$.

Q1449. Equations: (1) $x + 7y = 98$, (2) $x + 29y = 252$. Subtracting gives $22y = 154 \implies y = 7$. Then $x = 49$.

Q1450. Equations: (1) $x + 9y = 133$, (2) $x + 23y = 287$. Subtracting gives $14y = 154 \implies y = 11$. Then $x = 34$.

Q1451. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 5$. Eq 2: $(10x + y) - (10y + x) = 9 \implies 9(x - y) = 9 \implies x - y = 1$. Solving gives $x = 3, y = 2$, so number is 32.

Q1452. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 8$. Eq 2: $(10x + y) - (10y + x) = 36 \implies 9(x - y) = 36 \implies x - y = 4$. Solving gives $x = 6, y = 2$, so number is 62.

Q1453. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x - y) = -18 \implies x - y = -2$. Solving gives $x = 5, y = 7$, so number is 57.

Q1454. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -54 \implies 9(x - y) = -54 \implies x - y = -6$. Solving gives $x = 2, y = 8$, so number is 28.

Q1455. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = 0 \implies 9(x - y) = 0 \implies x - y = 0$. Solving gives $x = 6, y = 6$, so number is 66.

Q1456. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 16$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x - y) = -18 \implies x - y = -2$. Solving gives $x = 7, y = 9$, so number is 79.

Q1457. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 8$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x - y) = -36 \implies x - y = -4$. Solving gives $x = 2, y = 6$, so number is 26.

Q1458. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 15$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 7, y = 8$, so number is 78.

Q1459. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 8$. Eq 2: $(10x + y) - (10y + x) = 18 \implies 9(x - y) = 18 \implies x - y = 2$. Solving gives $x = 5, y = 3$, so number is 53.

Q1460. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 9$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 4, y = 5$, so number is 45.

Q1461. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 36 \implies 9(x - y) = 36 \implies x - y = 4$. Solving gives $x = 7, y = 3$, so number is 73.

Q1462. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 17$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 8, y = 9$, so number is 89.

Q1463. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = 9 \implies 9(x - y) = 9 \implies x - y = 1$. Solving gives $x = 6, y = 5$, so number is 65.

Q1464. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 7$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x - y) = -27 \implies x - y = -3$. Solving gives $x = 2, y = 5$, so number is 25.

Q1465. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 6, y = 7$, so number is 67.

Q1466. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x - y) = -36 \implies x - y = -4$. Solving gives $x = 3, y = 7$, so number is 37.

Q1467. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x - y) = -18 \implies x - y = -2$. Solving gives $x = 5, y = 7$, so number is 57.

Q1468. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 7$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 5, y = 2$, so number is 52.

Q1469. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -45 \implies 9(x-y) = -45 \implies x-y = -5$. Solving gives $x = 4, y = 9$, so number is 49.

Q1470. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 7$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 5, y = 2$, so number is 52.

Q1471. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 6$. Eq 2: $(10x+y)-(10y+x) = -18 \implies 9(x-y) = -18 \implies x - y = -2$. Solving gives $x = 2, y = 4$, so number is 24.

Q1472. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = -45 \implies 9(x-y) = -45 \implies x - y = -5$. Solving gives $x = 2, y = 7$, so number is 27.

Q1473. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = -45 \implies 9(x-y) = -45 \implies x - y = -5$. Solving gives $x = 2, y = 7$, so number is 27.

Q1474. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = -27 \implies 9(x-y) = -27 \implies x - y = -3$. Solving gives $x = 3, y = 6$, so number is 36.

Q1475. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 54 \implies 9(x-y) = 54 \implies x - y = 6$. Solving gives $x = 7, y = 1$, so number is 71.

Q1476. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y) - (10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 4, y = 4$, so number is 44.

Q1477. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 4, y = 8$, so number is 48.

Q1478. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 4, y = 8$, so number is 48.

Q1479. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 6, y = 3$, so number is 63.

Q1480. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 15$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 7, y = 8$, so number is 78.

Q1481. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x-y) = -18 \implies x-y = -2$. Solving gives $x = 4, y = 6$, so number is 46.

Q1482. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = 36 \implies 9(x - y) = 36 \implies x - y = 4$. Solving gives $x = 8, y = 4$, so number is 84.

Q1483. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 7$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 5, y = 2$, so number is 52.

Q1484. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 3, y = 7$, so number is 37.

Q1485. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 54 \implies 9(x-y) = 54 \implies x - y = 6$. Solving gives $x = 7, y = 1$, so number is 71.

Q1486. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -54 \implies 9(x-y) = -54 \implies x-y = -6$. Solving gives $x = 3, y = 9$, so number is 39.

Q1487. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 5, y = 4$, so number is 54.

Q1488. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 8, y = 5$, so number is 85.

Q1489. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 6$. Eq 2: $(10x+y)-(10y+x) = 36 \implies 9(x-y) = 36 \implies x - y = 4$. Solving gives $x = 5, y = 1$, so number is 51.

Q1490. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 15$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 8, y = 7$, so number is 87.

Q1491. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x-y) = -18 \implies x-y = -2$. Solving gives $x = 5, y = 7$, so number is 57.

Q1492. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 13$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 7, y = 6$, so number is 76.

Q1493. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 10$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 5, y = 5$, so number is 55.

Q1494. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 54 \implies 9(x - y) = 54 \implies x - y = 6$. Solving gives $x = 8, y = 2$, so number is 82.

Q1495. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 5$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 4, y = 1$, so number is 41.

Q1496. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 4, y = 7$, so number is 47.

Q1497. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 36 \implies 9(x-y) = 36 \implies x - y = 4$. Solving gives $x = 6, y = 2$, so number is 62.

Q1498. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 18 \implies$

$9(x-y)=18 \implies x-y=2$. Solving gives $x=6, y=4$, so number is 64.

Q1499. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=10$. Eq 2: $(10x+y)-(10y+x)=18 \implies 9(x-y)=18 \implies x-y=2$. Solving gives $x=6, y=4$, so number is 64.

Q1500. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=16$. Eq 2: $(10x+y)-(10y+x)=0 \implies 9(x-y)=0 \implies x-y=0$. Solving gives $x=8, y=8$, so number is 88.

Q1501. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=16$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=7, y=9$, so number is 79.

Q1502. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=10$. Eq 2: $(10x+y)-(10y+x)=-54 \implies 9(x-y)=-54 \implies x-y=-6$. Solving gives $x=2, y=8$, so number is 28.

Q1503. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=11$. Eq 2: $(10x+y)-(10y+x)=-45 \implies 9(x-y)=-45 \implies x-y=-5$. Solving gives $x=3, y=8$, so number is 38.

Q1504. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=15$. Eq 2: $(10x+y)-(10y+x)=9 \implies 9(x-y)=9 \implies x-y=1$. Solving gives $x=8, y=7$, so number is 87.

Q1505. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=13$. Eq 2: $(10x+y)-(10y+x)=9 \implies 9(x-y)=9 \implies x-y=1$. Solving gives $x=7, y=6$, so number is 76.

Q1506. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=12$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=5, y=7$, so number is 57.

Q1507. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=13$. Eq 2: $(10x+y)-(10y+x)=9 \implies 9(x-y)=9 \implies x-y=1$. Solving gives $x=7, y=6$, so number is 76.

Q1508. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=15$. Eq 2: $(10x+y)-(10y+x)=-27 \implies 9(x-y)=-27 \implies x-y=-3$. Solving gives $x=6, y=9$, so number is 69.

Q1509. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=11$. Eq 2: $(10x+y)-(10y+x)=-45 \implies 9(x-y)=-45 \implies x-y=-5$. Solving gives $x=3, y=8$, so number is 38.

Q1510. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=14$. Eq 2: $(10x+y)-(10y+x)=-36 \implies 9(x-y)=-36 \implies x-y=-4$. Solving gives $x=5, y=9$, so number is 59.

Q1511. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=16$. Eq 2: $(10x+y)-(10y+x)=0 \implies 9(x-y)=0 \implies x-y=0$. Solving gives $x=8, y=8$, so number is 88.

Q1512. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=12$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=5, y=7$, so number is 57.

Q1513. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=9$. Eq 2: $(10x+y)-(10y+x)=-9 \implies 9(x-y)=-9 \implies x-y=-1$. Solving gives $x=4, y=5$, so number is 45.

Q1514. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=6$. Eq 2: $(10x+y)-(10y+x)=36 \implies 9(x-y)=36 \implies x-y=4$. Solving gives $x=5, y=1$, so number is 51.

Q1515. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=7$. Eq 2: $(10x+y)-(10y+x)=9 \implies 9(x-y)=9 \implies x-y=1$. Solving gives $x=4, y=3$, so number is 43.

Q1516. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=13$. Eq 2: $(10x+y)-(10y+x)=-45 \implies 9(x-y)=-45 \implies x-y=-5$. Solving gives $x=4, y=9$, so number is 49.

Q1517. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=11$. Eq 2: $(10x+y)-(10y+x)=-45 \implies 9(x-y)=-45 \implies x-y=-5$. Solving gives $x=3, y=8$, so number is 38.

Q1518. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=4$. Eq 2: $(10x+y)-(10y+x)=18 \implies 9(x-y)=18 \implies x-y=2$. Solving gives $x=3, y=1$, so number is 31.

Q1519. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=16$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=7, y=9$, so number is 79.

Q1520. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=9$. Eq 2: $(10x+y)-(10y+x)=-45 \implies 9(x-y)=-45 \implies x-y=-5$. Solving gives $x=2, y=7$, so number is 27.

Q1521. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=12$. Eq 2: $(10x+y)-(10y+x)=-36 \implies 9(x-y)=-36 \implies x-y=-4$. Solving gives $x=4, y=8$, so number is 48.

Q1522. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=14$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=6, y=8$, so number is 68.

Q1523. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=6$. Eq 2: $(10x+y)-(10y+x)=-18 \implies 9(x-y)=-18 \implies x-y=-2$. Solving gives $x=2, y=4$, so number is 24.

Q1524. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=11$. Eq 2: $(10x+y)-(10y+x)=45 \implies 9(x-y)=45 \implies x-y=5$. Solving gives $x=8, y=3$, so number is 83.

Q1525. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=9$. Eq 2: $(10x+y)-(10y+x)=-27 \implies 9(x-y)=-27 \implies x-y=-3$. Solving gives $x=3, y=6$, so number is 36.

Q1526. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=9$. Eq 2: $(10x+y)-(10y+x)=45 \implies 9(x-y)=45 \implies x-y=5$. Solving gives $x=7, y=2$, so number is 72.

Q1527. Original $=10x+y$, Reversed $=10y+x$. Eq 1: $x+y=8$. Eq 2: $(10x+y)-(10y+x)=-36 \implies 9(x-y)=-36 \implies x-y=-4$. Solving gives $x=2, y=6$, so number is 26.

Q1528. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -45 \implies 9(x-y) = -45 \implies x-y = -5$. Solving gives $x = 4, y = 9$, so number is 49.

Q1529. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = 18 \implies 9(x - y) = 18 \implies x - y = 2$. Solving gives $x = 7, y = 5$, so number is 75.

Q1530. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 5, y = 8$, so number is 58.

Q1531. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -45 \implies 9(x-y) = -45 \implies x-y = -5$. Solving gives $x = 4, y = 9$, so number is 49.

Q1532. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -45 \implies 9(x-y) = -45 \implies x-y = -5$. Solving gives $x = 3, y = 8$, so number is 38.

Q1533. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 13$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 7, y = 6$, so number is 76.

Q1534. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 12$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 6, y = 6$, so number is 66.

Q1535. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 5$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 4, y = 1$, so number is 41.

Q1536. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = 45 \implies 9(x - y) = 45 \implies x - y = 5$. Solving gives $x = 8, y = 3$, so number is 83.

Q1537. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 54 \implies 9(x - y) = 54 \implies x - y = 6$. Solving gives $x = 8, y = 2$, so number is 82.

Q1538. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 4, y = 7$, so number is 47.

Q1539. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 4, y = 4$, so number is 44.

Q1540. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 17$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 8, y = 9$, so number is 89.

Q1541. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -54 \implies 9(x-y) = -54 \implies x-y = -6$. Solving gives $x = 3, y = 9$, so number is 39.

Q1542. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 6$. Eq 2: $(10x+y)-(10y+x) = 18 \implies 9(x-y) = 18 \implies x - y = 2$. Solving gives $x = 4, y = 2$, so number is 42.

Q1543. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -45 \implies 9(x-y) = -45 \implies x-y = -5$. Solving gives $x = 4, y = 9$, so number is 49.

Q1544. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 54 \implies 9(x - y) = 54 \implies x - y = 6$. Solving gives $x = 8, y = 2$, so number is 82.

Q1545. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 8, y = 5$, so number is 85.

Q1546. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 5$. Eq 2: $(10x+y)-(10y+x) = 27 \implies 9(x-y) = 27 \implies x - y = 3$. Solving gives $x = 4, y = 1$, so number is 41.

Q1547. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 5, y = 6$, so number is 56.

Q1548. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -63 \implies 9(x-y) = -63 \implies x-y = -7$. Solving gives $x = 2, y = 9$, so number is 29.

Q1549. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 4, y = 7$, so number is 47.

Q1550. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 14$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 5, y = 9$, so number is 59.

Q1551. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 14$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 7, y = 7$, so number is 77.

Q1552. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 3, y = 7$, so number is 37.

Q1553. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 11$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 6, y = 5$, so number is 65.

Q1554. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 10$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 5, y = 5$, so number is 55.

Q1555. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = -36 \implies 9(x-y) = -36 \implies x - y = -4$. Solving gives $x = 2, y = 6$, so number is 26.

Q1556. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 8, y = 5$, so number is 85.

Q1557. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -63 \implies$

$9(x-y) = -63 \implies x-y = -7$. Solving gives $x = 2, y = 9$, so number is 29.

Q1558. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 54 \implies 9(x-y) = 54 \implies x - y = 6$. Solving gives $x = 7, y = 1$, so number is 71.

Q1559. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 4, y = 5$, so number is 45.

Q1560. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 6$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 3, y = 3$, so number is 33.

Q1561. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 5$. Eq 2: $(10x+y)-(10y+x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 2, y = 3$, so number is 23.

Q1562. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 8$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 4, y = 4$, so number is 44.

Q1563. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 14$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 5, y = 9$, so number is 59.

Q1564. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 10$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 5, y = 5$, so number is 55.

Q1565. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = 54 \implies 9(x - y) = 54 \implies x - y = 6$. Solving gives $x = 8, y = 2$, so number is 82.

Q1566. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 5, y = 8$, so number is 58.

Q1567. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 14$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 7, y = 7$, so number is 77.

Q1568. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 8, y = 5$, so number is 85.

Q1569. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 6$. Eq 2: $(10x+y)-(10y+x) = 18 \implies 9(x-y) = 18 \implies x - y = 2$. Solving gives $x = 4, y = 2$, so number is 42.

Q1570. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 16$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 8, y = 8$, so number is 88.

Q1571. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 14$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x-y) = -18 \implies x-y = -2$. Solving gives $x = 6, y = 8$, so number is 68.

Q1572. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 7, y = 4$, so number is 74.

Q1573. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 16$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x-y) = -18 \implies x-y = -2$. Solving gives $x = 7, y = 9$, so number is 79.

Q1574. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x-y) = -27 \implies x-y = -3$. Solving gives $x = 5, y = 8$, so number is 58.

Q1575. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 7$. Eq 2: $(10x+y)-(10y+x) = -27 \implies 9(x-y) = -27 \implies x - y = -3$. Solving gives $x = 2, y = 5$, so number is 25.

Q1576. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 11$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 6, y = 5$, so number is 65.

Q1577. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 13$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 7, y = 6$, so number is 76.

Q1578. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 12$. Eq 2: $(10x + y) - (10y + x) = -36 \implies 9(x-y) = -36 \implies x-y = -4$. Solving gives $x = 4, y = 8$, so number is 48.

Q1579. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 14$. Eq 2: $(10x + y) - (10y + x) = 18 \implies 9(x - y) = 18 \implies x - y = 2$. Solving gives $x = 8, y = 6$, so number is 86.

Q1580. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 17$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x-y) = -9 \implies x-y = -1$. Solving gives $x = 8, y = 9$, so number is 89.

Q1581. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = 45 \implies 9(x - y) = 45 \implies x - y = 5$. Solving gives $x = 8, y = 3$, so number is 83.

Q1582. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 9$. Eq 2: $(10x+y)-(10y+x) = 63 \implies 9(x-y) = 63 \implies x - y = 7$. Solving gives $x = 8, y = 1$, so number is 81.

Q1583. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 12$. Eq 2: $(10x+y)-(10y+x) = 0 \implies 9(x-y) = 0 \implies x - y = 0$. Solving gives $x = 6, y = 6$, so number is 66.

Q1584. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 10$. Eq 2: $(10x + y) - (10y + x) = -54 \implies 9(x-y) = -54 \implies x-y = -6$. Solving gives $x = 2, y = 8$, so number is 28.

Q1585. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 11$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 6, y = 5$, so number is 65.

Q1586. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x+y = 7$. Eq 2: $(10x+y)-(10y+x) = 9 \implies 9(x-y) = 9 \implies x - y = 1$. Solving gives $x = 4, y = 3$, so number is 43.

Q1587. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 9$. Eq 2: $(10x + y) - (10y + x) = 9 \implies 9(x - y) = 9 \implies x - y = 1$. Solving gives $x = 5, y = 4$, so number is 54.

Q1588. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 6$. Eq 2: $(10x + y) - (10y + x) = 18 \implies 9(x - y) = 18 \implies x - y = 2$. Solving gives $x = 4, y = 2$, so number is 42.

Q1589. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 13$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 8, y = 5$, so number is 85.

Q1590. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 7$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 3, y = 4$, so number is 34.

Q1591. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 16$. Eq 2: $(10x + y) - (10y + x) = -18 \implies 9(x - y) = -18 \implies x - y = -2$. Solving gives $x = 7, y = 9$, so number is 79.

Q1592. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -63 \implies 9(x - y) = -63 \implies x - y = -7$. Solving gives $x = 2, y = 9$, so number is 29.

Q1593. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 4$. Eq 2: $(10x + y) - (10y + x) = 18 \implies 9(x - y) = 18 \implies x - y = 2$. Solving gives $x = 3, y = 1$, so number is 31.

Q1594. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x - y) = -27 \implies x - y = -3$. Solving gives $x = 4, y = 7$, so number is 47.

Q1595. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 9$. Eq 2: $(10x + y) - (10y + x) = 63 \implies 9(x - y) = 63 \implies x - y = 7$. Solving gives $x = 8, y = 1$, so number is 81.

Q1596. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 15$. Eq 2: $(10x + y) - (10y + x) = 9 \implies 9(x - y) = 9 \implies x - y = 1$. Solving gives $x = 8, y = 7$, so number is 87.

Q1597. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 5$. Eq 2: $(10x + y) - (10y + x) = -9 \implies 9(x - y) = -9 \implies x - y = -1$. Solving gives $x = 2, y = 3$, so number is 23.

Q1598. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 7$. Eq 2: $(10x + y) - (10y + x) = -27 \implies 9(x - y) = -27 \implies x - y = -3$. Solving gives $x = 2, y = 5$, so number is 25.

Q1599. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 11$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 7, y = 4$, so number is 74.

Q1600. Original $= 10x + y$, Reversed $= 10y + x$. Eq 1: $x + y = 7$. Eq 2: $(10x + y) - (10y + x) = 27 \implies 9(x - y) = 27 \implies x - y = 3$. Solving gives $x = 5, y = 2$, so number is 52.

Q1601. Cross-multiplying $\frac{x+5}{y-3} = \frac{1}{3} \implies 3(x + 5) = 1(y - 3) \implies 3x - 1y = -18$.

Q1602. Cross-multiplying $\frac{x+1}{y-3} = \frac{1}{2} \implies 2(x + 1) = 1(y - 3) \implies 2x - 1y = -5$.

Q1603. Cross-multiplying $\frac{x+2}{y-3} = \frac{3}{5} \implies 5(x + 2) = 3(y - 3) \implies 5x - 3y = -19$.

Q1604. Cross-multiplying $\frac{x+2}{y-5} = \frac{1}{2} \implies 2(x + 2) = 1(y - 5) \implies 2x - 1y = -9$.

Q1605. Cross-multiplying $\frac{x+4}{y-3} = \frac{3}{3} \implies 3(x + 4) = 3(y - 3) \implies 3x - 3y = -21$.

Q1606. Cross-multiplying $\frac{x+5}{y-5} = \frac{1}{4} \implies 4(x + 5) = 1(y - 5) \implies 4x - 1y = -25$.

Q1607. Cross-multiplying $\frac{x+3}{y-3} = \frac{1}{3} \implies 3(x + 3) = 1(y - 3) \implies 3x - 1y = -12$.

Q1608. Cross-multiplying $\frac{x+4}{y-1} = \frac{2}{4} \implies 4(x + 4) = 2(y - 1) \implies 4x - 2y = -18$.

Q1609. Cross-multiplying $\frac{x+4}{y-3} = \frac{1}{5} \implies 5(x + 4) = 1(y - 3) \implies 5x - 1y = -23$.

Q1610. Cross-multiplying $\frac{x+3}{y-2} = \frac{1}{5} \implies 5(x + 3) = 1(y - 2) \implies 5x - 1y = -17$.

Q1611. Cross-multiplying $\frac{x+5}{y-2} = \frac{1}{3} \implies 3(x + 5) = 1(y - 2) \implies 3x - 1y = -17$.

Q1612. Cross-multiplying $\frac{x+4}{y-4} = \frac{1}{5} \implies 5(x + 4) = 1(y - 4) \implies 5x - 1y = -24$.

Q1613. Cross-multiplying $\frac{x+3}{y-4} = \frac{2}{2} \implies 2(x + 3) = 2(y - 4) \implies 2x - 2y = -14$.

Q1614. Cross-multiplying $\frac{x+4}{y-5} = \frac{2}{4} \implies 4(x + 4) = 2(y - 5) \implies 4x - 2y = -26$.

Q1615. Cross-multiplying $\frac{x+3}{y-5} = \frac{3}{3} \implies 3(x + 3) = 3(y - 5) \implies 3x - 3y = -24$.

Q1616. Cross-multiplying $\frac{x+3}{y-2} = \frac{1}{3} \implies 3(x + 3) = 1(y - 2) \implies 3x - 1y = -11$.

Q1617. Cross-multiplying $\frac{x+5}{y-5} = \frac{2}{3} \implies 3(x + 5) = 2(y - 5) \implies 3x - 2y = -25$.

Q1618. Cross-multiplying $\frac{x+5}{y-4} = \frac{2}{2} \implies 2(x + 5) = 2(y - 4) \implies 2x - 2y = -18$.

Q1619. Cross-multiplying $\frac{x+1}{y-2} = \frac{2}{2} \implies 2(x + 1) = 2(y - 2) \implies 2x - 2y = -6$.

Q1620. Cross-multiplying $\frac{x+3}{y-4} = \frac{3}{2} \implies 2(x + 3) = 3(y - 4) \implies 2x - 3y = -18$.

Q1621. Cross-multiplying $\frac{x+3}{y-3} = \frac{1}{2} \implies 2(x + 3) = 1(y - 3) \implies 2x - 1y = -9$.

Q1622. Cross-multiplying $\frac{x+5}{y-1} = \frac{2}{3} \implies 3(x + 5) = 2(y - 1) \implies 3x - 2y = -17$.

Q1623. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{2} \implies 2(x + 3) = 3(y - 1) \implies 2x - 3y = -9$.

Q1624. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{5} \implies 5(x + 3) = 3(y - 1) \implies 5x - 3y = -18$.

Q1625. Cross-multiplying $\frac{x+5}{y-2} = \frac{1}{4} \implies 4(x + 5) = 1(y - 2) \implies 4x - 1y = -22$.

Q1626. Cross-multiplying $\frac{x+1}{y-3} = \frac{3}{2} \implies 2(x + 1) = 3(y - 3) \implies 2x - 3y = -11$.

Q1627. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{5} \implies 5(x + 3) = 3(y - 1) \implies 5x - 3y = -18$.

Q1628. Cross-multiplying $\frac{x+1}{y-1} = \frac{2}{4} \implies 4(x + 1) = 2(y - 1) \implies 4x - 2y = -6$.

Q1629. Cross-multiplying $\frac{x+1}{y-3} = \frac{1}{4} \implies 4(x+1) = 1(y-3) \implies 4x - 1y = -7$.

Q1630. Cross-multiplying $\frac{x+2}{y-3} = \frac{2}{5} \implies 5(x+2) = 2(y-3) \implies 5x - 2y = -16$.

Q1631. Cross-multiplying $\frac{x+4}{y-3} = \frac{3}{3} \implies 3(x+4) = 3(y-3) \implies 3x - 3y = -21$.

Q1632. Cross-multiplying $\frac{x+5}{y-3} = \frac{3}{4} \implies 4(x+5) = 3(y-3) \implies 4x - 3y = -29$.

Q1633. Cross-multiplying $\frac{x+5}{y-4} = \frac{1}{2} \implies 2(x+5) = 1(y-4) \implies 2x - 1y = -14$.

Q1634. Cross-multiplying $\frac{x+5}{y-3} = \frac{1}{4} \implies 4(x+5) = 1(y-3) \implies 4x - 1y = -23$.

Q1635. Cross-multiplying $\frac{x+1}{y-4} = \frac{2}{2} \implies 2(x+1) = 2(y-4) \implies 2x - 2y = -10$.

Q1636. Cross-multiplying $\frac{x+2}{y-5} = \frac{3}{5} \implies 5(x+2) = 3(y-5) \implies 5x - 3y = -25$.

Q1637. Cross-multiplying $\frac{x+5}{y-3} = \frac{1}{4} \implies 4(x+5) = 1(y-3) \implies 4x - 1y = -23$.

Q1638. Cross-multiplying $\frac{x+5}{y-1} = \frac{2}{4} \implies 4(x+5) = 2(y-1) \implies 4x - 2y = -22$.

Q1639. Cross-multiplying $\frac{x+2}{y-1} = \frac{2}{5} \implies 5(x+2) = 2(y-1) \implies 5x - 2y = -12$.

Q1640. Cross-multiplying $\frac{x+5}{y-2} = \frac{2}{5} \implies 5(x+5) = 2(y-2) \implies 5x - 2y = -29$.

Q1641. Cross-multiplying $\frac{x+2}{y-4} = \frac{1}{5} \implies 5(x+2) = 1(y-4) \implies 5x - 1y = -14$.

Q1642. Cross-multiplying $\frac{x+2}{y-1} = \frac{3}{4} \implies 4(x+2) = 3(y-1) \implies 4x - 3y = -11$.

Q1643. Cross-multiplying $\frac{x+2}{y-5} = \frac{1}{5} \implies 5(x+2) = 1(y-5) \implies 5x - 1y = -15$.

Q1644. Cross-multiplying $\frac{x+2}{y-5} = \frac{1}{3} \implies 3(x+2) = 1(y-5) \implies 3x - 1y = -11$.

Q1645. Cross-multiplying $\frac{x+5}{y-2} = \frac{2}{5} \implies 5(x+5) = 2(y-2) \implies 5x - 2y = -29$.

Q1646. Cross-multiplying $\frac{x+4}{y-4} = \frac{1}{5} \implies 5(x+4) = 1(y-4) \implies 5x - 1y = -24$.

Q1647. Cross-multiplying $\frac{x+1}{y-3} = \frac{1}{4} \implies 4(x+1) = 1(y-3) \implies 4x - 1y = -7$.

Q1648. Cross-multiplying $\frac{x+5}{y-1} = \frac{2}{3} \implies 3(x+5) = 2(y-1) \implies 3x - 2y = -17$.

Q1649. Cross-multiplying $\frac{x+5}{y-4} = \frac{3}{5} \implies 5(x+5) = 3(y-4) \implies 5x - 3y = -37$.

Q1650. Cross-multiplying $\frac{x+4}{y-4} = \frac{2}{2} \implies 2(x+4) = 3(y-4) \implies 2x - 3y = -20$.

Q1651. Cross-multiplying $\frac{x+3}{y-3} = \frac{2}{3} \implies 3(x+3) = 2(y-3) \implies 3x - 2y = -15$.

Q1652. Cross-multiplying $\frac{x+2}{y-5} = \frac{3}{5} \implies 5(x+2) = 3(y-5) \implies 5x - 3y = -25$.

Q1653. Cross-multiplying $\frac{x+5}{y-4} = \frac{1}{2} \implies 2(x+5) = 1(y-4) \implies 2x - 1y = -14$.

Q1654. Cross-multiplying $\frac{x+1}{y-1} = \frac{3}{3} \implies 3(x+1) = 3(y-1) \implies 3x - 3y = -6$.

Q1655. Cross-multiplying $\frac{x+5}{y-4} = \frac{1}{2} \implies 2(x+5) = 1(y-4) \implies 2x - 1y = -14$.

Q1656. Cross-multiplying $\frac{x+2}{y-2} = \frac{2}{2} \implies 2(x+2) = 2(y-2) \implies 2x - 2y = -8$.

Q1657. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{4} \implies 4(x+3) = 3(y-1) \implies 4x - 3y = -15$.

Q1658. Cross-multiplying $\frac{x+4}{y-1} = \frac{3}{3} \implies 3(x+4) = 3(y-1) \implies 3x - 3y = -15$.

Q1659. Cross-multiplying $\frac{x+1}{y-3} = \frac{3}{4} \implies 4(x+1) = 3(y-3) \implies 4x - 3y = -13$.

Q1660. Cross-multiplying $\frac{x+4}{y-4} = \frac{2}{5} \implies 5(x+4) = 2(y-4) \implies 5x - 2y = -28$.

Q1661. Cross-multiplying $\frac{x+4}{y-5} = \frac{3}{5} \implies 5(x+4) = 3(y-5) \implies 5x - 3y = -35$.

Q1662. Cross-multiplying $\frac{x+2}{y-4} = \frac{2}{4} \implies 4(x+2) = 2(y-4) \implies 4x - 2y = -16$.

Q1663. Cross-multiplying $\frac{x+2}{y-1} = \frac{3}{5} \implies 5(x+2) = 3(y-1) \implies 5x - 3y = -13$.

Q1664. Cross-multiplying $\frac{x+3}{y-2} = \frac{3}{5} \implies 5(x+3) = 3(y-2) \implies 5x - 3y = -21$.

Q1665. Cross-multiplying $\frac{x+2}{y-3} = \frac{1}{3} \implies 3(x+2) = 1(y-3) \implies 3x - 1y = -9$.

Q1666. Cross-multiplying $\frac{x+4}{y-5} = \frac{1}{3} \implies 3(x+4) = 1(y-5) \implies 3x - 1y = -17$.

Q1667. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{4} \implies 4(x+3) = 3(y-1) \implies 4x - 3y = -15$.

Q1668. Cross-multiplying $\frac{x+4}{y-2} = \frac{2}{5} \implies 5(x+4) = 2(y-2) \implies 5x - 2y = -24$.

Q1669. Cross-multiplying $\frac{x+2}{y-1} = \frac{1}{2} \implies 2(x+2) = 1(y-1) \implies 2x - 1y = -5$.

Q1670. Cross-multiplying $\frac{x+2}{y-5} = \frac{2}{5} \implies 5(x+2) = 2(y-5) \implies 5x - 2y = -20$.

Q1671. Cross-multiplying $\frac{x+5}{y-5} = \frac{2}{3} \implies 3(x+5) = 2(y-5) \implies 3x - 2y = -25$.

Q1672. Cross-multiplying $\frac{x+1}{y-5} = \frac{2}{4} \implies 4(x+1) = 2(y-5) \implies 4x - 2y = -14$.

Q1673. Cross-multiplying $\frac{x+4}{y-4} = \frac{1}{2} \implies 2(x+4) = 1(y-4) \implies 2x - 1y = -12$.

Q1674. Cross-multiplying $\frac{x+1}{y-1} = \frac{2}{5} \implies 5(x+1) = 2(y-1) \implies 5x - 2y = -7$.

Q1675. Cross-multiplying $\frac{x+5}{y-4} = \frac{3}{3} \implies 3(x+5) = 3(y-4) \implies 3x - 3y = -27$.

Q1676. Cross-multiplying $\frac{x+3}{y-4} = \frac{3}{4} \implies 4(x+3) = 3(y-4) \implies 4x - 3y = -24$.

Q1677. Cross-multiplying $\frac{x+2}{y-2} = \frac{2}{3} \implies 3(x+2) = 2(y-2) \implies 3x - 2y = -10$.

Q1678. Cross-multiplying $\frac{x+5}{y-1} = \frac{2}{5} \implies 5(x+5) = 2(y-1) \implies 5x - 2y = -27$.

Q1679. Cross-multiplying $\frac{x+2}{y-2} = \frac{1}{4} \implies 4(x+2) = 1(y-2) \implies 4x - 1y = -10$.

Q1680. Cross-multiplying $\frac{x+4}{y-1} = \frac{2}{4} \implies 4(x+4) = 2(y-1) \implies 4x - 2y = -18$.

Q1681. Cross-multiplying $\frac{x+3}{y-3} = \frac{1}{4} \implies 4(x+3) = 1(y-3) \implies 4x - 1y = -15$.

Q1682. Cross-multiplying $\frac{x+2}{y-4} = \frac{2}{5} \implies 5(x+2) = 2(y-4) \implies 5x - 2y = -18$.

Q1683. Cross-multiplying $\frac{x+2}{y-1} = \frac{1}{2} \implies 2(x+2) = 1(y-1) \implies 2x - 1y = -5$.

Q1684. Cross-multiplying $\frac{x+5}{y-3} = \frac{2}{2} \implies 2(x+5) = 2(y-3) \implies 2x - 2y = -16$.

Q1685. Cross-multiplying $\frac{x+5}{y-3} = \frac{2}{2} \implies 2(x+5) = 2(y-3) \implies 2x - 2y = -16$.

Q1686. Cross-multiplying $\frac{x+5}{y-4} = \frac{1}{2} \implies 2(x+5) = 1(y-4) \implies 2x - 1y = -14$.

Q1687. Cross-multiplying $\frac{x+5}{y-4} = \frac{3}{4} \implies 4(x+5) = 3(y-4) \implies 4x - 3y = -32$.

Q1688. Cross-multiplying $\frac{x+2}{y-2} = \frac{2}{5} \implies 5(x+2) = 2(y-2) \implies 5x - 2y = -14$.

Q1689. Cross-multiplying $\frac{x+4}{y-2} = \frac{2}{3} \implies 3(x+4) = 2(y-2) \implies 3x - 2y = -16$.

Q1690. Cross-multiplying $\frac{x+3}{y-2} = \frac{3}{4} \implies 4(x+3) = 3(y-2) \implies 4x - 3y = -18$.

Q1691. Cross-multiplying $\frac{x+3}{y-3} = \frac{1}{5} \implies 5(x+3) = 1(y-3) \implies 5x - 1y = -18$.

Q1692. Cross-multiplying $\frac{x+5}{y-3} = \frac{2}{2} \implies 2(x+5) = 2(y-3) \implies 2x - 2y = -16$.

Q1693. Cross-multiplying $\frac{x+5}{y-2} = \frac{3}{3} \implies 3(x+5) = 3(y-2) \implies 3x - 3y = -21$.

Q1694. Cross-multiplying $\frac{x+5}{y-1} = \frac{1}{2} \implies 2(x+5) = 1(y-1) \implies 2x - 1y = -11$.

Q1695. Cross-multiplying $\frac{x+2}{y-2} = \frac{3}{4} \implies 4(x+2) = 3(y-2) \implies 4x - 3y = -14$.

Q1696. Cross-multiplying $\frac{x+5}{y-5} = \frac{1}{5} \implies 5(x+5) = 1(y-5) \implies 5x - 1y = -30$.

Q1697. Cross-multiplying $\frac{x+2}{y-4} = \frac{3}{5} \implies 5(x+2) = 3(y-4) \implies 5x - 3y = -22$.

Q1698. Cross-multiplying $\frac{x+1}{y-1} = \frac{2}{5} \implies 5(x+1) = 2(y-1) \implies 5x - 2y = -7$.

Q1699. Cross-multiplying $\frac{x+5}{y-5} = \frac{1}{5} \implies 5(x+5) = 1(y-5) \implies 5x - 1y = -30$.

Q1700. Cross-multiplying $\frac{x+1}{y-1} = \frac{1}{3} \implies 3(x+1) = 1(y-1) \implies 3x - 1y = -4$.

Q1701. Cross-multiplying $\frac{x+1}{y-4} = \frac{1}{4} \implies 4(x+1) = 1(y-4) \implies 4x - 1y = -8$.

Q1702. Cross-multiplying $\frac{x+3}{y-4} = \frac{2}{2} \implies 2(x+3) = 2(y-4) \implies 2x - 2y = -14$.

Q1703. Cross-multiplying $\frac{x+4}{y-1} = \frac{3}{3} \implies 3(x+4) = 3(y-1) \implies 3x - 3y = -15$.

Q1704. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{4} \implies 4(x+3) = 3(y-1) \implies 4x - 3y = -15$.

Q1705. Cross-multiplying $\frac{x+4}{y-1} = \frac{3}{2} \implies 2(x+4) = 3(y-1) \implies 2x - 3y = -11$.

Q1706. Cross-multiplying $\frac{x+1}{y-2} = \frac{3}{3} \implies 3(x+1) = 3(y-2) \implies 3x - 3y = -9$.

Q1707. Cross-multiplying $\frac{x+2}{y-2} = \frac{3}{2} \implies 2(x+2) = 3(y-2) \implies 2x - 3y = -10$.

Q1708. Cross-multiplying $\frac{x+1}{y-1} = \frac{3}{5} \implies 5(x+1) = 3(y-1) \implies 5x - 3y = -8$.

Q1709. Cross-multiplying $\frac{x+2}{y-3} = \frac{3}{2} \implies 2(x+2) = 3(y-3) \implies 2x - 3y = -13$.

Q1710. Cross-multiplying $\frac{x+3}{y-4} = \frac{2}{3} \implies 3(x+3) = 2(y-4) \implies 3x - 2y = -17$.

Q1711. Cross-multiplying $\frac{x+1}{y-3} = \frac{2}{2} \implies 2(x+1) = 2(y-3) \implies 2x - 2y = -8$.

Q1712. Cross-multiplying $\frac{x+4}{y-3} = \frac{1}{2} \implies 2(x+4) = 1(y-3) \implies 2x - 1y = -11$.

Q1713. Cross-multiplying $\frac{x+1}{y-5} = \frac{2}{3} \implies 3(x+1) = 2(y-5) \implies 3x - 2y = -13$.

Q1714. Cross-multiplying $\frac{x+1}{y-1} = \frac{1}{2} \implies 2(x+1) = 1(y-1) \implies 2x - 1y = -3$.

Q1715. Cross-multiplying $\frac{x+3}{y-4} = \frac{2}{2} \implies 2(x+3) = 2(y-4) \implies 2x - 2y = -14$.

Q1716. Cross-multiplying $\frac{x+4}{y-2} = \frac{3}{5} \implies 5(x+4) = 3(y-2) \implies 5x - 3y = -26$.

Q1717. Cross-multiplying $\frac{x+2}{y-4} = \frac{2}{5} \implies 5(x+2) = 2(y-4) \implies 5x - 2y = -18$.

Q1718. Cross-multiplying $\frac{x+4}{y-4} = \frac{3}{2} \implies 2(x+4) = 3(y-4) \implies 2x - 3y = -20$.

Q1719. Cross-multiplying $\frac{x+4}{y-2} = \frac{1}{2} \implies 2(x+4) = 1(y-2) \implies 2x - 1y = -10$.

Q1720. Cross-multiplying $\frac{x+1}{y-1} = \frac{3}{5} \implies 5(x+1) = 3(y-1) \implies 5x - 3y = -8$.

Q1721. Cross-multiplying $\frac{x+3}{y-5} = \frac{2}{3} \implies 3(x+3) = 2(y-5) \implies 3x - 2y = -19$.

Q1722. Cross-multiplying $\frac{x+3}{y-3} = \frac{3}{2} \implies 2(x+3) = 3(y-3) \implies 2x - 3y = -15$.

Q1723. Cross-multiplying $\frac{x+3}{y-2} = \frac{3}{3} \implies 3(x+3) = 3(y-2) \implies 3x - 3y = -15$.

Q1724. Cross-multiplying $\frac{x+5}{y-1} = \frac{3}{3} \implies 3(x+5) = 3(y-1) \implies 3x - 3y = -18$.

Q1725. Cross-multiplying $\frac{x+5}{y-3} = \frac{1}{2} \implies 2(x+5) = 1(y-3) \implies 2x - 1y = -13$.

Q1726. Cross-multiplying $\frac{x+2}{y-5} = \frac{1}{3} \implies 3(x+2) = 1(y-5) \implies 3x - 1y = -11$.

Q1727. Cross-multiplying $\frac{x+1}{y-3} = \frac{2}{5} \implies 5(x+1) = 2(y-3) \implies 5x - 2y = -11$.

Q1728. Cross-multiplying $\frac{x+2}{y-1} = \frac{3}{5} \implies 5(x+2) = 3(y-1) \implies 5x - 3y = -13$.

Q1729. Cross-multiplying $\frac{x+1}{y-4} = \frac{1}{3} \implies 3(x+1) = 1(y-4) \implies 3x - 1y = -7$.

Q1730. Cross-multiplying $\frac{x+5}{y-3} = \frac{2}{4} \implies 4(x+5) = 2(y-3) \implies 4x - 2y = -26$.

Q1731. Cross-multiplying $\frac{x+4}{y-5} = \frac{3}{2} \implies 2(x+4) = 3(y-5) \implies 2x - 3y = -23$.

Q1732. Cross-multiplying $\frac{x+2}{y-2} = \frac{1}{5} \implies 5(x+2) = 1(y-2) \implies 5x - 1y = -12$.

Q1733. Cross-multiplying $\frac{x+4}{y-1} = \frac{1}{3} \implies 3(x+4) = 1(y-1) \implies 3x - 1y = -13$.

Q1734. Cross-multiplying $\frac{x+5}{y-3} = \frac{2}{4} \implies 4(x+5) = 2(y-3) \implies 4x - 2y = -26$.

Q1735. Cross-multiplying $\frac{x+3}{y-5} = \frac{2}{5} \implies 5(x+3) = 2(y-5) \implies 5x - 2y = -25$.

Q1736. Cross-multiplying $\frac{x+2}{y-5} = \frac{2}{4} \implies 4(x+2) = 2(y-5) \implies 4x - 2y = -18$.

Q1737. Cross-multiplying $\frac{x+3}{y-3} = \frac{3}{2} \implies 2(x+3) = 3(y-3) \implies 2x - 3y = -15$.

Q1738. Cross-multiplying $\frac{x+4}{y-2} = \frac{1}{5} \implies 5(x+4) = 1(y-2) \implies 5x - 1y = -22$.

Q1739. Cross-multiplying $\frac{x+4}{y-4} = \frac{3}{5} \implies 5(x+4) = 3(y-4) \implies 5x - 3y = -32$.

Q1740. Cross-multiplying $\frac{x+3}{y-1} = \frac{3}{3} \implies 3(x+3) = 3(y-1) \implies 3x - 3y = -12$.

Q1741. Cross-multiplying $\frac{x+4}{y-4} = \frac{1}{2} \implies 2(x+4) = 1(y-4) \implies 2x - 1y = -12$.

Q1742. Cross-multiplying $\frac{x+5}{y-5} = \frac{1}{2} \implies 2(x+5) = 1(y-5) \implies 2x - 1y = -15$.

Q1743. Cross-multiplying $\frac{x+3}{y-2} = \frac{2}{5} \implies 5(x+3) = 2(y-2) \implies 5x - 2y = -19$.

Q1744. Cross-multiplying $\frac{x+1}{y-5} = \frac{3}{2} \implies 2(x+1) = 3(y-5) \implies 2x - 3y = -17$.

Q1745. Cross-multiplying $\frac{x+2}{y-2} = \frac{3}{5} \implies 5(x+2) = 3(y-2) \implies 5x - 3y = -16$.

Q1746. Cross-multiplying $\frac{x+1}{y-3} = \frac{2}{3} \implies 3(x+1) = 2(y-3) \implies 3x - 2y = -9$.

Q1747. Cross-multiplying $\frac{x+4}{y-5} = \frac{2}{2} \implies 2(x+4) = 2(y-5) \implies 2x - 2y = -18$.

Q1748. Cross-multiplying $\frac{x+1}{y-5} = \frac{3}{5} \implies 5(x+1) = 3(y-5) \implies 5x - 3y = -20$.

Q1749. Cross-multiplying $\frac{x+3}{y-3} = \frac{3}{2} \implies 2(x+3) = 3(y-3) \implies 2x - 3y = -15$.

Q1750. Cross-multiplying $\frac{x+1}{y-3} = \frac{1}{4} \implies 4(x+1) = 1(y-3) \implies 4x - 1y = -7$.

Q1751. Upstream speed $= x - y \implies T_1 = \frac{30}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{30}{x+y}$. Total time $T = \frac{30}{x-y} + \frac{30}{x+y}$.

Q1752. Upstream speed $= x - y \implies T_1 = \frac{29}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{29}{x+y}$. Total time $T = \frac{29}{x-y} + \frac{29}{x+y}$.

Q1753. Upstream speed $= x - y \implies T_1 = \frac{23}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{23}{x+y}$. Total time $T = \frac{23}{x-y} + \frac{23}{x+y}$.

Q1754. Upstream speed $= x - y \implies T_1 = \frac{56}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{56}{x+y}$. Total time $T = \frac{56}{x-y} + \frac{56}{x+y}$.

Q1755. Upstream speed $= x - y \implies T_1 = \frac{34}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{34}{x+y}$. Total time $T = \frac{34}{x-y} + \frac{34}{x+y}$.

Q1756. Upstream speed $= x - y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1757. Upstream speed $= x - y \implies T_1 = \frac{29}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{29}{x+y}$. Total time $T = \frac{29}{x-y} + \frac{29}{x+y}$.

Q1758. Upstream speed $= x - y \implies T_1 = \frac{38}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{38}{x+y}$. Total time $T = \frac{38}{x-y} + \frac{38}{x+y}$.

Q1759. Upstream speed $= x - y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1760. Upstream speed $= x - y \implies T_1 = \frac{35}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{35}{x+y}$. Total time $T = \frac{35}{x-y} + \frac{35}{x+y}$.

Q1761. Upstream speed $= x - y \implies T_1 = \frac{49}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{49}{x+y}$. Total time $T = \frac{49}{x-y} + \frac{49}{x+y}$.

Q1762. Upstream speed $= x - y \implies T_1 = \frac{25}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{25}{x+y}$. Total time $T = \frac{25}{x-y} + \frac{25}{x+y}$.

Q1763. Upstream speed $= x - y \implies T_1 = \frac{39}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{39}{x+y}$. Total time $T = \frac{39}{x-y} + \frac{39}{x+y}$.

Q1764. Upstream speed $= x - y \implies T_1 = \frac{60}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{60}{x+y}$. Total time $T = \frac{60}{x-y} + \frac{60}{x+y}$.

Q1765. Upstream speed $= x - y \implies T_1 = \frac{21}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{21}{x+y}$. Total time $T = \frac{21}{x-y} + \frac{21}{x+y}$.

Q1766. Upstream speed $= x - y \implies T_1 = \frac{34}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{34}{x+y}$. Total time $T = \frac{34}{x-y} + \frac{34}{x+y}$.

Q1767. Upstream speed $= x - y \implies T_1 = \frac{26}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{26}{x+y}$. Total time $T = \frac{26}{x-y} + \frac{26}{x+y}$.

Q1768. Upstream speed $= x - y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1769. Upstream speed $= x - y \implies T_1 = \frac{39}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{39}{x+y}$. Total time $T = \frac{39}{x-y} + \frac{39}{x+y}$.

Q1770. Upstream speed $= x - y \implies T_1 = \frac{45}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{45}{x+y}$. Total time $T = \frac{45}{x-y} + \frac{45}{x+y}$.

Q1771. Upstream speed $= x - y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1772. Upstream speed $= x - y \implies T_1 = \frac{28}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{28}{x+y}$. Total time $T = \frac{28}{x-y} + \frac{28}{x+y}$.

Q1773. Upstream speed $= x - y \implies T_1 = \frac{21}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{21}{x+y}$. Total time $T = \frac{21}{x-y} + \frac{21}{x+y}$.

Q1774. Upstream speed $= x-y \implies T_1 = \frac{55}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{55}{x+y}$. Total time $T = \frac{55}{x-y} + \frac{55}{x+y}$.

Q1775. Upstream speed $= x-y \implies T_1 = \frac{56}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{56}{x+y}$. Total time $T = \frac{56}{x-y} + \frac{56}{x+y}$.

Q1776. Upstream speed $= x-y \implies T_1 = \frac{59}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{59}{x+y}$. Total time $T = \frac{59}{x-y} + \frac{59}{x+y}$.

Q1777. Upstream speed $= x-y \implies T_1 = \frac{24}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{24}{x+y}$. Total time $T = \frac{24}{x-y} + \frac{24}{x+y}$.

Q1778. Upstream speed $= x-y \implies T_1 = \frac{34}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{34}{x+y}$. Total time $T = \frac{34}{x-y} + \frac{34}{x+y}$.

Q1779. Upstream speed $= x-y \implies T_1 = \frac{25}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{25}{x+y}$. Total time $T = \frac{25}{x-y} + \frac{25}{x+y}$.

Q1780. Upstream speed $= x-y \implies T_1 = \frac{33}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{33}{x+y}$. Total time $T = \frac{33}{x-y} + \frac{33}{x+y}$.

Q1781. Upstream speed $= x-y \implies T_1 = \frac{29}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{29}{x+y}$. Total time $T = \frac{29}{x-y} + \frac{29}{x+y}$.

Q1782. Upstream speed $= x-y \implies T_1 = \frac{55}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{55}{x+y}$. Total time $T = \frac{55}{x-y} + \frac{55}{x+y}$.

Q1783. Upstream speed $= x-y \implies T_1 = \frac{39}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{39}{x+y}$. Total time $T = \frac{39}{x-y} + \frac{39}{x+y}$.

Q1784. Upstream speed $= x-y \implies T_1 = \frac{57}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{57}{x+y}$. Total time $T = \frac{57}{x-y} + \frac{57}{x+y}$.

Q1785. Upstream speed $= x-y \implies T_1 = \frac{32}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{32}{x+y}$. Total time $T = \frac{32}{x-y} + \frac{32}{x+y}$.

Q1786. Upstream speed $= x-y \implies T_1 = \frac{47}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{47}{x+y}$. Total time $T = \frac{47}{x-y} + \frac{47}{x+y}$.

Q1787. Upstream speed $= x-y \implies T_1 = \frac{54}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{54}{x+y}$. Total time $T = \frac{54}{x-y} + \frac{54}{x+y}$.

Q1788. Upstream speed $= x-y \implies T_1 = \frac{41}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{41}{x+y}$. Total time $T = \frac{41}{x-y} + \frac{41}{x+y}$.

Q1789. Upstream speed $= x-y \implies T_1 = \frac{24}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{24}{x+y}$. Total time $T = \frac{24}{x-y} + \frac{24}{x+y}$.

Q1790. Upstream speed $= x-y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1791. Upstream speed $= x-y \implies T_1 = \frac{54}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{54}{x+y}$. Total time $T = \frac{54}{x-y} + \frac{54}{x+y}$.

Q1792. Upstream speed $= x-y \implies T_1 = \frac{36}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{36}{x+y}$. Total time $T = \frac{36}{x-y} + \frac{36}{x+y}$.

Q1793. Upstream speed $= x-y \implies T_1 = \frac{32}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{32}{x+y}$. Total time $T = \frac{32}{x-y} + \frac{32}{x+y}$.

Q1794. Upstream speed $= x-y \implies T_1 = \frac{28}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{28}{x+y}$. Total time $T = \frac{28}{x-y} + \frac{28}{x+y}$.

Q1795. Upstream speed $= x-y \implies T_1 = \frac{58}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{58}{x+y}$. Total time $T = \frac{58}{x-y} + \frac{58}{x+y}$.

Q1796. Upstream speed $= x-y \implies T_1 = \frac{30}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{30}{x+y}$. Total time $T = \frac{30}{x-y} + \frac{30}{x+y}$.

Q1797. Upstream speed $= x-y \implies T_1 = \frac{58}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{58}{x+y}$. Total time $T = \frac{58}{x-y} + \frac{58}{x+y}$.

Q1798. Upstream speed $= x-y \implies T_1 = \frac{45}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{45}{x+y}$. Total time $T = \frac{45}{x-y} + \frac{45}{x+y}$.

Q1799. Upstream speed $= x-y \implies T_1 = \frac{41}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{41}{x+y}$. Total time $T = \frac{41}{x-y} + \frac{41}{x+y}$.

Q1800. Upstream speed $= x-y \implies T_1 = \frac{29}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{29}{x+y}$. Total time $T = \frac{29}{x-y} + \frac{29}{x+y}$.

Q1801. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1802. Upstream speed $= x-y \implies T_1 = \frac{31}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{31}{x+y}$. Total time $T = \frac{31}{x-y} + \frac{31}{x+y}$.

Q1803. Upstream speed $= x-y \implies T_1 = \frac{44}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{44}{x+y}$. Total time $T = \frac{44}{x-y} + \frac{44}{x+y}$.

Q1804. Upstream speed $= x-y \implies T_1 = \frac{31}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{31}{x+y}$. Total time $T = \frac{31}{x-y} + \frac{31}{x+y}$.

Q1805. Upstream speed $= x-y \implies T_1 = \frac{31}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{31}{x+y}$. Total time $T = \frac{31}{x-y} + \frac{31}{x+y}$.

Q1806. Upstream speed $= x-y \implies T_1 = \frac{56}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{56}{x+y}$. Total time $T = \frac{56}{x-y} + \frac{56}{x+y}$.

Q1807. Upstream speed $= x-y \implies T_1 = \frac{40}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{40}{x+y}$. Total time $T = \frac{40}{x-y} + \frac{40}{x+y}$.

Q1808. Upstream speed $= x-y \implies T_1 = \frac{27}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{27}{x+y}$. Total time $T = \frac{27}{x-y} + \frac{27}{x+y}$.

Q1809. Upstream speed $= x-y \implies T_1 = \frac{50}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{50}{x+y}$. Total time $T = \frac{50}{x-y} + \frac{50}{x+y}$.

Q1810. Upstream speed $= x-y \implies T_1 = \frac{35}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{35}{x+y}$. Total time $T = \frac{35}{x-y} + \frac{35}{x+y}$.

Q1811. Upstream speed $= x-y \implies T_1 = \frac{39}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{39}{x+y}$. Total time $T = \frac{39}{x-y} + \frac{39}{x+y}$.

Q1812. Upstream speed $= x-y \implies T_1 = \frac{54}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{54}{x+y}$. Total time $T = \frac{54}{x-y} + \frac{54}{x+y}$.

Q1813. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1814. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1815. Upstream speed $= x-y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1816. Upstream speed $= x-y \implies T_1 = \frac{33}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{33}{x+y}$. Total time $T = \frac{33}{x-y} + \frac{33}{x+y}$.

Q1817. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1818. Upstream speed $= x-y \implies T_1 = \frac{55}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{55}{x+y}$. Total time $T = \frac{55}{x-y} + \frac{55}{x+y}$.

Q1819. Upstream speed $= x-y \implies T_1 = \frac{34}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{34}{x+y}$. Total time $T = \frac{34}{x-y} + \frac{34}{x+y}$.

Q1820. Upstream speed $= x-y \implies T_1 = \frac{52}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{52}{x+y}$. Total time $T = \frac{52}{x-y} + \frac{52}{x+y}$.

Q1821. Upstream speed $= x-y \implies T_1 = \frac{33}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{33}{x+y}$. Total time $T = \frac{33}{x-y} + \frac{33}{x+y}$.

Q1822. Upstream speed $= x-y \implies T_1 = \frac{33}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{33}{x+y}$. Total time $T = \frac{33}{x-y} + \frac{33}{x+y}$.

Q1823. Upstream speed $= x-y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1824. Upstream speed $= x-y \implies T_1 = \frac{34}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{34}{x+y}$. Total time $T = \frac{34}{x-y} + \frac{34}{x+y}$.

Q1825. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1826. Upstream speed $= x-y \implies T_1 = \frac{22}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{22}{x+y}$. Total time $T = \frac{22}{x-y} + \frac{22}{x+y}$.

Q1827. Upstream speed $= x-y \implies T_1 = \frac{25}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{25}{x+y}$. Total time $T = \frac{25}{x-y} + \frac{25}{x+y}$.

Q1828. Upstream speed $= x-y \implies T_1 = \frac{49}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{49}{x+y}$. Total time $T = \frac{49}{x-y} + \frac{49}{x+y}$.

Q1829. Upstream speed $= x-y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1830. Upstream speed $= x-y \implies T_1 = \frac{37}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{37}{x+y}$. Total time $T = \frac{37}{x-y} + \frac{37}{x+y}$.

Q1831. Upstream speed $= x-y \implies T_1 = \frac{25}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{25}{x+y}$. Total time $T = \frac{25}{x-y} + \frac{25}{x+y}$.

Q1832. Upstream speed $= x-y \implies T_1 = \frac{51}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{51}{x+y}$. Total time $T = \frac{51}{x-y} + \frac{51}{x+y}$.

Q1833. Upstream speed $= x-y \implies T_1 = \frac{56}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{56}{x+y}$. Total time $T = \frac{56}{x-y} + \frac{56}{x+y}$.

Q1834. Upstream speed $= x-y \implies T_1 = \frac{50}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{50}{x+y}$. Total time $T = \frac{50}{x-y} + \frac{50}{x+y}$.

Q1835. Upstream speed $= x-y \implies T_1 = \frac{40}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{40}{x+y}$. Total time $T = \frac{40}{x-y} + \frac{40}{x+y}$.

Q1836. Upstream speed $= x-y \implies T_1 = \frac{49}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{49}{x+y}$. Total time $T = \frac{49}{x-y} + \frac{49}{x+y}$.

Q1837. Upstream speed $= x-y \implies T_1 = \frac{56}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{56}{x+y}$. Total time $T = \frac{56}{x-y} + \frac{56}{x+y}$.

Q1838. Upstream speed $= x-y \implies T_1 = \frac{24}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{24}{x+y}$. Total time $T = \frac{24}{x-y} + \frac{24}{x+y}$.

Q1839. Upstream speed $= x-y \implies T_1 = \frac{36}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{36}{x+y}$. Total time $T = \frac{36}{x-y} + \frac{36}{x+y}$.

Q1840. Upstream speed $= x-y \implies T_1 = \frac{29}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{29}{x+y}$. Total time $T = \frac{29}{x-y} + \frac{29}{x+y}$.

Q1841. Upstream speed $= x-y \implies T_1 = \frac{50}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{50}{x+y}$. Total time $T = \frac{50}{x-y} + \frac{50}{x+y}$.

Q1842. Upstream speed $= x-y \implies T_1 = \frac{55}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{55}{x+y}$. Total time $T = \frac{55}{x-y} + \frac{55}{x+y}$.

Q1843. Upstream speed $= x-y \implies T_1 = \frac{36}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{36}{x+y}$. Total time $T = \frac{36}{x-y} + \frac{36}{x+y}$.

Q1844. Upstream speed $= x-y \implies T_1 = \frac{37}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{37}{x+y}$. Total time $T = \frac{37}{x-y} + \frac{37}{x+y}$.

Q1845. Upstream speed $= x-y \implies T_1 = \frac{28}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{28}{x+y}$. Total time $T = \frac{28}{x-y} + \frac{28}{x+y}$.

Q1846. Upstream speed $= x-y \implies T_1 = \frac{20}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{20}{x+y}$. Total time $T = \frac{20}{x-y} + \frac{20}{x+y}$.

Q1847. Upstream speed $= x-y \implies T_1 = \frac{39}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{39}{x+y}$. Total time $T = \frac{39}{x-y} + \frac{39}{x+y}$.

Q1848. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1849. Upstream speed $= x-y \implies T_1 = \frac{45}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{45}{x+y}$. Total time $T = \frac{45}{x-y} + \frac{45}{x+y}$.

Q1850. Upstream speed $= x-y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1851. Upstream speed $= x-y \implies T_1 = \frac{50}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{50}{x+y}$. Total time $T = \frac{50}{x-y} + \frac{50}{x+y}$.

Q1852. Upstream speed $= x-y \implies T_1 = \frac{32}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{32}{x+y}$. Total time $T = \frac{32}{x-y} + \frac{32}{x+y}$.

Q1853. Upstream speed $= x-y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1854. Upstream speed $= x-y \implies T_1 = \frac{33}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{33}{x+y}$. Total time $T = \frac{33}{x-y} + \frac{33}{x+y}$.

Q1855. Upstream speed $= x-y \implies T_1 = \frac{49}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{49}{x+y}$. Total time $T = \frac{49}{x-y} + \frac{49}{x+y}$.

Q1856. Upstream speed $= x-y \implies T_1 = \frac{32}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{32}{x+y}$. Total time $T = \frac{32}{x-y} + \frac{32}{x+y}$.

Q1857. Upstream speed $= x-y \implies T_1 = \frac{54}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{54}{x+y}$. Total time $T = \frac{54}{x-y} + \frac{54}{x+y}$.

Q1858. Upstream speed $= x-y \implies T_1 = \frac{49}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{49}{x+y}$. Total time $T = \frac{49}{x-y} + \frac{49}{x+y}$.

Q1859. Upstream speed $= x-y \implies T_1 = \frac{37}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{37}{x+y}$. Total time $T = \frac{37}{x-y} + \frac{37}{x+y}$.

Q1860. Upstream speed $= x-y \implies T_1 = \frac{47}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{47}{x+y}$. Total time $T = \frac{47}{x-y} + \frac{47}{x+y}$.

Q1861. Upstream speed $= x-y \implies T_1 = \frac{50}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{50}{x+y}$. Total time $T = \frac{50}{x-y} + \frac{50}{x+y}$.

Q1862. Upstream speed $= x-y \implies T_1 = \frac{27}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{27}{x+y}$. Total time $T = \frac{27}{x-y} + \frac{27}{x+y}$.

Q1863. Upstream speed $= x-y \implies T_1 = \frac{32}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{32}{x+y}$. Total time $T = \frac{32}{x-y} + \frac{32}{x+y}$.

Q1864. Upstream speed $= x-y \implies T_1 = \frac{23}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{23}{x+y}$. Total time $T = \frac{23}{x-y} + \frac{23}{x+y}$.

Q1865. Upstream speed $= x-y \implies T_1 = \frac{60}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{60}{x+y}$. Total time $T = \frac{60}{x-y} + \frac{60}{x+y}$.

Q1866. Upstream speed $= x-y \implies T_1 = \frac{48}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{48}{x+y}$. Total time $T = \frac{48}{x-y} + \frac{48}{x+y}$.

Q1867. Upstream speed $= x-y \implies T_1 = \frac{41}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{41}{x+y}$. Total time $T = \frac{41}{x-y} + \frac{41}{x+y}$.

Q1868. Upstream speed $= x-y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1869. Upstream speed $= x-y \implies T_1 = \frac{48}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{48}{x+y}$. Total time $T = \frac{48}{x-y} + \frac{48}{x+y}$.

Q1870. Upstream speed $= x-y \implies T_1 = \frac{24}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{24}{x+y}$. Total time $T = \frac{24}{x-y} + \frac{24}{x+y}$.

Q1871. Upstream speed $= x-y \implies T_1 = \frac{53}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{53}{x+y}$. Total time $T = \frac{53}{x-y} + \frac{53}{x+y}$.

Q1872. Upstream speed $= x-y \implies T_1 = \frac{42}{x-y}$. Downstream speed $= x+y \implies T_2 = \frac{42}{x+y}$. Total time $T = \frac{42}{x-y} + \frac{42}{x+y}$.

Q1873. Upstream speed $= x - y \implies T_1 = \frac{43}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{43}{x+y}$. Total time $T = \frac{43}{x-y} + \frac{43}{x+y}$.

Q1874. Upstream speed $= x - y \implies T_1 = \frac{24}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{24}{x+y}$. Total time $T = \frac{24}{x-y} + \frac{24}{x+y}$.

Q1875. Upstream speed $= x - y \implies T_1 = \frac{59}{x-y}$. Downstream speed $= x + y \implies T_2 = \frac{59}{x+y}$. Total time $T = \frac{59}{x-y} + \frac{59}{x+y}$.

Q1876. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{5}{x} + \frac{4}{y} = \frac{1}{9}$.

Q1877. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{3}{y} = \frac{1}{4}$.

Q1878. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{7}$.

Q1879. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{5}{y} = \frac{1}{4}$.

Q1880. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{2}{x} + \frac{7}{y} = \frac{1}{6}$.

Q1881. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{2}{x} + \frac{8}{y} = \frac{1}{4}$.

Q1882. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{3}{x} + \frac{5}{y} = \frac{1}{8}$.

Q1883. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1884. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{5}$.

Q1885. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{4}{x} + \frac{4}{y} = \frac{1}{11}$.

Q1886. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{3}{x} + \frac{4}{y} = \frac{1}{9}$.

Q1887. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{3}{x} + \frac{3}{y} = \frac{1}{11}$.

Q1888. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{7}$.

Q1889. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{3}{y} = \frac{1}{7}$.

Q1890. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{2}{x} + \frac{7}{y} = \frac{1}{11}$.

Q1891. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{4}{x} + \frac{7}{y} = \frac{1}{7}$.

Q1892. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{2}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1893. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{5}{x} + \frac{4}{y} = \frac{1}{5}$.

Q1894. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{3}{x} + \frac{7}{y} = \frac{1}{4}$.

Q1895. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{2}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1896. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{4}$.

Q1897. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{11}$.

Q1898. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{4}$.

Q1899. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{8}$.

Q1900. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{2}{x} + \frac{4}{y} = \frac{1}{8}$.

Q1901. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{2}{x} + \frac{7}{y} = \frac{1}{11}$.

Q1902. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{2}{x} + \frac{8}{y} = \frac{1}{7}$.

Q1903. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{5}$.

Q1904. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{3}{x} + \frac{6}{y} = \frac{1}{11}$.

Q1905. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{4}{y} = \frac{1}{4}$.

Q1906. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{4}{x} + \frac{8}{y} = \frac{1}{6}$.

Q1907. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{9}$.

Q1908. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{5}{x} + \frac{7}{y} = \frac{1}{11}$.

Q1909. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{4}{x} + \frac{7}{y} = \frac{1}{12}$.

Q1910. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{6}{x} + \frac{7}{y} = \frac{1}{9}$.

Q1911. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{3}{x} + \frac{5}{y} = \frac{1}{6}$.

Q1912. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1913. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{5}$.

Q1914. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{5}{x} + \frac{7}{y} = \frac{1}{8}$.

Q1915. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{3}{x} + \frac{8}{y} = \frac{1}{7}$.

Q1916. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{5}{x} + \frac{5}{y} = \frac{1}{6}$.

Q1917. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{5}{x} + \frac{7}{y} = \frac{1}{4}$.

Q1918. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{6}{x} + \frac{6}{y} = \frac{1}{12}$.

Q1919. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{5}{x} + \frac{4}{y} = \frac{1}{12}$.

Q1920. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{2}{x} + \frac{6}{y} = \frac{1}{4}$.

Q1921. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{4}$.

Q1922. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{4}{y} = \frac{1}{4}$.

Q1923. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{2}{x} + \frac{6}{y} = \frac{1}{5}$.

Q1924. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{2}{x} + \frac{4}{y} = \frac{1}{8}$.

Q1925. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{7}{y} = \frac{1}{7}$.

Q1926. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{5}{x} + \frac{3}{y} = \frac{1}{5}$.

Q1927. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{10}$.

Q1928. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{9}$.

Q1929. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{3}{y} = \frac{1}{7}$.

Q1930. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{4}{x} + \frac{7}{y} = \frac{1}{6}$.

Q1931. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{11}$.

Q1932. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1933. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{5}{x} + \frac{3}{y} = \frac{1}{12}$.

Q1934. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{3}{x} + \frac{3}{y} = \frac{1}{10}$.

Q1935. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{4}{x} + \frac{7}{y} = \frac{1}{12}$.

Q1936. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{5}{x} + \frac{8}{y} = \frac{1}{6}$.

Q1937. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{6}{x} + \frac{7}{y} = \frac{1}{10}$.

Q1938. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1939. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{2}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1940. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{2}{x} + \frac{6}{y} = \frac{1}{8}$.

Q1941. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{4}{x} + \frac{8}{y} = \frac{1}{6}$.

Q1942. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{6}{x} + \frac{7}{y} = \frac{1}{4}$.

Q1943. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{3}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1944. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{3}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1945. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{2}{x} + \frac{5}{y} = \frac{1}{11}$.

Q1946. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{10}$.

Q1947. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{5}{x} + \frac{8}{y} = \frac{1}{4}$.

Q1948. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{4}{x} + \frac{5}{y} = \frac{1}{6}$.

Q1949. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{3}{x} + \frac{3}{y} = \frac{1}{5}$.

Q1950. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{2}{x} + \frac{3}{y} = \frac{1}{6}$.

Q1951. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{3}{x} + \frac{4}{y} = \frac{1}{11}$.

Q1952. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{5}$.

Q1953. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{2}{x} + \frac{8}{y} = \frac{1}{4}$.

Q1954. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{6}{x} + \frac{4}{y} = \frac{1}{11}$.

Q1955. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{11}$.

Q1956. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{10}$.

Q1957. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{2}{x} + \frac{5}{y} = \frac{1}{12}$.

Q1958. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{3}{x} + \frac{6}{y} = \frac{1}{4}$.

Q1959. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{5}{x} + \frac{4}{y} = \frac{1}{6}$.

Q1960. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{3}{x} + \frac{4}{y} = \frac{1}{6}$.

Q1961. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{2}{x} + \frac{4}{y} = \frac{1}{9}$.

Q1962. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{3}{x} + \frac{5}{y} = \frac{1}{10}$.

Q1963. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{6}{x} + \frac{7}{y} = \frac{1}{5}$.

Q1964. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{3}{x} + \frac{3}{y} = \frac{1}{10}$.

Q1965. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{4}$.

Q1966. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1967. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{2}{x} + \frac{7}{y} = \frac{1}{7}$.

Q1968. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{6}$.

Q1969. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{4}{x} + \frac{7}{y} = \frac{1}{12}$.

Q1970. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{5}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1971. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 7\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{5}{x} + \frac{7}{y} = \frac{1}{5}$.

Q1972. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{7}$.

Q1973. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{5}{y} = \frac{1}{7}$.

Q1974. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1975. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{2}{x} + \frac{3}{y} = \frac{1}{4}$.

Q1976. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{5}{x} + \frac{5}{y} = \frac{1}{9}$.

Q1977. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{5}{x} + \frac{8}{y} = \frac{1}{7}$.

Q1978. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{6}{x} + \frac{3}{y} = \frac{1}{7}$.

Q1979. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{6}{x} + \frac{6}{y} = \frac{1}{6}$.

Q1980. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{2}{x} + \frac{4}{y} = \frac{1}{7}$.

Q1981. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{5}{x} + \frac{6}{y} = \frac{1}{4}$.

Q1982. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 5\left(\frac{1}{y}\right) = \frac{1}{5} \implies \frac{6}{x} + \frac{5}{y} = \frac{1}{5}$.

Q1983. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{3}{x} + \frac{8}{y} = \frac{1}{12}$.

Q1984. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{11}$.

Q1985. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{9} \implies \frac{3}{x} + \frac{3}{y} = \frac{1}{9}$.

Q1986. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{10} \implies \frac{6}{x} + \frac{8}{y} = \frac{1}{10}$.

Q1987. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{2}{x} + \frac{4}{y} = \frac{1}{4}$.

Q1988. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{4}$.

Q1989. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $2\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{11} \implies \frac{2}{x} + \frac{8}{y} = \frac{1}{11}$.

Q1990. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{4}$.

Q1991. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{7} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{7}$.

Q1992. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{4}{x} + \frac{3}{y} = \frac{1}{8}$.

Q1993. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{3}{x} + \frac{6}{y} = \frac{1}{6}$.

Q1994. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 4\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{6}{x} + \frac{4}{y} = \frac{1}{6}$.

Q1995. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 6\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{4}{x} + \frac{6}{y} = \frac{1}{8}$.

Q1996. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $5\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{6} \implies \frac{5}{x} + \frac{3}{y} = \frac{1}{6}$.

Q1997. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{3}{x} + \frac{8}{y} = \frac{1}{4}$.

Q1998. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $3\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{8} \implies \frac{3}{x} + \frac{8}{y} = \frac{1}{8}$.

Q1999. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $6\left(\frac{1}{x}\right) + 3\left(\frac{1}{y}\right) = \frac{1}{12} \implies \frac{6}{x} + \frac{3}{y} = \frac{1}{12}$.

Q2000. Work done by 1 man in 1 day $= \frac{1}{x}$, by 1 woman $= \frac{1}{y}$. Equation: $4\left(\frac{1}{x}\right) + 8\left(\frac{1}{y}\right) = \frac{1}{4} \implies \frac{4}{x} + \frac{8}{y} = \frac{1}{4}$.

MODULE 3: METHOD SELECTION STRATEGY (WHAT METHOD WHEN AND WHERE) (Q2001 – Q3000) — Answer Key

Q2001. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2002. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2003. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2004. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2005. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2006. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2007. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2008. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2009. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{7}$, $\frac{b_1}{b_2} = \frac{5}{10}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2010. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2011. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2012. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2013. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2014. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2015. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{5}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2016. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2017. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2018. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2019. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2020. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2021. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{9}$, $\frac{b_1}{b_2} = \frac{6}{24}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2022. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2023. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2024. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{13}$, $\frac{b_1}{b_2} = \frac{6}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2025. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2026. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2027. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{7}$, $\frac{b_1}{b_2} = \frac{2}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2028. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2029. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2030. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2031. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2032. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2033. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{9}$, $\frac{b_1}{b_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2034. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2035. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2036. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{9}$, $\frac{b_1}{b_2} = \frac{4}{16}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2037. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2038. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2039. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{21}$, $\frac{b_1}{b_2} = \frac{3}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2040. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2041. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2042. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2043. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2044. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2045. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{19}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2046. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2047. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2048. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{7}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2049. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2050. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2051. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{21}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2052. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2053. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2054. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{13}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2055. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2056. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2057. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2058. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2059. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2060. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{7}$, $\frac{b_1}{b_2} = \frac{2}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2061. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2062. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2063. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{13}$, $\frac{b_1}{b_2} = \frac{6}{18}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2064. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2065. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2066. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{21}$, $\frac{b_1}{b_2} = \frac{3}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2067. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2068. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2069. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{13}$, $\frac{b_1}{b_2} = \frac{3}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2070. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2071. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2072. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{3}{9}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2073. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2074. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2075. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2076. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2077. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2078. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2079. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2080. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2081. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2082. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2083. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2084. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{2}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2085. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2086. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2087. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2088. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2089. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2090. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{13}$, $\frac{b_1}{b_2} = \frac{6}{18}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2091. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2092. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2093. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{9}$, $\frac{b_1}{b_2} = \frac{4}{16}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2094. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2095. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2096. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{11}$, $\frac{b_1}{b_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2097. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2098. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2099. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{13}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2100. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2101. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2102. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{13}$, $\frac{b_1}{b_2} = \frac{4}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2103. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2104. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2105. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{11}$, $\frac{b_1}{b_2} = \frac{6}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2106. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2107. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2108. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{4}{16}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2109. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2110. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2111. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{4}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2112. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2113. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2114. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2115. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2116. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2117. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{5}$, $\frac{b_1}{b_2} = \frac{6}{24}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2118. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2119. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2120. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{16}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2121. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2122. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2123. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{19}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2124. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2125. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2126. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{2}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2127. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2128. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2129. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{7}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2130. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2131. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2132. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{11}$, $\frac{b_1}{b_2} = \frac{2}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2133. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2134. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2135. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{21}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2136. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2137. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2138. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{9}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2139. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2140. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2141. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{5}$, $\frac{b_1}{b_2} = \frac{5}{10}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2142. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2143. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2144. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2145. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2146. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2147. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{19}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2148. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2149. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2150. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{16}$, $\frac{b_1}{b_2} = \frac{3}{9}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2151. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2152. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2153. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{7}$, $\frac{b_1}{b_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2154. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2155. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2156. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{7}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2157. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2158. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2159. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2160. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2161. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2162. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2163. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2164. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2165. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{3}{9}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2166. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2167. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2168. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{13}$, $\frac{b_1}{b_2} = \frac{3}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2169. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2170. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2171. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2172. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2173. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2174. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{9}$, $\frac{b_1}{b_2} = \frac{3}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2175. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2176. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2177. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{9}$, $\frac{b_1}{b_2} = \frac{3}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2178. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2179. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2180. Compare ratios: $\frac{a_1}{a_2} = \frac{5}{16}$, $\frac{b_1}{b_2} = \frac{4}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2181. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2182. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2183. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{9}$, $\frac{b_1}{b_2} = \frac{3}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2184. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2185. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2186. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{19}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2187. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2188. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2189. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{6}{18}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2190. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2191. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2192. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{6}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2193. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2194. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2195. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2196. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2197. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2198. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{9}$, $\frac{b_1}{b_2} = \frac{5}{20}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2199. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2200. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2201. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{5}{10}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2202. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2203. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2204. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{5}$, $\frac{b_1}{b_2} = \frac{4}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2205. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2206. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2207. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{7}$, $\frac{b_1}{b_2} = \frac{2}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2208. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2209. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2210. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{2}{8}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2211. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2212. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2213. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{7}$, $\frac{b_1}{b_2} = \frac{2}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2214. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2215. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2216. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{19}$, $\frac{b_1}{b_2} = \frac{2}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2217. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2218. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2219. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{4}{16}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2220. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2221. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2222. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{5}$, $\frac{b_1}{b_2} = \frac{6}{24}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2223. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2224. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2225. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{17}$, $\frac{b_1}{b_2} = \frac{6}{24}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2226. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2227. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2228. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2229. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2230. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2231. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{2}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2232. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2233. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2234. Compare ratios: $\frac{a_1}{a_2} = \frac{6}{25}$, $\frac{b_1}{b_2} = \frac{3}{12}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2235. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2236. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2237. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{10}$, $\frac{b_1}{b_2} = \frac{5}{15}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2238. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, $\frac{c_1}{c_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2239. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, but $\frac{c_1}{c_2} \neq \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2240. Compare ratios: $\frac{a_1}{a_2} = \frac{4}{13}$, $\frac{b_1}{b_2} = \frac{2}{6}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2241. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2242. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{2}$, $\frac{b_1}{b_2} = \frac{1}{2}$, but $\frac{c_1}{c_2} \neq \frac{1}{2}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2243. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{13}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2244. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2245. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2246. Compare ratios: $\frac{a_1}{a_2} = \frac{2}{5}$, $\frac{b_1}{b_2} = \frac{1}{2}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2247. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{3}$, $\frac{b_1}{b_2} = \frac{1}{3}$, $\frac{c_1}{c_2} = \frac{1}{3}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2248. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, but $\frac{c_1}{c_2} \neq \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$, the system has **No Solution** (Parallel lines).

Q2249. Compare ratios: $\frac{a_1}{a_2} = \frac{3}{13}$, $\frac{b_1}{b_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$, the system has a **Unique Solution** (Intersecting lines).

Q2250. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2251. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2252. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2253. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2254. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2255. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2256. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2257. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2258. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2259. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2260. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2261. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2262. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2263. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2264. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2265. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2266. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2267. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2268. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2269. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2270. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2271. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2272. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2273. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2274. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2275. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2276. **Substitution Method** is optimal here because

Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2277. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2278. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2279. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2280. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2281. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2282. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2283. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2284. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2285. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2286. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2287. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2288. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2289. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2290. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2291. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2292. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2293. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2294. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2295. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2296. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2297. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2298. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2299. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2300. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2301. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2302. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2303. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2304. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2305. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly elim-

inates x in one step.

Q2306. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2307. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2308. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2309. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2310. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2311. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2312. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2313. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2314. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2315. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2316. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2317. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2318. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2319. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2320. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2321. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2322. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2323. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2324. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2325. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2326. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2327. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2328. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2329. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2330. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2331. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2332. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2333. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2334. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2335. **Elimination Method** (by subtracting Eq 2

from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2336. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2337. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2338. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2339. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2340. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2341. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2342. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2343. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2344. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2345. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2346. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2347. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2348. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2349. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2350. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2351. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2352. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2353. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2354. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2355. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2356. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2357. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2358. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2359. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2360. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2361. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2362. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2363. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2364. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra

multiplication steps.

Q2365. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2366. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2367. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2368. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2369. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2370. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2371. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2372. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2373. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2374. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2375. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2376. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2377. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2378. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2379. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2380. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2381. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2382. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2383. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2384. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2385. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2386. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2387. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2388. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2389. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2390. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2391. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2392. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2393. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2394. **Substitution Method** is optimal here because

Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2395. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2396. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2397. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2398. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2399. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2400. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2401. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2402. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2403. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2404. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2405. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2406. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2407. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2408. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2409. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2410. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2411. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2412. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2413. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2414. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2415. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2416. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2417. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2418. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2419. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2420. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2421. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2422. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2423. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly elim-

inates x in one step.

Q2424. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2425. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2426. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2427. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2428. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2429. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2430. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2431. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2432. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2433. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2434. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2435. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2436. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2437. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2438. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2439. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2440. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2441. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2442. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2443. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2444. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2445. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2446. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2447. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2448. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2449. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2450. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2451. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2452. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2453. **Elimination Method** (by subtracting Eq 2

from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2454. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2455. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2456. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2457. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2458. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2459. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2460. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2461. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2462. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2463. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2464. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2465. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2466. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2467. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2468. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2469. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2470. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2471. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (3) for x. Subtracting directly eliminates x in one step.

Q2472. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2473. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2474. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2475. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2476. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2477. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2478. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2479. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2480. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2481. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2482. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra

multiplication steps.

Q2483. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2484. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2485. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2486. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2487. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2488. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2489. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (5) for x. Subtracting directly eliminates x in one step.

Q2490. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2491. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (8) for x. Subtracting directly eliminates x in one step.

Q2492. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2493. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2494. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2495. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (4) for x. Subtracting directly eliminates x in one step.

Q2496. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2497. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (7) for x. Subtracting directly eliminates x in one step.

Q2498. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2499. **Elimination Method** (by subtracting Eq 2 from Eq 1) is optimal because both equations share the exact same coefficient (6) for x. Subtracting directly eliminates x in one step.

Q2500. **Substitution Method** is optimal here because Equation (1) is already explicitly solved for x in terms of y. Substituting x directly into Equation (2) avoids extra multiplication steps.

Q2501. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2502. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(3) \times (2)$ whose sum is $b = 5$. Factoring yields $(x + 3)(x + 2) = 0$, giving $x = -3, -2$ instantly.

Q2503. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2504. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(2) \times (1)$ whose sum is $b = 3$. Factoring yields $(x + 2)(x + 1) = 0$, giving $x = -2, -1$ instantly.

Q2505. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2506. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(2) \times (2)$ whose sum is $b = 4$. Factoring yields $(x + 2)(x + 2) = 0$, giving $x = -2, -2$ instantly.

Q2507. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2508. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2509. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2510. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2511. Both work, but **Quadratic Formula** ($x = \frac{-b \pm \sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2512. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2513. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2514. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(6) \times (5)$ whose sum is $b = 11$. Factoring yields $(x + 6)(x + 5) = 0$, giving $x = -6, -5$ instantly.

Q2515. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2516. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(1) \times (4)$ whose sum is $b = 5$. Factoring yields $(x + 1)(x + 4) = 0$, giving $x = -1, -4$ instantly.

Q2517. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2518. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(6) \times (4)$ whose sum is $b = 10$. Factoring yields $(x + 6)(x + 4) = 0$, giving $x = -6, -4$ instantly.

Q2519. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2520. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2521. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2522. **Middle-Term Splitting** is ideal because $c = 36$ easily factors into $(6) \times (6)$ whose sum is $b = 12$. Factoring yields $(x + 6)(x + 6) = 0$, giving $x = -6, -6$ instantly.

Q2523. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2524. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2525. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2526. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(6) \times (1)$ whose sum is $b = 7$. Factoring yields $(x + 6)(x + 1) = 0$, giving $x = -6, -1$ instantly.

Q2527. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2528. **Middle-Term Splitting** is ideal because $c = 16$ easily factors into $(4) \times (4)$ whose sum is $b = 8$. Factoring yields $(x + 4)(x + 4) = 0$, giving $x = -4, -4$ instantly.

Q2529. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2530. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(4) \times (1)$ whose sum is $b = 5$. Factoring yields $(x + 4)(x + 1) = 0$, giving $x = -4, -1$ instantly.

Q2531. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2532. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(2) \times (3)$ whose sum is $b = 5$. Factoring yields $(x + 2)(x + 3) = 0$, giving $x = -2, -3$ instantly.

Q2533. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2534. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(2) \times (6)$ whose sum is $b = 8$. Factoring yields $(x + 2)(x + 6) = 0$, giving $x = -2, -6$ instantly.

Q2535. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2536. **Middle-Term Splitting** is ideal because $c = 1$ easily factors into $(1) \times (1)$ whose sum is $b = 2$. Factoring yields $(x + 1)(x + 1) = 0$, giving $x = -1, -1$ instantly.

Q2537. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2538. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2539. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2540. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2541. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2542. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2543. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2544. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(4) \times (6)$ whose sum is $b = 10$. Factoring yields $(x + 4)(x + 6) = 0$, giving $x = -4, -6$ instantly.

Q2545. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2546. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(1) \times (3)$ whose sum is $b = 4$. Factoring yields $(x + 1)(x + 3) = 0$, giving $x = -1, -3$ instantly.

Q2547. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2548. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2549. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2550. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2551. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2552. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2553. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2554. **Middle-Term Splitting** is ideal because $c = 25$ easily factors into $(5) \times (5)$ whose sum is $b = 10$. Factoring yields $(x + 5)(x + 5) = 0$, giving $x = -5, -5$ instantly.

Q2555. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2556. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2557. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2558. **Middle-Term Splitting** is ideal because $c = 10$ easily factors into $(2) \times (5)$ whose sum is $b = 7$. Factoring yields $(x + 2)(x + 5) = 0$, giving $x = -2, -5$ instantly.

Q2559. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2560. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(3) \times (4)$ whose sum is $b = 7$. Factoring yields $(x + 3)(x + 4) = 0$, giving $x = -3, -4$ instantly.

Q2561. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2562. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(1) \times (4)$ whose sum is $b = 5$. Factoring yields $(x + 1)(x + 4) = 0$, giving $x = -1, -4$ instantly.

Q2563. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2564. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(6) \times (3)$ whose sum is $b = 9$. Factoring yields $(x + 6)(x + 3) = 0$, giving $x = -6, -3$ instantly.

Q2565. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2566. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(2) \times (6)$ whose sum is $b = 8$. Factoring yields $(x + 2)(x + 6) = 0$, giving $x = -2, -6$ instantly.

Q2567. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2568. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(1) \times (5)$ whose sum is $b = 6$. Factoring yields $(x + 1)(x + 5) = 0$, giving $x = -1, -5$ instantly.

Q2569. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2570. **Middle-Term Splitting** is ideal because $c = 10$ easily factors into $(5) \times (2)$ whose sum is $b = 7$. Factoring yields $(x + 5)(x + 2) = 0$, giving $x = -5, -2$ instantly.

Q2571. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2572. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2573. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2574. **Middle-Term Splitting** is ideal because $c = 15$ easily factors into $(3) \times (5)$ whose sum is $b = 8$. Factoring yields $(x + 3)(x + 5) = 0$, giving $x = -3, -5$ instantly.

Q2575. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2576. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2577. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2578. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(3) \times (2)$ whose sum is $b = 5$. Factoring yields $(x + 3)(x + 2) = 0$, giving $x = -3, -2$ instantly.

Q2579. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2580. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2581. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2582. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(2) \times (3)$ whose sum is $b = 5$. Factoring yields $(x + 2)(x + 3) = 0$, giving $x = -2, -3$ instantly.

Q2583. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2584. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(1) \times (5)$ whose sum is $b = 6$. Factoring yields $(x + 1)(x + 5) = 0$, giving $x = -1, -5$ instantly.

Q2585. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2586. **Middle-Term Splitting** is ideal because $c = 1$ easily factors into $(1) \times (1)$ whose sum is $b = 2$. Factoring yields $(x + 1)(x + 1) = 0$, giving $x = -1, -1$ instantly.

Q2587. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2588. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2589. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2590. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(5) \times (4)$ whose sum is $b = 9$. Factoring yields $(x + 5)(x + 4) = 0$, giving $x = -5, -4$ instantly.

Q2591. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2592. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(5) \times (4)$ whose sum is $b = 9$. Factoring yields $(x + 5)(x + 4) = 0$, giving $x = -5, -4$ instantly.

Q2593. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2594. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(4) \times (5)$ whose sum is $b = 9$. Factoring yields $(x + 4)(x + 5) = 0$, giving $x = -4, -5$ instantly.

Q2595. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2596. **Middle-Term Splitting** is ideal because $c = 15$ easily factors into $(3) \times (5)$ whose sum is $b = 8$. Factoring yields $(x + 3)(x + 5) = 0$, giving $x = -3, -5$ instantly.

Q2597. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2598. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(1) \times (5)$ whose sum is $b = 6$. Factoring yields $(x + 1)(x + 5) = 0$, giving $x = -1, -5$ instantly.

Q2599. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2600. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2601. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2602. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2603. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2604. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2605. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2606. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2607. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2608. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(4) \times (6)$ whose sum is $b = 10$. Factoring yields $(x + 4)(x + 6) = 0$, giving $x = -4, -6$ instantly.

Q2609. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2610. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(1) \times (6)$ whose sum is $b = 7$. Factoring yields $(x + 1)(x + 6) = 0$, giving $x = -1, -6$ instantly.

Q2611. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2612. **Middle-Term Splitting** is ideal because $c = 1$ easily factors into $(1) \times (1)$ whose sum is $b = 2$. Factoring yields $(x + 1)(x + 1) = 0$, giving $x = -1, -1$ instantly.

Q2613. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2614. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(2) \times (1)$ whose sum is $b = 3$. Factoring yields $(x + 2)(x + 1) = 0$, giving $x = -2, -1$ instantly.

Q2615. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2616. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2617. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2618. **Middle-Term Splitting** is ideal because $c = 16$ easily factors into $(4) \times (4)$ whose sum is $b = 8$. Factoring yields $(x + 4)(x + 4) = 0$, giving $x = -4, -4$ instantly.

Q2619. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2620. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2621. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2622. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2623. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2624. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(5) \times (4)$ whose sum is $b = 9$. Factoring yields $(x + 5)(x + 4) = 0$, giving $x = -5, -4$ instantly.

Q2625. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2626. **Middle-Term Splitting** is ideal because $c = 10$ easily factors into $(2) \times (5)$ whose sum is $b = 7$. Factoring yields $(x + 2)(x + 5) = 0$, giving $x = -2, -5$ instantly.

Q2627. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2628. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2629. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2630. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(4) \times (1)$ whose sum is $b = 5$. Factoring yields $(x + 4)(x + 1) = 0$, giving $x = -4, -1$ instantly.

Q2631. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2632. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(6) \times (4)$ whose sum is $b = 10$. Factoring yields $(x + 6)(x + 4) = 0$, giving $x = -6, -4$ instantly.

Q2633. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2634. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(1) \times (6)$ whose sum is $b = 7$. Factoring yields $(x + 1)(x + 6) = 0$, giving $x = -1, -6$ instantly.

Q2635. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2636. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2637. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2638. **Middle-Term Splitting** is ideal because $c = 16$ easily factors into $(4) \times (4)$ whose sum is $b = 8$. Factoring yields $(x + 4)(x + 4) = 0$, giving $x = -4, -4$ instantly.

Q2639. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2640. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(2) \times (6)$ whose sum is $b = 8$. Factoring yields $(x + 2)(x + 6) = 0$, giving $x = -2, -6$ instantly.

Q2641. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2642. **Middle-Term Splitting** is ideal because $c = 10$ easily factors into $(5) \times (2)$ whose sum is $b = 7$. Factoring yields $(x + 5)(x + 2) = 0$, giving $x = -5, -2$ instantly.

Q2643. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2644. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(6) \times (1)$ whose sum is $b = 7$. Factoring yields $(x + 6)(x + 1) = 0$, giving $x = -6, -1$ instantly.

Q2645. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2646. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2647. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2648. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2649. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2650. **Middle-Term Splitting** is ideal because $c = 1$ easily factors into $(1) \times (1)$ whose sum is $b = 2$. Factoring yields $(x + 1)(x + 1) = 0$, giving $x = -1, -1$ instantly.

Q2651. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2652. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2653. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2654. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(4) \times (6)$ whose sum is $b = 10$. Factoring yields $(x + 4)(x + 6) = 0$, giving $x = -4, -6$ instantly.

Q2655. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2656. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(5) \times (1)$ whose sum is $b = 6$. Factoring yields $(x + 5)(x + 1) = 0$, giving $x = -5, -1$ instantly.

Q2657. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2658. **Middle-Term Splitting** is ideal because $c = 15$ easily factors into $(5) \times (3)$ whose sum is $b = 8$. Factoring yields $(x + 5)(x + 3) = 0$, giving $x = -5, -3$ instantly.

Q2659. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2660. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(6) \times (5)$ whose sum is $b = 11$. Factoring yields $(x + 6)(x + 5) = 0$, giving $x = -6, -5$ instantly.

Q2661. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2662. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(1) \times (4)$ whose sum is $b = 5$. Factoring yields $(x + 1)(x + 4) = 0$, giving $x = -1, -4$ instantly.

Q2663. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2664. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(3) \times (4)$ whose sum is $b = 7$. Factoring yields $(x + 3)(x + 4) = 0$, giving $x = -3, -4$ instantly.

Q2665. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2666. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(3) \times (4)$ whose sum is $b = 7$. Factoring yields $(x + 3)(x + 4) = 0$, giving $x = -3, -4$ instantly.

Q2667. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2668. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(3) \times (2)$ whose sum is $b = 5$. Factoring yields $(x + 3)(x + 2) = 0$, giving $x = -3, -2$ instantly.

Q2669. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2670. **Middle-Term Splitting** is ideal because $c = 15$ easily factors into $(5) \times (3)$ whose sum is $b = 8$. Factoring yields $(x + 5)(x + 3) = 0$, giving $x = -5, -3$ instantly.

Q2671. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2672. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(6) \times (5)$ whose sum is $b = 11$. Factoring yields $(x + 6)(x + 5) = 0$, giving $x = -6, -5$ instantly.

Q2673. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2674. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(4) \times (5)$ whose sum is $b = 9$. Factoring yields $(x + 4)(x + 5) = 0$, giving $x = -4, -5$ instantly.

Q2675. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2676. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2677. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2678. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(4) \times (3)$ whose sum is $b = 7$. Factoring yields $(x + 4)(x + 3) = 0$, giving $x = -4, -3$ instantly.

Q2679. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2680. **Middle-Term Splitting** is ideal because $c = 10$ easily factors into $(2) \times (5)$ whose sum is $b = 7$. Factoring yields $(x + 2)(x + 5) = 0$, giving $x = -2, -5$ instantly.

Q2681. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2682. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(5) \times (1)$ whose sum is $b = 6$. Factoring yields $(x + 5)(x + 1) = 0$, giving $x = -5, -1$ instantly.

Q2683. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2684. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(3) \times (4)$ whose sum is $b = 7$. Factoring yields $(x + 3)(x + 4) = 0$, giving $x = -3, -4$ instantly.

Q2685. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2686. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(2) \times (1)$ whose sum is $b = 3$. Factoring yields $(x + 2)(x + 1) = 0$, giving $x = -2, -1$ instantly.

Q2687. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2688. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(4) \times (6)$ whose sum is $b = 10$. Factoring yields $(x + 4)(x + 6) = 0$, giving $x = -4, -6$ instantly.

Q2689. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2690. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2691. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2692. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(5) \times (4)$ whose sum is $b = 9$. Factoring yields $(x + 5)(x + 4) = 0$, giving $x = -5, -4$ instantly.

Q2693. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2694. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(4) \times (6)$ whose sum is $b = 10$. Factoring yields $(x + 4)(x + 6) = 0$, giving $x = -4, -6$ instantly.

Q2695. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2696. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(3) \times (4)$ whose sum is $b = 7$. Factoring yields $(x + 3)(x + 4) = 0$, giving $x = -3, -4$ instantly.

Q2697. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2698. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2699. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2700. **Middle-Term Splitting** is ideal because $c = 5$ easily factors into $(1) \times (5)$ whose sum is $b = 6$. Factoring yields $(x + 1)(x + 5) = 0$, giving $x = -1, -5$ instantly.

Q2701. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2702. **Middle-Term Splitting** is ideal because $c = 20$ easily factors into $(5) \times (4)$ whose sum is $b = 9$. Factoring yields $(x + 5)(x + 4) = 0$, giving $x = -5, -4$ instantly.

Q2703. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2704. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(6) \times (3)$ whose sum is $b = 9$. Factoring yields $(x + 6)(x + 3) = 0$, giving $x = -6, -3$ instantly.

Q2705. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2706. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(6) \times (3)$ whose sum is $b = 9$. Factoring yields $(x + 6)(x + 3) = 0$, giving $x = -6, -3$ instantly.

Q2707. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2708. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2709. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2710. **Middle-Term Splitting** is ideal because $c = 3$ easily factors into $(3) \times (1)$ whose sum is $b = 4$. Factoring yields $(x + 3)(x + 1) = 0$, giving $x = -3, -1$ instantly.

Q2711. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2712. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2713. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2714. **Middle-Term Splitting** is ideal because $c = 25$ easily factors into $(5) \times (5)$ whose sum is $b = 10$. Factoring yields $(x + 5)(x + 5) = 0$, giving $x = -5, -5$ instantly.

Q2715. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2716. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2717. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2718. **Middle-Term Splitting** is ideal because $c = 1$ easily factors into $(1) \times (1)$ whose sum is $b = 2$. Factoring yields $(x + 1)(x + 1) = 0$, giving $x = -1, -1$ instantly.

Q2719. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2720. **Middle-Term Splitting** is ideal because $c = 4$ easily factors into $(4) \times (1)$ whose sum is $b = 5$. Factoring yields $(x + 4)(x + 1) = 0$, giving $x = -4, -1$ instantly.

Q2721. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2722. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2723. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2724. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(2) \times (3)$ whose sum is $b = 5$. Factoring yields $(x + 2)(x + 3) = 0$, giving $x = -2, -3$ instantly.

Q2725. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2726. **Middle-Term Splitting** is ideal because $c = 24$ easily factors into $(6) \times (4)$ whose sum is $b = 10$. Factoring yields $(x + 6)(x + 4) = 0$, giving $x = -6, -4$ instantly.

Q2727. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2728. **Middle-Term Splitting** is ideal because $c = 18$ easily factors into $(3) \times (6)$ whose sum is $b = 9$. Factoring yields $(x + 3)(x + 6) = 0$, giving $x = -3, -6$ instantly.

Q2729. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2730. **Middle-Term Splitting** is ideal because $c = 16$ easily factors into $(4) \times (4)$ whose sum is $b = 8$. Factoring yields $(x + 4)(x + 4) = 0$, giving $x = -4, -4$ instantly.

Q2731. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2732. **Middle-Term Splitting** is ideal because $c = 15$ easily factors into $(5) \times (3)$ whose sum is $b = 8$. Factoring yields $(x + 5)(x + 3) = 0$, giving $x = -5, -3$ instantly.

Q2733. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{5})(3\sqrt{5}) = 49 - 12(5) = -11$. Since $D > 0$, roots are real.

Q2734. **Middle-Term Splitting** is ideal because $c = 8$ easily factors into $(4) \times (2)$ whose sum is $b = 6$. Factoring yields $(x + 4)(x + 2) = 0$, giving $x = -4, -2$ instantly.

Q2735. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2736. **Middle-Term Splitting** is ideal because $c = 12$ easily factors into $(6) \times (2)$ whose sum is $b = 8$. Factoring yields $(x + 6)(x + 2) = 0$, giving $x = -6, -2$ instantly.

Q2737. Both work, but **Quadratic Formula** ($x =$

$\frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2738. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(1) \times (2)$ whose sum is $b = 3$. Factoring yields $(x + 1)(x + 2) = 0$, giving $x = -1, -2$ instantly.

Q2739. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2740. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(2) \times (3)$ whose sum is $b = 5$. Factoring yields $(x + 2)(x + 3) = 0$, giving $x = -2, -3$ instantly.

Q2741. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2742. **Middle-Term Splitting** is ideal because $c = 9$ easily factors into $(3) \times (3)$ whose sum is $b = 6$. Factoring yields $(x + 3)(x + 3) = 0$, giving $x = -3, -3$ instantly.

Q2743. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2744. **Middle-Term Splitting** is ideal because $c = 25$ easily factors into $(5) \times (5)$ whose sum is $b = 10$. Factoring yields $(x + 5)(x + 5) = 0$, giving $x = -5, -5$ instantly.

Q2745. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2746. **Middle-Term Splitting** is ideal because $c = 30$ easily factors into $(5) \times (6)$ whose sum is $b = 11$. Factoring yields $(x + 5)(x + 6) = 0$, giving $x = -5, -6$ instantly.

Q2747. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{2})(3\sqrt{2}) = 49 - 12(2) = 25$. Since $D > 0$, roots are real.

Q2748. **Middle-Term Splitting** is ideal because $c = 2$ easily factors into $(2) \times (1)$ whose sum is $b = 3$. Factoring yields $(x + 2)(x + 1) = 0$, giving $x = -2, -1$ instantly.

Q2749. Both work, but **Quadratic Formula** ($x = \frac{-b\pm\sqrt{D}}{2a}$) is less prone to error with radical coefficients. Here $D = 49 - 4(\sqrt{3})(3\sqrt{3}) = 49 - 12(3) = 13$. Since $D > 0$, roots are real.

Q2750. **Middle-Term Splitting** is ideal because $c = 6$ easily factors into $(2) \times (3)$ whose sum is $b = 5$. Factoring yields $(x + 2)(x + 3) = 0$, giving $x = -2, -3$ instantly.

Q2751. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2752. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2753. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2754. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2755. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2756. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2757. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2758. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2759. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2760. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2761. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2762. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2763. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2764. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2765. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2766. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2767. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2768. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2769. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2770. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2771. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solu-

tions** (Coincident lines).

Q2772. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2773. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2774. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2775. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2776. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2777. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2778. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2779. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2780. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2781. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2782. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2783. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2784. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2785. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2786. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2787. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2788. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2789. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2790. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2791. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2792. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2793. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2794. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2795. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2796. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2797. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2798. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2799. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2800. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2801. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2802. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2803. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2804. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2805. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2806. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2807. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2808. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2809. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2810. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2811. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2812. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2813. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2814. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2815. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2816. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2817. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2818. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2819. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2820. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2821. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2822. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2823. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2824. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2825. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2826. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2827. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2828. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2829. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2830. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2831. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2832. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2833. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2834. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2835. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2836. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2837. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2838. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2839. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2840. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2841. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2842. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2843. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2844. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2845. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2846. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2847. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2848. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2849. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2850. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2851. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2852. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2853. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2854. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2855. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2856. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2857. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2858. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2859. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2860. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2861. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2862. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2863. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2864. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2865. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2866. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2867. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2868. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2869. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2870. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2871. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2872. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2873. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2874. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2875. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2876. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2877. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2878. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2879. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2880. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2881. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2882. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2883. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2884. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2885. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2886. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2887. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2888. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2889. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2890. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2891. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2892. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2893. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2894. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2895. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2896. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2897. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2898. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2899. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2900. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2901. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2902. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2903. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2904. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2905. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2906. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2907. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2908. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2909. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2910. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2911. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2912. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2913. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2914. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2915. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2916. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2917. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2918. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2919. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2920. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2921. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2922. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2923. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2924. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2925. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2926. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2927. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2928. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2929. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2930. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2931. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2932. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2933. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2934. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2935. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2936. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2937. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2938. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2939. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2940. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2941. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2942. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2943. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2944. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2945. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2946. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2947. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2948. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2949. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2950. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2951. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2952. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2953. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2954. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2955. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2956. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2957. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2958. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2959. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2960. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2961. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2962. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2963. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2964. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2965. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2966. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2967. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2968. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2969. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2970. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2971. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2972. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2973. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2974. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2975. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2976. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2977. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2978. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2979. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2980. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2981. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2982. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2983. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2984. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2985. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2986. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2987. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

tions** (Coincident lines).

Q2988. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2989. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2990. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2991. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2992. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2993. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2994. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2995. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2996. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2997. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2998. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q2999. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

Q3000. Compare ratios: $\frac{a_1}{a_2} = \frac{1}{4}$, $\frac{b_1}{b_2} = \frac{1}{4}$, $\frac{c_1}{c_2} = \frac{1}{4}$. Since $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$, the system has **Infinitely Many Solutions** (Coincident lines).

MODULE 4: DENOMINATOR RULES AND FRACTIONAL EQUATIONS (Q3001 – Q4000) — Answer Key

Q3001. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 8 = 24 \times 2 = 48$. In (B), 2 only divides x, so you must transpose +8 FIRST before moving 2: $\frac{x}{2} = 24 - 8 = 16 \implies x = 32$.

Q3002. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 11 \times 5 = 55$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 11 - 8 = 3 \implies x = 15$.

Q3003. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 8 = 29 \times 2 = 58$. In (B), 2 only divides x, so you must transpose +8 FIRST before moving 2: $\frac{x}{2} = 29 - 8 = 21 \implies x = 42$.

Q3004. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 29 \times 5 = 145$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 29 - 6 = 23 \implies x = 115$.

Q3005. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 3 = 22 \times 6 = 132$. In (B), 6 only divides x, so you must transpose +3 FIRST before moving 6: $\frac{x}{6} = 22 - 3 = 19 \implies x = 114$.

Q3006. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 26 \times 2 = 52$. In (B), 2 only divides x, so you must transpose +2 FIRST before moving 2: $\frac{x}{2} = 26 - 2 = 24 \implies x = 48$.

Q3007. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 9 = 20 \times 5 = 100$. In (B), 5 only divides x, so you must transpose +9 FIRST before moving 5: $\frac{x}{5} = 20 - 9 = 11 \implies x = 55$.

Q3008. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 6 = 19 \times 3 = 57$. In (B), 3 only divides x, so you must transpose +6 FIRST before moving 3: $\frac{x}{3} = 19 - 6 = 13 \implies x = 39$.

Q3009. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 16 \times 4 = 64$. In (B), 4 only divides x, so you must transpose +9 FIRST before moving 4: $\frac{x}{4} = 16 - 9 = 7 \implies x = 28$.

Q3010. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 11 \times 5 = 55$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 11 - 2 = 9 \implies x = 45$.

Q3011. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 8 = 11 \times 6 = 66$. In (B), 6 only divides x, so you must transpose +8 FIRST before moving 6: $\frac{x}{6} = 11 - 8 = 3 \implies x = 18$.

Q3012. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 2 = 28 \times 3 = 84$. In (B), 3 only divides x, so you must transpose +2 FIRST before moving 3: $\frac{x}{3} = 28 - 2 = 26 \implies x = 78$.

Q3013. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 24 \times 3 = 72$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 24 - 5 = 19 \implies x = 57$.

Q3014. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 2 = 22 \times 3 = 66$. In (B), 3 only divides x, so you must transpose +2 FIRST before moving 3: $\frac{x}{3} = 22 - 2 = 20 \implies x = 60$.

Q3015. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 5 = 30 \times 5 = 150$. In (B), 5 only divides x, so you must transpose +5 FIRST before moving 5: $\frac{x}{5} = 30 - 5 = 25 \implies x = 125$.

Q3016. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 17 \times 5 = 85$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 17 - 6 = 11 \implies x = 55$.

Q3017. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 16 \times 5 = 80$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 16 - 6 = 10 \implies x = 50$.

Q3018. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 9 = 26 \times 7 = 182$. In (B), 7 only divides x, so you must transpose +9 FIRST before moving 7: $\frac{x}{7} = 26 - 9 = 17 \implies x = 119$.

Q3019. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 18 \times 4 = 72$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 18 - 5 = 13 \implies x = 52$.

Q3020. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 6 = 30 \times 7 = 210$. In (B), 7 only divides x, so you must transpose +6 FIRST before moving 7: $\frac{x}{7} = 30 - 6 = 24 \implies x = 168$.

Q3021. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 5 = 22 \times 7 = 154$. In (B), 7 only divides x, so you must transpose +5 FIRST before moving 7: $\frac{x}{7} = 22 - 5 = 17 \implies x = 119$.

Q3022. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 11 \times 3 = 33$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 11 - 5 = 6 \implies x = 18$.

Q3023. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 19 - 8 = 11 \implies x = 55$.

Q3024. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 12 \times 3 = 36$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 12 - 9 = 3 \implies x = 9$.

Q3025. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 7 = 12 \times 7 = 84$. In (B), 7 only divides x, so you must transpose +7 FIRST before moving 7: $\frac{x}{7} = 12 - 7 = 5 \implies x = 35$.

Q3026. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 27 \times 2 = 54$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 27 - 4 = 23 \implies x = 46$.

Q3027. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 6 = 17 \times 4 = 68$. In (B), 4 only divides x, so you must transpose +6 FIRST before moving 4: $\frac{x}{4} = 17 - 6 = 11 \implies x = 44$.

Q3028. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 22 \times 3 = 66$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 22 - 9 = 13 \implies x = 39$.

Q3029. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 5 = 29 \times 5 = 145$. In (B), 5 only divides x, so you must transpose +5 FIRST before moving 5: $\frac{x}{5} = 29 - 5 = 24 \implies x = 120$.

Q3030. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 19 - 2 = 17 \implies x = 85$.

Q3031. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 19 \times 4 = 76$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 19 - 5 = 14 \implies x = 56$.

Q3032. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 19 - 8 = 11 \implies x = 55$.

Q3033. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 25 \times 2 = 50$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 25 - 9 = 16 \implies x = 32$.

Q3034. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 8 = 28 \times 3 = 84$. In (B), 3 only divides x, so you must transpose +8 FIRST before moving 3: $\frac{x}{3} = 28 - 8 = 20 \implies x = 60$.

Q3035. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 23 \times 2 = 46$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 23 - 6 = 17 \implies x = 34$.

Q3036. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 2 = 28 \times 7 = 196$. In (B), 7 only divides x, so you must transpose +2 FIRST before moving 7: $\frac{x}{7} = 28 - 2 = 26 \implies x = 182$.

Q3037. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 9 = 16 \times 5 = 80$. In (B), 5 only divides x, so you must transpose +9 FIRST before moving 5: $\frac{x}{5} = 16 - 9 = 7 \implies x = 35$.

Q3038. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 13 \times 6 = 78$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 13 - 6 = 7 \implies x = 42$.

Q3039. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 13 \times 4 = 52$. In (B), 4 only divides x, so you must transpose +9 FIRST before moving 4: $\frac{x}{4} = 13 - 9 = 4 \implies x = 16$.

Q3040. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 4 = 25 \times 7 = 175$. In (B), 7 only divides x, so you must transpose +4 FIRST before moving 7: $\frac{x}{7} = 25 - 4 = 21 \implies x = 147$.

Q3041. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 27 \times 4 = 108$. In (B), 4 only divides x, so you must transpose +9 FIRST before moving 4: $\frac{x}{4} = 27 - 9 = 18 \implies x = 72$.

Q3042. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 16 \times 5 = 80$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 16 - 6 = 10 \implies x = 50$.

Q3043. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 24 \times 3 = 72$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 24 - 9 = 15 \implies x = 45$.

Q3044. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 16 \times 6 = 96$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 16 - 2 = 14 \implies x = 84$.

Q3045. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 21 \times 3 = 63$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 21 - 9 = 12 \implies x = 36$.

Q3046. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 27 \times 2 = 54$. In (B), 2 only divides x, so you must transpose +2 FIRST before moving 2: $\frac{x}{2} = 27 - 2 = 25 \implies x = 50$.

Q3047. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 14 \times 6 = 84$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 14 - 2 = 12 \implies x = 72$.

Q3048. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 7 = 29 \times 2 = 58$. In (B), 2 only divides x, so you must transpose +7 FIRST before moving 2: $\frac{x}{2} = 29 - 7 = 22 \implies x = 44$.

Q3049. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 2 = 12 \times 7 = 84$. In (B), 7 only divides x, so you must transpose +2 FIRST before moving 7: $\frac{x}{7} = 12 - 2 = 10 \implies x = 70$.

Q3050. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 3 = 12 \times 6 = 72$. In (B), 6 only divides x, so you must transpose +3 FIRST before moving 6: $\frac{x}{6} = 12 - 3 = 9 \implies x = 54$.

Q3051. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 14 \times 5 = 70$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 14 - 3 = 11 \implies x = 55$.

Q3052. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 2 = 14 \times 7 = 98$. In (B), 7 only divides x, so you must transpose +2 FIRST before moving 7: $\frac{x}{7} = 14 - 2 = 12 \implies x = 84$.

Q3053. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 16 \times 6 = 96$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 16 - 4 = 12 \implies x = 72$.

Q3054. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 24 - 4 = 20 \implies x = 120$.

Q3055. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 17 \times 4 = 68$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 17 - 2 = 15 \implies x = 60$.

Q3056. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 5 = 27 \times 6 = 162$. In (B), 6 only divides x, so you must transpose +5 FIRST before moving 6: $\frac{x}{6} = 27 - 5 = 22 \implies x = 132$.

Q3057. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 23 \times 4 = 92$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 23 - 2 = 21 \implies x = 84$.

Q3058. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 19 \times 4 = 76$. In (B),

4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 19 - 5 = 14 \implies x = 56$.

Q3059. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 15 \times 2 = 30$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 15 - 6 = 9 \implies x = 18$.

Q3060. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 25 \times 6 = 150$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 25 - 4 = 21 \implies x = 126$.

Q3061. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 3 = 24 \times 4 = 96$. In (B), 4 only divides x, so you must transpose +3 FIRST before moving 4: $\frac{x}{4} = 24 - 3 = 21 \implies x = 84$.

Q3062. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 6 = 11 \times 3 = 33$. In (B), 3 only divides x, so you must transpose +6 FIRST before moving 3: $\frac{x}{3} = 11 - 6 = 5 \implies x = 15$.

Q3063. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 24 \times 2 = 48$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 24 - 9 = 15 \implies x = 30$.

Q3064. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 7 = 28 \times 2 = 56$. In (B), 2 only divides x, so you must transpose +7 FIRST before moving 2: $\frac{x}{2} = 28 - 7 = 21 \implies x = 42$.

Q3065. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 3 = 11 \times 3 = 33$. In (B), 3 only divides x, so you must transpose +3 FIRST before moving 3: $\frac{x}{3} = 11 - 3 = 8 \implies x = 24$.

Q3066. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 3 = 30 \times 2 = 60$. In (B), 2 only divides x, so you must transpose +3 FIRST before moving 2: $\frac{x}{2} = 30 - 3 = 27 \implies x = 54$.

Q3067. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 6 = 22 \times 3 = 66$. In (B), 3 only divides x, so you must transpose +6 FIRST before moving 3: $\frac{x}{3} = 22 - 6 = 16 \implies x = 48$.

Q3068. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 6 = 12 \times 7 = 84$. In (B), 7 only divides x, so you must transpose +6 FIRST before moving 7: $\frac{x}{7} = 12 - 6 = 6 \implies x = 42$.

Q3069. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 13 \times 5 = 65$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 13 - 3 = 10 \implies x = 50$.

Q3070. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 17 \times 2 = 34$. In (B), 2 only divides x, so you must transpose +5 FIRST before moving 2: $\frac{x}{2} = 17 - 5 = 12 \implies x = 24$.

Q3071. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 4 = 20 \times 3 = 60$. In (B), 3 only divides x, so you must transpose +4 FIRST before moving 3: $\frac{x}{3} = 20 - 4 = 16 \implies x = 48$.

Q3072. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 10 \times 2 = 20$. In (B), 2 only divides x, so you must transpose +5 FIRST before moving 2: $\frac{x}{2} = 10 - 5 = 5 \implies x = 10$.

Q3073. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 7 = 11 \times 2 = 22$. In (B), 2 only divides x, so you must transpose +7 FIRST before moving 2: $\frac{x}{2} = 11 - 7 = 4 \implies x = 8$.

Q3074. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 21 \times 2 = 42$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 21 - 9 = 12 \implies x = 24$.

Q3075. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 22 \times 2 = 44$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 22 - 9 = 13 \implies x = 26$.

Q3076. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 3 = 17 \times 2 = 34$. In (B), 2 only divides x, so you must transpose +3 FIRST before moving 2: $\frac{x}{2} = 17 - 3 = 14 \implies x = 28$.

Q3077. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 12 \times 6 = 72$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 12 - 6 = 6 \implies x = 36$.

Q3078. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 17 \times 6 = 102$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 17 - 6 = 11 \implies x = 66$.

Q3079. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 8 = 25 \times 2 = 50$. In (B), 2 only divides x, so you must transpose +8 FIRST before moving 2: $\frac{x}{2} = 25 - 8 = 17 \implies x = 34$.

Q3080. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 8 = 24 \times 4 = 96$. In (B), 4 only divides x, so you must transpose +8 FIRST before moving 4: $\frac{x}{4} = 24 - 8 = 16 \implies x = 64$.

Q3081. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 8 = 27 \times 6 = 162$. In (B), 6 only divides x, so you must transpose +8 FIRST before moving 6: $\frac{x}{6} = 27 - 8 = 19 \implies x = 114$.

Q3082. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 4 = 14 \times 3 = 42$. In (B), 3 only divides x, so you must transpose +4 FIRST before moving 3: $\frac{x}{3} = 14 - 4 = 10 \implies x = 30$.

Q3083. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 27 \times 2 = 54$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 27 - 4 = 23 \implies x = 46$.

Q3084. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 14 \times 2 = 28$. In (B), 2 only divides x, so you must transpose +5 FIRST before moving 2: $\frac{x}{2} = 14 - 5 = 9 \implies x = 18$.

Q3085. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 17 \times 5 = 85$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 17 - 3 = 14 \implies x = 70$.

Q3086. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 3 = 21 \times 7 = 147$. In (B), 7 only divides x, so you must transpose +3 FIRST before moving 7: $\frac{x}{7} = 21 - 3 = 18 \implies x = 126$.

Q3087. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 11 \times 5 = 55$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 11 - 3 = 8 \implies x = 40$.

Q3088. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 20 \times 2 = 40$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 20 - 6 = 14 \implies x = 28$.

Q3089. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 9 = 29 \times 7 = 203$. In (B), 7 only divides x, so you must transpose +9 FIRST before moving 7: $\frac{x}{7} = 29 - 9 = 20 \implies x = 140$.

Q3090. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 3 = 28 \times 7 = 196$. In (B), 7 only divides x, so you must transpose +3 FIRST before moving 7: $\frac{x}{7} = 28 - 3 = 25 \implies x = 175$.

Q3091. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 5 = 17 \times 7 = 119$. In (B), 7 only divides x, so you must transpose +5 FIRST before moving 7: $\frac{x}{7} = 17 - 5 = 12 \implies x = 84$.

Q3092. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 20 \times 2 = 40$. In (B), 2 only divides x, so you must transpose +2 FIRST before moving 2: $\frac{x}{2} = 20 - 2 = 18 \implies x = 36$.

Q3093. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 12 \times 2 = 24$. In (B), 2 only divides x, so you must transpose +2 FIRST before moving 2: $\frac{x}{2} = 12 - 2 = 10 \implies x = 20$.

Q3094. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 25 \times 5 = 125$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 25 - 8 = 17 \implies x = 85$.

Q3095. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 17 \times 6 = 102$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 17 - 6 = 11 \implies x = 66$.

Q3096. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 7 = 10 \times 7 = 70$. In (B), 7 only divides x, so you must transpose +7 FIRST before moving 7: $\frac{x}{7} = 10 - 7 = 3 \implies x = 21$.

Q3097. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 11 \times 6 = 66$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 11 - 4 = 7 \implies x = 42$.

Q3098. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 28 \times 6 = 168$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 28 - 6 = 22 \implies x = 132$.

Q3099. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 19 \times 2 = 38$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 19 - 9 = 10 \implies x = 20$.

Q3100. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 27 \times 6 = 162$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 27 - 6 = 21 \implies x = 126$.

Q3101. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 6 = 24 \times 7 = 168$. In (B), 7 only divides x, so you must transpose +6 FIRST before moving 7: $\frac{x}{7} = 24 - 6 = 18 \implies x = 126$.

Q3102. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 7 = 30 \times 2 = 60$. In (B), 2 only divides x, so you must transpose +7 FIRST before moving 2: $\frac{x}{2} = 30 - 7 = 23 \implies x = 46$.

Q3103. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 3 = 10 \times 6 = 60$. In (B), 6 only divides x, so you must transpose +3 FIRST before moving 6: $\frac{x}{6} = 10 - 3 = 7 \implies x = 42$.

Q3104. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 15 \times 6 = 90$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 15 - 2 = 13 \implies x = 78$.

Q3105. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 8 = 26 \times 3 = 78$. In (B), 3 only divides x, so you must transpose +8 FIRST before moving 3: $\frac{x}{3} = 26 - 8 = 18 \implies x = 54$.

Q3106. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 17 \times 6 = 102$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 17 - 4 = 13 \implies x = 78$.

Q3107. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 4 = 29 \times 7 = 203$. In (B), 7 only divides x, so you must transpose +4 FIRST before moving 7: $\frac{x}{7} = 29 - 4 = 25 \implies x = 175$.

Q3108. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 14 \times 5 = 70$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 14 - 3 = 11 \implies x = 55$.

Q3109. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 18 \times 6 = 108$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 18 - 2 = 16 \implies x = 96$.

Q3110. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 27 \times 4 = 108$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 27 - 2 = 25 \implies x = 100$.

Q3111. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 18 \times 2 = 36$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 18 - 4 = 14 \implies x = 28$.

Q3112. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 4 = 12 \times 3 = 36$. In (B), 3 only divides x, so you must transpose +4 FIRST before moving 3: $\frac{x}{3} = 12 - 4 = 8 \implies x = 24$.

Q3113. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 25 \times 5 = 125$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 25 - 2 = 23 \implies x = 115$.

Q3114. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 8 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +8 FIRST before moving 6: $\frac{x}{6} = 24 - 8 = 16 \implies x = 96$.

Q3115. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 28 \times 3 = 84$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 28 - 9 = 19 \implies x = 57$.

Q3116. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 2 = 13 \times 3 = 39$. In (B), 3 only divides x, so you must transpose +2 FIRST before moving 3: $\frac{x}{3} = 13 - 2 = 11 \implies x = 33$.

Q3117. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 23 \times 5 = 115$. In (B),

5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 23 - 3 = 20 \implies x = 100$.

Q3118. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 13 \times 6 = 78$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 13 - 6 = 7 \implies x = 42$.

Q3119. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 26 \times 2 = 52$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 26 - 4 = 22 \implies x = 44$.

Q3120. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 7 = 12 \times 5 = 60$. In (B), 5 only divides x, so you must transpose +7 FIRST before moving 5: $\frac{x}{5} = 12 - 7 = 5 \implies x = 25$.

Q3121. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 26 \times 5 = 130$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 26 - 6 = 20 \implies x = 100$.

Q3122. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 23 \times 2 = 46$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 23 - 4 = 19 \implies x = 38$.

Q3123. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 25 \times 4 = 100$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 25 - 2 = 23 \implies x = 92$.

Q3124. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 9 = 29 \times 7 = 203$. In (B), 7 only divides x, so you must transpose +9 FIRST before moving 7: $\frac{x}{7} = 29 - 9 = 20 \implies x = 140$.

Q3125. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 8 = 23 \times 7 = 161$. In (B), 7 only divides x, so you must transpose +8 FIRST before moving 7: $\frac{x}{7} = 23 - 8 = 15 \implies x = 105$.

Q3126. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 16 \times 3 = 48$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 16 - 9 = 7 \implies x = 21$.

Q3127. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 9 = 29 \times 6 = 174$. In (B), 6 only divides x, so you must transpose +9 FIRST before moving 6: $\frac{x}{6} = 29 - 9 = 20 \implies x = 120$.

Q3128. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 24 \times 2 = 48$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 24 - 4 = 20 \implies x = 40$.

Q3129. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 2 = 22 \times 7 = 154$. In (B), 7 only divides x, so you must transpose +2 FIRST before moving 7: $\frac{x}{7} = 22 - 2 = 20 \implies x = 140$.

Q3130. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 11 \times 2 = 22$. In (B), 2 only divides x, so you must transpose +5 FIRST before moving 2: $\frac{x}{2} = 11 - 5 = 6 \implies x = 12$.

Q3131. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 14 \times 6 = 84$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 14 - 2 = 12 \implies x = 72$.

Q3132. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 9 = 17 \times 6 = 102$. In (B), 6 only divides x, so you must transpose +9 FIRST before moving 6: $\frac{x}{6} = 17 - 9 = 8 \implies x = 48$.

Q3133. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 2 = 21 \times 7 = 147$. In (B), 7 only divides x, so you must transpose +2 FIRST before moving 7: $\frac{x}{7} = 21 - 2 = 19 \implies x = 133$.

Q3134. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 28 \times 4 = 112$. In (B), 4 only divides x, so you must transpose +9 FIRST before moving 4: $\frac{x}{4} = 28 - 9 = 19 \implies x = 76$.

Q3135. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 4 = 21 \times 5 = 105$. In (B), 5 only divides x, so you must transpose +4 FIRST before moving 5: $\frac{x}{5} = 21 - 4 = 17 \implies x = 85$.

Q3136. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 11 \times 5 = 55$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 11 - 6 = 5 \implies x = 25$.

Q3137. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 8 = 17 \times 7 = 119$. In (B), 7 only divides x, so you must transpose +8 FIRST before moving 7: $\frac{x}{7} = 17 - 8 = 9 \implies x = 63$.

Q3138. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 28 \times 5 = 140$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 28 - 6 = 22 \implies x = 110$.

Q3139. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 6 = 22 \times 4 = 88$. In (B), 4 only divides x, so you must transpose +6 FIRST before moving 4: $\frac{x}{4} = 22 - 6 = 16 \implies x = 64$.

Q3140. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 5 = 20 \times 6 = 120$. In (B), 6 only divides x, so you must transpose +5 FIRST before moving 6: $\frac{x}{6} = 20 - 5 = 15 \implies x = 90$.

Q3141. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 7 = 19 \times 3 = 57$. In (B), 3 only divides x, so you must transpose +7 FIRST before moving 3: $\frac{x}{3} = 19 - 7 = 12 \implies x = 36$.

Q3142. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 3 = 21 \times 3 = 63$. In (B), 3 only divides x, so you must transpose +3 FIRST before moving 3: $\frac{x}{3} = 21 - 3 = 18 \implies x = 54$.

Q3143. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 19 - 3 = 16 \implies x = 80$.

Q3144. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 5 = 21 \times 7 = 147$. In (B), 7 only divides x, so you must transpose +5 FIRST before moving 7: $\frac{x}{7} = 21 - 5 = 16 \implies x = 112$.

Q3145. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 4 = 13 \times 4 = 52$. In (B), 4 only divides x, so you must transpose +4 FIRST before moving 4: $\frac{x}{4} = 13 - 4 = 9 \implies x = 36$.

Q3146. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 19 - 2 = 17 \implies x = 85$.

Q3147. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 8 = 10 \times 7 = 70$. In (B), 7 only divides x, so you must transpose +8 FIRST before moving 7: $\frac{x}{7} = 10 - 8 = 2 \implies x = 14$.

Q3148. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 4 = 23 \times 5 = 115$. In (B), 5 only divides x, so you must transpose +4 FIRST before moving 5: $\frac{x}{5} = 23 - 4 = 19 \implies x = 95$.

Q3149. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 7 = 19 \times 3 = 57$. In (B), 3 only divides x, so you must transpose +7 FIRST before moving 3: $\frac{x}{3} = 19 - 7 = 12 \implies x = 36$.

Q3150. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 8 = 16 \times 6 = 96$. In (B), 6 only divides x, so you must transpose +8 FIRST before moving 6: $\frac{x}{6} = 16 - 8 = 8 \implies x = 48$.

Q3151. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 29 \times 3 = 87$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 29 - 5 = 24 \implies x = 72$.

Q3152. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 24 - 2 = 22 \implies x = 132$.

Q3153. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 6 = 20 \times 3 = 60$. In (B), 3 only divides x, so you must transpose +6 FIRST before moving 3: $\frac{x}{3} = 20 - 6 = 14 \implies x = 42$.

Q3154. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 27 \times 4 = 108$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 27 - 5 = 22 \implies x = 88$.

Q3155. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 3 = 18 \times 2 = 36$. In (B), 2 only divides x, so you must transpose +3 FIRST before moving 2: $\frac{x}{2} = 18 - 3 = 15 \implies x = 30$.

Q3156. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 4 = 13 \times 4 = 52$. In (B), 4 only divides x, so you must transpose +4 FIRST before moving 4: $\frac{x}{4} = 13 - 4 = 9 \implies x = 36$.

Q3157. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 21 \times 5 = 105$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 21 - 8 = 13 \implies x = 65$.

Q3158. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 17 \times 2 = 34$. In (B), 2 only divides x, so you must transpose +5 FIRST before moving 2: $\frac{x}{2} = 17 - 5 = 12 \implies x = 24$.

Q3159. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 4 = 25 \times 5 = 125$. In (B), 5 only divides x, so you must transpose +4 FIRST before moving 5: $\frac{x}{5} = 25 - 4 = 21 \implies x = 105$.

Q3160. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 24 \times 5 = 120$. In (B), 5 only divides x, so you must transpose +3 FIRST before moving 5: $\frac{x}{5} = 24 - 3 = 21 \implies x = 105$.

Q3161. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 13 \times 4 = 52$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 13 - 5 = 8 \implies x = 32$.

Q3162. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 3 = 25 \times 4 = 100$. In (B), 4 only divides x, so you must transpose +3 FIRST before moving 4: $\frac{x}{4} = 25 - 3 = 22 \implies x = 88$.

Q3163. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 6 = 13 \times 5 = 65$. In (B), 5 only divides x, so you must transpose +6 FIRST before moving 5: $\frac{x}{5} = 13 - 6 = 7 \implies x = 35$.

Q3164. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 7 = 27 \times 7 = 189$. In (B), 7 only divides x, so you must transpose +7 FIRST before moving 7: $\frac{x}{7} = 27 - 7 = 20 \implies x = 140$.

Q3165. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 17 \times 4 = 68$. In (B), 4 only divides x, so you must transpose +9 FIRST before moving 4: $\frac{x}{4} = 17 - 9 = 8 \implies x = 32$.

Q3166. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 18 \times 5 = 90$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 18 - 8 = 10 \implies x = 50$.

Q3167. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 20 \times 5 = 100$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 20 - 8 = 12 \implies x = 60$.

Q3168. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 7 = 13 \times 4 = 52$. In (B), 4 only divides x, so you must transpose +7 FIRST before moving 4: $\frac{x}{4} = 13 - 7 = 6 \implies x = 24$.

Q3169. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 4 = 29 \times 4 = 116$. In (B), 4 only divides x, so you must transpose +4 FIRST before moving 4: $\frac{x}{4} = 29 - 4 = 25 \implies x = 100$.

Q3170. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 9 = 14 \times 7 = 98$. In (B), 7 only divides x, so you must transpose +9 FIRST before moving 7: $\frac{x}{7} = 14 - 9 = 5 \implies x = 35$.

Q3171. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 3 = 19 \times 3 = 57$. In (B), 3 only divides x, so you must transpose +3 FIRST before moving 3: $\frac{x}{3} = 19 - 3 = 16 \implies x = 48$.

Q3172. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 23 \times 6 = 138$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 23 - 4 = 19 \implies x = 114$.

Q3173. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 9 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +9 FIRST before moving 6: $\frac{x}{6} = 24 - 9 = 15 \implies x = 90$.

Q3174. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 28 \times 3 = 84$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 28 - 5 = 23 \implies x = 69$.

Q3175. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 25 \times 2 = 50$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 25 - 6 = 19 \implies x = 38$.

Q3176. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 4 = 27 \times 5 = 135$. In (B),

5 only divides x, so you must transpose +4 FIRST before moving 5: $\frac{x}{5} = 27 - 4 = 23 \implies x = 115$.

Q3177. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 8 = 30 \times 7 = 210$. In (B), 7 only divides x, so you must transpose +8 FIRST before moving 7: $\frac{x}{7} = 30 - 8 = 22 \implies x = 154$.

Q3178. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 5 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +5 FIRST before moving 5: $\frac{x}{5} = 19 - 5 = 14 \implies x = 70$.

Q3179. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 24 - 4 = 20 \implies x = 120$.

Q3180. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 14 \times 2 = 28$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 14 - 9 = 5 \implies x = 10$.

Q3181. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 8 = 14 \times 3 = 42$. In (B), 3 only divides x, so you must transpose +8 FIRST before moving 3: $\frac{x}{3} = 14 - 8 = 6 \implies x = 18$.

Q3182. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 16 \times 3 = 48$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 16 - 5 = 11 \implies x = 33$.

Q3183. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 4 = 23 \times 5 = 115$. In (B), 5 only divides x, so you must transpose +4 FIRST before moving 5: $\frac{x}{5} = 23 - 4 = 19 \implies x = 95$.

Q3184. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 23 \times 4 = 92$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 23 - 2 = 21 \implies x = 84$.

Q3185. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 5 = 15 \times 3 = 45$. In (B), 3 only divides x, so you must transpose +5 FIRST before moving 3: $\frac{x}{3} = 15 - 5 = 10 \implies x = 30$.

Q3186. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 21 \times 6 = 126$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 21 - 6 = 15 \implies x = 90$.

Q3187. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 2 = 14 \times 4 = 56$. In (B), 4 only divides x, so you must transpose +2 FIRST before moving 4: $\frac{x}{4} = 14 - 2 = 12 \implies x = 48$.

Q3188. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 8 = 26 \times 4 = 104$. In (B), 4 only divides x, so you must transpose +8 FIRST before moving 4: $\frac{x}{4} = 26 - 8 = 18 \implies x = 72$.

Q3189. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 16 \times 5 = 80$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 16 - 2 = 14 \implies x = 70$.

Q3190. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 28 \times 2 = 56$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 28 - 6 = 22 \implies x = 44$.

Q3191. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 9 = 20 \times 7 = 140$. In (B), 7 only divides x, so you must transpose +9 FIRST before moving 7: $\frac{x}{7} = 20 - 9 = 11 \implies x = 77$.

Q3192. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 20 \times 3 = 60$. In (B), 3 only divides x, so you must transpose +9 FIRST before moving 3: $\frac{x}{3} = 20 - 9 = 11 \implies x = 33$.

Q3193. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 7 = 21 \times 5 = 105$. In (B), 5 only divides x, so you must transpose +7 FIRST before moving 5: $\frac{x}{5} = 21 - 7 = 14 \implies x = 70$.

Q3194. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 7 = 30 \times 6 = 180$. In (B), 6 only divides x, so you must transpose +7 FIRST before moving 6: $\frac{x}{6} = 30 - 7 = 23 \implies x = 138$.

Q3195. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 2 = 22 \times 6 = 132$. In (B), 6 only divides x, so you must transpose +2 FIRST before moving 6: $\frac{x}{6} = 22 - 2 = 20 \implies x = 120$.

Q3196. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 16 \times 5 = 80$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 16 - 2 = 14 \implies x = 70$.

Q3197. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 4 = 22 \times 7 = 154$. In (B), 7 only divides x, so you must transpose +4 FIRST before moving 7: $\frac{x}{7} = 22 - 4 = 18 \implies x = 126$.

Q3198. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 8 = 18 \times 2 = 36$. In (B), 2 only divides x, so you must transpose +8 FIRST before moving 2: $\frac{x}{2} = 18 - 8 = 10 \implies x = 20$.

Q3199. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 10 \times 2 = 20$. In (B), 2 only divides x, so you must transpose +6 FIRST before moving 2: $\frac{x}{2} = 10 - 6 = 4 \implies x = 8$.

Q3200. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 5 = 28 \times 6 = 168$. In (B), 6 only divides x, so you must transpose +5 FIRST before moving 6: $\frac{x}{6} = 28 - 5 = 23 \implies x = 138$.

Q3201. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 26 \times 6 = 156$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 26 - 6 = 20 \implies x = 120$.

Q3202. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 19 \times 5 = 95$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 19 - 8 = 11 \implies x = 55$.

Q3203. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 24 \times 5 = 120$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 24 - 2 = 22 \implies x = 110$.

Q3204. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 6 = 11 \times 6 = 66$. In (B), 6 only divides x, so you must transpose +6 FIRST before moving 6: $\frac{x}{6} = 11 - 6 = 5 \implies x = 30$.

Q3205. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 27 \times 4 = 108$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 27 - 5 = 22 \implies x = 88$.

Q3206. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 5 = 15 \times 6 = 90$. In (B), 6 only divides x, so you must transpose $+5$ FIRST before moving 6: $\frac{x}{6} = 15 - 5 = 10 \implies x = 60$.

Q3207. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 7 = 23 \times 6 = 138$. In (B), 6 only divides x, so you must transpose $+7$ FIRST before moving 6: $\frac{x}{6} = 23 - 7 = 16 \implies x = 96$.

Q3208. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 10 \times 2 = 20$. In (B), 2 only divides x, so you must transpose $+2$ FIRST before moving 2: $\frac{x}{2} = 10 - 2 = 8 \implies x = 16$.

Q3209. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 4 = 28 \times 3 = 84$. In (B), 3 only divides x, so you must transpose $+4$ FIRST before moving 3: $\frac{x}{3} = 28 - 4 = 24 \implies x = 72$.

Q3210. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 23 \times 2 = 46$. In (B), 2 only divides x, so you must transpose $+4$ FIRST before moving 2: $\frac{x}{2} = 23 - 4 = 19 \implies x = 38$.

Q3211. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 12 \times 5 = 60$. In (B), 5 only divides x, so you must transpose $+8$ FIRST before moving 5: $\frac{x}{5} = 12 - 8 = 4 \implies x = 20$.

Q3212. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 7 = 16 \times 6 = 96$. In (B), 6 only divides x, so you must transpose $+7$ FIRST before moving 6: $\frac{x}{6} = 16 - 7 = 9 \implies x = 54$.

Q3213. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 5 = 17 \times 2 = 34$. In (B), 2 only divides x, so you must transpose $+5$ FIRST before moving 2: $\frac{x}{2} = 17 - 5 = 12 \implies x = 24$.

Q3214. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 4 = 22 \times 7 = 154$. In (B), 7 only divides x, so you must transpose $+4$ FIRST before moving 7: $\frac{x}{7} = 22 - 4 = 18 \implies x = 126$.

Q3215. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 5 = 14 \times 7 = 98$. In (B), 7 only divides x, so you must transpose $+5$ FIRST before moving 7: $\frac{x}{7} = 14 - 5 = 9 \implies x = 63$.

Q3216. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 11 \times 2 = 22$. In (B), 2 only divides x, so you must transpose $+6$ FIRST before moving 2: $\frac{x}{2} = 11 - 6 = 5 \implies x = 10$.

Q3217. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 9 = 18 \times 3 = 54$. In (B), 3 only divides x, so you must transpose $+9$ FIRST before moving 3: $\frac{x}{3} = 18 - 9 = 9 \implies x = 27$.

Q3218. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 9 = 30 \times 4 = 120$. In (B), 4 only divides x, so you must transpose $+9$ FIRST before moving 4: $\frac{x}{4} = 30 - 9 = 21 \implies x = 84$.

Q3219. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 9 = 26 \times 6 = 156$. In (B), 6 only divides x, so you must transpose $+9$ FIRST before moving 6: $\frac{x}{6} = 26 - 9 = 17 \implies x = 102$.

Q3220. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 2 = 30 \times 2 = 60$. In (B), 2 only divides x, so you must transpose $+2$ FIRST before moving 2: $\frac{x}{2} = 30 - 2 = 28 \implies x = 56$.

Q3221. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 6 = 12 \times 2 = 24$. In (B), 2 only divides x, so you must transpose $+6$ FIRST before moving 2: $\frac{x}{2} = 12 - 6 = 6 \implies x = 12$.

Q3222. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 9 = 13 \times 6 = 78$. In (B), 6 only divides x, so you must transpose $+9$ FIRST before moving 6: $\frac{x}{6} = 13 - 9 = 4 \implies x = 24$.

Q3223. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 3 = 22 \times 5 = 110$. In (B), 5 only divides x, so you must transpose $+3$ FIRST before moving 5: $\frac{x}{5} = 22 - 3 = 19 \implies x = 95$.

Q3224. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 18 \times 2 = 36$. In (B), 2 only divides x, so you must transpose $+4$ FIRST before moving 2: $\frac{x}{2} = 18 - 4 = 14 \implies x = 28$.

Q3225. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 8 = 22 \times 6 = 132$. In (B), 6 only divides x, so you must transpose $+8$ FIRST before moving 6: $\frac{x}{6} = 22 - 8 = 14 \implies x = 84$.

Q3226. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 4 = 30 \times 4 = 120$. In (B), 4 only divides x, so you must transpose $+4$ FIRST before moving 4: $\frac{x}{4} = 30 - 4 = 26 \implies x = 104$.

Q3227. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 8 = 12 \times 2 = 24$. In (B), 2 only divides x, so you must transpose $+8$ FIRST before moving 2: $\frac{x}{2} = 12 - 8 = 4 \implies x = 8$.

Q3228. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 6 = 24 \times 3 = 72$. In (B), 3 only divides x, so you must transpose $+6$ FIRST before moving 3: $\frac{x}{3} = 24 - 6 = 18 \implies x = 54$.

Q3229. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 28 \times 5 = 140$. In (B), 5 only divides x, so you must transpose $+2$ FIRST before moving 5: $\frac{x}{5} = 28 - 2 = 26 \implies x = 130$.

Q3230. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 25 \times 2 = 50$. In (B), 2 only divides x, so you must transpose $+4$ FIRST before moving 2: $\frac{x}{2} = 25 - 4 = 21 \implies x = 42$.

Q3231. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 5 = 29 \times 6 = 174$. In (B), 6 only divides x, so you must transpose $+5$ FIRST before moving 6: $\frac{x}{6} = 29 - 5 = 24 \implies x = 144$.

Q3232. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 9 = 30 \times 5 = 150$. In (B), 5 only divides x, so you must transpose $+9$ FIRST before moving 5: $\frac{x}{5} = 30 - 9 = 21 \implies x = 105$.

Q3233. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 8 = 21 \times 7 = 147$. In (B), 7 only divides x, so you must transpose $+8$ FIRST before moving 7: $\frac{x}{7} = 21 - 8 = 13 \implies x = 91$.

Q3234. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 7 = 14 \times 4 = 56$. In (B), 4 only divides x, so you must transpose $+7$ FIRST before moving 4: $\frac{x}{4} = 14 - 7 = 7 \implies x = 28$.

Q3235. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 7 = 29 \times 6 = 174$. In (B),

6 only divides x, so you must transpose +7 FIRST before moving 6: $\frac{x}{6} = 29 - 7 = 22 \implies x = 132$.

Q3236. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 8 = 26 \times 3 = 78$. In (B), 3 only divides x, so you must transpose +8 FIRST before moving 3: $\frac{x}{3} = 26 - 8 = 18 \implies x = 54$.

Q3237. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 7 = 21 \times 7 = 147$. In (B), 7 only divides x, so you must transpose +7 FIRST before moving 7: $\frac{x}{7} = 21 - 7 = 14 \implies x = 98$.

Q3238. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 4 = 12 \times 2 = 24$. In (B), 2 only divides x, so you must transpose +4 FIRST before moving 2: $\frac{x}{2} = 12 - 4 = 8 \implies x = 16$.

Q3239. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 8 = 11 \times 4 = 44$. In (B), 4 only divides x, so you must transpose +8 FIRST before moving 4: $\frac{x}{4} = 11 - 8 = 3 \implies x = 12$.

Q3240. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 6 = 16 \times 4 = 64$. In (B), 4 only divides x, so you must transpose +6 FIRST before moving 4: $\frac{x}{4} = 16 - 6 = 10 \implies x = 40$.

Q3241. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 4 = 24 \times 6 = 144$. In (B), 6 only divides x, so you must transpose +4 FIRST before moving 6: $\frac{x}{6} = 24 - 4 = 20 \implies x = 120$.

Q3242. In (A), 3 divides the entire left side, so you multiply the right side directly: $x + 4 = 10 \times 3 = 30$. In (B), 3 only divides x, so you must transpose +4 FIRST before moving 3: $\frac{x}{3} = 10 - 4 = 6 \implies x = 18$.

Q3243. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 3 = 29 \times 4 = 116$. In (B), 4 only divides x, so you must transpose +3 FIRST before moving 4: $\frac{x}{4} = 29 - 3 = 26 \implies x = 104$.

Q3244. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 8 = 12 \times 5 = 60$. In (B), 5 only divides x, so you must transpose +8 FIRST before moving 5: $\frac{x}{5} = 12 - 8 = 4 \implies x = 20$.

Q3245. In (A), 4 divides the entire left side, so you multiply the right side directly: $x + 5 = 20 \times 4 = 80$. In (B), 4 only divides x, so you must transpose +5 FIRST before moving 4: $\frac{x}{4} = 20 - 5 = 15 \implies x = 60$.

Q3246. In (A), 5 divides the entire left side, so you multiply the right side directly: $x + 2 = 22 \times 5 = 110$. In (B), 5 only divides x, so you must transpose +2 FIRST before moving 5: $\frac{x}{5} = 22 - 2 = 20 \implies x = 100$.

Q3247. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 3 = 17 \times 2 = 34$. In (B), 2 only divides x, so you must transpose +3 FIRST before moving 2: $\frac{x}{2} = 17 - 3 = 14 \implies x = 28$.

Q3248. In (A), 7 divides the entire left side, so you multiply the right side directly: $x + 4 = 19 \times 7 = 133$. In (B), 7 only divides x, so you must transpose +4 FIRST before moving 7: $\frac{x}{7} = 19 - 4 = 15 \implies x = 105$.

Q3249. In (A), 2 divides the entire left side, so you multiply the right side directly: $x + 9 = 13 \times 2 = 26$. In (B), 2 only divides x, so you must transpose +9 FIRST before moving 2: $\frac{x}{2} = 13 - 9 = 4 \implies x = 8$.

Q3250. In (A), 6 divides the entire left side, so you multiply the right side directly: $x + 3 = 10 \times 6 = 60$. In (B), 6 only divides x, so you must transpose +3 FIRST before moving 6: $\frac{x}{6} = 10 - 3 = 7 \implies x = 42$.

Q3251. Moving denominators yields $q(3x + 9) = p(5x - 5) \implies 15x + 45 = 10x - 10$. Transposing x-terms gives $(15 - 10)x = -10 - 45 \implies x = \frac{-55}{5}$.

Q3252. Moving denominators yields $q(2x + 7) = p(5x - 8) \implies 14x + 49 = 10x - 16$. Transposing x-terms gives $(14 - 10)x = -16 - 49 \implies x = \frac{-65}{4}$.

Q3253. Moving denominators yields $q(4x + 7) = p(2x - 2) \implies 8x + 14 = 12x - 12$. Transposing x-terms gives $(8 - 12)x = -12 - 14 \implies x = \frac{-26}{-4}$.

Q3254. Moving denominators yields $q(2x + 8) = p(5x - 5) \implies 10x + 40 = 35x - 35$. Transposing x-terms gives $(10 - 35)x = -35 - 40 \implies x = \frac{-75}{-25}$.

Q3255. Moving denominators yields $q(3x + 9) = p(5x - 3) \implies 12x + 36 = 35x - 21$. Transposing x-terms gives $(12 - 35)x = -21 - 36 \implies x = \frac{-57}{-23}$.

Q3256. Moving denominators yields $q(3x + 9) = p(5x - 2) \implies 15x + 45 = 20x - 8$. Transposing x-terms gives $(15 - 20)x = -8 - 45 \implies x = \frac{-53}{-5}$.

Q3257. Moving denominators yields $q(3x + 2) = p(5x - 5) \implies 9x + 6 = 15x - 15$. Transposing x-terms gives $(9 - 15)x = -15 - 6 \implies x = \frac{-21}{-6}$.

Q3258. Moving denominators yields $q(4x + 7) = p(4x - 7) \implies 28x + 49 = 8x - 14$. Transposing x-terms gives $(28 - 8)x = -14 - 49 \implies x = \frac{-63}{20}$.

Q3259. Moving denominators yields $q(5x + 4) = p(5x - 9) \implies 10x + 8 = 25x - 45$. Transposing x-terms gives $(10 - 25)x = -45 - 8 \implies x = \frac{-53}{-15}$.

Q3260. Moving denominators yields $q(5x + 3) = p(2x - 3) \implies 15x + 9 = 10x - 15$. Transposing x-terms gives $(15 - 10)x = -15 - 9 \implies x = \frac{-24}{5}$.

Q3261. Moving denominators yields $q(5x + 3) = p(2x - 3) \implies 35x + 21 = 8x - 12$. Transposing x-terms gives $(35 - 8)x = -12 - 21 \implies x = \frac{-33}{27}$.

Q3262. Moving denominators yields $q(4x + 8) = p(4x - 5) \implies 12x + 24 = 12x - 15$. Transposing x-terms gives $(12 - 12)x = -15 - 24 \implies x = \frac{-39}{0}$.

Q3263. Moving denominators yields $q(4x + 5) = p(2x - 4) \implies 12x + 15 = 12x - 24$. Transposing x-terms gives $(12 - 12)x = -24 - 15 \implies x = \frac{-39}{0}$.

Q3264. Moving denominators yields $q(5x + 3) = p(3x - 5) \implies 25x + 15 = 15x - 25$. Transposing x-terms gives $(25 - 15)x = -25 - 15 \implies x = \frac{-40}{10}$.

Q3265. Moving denominators yields $q(2x + 2) = p(2x - 5) \implies 10x + 10 = 14x - 35$. Transposing x-terms gives $(10 - 14)x = -35 - 10 \implies x = \frac{-45}{-4}$.

Q3266. Moving denominators yields $q(5x + 2) = p(5x - 1) \implies 20x + 8 = 30x - 6$. Transposing x-terms gives $(20 - 30)x = -6 - 8 \implies x = \frac{-14}{-10}$.

Q3267. Moving denominators yields $q(5x + 9) = p(3x - 2) \implies 25x + 45 = 21x - 14$. Transposing x-terms gives $(25 - 21)x = -14 - 45 \implies x = \frac{-59}{4}$.

Q3268. Moving denominators yields $q(4x + 8) = p(2x - 4) \implies 12x + 24 = 12x - 24$. Transposing x-terms gives $(12 - 12)x = -24 - 24 \implies x = \frac{-48}{0}$.

Q3269. Moving denominators yields $q(5x + 3) = p(4x -$

6) $\implies 25x + 15 = 28x - 42$. Transposing x-terms gives $(25 - 28)x = -42 - 15 \implies x = \frac{-57}{-3}$.

Q3270. Moving denominators yields $q(4x + 7) = p(3x - 7) \implies 12x + 21 = 21x - 49$. Transposing x-terms gives $(12 - 21)x = -49 - 21 \implies x = \frac{-70}{-9}$.

Q3271. Moving denominators yields $q(2x + 9) = p(2x - 3) \implies 8x + 36 = 4x - 6$. Transposing x-terms gives $(8 - 4)x = -6 - 36 \implies x = \frac{-42}{4}$.

Q3272. Moving denominators yields $q(5x + 4) = p(5x - 1) \implies 30x + 24 = 10x - 2$. Transposing x-terms gives $(30 - 10)x = -2 - 24 \implies x = \frac{-26}{20}$.

Q3273. Moving denominators yields $q(3x + 1) = p(3x - 8) \implies 12x + 4 = 15x - 40$. Transposing x-terms gives $(12 - 15)x = -40 - 4 \implies x = \frac{-44}{-3}$.

Q3274. Moving denominators yields $q(4x + 1) = p(5x - 1) \implies 16x + 4 = 30x - 6$. Transposing x-terms gives $(16 - 30)x = -6 - 4 \implies x = \frac{-10}{-14}$.

Q3275. Moving denominators yields $q(2x + 6) = p(5x - 6) \implies 14x + 42 = 35x - 42$. Transposing x-terms gives $(14 - 35)x = -42 - 42 \implies x = \frac{-84}{-21}$.

Q3276. Moving denominators yields $q(2x + 5) = p(4x - 8) \implies 14x + 35 = 24x - 48$. Transposing x-terms gives $(14 - 24)x = -48 - 35 \implies x = \frac{-83}{-10}$.

Q3277. Moving denominators yields $q(2x + 2) = p(4x - 3) \implies 8x + 8 = 12x - 9$. Transposing x-terms gives $(8 - 12)x = -9 - 8 \implies x = \frac{-17}{-4}$.

Q3278. Moving denominators yields $q(4x + 7) = p(3x - 6) \implies 16x + 28 = 18x - 36$. Transposing x-terms gives $(16 - 18)x = -36 - 28 \implies x = \frac{-64}{-2}$.

Q3279. Moving denominators yields $q(4x + 5) = p(2x - 6) \implies 16x + 20 = 4x - 12$. Transposing x-terms gives $(16 - 4)x = -12 - 20 \implies x = \frac{-32}{12}$.

Q3280. Moving denominators yields $q(4x + 2) = p(2x - 5) \implies 20x + 10 = 12x - 30$. Transposing x-terms gives $(20 - 12)x = -30 - 10 \implies x = \frac{-40}{8}$.

Q3281. Moving denominators yields $q(2x + 1) = p(5x - 3) \implies 14x + 7 = 10x - 6$. Transposing x-terms gives $(14 - 10)x = -6 - 7 \implies x = \frac{-13}{4}$.

Q3282. Moving denominators yields $q(5x + 7) = p(2x - 3) \implies 30x + 42 = 12x - 18$. Transposing x-terms gives $(30 - 12)x = -18 - 42 \implies x = \frac{-60}{18}$.

Q3283. Moving denominators yields $q(4x + 7) = p(2x - 9) \implies 16x + 28 = 12x - 54$. Transposing x-terms gives $(16 - 12)x = -54 - 28 \implies x = \frac{-82}{4}$.

Q3284. Moving denominators yields $q(5x + 5) = p(3x - 3) \implies 20x + 20 = 15x - 15$. Transposing x-terms gives $(20 - 15)x = -15 - 20 \implies x = \frac{-35}{5}$.

Q3285. Moving denominators yields $q(3x + 1) = p(4x - 7) \implies 15x + 5 = 24x - 42$. Transposing x-terms gives $(15 - 24)x = -42 - 5 \implies x = \frac{-47}{-9}$.

Q3286. Moving denominators yields $q(5x + 7) = p(3x - 2) \implies 20x + 28 = 6x - 4$. Transposing x-terms gives $(20 - 6)x = -4 - 28 \implies x = \frac{-32}{14}$.

Q3287. Moving denominators yields $q(4x + 8) = p(4x - 3) \implies 28x + 56 = 16x - 12$. Transposing x-terms gives $(28 - 16)x = -12 - 56 \implies x = \frac{-68}{12}$.

Q3288. Moving denominators yields $q(5x + 7) = p(2x - 7) \implies 25x + 35 = 6x - 21$. Transposing x-terms gives $(25 - 6)x = -21 - 35 \implies x = \frac{-56}{19}$.

Q3289. Moving denominators yields $q(2x + 7) = p(5x - 3) \implies 10x + 35 = 20x - 12$. Transposing x-terms gives $(10 - 20)x = -12 - 35 \implies x = \frac{-47}{-10}$.

Q3290. Moving denominators yields $q(5x + 7) = p(5x - 4) \implies 10x + 14 = 30x - 24$. Transposing x-terms gives $(10 - 30)x = -24 - 14 \implies x = \frac{-38}{-20}$.

Q3291. Moving denominators yields $q(3x + 4) = p(3x - 9) \implies 15x + 20 = 18x - 54$. Transposing x-terms gives $(15 - 18)x = -54 - 20 \implies x = \frac{-74}{-3}$.

Q3292. Moving denominators yields $q(2x + 3) = p(2x - 2) \implies 10x + 15 = 14x - 14$. Transposing x-terms gives $(10 - 14)x = -14 - 15 \implies x = \frac{-29}{-4}$.

Q3293. Moving denominators yields $q(3x + 6) = p(3x - 9) \implies 6x + 12 = 18x - 54$. Transposing x-terms gives $(6 - 18)x = -54 - 12 \implies x = \frac{-66}{-12}$.

Q3294. Moving denominators yields $q(2x + 9) = p(4x - 8) \implies 6x + 27 = 20x - 40$. Transposing x-terms gives $(6 - 20)x = -40 - 27 \implies x = \frac{-67}{-14}$.

Q3295. Moving denominators yields $q(2x + 2) = p(2x - 2) \implies 6x + 6 = 14x - 14$. Transposing x-terms gives $(6 - 14)x = -14 - 6 \implies x = \frac{-20}{-8}$.

Q3296. Moving denominators yields $q(3x + 4) = p(5x - 8) \implies 18x + 24 = 20x - 32$. Transposing x-terms gives $(18 - 20)x = -32 - 24 \implies x = \frac{-56}{-2}$.

Q3297. Moving denominators yields $q(2x + 7) = p(3x - 5) \implies 4x + 14 = 6x - 10$. Transposing x-terms gives $(4 - 6)x = -10 - 14 \implies x = \frac{-24}{-2}$.

Q3298. Moving denominators yields $q(3x + 5) = p(4x - 3) \implies 18x + 30 = 12x - 9$. Transposing x-terms gives $(18 - 12)x = -9 - 30 \implies x = \frac{-39}{6}$.

Q3299. Moving denominators yields $q(3x + 5) = p(2x - 1) \implies 6x + 10 = 12x - 6$. Transposing x-terms gives $(6 - 12)x = -6 - 10 \implies x = \frac{-16}{-6}$.

Q3300. Moving denominators yields $q(2x + 1) = p(4x - 1) \implies 12x + 6 = 12x - 3$. Transposing x-terms gives $(12 - 12)x = -3 - 6 \implies x = \frac{-9}{0}$.

Q3301. Moving denominators yields $q(2x + 7) = p(4x - 7) \implies 8x + 28 = 8x - 14$. Transposing x-terms gives $(8 - 8)x = -14 - 28 \implies x = \frac{-42}{0}$.

Q3302. Moving denominators yields $q(5x + 1) = p(4x - 1) \implies 25x + 5 = 20x - 5$. Transposing x-terms gives $(25 - 20)x = -5 - 5 \implies x = \frac{-10}{5}$.

Q3303. Moving denominators yields $q(4x + 1) = p(3x - 1) \implies 8x + 2 = 12x - 4$. Transposing x-terms gives $(8 - 12)x = -4 - 2 \implies x = \frac{-6}{-4}$.

Q3304. Moving denominators yields $q(4x + 7) = p(5x - 2) \implies 16x + 28 = 30x - 12$. Transposing x-terms gives $(16 - 30)x = -12 - 28 \implies x = \frac{-40}{-14}$.

Q3305. Moving denominators yields $q(4x + 5) = p(2x - 4) \implies 24x + 30 = 6x - 12$. Transposing x-terms gives $(24 - 6)x = -12 - 30 \implies x = \frac{-42}{18}$.

Q3306. Moving denominators yields $q(3x + 3) = p(2x - 9) \implies 18x + 18 = 12x - 54$. Transposing x-terms gives $(18 - 12)x = -54 - 18 \implies x = \frac{-72}{6}$.

Q3307. Moving denominators yields $q(3x + 6) = p(4x - 3) \implies 6x + 12 = 28x - 21$. Transposing x-terms gives $(6 - 28)x = -21 - 12 \implies x = \frac{-33}{-22}$.

Q3308. Moving denominators yields $q(4x+4) = p(3x-1) \implies 12x+12 = 12x-4$. Transposing x-terms gives $(12-12)x = -4-12 \implies x = \frac{-16}{0}$.

Q3309. Moving denominators yields $q(4x+2) = p(5x-4) \implies 12x+6 = 30x-24$. Transposing x-terms gives $(12-30)x = -24-6 \implies x = \frac{-30}{-18}$.

Q3310. Moving denominators yields $q(5x+4) = p(5x-8) \implies 10x+8 = 30x-48$. Transposing x-terms gives $(10-30)x = -48-8 \implies x = \frac{-56}{-20}$.

Q3311. Moving denominators yields $q(3x+9) = p(3x-4) \implies 21x+63 = 21x-28$. Transposing x-terms gives $(21-21)x = -28-63 \implies x = \frac{-91}{0}$.

Q3312. Moving denominators yields $q(4x+8) = p(4x-3) \implies 24x+48 = 20x-15$. Transposing x-terms gives $(24-20)x = -15-48 \implies x = \frac{-63}{4}$.

Q3313. Moving denominators yields $q(4x+1) = p(3x-4) \implies 24x+6 = 15x-20$. Transposing x-terms gives $(24-15)x = -20-6 \implies x = \frac{-26}{9}$.

Q3314. Moving denominators yields $q(2x+3) = p(2x-6) \implies 12x+18 = 4x-12$. Transposing x-terms gives $(12-4)x = -12-18 \implies x = \frac{-30}{8}$.

Q3315. Moving denominators yields $q(2x+7) = p(2x-7) \implies 4x+14 = 6x-21$. Transposing x-terms gives $(4-6)x = -21-14 \implies x = \frac{-35}{-2}$.

Q3316. Moving denominators yields $q(5x+5) = p(2x-7) \implies 35x+35 = 8x-28$. Transposing x-terms gives $(35-8)x = -28-35 \implies x = \frac{-63}{27}$.

Q3317. Moving denominators yields $q(5x+7) = p(4x-7) \implies 35x+49 = 20x-35$. Transposing x-terms gives $(35-20)x = -35-49 \implies x = \frac{-84}{15}$.

Q3318. Moving denominators yields $q(5x+3) = p(2x-9) \implies 25x+15 = 12x-54$. Transposing x-terms gives $(25-12)x = -54-15 \implies x = \frac{-69}{13}$.

Q3319. Moving denominators yields $q(4x+7) = p(5x-1) \implies 24x+42 = 10x-2$. Transposing x-terms gives $(24-10)x = -2-42 \implies x = \frac{-44}{14}$.

Q3320. Moving denominators yields $q(2x+3) = p(5x-9) \implies 12x+18 = 15x-27$. Transposing x-terms gives $(12-15)x = -27-18 \implies x = \frac{-45}{-3}$.

Q3321. Moving denominators yields $q(4x+3) = p(4x-7) \implies 28x+21 = 20x-35$. Transposing x-terms gives $(28-20)x = -35-21 \implies x = \frac{-56}{8}$.

Q3322. Moving denominators yields $q(5x+2) = p(5x-4) \implies 20x+8 = 30x-24$. Transposing x-terms gives $(20-30)x = -24-8 \implies x = \frac{-32}{-10}$.

Q3323. Moving denominators yields $q(5x+8) = p(5x-6) \implies 20x+32 = 30x-36$. Transposing x-terms gives $(20-30)x = -36-32 \implies x = \frac{-68}{-10}$.

Q3324. Moving denominators yields $q(5x+1) = p(3x-5) \implies 30x+6 = 18x-30$. Transposing x-terms gives $(30-18)x = -30-6 \implies x = \frac{-36}{12}$.

Q3325. Moving denominators yields $q(3x+6) = p(4x-5) \implies 21x+42 = 16x-20$. Transposing x-terms gives $(21-16)x = -20-42 \implies x = \frac{-62}{5}$.

Q3326. Moving denominators yields $q(3x+7) = p(2x-7) \implies 6x+14 = 12x-42$. Transposing x-terms gives $(6-12)x = -42-14 \implies x = \frac{-56}{-6}$.

Q3327. Moving denominators yields $q(5x+4) = p(4x-1) \implies 30x+24 = 12x-3$. Transposing x-terms gives $(30-12)x = -3-24 \implies x = \frac{-27}{18}$.

Q3328. Moving denominators yields $q(2x+7) = p(5x-3) \implies 4x+14 = 15x-9$. Transposing x-terms gives $(4-15)x = -9-14 \implies x = \frac{-23}{-11}$.

Q3329. Moving denominators yields $q(2x+1) = p(5x-9) \implies 6x+3 = 15x-27$. Transposing x-terms gives $(6-15)x = -27-3 \implies x = \frac{-30}{-9}$.

Q3330. Moving denominators yields $q(2x+1) = p(2x-3) \implies 4x+2 = 8x-12$. Transposing x-terms gives $(4-8)x = -12-2 \implies x = \frac{-14}{-4}$.

Q3331. Moving denominators yields $q(2x+3) = p(5x-7) \implies 4x+6 = 15x-21$. Transposing x-terms gives $(4-15)x = -21-6 \implies x = \frac{-27}{-11}$.

Q3332. Moving denominators yields $q(4x+4) = p(3x-4) \implies 12x+12 = 21x-28$. Transposing x-terms gives $(12-21)x = -28-12 \implies x = \frac{-40}{-9}$.

Q3333. Moving denominators yields $q(4x+3) = p(2x-7) \implies 8x+6 = 6x-21$. Transposing x-terms gives $(8-6)x = -21-6 \implies x = \frac{-27}{2}$.

Q3334. Moving denominators yields $q(3x+4) = p(3x-1) \implies 18x+24 = 9x-3$. Transposing x-terms gives $(18-9)x = -3-24 \implies x = \frac{-27}{9}$.

Q3335. Moving denominators yields $q(3x+8) = p(2x-8) \implies 21x+56 = 8x-32$. Transposing x-terms gives $(21-8)x = -32-56 \implies x = \frac{-88}{13}$.

Q3336. Moving denominators yields $q(5x+6) = p(4x-2) \implies 20x+24 = 28x-14$. Transposing x-terms gives $(20-28)x = -14-24 \implies x = \frac{-38}{-8}$.

Q3337. Moving denominators yields $q(4x+3) = p(4x-4) \implies 16x+12 = 20x-20$. Transposing x-terms gives $(16-20)x = -20-12 \implies x = \frac{-32}{-4}$.

Q3338. Moving denominators yields $q(4x+4) = p(4x-8) \implies 16x+16 = 16x-32$. Transposing x-terms gives $(16-16)x = -32-16 \implies x = \frac{-48}{0}$.

Q3339. Moving denominators yields $q(2x+7) = p(4x-5) \implies 8x+28 = 20x-25$. Transposing x-terms gives $(8-20)x = -25-28 \implies x = \frac{-53}{-12}$.

Q3340. Moving denominators yields $q(2x+5) = p(2x-3) \implies 12x+30 = 14x-21$. Transposing x-terms gives $(12-14)x = -21-30 \implies x = \frac{-51}{-2}$.

Q3341. Moving denominators yields $q(4x+9) = p(4x-3) \implies 24x+54 = 12x-9$. Transposing x-terms gives $(24-12)x = -9-54 \implies x = \frac{-63}{12}$.

Q3342. Moving denominators yields $q(3x+1) = p(3x-7) \implies 6x+2 = 12x-28$. Transposing x-terms gives $(6-12)x = -28-2 \implies x = \frac{-30}{-6}$.

Q3343. Moving denominators yields $q(2x+7) = p(5x-4) \implies 14x+49 = 30x-24$. Transposing x-terms gives $(14-30)x = -24-49 \implies x = \frac{-73}{-16}$.

Q3344. Moving denominators yields $q(5x+5) = p(2x-6) \implies 20x+20 = 4x-12$. Transposing x-terms gives $(20-4)x = -12-20 \implies x = \frac{-32}{16}$.

Q3345. Moving denominators yields $q(5x+7) = p(3x-9) \implies 25x+35 = 18x-54$. Transposing x-terms gives $(25-18)x = -54-35 \implies x = \frac{-89}{7}$.

Q3346. Moving denominators yields $q(2x+1) = p(3x-6) \implies 4x+2 = 15x-30$. Transposing x-terms gives

$(4-15)x = -30-2 \implies x = \frac{-32}{-11}$.

Q3347. Moving denominators yields $q(3x+6) = p(3x-5) \implies 18x+36 = 15x-25$. Transposing x-terms gives $(18-15)x = -25-36 \implies x = \frac{-61}{3}$.

Q3348. Moving denominators yields $q(4x+8) = p(2x-7) \implies 16x+32 = 8x-28$. Transposing x-terms gives $(16-8)x = -28-32 \implies x = \frac{-60}{8}$.

Q3349. Moving denominators yields $q(4x+4) = p(3x-9) \implies 28x+28 = 21x-63$. Transposing x-terms gives $(28-21)x = -63-28 \implies x = \frac{-91}{7}$.

Q3350. Moving denominators yields $q(4x+4) = p(5x-3) \implies 16x+16 = 25x-15$. Transposing x-terms gives $(16-25)x = -15-16 \implies x = \frac{-31}{-9}$.

Q3351. Moving denominators yields $q(2x+4) = p(3x-3) \implies 12x+24 = 18x-18$. Transposing x-terms gives $(12-18)x = -18-24 \implies x = \frac{-42}{-6}$.

Q3352. Moving denominators yields $q(5x+9) = p(4x-1) \implies 30x+54 = 24x-6$. Transposing x-terms gives $(30-24)x = -6-54 \implies x = \frac{-60}{6}$.

Q3353. Moving denominators yields $q(3x+3) = p(5x-5) \implies 9x+9 = 25x-25$. Transposing x-terms gives $(9-25)x = -25-9 \implies x = \frac{-34}{-16}$.

Q3354. Moving denominators yields $q(2x+5) = p(2x-7) \implies 4x+10 = 12x-42$. Transposing x-terms gives $(4-12)x = -42-10 \implies x = \frac{-52}{-8}$.

Q3355. Moving denominators yields $q(5x+6) = p(3x-3) \implies 15x+18 = 12x-12$. Transposing x-terms gives $(15-12)x = -12-18 \implies x = \frac{-30}{3}$.

Q3356. Moving denominators yields $q(2x+8) = p(3x-7) \implies 12x+48 = 9x-21$. Transposing x-terms gives $(12-9)x = -21-48 \implies x = \frac{-69}{3}$.

Q3357. Moving denominators yields $q(5x+2) = p(2x-4) \implies 15x+6 = 12x-24$. Transposing x-terms gives $(15-12)x = -24-6 \implies x = \frac{-30}{3}$.

Q3358. Moving denominators yields $q(3x+7) = p(3x-9) \implies 12x+28 = 21x-63$. Transposing x-terms gives $(12-21)x = -63-28 \implies x = \frac{-91}{-9}$.

Q3359. Moving denominators yields $q(3x+4) = p(5x-9) \implies 9x+12 = 15x-27$. Transposing x-terms gives $(9-15)x = -27-12 \implies x = \frac{-39}{-6}$.

Q3360. Moving denominators yields $q(3x+9) = p(3x-7) \implies 6x+18 = 15x-35$. Transposing x-terms gives $(6-15)x = -35-18 \implies x = \frac{-53}{-9}$.

Q3361. Moving denominators yields $q(3x+4) = p(5x-2) \implies 15x+20 = 30x-12$. Transposing x-terms gives $(15-30)x = -12-20 \implies x = \frac{-32}{-15}$.

Q3362. Moving denominators yields $q(3x+5) = p(4x-9) \implies 9x+15 = 8x-18$. Transposing x-terms gives $(9-8)x = -18-15 \implies x = \frac{-33}{1}$.

Q3363. Moving denominators yields $q(2x+1) = p(4x-9) \implies 4x+2 = 12x-27$. Transposing x-terms gives $(4-12)x = -27-2 \implies x = \frac{-29}{-8}$.

Q3364. Moving denominators yields $q(2x+7) = p(3x-7) \implies 8x+28 = 21x-49$. Transposing x-terms gives $(8-21)x = -49-28 \implies x = \frac{-77}{-13}$.

Q3365. Moving denominators yields $q(2x+2) = p(5x-7) \implies 4x+4 = 30x-42$. Transposing x-terms gives $(4-30)x = -42-4 \implies x = \frac{-46}{-26}$.

Q3366. Moving denominators yields $q(2x+7) = p(5x-3) \implies 12x+42 = 10x-6$. Transposing x-terms gives $(12-10)x = -6-42 \implies x = \frac{-48}{2}$.

Q3367. Moving denominators yields $q(2x+1) = p(2x-1) \implies 8x+4 = 12x-6$. Transposing x-terms gives $(8-12)x = -6-4 \implies x = \frac{-10}{-4}$.

Q3368. Moving denominators yields $q(2x+2) = p(2x-7) \implies 8x+8 = 10x-35$. Transposing x-terms gives $(8-10)x = -35-8 \implies x = \frac{-43}{-2}$.

Q3369. Moving denominators yields $q(4x+7) = p(4x-4) \implies 24x+42 = 16x-16$. Transposing x-terms gives $(24-16)x = -16-42 \implies x = \frac{-58}{8}$.

Q3370. Moving denominators yields $q(4x+8) = p(5x-7) \implies 12x+24 = 10x-14$. Transposing x-terms gives $(12-10)x = -14-24 \implies x = \frac{-38}{2}$.

Q3371. Moving denominators yields $q(4x+6) = p(4x-8) \implies 8x+12 = 12x-24$. Transposing x-terms gives $(8-12)x = -24-12 \implies x = \frac{-36}{-4}$.

Q3372. Moving denominators yields $q(3x+1) = p(2x-5) \implies 12x+4 = 6x-15$. Transposing x-terms gives $(12-6)x = -15-4 \implies x = \frac{-19}{6}$.

Q3373. Moving denominators yields $q(5x+3) = p(5x-7) \implies 10x+6 = 30x-42$. Transposing x-terms gives $(10-30)x = -42-6 \implies x = \frac{-48}{-20}$.

Q3374. Moving denominators yields $q(5x+2) = p(3x-9) \implies 15x+6 = 6x-18$. Transposing x-terms gives $(15-6)x = -18-6 \implies x = \frac{-24}{9}$.

Q3375. Moving denominators yields $q(3x+5) = p(2x-5) \implies 9x+15 = 10x-25$. Transposing x-terms gives $(9-10)x = -25-15 \implies x = \frac{-40}{-1}$.

Q3376. Moving denominators yields $q(3x+5) = p(3x-6) \implies 6x+10 = 12x-24$. Transposing x-terms gives $(6-12)x = -24-10 \implies x = \frac{-34}{-6}$.

Q3377. Moving denominators yields $q(2x+3) = p(2x-3) \implies 10x+15 = 14x-21$. Transposing x-terms gives $(10-14)x = -21-15 \implies x = \frac{-36}{-4}$.

Q3378. Moving denominators yields $q(3x+8) = p(4x-3) \implies 15x+40 = 16x-12$. Transposing x-terms gives $(15-16)x = -12-40 \implies x = \frac{-52}{-1}$.

Q3379. Moving denominators yields $q(4x+8) = p(4x-7) \implies 8x+16 = 20x-35$. Transposing x-terms gives $(8-20)x = -35-16 \implies x = \frac{-51}{-12}$.

Q3380. Moving denominators yields $q(5x+7) = p(5x-7) \implies 25x+35 = 25x-35$. Transposing x-terms gives $(25-25)x = -35-35 \implies x = \frac{-70}{0}$.

Q3381. Moving denominators yields $q(2x+5) = p(2x-6) \implies 8x+20 = 12x-36$. Transposing x-terms gives $(8-12)x = -36-20 \implies x = \frac{-56}{-4}$.

Q3382. Moving denominators yields $q(3x+8) = p(5x-7) \implies 18x+48 = 35x-49$. Transposing x-terms gives $(18-35)x = -49-48 \implies x = \frac{-97}{-17}$.

Q3383. Moving denominators yields $q(5x+3) = p(5x-6) \implies 15x+9 = 20x-24$. Transposing x-terms gives $(15-20)x = -24-9 \implies x = \frac{-33}{-5}$.

Q3384. Moving denominators yields $q(3x+4) = p(2x-4) \implies 6x+8 = 10x-20$. Transposing x-terms gives $(6-10)x = -20-8 \implies x = \frac{-28}{-4}$.

Q3385. Moving denominators yields $q(5x+4) = p(5x-$

2) $\implies 30x + 24 = 15x - 6$. Transposing x-terms gives $(30 - 15)x = -6 - 24 \implies x = \frac{-30}{15}$.

Q3386. Moving denominators yields $q(2x + 8) = p(2x - 5) \implies 6x + 24 = 4x - 10$. Transposing x-terms gives $(6 - 4)x = -10 - 24 \implies x = \frac{-34}{2}$.

Q3387. Moving denominators yields $q(3x + 9) = p(3x - 2) \implies 21x + 63 = 15x - 10$. Transposing x-terms gives $(21 - 15)x = -10 - 63 \implies x = \frac{-73}{6}$.

Q3388. Moving denominators yields $q(5x + 5) = p(4x - 2) \implies 35x + 35 = 24x - 12$. Transposing x-terms gives $(35 - 24)x = -12 - 35 \implies x = \frac{-47}{11}$.

Q3389. Moving denominators yields $q(4x + 1) = p(2x - 7) \implies 8x + 2 = 10x - 35$. Transposing x-terms gives $(8 - 10)x = -35 - 2 \implies x = \frac{-37}{-2}$.

Q3390. Moving denominators yields $q(4x + 2) = p(4x - 5) \implies 12x + 6 = 12x - 15$. Transposing x-terms gives $(12 - 12)x = -15 - 6 \implies x = \frac{-21}{0}$.

Q3391. Moving denominators yields $q(5x + 1) = p(2x - 3) \implies 15x + 3 = 8x - 12$. Transposing x-terms gives $(15 - 8)x = -12 - 3 \implies x = \frac{-15}{7}$.

Q3392. Moving denominators yields $q(3x + 5) = p(5x - 5) \implies 15x + 25 = 30x - 30$. Transposing x-terms gives $(15 - 30)x = -30 - 25 \implies x = \frac{-55}{-15}$.

Q3393. Moving denominators yields $q(3x + 5) = p(3x - 6) \implies 9x + 15 = 15x - 30$. Transposing x-terms gives $(9 - 15)x = -30 - 15 \implies x = \frac{-45}{-6}$.

Q3394. Moving denominators yields $q(2x + 7) = p(2x - 7) \implies 6x + 21 = 4x - 14$. Transposing x-terms gives $(6 - 4)x = -14 - 21 \implies x = \frac{-35}{2}$.

Q3395. Moving denominators yields $q(5x + 7) = p(4x - 6) \implies 30x + 42 = 12x - 18$. Transposing x-terms gives $(30 - 12)x = -18 - 42 \implies x = \frac{-60}{18}$.

Q3396. Moving denominators yields $q(2x + 5) = p(4x - 6) \implies 12x + 30 = 20x - 30$. Transposing x-terms gives $(12 - 20)x = -30 - 30 \implies x = \frac{-60}{-8}$.

Q3397. Moving denominators yields $q(5x + 4) = p(2x - 6) \implies 25x + 20 = 14x - 42$. Transposing x-terms gives $(25 - 14)x = -42 - 20 \implies x = \frac{-62}{11}$.

Q3398. Moving denominators yields $q(2x + 2) = p(5x - 9) \implies 12x + 12 = 15x - 27$. Transposing x-terms gives $(12 - 15)x = -27 - 12 \implies x = \frac{-39}{-3}$.

Q3399. Moving denominators yields $q(3x + 6) = p(2x - 2) \implies 21x + 42 = 14x - 14$. Transposing x-terms gives $(21 - 14)x = -14 - 42 \implies x = \frac{-56}{7}$.

Q3400. Moving denominators yields $q(2x + 5) = p(3x - 7) \implies 10x + 25 = 9x - 21$. Transposing x-terms gives $(10 - 9)x = -21 - 25 \implies x = \frac{-46}{1}$.

Q3401. Moving denominators yields $q(2x + 1) = p(3x - 2) \implies 12x + 6 = 12x - 8$. Transposing x-terms gives $(12 - 12)x = -8 - 6 \implies x = \frac{-14}{0}$.

Q3402. Moving denominators yields $q(2x + 3) = p(4x - 2) \implies 14x + 21 = 20x - 10$. Transposing x-terms gives $(14 - 20)x = -10 - 21 \implies x = \frac{-31}{-6}$.

Q3403. Moving denominators yields $q(3x + 7) = p(5x - 6) \implies 18x + 42 = 30x - 36$. Transposing x-terms gives $(18 - 30)x = -36 - 42 \implies x = \frac{-78}{-12}$.

Q3404. Moving denominators yields $q(3x + 2) = p(5x - 6) \implies 21x + 14 = 15x - 18$. Transposing x-terms gives $(21 - 15)x = -18 - 14 \implies x = \frac{-32}{6}$.

Q3405. Moving denominators yields $q(3x + 1) = p(4x - 3) \implies 15x + 5 = 20x - 15$. Transposing x-terms gives $(15 - 20)x = -15 - 5 \implies x = \frac{-20}{-5}$.

Q3406. Moving denominators yields $q(4x + 3) = p(2x - 3) \implies 16x + 12 = 12x - 18$. Transposing x-terms gives $(16 - 12)x = -18 - 12 \implies x = \frac{-30}{4}$.

Q3407. Moving denominators yields $q(2x + 2) = p(2x - 8) \implies 14x + 14 = 6x - 24$. Transposing x-terms gives $(14 - 6)x = -24 - 14 \implies x = \frac{-38}{8}$.

Q3408. Moving denominators yields $q(4x + 4) = p(4x - 6) \implies 28x + 28 = 28x - 42$. Transposing x-terms gives $(28 - 28)x = -42 - 28 \implies x = \frac{-70}{0}$.

Q3409. Moving denominators yields $q(5x + 7) = p(4x - 8) \implies 35x + 49 = 16x - 32$. Transposing x-terms gives $(35 - 16)x = -32 - 49 \implies x = \frac{-81}{19}$.

Q3410. Moving denominators yields $q(2x + 4) = p(4x - 9) \implies 4x + 8 = 24x - 54$. Transposing x-terms gives $(4 - 24)x = -54 - 8 \implies x = \frac{-62}{-20}$.

Q3411. Moving denominators yields $q(2x + 9) = p(4x - 4) \implies 12x + 54 = 28x - 28$. Transposing x-terms gives $(12 - 28)x = -28 - 54 \implies x = \frac{-82}{-16}$.

Q3412. Moving denominators yields $q(5x + 1) = p(2x - 4) \implies 10x + 2 = 4x - 8$. Transposing x-terms gives $(10 - 4)x = -8 - 2 \implies x = \frac{-10}{6}$.

Q3413. Moving denominators yields $q(4x + 3) = p(4x - 9) \implies 16x + 12 = 24x - 54$. Transposing x-terms gives $(16 - 24)x = -54 - 12 \implies x = \frac{-66}{-8}$.

Q3414. Moving denominators yields $q(5x + 8) = p(4x - 1) \implies 20x + 32 = 12x - 3$. Transposing x-terms gives $(20 - 12)x = -3 - 32 \implies x = \frac{-35}{8}$.

Q3415. Moving denominators yields $q(5x + 3) = p(5x - 7) \implies 10x + 6 = 25x - 35$. Transposing x-terms gives $(10 - 25)x = -35 - 6 \implies x = \frac{-41}{-15}$.

Q3416. Moving denominators yields $q(3x + 2) = p(5x - 1) \implies 6x + 4 = 20x - 4$. Transposing x-terms gives $(6 - 20)x = -4 - 4 \implies x = \frac{-8}{-14}$.

Q3417. Moving denominators yields $q(4x + 6) = p(4x - 7) \implies 24x + 36 = 12x - 21$. Transposing x-terms gives $(24 - 12)x = -21 - 36 \implies x = \frac{-57}{12}$.

Q3418. Moving denominators yields $q(2x + 1) = p(4x - 9) \implies 4x + 2 = 20x - 45$. Transposing x-terms gives $(4 - 20)x = -45 - 2 \implies x = \frac{-47}{-16}$.

Q3419. Moving denominators yields $q(4x + 6) = p(3x - 4) \implies 20x + 30 = 15x - 20$. Transposing x-terms gives $(20 - 15)x = -20 - 30 \implies x = \frac{-50}{5}$.

Q3420. Moving denominators yields $q(5x + 6) = p(4x - 8) \implies 10x + 12 = 12x - 24$. Transposing x-terms gives $(10 - 12)x = -24 - 12 \implies x = \frac{-36}{-2}$.

Q3421. Moving denominators yields $q(3x + 3) = p(2x - 4) \implies 6x + 6 = 14x - 28$. Transposing x-terms gives $(6 - 14)x = -28 - 6 \implies x = \frac{-34}{-8}$.

Q3422. Moving denominators yields $q(2x + 7) = p(2x - 5) \implies 10x + 35 = 4x - 10$. Transposing x-terms gives $(10 - 4)x = -10 - 35 \implies x = \frac{-45}{6}$.

Q3423. Moving denominators yields $q(4x + 2) = p(4x - 3) \implies 24x + 12 = 28x - 21$. Transposing x-terms gives $(24 - 28)x = -21 - 12 \implies x = \frac{-33}{-4}$.

Q3424. Moving denominators yields $q(3x+9) = p(2x-5) \implies 9x+27 = 8x-20$. Transposing x-terms gives $(9-8)x = -20-27 \implies x = \frac{-47}{1}$.

Q3425. Moving denominators yields $q(3x+1) = p(3x-9) \implies 12x+4 = 18x-54$. Transposing x-terms gives $(12-18)x = -54-4 \implies x = \frac{-58}{-6}$.

Q3426. Moving denominators yields $q(4x+7) = p(3x-1) \implies 24x+42 = 12x-4$. Transposing x-terms gives $(24-12)x = -4-42 \implies x = \frac{-46}{12}$.

Q3427. Moving denominators yields $q(2x+1) = p(5x-3) \implies 6x+3 = 25x-15$. Transposing x-terms gives $(6-25)x = -15-3 \implies x = \frac{-18}{-19}$.

Q3428. Moving denominators yields $q(5x+2) = p(3x-9) \implies 30x+12 = 9x-27$. Transposing x-terms gives $(30-9)x = -27-12 \implies x = \frac{-39}{21}$.

Q3429. Moving denominators yields $q(3x+8) = p(3x-6) \implies 9x+24 = 6x-12$. Transposing x-terms gives $(9-6)x = -12-24 \implies x = \frac{-36}{3}$.

Q3430. Moving denominators yields $q(5x+6) = p(3x-9) \implies 20x+24 = 12x-36$. Transposing x-terms gives $(20-12)x = -36-24 \implies x = \frac{-60}{8}$.

Q3431. Moving denominators yields $q(5x+5) = p(2x-5) \implies 30x+30 = 8x-20$. Transposing x-terms gives $(30-8)x = -20-30 \implies x = \frac{-50}{22}$.

Q3432. Moving denominators yields $q(3x+4) = p(2x-9) \implies 21x+28 = 8x-36$. Transposing x-terms gives $(21-8)x = -36-28 \implies x = \frac{-64}{13}$.

Q3433. Moving denominators yields $q(2x+2) = p(4x-6) \implies 12x+12 = 8x-12$. Transposing x-terms gives $(12-8)x = -12-12 \implies x = \frac{-24}{4}$.

Q3434. Moving denominators yields $q(4x+2) = p(2x-6) \implies 12x+6 = 4x-12$. Transposing x-terms gives $(12-4)x = -12-6 \implies x = \frac{-18}{8}$.

Q3435. Moving denominators yields $q(5x+4) = p(5x-7) \implies 30x+24 = 10x-14$. Transposing x-terms gives $(30-10)x = -14-24 \implies x = \frac{-38}{20}$.

Q3436. Moving denominators yields $q(5x+2) = p(2x-4) \implies 30x+12 = 6x-12$. Transposing x-terms gives $(30-6)x = -12-12 \implies x = \frac{-24}{24}$.

Q3437. Moving denominators yields $q(2x+2) = p(2x-9) \implies 8x+8 = 10x-45$. Transposing x-terms gives $(8-10)x = -45-8 \implies x = \frac{-53}{-2}$.

Q3438. Moving denominators yields $q(5x+8) = p(3x-4) \implies 10x+16 = 9x-12$. Transposing x-terms gives $(10-9)x = -12-16 \implies x = \frac{-28}{1}$.

Q3439. Moving denominators yields $q(3x+6) = p(3x-1) \implies 18x+36 = 18x-6$. Transposing x-terms gives $(18-18)x = -6-36 \implies x = \frac{-42}{0}$.

Q3440. Moving denominators yields $q(4x+5) = p(3x-8) \implies 24x+30 = 12x-32$. Transposing x-terms gives $(24-12)x = -32-30 \implies x = \frac{-62}{12}$.

Q3441. Moving denominators yields $q(5x+5) = p(5x-6) \implies 15x+15 = 35x-42$. Transposing x-terms gives $(15-35)x = -42-15 \implies x = \frac{-57}{-20}$.

Q3442. Moving denominators yields $q(3x+8) = p(5x-5) \implies 9x+24 = 20x-20$. Transposing x-terms gives $(9-20)x = -20-24 \implies x = \frac{-44}{-11}$.

Q3443. Moving denominators yields $q(4x+5) = p(3x-3) \implies 8x+10 = 9x-9$. Transposing x-terms gives $(8-9)x = -9-10 \implies x = \frac{-19}{-1}$.

Q3444. Moving denominators yields $q(4x+9) = p(4x-4) \implies 20x+45 = 24x-24$. Transposing x-terms gives $(20-24)x = -24-45 \implies x = \frac{-69}{-4}$.

Q3445. Moving denominators yields $q(4x+7) = p(3x-3) \implies 8x+14 = 21x-21$. Transposing x-terms gives $(8-21)x = -21-14 \implies x = \frac{-35}{-13}$.

Q3446. Moving denominators yields $q(3x+6) = p(4x-4) \implies 21x+42 = 16x-16$. Transposing x-terms gives $(21-16)x = -16-42 \implies x = \frac{-58}{5}$.

Q3447. Moving denominators yields $q(3x+9) = p(3x-4) \implies 6x+18 = 15x-20$. Transposing x-terms gives $(6-15)x = -20-18 \implies x = \frac{-38}{-9}$.

Q3448. Moving denominators yields $q(2x+6) = p(4x-4) \implies 10x+30 = 8x-8$. Transposing x-terms gives $(10-8)x = -8-30 \implies x = \frac{-38}{2}$.

Q3449. Moving denominators yields $q(2x+6) = p(3x-1) \implies 6x+18 = 18x-6$. Transposing x-terms gives $(6-18)x = -6-18 \implies x = \frac{-24}{-12}$.

Q3450. Moving denominators yields $q(4x+7) = p(5x-2) \implies 28x+49 = 10x-4$. Transposing x-terms gives $(28-10)x = -4-49 \implies x = \frac{-53}{18}$.

Q3451. Moving denominators yields $q(3x+5) = p(3x-2) \implies 9x+15 = 18x-12$. Transposing x-terms gives $(9-18)x = -12-15 \implies x = \frac{-27}{-9}$.

Q3452. Moving denominators yields $q(3x+8) = p(2x-3) \implies 6x+16 = 14x-21$. Transposing x-terms gives $(6-14)x = -21-16 \implies x = \frac{-37}{-8}$.

Q3453. Moving denominators yields $q(5x+7) = p(2x-6) \implies 15x+21 = 8x-24$. Transposing x-terms gives $(15-8)x = -24-21 \implies x = \frac{-45}{7}$.

Q3454. Moving denominators yields $q(4x+3) = p(5x-7) \implies 16x+12 = 20x-28$. Transposing x-terms gives $(16-20)x = -28-12 \implies x = \frac{-40}{-4}$.

Q3455. Moving denominators yields $q(2x+1) = p(4x-9) \implies 12x+6 = 8x-18$. Transposing x-terms gives $(12-8)x = -18-6 \implies x = \frac{-24}{4}$.

Q3456. Moving denominators yields $q(5x+6) = p(4x-2) \implies 35x+42 = 16x-8$. Transposing x-terms gives $(35-16)x = -8-42 \implies x = \frac{-50}{19}$.

Q3457. Moving denominators yields $q(3x+8) = p(2x-9) \implies 6x+16 = 10x-45$. Transposing x-terms gives $(6-10)x = -45-16 \implies x = \frac{-61}{-4}$.

Q3458. Moving denominators yields $q(3x+6) = p(2x-1) \implies 6x+12 = 6x-3$. Transposing x-terms gives $(6-6)x = -3-12 \implies x = \frac{-15}{0}$.

Q3459. Moving denominators yields $q(3x+7) = p(4x-3) \implies 18x+42 = 24x-18$. Transposing x-terms gives $(18-24)x = -18-42 \implies x = \frac{-60}{-6}$.

Q3460. Moving denominators yields $q(2x+9) = p(5x-9) \implies 14x+63 = 15x-27$. Transposing x-terms gives $(14-15)x = -27-63 \implies x = \frac{-90}{-1}$.

Q3461. Moving denominators yields $q(3x+3) = p(3x-6) \implies 9x+9 = 21x-42$. Transposing x-terms gives $(9-21)x = -42-9 \implies x = \frac{-51}{-12}$.

Q3462. Moving denominators yields $q(3x+9) = p(3x-7) \implies 9x+27 = 6x-14$. Transposing x-terms gives

$(9-6)x = -14-27 \implies x = \frac{-41}{3}$.

Q3463. Moving denominators yields $q(3x+8) = p(4x-2) \implies 6x+16 = 8x-4$. Transposing x-terms gives $(6-8)x = -4-16 \implies x = \frac{-20}{-2}$.

Q3464. Moving denominators yields $q(3x+6) = p(2x-6) \implies 6x+12 = 12x-36$. Transposing x-terms gives $(6-12)x = -36-12 \implies x = \frac{-48}{-6}$.

Q3465. Moving denominators yields $q(5x+6) = p(3x-9) \implies 15x+18 = 6x-18$. Transposing x-terms gives $(15-6)x = -18-18 \implies x = \frac{-36}{9}$.

Q3466. Moving denominators yields $q(5x+3) = p(5x-9) \implies 20x+12 = 35x-63$. Transposing x-terms gives $(20-35)x = -63-12 \implies x = \frac{-75}{-15}$.

Q3467. Moving denominators yields $q(4x+7) = p(3x-7) \implies 16x+28 = 12x-28$. Transposing x-terms gives $(16-12)x = -28-28 \implies x = \frac{-56}{4}$.

Q3468. Moving denominators yields $q(2x+7) = p(2x-3) \implies 4x+14 = 4x-6$. Transposing x-terms gives $(4-4)x = -6-14 \implies x = \frac{-20}{0}$.

Q3469. Moving denominators yields $q(3x+1) = p(5x-9) \implies 15x+5 = 30x-54$. Transposing x-terms gives $(15-30)x = -54-5 \implies x = \frac{-59}{-15}$.

Q3470. Moving denominators yields $q(5x+2) = p(5x-9) \implies 30x+12 = 20x-36$. Transposing x-terms gives $(30-20)x = -36-12 \implies x = \frac{-48}{10}$.

Q3471. Moving denominators yields $q(5x+3) = p(2x-6) \implies 25x+15 = 12x-36$. Transposing x-terms gives $(25-12)x = -36-15 \implies x = \frac{-51}{13}$.

Q3472. Moving denominators yields $q(5x+9) = p(5x-8) \implies 20x+36 = 10x-16$. Transposing x-terms gives $(20-10)x = -16-36 \implies x = \frac{-52}{10}$.

Q3473. Moving denominators yields $q(4x+7) = p(3x-9) \implies 16x+28 = 18x-54$. Transposing x-terms gives $(16-18)x = -54-28 \implies x = \frac{-82}{-2}$.

Q3474. Moving denominators yields $q(5x+4) = p(4x-1) \implies 25x+20 = 28x-7$. Transposing x-terms gives $(25-28)x = -7-20 \implies x = \frac{-27}{-3}$.

Q3475. Moving denominators yields $q(3x+5) = p(5x-5) \implies 6x+10 = 20x-20$. Transposing x-terms gives $(6-20)x = -20-10 \implies x = \frac{-30}{-14}$.

Q3476. Moving denominators yields $q(4x+4) = p(5x-4) \implies 12x+12 = 20x-16$. Transposing x-terms gives $(12-20)x = -16-12 \implies x = \frac{-28}{-8}$.

Q3477. Moving denominators yields $q(5x+2) = p(3x-8) \implies 25x+10 = 9x-24$. Transposing x-terms gives $(25-9)x = -24-10 \implies x = \frac{-34}{16}$.

Q3478. Moving denominators yields $q(4x+3) = p(4x-9) \implies 20x+15 = 8x-18$. Transposing x-terms gives $(20-8)x = -18-15 \implies x = \frac{-33}{12}$.

Q3479. Moving denominators yields $q(4x+5) = p(5x-2) \implies 20x+25 = 25x-10$. Transposing x-terms gives $(20-25)x = -10-25 \implies x = \frac{-35}{-5}$.

Q3480. Moving denominators yields $q(5x+4) = p(4x-7) \implies 15x+12 = 20x-35$. Transposing x-terms gives $(15-20)x = -35-12 \implies x = \frac{-47}{-5}$.

Q3481. Moving denominators yields $q(3x+9) = p(4x-6) \implies 18x+54 = 12x-18$. Transposing x-terms gives $(18-12)x = -18-54 \implies x = \frac{-72}{6}$.

Q3482. Moving denominators yields $q(5x+5) = p(2x-1) \implies 15x+15 = 10x-5$. Transposing x-terms gives $(15-10)x = -5-15 \implies x = \frac{-20}{5}$.

Q3483. Moving denominators yields $q(2x+5) = p(2x-3) \implies 12x+30 = 12x-18$. Transposing x-terms gives $(12-12)x = -18-30 \implies x = \frac{-48}{0}$.

Q3484. Moving denominators yields $q(4x+8) = p(5x-6) \implies 24x+48 = 35x-42$. Transposing x-terms gives $(24-35)x = -42-48 \implies x = \frac{-90}{-11}$.

Q3485. Moving denominators yields $q(5x+5) = p(2x-3) \implies 35x+35 = 14x-21$. Transposing x-terms gives $(35-14)x = -21-35 \implies x = \frac{-56}{21}$.

Q3486. Moving denominators yields $q(5x+6) = p(2x-4) \implies 25x+30 = 12x-24$. Transposing x-terms gives $(25-12)x = -24-30 \implies x = \frac{-54}{13}$.

Q3487. Moving denominators yields $q(4x+2) = p(5x-9) \implies 8x+4 = 35x-63$. Transposing x-terms gives $(8-35)x = -63-4 \implies x = \frac{-67}{-27}$.

Q3488. Moving denominators yields $q(3x+7) = p(2x-3) \implies 6x+14 = 4x-6$. Transposing x-terms gives $(6-4)x = -6-14 \implies x = \frac{-20}{2}$.

Q3489. Moving denominators yields $q(4x+1) = p(2x-5) \implies 16x+4 = 4x-10$. Transposing x-terms gives $(16-4)x = -10-4 \implies x = \frac{-14}{12}$.

Q3490. Moving denominators yields $q(5x+2) = p(4x-8) \implies 20x+8 = 24x-48$. Transposing x-terms gives $(20-24)x = -48-8 \implies x = \frac{-56}{-4}$.

Q3491. Moving denominators yields $q(3x+2) = p(5x-3) \implies 15x+10 = 25x-15$. Transposing x-terms gives $(15-25)x = -15-10 \implies x = \frac{-25}{-10}$.

Q3492. Moving denominators yields $q(4x+9) = p(4x-2) \implies 16x+36 = 28x-14$. Transposing x-terms gives $(16-28)x = -14-36 \implies x = \frac{-50}{-12}$.

Q3493. Moving denominators yields $q(3x+8) = p(4x-7) \implies 15x+40 = 28x-49$. Transposing x-terms gives $(15-28)x = -49-40 \implies x = \frac{-89}{-13}$.

Q3494. Moving denominators yields $q(2x+7) = p(4x-4) \implies 14x+49 = 20x-20$. Transposing x-terms gives $(14-20)x = -20-49 \implies x = \frac{-69}{-6}$.

Q3495. Moving denominators yields $q(4x+4) = p(3x-2) \implies 16x+16 = 21x-14$. Transposing x-terms gives $(16-21)x = -14-16 \implies x = \frac{-30}{-5}$.

Q3496. Moving denominators yields $q(3x+1) = p(4x-2) \implies 21x+7 = 8x-4$. Transposing x-terms gives $(21-8)x = -4-7 \implies x = \frac{-11}{13}$.

Q3497. Moving denominators yields $q(4x+7) = p(5x-1) \implies 20x+35 = 35x-7$. Transposing x-terms gives $(20-35)x = -7-35 \implies x = \frac{-42}{-15}$.

Q3498. Moving denominators yields $q(4x+1) = p(4x-7) \implies 28x+7 = 16x-28$. Transposing x-terms gives $(28-16)x = -28-7 \implies x = \frac{-35}{12}$.

Q3499. Moving denominators yields $q(3x+4) = p(3x-1) \implies 21x+28 = 9x-3$. Transposing x-terms gives $(21-9)x = -3-28 \implies x = \frac{-31}{12}$.

Q3500. Moving denominators yields $q(3x+3) = p(4x-8) \implies 18x+18 = 28x-56$. Transposing x-terms gives $(18-28)x = -56-18 \implies x = \frac{-74}{-10}$.

Q3501. The LCM of 5 and 4 is 20. Multiplying every

term by 20 gives: $4x + 5(x-1) = 200 \implies 9x - 5 = 200 \implies 9x = 205 \implies x = \frac{205}{9}$.

Q3502. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 20 \implies 7x - 2 = 20 \implies 7x = 22 \implies x = \frac{22}{7}$.

Q3503. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 350 \implies 12x - 5 = 350 \implies 12x = 355 \implies x = \frac{355}{12}$.

Q3504. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 72 \implies 10x - 4 = 72 \implies 10x = 76 \implies x = \frac{76}{10}$.

Q3505. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 140 \implies 9x - 4 = 140 \implies 9x = 144 \implies x = \frac{144}{9}$.

Q3506. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 60 \implies 9x - 5 = 60 \implies 9x = 65 \implies x = \frac{65}{9}$.

Q3507. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 36 \implies 7x - 3 = 36 \implies 7x = 39 \implies x = \frac{39}{7}$.

Q3508. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 144 \implies 10x - 4 = 144 \implies 10x = 148 \implies x = \frac{148}{10}$.

Q3509. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 75 \implies 8x - 5 = 75 \implies 8x = 80 \implies x = \frac{80}{8}$.

Q3510. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 168 \implies 11x - 4 = 168 \implies 11x = 172 \implies x = \frac{172}{11}$.

Q3511. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 60 \implies 7x - 3 = 60 \implies 7x = 63 \implies x = \frac{63}{7}$.

Q3512. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 120 \implies 7x - 3 = 120 \implies 7x = 123 \implies x = \frac{123}{7}$.

Q3513. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 180 \implies 11x - 5 = 180 \implies 11x = 185 \implies x = \frac{185}{11}$.

Q3514. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 162 \implies 9x - 3 = 162 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3515. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 40 \implies 6x - 2 = 40 \implies 6x = 42 \implies x = \frac{42}{6}$.

Q3516. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 200 \implies 9x - 4 = 200 \implies 9x = 204 \implies x = \frac{204}{9}$.

Q3517. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 84 \implies 8x - 2 = 84 \implies 8x = 86 \implies x = \frac{86}{8}$.

Q3518. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 48 \implies 8x - 2 = 48 \implies 8x = 50 \implies x = \frac{50}{8}$.

Q3519. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 120 \implies 8x - 5 = 120 \implies 8x = 125 \implies x = \frac{125}{8}$.

Q3520. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 120 \implies 8x - 2 = 120 \implies 8x = 122 \implies x = \frac{122}{8}$.

Q3521. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 216 \implies 10x - 4 = 216 \implies 10x = 220 \implies x = \frac{220}{10}$.

Q3522. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 45 \implies 8x - 5 = 45 \implies 8x = 50 \implies x = \frac{50}{8}$.

Q3523. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 168 \implies 11x - 4 = 168 \implies 11x = 172 \implies x = \frac{172}{11}$.

Q3524. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 56 \implies 6x - 2 = 56 \implies 6x = 58 \implies x = \frac{58}{6}$.

Q3525. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 210 \implies 10x - 3 = 210 \implies 10x = 213 \implies x = \frac{213}{10}$.

Q3526. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 160 \implies 9x - 4 = 160 \implies 9x = 164 \implies x = \frac{164}{9}$.

Q3527. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 36 \implies 7x - 4 = 36 \implies 7x = 40 \implies x = \frac{40}{7}$.

Q3528. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 280 \implies 12x - 5 = 280 \implies 12x = 285 \implies x = \frac{285}{12}$.

Q3529. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 189 \implies 10x - 3 = 189 \implies 10x = 192 \implies x = \frac{192}{10}$.

Q3530. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 90 \implies 11x - 5 = 90 \implies 11x = 95 \implies x = \frac{95}{11}$.

Q3531. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 63 \implies 10x - 3 = 63 \implies 10x = 66 \implies x = \frac{66}{10}$.

Q3532. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 98 \implies 9x - 2 = 98 \implies 9x = 100 \implies x = \frac{100}{9}$.

Q3533. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 120 \implies 7x - 3 = 120 \implies 7x = 123 \implies x = \frac{123}{7}$.

Q3534. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 180 \implies 9x - 3 = 180 \implies 9x = 183 \implies x = \frac{183}{9}$.

Q3535. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 147 \implies 10x - 3 = 147 \implies 10x = 150 \implies x = \frac{150}{10}$.

Q3536. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 160 \implies 9x - 4 = 160 \implies 9x = 164 \implies x = \frac{164}{9}$.

Q3537. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 54 \implies 9x - 3 = 54 \implies 9x = 57 \implies x = \frac{57}{9}$.

Q3538. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 56 \implies 11x - 4 = 56 \implies 11x = 60 \implies x = \frac{60}{11}$.

Q3539. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 32 \implies 6x - 2 = 32 \implies 6x = 34 \implies x = \frac{34}{6}$.

Q3540. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 60 \implies 11x - 5 = 60 \implies 11x = 65 \implies x = \frac{65}{11}$.

Q3541. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 270 \implies 11x - 5 = 270 \implies 11x = 275 \implies x = \frac{275}{11}$.

Q3542. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 48 \implies 10x - 4 = 48 \implies 10x = 52 \implies x = \frac{52}{10}$.

Q3543. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x+5(x-1) = 75 \implies 8x-5 = 75 \implies 8x = 80 \implies x = \frac{80}{8}$.

Q3544. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1) = 60 \implies 9x-4 = 60 \implies 9x = 64 \implies x = \frac{64}{9}$.

Q3545. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 280 \implies 11x - 4 = 280 \implies 11x = 284 \implies x = \frac{284}{11}$.

Q3546. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1) = 48 \implies 7x-4 = 48 \implies 7x = 52 \implies x = \frac{52}{7}$.

Q3547. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 180 \implies 9x - 5 = 180 \implies 9x = 185 \implies x = \frac{185}{9}$.

Q3548. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 120 \implies 8x - 5 = 120 \implies 8x = 125 \implies x = \frac{125}{8}$.

Q3549. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 160 \implies 9x - 4 = 160 \implies 9x = 164 \implies x = \frac{164}{9}$.

Q3550. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1) = 45 \implies 8x-3 = 45 \implies 8x = 48 \implies x = \frac{48}{8}$.

Q3551. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 240 \implies 11x - 5 = 240 \implies 11x = 245 \implies x = \frac{245}{11}$.

Q3552. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 224 \implies 11x - 4 = 224 \implies 11x = 228 \implies x = \frac{228}{11}$.

Q3553. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1) = 80 \implies 9x-4 = 80 \implies 9x = 84 \implies x = \frac{84}{9}$.

Q3554. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1) = 24 \implies 6x-2 = 24 \implies 6x = 26 \implies x = \frac{26}{6}$.

Q3555. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1) = 48 \implies 7x-4 = 48 \implies 7x = 52 \implies x = \frac{52}{7}$.

Q3556. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x+2(x-1) = 98 \implies 9x-2 = 98 \implies 9x = 100 \implies x = \frac{100}{9}$.

Q3557. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 56 \implies 11x - 4 = 56 \implies 11x = 60 \implies x = \frac{60}{11}$.

Q3558. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1) = 32 \implies 6x-2 = 32 \implies 6x = 34 \implies x = \frac{34}{6}$.

Q3559. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 120 \implies 11x - 5 = 120 \implies 11x = 125 \implies x = \frac{125}{11}$.

Q3560. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x+3(x-1) = 48 \implies 7x-3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3561. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1) = 75 \implies 8x-3 = 75 \implies 8x = 78 \implies x = \frac{78}{8}$.

Q3562. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1) = 80 \implies 6x-2 = 80 \implies 6x = 82 \implies x = \frac{82}{6}$.

Q3563. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 84 \implies 10x - 3 = 84 \implies 10x = 87 \implies x = \frac{87}{10}$.

Q3564. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 120 \implies 9x - 4 = 120 \implies 9x = 124 \implies x = \frac{124}{9}$.

Q3565. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 200 \implies 9x - 4 = 200 \implies 9x = 204 \implies x = \frac{204}{9}$.

Q3566. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 120 \implies 7x - 3 = 120 \implies 7x = 123 \implies x = \frac{123}{7}$.

Q3567. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1) = 45 \implies 8x-3 = 45 \implies 8x = 48 \implies x = \frac{48}{8}$.

Q3568. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1) = 16 \implies 6x-2 = 16 \implies 6x = 18 \implies x = \frac{18}{6}$.

Q3569. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 160 \implies 9x - 4 = 160 \implies 9x = 164 \implies x = \frac{164}{9}$.

Q3570. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1) = 32 \implies 6x-2 = 32 \implies 6x = 34 \implies x = \frac{34}{6}$.

Q3571. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x+5(x-1) = 80 \implies 9x-5 = 80 \implies 9x = 85 \implies x = \frac{85}{9}$.

Q3572. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 210 \implies 11x - 5 = 210 \implies 11x = 215 \implies x = \frac{215}{11}$.

Q3573. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x+2(x-1) = 42 \implies 9x-2 = 42 \implies 9x = 44 \implies x = \frac{44}{9}$.

Q3574. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 96 \implies 10x - 4 = 96 \implies 10x = 100 \implies x = \frac{100}{10}$.

Q3575. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x+2(x-1) = 84 \implies 9x-2 = 84 \implies 9x = 86 \implies x = \frac{86}{9}$.

Q3576. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 70 \implies 12x - 5 = 70 \implies 12x = 75 \implies x = \frac{75}{12}$.

Q3577. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x+2(x-1) = 36 \implies 8x-2 = 36 \implies 8x = 38 \implies x = \frac{38}{8}$.

Q3578. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 108 \implies 7x - 3 =$

$108 \implies 7x = 111 \implies x = \frac{111}{7}$.

Q3579. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 150 \implies 11x - 5 = 150 \implies 11x = 155 \implies x = \frac{155}{11}$.

Q3580. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 45 \implies 8x - 3 = 45 \implies 8x = 48 \implies x = \frac{48}{8}$.

Q3581. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 144 \implies 9x - 3 = 144 \implies 9x = 147 \implies x = \frac{147}{9}$.

Q3582. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 150 \implies 11x - 5 = 150 \implies 11x = 155 \implies x = \frac{155}{11}$.

Q3583. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 160 \implies 9x - 5 = 160 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3584. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 96 \implies 10x - 4 = 96 \implies 10x = 100 \implies x = \frac{100}{10}$.

Q3585. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 48 \implies 7x - 4 = 48 \implies 7x = 52 \implies x = \frac{52}{7}$.

Q3586. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 48 \implies 5x - 2 = 48 \implies 5x = 50 \implies x = \frac{50}{5}$.

Q3587. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 126 \implies 9x - 2 = 126 \implies 9x = 128 \implies x = \frac{128}{9}$.

Q3588. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 72 \implies 6x - 2 = 72 \implies 6x = 74 \implies x = \frac{74}{6}$.

Q3589. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 300 \implies 11x - 5 = 300 \implies 11x = 305 \implies x = \frac{305}{11}$.

Q3590. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 48 \implies 7x - 3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3591. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 36 \implies 7x - 3 = 36 \implies 7x = 39 \implies x = \frac{39}{7}$.

Q3592. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 48 \implies 7x - 3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3593. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 48 \implies 7x - 4 = 48 \implies 7x = 52 \implies x = \frac{52}{7}$.

Q3594. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 150 \implies 11x - 5 = 150 \implies 11x = 155 \implies x = \frac{155}{11}$.

Q3595. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 45 \implies 8x - 3 = 45 \implies 8x = 48 \implies x = \frac{48}{8}$.

Q3596. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 180 \implies 9x - 4 = 180 \implies 9x = 184 \implies x = \frac{184}{9}$.

Q3597. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 90 \implies 8x - 5 = 90 \implies 8x = 95 \implies x = \frac{95}{8}$.

Q3598. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 56 \implies 9x - 2 = 56 \implies 9x = 58 \implies x = \frac{58}{9}$.

Q3599. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 54 \implies 5x - 2 = 54 \implies 5x = 56 \implies x = \frac{56}{5}$.

Q3600. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 270 \implies 11x - 5 = 270 \implies 11x = 275 \implies x = \frac{275}{11}$.

Q3601. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 90 \implies 11x - 5 = 90 \implies 11x = 95 \implies x = \frac{95}{11}$.

Q3602. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 216 \implies 10x - 4 = 216 \implies 10x = 220 \implies x = \frac{220}{10}$.

Q3603. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 80 \implies 9x - 4 = 80 \implies 9x = 84 \implies x = \frac{84}{9}$.

Q3604. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 96 \implies 10x - 4 = 96 \implies 10x = 100 \implies x = \frac{100}{10}$.

Q3605. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 60 \implies 8x - 3 = 60 \implies 8x = 63 \implies x = \frac{63}{8}$.

Q3606. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 120 \implies 9x - 5 = 120 \implies 9x = 125 \implies x = \frac{125}{9}$.

Q3607. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 48 \implies 7x - 3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3608. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 28 \implies 9x - 2 = 28 \implies 9x = 30 \implies x = \frac{30}{9}$.

Q3609. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 160 \implies 9x - 5 = 160 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3610. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 135 \implies 8x - 3 = 135 \implies 8x = 138 \implies x = \frac{138}{8}$.

Q3611. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 80 \implies 7x - 2 = 80 \implies 7x = 82 \implies x = \frac{82}{7}$.

Q3612. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 30 \implies 8x - 5 = 30 \implies 8x = 35 \implies x = \frac{35}{8}$.

Q3613. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 70 \implies 7x - 2 = 70 \implies 7x = 72 \implies x = \frac{72}{7}$.

Q3614. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 48 \implies 7x - 4 = 48 \implies 7x = 52 \implies x = \frac{52}{7}$.

Q3615. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 84 \implies 9x - 2 = 84 \implies 9x = 86 \implies x = \frac{86}{9}$.

Q3616. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 63 \implies 10x - 3 = 63 \implies 10x = 66 \implies x = \frac{66}{10}$.

Q3617. The LCM of 4 and 5 is 20. Multiplying every

term by 20 gives: $5x+4(x-1)=60 \implies 9x-4=60 \implies 9x=64 \implies x=\frac{64}{9}$.

Q3618. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1)=24 \implies 7x-4=24 \implies 7x=28 \implies x=\frac{28}{7}$.

Q3619. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x+2(x-1)=56 \implies 9x-2=56 \implies 9x=58 \implies x=\frac{58}{9}$.

Q3620. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x+2(x-1)=20 \implies 7x-2=20 \implies 7x=22 \implies x=\frac{22}{7}$.

Q3621. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x+3(x-1)=63 \implies 10x-3=63 \implies 10x=66 \implies x=\frac{66}{10}$.

Q3622. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=160 \implies 9x-4=160 \implies 9x=164 \implies x=\frac{164}{9}$.

Q3623. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=180 \implies 9x-4=180 \implies 9x=184 \implies x=\frac{184}{9}$.

Q3624. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1)=108 \implies 7x-4=108 \implies 7x=112 \implies x=\frac{112}{7}$.

Q3625. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x+2(x-1)=50 \implies 7x-2=50 \implies 7x=52 \implies x=\frac{52}{7}$.

Q3626. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x+5(x-1)=105 \implies 8x-5=105 \implies 8x=110 \implies x=\frac{110}{8}$.

Q3627. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=160 \implies 9x-4=160 \implies 9x=164 \implies x=\frac{164}{9}$.

Q3628. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x+5(x-1)=30 \implies 8x-5=30 \implies 8x=35 \implies x=\frac{35}{8}$.

Q3629. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x+3(x-1)=168 \implies 10x-3=168 \implies 10x=171 \implies x=\frac{171}{10}$.

Q3630. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x+3(x-1)=24 \implies 7x-3=24 \implies 7x=27 \implies x=\frac{27}{7}$.

Q3631. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x+2(x-1)=24 \implies 5x-2=24 \implies 5x=26 \implies x=\frac{26}{5}$.

Q3632. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=100 \implies 9x-4=100 \implies 9x=104 \implies x=\frac{104}{9}$.

Q3633. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x+3(x-1)=24 \implies 7x-3=24 \implies 7x=27 \implies x=\frac{27}{7}$.

Q3634. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x+5(x-1)=210 \implies 11x-5=210 \implies 11x=215 \implies x=\frac{215}{11}$.

Q3635. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x+5(x-1)=315 \implies 12x-5=315 \implies 12x=320 \implies x=\frac{320}{12}$.

Q3636. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1)=84 \implies 7x-4=84 \implies 7x=88 \implies x=\frac{88}{7}$.

Q3637. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x+2(x-1)=60 \implies 7x-2=60 \implies 7x=62 \implies x=\frac{62}{7}$.

Q3638. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1)=90 \implies 8x-3=90 \implies 8x=93 \implies x=\frac{93}{8}$.

Q3639. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x+3(x-1)=72 \implies 7x-3=72 \implies 7x=75 \implies x=\frac{75}{7}$.

Q3640. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=60 \implies 9x-4=60 \implies 9x=64 \implies x=\frac{64}{9}$.

Q3641. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x+2(x-1)=70 \implies 9x-2=70 \implies 9x=72 \implies x=\frac{72}{9}$.

Q3642. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1)=90 \implies 8x-3=90 \implies 8x=93 \implies x=\frac{93}{8}$.

Q3643. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x+3(x-1)=72 \implies 7x-3=72 \implies 7x=75 \implies x=\frac{75}{7}$.

Q3644. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x+3(x-1)=60 \implies 8x-3=60 \implies 8x=63 \implies x=\frac{63}{8}$.

Q3645. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1)=80 \implies 6x-2=80 \implies 6x=82 \implies x=\frac{82}{6}$.

Q3646. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x+2(x-1)=108 \implies 8x-2=108 \implies 8x=110 \implies x=\frac{110}{8}$.

Q3647. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=160 \implies 9x-4=160 \implies 9x=164 \implies x=\frac{164}{9}$.

Q3648. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x+4(x-1)=40 \implies 9x-4=40 \implies 9x=44 \implies x=\frac{44}{9}$.

Q3649. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x+3(x-1)=105 \implies 10x-3=105 \implies 10x=108 \implies x=\frac{108}{10}$.

Q3650. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x+4(x-1)=280 \implies 11x-4=280 \implies 11x=284 \implies x=\frac{284}{11}$.

Q3651. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x+2(x-1)=64 \implies 6x-2=64 \implies 6x=66 \implies x=\frac{66}{6}$.

Q3652. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x+2(x-1)=24 \implies 5x-2=24 \implies 5x=26 \implies x=\frac{26}{5}$.

Q3653. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x+5(x-1)=120 \implies 8x-5=120 \implies 8x=125 \implies x=\frac{125}{8}$.

Q3654. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1)=108 \implies 7x-4=108 \implies 7x=112 \implies x=\frac{112}{7}$.

Q3655. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x+4(x-1)=84 \implies 7x-4=84 \implies 7x=88 \implies x=\frac{88}{7}$.

Q3656. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 140 \implies 9x - 5 = 140 \implies 9x = 145 \implies x = \frac{145}{9}$.

Q3657. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 30 \implies 7x - 2 = 30 \implies 7x = 32 \implies x = \frac{32}{7}$.

Q3658. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 100 \implies 9x - 4 = 100 \implies 9x = 104 \implies x = \frac{104}{9}$.

Q3659. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 144 \implies 9x - 3 = 144 \implies 9x = 147 \implies x = \frac{147}{9}$.

Q3660. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 162 \implies 9x - 3 = 162 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3661. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 36 \implies 9x - 3 = 36 \implies 9x = 39 \implies x = \frac{39}{9}$.

Q3662. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 180 \implies 9x - 5 = 180 \implies 9x = 185 \implies x = \frac{185}{9}$.

Q3663. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 60 \implies 7x - 3 = 60 \implies 7x = 63 \implies x = \frac{63}{7}$.

Q3664. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 126 \implies 9x - 3 = 126 \implies 9x = 129 \implies x = \frac{129}{9}$.

Q3665. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 12 \implies 5x - 2 = 12 \implies 5x = 14 \implies x = \frac{14}{5}$.

Q3666. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 140 \implies 11x - 4 = 140 \implies 11x = 144 \implies x = \frac{144}{11}$.

Q3667. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 84 \implies 10x - 3 = 84 \implies 10x = 87 \implies x = \frac{87}{10}$.

Q3668. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 252 \implies 11x - 4 = 252 \implies 11x = 256 \implies x = \frac{256}{11}$.

Q3669. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 252 \implies 11x - 4 = 252 \implies 11x = 256 \implies x = \frac{256}{11}$.

Q3670. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 72 \implies 8x - 2 = 72 \implies 8x = 74 \implies x = \frac{74}{8}$.

Q3671. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 135 \implies 8x - 3 = 135 \implies 8x = 138 \implies x = \frac{138}{8}$.

Q3672. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 75 \implies 8x - 5 = 75 \implies 8x = 80 \implies x = \frac{80}{8}$.

Q3673. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 120 \implies 10x - 4 = 120 \implies 10x = 124 \implies x = \frac{124}{10}$.

Q3674. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 18 \implies 5x - 2 = 18 \implies 5x = 20 \implies x = \frac{20}{5}$.

Q3675. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 180 \implies 9x - 4 = 180 \implies 9x = 184 \implies x = \frac{184}{9}$.

Q3676. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 96 \implies 7x - 3 = 96 \implies 7x = 99 \implies x = \frac{99}{7}$.

Q3677. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 96 \implies 10x - 4 = 96 \implies 10x = 100 \implies x = \frac{100}{10}$.

Q3678. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 162 \implies 9x - 3 = 162 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3679. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 140 \implies 12x - 5 = 140 \implies 12x = 145 \implies x = \frac{145}{12}$.

Q3680. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 180 \implies 9x - 3 = 180 \implies 9x = 183 \implies x = \frac{183}{9}$.

Q3681. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 160 \implies 9x - 4 = 160 \implies 9x = 164 \implies x = \frac{164}{9}$.

Q3682. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 160 \implies 9x - 5 = 160 \implies 9x = 165 \implies x = \frac{165}{9}$.

Q3683. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 245 \implies 12x - 5 = 245 \implies 12x = 250 \implies x = \frac{250}{12}$.

Q3684. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 200 \implies 9x - 4 = 200 \implies 9x = 204 \implies x = \frac{204}{9}$.

Q3685. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 112 \implies 11x - 4 = 112 \implies 11x = 116 \implies x = \frac{116}{11}$.

Q3686. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 120 \implies 9x - 4 = 120 \implies 9x = 124 \implies x = \frac{124}{9}$.

Q3687. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 36 \implies 5x - 2 = 36 \implies 5x = 38 \implies x = \frac{38}{5}$.

Q3688. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 270 \implies 11x - 5 = 270 \implies 11x = 275 \implies x = \frac{275}{11}$.

Q3689. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 120 \implies 10x - 4 = 120 \implies 10x = 124 \implies x = \frac{124}{10}$.

Q3690. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 72 \implies 7x - 3 = 72 \implies 7x = 75 \implies x = \frac{75}{7}$.

Q3691. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 210 \implies 11x - 5 = 210 \implies 11x = 215 \implies x = \frac{215}{11}$.

Q3692. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 135 \implies 8x - 5 = 135 \implies 8x = 140 \implies x = \frac{140}{8}$.

Q3693. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 108 \implies 7x - 3 = 108 \implies 7x = 111 \implies x = \frac{111}{7}$.

Q3694. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 30 \implies 7x - 2 = 30 \implies$

$7x = 32 \implies x = \frac{32}{7}$.

Q3695. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 42 \implies 10x - 3 = 42 \implies 10x = 45 \implies x = \frac{45}{10}$.

Q3696. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 60 \implies 5x - 2 = 60 \implies 5x = 62 \implies x = \frac{62}{5}$.

Q3697. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 72 \implies 7x - 3 = 72 \implies 7x = 75 \implies x = \frac{75}{7}$.

Q3698. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 36 \implies 7x - 3 = 36 \implies 7x = 39 \implies x = \frac{39}{7}$.

Q3699. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 90 \implies 8x - 3 = 90 \implies 8x = 93 \implies x = \frac{93}{8}$.

Q3700. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 105 \implies 8x - 5 = 105 \implies 8x = 110 \implies x = \frac{110}{8}$.

Q3701. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 240 \implies 10x - 4 = 240 \implies 10x = 244 \implies x = \frac{244}{10}$.

Q3702. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 120 \implies 9x - 4 = 120 \implies 9x = 124 \implies x = \frac{124}{9}$.

Q3703. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 80 \implies 9x - 5 = 80 \implies 9x = 85 \implies x = \frac{85}{9}$.

Q3704. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 72 \implies 10x - 4 = 72 \implies 10x = 76 \implies x = \frac{76}{10}$.

Q3705. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 105 \implies 12x - 5 = 105 \implies 12x = 110 \implies x = \frac{110}{12}$.

Q3706. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 48 \implies 7x - 3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3707. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 32 \implies 6x - 2 = 32 \implies 6x = 34 \implies x = \frac{34}{6}$.

Q3708. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 280 \implies 11x - 4 = 280 \implies 11x = 284 \implies x = \frac{284}{11}$.

Q3709. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 45 \implies 8x - 5 = 45 \implies 8x = 50 \implies x = \frac{50}{8}$.

Q3710. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 112 \implies 11x - 4 = 112 \implies 11x = 116 \implies x = \frac{116}{11}$.

Q3711. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 64 \implies 6x - 2 = 64 \implies 6x = 66 \implies x = \frac{66}{6}$.

Q3712. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 189 \implies 10x - 3 = 189 \implies 10x = 192 \implies x = \frac{192}{10}$.

Q3713. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 70 \implies 7x - 2 = 70 \implies 7x = 72 \implies x = \frac{72}{7}$.

Q3714. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 96 \implies 8x - 2 = 96 \implies 8x = 98 \implies x = \frac{98}{8}$.

Q3715. The LCM of 2 and 3 is 6. Multiplying every term by 6 gives: $3x + 2(x-1) = 18 \implies 5x - 2 = 18 \implies 5x = 20 \implies x = \frac{20}{5}$.

Q3716. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 120 \implies 7x - 3 = 120 \implies 7x = 123 \implies x = \frac{123}{7}$.

Q3717. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 150 \implies 8x - 5 = 150 \implies 8x = 155 \implies x = \frac{155}{8}$.

Q3718. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 60 \implies 8x - 5 = 60 \implies 8x = 65 \implies x = \frac{65}{8}$.

Q3719. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 315 \implies 12x - 5 = 315 \implies 12x = 320 \implies x = \frac{320}{12}$.

Q3720. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 180 \implies 9x - 4 = 180 \implies 9x = 184 \implies x = \frac{184}{9}$.

Q3721. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 48 \implies 10x - 4 = 48 \implies 10x = 52 \implies x = \frac{52}{10}$.

Q3722. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 80 \implies 9x - 5 = 80 \implies 9x = 85 \implies x = \frac{85}{9}$.

Q3723. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 98 \implies 9x - 2 = 98 \implies 9x = 100 \implies x = \frac{100}{9}$.

Q3724. The LCM of 2 and 4 is 8. Multiplying every term by 8 gives: $4x + 2(x-1) = 40 \implies 6x - 2 = 40 \implies 6x = 42 \implies x = \frac{42}{6}$.

Q3725. The LCM of 5 and 4 is 20. Multiplying every term by 20 gives: $4x + 5(x-1) = 120 \implies 9x - 5 = 120 \implies 9x = 125 \implies x = \frac{125}{9}$.

Q3726. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 36 \implies 8x - 2 = 36 \implies 8x = 38 \implies x = \frac{38}{8}$.

Q3727. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 120 \implies 11x - 5 = 120 \implies 11x = 125 \implies x = \frac{125}{11}$.

Q3728. The LCM of 2 and 5 is 10. Multiplying every term by 10 gives: $5x + 2(x-1) = 40 \implies 7x - 2 = 40 \implies 7x = 42 \implies x = \frac{42}{7}$.

Q3729. The LCM of 4 and 6 is 24. Multiplying every term by 24 gives: $6x + 4(x-1) = 192 \implies 10x - 4 = 192 \implies 10x = 196 \implies x = \frac{196}{10}$.

Q3730. The LCM of 5 and 3 is 15. Multiplying every term by 15 gives: $3x + 5(x-1) = 120 \implies 8x - 5 = 120 \implies 8x = 125 \implies x = \frac{125}{8}$.

Q3731. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 280 \implies 12x - 5 = 280 \implies 12x = 285 \implies x = \frac{285}{12}$.

Q3732. The LCM of 3 and 6 is 18. Multiplying every term by 18 gives: $6x + 3(x-1) = 54 \implies 9x - 3 = 54 \implies 9x = 57 \implies x = \frac{57}{9}$.

Q3733. The LCM of 4 and 6 is 24. Multiplying every

term by 24 gives: $6x + 4(x-1) = 48 \implies 10x - 4 = 48 \implies 10x = 52 \implies x = \frac{52}{10}$.

Q3734. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 240 \implies 11x - 5 = 240 \implies 11x = 245 \implies x = \frac{245}{11}$.

Q3735. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 60 \implies 9x - 4 = 60 \implies 9x = 64 \implies x = \frac{64}{9}$.

Q3736. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 60 \implies 7x - 3 = 60 \implies 7x = 63 \implies x = \frac{63}{7}$.

Q3737. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 42 \implies 10x - 3 = 42 \implies 10x = 45 \implies x = \frac{45}{10}$.

Q3738. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 84 \implies 7x - 4 = 84 \implies 7x = 88 \implies x = \frac{88}{7}$.

Q3739. The LCM of 5 and 7 is 35. Multiplying every term by 35 gives: $7x + 5(x-1) = 175 \implies 12x - 5 = 175 \implies 12x = 180 \implies x = \frac{180}{12}$.

Q3740. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 80 \implies 9x - 4 = 80 \implies 9x = 84 \implies x = \frac{84}{9}$.

Q3741. The LCM of 5 and 6 is 30. Multiplying every term by 30 gives: $6x + 5(x-1) = 60 \implies 11x - 5 = 60 \implies 11x = 65 \implies x = \frac{65}{11}$.

Q3742. The LCM of 2 and 7 is 14. Multiplying every term by 14 gives: $7x + 2(x-1) = 70 \implies 9x - 2 = 70 \implies 9x = 72 \implies x = \frac{72}{9}$.

Q3743. The LCM of 2 and 6 is 12. Multiplying every term by 12 gives: $6x + 2(x-1) = 60 \implies 8x - 2 = 60 \implies 8x = 62 \implies x = \frac{62}{8}$.

Q3744. The LCM of 3 and 5 is 15. Multiplying every term by 15 gives: $5x + 3(x-1) = 105 \implies 8x - 3 = 105 \implies 8x = 108 \implies x = \frac{108}{8}$.

Q3745. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 140 \implies 11x - 4 = 140 \implies 11x = 144 \implies x = \frac{144}{11}$.

Q3746. The LCM of 3 and 4 is 12. Multiplying every term by 12 gives: $4x + 3(x-1) = 48 \implies 7x - 3 = 48 \implies 7x = 51 \implies x = \frac{51}{7}$.

Q3747. The LCM of 4 and 5 is 20. Multiplying every term by 20 gives: $5x + 4(x-1) = 180 \implies 9x - 4 = 180 \implies 9x = 184 \implies x = \frac{184}{9}$.

Q3748. The LCM of 3 and 7 is 21. Multiplying every term by 21 gives: $7x + 3(x-1) = 126 \implies 10x - 3 = 126 \implies 10x = 129 \implies x = \frac{129}{10}$.

Q3749. The LCM of 4 and 3 is 12. Multiplying every term by 12 gives: $3x + 4(x-1) = 60 \implies 7x - 4 = 60 \implies 7x = 64 \implies x = \frac{64}{7}$.

Q3750. The LCM of 4 and 7 is 28. Multiplying every term by 28 gives: $7x + 4(x-1) = 280 \implies 11x - 4 = 280 \implies 11x = 284 \implies x = \frac{284}{11}$.

Q3751. The denominator $x - 3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x + 3 = 23 \implies x = 3$. However, substituting $x = 3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 3$ is an **extraneous root** and the equation has no solution.

Q3752. The denominator $x - 3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x + 3 = 23 \implies x = 3$. However, substituting $x = 3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 3$ is an **extraneous root** and the equation has no solution.

Q3753. The denominator $x - 3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x + 3 = 23 \implies x = 3$. However, substituting $x = 3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 3$ is an **extraneous root** and the equation has no solution.

Q3754. The denominator $x - 6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x + 6 = 26 \implies x = 6$. However, substituting $x = 6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 6$ is an **extraneous root** and the equation has no solution.

Q3755. The denominator $x - 8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x + 8 = 28 \implies x = 8$. However, substituting $x = 8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 8$ is an **extraneous root** and the equation has no solution.

Q3756. The denominator $x - 2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x + 2 = 22 \implies x = 2$. However, substituting $x = 2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 2$ is an **extraneous root** and the equation has no solution.

Q3757. The denominator $x - 4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x + 4 = 24 \implies x = 4$. However, substituting $x = 4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 4$ is an **extraneous root** and the equation has no solution.

Q3758. The denominator $x - 5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x + 5 = 25 \implies x = 5$. However, substituting $x = 5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 5$ is an **extraneous root** and the equation has no solution.

Q3759. The denominator $x - 3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x + 3 = 23 \implies x = 3$. However, substituting $x = 3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 3$ is an **extraneous root** and the equation has no solution.

Q3760. The denominator $x - 4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x + 4 = 24 \implies x = 4$. However, substituting $x = 4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 4$ is an **extraneous root** and the equation has no solution.

Q3761. The denominator $x - 5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x + 5 = 25 \implies x = 5$. However, substituting $x = 5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x = 5$ is an **extraneous root** and the equation

has no solution.

Q3762. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3763. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3764. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3765. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3766. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3767. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3768. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3769. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3770. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3771. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3772. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3773. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3774. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3775. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3776. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3777. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3778. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3779. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3780. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3781. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation

has no solution.

Q3782. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3783. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3784. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3785. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3786. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3787. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3788. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3789. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3790. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3791. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3792. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3793. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3794. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3795. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3796. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3797. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3798. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3799. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3800. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3801. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation

has no solution.

Q3802. The denominator $x-4$ cannot be zero, so the domain restriction is **$x \neq 4$**. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3803. The denominator $x-5$ cannot be zero, so the domain restriction is **$x \neq 5$**. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3804. The denominator $x-1$ cannot be zero, so the domain restriction is **$x \neq 1$**. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3805. The denominator $x-5$ cannot be zero, so the domain restriction is **$x \neq 5$**. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3806. The denominator $x-5$ cannot be zero, so the domain restriction is **$x \neq 5$**. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3807. The denominator $x-7$ cannot be zero, so the domain restriction is **$x \neq 7$**. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3808. The denominator $x-2$ cannot be zero, so the domain restriction is **$x \neq 2$**. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3809. The denominator $x-7$ cannot be zero, so the domain restriction is **$x \neq 7$**. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3810. The denominator $x-9$ cannot be zero, so the domain restriction is **$x \neq 9$**. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3811. The denominator $x-6$ cannot be zero, so the domain restriction is **$x \neq 6$**. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3812. The denominator $x-7$ cannot be zero, so the domain restriction is **$x \neq 7$**. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3813. The denominator $x-5$ cannot be zero, so the domain restriction is **$x \neq 5$**. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3814. The denominator $x-1$ cannot be zero, so the domain restriction is **$x \neq 1$**. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3815. The denominator $x-8$ cannot be zero, so the domain restriction is **$x \neq 8$**. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3816. The denominator $x-4$ cannot be zero, so the domain restriction is **$x \neq 4$**. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3817. The denominator $x-9$ cannot be zero, so the domain restriction is **$x \neq 9$**. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3818. The denominator $x-4$ cannot be zero, so the domain restriction is **$x \neq 4$**. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3819. The denominator $x-2$ cannot be zero, so the domain restriction is **$x \neq 2$**. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3820. The denominator $x-3$ cannot be zero, so the domain restriction is **$x \neq 3$**. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3821. The denominator $x-5$ cannot be zero, so the domain restriction is **$x \neq 5$**. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation

has no solution.

Q3822. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3823. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3824. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3825. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3826. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3827. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3828. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3829. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3830. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3831. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3832. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3833. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3834. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3835. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3836. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3837. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3838. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3839. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3840. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3841. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation

has no solution.

Q3842. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3843. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3844. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3845. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3846. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3847. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3848. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3849. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3850. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3851. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3852. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3853. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3854. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3855. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3856. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3857. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3858. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3859. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3860. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3861. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation

has no solution.

Q3862. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3863. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3864. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3865. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3866. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3867. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3868. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3869. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3870. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3871. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3872. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3873. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3874. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3875. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3876. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3877. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3878. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3879. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3880. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3881. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation

has no solution.

Q3882. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3883. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3884. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3885. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3886. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3887. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3888. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3889. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3890. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3891. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3892. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3893. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3894. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3895. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3896. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3897. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3898. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3899. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3900. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3901. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation

has no solution.

Q3902. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3903. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3904. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3905. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3906. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3907. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3908. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3909. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3910. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3911. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3912. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3913. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3914. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3915. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3916. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3917. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3918. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3919. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3920. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3921. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation

has no solution.

Q3922. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3923. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3924. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3925. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3926. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3927. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3928. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3929. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3930. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3931. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3932. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3933. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3934. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3935. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3936. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3937. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3938. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3939. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3940. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3941. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation

has no solution.

Q3942. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3943. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3944. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3945. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3946. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3947. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3948. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3949. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3950. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3951. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3952. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3953. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3954. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3955. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3956. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3957. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3958. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3959. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3960. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3961. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation

has no solution.

Q3962. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3963. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3964. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3965. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3966. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3967. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3968. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3969. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3970. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3971. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3972. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3973. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3974. The denominator $x-2$ cannot be zero, so the domain restriction is $**x \neq 2**$. Simplifying $\frac{(x-2)(x+2)}{x-2} = 22 \implies x+2=22 \implies x=2$. However, substituting $x=2$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=2$ is an **extraneous root** and the equation has no solution.

Q3975. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3976. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3977. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3978. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3979. The denominator $x-5$ cannot be zero, so the domain restriction is $**x \neq 5**$. Simplifying $\frac{(x-5)(x+5)}{x-5} = 25 \implies x+5=25 \implies x=5$. However, substituting $x=5$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=5$ is an **extraneous root** and the equation has no solution.

Q3980. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3981. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation

has no solution.

Q3982. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3983. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3984. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3985. The denominator $x-7$ cannot be zero, so the domain restriction is $**x \neq 7**$. Simplifying $\frac{(x-7)(x+7)}{x-7} = 27 \implies x+7=27 \implies x=7$. However, substituting $x=7$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=7$ is an **extraneous root** and the equation has no solution.

Q3986. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3987. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3988. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3989. The denominator $x-6$ cannot be zero, so the domain restriction is $**x \neq 6**$. Simplifying $\frac{(x-6)(x+6)}{x-6} = 26 \implies x+6=26 \implies x=6$. However, substituting $x=6$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=6$ is an **extraneous root** and the equation has no solution.

Q3990. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3991. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3992. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3993. The denominator $x-3$ cannot be zero, so the domain restriction is $**x \neq 3**$. Simplifying $\frac{(x-3)(x+3)}{x-3} = 23 \implies x+3=23 \implies x=3$. However, substituting $x=3$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=3$ is an **extraneous root** and the equation has no solution.

Q3994. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q3995. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3996. The denominator $x-1$ cannot be zero, so the domain restriction is $**x \neq 1**$. Simplifying $\frac{(x-1)(x+1)}{x-1} = 21 \implies x+1=21 \implies x=1$. However, substituting $x=1$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=1$ is an **extraneous root** and the equation has no solution.

Q3997. The denominator $x-9$ cannot be zero, so the domain restriction is $**x \neq 9**$. Simplifying $\frac{(x-9)(x+9)}{x-9} = 29 \implies x+9=29 \implies x=9$. However, substituting $x=9$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=9$ is an **extraneous root** and the equation has no solution.

Q3998. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.

Q3999. The denominator $x-8$ cannot be zero, so the domain restriction is $**x \neq 8**$. Simplifying $\frac{(x-8)(x+8)}{x-8} = 28 \implies x+8=28 \implies x=8$. However, substituting $x=8$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=8$ is an **extraneous root** and the equation has no solution.

Q4000. The denominator $x-4$ cannot be zero, so the domain restriction is $**x \neq 4**$. Simplifying $\frac{(x-4)(x+4)}{x-4} = 24 \implies x+4=24 \implies x=4$. However, substituting $x=4$ into the original equation creates $\frac{0}{0}$ (undefined). Thus $x=4$ is an **extraneous root** and the equation has no solution.