

# Subspaces

---

Nachiketa Mishra  
IIITDM Kancheepuram, Chennai

# Subspace

Let  $V$  be a vector space over a field  $F$ .

# Subspace

Let  $V$  be a vector space over a field  $F$ . A subspace of  $V$  is a subset  $W$  of  $V$

# Subspace


Let  $V$  be a vector space over a field  $F$ . A subspace of  $V$  is a subset  $W$  of  $V$  which is itself a vector space over  $F$

# Subspace

Let  $V$  be a vector space over a field  $F$ . A **subspace** of  $V$  is a **subset**  $W$  of  $V$  which is itself a vector space over  $F$  with **the operations of vector addition and scalar multiplication defined on  $V$ .**


# Subspace

Let  $V$  be a vector space over a field  $F$ . A **subspace** of  $V$  is a **subset**  $W$  of  $V$  which is itself a vector space over  $F$  with **the operations of vector addition and scalar multiplication defined on  $V$ .**



# Subspace

Let  $V$  be a vector space over a field  $F$ . A **subspace** of  $V$  is a **subset**  $W$  of  $V$  which is itself a vector space over  $F$  with **the operations of vector addition and scalar multiplication defined on  $V$ .**



## Remark

If  $\langle V, F, +, \cdot \rangle$  is a vector space, then

- (i)  $\forall \alpha, \beta \in V, \alpha + \beta \in V$  ( $V$  is closed under vector addition)
- (ii)  $\forall c \in F$  and  $\alpha \in V, c\alpha \in V$  ( $V$  is closed under scalar multiplication)
- (iii) If  $\alpha_1, \dots, \alpha_n \in V$ , then  $c_1\alpha_1 + \dots + c_n\alpha_n \in V$  where  $c_i \in F$ .



# Theorem 1

Let  $V$  be a vector space over the field  $F$ .

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

**Proof:**

Case 1 : Suppose that  $W$  is a subspace of  $V$ .

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

**Proof:**

Case 1 : Suppose that  $W$  is a subspace of  $V$ .  $\implies W$  is a vector space over the field  $F$ .

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

## Proof:

Case 1 : Suppose that  $W$  is a subspace of  $V$ .  $\implies W$  is a vector space over the field  $F$ .

If  $c \in F, \alpha, \beta \in W$ ,

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

## Proof:

Case 1 : Suppose that  $W$  is a subspace of  $V$ .  $\implies W$  is a vector space over the field  $F$ .

If  $c \in F, \alpha, \beta \in W$ , then  $c\alpha \in W$  (closed under scalar multiplication) and

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

## Proof:

Case 1 : Suppose that  $W$  is a subspace of  $V$ .  $\implies W$  is a vector space over the field  $F$ .

If  $c \in F, \alpha, \beta \in W$ , then  $c\alpha \in W$  (closed under scalar multiplication) and  $c\alpha + \beta \in W$  (closed under vector addition).

# Theorem 1

Let  $V$  be a vector space over the field  $F$ . A non-empty subset  $W$  of  $V$  is a subspace of  $V$  if and only if

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$

## Proof:

Case 1 : Suppose that  $W$  is a subspace of  $V$ .  $\implies W$  is a vector space over the field  $F$ .

If  $c \in F, \alpha, \beta \in W$ , then  $c\alpha \in W$  (closed under scalar multiplication) and  $c\alpha + \beta \in W$  (closed under vector addition).

Hence

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W.$$



## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a).

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a). For all  $\alpha \in W \subseteq V$ ,  $1.\alpha = \alpha$  ( $V$  is a vector space).

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a). For all  $\alpha \in W \subseteq V$ ,  $1.\alpha = \alpha$  ( $V$  is a vector space). For all  $\alpha, \beta \in W$ ,  $1.\alpha + \beta = \alpha + \beta \in W$  by (a).

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a). For all  $\alpha \in W \subseteq V$ ,  $1.\alpha = \alpha$  ( $V$  is a vector space). For all  $\alpha, \beta \in W$ ,  $1.\alpha + \beta = \alpha + \beta \in W$  by (a). For all  $c \in F$  and  $\alpha \in W$ ,  $c\alpha + 0 = c\alpha \in W$ , by (a).

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a). For all  $\alpha \in W \subseteq V$ ,  $1.\alpha = \alpha$  ( $V$  is a vector space). For all  $\alpha, \beta \in W$ ,  $1.\alpha + \beta = \alpha + \beta \in W$  by (a). For all  $c \in F$  and  $\alpha \in W$ ,  $c\alpha + 0 = c\alpha \in W$ , by (a). In addition,  $(-1)\alpha + 0 = -\alpha \in W$  for all  $\alpha \in W$  by (a).

## Theorem 1 contd.

Case 2 : Suppose that  $W$  is a non-empty subset of  $V$  such that

$$\forall \alpha, \beta \in W, c \in F \implies c\alpha + \beta \in W. \text{ --- (a)}$$

Since  $W \neq \phi$ , there exists  $\rho \in W$  and hence  $(-1)\rho + \rho = 0 \in W$ , by (a). For all  $\alpha \in W \subseteq V$ ,  $1.\alpha = \alpha$  ( $V$  is a vector space). For all  $\alpha, \beta \in W$ ,  $1.\alpha + \beta = \alpha + \beta \in W$  by (a). For all  $c \in F$  and  $\alpha \in W$ ,  $c\alpha + 0 = c\alpha \in W$ , by (a). In addition,  $(-1)\alpha + 0 = -\alpha \in W$  for all  $\alpha \in W$  by (a). Since  $W \subseteq V$ ,  $\langle W, F, +, . \rangle$  satisfies the rest of the axioms (**verify!**) of a vector space and thus  $W$  is a subspace of  $V$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.



## Examples of subspaces

- (1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.
- (2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\phi \neq W \subseteq F^n$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\emptyset \neq W \subseteq F^n$ . Let  $\alpha = (0, x_2, \dots, x_n)$ ,  $\beta = (0, y_2, \dots, y_n) \in W$  and  $c \in F$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\emptyset \neq W \subseteq F^n$ . Let  $\alpha = (0, x_2, \dots, x_n)$ ,  $\beta = (0, y_2, \dots, y_n) \in W$  and  $c \in F$ .

$$c\alpha + \beta = (0, cx_2 + y_2, \dots, cx_n + y_n) \in W$$

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\emptyset \neq W \subseteq F^n$ . Let  $\alpha = (0, x_2, \dots, x_n)$ ,  $\beta = (0, y_2, \dots, y_n) \in W$  and  $c \in F$ .

$$c\alpha + \beta = (0, cx_2 + y_2, \dots, cx_n + y_n) \in W$$

By Theorem 1,  $W$  is a subspace of  $F^n$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\emptyset \neq W \subseteq F^n$ . Let  $\alpha = (0, x_2, \dots, x_n)$ ,  $\beta = (0, y_2, \dots, y_n) \in W$  and  $c \in F$ .

$$c\alpha + \beta = (0, cx_2 + y_2, \dots, cx_n + y_n) \in W$$

By Theorem 1,  $W$  is a subspace of  $F^n$ .

(3) Prove that  $W = \{(1 + x_2, x_2, x_3, \dots, x_n) : x_i \in F\}$  is not a subspace of  $F^n$ .

## Examples of subspaces

(1) Let  $V$  be a vector space over the field  $F$ . Then the subset  $\{0\}$  of  $V$  is a subspace of  $V$  and it is called **the zero subspace**.

(2) Note that  $W = \{(0, x_2, \dots, x_n) : x_i \in F\}$  is a subspace of  $F^n$ .

**Proof:** Clearly  $0 = (0, 0, \dots, 0) \in W$ . So  $\emptyset \neq W \subseteq F^n$ . Let  $\alpha = (0, x_2, \dots, x_n)$ ,  $\beta = (0, y_2, \dots, y_n) \in W$  and  $c \in F$ .

$$c\alpha + \beta = (0, cx_2 + y_2, \dots, cx_n + y_n) \in W$$

By Theorem 1,  $W$  is a subspace of  $F^n$ .

(3) Prove that  $W = \{(1 + x_2, x_2, x_3, \dots, x_n) : x_i \in F\}$  is not a subspace of  $F^n$ .

Reason :  $0 = (0, 0, \dots, 0) \notin W$



(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .  
Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ .**

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .  
Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .**

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .  $\implies AX_1 = AX_2 = 0$ .**

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .  $\implies AX_1 = AX_2 = 0$ .**

$$\implies A(cX_1 + X_2) = cAX_1 + AX_2 = 0,$$

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .  $\implies AX_1 = AX_2 = 0$ .**

$$\implies A(cX_1 + X_2) = cAX_1 + AX_2 = 0, \quad \implies cX_1 + X_2 \in S$$

## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .  $\implies AX_1 = AX_2 = 0$ .**

$$\implies A(cX_1 + X_2) = cAX_1 + AX_2 = 0, \quad \implies cX_1 + X_2 \in S$$

$$\forall X_1, X_2 \in S, c \in F \implies cX_1 + X_2 \in S$$



## Examples contd.

**(4) Prove that the solution set of the homogeneous system  $AX = 0$  is subspace of  $F^{n \times 1}$  where  $A \in F^{m \times n}$ .**

**Let  $S = \{X \in F^{n \times 1} : AX = 0\}$ . Clearly  $0 \in S \neq \phi$ .**

**Let  $X_1, X_2 \in S$  and  $c \in F$ .  $\implies AX_1 = AX_2 = 0$ .**

$$\implies A(cX_1 + X_2) = cAX_1 + AX_2 = 0, \quad \implies cX_1 + X_2 \in S$$

$$\forall X_1, X_2 \in S, c \in F \implies cX_1 + X_2 \in S$$

**Hence,  $S$  is a subspace of  $F^{n \times 1}$ .**

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and  
(b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and  
(b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and  
(b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F$ .

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and  
(b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F. \implies \alpha, \beta \in W_i$  for  $i = 1, 2$ .

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and  
(b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F. \implies \alpha, \beta \in W_i$  for  $i = 1, 2$ .  
 $\implies c\alpha + \beta \in W_i$  for  $i = 1, 2$  by (b).

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and (b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F. \implies \alpha, \beta \in W_i$  for  $i = 1, 2$ .

$\implies c\alpha + \beta \in W_i$  for  $i = 1, 2$  by (b).

$\implies c\alpha + \beta \in W_1 \cap W_2$ .



## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and (b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F. \implies \alpha, \beta \in W_i$  for  $i = 1, 2$ .

$\implies c\alpha + \beta \in W_i$  for  $i = 1, 2$  by (b).

$\implies c\alpha + \beta \in W_1 \cap W_2$ .

By Theorem 1,  $W_1 \cap W_2$  is a subspace of  $V$ .

## Theorem 2

**Let  $V$  be a vector space over the field  $F$ . Let  $W_1, W_2$  be two subspaces of  $V$ . Then  $W_1 \cap W_2$  is a subspace of  $V$ .**

**Proof :** Since  $W_1$  and  $W_2$  are subspace of  $V$ , (a)  $0 \in W_i \neq \phi$  and (b)  $\forall \alpha, \beta \in W_i, c \in F \implies c\alpha + \beta \in W_i$  for  $i = 1, 2$

By (a),  $0 \in W_1 \cap W_2 \neq \phi$ .

Let  $\alpha, \beta \in W_1 \cap W_2, c \in F. \implies \alpha, \beta \in W_i$  for  $i = 1, 2$ .

$\implies c\alpha + \beta \in W_i$  for  $i = 1, 2$  by (b).

$\implies c\alpha + \beta \in W_1 \cap W_2$ .

By Theorem 1,  $W_1 \cap W_2$  is a subspace of  $V$ .

**Corollary :** Intersection of any collection of subspaces of a vector space  $V$  is a subspace of  $V$

## The subspace spanned by $S$

Let  $S$  be a subset of a vector space  $V$ .

## The subspace spanned by $S$

Let  $S$  be a subset of a vector space  $V$ . The subspace spanned by  $S$  is defined as the intersection all subspaces of  $V$  which contains  $S$ .

## The subspace spanned by $S$

Let  $S$  be a **subset** of a vector space  $V$ . The subspace spanned by  $S$  is defined as the intersection all subspaces of  $V$  which contains  $S$ .

Subspace spanned by  $S = \cap \{W : S \subseteq W, W \text{ is a subspace of } V\}$

## The subspace spanned by $S$

Let  $S$  be a **subset** of a vector space  $V$ . The subspace spanned by  $S$  is defined as the intersection all subspaces of  $V$  which contains  $S$ .

Subspace spanned by  $S = \cap \{W : S \subseteq W, W \text{ is a subspace of } V\}$

**Note (1) : Subspace spanned by  $S$  is the smallest subspace which contains  $S$ .**

## The subspace spanned by $S$

Let  $S$  be a **subset** of a vector space  $V$ . The subspace spanned by  $S$  is defined as the intersection all subspaces of  $V$  which contains  $S$ .

Subspace spanned by  $S = \cap \{W : S \subseteq W, W \text{ is a subspace of } V\}$

**Note (1) : Subspace spanned by  $S$  is the smallest subspace which contains  $S$ .**

**Note (2) : If  $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ , we call the subspace spanned by  $S$  as **the subspace spanned by the vectors  $\alpha_1, \alpha_2, \dots, \alpha_n$ .****

$L(S)$  = the set of all linear combinations of vectors in  $S$

Let  $S$  be a non-empty subset of a vector space  $V$ . The set of all linear combinations of vectors in  $S$  is denoted by  $L(S)$ .



$L(S)$  = the set of all linear combinations of vectors in  $S$

Let  $S$  be a **non-empty subset** of a vector space  $V$ . The set of all linear combinations of vectors in  $S$  is denoted by  $L(S)$ .

$$L(S) = \left\{ \sum_{i=1}^n c_i \alpha_i \quad : \quad c_i \in F, \alpha_i \in S, n \in \mathbb{N} \right\}$$

$L(S)$  = the set of all linear combinations of vectors in  $S$

Let  $S$  be a **non-empty subset** of a vector space  $V$ . The set of all linear combinations of vectors in  $S$  is denoted by  $L(S)$ .

$$L(S) = \left\{ \sum_{i=1}^n c_i \alpha_i \quad : \quad c_i \in F, \alpha_i \in S, n \in \mathbb{N} \right\}$$

**(1)** Let  $S = \{(1, 0, 0), (0, 0, 1)\}$

## $L(S)$ = the set of all linear combinations of vectors in $S$

Let  $S$  be a **non-empty subset** of a vector space  $V$ . The set of all linear combinations of vectors in  $S$  is denoted by  $L(S)$ .

$$L(S) = \left\{ \sum_{i=1}^n c_i \alpha_i \quad : \quad c_i \in F, \alpha_i \in S, n \in \mathbb{N} \right\}$$

**(1)** Let  $S = \{(1, 0, 0), (0, 0, 1)\}$

$$L(S) = \{a(1, 0, 0) + b(0, 0, 1) \quad : \quad a, b \in \mathbb{R}\}$$

## $L(S)$ = the set of all linear combinations of vectors in $S$

Let  $S$  be a **non-empty subset** of a vector space  $V$ . The set of all linear combinations of vectors in  $S$  is denoted by  $L(S)$ .

$$L(S) = \left\{ \sum_{i=1}^n c_i \alpha_i \quad : \quad c_i \in F, \alpha_i \in S, n \in \mathbb{N} \right\}$$

**(1)** Let  $S = \{(1, 0, 0), (0, 0, 1)\}$

$$L(S) = \{a(1, 0, 0) + b(0, 0, 1) \quad : \quad a, b \in \mathbb{R}\}$$

$$L(S) = \{(a, 0, b) \quad : \quad a, b \in \mathbb{R}\}$$

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ .

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ .

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ . In addition  $\forall \alpha \in S$ ,  $1.\alpha = \alpha \in L(S)$  and thus  $S \subseteq L(S)$ .



## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ . In addition  $\forall \alpha \in S$ ,  $1.\alpha = \alpha \in L(S)$  and thus  $S \subseteq L(S)$ .

Let  $x, y \in L(S)$ .

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ . In addition  $\forall \alpha \in S$ ,  $1.\alpha = \alpha \in L(S)$  and thus  $S \subseteq L(S)$ .

Let  $x, y \in L(S)$ .  $\implies x = \sum_{i=1}^m c_i \alpha_i, y = \sum_{j=1}^n d_j \beta_j$

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ . In addition  $\forall \alpha \in S$ ,  $1.\alpha = \alpha \in L(S)$  and thus  $S \subseteq L(S)$ .

Let  $x, y \in L(S)$ .  $\implies x = \sum_{i=1}^m c_i \alpha_i, y = \sum_{j=1}^n d_j \beta_j$

$\implies cx + y = \sum_{i=1}^m cc_i \alpha_i + \sum_{j=1}^n d_j \beta_j$  is a linear combination of vectors

in  $S$ . Thus  $cx + y \in L(S)$ .

## Lemma

$L(S)$  is a subspace of  $V$  and  $S \subseteq L(S)$ .

**Proof:** Since  $S \neq \phi$ , there exists  $\alpha \in S \subseteq V$ . By Note 2,  $0\alpha = 0 \in L(S)$  ( $0\alpha$  is a linear combination of  $\alpha$ ).  $L(S) \neq \phi$ . In addition  $\forall \alpha \in S$ ,  $1.\alpha = \alpha \in L(S)$  and thus  $S \subseteq L(S)$ .

Let  $x, y \in L(S)$ .  $\implies x = \sum_{i=1}^m c_i \alpha_i, y = \sum_{j=1}^n d_j \beta_j$

$\implies cx + y = \sum_{i=1}^m cc_i \alpha_i + \sum_{j=1}^n d_j \beta_j$  is a linear combination of vectors

in  $S$ . Thus  $cx + y \in L(S)$ .

$\implies L(S)$  is a subspace of  $V$  by Theorem 1.

## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ .  
Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ . Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

**Proof.**

It is enough to prove that the subspace spanned by  $S = L(S)$ .

## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ . Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

**Proof.**

It is enough to prove that the subspace spanned by  $S = L(S)$ . Prove that

$$W^* = \cap \{W : S \subseteq W, \text{ } W \text{ is a subspace of } V\}$$

## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ . Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

**Proof.**

It is enough to prove that the subspace spanned by  $S = L(S)$ . Prove that

$$W^* = \cap \{W : S \subseteq W, \text{ } W \text{ is a subspace of } V\} = L(S) \text{ --- (a)}$$



## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ . Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

**Proof.**

It is enough to prove that the subspace spanned by  $S = L(S)$ . Prove that

$$W^* = \cap \{W : S \subseteq W, \text{ } W \text{ is a subspace of } V\} = L(S) \text{ --- (a)}$$

By the previous lemma,  $S \subseteq L(S)$  and  $L(S)$  is a subspace of  $V$ ,

## Theorem 3

Let  $S$  be a non-empty subset of a vector space  $V$  over the field  $F$ . Then the subspace spanned by the set  $S$  is the set of all linear combinations of vectors in  $S$ .

**Proof.**

It is enough to prove that the subspace spanned by  $S = L(S)$ . Prove that

$$W^* = \cap \{ W : S \subseteq W, \text{ } W \text{ is a subspace of } V \} = L(S) \text{ --- (a)}$$

By the previous lemma,  $S \subseteq L(S)$  and  $L(S)$  is a subspace of  $V$ , and thus  $W^* \subseteq L(S) \text{ --- (i)}$ .

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

$$x \in L(S) \implies x \in W.$$

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

$$x \in L(S) \implies x \in W. \text{ Thus } L(S) \subseteq W.$$

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

$$x \in L(S) \implies x \in W. \text{ Thus } L(S) \subseteq W.$$

By above claim,

$$L(S) \subseteq \cap \{W : S \subseteq W, \text{ } W \text{ is a subspace of } V\}$$

## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

$$x \in L(S) \implies x \in W. \text{ Thus } L(S) \subseteq W.$$

By above claim,

$$L(S) \subseteq \cap \{W : S \subseteq W, W \text{ is a subspace of } V\} = W^* \text{ --- (ii)}$$



## Theorem 3 contd.

**Claim :** If  $W$  is a subspace of  $V$  and  $S \subseteq W$ , then  $L(S) \subseteq W$ .

Let  $x \in L(S)$ .  $\implies$   $x$  is a linear combination of vectors in  $S$ . Since  $W$  is a subspace and  $S \subseteq W$ , every linear combination of vectors in  $S$  is also a member of  $W$  and thus  $x \in W$ .

$$x \in L(S) \implies x \in W. \text{ Thus } L(S) \subseteq W.$$

By above claim,

$$L(S) \subseteq \cap \{W : S \subseteq W, W \text{ is a subspace of } V\} = W^* \text{ --- (ii)}$$

By (i) and (ii),

$$W^* = \cap \{W : S \subseteq W, W \text{ is a subspace of } V\} = L(S) \text{ --- (a)}$$

## Row space and Column space of a matrix

Let  $A \in F^{m \times n}$  with rows  $\{R_1, R_2, \dots, R_m\}$  and columns  $\{C_1, C_2, \dots, C_n\}$ .

## Row space and Column space of a matrix

Let  $A \in F^{m \times n}$  with rows  $\{R_1, R_2, \dots, R_m\}$  and columns  $\{C_1, C_2, \dots, C_n\}$ . Then

Row space of  $A =$  The subspace spanned by  $R_1, R_2, \dots, R_m$

## Row space and Column space of a matrix

Let  $A \in F^{m \times n}$  with rows  $\{R_1, R_2, \dots, R_m\}$  and columns  $\{C_1, C_2, \dots, C_n\}$ . Then

Row space of  $A =$  The subspace spanned by  $R_1, R_2, \dots, R_m$

Column space of  $A =$  The subspace spanned by  $C_1, C_2, \dots, C_n$

## Row space and Column space of a matrix

Let  $A \in F^{m \times n}$  with rows  $\{R_1, R_2, \dots, R_m\}$  and columns  $\{C_1, C_2, \dots, C_n\}$ . Then

Row space of  $A =$  The subspace spanned by  $R_1, R_2, \dots, R_m$

Column space of  $A =$  The subspace spanned by  $C_1, C_2, \dots, C_n$

**Note :** Row space of  $A \subseteq F^{1 \times n}$  and Column space of  $A \subseteq F^{m \times 1}$ .

## Example

$$\text{Let } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

where

$$R_1 = (1, 0, 0), R_2 = (0, 1, 0), C_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, C_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, C_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

## Example

$$\text{Let } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

where

$$R_1 = (1, 0, 0), R_2 = (0, 1, 0), C_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, C_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, C_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Row space of  $A$

$$= \{x(1, 0, 0) + y(0, 1, 0) : x, y \in F\}$$

## Example

$$\text{Let } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

where

$$R_1 = (1, 0, 0), R_2 = (0, 1, 0), C_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, C_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, C_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Row space of  $A$

$$= \{x(1, 0, 0) + y(0, 1, 0) : x, y \in F\} = \{(x, y, 0) : x, y \in F\}$$



## Example

$$\text{Let } A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

where

$$R_1 = (1, 0, 0), R_2 = (0, 1, 0), C_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, C_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, C_3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Row space of  $A$

$$= \{x(1, 0, 0) + y(0, 1, 0) : x, y \in F\} = \{(x, y, 0) : x, y \in F\}$$

Column Space of  $A$

$$= \left\{ x \begin{pmatrix} 1 \\ 0 \end{pmatrix} + y \begin{pmatrix} 0 \\ 1 \end{pmatrix} + z \begin{pmatrix} 0 \\ 0 \end{pmatrix} : x, y, z \in F \right\} = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \right\}_{14}$$

# Assignment

Prove or disprove that

- (i) column space of  $AB$  is same as column space of  $A$  and
- (ii) row space of  $AB$  is same as row space of  $B$ .

## Note 1: ( Visit previous lecture notes)

Find the solution space of the system  $RX = 0$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Note 1: ( Visit previous lecture notes)

Find the solution space of the system  $RX = 0$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

**No. of non-zero rows of  $R$ ,  $r = 2$ , No. of variables,  $n = 5$**

$k_1 = 2, k_2 = 4 \implies$  Pivot variables  $= \{x_{k_1}, x_{k_2}\} = \{x_2, x_4\}$

No. of free variables  $= n - r = 5 - 2 = 3$ ,

Free variables  $= \{u_1, u_2, u_3\} = \{x_1, x_3, x_5\}$

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \implies \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j}u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j}u_j = 0 \end{array} \right\} \text{ (general expression)}$$

## Note 1 contd.

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0 \end{array} \right\} \text{general expression)}$$

**Set the free variables as :**

$$u_1 = x_1 = a, \quad u_2 = x_3 = b, \quad u_3 = x_5 = c$$

$$\Rightarrow x_2 = 3b - \frac{1}{2}c, \quad x_4 = -2c$$

$$\text{Solution set } S = \left\{ (a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R} \right\}$$

## Note 1 contd. (back to chapter one !)

**Solution set**  $S = \{(a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R}\}$

$$S = \left\{ a(1, 0, 0, 0, 0) + b(0, 3, 1, 0, 0) + c(0, -\frac{1}{2}, 0, -2, 1) : a, b, c \in \mathbb{R} \right\}$$

$$= \text{Span of } \left\{ (1, 0, 0, 0, 0), (0, 3, 1, 0, 0), (0, -\frac{1}{2}, 0, -2, 1) \right\}$$

**Dimension of  $S = \dim S = 3 = n - r$  (Information for future)**

## Problem

Let  $W$  be set of all  $(x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^5$  which satisfies

$$2x_1 - x_2 + \frac{4}{3}x_3 - x_4 = 0$$

$$x_1 + \frac{2}{3}x_3 - x_5 = 0$$

$$9x_1 - 3x_2 + 6x_3 - 3x_4 - 3x_5 = 0$$

Find a finite set of vectors which spans  $W$ .

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ .



Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

## Note

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

## Note

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

## Note

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\gamma = (3, 4)$ .

## Note

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\gamma = (3, 4)$ . Prove that there is no  $c \in \mathbb{R}$  such that  $\gamma = c\alpha$

## Note

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\gamma = (3, 4)$ . Prove that there is no  $c \in \mathbb{R}$  such that  $\gamma = c\alpha$

$$c_1\alpha + c_2\gamma = 0 \implies c_1 = c_2 = 0$$

Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\gamma = (3, 4)$ . Prove that there is no  $c \in \mathbb{R}$  such that  $\gamma = c\alpha$

$$c_1\alpha + c_2\gamma = 0 \implies c_1 = c_2 = 0$$

We say  $\{\alpha, \gamma\}$  is a linearly independent set.



Let  $\alpha = (2, 3)$  and  $\beta = (6, 9)$ . Then  $\beta = 3\alpha$ .

$$\implies 3\alpha + (-1)\beta = 0.$$

$$\implies c_1\alpha + c_2\beta = 0 \text{ where } c_i \neq 0 \text{ for at least one } i.$$

We say  $\{\alpha, \beta\}$  is a linearly dependent set.

Let  $\gamma = (3, 4)$ . Prove that there is no  $c \in \mathbb{R}$  such that  $\gamma = c\alpha$

$$c_1\alpha + c_2\gamma = 0 \implies c_1 = c_2 = 0$$

We say  $\{\alpha, \gamma\}$  is a linearly independent set.

Since  $3\alpha + (-1)\beta + 0\gamma = 0$ ,  $\{\alpha, \beta, \gamma\}$  is a linearly dependent set.