

Absolute and Conditional Convergence

Definition (Absolute Convergence)

A series $\sum a_n$ **converges absolutely** if the corresponding series of absolute values $\sum |a_n|$ converges.

Absolute and Conditional Convergence

Definition (Absolute Convergence)

A series $\sum a_n$ **converges absolutely** if the corresponding series of absolute values $\sum |a_n|$ converges.

Example:

The geometric series

$$1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$$

converges absolutely because the corresponding series of absolute values

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$$

converges.

Absolute and Conditional Convergence

Definition (Absolute Convergence)

A series $\sum a_n$ **converges absolutely** if the corresponding series of absolute values $\sum |a_n|$ converges.

Example:

The geometric series

$$1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots$$

converges absolutely because the corresponding series of absolute values

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$$

converges.

In contrast, the alternating harmonic series does not converge absolutely: The corresponding series of absolute values is the divergent harmonic series.

Definition (Conditional Convergence)

A series that converges but does not converge absolutely is said to **converge conditionally**.

Example: The alternating harmonic series converges conditionally.

The Absolute Convergence Test

Theorem

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} a_n$ converges.

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} 2|a_n|$ converges.

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} 2|a_n|$ converges.

So, by the Comparison Test, $\sum_{n=1}^{\infty} (a_n + |a_n|)$ converges.

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} 2|a_n|$ converges.

So, by the Comparison Test, $\sum_{n=1}^{\infty} (a_n + |a_n|)$ converges.

But $a_n = (a_n + |a_n|) - |a_n|$.

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} 2|a_n|$ converges.

So, by the Comparison Test, $\sum_{n=1}^{\infty} (a_n + |a_n|)$ converges.

But $a_n = (a_n + |a_n|) - |a_n|$. So, $\sum_{n=1}^{\infty} a_n$ can be expressed as the difference of two convergent series:

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (a_n + |a_n|) - \sum_{n=1}^{\infty} |a_n|.$$

Proof:

For each n ,

$$-|a_n| \leq a_n \leq |a_n| \quad \text{so} \quad 0 \leq a_n + |a_n| \leq 2|a_n|.$$

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} 2|a_n|$ converges.

So, by the Comparison Test, $\sum_{n=1}^{\infty} (a_n + |a_n|)$ converges.

But $a_n = (a_n + |a_n|) - |a_n|$. So, $\sum_{n=1}^{\infty} a_n$ can be expressed as the difference of two convergent series:

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (a_n + |a_n| - |a_n|) = \sum_{n=1}^{\infty} (a_n + |a_n|) - \sum_{n=1}^{\infty} |a_n|.$$

Therefore, $\sum_{n=1}^{\infty} a_n$ converges.

Examples

- (a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series
- $$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

Examples

(a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series

$$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges.

Examples

(a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series

$$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely.

Examples

(a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series

$$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely. Hence it converges.

Examples

(a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series

$$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely. Hence it converges.

(b) For $\sum_{n=1}^{\infty} \frac{\sin n}{n^2} = \frac{\sin 1}{1} + \frac{\sin 2}{4} + \frac{\sin 3}{9} + \dots$, the corresponding series of absolute values is

$$\sum_{n=1}^{\infty} \left| \frac{\sin n}{n^2} \right| = \left| \frac{\sin 1}{1} \right| + \left| \frac{\sin 2}{4} \right| + \left| \frac{\sin 3}{9} \right| + \dots$$

Examples

- (a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series
- $$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely. Hence it converges.

- (b) For $\sum_{n=1}^{\infty} \frac{\sin n}{n^2} = \frac{\sin 1}{1} + \frac{\sin 2}{4} + \frac{\sin 3}{9} + \dots$, the corresponding series of absolute values is

$$\sum_{n=1}^{\infty} \left| \frac{\sin n}{n^2} \right| = \left| \frac{\sin 1}{1} \right| + \left| \frac{\sin 2}{4} \right| + \left| \frac{\sin 3}{9} \right| + \dots$$

Since $|\sin n| \leq 1$, the latter converges by comparison with $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

Examples

- (a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series
- $$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely. Hence it converges.

- (b) For $\sum_{n=1}^{\infty} \frac{\sin n}{n^2} = \frac{\sin 1}{1} + \frac{\sin 2}{4} + \frac{\sin 3}{9} + \dots$, the corresponding series of absolute values is

$$\sum_{n=1}^{\infty} \left| \frac{\sin n}{n^2} \right| = \left| \frac{\sin 1}{1} \right| + \left| \frac{\sin 2}{4} \right| + \left| \frac{\sin 3}{9} \right| + \dots$$

Since $|\sin n| \leq 1$, the latter converges by comparison with $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

Thus the original series converges absolutely.

Examples

- (a) For $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^3} = 1 - \frac{1}{8} + \frac{1}{27} - \frac{1}{64} + \dots$, the corresponding series of absolute values is the series
- $$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots$$

The latter converges. Thus the original series converges absolutely. Hence it converges.

- (b) For $\sum_{n=1}^{\infty} \frac{\sin n}{n^2} = \frac{\sin 1}{1} + \frac{\sin 2}{4} + \frac{\sin 3}{9} + \dots$, the corresponding series of absolute values is

$$\sum_{n=1}^{\infty} \left| \frac{\sin n}{n^2} \right| = \left| \frac{\sin 1}{1} \right| + \left| \frac{\sin 2}{4} \right| + \left| \frac{\sin 3}{9} \right| + \dots$$

Since $|\sin n| \leq 1$, the latter converges by comparison with $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

Thus the original series converges absolutely. Hence it converges.

Power Series

Definition (Power Series, Center, Coefficients)

A **power series about** $x = 0$ is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

Power Series

Definition (Power Series, Center, Coefficients)

A **power series about** $x = 0$ is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

A **power series about** $x = a$ is a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \dots + c_n (x - a)^n + \dots$$

Power Series

Definition (Power Series, Center, Coefficients)

A **power series about** $x = 0$ is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

A **power series about** $x = a$ is a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \dots + c_n (x - a)^n + \dots$$

Here a is the **center**

Power Series

Definition (Power Series, Center, Coefficients)

A **power series about** $x = 0$ is a series of the form

$$\sum_{n=0}^{\infty} c_n x^n = c_0 + c_1 x + c_2 x^2 + \dots + c_n x^n + \dots$$

A **power series about** $x = a$ is a series of the form

$$\sum_{n=0}^{\infty} c_n (x - a)^n = c_0 + c_1 (x - a) + c_2 (x - a)^2 + \dots + c_n (x - a)^n + \dots$$

Here a is the **center** and $c_0, c_1, c_2, \dots, c_n, \dots$ are the **coefficients** of the power series. These are constants.

Example: A geometric series

Taking all the coefficients to be 1 gives the geometric power series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots + x^n + \dots$$

Example: A geometric series

Taking all the coefficients to be 1 gives the geometric power series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots + x^n + \dots$$

This the geometric series with first term 1 and common ratio x .

Example: A geometric series

Taking all the coefficients to be 1 gives the geometric power series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots + x^n + \dots$$

This the geometric series with first term 1 and common ratio x .
It converges to $1/(1 - x)$ for $|x| < 1$. We express this fact by writing

$$\frac{1}{1 - x} = 1 + x + x^2 + \dots + x^n + \dots, \quad -1 < x < 1.$$

Note

The power series

$$\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + \dots, \quad -1 < x < 1,$$

gives the following polynomial approximations for the non-polynomial function $\frac{1}{1-x}$ for values of x near 0:

$$P_0(x) = 1$$

Note

The power series

$$\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + \dots, \quad -1 < x < 1,$$

gives the following polynomial approximations for the non-polynomial function $\frac{1}{1-x}$ for values of x near 0:

$$\begin{aligned} P_0(x) &= 1 \\ P_1(x) &= 1 + x \end{aligned}$$

Note

The power series

$$\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + \dots, \quad -1 < x < 1,$$

gives the following polynomial approximations for the non-polynomial function $\frac{1}{1-x}$ for values of x near 0:

$$\begin{aligned} P_0(x) &= 1 \\ P_1(x) &= 1 + x \\ P_2(x) &= 1 + x + x^2 \end{aligned}$$

Note

The power series

$$\frac{1}{1-x} = 1 + x + x^2 + \dots + x^n + \dots, \quad -1 < x < 1,$$

gives the following polynomial approximations for the non-polynomial function $\frac{1}{1-x}$ for values of x near 0:

$$\begin{aligned} P_0(x) &= 1 \\ P_1(x) &= 1 + x \\ P_2(x) &= 1 + x + x^2 \\ &\vdots \end{aligned}$$

Example: Another geometric series

The power series

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

has center $a = 2$ and coefficients $c_0 = 1, c_1 = -1/2, c_2 = 1/4, \dots, c_n = (-1/2)^n, \dots$

Example: Another geometric series

The power series

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

has center $a = 2$ and coefficients $c_0 = 1, c_1 = -1/2, c_2 = 1/4, \dots, c_n = (-1/2)^n, \dots$

This is a geometric series with first term 1 and common ratio $r = -\frac{x-2}{2}$.

Example: Another geometric series

The power series

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

has center $a = 2$ and coefficients $c_0 = 1, c_1 = -1/2, c_2 = 1/4, \dots, c_n = (-1/2)^n, \dots$

This is a geometric series with first term 1 and common ratio $r = -\frac{x-2}{2}$.

The series converges for $|r| = \left| -\frac{x-2}{2} \right| < 1$ or $0 < x < 4$.

The sum is

$$\frac{1}{1-r} = \frac{1}{1 + \frac{x-2}{2}} = \frac{2}{x}.$$

Example: Another geometric series

The power series

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

has center $a = 2$ and coefficients $c_0 = 1, c_1 = -1/2, c_2 = 1/4, \dots, c_n = (-1/2)^n, \dots$

This is a geometric series with first term 1 and common ratio $r = -\frac{x-2}{2}$.

The series converges for $|r| = \left| -\frac{x-2}{2} \right| < 1$ or $0 < x < 4$.

The sum is

$$\frac{1}{1-r} = \frac{1}{1 + \frac{x-2}{2}} = \frac{2}{x}.$$

So,

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots, \quad 0 < x < 4.$$

Note

The power series

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots, \quad 0 < x < 4,$$

gives the following polynomial approximations for the non-polynomial function $\frac{2}{x}$ for values of x near 2:

$$P_0(x) = 1$$

Note

The power series

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots, \quad 0 < x < 4,$$

gives the following polynomial approximations for the non-polynomial function $\frac{2}{x}$ for values of x near 2:

$$P_0(x) = 1$$

$$P_1(x) = 1 - \frac{1}{2}(x-2)$$

Note

The power series

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots, \quad 0 < x < 4,$$

gives the following polynomial approximations for the non-polynomial function $\frac{2}{x}$ for values of x near 2:

$$P_0(x) = 1$$

$$P_1(x) = 1 - \frac{1}{2}(x-2)$$

$$P_2(x) = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2$$

Note

The power series

$$\frac{2}{x} = 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots, \quad 0 < x < 4,$$

gives the following polynomial approximations for the non-polynomial function $\frac{2}{x}$ for values of x near 2:

$$\begin{aligned} P_0(x) &= 1 \\ P_1(x) &= 1 - \frac{1}{2}(x-2) \\ P_2(x) &= 1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 \\ &\vdots \end{aligned}$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Thus the given power series converges absolutely for $|x| < 1$.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Thus the given power series converges absolutely for $|x| < 1$.
It diverges if $|x| > 1$ since the n th term does not converge to zero (?).

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Thus the given power series converges absolutely for $|x| < 1$.
It diverges if $|x| > 1$ since the n th term does not converge to zero (?).
At $x = 1$, it becomes the alternating harmonic series. So, it converges (conditionally).

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Thus the given power series converges absolutely for $|x| < 1$.

It diverges if $|x| > 1$ since the n th term does not converge to zero (?).

At $x = 1$, it becomes the alternating harmonic series. So, it converges (conditionally).

At $x = -1$, we get the negative of the harmonic series. So, it diverges.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{n+1} \frac{n}{x^n} \right| = \frac{n}{n+1} |x| \rightarrow |x|.$$

Thus the given power series converges absolutely for $|x| < 1$.

It diverges if $|x| > 1$ since the n th term does not converge to zero (?).

At $x = 1$, it becomes the alternating harmonic series. So, it converges (conditionally).

At $x = -1$, we get the negative of the harmonic series. So, it diverges.

Summary: the series converges for $-1 < x \leq 1$ and diverges for other values of x .

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Thus the given power series converges absolutely for $x^2 < 1$.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Thus the given power series converges absolutely for $x^2 < 1$.
It diverges for $x^2 > 1$ since the n th term does not converge to zero.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Thus the given power series converges absolutely for $x^2 < 1$.

It diverges for $x^2 > 1$ since the n th term does not converge to zero.

At $x = 1$, the series becomes the alternating series $1 - 1/3 + 1/5 - 1/7 + \dots$, which converges by the Alternating Series Test.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Thus the given power series converges absolutely for $x^2 < 1$.

It diverges for $x^2 > 1$ since the n th term does not converge to zero.

At $x = 1$, the series becomes the alternating series $1 - 1/3 + 1/5 - 1/7 + \dots$, which converges by the Alternating Series Test.

The value at $x = -1$ is the negative of the value at $x = 1$. So, it converges for $x = -1$ as well.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^{2n-1}}{2n-1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{2n+1}}{2n+1} \frac{2n-1}{x^{2n-1}} \right| = \frac{2n-1}{2n+1} x^2 \rightarrow x^2.$$

Thus the given power series converges absolutely for $x^2 < 1$.

It diverges for $x^2 > 1$ since the n th term does not converge to zero.

At $x = 1$, the series becomes the alternating series $1 - 1/3 + 1/5 - 1/7 + \dots$, which converges by the Alternating Series Test.

The value at $x = -1$ is the negative of the value at $x = 1$. So, it converges for $x = -1$ as well.

Summary: The series converges for $|x| \leq 1$ and diverges otherwise.

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{(n+1)!} \frac{n!}{x^n} \right| = \frac{|x|}{n+1} \rightarrow 0 \quad \text{for every } x.$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=1}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{x^{n+1}}{(n+1)!} \cdot \frac{n!}{x^n} \right| = \frac{|x|}{n+1} \rightarrow 0 \quad \text{for every } x.$$

So, the series converges absolutely for all values of x .

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} n!x^n = 1 + x + 2!x^2 + 3!x^3 + \dots$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} n!x^n = 1 + x + 2!x^2 + 3!x^3 + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=0}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} n!x^n = 1 + x + 2!x^2 + 3!x^3 + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=0}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| = (n+1)|x| \rightarrow \infty \quad \text{for every } x \neq 0.$$

Testing Power Series for Convergence Using the Ratio Test

For what values of x does the following series converge?

$$\sum_{n=0}^{\infty} n!x^n = 1 + x + 2!x^2 + 3!x^3 + \dots$$

Solution: We apply the Ratio Test to the series $\sum_{n=0}^{\infty} |u_n|$, where u_n is the n th term of the given power series:

$$\left| \frac{u_{n+1}}{u_n} \right| = \left| \frac{(n+1)!x^{n+1}}{n!x^n} \right| = (n+1)|x| \rightarrow \infty \quad \text{for every } x \neq 0.$$

So, the n th term of the series does not converge to zero for $x \neq 0$. Hence the series diverges for all values of x except $x = 0$.

Theorem (The Convergence Theorem for Power Series)

If the power series $\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$ converges for $x = c \neq 0$, then it converges absolutely for all x with $|x| < |c|$.

Theorem (The Convergence Theorem for Power Series)

If the power series $\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots$ converges for $x = c \neq 0$, then it converges absolutely for all x with $|x| < |c|$. If the series diverges for $x = d$, then it diverges for all x with $|x| > |d|$.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

There are only a finite number of terms prior to $|a_N x^N|$ and so their sum is finite.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

There are only a finite number of terms prior to $|a_N x^N|$ and so their sum is finite. Starting from $|a_N x^N|$, the sum of the terms is less than

$$\left| \frac{x}{c} \right|^N + \left| \frac{x}{c} \right|^{N+1} + \left| \frac{x}{c} \right|^{N+2} + \dots$$

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

There are only a finite number of terms prior to $|a_N x^N|$ and so their sum is finite. Starting from $|a_N x^N|$, the sum of the terms is less than

$$\left| \frac{x}{c} \right|^N + \left| \frac{x}{c} \right|^{N+1} + \left| \frac{x}{c} \right|^{N+2} + \dots$$

But the above series is a geometric series with common ratio less than 1 since $|x| < |c|$.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

There are only a finite number of terms prior to $|a_N x^N|$ and so their sum is finite. Starting from $|a_N x^N|$, the sum of the terms is less than

$$\left| \frac{x}{c} \right|^N + \left| \frac{x}{c} \right|^{N+1} + \left| \frac{x}{c} \right|^{N+2} + \dots$$

But the above series is a geometric series with common ratio less than 1 since $|x| < |c|$. So, it converges.

Proof:

Suppose the series $\sum_{n=0}^{\infty} a_n c^n$ converges.

Then $a_n c^n \rightarrow 0$. Therefore, corresponding to $\epsilon = 1$, there is an integer N such that, for $n \geq N$,

$$|a_n c^n| < 1 \quad \text{or} \quad |a_n| < \frac{1}{|c|^n}.$$

Now take any x such that $|x| < |c|$ and consider

$$|a_0| + |a_1 x| + \dots + |a_{N-1} x^{N-1}| + |a_N x^N| + |a_{N+1} x^{N+1}| + \dots$$

There are only a finite number of terms prior to $|a_N x^N|$ and so their sum is finite. Starting from $|a_N x^N|$, the sum of the terms is less than

$$\left| \frac{x}{c} \right|^N + \left| \frac{x}{c} \right|^{N+1} + \left| \frac{x}{c} \right|^{N+2} + \dots$$

But the above series is a geometric series with common ratio less than 1 since $|x| < |c|$. So, it converges. Thus it follows that the given power series converges absolutely for $|x| < |c|$.

To prove the second half of the theorem, we use the first half.

To prove the second half of the theorem, we use the first half.

Suppose, for contradiction, that the power series diverges at $x = d$ and converges at a value x_0 with $|x_0| > |d|$.

To prove the second half of the theorem, we use the first half.

Suppose, for contradiction, that the power series diverges at $x = d$ and converges at a value x_0 with $|x_0| > |d|$.

Then by taking $c = x_0$, we can conclude by the first half of the theorem that the power series converges at $x = d$, which is a contradiction.

To prove the second half of the theorem, we use the first half.

Suppose, for contradiction, that the power series diverges at $x = d$ and converges at a value x_0 with $|x_0| > |d|$.

Then by taking $c = x_0$, we can conclude by the first half of the theorem that the power series converges at $x = d$, which is a contradiction.

Thus, it follows that if the power series diverges for $x = d$, then it diverges for all x with $|x| > |d|$.

Note

We prove a similar theorem for power series of the form

$$\sum_{n=0}^{\infty} a_n (x - a)^n$$

by considering the power series $\sum_{n=0}^{\infty} a_n y^n$:

From the theorem, the latter series converges for $y = c \neq 0$ implies that it converges for

$$|y| < |c|.$$

This means that the given power series converges for x with $|x - a| < |c|$.

And so on.

The Radius of Convergence

Theorem

The convergence of the series $\sum_{n=0}^{\infty} c_n(x-a)^n$ is described by one of the following three possibilities:

1. There is a positive number R such that the series diverges for x with $|x-a| > R$ but converges absolutely for x with $|x-a| < R$ (i.e., for $a-R < x < a+R$). The series may or may not converge at either of the end points $x = a-R$ and $x = a+R$.

The Radius of Convergence

Theorem

The convergence of the series $\sum_{n=0}^{\infty} c_n(x-a)^n$ is described by one of the following three possibilities:

1. There is a positive number R such that the series diverges for x with $|x-a| > R$ but converges absolutely for x with $|x-a| < R$ (i.e., for $a-R < x < a+R$). The series may or may not converge at either of the end points $x = a-R$ and $x = a+R$.
2. The series converges absolutely for every x ($R = \infty$).

The Radius of Convergence

Theorem

The convergence of the series $\sum_{n=0}^{\infty} c_n(x-a)^n$ is described by one of the following three possibilities:

1. There is a positive number R such that the series diverges for x with $|x-a| > R$ but converges absolutely for x with $|x-a| < R$ (i.e., for $a-R < x < a+R$). The series may or may not converge at either of the end points $x = a-R$ and $x = a+R$.
2. The series converges absolutely for every x ($R = \infty$).
3. The series converges at $x = a$ and diverges for all $x \neq a$ ($R = 0$).

Proof

- We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.
- ▶ Then S is nonempty as it contains 0.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.
- ▶ Then S is nonempty as it contains 0. Also it contains the numbers $\pm p$:

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.
- ▶ Then S is nonempty as it contains 0. Also it contains the numbers $\pm p$: $0 < p < |c|$.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.
- ▶ Then S is nonempty as it contains 0. Also it contains the numbers $\pm p$: $0 < p < |c|$.
- ▶ Indeed S contains all nonzero real numbers x with $|x| \leq p$ by the Convergence Theorem.

Proof

- ▶ We will prove the theorem for power series of the form $\sum_{n=0}^{\infty} c_n x^n$.
- ▶ If the series converges for every x , then we are in Case 2.
- ▶ If the series converges only at $x = 0$, then we are in Case 3.
- ▶ Otherwise there is a nonzero number d such that $\sum c_n d^n$ diverges and there is a nonzero number c such that $\sum c_n c^n$ converges.
- ▶ Let S denote the set of all numbers x such that $\sum c_n x^n$ converges.
- ▶ Then S is nonempty as it contains 0. Also it contains the numbers $\pm p$: $0 < p < |c|$.
- ▶ Indeed S contains all nonzero real numbers x with $|x| \leq p$ by the Convergence Theorem.

Proof...

- Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$.
So, $S \subseteq [-d, d]$.

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$.
So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$.
So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.
- ▶ Then by the Completeness Property of the real numbers, S has a least upper bound R .

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$.
So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.
- ▶ Then by the Completeness Property of the real numbers, S has a least upper bound R .
- ▶ If $|x| > R$, then $x \notin S$ and so $\sum c_n x^n$ diverges.

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$.
So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.
- ▶ Then by the Completeness Property of the real numbers, S has a least upper bound R .
- ▶ If $|x| > R$, then $x \notin S$ and so $\sum c_n x^n$ diverges.
- ▶ If $|x| < R$, then there is a number $b \in S$ such that $|x| < b < R$.

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$. So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.
- ▶ Then by the Completeness Property of the real numbers, S has a least upper bound R .
- ▶ If $|x| > R$, then $x \notin S$ and so $\sum c_n x^n$ diverges.
- ▶ If $|x| < R$, then there is a number $b \in S$ such that $|x| < b < R$. So, $\sum c_n b^n$ converges.

Proof...

- ▶ Also by the Convergence Theorem again, the power series diverges for all x with $|x| > |d|$. So, $S \subseteq [-d, d]$.
- ▶ Thus S is a bounded set.
- ▶ Then by the Completeness Property of the real numbers, S has a least upper bound R .
- ▶ If $|x| > R$, then $x \notin S$ and so $\sum c_n x^n$ diverges.
- ▶ If $|x| < R$, then there is a number $b \in S$ such that $|x| < b < R$. So, $\sum c_n b^n$ converges. Therefore $\sum c_n x^n$ converges absolutely by the Convergence Theorem.

The Radius of Convergence

The power series $\sum_{n=0}^{\infty} a_n x^n$ converges if

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1} x^{n+1}}{a_n x^n} \right| = |x| \cdot \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1.$$

The Radius of Convergence

The power series $\sum_{n=0}^{\infty} a_n x^n$ converges if

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1} x^{n+1}}{a_n x^n} \right| = |x| \cdot \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1.$$

That is, if

$$|x| < \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|.$$

The Radius of Convergence

The power series $\sum_{n=0}^{\infty} a_n x^n$ converges if

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{a_{n+1} x^{n+1}}{a_n x^n} \right| = |x| \cdot \lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1.$$

That is, if

$$|x| < \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|.$$

Definition

The **radius of convergence** of the power series $\sum_{n=0}^{\infty} a_n x^n$ is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|.$$

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{1/n}{1/(n+1)} = \lim_{n \rightarrow \infty} \frac{n+1}{n} = 1.$$

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{1/n}{1/(n+1)} = \lim_{n \rightarrow \infty} \frac{n+1}{n} = 1.$$

Thus the given power series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{1/n}{1/(n+1)} = \lim_{n \rightarrow \infty} \frac{n+1}{n} = 1.$$

Thus the given power series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.

At $x = 1$, it becomes the alternating harmonic series. So, it converges.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.
So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{1/n}{1/(n+1)} = \lim_{n \rightarrow \infty} \frac{n+1}{n} = 1.$$

Thus the given power series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.

At $x = 1$, it becomes the alternating harmonic series. So, it converges.
At $x = -1$, we get the negative of the harmonic series. So, it diverges.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$\sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$$

Solution: Here $a_n = \frac{(-1)^{n-1}}{n}$.
So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{1/n}{1/(n+1)} = \lim_{n \rightarrow \infty} \frac{n+1}{n} = 1.$$

Thus the given power series converges absolutely for $|x| < 1$ and diverges for $|x| > 1$.

At $x = 1$, it becomes the alternating harmonic series. So, it converges.
At $x = -1$, we get the negative of the harmonic series. So, it diverges.

Thus the interval of convergence of the power series is $-1 < x \leq 1$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Solution: Here $a_n = \left(-\frac{1}{2}\right)^n$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Solution: Here $a_n = \left(-\frac{1}{2}\right)^n$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{(1/2)^n}{(1/2)^{n+1}} = 2.$$

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Solution: Here $a_n = \left(-\frac{1}{2}\right)^n$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{(1/2)^n}{(1/2)^{n+1}} = 2.$$

Thus the given power series converges absolutely for $|x-2| < 2$ and diverges for $|x-2| > 2$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Solution: Here $a_n = \left(-\frac{1}{2}\right)^n$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{(1/2)^n}{(1/2)^{n+1}} = 2.$$

Thus the given power series converges absolutely for $|x-2| < 2$ and diverges for $|x-2| > 2$.

As we know that it is a geometric ratio $r = -\frac{x-2}{2}$, it diverges for x with $|x-2| = 2$.

Example: Computing the Radius and Interval of Convergence

Find the radius and interval of convergence of

$$1 - \frac{1}{2}(x-2) + \frac{1}{4}(x-2)^2 + \dots + \left(-\frac{1}{2}\right)^n (x-2)^n + \dots$$

Solution: Here $a_n = \left(-\frac{1}{2}\right)^n$.

So, the radius of convergence is

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{(1/2)^n}{(1/2)^{n+1}} = 2.$$

Thus the given power series converges absolutely for $|x-2| < 2$ and diverges for $|x-2| > 2$.

As we know that it is a geometric ratio $r = -\frac{x-2}{2}$, it diverges for x with $|x-2| = 2$.

Thus the interval of convergence is $|x-2| < 2$ or $0 < x < 4$.

Theorem (The Term-by-Term Differentiation Theorem)

If $\sum_{n=0}^{\infty} c_n(x-a)^n$ converges for $a-R < x < a+R$ for some $R > 0$, it defines a function f :

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n, \quad a-R < x < a+R.$$

Theorem (The Term-by-Term Differentiation Theorem)

If $\sum_{n=0}^{\infty} c_n(x-a)^n$ converges for $a-R < x < a+R$ for some $R > 0$, it defines a function f :

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n, \quad a-R < x < a+R.$$

This function f has derivatives all orders inside the interval of convergence. We can obtain the derivatives by differentiating the original series term by term:

Theorem (The Term-by-Term Differentiation Theorem)

If $\sum_{n=0}^{\infty} c_n(x-a)^n$ converges for $a-R < x < a+R$ for some $R > 0$, it defines a function f :

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n, \quad a-R < x < a+R.$$

This function f has derivatives all orders inside the interval of convergence. We can obtain the derivatives by differentiating the original series term by term:

$$f'(x) = \sum_{n=1}^{\infty} n c_n(x-a)^{n-1},$$

$$f''(x) = \sum_{n=2}^{\infty} n(n-1) c_n(x-a)^{n-2},$$

and so on.

Theorem (The Term-by-Term Differentiation Theorem)

If $\sum_{n=0}^{\infty} c_n(x-a)^n$ converges for $a-R < x < a+R$ for some $R > 0$, it defines a function f :

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n, \quad a-R < x < a+R.$$

This function f has derivatives all orders inside the interval of convergence. We can obtain the derivatives by differentiating the original series term by term:

$$f'(x) = \sum_{n=1}^{\infty} n c_n(x-a)^{n-1},$$

$$f''(x) = \sum_{n=2}^{\infty} n(n-1) c_n(x-a)^{n-2},$$

and so on. Each of these derived series converges at every interior point of the interval of convergence of the original series.

Example

Find series for $f'(x)$ and $f''(x)$ if

$$f(x) = \frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots, \quad -1 < x < 1.$$

Example

Find series for $f'(x)$ and $f''(x)$ if

$$f(x) = \frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots, \quad -1 < x < 1.$$

Solution:

$$f'(x) = \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots + nx^{n-1} + \dots, \quad -1 < x < 1.$$

Example

Find series for $f'(x)$ and $f''(x)$ if

$$f(x) = \frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots, \quad -1 < x < 1.$$

Solution:

$$f'(x) = \frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots + nx^{n-1} + \dots, \quad -1 < x < 1.$$

$$f''(x) = \frac{2}{(1-x)^3} = 2 + 6x + 12x^2 + \dots + n(n-1)x^{n-2} + \dots, \quad -1 < x < 1.$$

Theorem (The Term-by-Term Integration Theorem)

Suppose that

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n$$

converges for $a - R < x < a + R$ ($R > 0$).

Theorem (The Term-by-Term Integration Theorem)

Suppose that

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n$$

converges for $a - R < x < a + R$ ($R > 0$). Then

$$\sum_{n=0}^{\infty} c_n \frac{(x - a)^{n+1}}{n + 1}$$

converges for $a - R < x < a + R$ and

$$\int f(x) dx = \sum_{n=0}^{\infty} c_n \frac{(x - a)^{n+1}}{n + 1} + C$$

for $a - R < x < a + R$.

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Solution: Differentiating the given series, we get

$$f'(x) = 1 - x^2 + x^3 - x^4 + \dots, \quad -1 < x < 1.$$

This is a geometric series with first term 1 and common ratio $-x^2$.

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Solution: Differentiating the given series, we get

$$f'(x) = 1 - x^2 + x^3 - x^4 + \dots, \quad -1 < x < 1.$$

This is a geometric series with first term 1 and common ratio $-x^2$. So,

$$f'(x) = \frac{1}{1 - (-x^2)} = \frac{1}{1 + x^2}.$$

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Solution: Differentiating the given series, we get

$$f'(x) = 1 - x^2 + x^3 - x^4 + \dots, \quad -1 < x < 1.$$

This is a geometric series with first term 1 and common ratio $-x^2$. So,

$$f'(x) = \frac{1}{1 - (-x^2)} = \frac{1}{1 + x^2}.$$

Now integration gives

$$f(x) = \int f'(x) dx = \int \frac{dx}{1 + x^2} = \tan^{-1} x + C$$

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Solution: Differentiating the given series, we get

$$f'(x) = 1 - x^2 + x^3 - x^4 + \dots, \quad -1 < x < 1.$$

This is a geometric series with first term 1 and common ratio $-x^2$. So,

$$f'(x) = \frac{1}{1 - (-x^2)} = \frac{1}{1 + x^2}.$$

Now integration gives

$$f(x) = \int f'(x) dx = \int \frac{dx}{1 + x^2} = \tan^{-1} x + C$$

The series for $f(x)$ is zero when $x = 0$; so $C = 0$.

Example: A series for $\tan^{-1} x$

Identify the function

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots, \quad -1 \leq x \leq 1.$$

Solution: Differentiating the given series, we get

$$f'(x) = 1 - x^2 + x^3 - x^4 + \dots, \quad -1 < x < 1.$$

This is a geometric series with first term 1 and common ratio $-x^2$. So,

$$f'(x) = \frac{1}{1 - (-x^2)} = \frac{1}{1 + x^2}.$$

Now integration gives

$$f(x) = \int f'(x) dx = \int \frac{dx}{1 + x^2} = \tan^{-1} x + C$$

The series for $f(x)$ is zero when $x = 0$; so $C = 0$.

Hence

$$f(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots = \tan^{-1} x, \quad -1 < x < 1.$$

Example: A series for $\ln(1+x)$

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges for $-1 < t < 1$.

Example: A series for $\ln(1+x)$

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges for $-1 < t < 1$. Therefore

$$\ln(1+x) = \int_0^x \frac{1}{1+t} dt$$

Example: A series for $\ln(1+x)$

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges for $-1 < t < 1$. Therefore

$$\begin{aligned}\ln(1+x) &= \int_0^x \frac{1}{1+t} dt \\ &= \left[t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \dots \right]_0^x\end{aligned}$$

Example: A series for $\ln(1+x)$

The series

$$\frac{1}{1+t} = 1 - t + t^2 - t^3 + \dots$$

converges for $-1 < t < 1$. Therefore

$$\begin{aligned}\ln(1+x) &= \int_0^x \frac{1}{1+t} dt \\ &= \left[t - \frac{t^2}{2} + \frac{t^3}{3} - \frac{t^4}{4} + \dots \right]_0^x \\ &= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots, \quad -1 < x < 1.\end{aligned}$$

Theorem (The Multiplication Theorem for Power Series)

If $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} b_n x^n$ converge absolutely for $|x| < R$ ($R > 0$)

Theorem (The Multiplication Theorem for Power Series)

If $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} b_n x^n$ converge absolutely for $|x| < R$ ($R > 0$) and

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_{n-1} b_1 + a_n b_0 = \sum_{k=0}^n a_k b_{n-k},$$

Theorem (The Multiplication Theorem for Power Series)

If $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} b_n x^n$ converge absolutely for $|x| < R$ ($R > 0$) and

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_{n-1} b_1 + a_n b_0 = \sum_{k=0}^n a_k b_{n-k},$$

then $\sum_{n=0}^{\infty} c_n x^n$ converges absolutely to $A(x)B(x)$ for $|x| < R$:

Theorem (The Multiplication Theorem for Power Series)

If $A(x) = \sum_{n=0}^{\infty} a_n x^n$ and $B(x) = \sum_{n=0}^{\infty} b_n x^n$ converge absolutely for $|x| < R$ ($R > 0$) and

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_{n-1} b_1 + a_n b_0 = \sum_{k=0}^n a_k b_{n-k},$$

then $\sum_{n=0}^{\infty} c_n x^n$ converges absolutely to $A(x)B(x)$ for $|x| < R$:

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) \cdot \left(\sum_{n=0}^{\infty} b_n x^n \right) = \sum_{n=0}^{\infty} c_n x^n.$$

Example

Multiply the geometric series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots = \frac{1}{1-x} \quad (|x| < 1)$$

by itself to get a power series for $\frac{1}{(1-x)^2}$ for $|x| < 1$.

Example

Multiply the geometric series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots = \frac{1}{1-x} \quad (|x| < 1)$$

by itself to get a power series for $\frac{1}{(1-x)^2}$ for $|x| < 1$.

Solution: Let

$$A(x) = \sum_{n=0}^{\infty} a_n x^n = 1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}$$

Example

Multiply the geometric series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots = \frac{1}{1-x} \quad (|x| < 1)$$

by itself to get a power series for $\frac{1}{(1-x)^2}$ for $|x| < 1$.

Solution: Let

$$A(x) = \sum_{n=0}^{\infty} a_n x^n = 1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}$$

$$B(x) = \sum_{n=0}^{\infty} b_n x^n = 1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}$$

Example

Multiply the geometric series

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots = \frac{1}{1-x} \quad (|x| < 1)$$

by itself to get a power series for $\frac{1}{(1-x)^2}$ for $|x| < 1$.

Solution: Let

$$A(x) = \sum_{n=0}^{\infty} a_n x^n = 1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}$$

$$B(x) = \sum_{n=0}^{\infty} b_n x^n = 1 + x + x^2 + \dots + x^n + \dots = \frac{1}{1-x}$$

and

$$\begin{aligned} c_n &= a_n b_n = a_n b_n + a_1 b_{n-1} + a_2 b_{n-2} + \dots + a_{n-1} b_1 + a_n b_0 \\ &= 1 + 1 + \dots + 1 = n + 1 \end{aligned}$$

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$ and

$$f'(x) = a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots$$

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$ and

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots \\ f''(x) &= 1 \cdot 2a_2 + 2 \cdot 3a_3(x-a) + 3 \cdot 4a_4(x-a)^2 + \dots \end{aligned}$$

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$ and

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots \\ f''(x) &= 1 \cdot 2a_2 + 2 \cdot 3a_3(x-a) + 3 \cdot 4a_4(x-a)^2 + \dots \\ f'''(x) &= 1 \cdot 2 \cdot 3a_3 + 2 \cdot 3 \cdot 4a_4(x-a) + 3 \cdot 4 \cdot 5a_5(x-a)^2 + \dots \end{aligned}$$

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$ and

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots \\ f''(x) &= 1 \cdot 2a_2 + 2 \cdot 3a_3(x-a) + 3 \cdot 4a_4(x-a)^2 + \dots \\ f'''(x) &= 1 \cdot 2 \cdot 3a_3 + 2 \cdot 3 \cdot 4a_4(x-a) + 3 \cdot 4 \cdot 5a_5(x-a)^2 + \dots \end{aligned}$$

and so on.

Taylor and Maclaurin Series

Suppose that

$$f(x) = \sum_{n=0}^{\infty} a_n(x-a)^n = a_0 + a_1(x-a) + a_2(x-a)^2 + \dots + a_n(x-a)^n + \dots$$

converges for $a - R < x < a + R$ ($R > 0$).

Then the Term-by-Term Differentiation Theorem tells us that the sum function has derivatives of all orders within the interval of convergence $a - R < x < a + R$ and

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + \dots \\ f''(x) &= 1 \cdot 2a_2 + 2 \cdot 3a_3(x-a) + 3 \cdot 4a_4(x-a)^2 + \dots \\ f'''(x) &= 1 \cdot 2 \cdot 3a_3 + 2 \cdot 3 \cdot 4a_4(x-a) + 3 \cdot 4 \cdot 5a_5(x-a)^2 + \dots \end{aligned}$$

and so on. The above equations all hold, in particular, at $x = a$.

50

$$f(a) = a_0$$



5

$$\begin{array}{rcl} f(a) & = & a_0 \\ f'(a) & = & a_1 \end{array}$$

So,

$$\begin{aligned}f(a) &= a_0 \\f'(a) &= a_1 \\f''(a) &= 1 \cdot 2a_2\end{aligned}$$

So,

$$\begin{aligned} f(a) &= a_0 \\ f'(a) &= a_1 \\ f''(a) &= 1 \cdot 2a_2 \\ f'''(a) &= 1 \cdot 2 \cdot 3a_3 \end{aligned}$$

5

$$\begin{aligned} f(a) &= a_0 \\ f'(a) &= a_1 \\ f''(a) &= 1 \cdot 2a_2 \\ f'''(a) &= 1 \cdot 2 \cdot 3a_3 \\ &\vdots \\ f^{(n)}(a) &= n!a_n \end{aligned}$$

So,

$$\begin{aligned} f(a) &= a_0 \\ f'(a) &= a_1 \\ f''(a) &= 1 \cdot 2a_2 \\ f'''(a) &= 1 \cdot 2 \cdot 3a_3 \\ &\vdots \\ f^{(n)}(a) &= n!a_n. \end{aligned}$$

Thus

$$a_n = \frac{f^{(n)}(a)}{n!} \quad \text{for } n = 0, 1, 2, \dots$$

So,

$$\begin{aligned} f(a) &= a_0 \\ f'(a) &= a_1 \\ f''(a) &= 1 \cdot 2a_2 \\ f'''(a) &= 1 \cdot 2 \cdot 3a_3 \\ &\vdots \\ f^{(n)}(a) &= n!a_n. \end{aligned}$$

Thus

$$a_n = \frac{f^{(n)}(a)}{n!} \quad \text{for } n = 0, 1, 2, \dots$$

Hence

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots$$

So,


$$\begin{aligned}f(a) &= a_0 \\f'(a) &= a_1 \\f''(a) &= 1 \cdot 2a_2 \\f'''(a) &= 1 \cdot 2 \cdot 3a_3 \\&\vdots \\f^{(n)}(a) &= n!a_n.\end{aligned}$$

Thus

$$a_n = \frac{f^{(n)}(a)}{n!} \quad \text{for } n = 0, 1, 2, \dots$$

Hence

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots$$

This implies that the sum function $f(x)$ has a unique power series expansion. (Why?) 

Definition (Taylor Series, Maclaurin Series)

Let f be a function with derivatives of all orders throughout some interval containing a as an interior point.

Definition (Taylor Series, Maclaurin Series)

Let f be a function with derivatives of all orders throughout some interval containing a as an interior point. Then the **Taylor series generated by f at $x = a$** is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!} (x-a)^n + \dots$$

Definition (Taylor Series, Maclaurin Series)

Let f be a function with derivatives of all orders throughout some interval containing a as an interior point. Then the **Taylor series generated by f at $x = a$** is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!} (x-a)^n + \dots$$

The **Maclaurin series generated by f** is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + f'(0)x + \frac{f''(0)}{2!} x^2 + \dots + \frac{f^{(n)}(0)}{n!} x^n + \dots,$$

which is the Taylor series generated by f at $x = 0$.

Definition (Taylor Series, Maclaurin Series)

Let f be a function with derivatives of all orders throughout some interval containing a as an interior point. Then the **Taylor series generated by f at $x = a$** is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!} (x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!} (x-a)^n + \dots$$

The **Maclaurin series generated by f** is

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = f(0) + f'(0)x + \frac{f''(0)}{2!} x^2 + \dots + \frac{f^{(n)}(0)}{n!} x^n + \dots,$$

which is the Taylor series generated by f at $x = 0$.

Note: The Maclaurin series generated by f is often called the Taylor series of f .

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots,$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n!x^{-(n+1)}, \quad \dots$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2}, \quad f'(2) = -\frac{1}{2^2},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2}, \quad f'(2) = -\frac{1}{2^2}, \quad f''(2) = 2! \frac{1}{2^3},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2}, \quad f'(2) = -\frac{1}{2^2}, \quad f''(2) = 2! \frac{1}{2^3}, \quad f'''(2) = -3! \frac{1}{2^4},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2}, \quad f'(2) = -\frac{1}{2^2}, \quad f''(2) = 2! \frac{1}{2^3}, \quad f'''(2) = -3! \frac{1}{2^4}, \quad \dots, \quad f^{(n)}(2) = (-1)^n n! \frac{1}{2^{n+1}},$$

Example

Find the Taylor series generated by $f(x) = 1/x$ at $a = 2$. Where, if anywhere, does the series converge to $f(x)$?

Solution: We need to find $f(2), f'(2), f''(2), \dots$. Taking derivatives, we get

$$f(x) = x^{-1}, \quad f'(x) = -x^{-2}, \quad f''(x) = 2!x^{-3}, \quad f'''(x) = -3!x^{-4}, \quad \dots, \quad f^{(n)}(x) = (-1)^n n! x^{-(n+1)}, \quad \dots$$

So,

$$f(2) = \frac{1}{2}, \quad f'(2) = -\frac{1}{2^2}, \quad f''(2) = 2! \frac{1}{2^3}, \quad f'''(2) = -3! \frac{1}{2^4}, \quad \dots, \quad f^{(n)}(2) = (-1)^n n! \frac{1}{2^{n+1}}, \quad \dots$$

Hence the Taylor series is

$$f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots$$

Hence the Taylor series is

$$\begin{aligned} f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots \\ = \frac{1}{2} - \frac{(x-2)}{2^2} + \frac{(x-2)^2}{2^3} - \dots + (-1)^n \frac{(x-2)^n}{2^{n+1}} + \dots \end{aligned}$$

Hence the Taylor series is

$$\begin{aligned} f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots \\ = \frac{1}{2} - \frac{(x-2)}{2^2} + \frac{(x-2)^2}{2^3} - \dots + (-1)^n \frac{(x-2)^n}{2^{n+1}} + \dots \end{aligned}$$

This is a geometric series with first term $1/2$ and common ratio $r = -(x-2)/2$.

Hence the Taylor series is

$$\begin{aligned} f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots \\ = \frac{1}{2} - \frac{(x-2)}{2^2} + \frac{(x-2)^2}{2^3} - \dots + (-1)^n \frac{(x-2)^n}{2^{n+1}} + \dots \end{aligned}$$

This is a geometric series with first term $1/2$ and common ratio $r = -(x-2)/2$. It converges absolutely for $|x-2| < 2$.

Hence the Taylor series is

$$\begin{aligned} f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots \\ = \frac{1}{2} - \frac{(x-2)}{2^2} + \frac{(x-2)^2}{2^3} - \dots + (-1)^n \frac{(x-2)^n}{2^{n+1}} + \dots \end{aligned}$$

This is a geometric series with first term $1/2$ and common ratio $r = -(x-2)/2$. It converges absolutely for $|x-2| < 2$. The sum is

$$\frac{1/2}{1 + (x-2)/2} = \frac{1}{x}.$$

Hence the Taylor series is

$$\begin{aligned} f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \dots + \frac{f^{(n)}(2)}{n!}(x-2)^n + \dots \\ = \frac{1}{2} - \frac{(x-2)}{2^2} + \frac{(x-2)^2}{2^3} - \dots + (-1)^n \frac{(x-2)^n}{2^{n+1}} + \dots \end{aligned}$$

This is a geometric series with first term $1/2$ and common ratio $r = -(x-2)/2$. It converges absolutely for $|x-2| < 2$. The sum is

$$\frac{1/2}{1 + (x-2)/2} = \frac{1}{x}.$$

Thus the Taylor series generated by $f(x) = 1/x$ at $a = 2$ converges to $1/x$ for $|x-2| < 2$ or $0 < x < 4$.

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$.

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$. Indeed, $f^{(n)}(0) = 0$ for all n .

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$. Indeed, $f^{(n)}(0) = 0$ for all n .

Thus the Taylor series generated by f at $x = 0$ is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$. Indeed, $f^{(n)}(0) = 0$ for all n .

Thus the Taylor series generated by f at $x = 0$ is

$$\begin{aligned} f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots \\ = 0 + 0 \cdot x + 0 \cdot x^2 + \dots \cdot 0 \cdot x^n + \dots \end{aligned}$$

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$. Indeed, $f^{(n)}(0) = 0$ for all n .

Thus the Taylor series generated by f at $x = 0$ is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

$$= 0 + 0 \cdot x + 0 \cdot x^2 + \dots \cdot 0 \cdot x^n + \dots$$

$$= 0 + 0 + \dots + 0 + \dots$$

The series converges for every x (the sum is 0)

Example

The function

$$f(x) = \begin{cases} 0, & x = 0 \\ e^{-1/x^2}, & x \neq 0 \end{cases}$$

has derivatives of all orders at $x = 0$. Indeed, $f^{(n)}(0) = 0$ for all n .

Thus the Taylor series generated by f at $x = 0$ is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

$$= 0 + 0 \cdot x + 0 \cdot x^2 + \dots \cdot 0 \cdot x^n + \dots$$

$$= 0 + 0 + \dots + 0 + \dots$$

The series converges for every x (the sum is 0) but converges to $f(x)$ only at $x = 0$.

Homework

Find the Taylor series generated by the following functions at $x = 0$:

