

Inner product spaces

Dr Nachiketa Mishra
IIITDM Kancheepuram, Chennai

Introducing (1) length and (2) angle (orthogonal) on vector spaces over R or C .

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F .

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F . An **inner product** on V is a function $\langle \rangle : V \times V \longrightarrow F$ such a way that for all $\alpha, \beta, \gamma \in V$ and all $c \in F$,

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F . An **inner product** on V is a function $\langle \rangle : V \times V \longrightarrow F$ such a way that for all $\alpha, \beta, \gamma \in V$ and all $c \in F$,

$$(1) \quad \langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$$

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F . An **inner product** on V is a function $\langle \rangle : V \times V \longrightarrow F$ such a way that for all $\alpha, \beta, \gamma \in V$ and all $c \in F$,

$$(1) \quad \langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$$

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F . An **inner product** on V is a function $\langle \rangle : V \times V \longrightarrow F$ such a way that for all $\alpha, \beta, \gamma \in V$ and all $c \in F$,

- (1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle$;
- (2) $\langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle$;
- (3) $\langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle}$; (Complex conjugate)

Inner product

Let F be the field of real numbers or the field of complex numbers, and V a vector space over F . An **inner product** on V is a function $\langle \rangle : V \times V \longrightarrow F$ such a way that for all $\alpha, \beta, \gamma \in V$ and all $c \in F$,

- (1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle$;
- (2) $\langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle$;
- (3) $\langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle}$; (Complex conjugate)
- (4) $\langle \alpha, \alpha \rangle > 0$ if $\alpha \neq 0$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$.

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n}$$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

(1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

(1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$

$\alpha + \beta = (x_1 + y_1, \dots, x_n + y_n).$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

(1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$

$$\alpha + \beta = (x_1 + y_1, \dots, x_n + y_n).$$

$$\langle \alpha + \beta, \gamma \rangle = (x_1 + y_1) \overline{z_1} + (x_2 + y_2) \overline{z_2} + \dots + (x_n + y_n) \overline{z_n}$$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

(1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$

$$\alpha + \beta = (x_1 + y_1, \dots, x_n + y_n).$$

$$\langle \alpha + \beta, \gamma \rangle = (x_1 + y_1) \overline{z_1} + (x_2 + y_2) \overline{z_2} + \dots + (x_n + y_n) \overline{z_n} = \sum_{j=1}^n (x_j + y_j) \overline{z_j}$$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

(1) $\langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$

$$\alpha + \beta = (x_1 + y_1, \dots, x_n + y_n).$$

$$\langle \alpha + \beta, \gamma \rangle = (x_1 + y_1) \overline{z_1} + (x_2 + y_2) \overline{z_2} + \dots + (x_n + y_n) \overline{z_n} = \sum_{j=1}^n (x_j + y_j) \overline{z_j}$$

$$\langle \alpha + \beta, \gamma \rangle = \sum_{j=1}^n x_j \overline{z_j} + \sum_{j=1}^n y_j \overline{z_j}$$

Standard inner product

Let $V = F^n = \{(x_1, x_2, \dots, x_n) : x_i \in F\}$. Let $\alpha = (x_1, x_2, \dots, x_n)$, $\beta = (y_1, y_2, \dots, y_n)$ and $\gamma = (z_1, z_2, \dots, z_n)$ be vectors in V .

$$\langle \alpha, \beta \rangle = x_1 \overline{y_1} + x_2 \overline{y_2} + \dots + x_n \overline{y_n} = \sum_{j=1}^n x_j \overline{y_j}$$

Show that $\langle \rangle$ is an inner product.

$$(1) \quad \langle \alpha + \beta, \gamma \rangle = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle;$$

$$\alpha + \beta = (x_1 + y_1, \dots, x_n + y_n).$$

$$\langle \alpha + \beta, \gamma \rangle = (x_1 + y_1) \overline{z_1} + (x_2 + y_2) \overline{z_2} + \dots + (x_n + y_n) \overline{z_n} = \sum_{j=1}^n (x_j + y_j) \overline{z_j}$$

$$\langle \alpha + \beta, \gamma \rangle = \sum_{j=1}^n x_j \overline{z_j} + \sum_{j=1}^n y_j \overline{z_j} = \langle \alpha, \gamma \rangle + \langle \beta, \gamma \rangle$$

Standard inner product contd.

$$(2) \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j}$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j}$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

Standard inner product contd.

(2) $\langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

(3) $\langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}}$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j}$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j} = \langle \beta, \alpha \rangle$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j} = \langle \beta, \alpha \rangle$$

$$(4) \quad \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j} = \langle \beta, \alpha \rangle$$

$$(4) \quad \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0$$

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j}$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j} = \langle \beta, \alpha \rangle$$

$$(4) \quad \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0$$

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n |x_j|^2$$

Standard inner product contd.

$$(2) \quad \langle c\alpha, \beta \rangle = c\langle \alpha, \beta \rangle;$$

$$c\alpha = (cx_1, cx_2, \dots, cx_n)$$

$$\langle c\alpha, \beta \rangle = \sum_{j=1}^n cx_j \overline{y_j} = c \sum_{j=1}^n x_j \overline{y_j} = c\langle \alpha, \beta \rangle$$

$$(3) \quad \langle \beta, \alpha \rangle = \overline{\langle \alpha, \beta \rangle};$$

$$\overline{\langle \alpha, \beta \rangle} = \overline{\sum_{j=1}^n x_j \overline{y_j}} = \sum_{j=1}^n y_j \overline{x_j} = \langle \beta, \alpha \rangle$$

$$(4) \quad \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0$$

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n |x_j|^2 > 0, \quad \text{provided } \alpha \neq 0$$

Problem 1

Let $V = \mathbb{R}^2$. Let $\alpha = (x_1, x_2)$ and $\beta = (y_1, y_2)$ be two vectors in \mathbb{R}^2 .

Problem 1

Let $V = \mathbb{R}^2$. Let $\alpha = (x_1, x_2)$ and $\beta = (y_1, y_2)$ be two vectors in \mathbb{R}^2 .

$$\langle \alpha, \beta \rangle = x_1 y_1 - x_2 y_1 - x_1 y_2 + 4x_2 y_2$$

Prove that $\langle \rangle$ is an inner product. (Please do it now)

Problem 1

Let $V = R^2$. Let $\alpha = (x_1, x_2)$ and $\beta = (y_1, y_2)$ be two vectors in R^2 .

$$\langle \alpha, \beta \rangle = x_1 y_1 - x_2 y_1 - x_1 y_2 + 4x_2 y_2$$

Prove that $\langle \rangle$ is an inner product. (Please do it now)

Note that $\langle \alpha, \alpha \rangle = (x_1 - x_2)^2 + 3x_2^2$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle$$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle = \overline{\langle c\beta, \alpha \rangle}$$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle = \overline{\langle c\beta, \alpha \rangle} = \overline{c \langle \beta, \alpha \rangle}$$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle = \overline{\langle c\beta, \alpha \rangle} = \overline{c \langle \beta, \alpha \rangle} = \overline{c} \overline{\langle \beta, \alpha \rangle}$$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle = \overline{\langle c\beta, \alpha \rangle} = \overline{c \langle \beta, \alpha \rangle} = \overline{c} \overline{\langle \beta, \alpha \rangle} = \overline{c} \langle \alpha, \beta \rangle$$

Problem 2

(1) Prove that $\langle \alpha, c\beta \rangle = \overline{c} \langle \alpha, \beta \rangle$

$$\langle \alpha, c\beta \rangle = \overline{\langle c\beta, \alpha \rangle} = \overline{c \langle \beta, \alpha \rangle} = \overline{c} \overline{\langle \beta, \alpha \rangle} = \overline{c} \langle \alpha, \beta \rangle$$

(2) Prove that $c \langle \alpha, \beta \rangle = \langle \alpha, \overline{c}\beta \rangle$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle}$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle}$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

Let $z = x + iy$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

Let $z = x + iy = \operatorname{Re} z + i \operatorname{Im} z$,

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

Let $z = x + iy = \operatorname{Re} z + i \operatorname{Im} z$, $\implies z + \bar{z} = 2 \operatorname{Re} z$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

Let $z = x + iy = \operatorname{Re} z + i \operatorname{Im} z$, $\implies z + \bar{z} = 2 \operatorname{Re} z$

$-iz = -ix + y$, $\implies \operatorname{Re}(-iz) = \operatorname{Im} z$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

Let $z = x + iy = \operatorname{Re} z + i \operatorname{Im} z$, $\implies z + \bar{z} = 2 \operatorname{Re} z$

$-iz = -ix + y$, $\implies \operatorname{Re}(-iz) = \operatorname{Im} z$

$$\langle \alpha, \beta \rangle = \operatorname{Re} \langle \alpha, \beta \rangle + i \operatorname{Im} \langle \alpha, \beta \rangle$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

$$\text{Let } z = x + iy = \operatorname{Re} z + i \operatorname{Im} z, \implies z + \bar{z} = 2 \operatorname{Re} z$$

$$-iz = -ix + y, \implies \operatorname{Re}(-iz) = \operatorname{Im} z$$

$$\langle \alpha, \beta \rangle = \operatorname{Re} \langle \alpha, \beta \rangle + i \operatorname{Im} \langle \alpha, \beta \rangle$$

$$\operatorname{Im} \langle \alpha, \beta \rangle = \operatorname{Re}(-i \langle \alpha, \beta \rangle)$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

$$\text{Let } z = x + iy = \operatorname{Re} z + i \operatorname{Im} z, \implies z + \bar{z} = 2 \operatorname{Re} z$$

$$-iz = -ix + y, \implies \operatorname{Re}(-iz) = \operatorname{Im} z$$

$$\langle \alpha, \beta \rangle = \operatorname{Re} \langle \alpha, \beta \rangle + i \operatorname{Im} \langle \alpha, \beta \rangle$$

$$\operatorname{Im} \langle \alpha, \beta \rangle = \operatorname{Re}(-i \langle \alpha, \beta \rangle) = \operatorname{Re} \langle \alpha, i\beta \rangle$$

Note 1

$$\langle \alpha, \beta + \gamma \rangle = \overline{\langle \beta + \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle + \langle \gamma, \alpha \rangle} = \overline{\langle \beta, \alpha \rangle} + \overline{\langle \gamma, \alpha \rangle}$$

$$\langle \alpha, \beta + \gamma \rangle = \langle \alpha, \beta \rangle + \langle \alpha, \gamma \rangle$$

$$\text{Let } z = x + iy = \operatorname{Re} z + i \operatorname{Im} z, \implies z + \bar{z} = 2 \operatorname{Re} z$$

$$-iz = -ix + y, \implies \operatorname{Re}(-iz) = \operatorname{Im} z$$

$$\langle \alpha, \beta \rangle = \operatorname{Re} \langle \alpha, \beta \rangle + i \operatorname{Im} \langle \alpha, \beta \rangle$$

$$\operatorname{Im} \langle \alpha, \beta \rangle = \operatorname{Re}(-i \langle \alpha, \beta \rangle) = \operatorname{Re} \langle \alpha, i\beta \rangle$$

$$\implies \langle \alpha, \beta \rangle = \operatorname{Re} \langle \alpha, \beta \rangle + i \operatorname{Re} \langle \alpha, i\beta \rangle$$

Norm (length)

$$\|\alpha\|$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle} , \quad (\text{Positive square root})$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle} , \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}, \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j}$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}, \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n x_j x_j$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}, \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n x_j x_j = \sum_{j=1}^n x_j^2$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}, \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n x_j x_j = \sum_{j=1}^n x_j^2$$

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}$$

Norm (length)

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle}, \quad (\text{Positive square root})$$

Consider the standard inner product on R^n

$$\langle \alpha, \alpha \rangle = \sum_{j=1}^n x_j \overline{x_j} = \sum_{j=1}^n x_j x_j = \sum_{j=1}^n x_j^2$$

$$\|\alpha\| = \sqrt{\langle \alpha, \alpha \rangle} = \sqrt{\sum_{j=1}^n x_j^2} \quad (\text{length of the vector } \alpha)$$

$$\|\alpha\|^2$$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

Note 2

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2$$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle$$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle = \langle \alpha, \alpha + \beta \rangle + \langle \beta, \alpha + \beta \rangle$$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle = \langle \alpha, \alpha + \beta \rangle + \langle \beta, \alpha + \beta \rangle$$

$$= (\langle \alpha, \alpha \rangle + \langle \alpha, \beta \rangle) + (\langle \beta, \alpha \rangle + \langle \beta, \beta \rangle)$$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle = \langle \alpha, \alpha + \beta \rangle + \langle \beta, \alpha + \beta \rangle$$

$$= (\langle \alpha, \alpha \rangle + \langle \alpha, \beta \rangle) + (\langle \beta, \alpha \rangle + \langle \beta, \beta \rangle)$$

$$= \|\alpha\|^2 + \|\beta\|^2 + \langle \alpha, \beta \rangle + \langle \beta, \alpha \rangle$$

Note 2

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle = \langle \alpha, \alpha + \beta \rangle + \langle \beta, \alpha + \beta \rangle$$

$$= (\langle \alpha, \alpha \rangle + \langle \alpha, \beta \rangle) + (\langle \beta, \alpha \rangle + \langle \beta, \beta \rangle)$$

$$= \|\alpha\|^2 + \|\beta\|^2 + \langle \alpha, \beta \rangle + \langle \beta, \alpha \rangle$$

$$= \|\alpha\|^2 + \|\beta\|^2 + \langle \alpha, \beta \rangle + \overline{\langle \alpha, \beta \rangle}$$

Note 2

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle$$

$$\|\alpha + \beta\|^2 = \langle \alpha + \beta, \alpha + \beta \rangle = \langle \alpha, \alpha + \beta \rangle + \langle \beta, \alpha + \beta \rangle$$

$$= (\langle \alpha, \alpha \rangle + \langle \alpha, \beta \rangle) + (\langle \beta, \alpha \rangle + \langle \beta, \beta \rangle)$$

$$= \|\alpha\|^2 + \|\beta\|^2 + \langle \alpha, \beta \rangle + \langle \beta, \alpha \rangle$$

$$= \|\alpha\|^2 + \|\beta\|^2 + \langle \alpha, \beta \rangle + \overline{\langle \alpha, \beta \rangle}$$

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

Note 2 contd.

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

Using similar arguments,

$$\|\alpha - \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 - 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

Note 2 contd.

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

Using similar arguments,

$$\|\alpha - \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 - 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\|\alpha + \beta\|^2 + \|\alpha - \beta\|^2$$

Note 2 contd.

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

Using similar arguments,

$$\|\alpha - \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 - 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\|\alpha + \beta\|^2 + \|\alpha - \beta\|^2 = 2 (\|\alpha\|^2 + \|\beta\|^2)$$

(Parallelogram law)

Note 2 contd.

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$


Using similar arguments,

$$\|\alpha - \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 - 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\|\alpha + \beta\|^2 + \|\alpha - \beta\|^2 = 2 (\|\alpha\|^2 + \|\beta\|^2)$$

(Parallelogram law)

The sum of the squares of the lengths of the four sides of a parallelogram equals the sum of the squares of the lengths of the two diagonals



Problem

Show that if $F = R$,

$$\langle \alpha, \beta \rangle = \frac{1}{4} \|\alpha + \beta\|^2 - \frac{1}{4} \|\alpha - \beta\|^2$$

Inner Product Spaces

An **inner product space** is a real or complex vector space, together with a specified inner product on that space.

Inner Product Spaces

An **inner product space** is a real or complex vector space, together with a specified inner product on that space.

Note : A finite-dimensional real inner product space is called a **Euclidean space**.

Inner Product Spaces

An **inner product space** is a real or complex vector space, together with a specified inner product on that space.

Note : A finite-dimensional real inner product space is called a **Euclidean space**.

Note : A complex inner product space is called a **unitary space**.

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

$$(i) \quad \|c\alpha\| = |c| \|\alpha\|;$$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

- (i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

- (i) Prove that $\|c\alpha\| = |c| \|\alpha\|$
 $\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

- (i) Prove that $\|c\alpha\| = |c| \|\alpha\|$
 $\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle$$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle = |c|^2\|\alpha\|^2.$$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle = |c|^2\|\alpha\|^2.$$

$$\implies \|c\alpha\| = |c| \|\alpha\|$$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle = |c|^2\|\alpha\|^2.$$

$$\implies \|c\alpha\| = |c| \|\alpha\|$$

(ii) Prove that $\|\alpha\| > 0$ for $\alpha \neq 0$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle = |c|^2\|\alpha\|^2.$$

$$\implies \|c\alpha\| = |c| \|\alpha\|$$

(ii) Prove that $\|\alpha\| > 0$ for $\alpha \neq 0$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0;$$

Theorem 1

If V is an inner product space, then for any vectors $\alpha, \beta \in V$ and any scalar c ,

- (i) $\|c\alpha\| = |c| \|\alpha\|$;
- (ii) $\|\alpha\| > 0$ for $\alpha \neq 0$;
- (iii) $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$; (Cauchy -Schwarz inequality)
- (iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$.

Proof : Note that $\langle 0, \alpha \rangle = \langle 0\alpha, \alpha \rangle = 0\langle \alpha, \alpha \rangle = 0$, and $\|0\| = 0$

(i) Prove that $\|c\alpha\| = |c| \|\alpha\|$

$$\|c\alpha\|^2 = \langle c\alpha, c\alpha \rangle = c\langle \alpha, c\alpha \rangle = c\bar{c}\langle \alpha, \alpha \rangle = |c|^2\|\alpha\|^2.$$

$$\implies \|c\alpha\| = |c| \|\alpha\|$$

(ii) Prove that $\|\alpha\| > 0$ for $\alpha \neq 0$

$$\|\alpha\|^2 = \langle \alpha, \alpha \rangle > 0 \text{ if } \alpha \neq 0; \implies \|\alpha\| > 0 \text{ if } \alpha \neq 0.$$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0$.

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Let

$$\gamma = \beta - \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \alpha$$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Let

$$\gamma = \beta - \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \alpha = \beta - c\alpha$$

where

$$c = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2}$$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Let

$$\gamma = \beta - \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \alpha = \beta - c\alpha$$

where

$$c = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2}$$

$$c\bar{c} = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \frac{\overline{\langle \beta, \alpha \rangle}}{\|\alpha\|^2}$$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Let

$$\gamma = \beta - \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \alpha = \beta - c\alpha$$

where

$$c = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2}$$

$$c\bar{c} = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \frac{\overline{\langle \beta, \alpha \rangle}}{\|\alpha\|^2} = \frac{\overline{\langle \alpha, \beta \rangle}}{\|\alpha\|^2} \frac{\langle \alpha, \beta \rangle}{\|\alpha\|^2}$$

Theorem 1 contd.

(ii) Prove that $|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$

If $\alpha = 0$, then $\langle \alpha, \beta \rangle = \langle 0, \beta \rangle = 0$ and $\|\alpha\| = 0$.

$\implies |\langle \alpha, \beta \rangle| = \|\alpha\| \|\beta\| = 0$.

So $\alpha \neq 0 \implies \|\alpha\|^2 > 0$.

Let

$$\gamma = \beta - \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \alpha = \beta - c\alpha$$

where

$$c = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2}$$

$$c\bar{c} = \frac{\langle \beta, \alpha \rangle}{\|\alpha\|^2} \frac{\overline{\langle \beta, \alpha \rangle}}{\overline{\|\alpha\|^2}} = \frac{\overline{\langle \alpha, \beta \rangle}}{\|\alpha\|^2} \frac{\langle \alpha, \beta \rangle}{\|\alpha\|^2} = \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^4} \text{ --- (1)}$$

Theorem 1 contd.

$$c\langle\alpha,\beta\rangle = \frac{\langle\beta,\alpha\rangle}{\|\alpha\|^2}\langle\alpha,\beta\rangle$$

Theorem 1 contd.

$$c\langle\alpha,\beta\rangle = \frac{\langle\beta,\alpha\rangle}{\|\alpha\|^2}\langle\alpha,\beta\rangle = \frac{\overline{\langle\alpha,\beta\rangle}}{\|\alpha\|^2}\langle\alpha,\beta\rangle$$

Theorem 1 contd.

$$c\langle\alpha,\beta\rangle = \frac{\langle\beta,\alpha\rangle}{\|\alpha\|^2}\langle\alpha,\beta\rangle = \frac{\overline{\langle\alpha,\beta\rangle}}{\|\alpha\|^2}\langle\alpha,\beta\rangle = \frac{|\langle\alpha,\beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\overline{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\overline{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle}$$

Theorem 1 contd.

$$c\langle\alpha,\beta\rangle = \frac{\langle\beta,\alpha\rangle}{\|\alpha\|^2}\langle\alpha,\beta\rangle = \frac{\overline{\langle\alpha,\beta\rangle}}{\|\alpha\|^2}\langle\alpha,\beta\rangle = \frac{|\langle\alpha,\beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\overline{c}\langle\beta,\alpha\rangle = \frac{\overline{\langle\beta,\alpha\rangle}}{\|\alpha\|^2}\langle\beta,\alpha\rangle = \frac{\langle\alpha,\beta\rangle}{\|\alpha\|^2}\overline{\langle\alpha,\beta\rangle} = \frac{|\langle\alpha,\beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\overline{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle} = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Note that

$$0 \leq \|\gamma\|^2 = \|\beta - c\alpha\|^2$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\bar{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle} = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Note that

$$0 \leq \|\gamma\|^2 = \|\beta - c\alpha\|^2$$

$$0 \leq \langle\beta - c\alpha, \beta - c\alpha\rangle$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\overline{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle} = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Note that

$$0 \leq \|\gamma\|^2 = \|\beta - c\alpha\|^2$$

$$\begin{aligned} 0 &\leq \langle\beta - c\alpha, \beta - c\alpha\rangle \\ &\leq \langle\beta, \beta - c\alpha\rangle + \langle -c\alpha, \beta - c\alpha\rangle \end{aligned}$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\bar{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle} = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Note that

$$0 \leq \|\gamma\|^2 = \|\beta - c\alpha\|^2$$

$$\begin{aligned} 0 &\leq \langle\beta - c\alpha, \beta - c\alpha\rangle \\ &\leq \langle\beta, \beta - c\alpha\rangle + \langle -c\alpha, \beta - c\alpha\rangle \\ &\leq \langle\beta, \beta\rangle + \langle\beta, -c\alpha\rangle + \langle -c\alpha, \beta\rangle + \langle -c\alpha, -c\alpha\rangle \end{aligned}$$

Theorem 1 contd.

$$c\langle\alpha, \beta\rangle = \frac{\langle\beta, \alpha\rangle}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{\overline{\langle\alpha, \beta\rangle}}{\|\alpha\|^2} \langle\alpha, \beta\rangle = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (2)}$$

$$\bar{c}\langle\beta, \alpha\rangle = \frac{\overline{\langle\beta, \alpha\rangle}}{\|\alpha\|^2} \langle\beta, \alpha\rangle = \frac{\langle\alpha, \beta\rangle}{\|\alpha\|^2} \overline{\langle\alpha, \beta\rangle} = \frac{|\langle\alpha, \beta\rangle|^2}{\|\alpha\|^2} \text{ --- (3)}$$

Note that

$$0 \leq \|\gamma\|^2 = \|\beta - c\alpha\|^2$$

$$\begin{aligned} 0 &\leq \langle\beta - c\alpha, \beta - c\alpha\rangle \\ &\leq \langle\beta, \beta - c\alpha\rangle + \langle -c\alpha, \beta - c\alpha\rangle \\ &\leq \langle\beta, \beta\rangle + \langle\beta, -c\alpha\rangle + \langle -c\alpha, \beta\rangle + \langle -c\alpha, -c\alpha\rangle \\ &\leq \|\beta\|^2 - \bar{c}\langle\beta, \alpha\rangle - c\langle\alpha, \beta\rangle + c\bar{c}\langle\alpha, \alpha\rangle \end{aligned}$$

Theorem 1 contd.

$$0 \leq \|\beta\|^2 - 2 \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2} + \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^4} \|\alpha\|^2$$

Theorem 1 contd.

$$0 \leq \|\beta\|^2 - 2 \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2} + \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^4} \|\alpha\|^2$$

$$0 \leq \|\beta\|^2 - \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2}$$

Theorem 1 contd.

$$0 \leq \|\beta\|^2 - 2 \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2} + \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^4} \|\alpha\|^2$$

$$0 \leq \|\beta\|^2 - \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2}$$

$$|\langle \alpha, \beta \rangle|^2 \leq \|\alpha\|^2 \|\beta\|^2$$

Theorem 1 contd.

$$0 \leq \|\beta\|^2 - 2 \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2} + \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^4} \|\alpha\|^2$$

$$0 \leq \|\beta\|^2 - \frac{|\langle \alpha, \beta \rangle|^2}{\|\alpha\|^2}$$

$$|\langle \alpha, \beta \rangle|^2 \leq \|\alpha\|^2 \|\beta\|^2$$

$$|\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\| \quad (\text{Cauchy -Schwarz inequality})$$

$$\text{(iv) } \|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|.$$

(iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|.$

We know that

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$(iv) \|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|.$$

We know that

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\operatorname{Re} \langle \alpha, \beta \rangle \leq |\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$$

(iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|.$

We know that

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\operatorname{Re} \langle \alpha, \beta \rangle \leq |\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$$

$$\Rightarrow \|\alpha + \beta\|^2 \leq \|\alpha\|^2 + \|\beta\|^2 + 2\|\alpha\|\|\beta\| = (\|\alpha\| + \|\beta\|)^2$$

(iv) $\|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|.$

We know that

$$\|\alpha + \beta\|^2 = \|\alpha\|^2 + \|\beta\|^2 + 2 \operatorname{Re} \langle \alpha, \beta \rangle$$

$$\operatorname{Re} \langle \alpha, \beta \rangle \leq |\langle \alpha, \beta \rangle| \leq \|\alpha\| \|\beta\|$$

$$\implies \|\alpha + \beta\|^2 \leq \|\alpha\|^2 + \|\beta\|^2 + 2\|\alpha\|\|\beta\| = (\|\alpha\| + \|\beta\|)^2$$

$$\implies \|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

$$\langle \epsilon_1, \epsilon_3 \rangle = \langle \epsilon_2, \epsilon_3 \rangle = 0$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

$$\langle \epsilon_1, \epsilon_3 \rangle = \langle \epsilon_2, \epsilon_3 \rangle = 0$$

$\implies B$ is an orthogonal set.

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

$$\langle \epsilon_1, \epsilon_3 \rangle = \langle \epsilon_2, \epsilon_3 \rangle = 0$$

$\implies B$ is an orthogonal set.

$$\|\epsilon_1\| = \sqrt{\langle \epsilon_1, \epsilon_1 \rangle} = \sqrt{1 \times 1 + 0 \times 0 + 0 \times 0} = 1,$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

$$\langle \epsilon_1, \epsilon_3 \rangle = \langle \epsilon_2, \epsilon_3 \rangle = 0$$

$\implies B$ is an orthogonal set.

$$\begin{aligned}\|\epsilon_1\| &= \sqrt{\langle \epsilon_1, \epsilon_1 \rangle} = \sqrt{1 \times 1 + 0 \times 0 + 0 \times 0} = 1, \quad \|\epsilon_2\| = 1, \\ \|\epsilon_3\| &= 1.\end{aligned}$$

Orthogonal vectors

Let V be an inner product space and $\alpha, \beta \in V$.

We say α is orthogonal to β if $\langle \alpha, \beta \rangle = 0$

Example Consider the Euclidean space R^3 and a ordered basis $B = \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$.

$$\langle \alpha, \beta \rangle = x_1y_1 + x_2y_2 + x_3y_3, \text{ where } \alpha, \beta \in R^3$$

$$\langle \epsilon_1, \epsilon_2 \rangle = 1 \times 0 + 0 \times 1 + 0 \times 0 = 0$$

$$\langle \epsilon_1, \epsilon_3 \rangle = \langle \epsilon_2, \epsilon_3 \rangle = 0$$

$\implies B$ is an orthogonal set.

$$\|\epsilon_1\| = \sqrt{\langle \epsilon_1, \epsilon_1 \rangle} = \sqrt{1 \times 1 + 0 \times 0 + 0 \times 0} = 1, \|\epsilon_2\| = 1,$$

$\|\epsilon_3\| = 1. \implies B$ is an orthonormal set.

Orthogonal set and Orthonormal set

Let V be an inner product space. A set $S \subseteq V$ is called an **orthogonal set** if for all $\alpha, \beta \in S$ where $\alpha \neq \beta$, then $\langle \alpha, \beta \rangle = 0$.

Orthogonal set and Orthonormal set

Let V be an inner product space. A set $S \subseteq V$ is called an **orthogonal set** if for all $\alpha, \beta \in S$ where $\alpha \neq \beta$, then $\langle \alpha, \beta \rangle = 0$.

An **orthogonal set** S of V is **orthonormal** if $\|\alpha\| = 1$ for all $\alpha \in S$.

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Output : An orthogonal basis $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of V .

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Output : An orthogonal basis $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of V .

$$\alpha_1 = \beta_1$$

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Output : An orthogonal basis $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of V .

$$\alpha_1 = \beta_1$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Output : An orthogonal basis $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of V .

$$\alpha_1 = \beta_1$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

Gram - Schmidt orthogonalization process

Input : A basis $\{\beta_1, \beta_2, \dots, \beta_n\}$ of an inner product space V .

Output : An orthogonal basis $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ of V .

$$\alpha_1 = \beta_1$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

$$\alpha_4 = \beta_4 - \frac{\langle \beta_4, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_4, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2 - \frac{\langle \beta_4, \alpha_3 \rangle}{\|\alpha_3\|^2} \alpha_3$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4);$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_2 = (-1, 0, 7) - \frac{\langle (-1, 0, 7), (3, 0, 4) \rangle}{25} (3, 0, 4)$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_2 = (-1, 0, 7) - \frac{\langle (-1, 0, 7), (3, 0, 4) \rangle}{25} (3, 0, 4)$$

$$\alpha_2 = (-1, 0, 7) - \frac{(-1 \times 3 + 0 \times 0 + 7 \times 4)}{25} (3, 0, 4)$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_2 = (-1, 0, 7) - \frac{\langle (-1, 0, 7), (3, 0, 4) \rangle}{25} (3, 0, 4)$$

$$\alpha_2 = (-1, 0, 7) - \frac{(-1 \times 3 + 0 \times 0 + 7 \times 4)}{25} (3, 0, 4)$$

$$\alpha_2 = (-4, 0, 3);$$

Problem 3

Find an orthogonal basis of R^3 with standard inner product from the basis $B = \{\beta_1 = (3, 0, 4), \beta_2 = (-1, 0, 7), \beta_3 = (2, 9, 11)\}$ using Gram-Schmidt process.

$$\alpha_1 = \beta_1 = (3, 0, 4); \quad \|\alpha_1\|^2 = 3^2 + 0^2 + 4^2 = 25$$

$$\alpha_2 = \beta_2 - \frac{\langle \beta_2, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1$$

$$\alpha_2 = (-1, 0, 7) - \frac{\langle (-1, 0, 7), (3, 0, 4) \rangle}{25} (3, 0, 4)$$

$$\alpha_2 = (-1, 0, 7) - \frac{(-1 \times 3 + 0 \times 0 + 7 \times 4)}{25} (3, 0, 4)$$

$$\alpha_2 = (-4, 0, 3); \quad \|\alpha_2\|^2 = (-4)^2 + 0^2 + 3^2 = 25$$

Problem 3 contd.

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

Problem 3 contd.

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

$$\alpha_3 = (2, 9, 11) - \frac{\langle (2, 9, 11), (3, 0, 4) \rangle}{25} (3, 0, 4) - \frac{\langle (2, 9, 11), (-4, 0, 3) \rangle}{25} (-4, 0, 3)$$

Problem 3 contd.

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

$$\alpha_3 = (2, 9, 11) - \frac{\langle (2, 9, 11), (3, 0, 4) \rangle}{25} (3, 0, 4) - \frac{\langle (2, 9, 11), (-4, 0, 3) \rangle}{25} (-4, 0, 3)$$

$$\alpha_3 = (2, 9, 11) - 2(3, 0, 4) - (-4, 0, 3) = (0, 9, 0);$$

Problem 3 contd.

$$\alpha_3 = \beta_3 - \frac{\langle \beta_3, \alpha_1 \rangle}{\|\alpha_1\|^2} \alpha_1 - \frac{\langle \beta_3, \alpha_2 \rangle}{\|\alpha_2\|^2} \alpha_2$$

$$\alpha_3 = (2, 9, 11) - \frac{\langle (2, 9, 11), (3, 0, 4) \rangle}{25} (3, 0, 4) - \frac{\langle (2, 9, 11), (-4, 0, 3) \rangle}{25} (-4, 0, 3)$$

$$\alpha_3 = (2, 9, 11) - 2(3, 0, 4) - (-4, 0, 3) = (0, 9, 0); \quad \|\alpha_3\|^2 = 81$$

Problem 3 contd.

$$B' = \{\alpha_1 = (3, 0, 4), \alpha_2 = (-4, 0, 3), \alpha_3 = (0, 9, 0)\}$$

is an orthogonal basis of R^3 .

Problem 3 contd.

$$B' = \{\alpha_1 = (3, 0, 4), \alpha_2 = (-4, 0, 3), \alpha_3 = (0, 9, 0)\}$$

is an orthogonal basis of R^3 .

Verification

$$\langle \alpha_1, \alpha_2 \rangle = 3 \times (-4) + 0 \times 0 + 4 \times 3 = 0$$

Problem 3 contd.

$$B' = \{\alpha_1 = (3, 0, 4), \alpha_2 = (-4, 0, 3), \alpha_3 = (0, 9, 0)\}$$

is an orthogonal basis of R^3 .

Verification

$$\langle \alpha_1, \alpha_2 \rangle = 3 \times (-4) + 0 \times 0 + 4 \times 3 = 0$$

$$\langle \alpha_1, \alpha_3 \rangle = \langle \alpha_2, \alpha_3 \rangle = 0$$

Problem 3 contd.

$$B' = \{\alpha_1 = (3, 0, 4), \alpha_2 = (-4, 0, 3), \alpha_3 = (0, 9, 0)\}$$

is an orthogonal basis of R^3 .

Verification

$$\langle \alpha_1, \alpha_2 \rangle = 3 \times (-4) + 0 \times 0 + 4 \times 3 = 0$$

$$\langle \alpha_1, \alpha_3 \rangle = \langle \alpha_2, \alpha_3 \rangle = 0$$

Note that B' is a L.I. subset of R^3 and its an orthogonal basis of R^3 .

Problem 3 contd.

$$B'' = \left\{ \frac{\alpha_1}{\|\alpha_1\|}, \frac{\alpha_2}{\|\alpha_2\|}, \frac{\alpha_3}{\|\alpha_3\|} \right\}$$

is an orthonormal basis of R^3 .

Problem 3 contd.

$$B'' = \left\{ \frac{\alpha_1}{\|\alpha_1\|}, \frac{\alpha_2}{\|\alpha_2\|}, \frac{\alpha_3}{\|\alpha_3\|} \right\}$$

is an orthonormal basis of R^3 .

$$B'' = \left\{ \frac{1}{5}(3, 0, 4), \frac{1}{5}(-4, 0, 3), (0, 1, 0) \right\}$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1),$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1), \quad \|\alpha_1\|^2 = 3$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1), \quad \|\alpha_1\|^2 = 3$$

$$\alpha_2 = \left(-\frac{2}{3}, \frac{1}{3}, \frac{1}{3}\right),$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1), \quad \|\alpha_1\|^2 = 3$$

$$\alpha_2 = \left(-\frac{2}{3}, \frac{1}{3}, \frac{1}{3}\right), \quad \|\alpha_2\|^2 = \frac{2}{3}$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1), \quad \|\alpha_1\|^2 = 3$$

$$\alpha_2 = \left(-\frac{2}{3}, \frac{1}{3}, \frac{1}{3}\right), \quad \|\alpha_2\|^2 = \frac{2}{3}$$

$$\alpha_3 = \left(0, -\frac{1}{2}, \frac{1}{2}\right), \quad \|\alpha_3\|^2 = \frac{1}{2}$$

Problem 4

Using Gram-Schmidt process, find an orthonormal basis for the Euclidean space R^3 from a given ordered basis

$$B = \{\beta_1 = (1, 1, 1), \beta_2 = (0, 1, 1), \beta_3 = (0, 0, 1)\}.$$

$$\alpha_1 = \beta_1 = (1, 1, 1), \quad \|\alpha_1\|^2 = 3$$

$$\alpha_2 = \left(-\frac{2}{3}, \frac{1}{3}, \frac{1}{3}\right), \quad \|\alpha_2\|^2 = \frac{2}{3}$$

$$\alpha_3 = \left(0, -\frac{1}{2}, \frac{1}{2}\right), \quad \|\alpha_3\|^2 = \frac{1}{2}$$

The orthonormal basis :

$$\left(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right), \left(-\sqrt{\frac{2}{3}}, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}\right), \left(0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$$