

Elementary row operations

Nachiketa Mishra

IIITDM Kancheepuram, Chennai

Elementary row operations

Consider a matrix $A = [A_{ij}]$, where $A_{ij} \in F$, a field.

The i^{th} row of A is $R_i = [A_{i1}, A_{i2}, \dots, A_{in}]$

Elementary row operations

Consider a matrix $A = [A_{ij}]$, where $A_{ij} \in F$, a field.

The i^{th} row of A is $R_i = [A_{i1}, A_{i2}, \dots, A_{in}]$

Type 1: Multiplication of one row of A by a non-zero scalar

$c \in F$ ($e : R_i \longleftarrow cR_i$)

Elementary row operations

Consider a matrix $A = [A_{ij}]$, where $A_{ij} \in F$, a field.

The i^{th} row of A is $R_i = [A_{i1}, A_{i2}, \dots, A_{in}]$

Type 1: Multiplication of one row of A by a non-zero scalar

$$c \in F \quad (e : R_i \longleftarrow cR_i)$$

Type 2: Replacement of i^{th} row by row i plus c times of row

$$j \text{ where } c \in F \quad (e : R_i \longleftarrow R_i + cR_j)$$

Elementary row operations

Consider a matrix $A = [A_{ij}]$, where $A_{ij} \in F$, a field.

The i^{th} row of A is $R_i = [A_{i1}, A_{i2}, \dots, A_{in}]$

Type 1: Multiplication of one row of A by a non-zero scalar

$c \in F$ ($e : R_i \longleftarrow cR_i$)

Type 2: Replacement of i^{th} row by row i plus c times of row

j where $c \in F$ ($e : R_i \longleftarrow R_i + cR_j$)

Type 3: Interchange of two rows ($e : R_i \longleftrightarrow R_j$)

Elementary row operations

Consider a matrix $A = [A_{ij}]$, where $A_{ij} \in F$, a field.

The i^{th} row of A is $R_i = [A_{i1}, A_{i2}, \dots, A_{in}]$

Type 1: Multiplication of one row of A by a non-zero scalar

$c \in F$ ($e : R_i \longleftarrow cR_i$)

Type 2: Replacement of i^{th} row by row i plus c times of row

j where $c \in F$ ($e : R_i \longleftarrow R_i + cR_j$)

Type 3: Interchange of two rows ($e : R_i \longleftrightarrow R_j$)

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) =$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

$$e_1(e(A)) =$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Prove that $e(e_1(A)) = A$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \longleftarrow \frac{1}{c}R_1$

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Prove that $e(e_1(A)) = A \implies e_1(e(A)) = A = e(e_1(A))$

Inverse of the Type 1 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \leftarrow cR_1, \quad c \neq 0$

$$e(A) = \begin{bmatrix} cA_{11} & cA_{12} & cA_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

We define $e_1 : R_1 \leftarrow \frac{1}{c}R_1 \implies e_1$ is the inverse elementary operation of e

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Prove that $e(e_1(A)) = A \implies e_1(e(A)) = A = e(e_1(A))$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) =$$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

We define $e_1 : R_2 \longleftarrow R_2 - cR_1$.

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

We define $e_1 : R_2 \longleftarrow R_2 - cR_1. \implies e_1(e(A)) =$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

We define $e_1 : R_2 \longleftarrow R_2 - cR_1. \implies e_1(e(A)) = A$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

We define $e_1 : R_2 \longleftarrow R_2 - cR_1. \implies e_1(e(A)) = A$
Similarly, $e(e_1(A)) = A = e_1(e(A))$

Inverse of the Type 2 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_2 \longleftarrow R_2 + cR_1, \quad c \in F$

$$e(A) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} + cA_{11} & A_{22} + cA_{12} & A_{23} + cA_{13} \end{bmatrix}$$

We define $e_1 : R_2 \longleftarrow R_2 - cR_1. \implies e_1(e(A)) = A$
Similarly, $e(e_1(A)) = A = e_1(e(A))$

Note: e_1 is the inverse of e

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) =$$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

$$e_1(e(A)) =$$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Similarly, $e(e_1(A)) = A = e_1(e(A))$

Inverse of the Type 3 elementary row operation

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix}$$

Let $e : R_1 \longleftrightarrow R_2$.

$$e(A) = \begin{bmatrix} A_{21} & A_{22} & A_{23} \\ A_{11} & A_{12} & A_{13} \end{bmatrix}$$

We define $e_1 : R_1 \longleftrightarrow R_2$.

$$e_1(e(A)) = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \end{bmatrix} = A$$

Similarly, $e(e_1(A)) = A = e_1(e(A))$

Note: e_1 is the inverse of e

Theorem 2

To each elementary row operation e

Theorem 2

To each elementary row operation e there corresponds an elementary operation e_1 , of the same type as e ,

Theorem 2

To each elementary row operation e there corresponds an elementary operation e_1 , of the same type as e , such that $e(e_1(A)) = A = e_1(e(A))$.

Theorem 2

To each elementary row operation e there corresponds an elementary operation e_1 , of the same type as e , such that $e(e_1(A)) = A = e_1(e(A))$. In other words, inverse operation of an elementary operation exists and is of an elementary operation of the same type.

Theorem 2

To each elementary row operation e there corresponds an elementary operation e_1 , of the same type as e , such that $e(e_1(A)) = A = e_1(e(A))$. In other words, inverse operation of an elementary operation exists and is of an elementary operation of the same type.

Proof :

<i>Type 1</i>	<i>Inverse of the Type 1</i>
$e : R_i \longleftarrow cR_i, \ c \neq 0$	$e_1 : R_i \longleftarrow \frac{1}{c}R_i$

Proof of Theorem 2 contd.

<i>Type 2</i>	<i>Inverse of the Type 2</i>
$e : R_i \longleftarrow R_i + cR_j$	$e_1 : R_i \longleftarrow R_i - cR_j$

Proof of Theorem 2 contd.

<i>Type 2</i>	<i>Inverse of the Type 2</i>
$e : R_i \longleftarrow R_i + cR_j$	$e_1 : R_i \longleftarrow R_i - cR_j$

<i>Type 3</i>	<i>Inverse of the Type 3</i>
$e : R_i \longleftrightarrow R_j$	$e_1 : R_i \longleftrightarrow R_j$

Proof of Theorem 2 contd.

<i>Type 2</i>	<i>Inverse of the Type 2</i>
$e : R_i \longleftarrow R_i + cR_j$	$e_1 : R_i \longleftarrow R_i - cR_j$

<i>Type 3</i>	<i>Inverse of the Type 3</i>
$e : R_i \longleftrightarrow R_j$	$e_1 : R_i \longleftrightarrow R_j$

Note that for an $m \times n$ matrix A , $e(e_1(A)) = A = e_1(e(A))$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix}$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)}$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix}$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xrightarrow{(e_2)}$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xrightarrow{(e_2)} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

$$e_2(e_1(A)) = B$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xrightarrow{(e_2)} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

$$e_2(e_1(A)) = B$$

$$e_1^{-1} : R_2 \longleftarrow \frac{1}{2}R_2 \text{ and } e_2^{-1} : R_2 \longleftarrow R_2 + R_1$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xrightarrow{(e_2)} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

$$e_2(e_1(A)) = B$$

$$e_1^{-1} : R_2 \longleftarrow \frac{1}{2}R_2 \text{ and } e_2^{-1} : R_2 \longleftarrow R_2 + R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xleftarrow{(e_1^{-1})} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xleftarrow{(e_2^{-1})} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

Note

$$e_1 : R_2 \longleftarrow 2R_2 \text{ and } e_2 : R_2 \longleftarrow R_2 - R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xrightarrow{(e_1)} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xrightarrow{(e_2)} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

$$e_2(e_1(A)) = B$$

$$e_1^{-1} : R_2 \longleftarrow \frac{1}{2}R_2 \text{ and } e_2^{-1} : R_2 \longleftarrow R_2 + R_1$$

$$A = \begin{bmatrix} 4 & -1 & 5 \\ 2 & 1 & 7 \end{bmatrix} \xleftarrow{(e_1^{-1})} \begin{bmatrix} 4 & -1 & 5 \\ 4 & 2 & 14 \end{bmatrix} \xleftarrow{(e_2^{-1})} \begin{bmatrix} 4 & -1 & 5 \\ 0 & 3 & 9 \end{bmatrix} = B$$

$$e_1^{-1}(e_2^{-1}(B)) = A$$

A and B are called row-equivalent matrices.

Row-equivalent matrices

Definition : If A and B are $m \times n$ matrices over the field F , we say B is row-equivalent to A if B can be obtained from A by a **finite** sequence of elementary row operations.

Row-equivalent matrices

Definition : If A and B are $m \times n$ matrices over the field F , we say B is row-equivalent to A if B can be obtained from A by a **finite** sequence of elementary row operations.

Qn. Let $M_{m \times n}$ be the set of all $m \times n$ matrices defined on a field F .

Row-equivalent matrices

Definition : If A and B are $m \times n$ matrices over the field F , we say B is row-equivalent to A if B can be obtained from A by a **finite** sequence of elementary row operations.

Qn. Let $M_{m \times n}$ be the set of all $m \times n$ matrices defined on a field F . For $A, B \in M_{m \times n}$, we say $A \sim B$ if B is row - equivalent to A .

Row-equivalent matrices

Definition : If A and B are $m \times n$ matrices over the field F , we say B is row-equivalent to A if B can be obtained from A by **a finite** sequence of elementary row operations.

Qn. Let $M_{m \times n}$ be the set of all $m \times n$ matrices defined on a field F . For $A, B \in M_{m \times n}$, we say $A \sim B$ if B is row - equivalent to A . Show that \sim is an equivalence relation defined on $M_{m \times n}$

Theorem 3

If A and B are row-equivalent $m \times n$ matrices,

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

$$A = A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow A_k = B$$

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

$$A = A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow A_k = B$$

Note: If

(1) $A_0X = 0$ and $A_1X = 0$ have same solutions,

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

$$A = A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow A_k = B$$

Note: If

- (1) $A_0X = 0$ and $A_1X = 0$ have same solutions,
- (2) $A_1X = 0$ and $A_2X = 0$ have same solutions,

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

$$A = A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow A_k = B$$

Note: If

(1) $A_0X = 0$ and $A_1X = 0$ have same solutions,

(2) $A_1X = 0$ and $A_2X = 0$ have same solutions,

(j) $\dots, \dots,$

(k) $A_{k-1}X = 0$ and $A_kX = 0$ have same solutions,

then

Theorem 3

If A and B are row-equivalent $m \times n$ matrices, the homogeneous systems of linear equations $AX = 0$ and $BX = 0$ have exactly same solutions.

Proof: Suppose that we pass A to B by a finite sequence of elementary row operations :

$$A = A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow A_k = B$$

Note: If

(1) $A_0X = 0$ and $A_1X = 0$ have same solutions,

(2) $A_1X = 0$ and $A_2X = 0$ have same solutions,

(j) $\dots, \dots,$

(k) $A_{k-1}X = 0$ and $A_kX = 0$ have same solutions,

then $AX = 0$ and $BX = 0$ have same solutions.

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Suppose that B is obtained from A by a single elementary row operation, say e (i.e., $e(A) = B$).

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Suppose that B is obtained from A by a single elementary row operation, say e (i.e., $e(A) = B$). No matter which of the types the operation is : (1) , (2) or (3), each equation in the system $BX = 0$ is a linear combination of the equations in $AX = 0$.

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Suppose that B is obtained from A by a single elementary row operation, say e (i.e., $e(A) = B$). No matter which of the types the operation is : (1) , (2) or (3), each equation in the system $BX = 0$ is a linear combination of the equations in $AX = 0$. Since e^{-1} is an elementary row operation (i.e., $e^{-1}(B) = A$),

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Suppose that B is obtained from A by a single elementary row operation, say e (i.e., $e(A) = B$). No matter which of the types the operation is : (1) , (2) or (3), each equation in the system $BX = 0$ is a linear combination of the equations in $AX = 0$. Since e^{-1} is an elementary row operation (i.e., $e^{-1}(B) = A$), each equation in the system $AX = 0$ will also be a linear combination of equations in $BX = 0$.

Proof of Theorem 3 contd.

It is enough to prove that $A_j X = 0$ and $A_{j+1} X = 0$ have exactly the same solutions (that is one elementary row operation doesn't disturb the set of solutions).

Suppose that B is obtained from A by a single elementary row operation, say e (i.e., $e(A) = B$). No matter which of the types the operation is : (1) , (2) or (3), each equation in the system $BX = 0$ is a linear combination of the equations in $AX = 0$. Since e^{-1} is an elementary row operation (i.e., $e^{-1}(B) = A$), each equation in the system $AX = 0$ will also be a linear combination of equations in $BX = 0$. Hence these two systems are equivalent, and by Theorem 1, they have the same solutions.

Problem 1

Show that the following systems are equivalent.

$AX = 0$	$BX = 0$
$2x_1 - x_2 + 3x_3 + 2x_4 = 0$	$x_3 - \frac{11}{3}x_4 = 0$
$x_1 + 4x_2 - x_4 = 0$	$x_1 + \frac{17}{3}x_4 = 0$
$2x_1 + 6x_2 - x_3 + 5x_4 = 0$	$x_2 - \frac{5}{3}x_4 = 0$

Problem 1

Show that the following systems are equivalent.

$AX = 0$	$BX = 0$
$2x_1 - x_2 + 3x_3 + 2x_4 = 0$	$x_3 - \frac{11}{3}x_4 = 0$
$x_1 + 4x_2 - x_4 = 0$	$x_1 + \frac{17}{3}x_4 = 0$
$2x_1 + 6x_2 - x_3 + 5x_4 = 0$	$x_2 - \frac{5}{3}x_4 = 0$

Solution at page number 8(Hoffman and Kunz)

It's an assignment.

Problem 1

Show that the following systems are equivalent.

$AX = 0$	$BX = 0$
$2x_1 - x_2 + 3x_3 + 2x_4 = 0$	$x_3 - \frac{11}{3}x_4 = 0$
$x_1 + 4x_2 - x_4 = 0$	$x_1 + \frac{17}{3}x_4 = 0$
$2x_1 + 6x_2 - x_3 + 5x_4 = 0$	$x_2 - \frac{5}{3}x_4 = 0$

Solution at page number 8(Hoffman and Kunz)

It's an assignment.

Note that solving the second system is easy !

Note

Let us consider the matrix B from the previous problem.

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \end{bmatrix}$$

Note

Let us consider the matrix B from the previous problem.

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \end{bmatrix}$$

- Note that the first non-zero entry of each non-zero row of B is 1.

Note

Let us consider the matrix B from the previous problem.

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \end{bmatrix}$$

- Note that the first non-zero entry of each non-zero row of B is 1.
- Note that each column of B which contains the leading non-zero entry of some row has all its other entries 0.

Note

Let us consider the matrix B from the previous problem.

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \end{bmatrix}$$

- Note that the first non-zero entry of each non-zero row of B is 1.
- Note that each column of B which contains the leading non-zero entry of some row has all its other entries 0.

$$B = \begin{bmatrix} 0 & \textcolor{red}{0} & 1 & -\frac{11}{3} \\ 1 & \textcolor{red}{0} & 0 & \frac{17}{3} \\ 0 & \textcolor{red}{1} & 0 & -\frac{5}{3} \end{bmatrix}$$

Row-reduced matrix

An $m \times n$ matrix R is called **row-reduced** if :

Row-reduced matrix

An $m \times n$ matrix R is called **row-reduced** if :

- (a) the first non-zero entry of each non-zero row of R is 1;

Row-reduced matrix

An $m \times n$ matrix R is called row-reduced if :

- (a) the first non-zero entry of each non-zero row of R is 1;
- (b) each column of R which contains the leading non-zero entry of some row has all its other entries 0.

Row-reduced matrix

An $m \times n$ matrix R is called **row-reduced** if :

- (a) the first non-zero entry of each non-zero row of R is 1;
- (b) each column of R which contains the leading non-zero entry of some row has all its other entries 0.

Examples : (i) Identity matrix and (ii) the matrix B (previous problem)

Row-reduced matrix

An $m \times n$ matrix R is called **row-reduced** if :

- (a) the first non-zero entry of each non-zero row of R is 1;
- (b) each column of R which contains the leading non-zero entry of some row has all its other entries 0.

Examples : (i) Identity matrix and (ii) the matrix B (previous problem)

Examples of non row-reduced matrices

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix},$$

Row-reduced matrix

An $m \times n$ matrix R is called **row-reduced** if :

- (a) the first non-zero entry of each non-zero row of R is 1;
- (b) each column of R which contains the leading non-zero entry of some row has all its other entries 0.

Examples : (i) Identity matrix and (ii) the matrix B (previous problem)

Examples of non row-reduced matrices

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 2 & 1 \\ 1 & 0 & -3 \\ 0 & 0 & 0 \end{bmatrix}$$

Problem 1

Find all solutions of the following system of equations by row-reducing the coefficient matrix.

$$\frac{1}{3}x_1 + 2x_2 - 6x_3 = 0$$

$$-4x_1 \quad \quad + 5x_3 = 0$$

$$-3x_1 + 6x_2 - 13x_3 = 0$$

$$-\frac{7}{3}x_1 + 2x_2 - \frac{8}{3}x_3 = 0$$

Problem 1

Find all solutions of the following system of equations by row-reducing the coefficient matrix.

$$\frac{1}{3}x_1 + 2x_2 - 6x_3 = 0$$

$$-4x_1 + 5x_3 = 0$$

$$-3x_1 + 6x_2 - 13x_3 = 0$$

$$-\frac{7}{3}x_1 + 2x_2 - \frac{8}{3}x_3 = 0$$

Solution: The coefficient matrix of the system is

$$\begin{bmatrix} \frac{1}{3} & 2 & -6 \\ -4 & 0 & 5 \\ -3 & 6 & -13 \\ -\frac{7}{3} & 2 & -\frac{8}{3} \end{bmatrix}$$

Problem 1 contd.

$$R_1 \leftarrow 3R_1, R_4 \leftarrow 3R_4$$

Problem 1 contd.

$$R_1 \leftarrow 3R_1, R_4 \leftarrow 3R_4$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ -4 & 0 & 5 \\ -3 & 6 & -13 \\ -7 & 6 & -8 \end{bmatrix}$$

Problem 1 contd.

$$R_1 \longleftarrow 3R_1, R_4 \longleftarrow 3R_4$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ -4 & 0 & 5 \\ -3 & 6 & -13 \\ -7 & 6 & -8 \end{bmatrix}$$

$$R_2 \longleftarrow R_2 + 4R_1, R_3 \longleftarrow R_3 + 3R_1, R_4 \longleftarrow R_4 + 7R_1$$

Problem 1 contd.

$$R_1 \longleftarrow 3R_1, R_4 \longleftarrow 3R_4$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ -4 & 0 & 5 \\ -3 & 6 & -13 \\ -7 & 6 & -8 \end{bmatrix}$$

$$R_2 \longleftarrow R_2 + 4R_1, R_3 \longleftarrow R_3 + 3R_1, R_4 \longleftarrow R_4 + 7R_1$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 24 & -67 \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

Problem 1 contd.

$$R_2 \leftarrow \frac{1}{24} R_2$$

Problem 1 contd.

$$R_2 \leftarrow \frac{1}{24} R_2$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 1 & -\frac{67}{24} \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

Problem 1 contd.

$$R_2 \longleftarrow \frac{1}{24} R_2$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 1 & -\frac{67}{24} \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

$$R_1 \longleftarrow R_1 - 6R_2, R_3 \longleftarrow R_3 - 24R_2, R_4 \longleftarrow R_4 - 48R_2$$

Problem 1 contd.

$$R_2 \leftarrow \frac{1}{24} R_2$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 1 & -\frac{67}{24} \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

$$R_1 \leftarrow R_1 - 6R_2, R_3 \leftarrow R_3 - 24R_2, R_4 \leftarrow R_4 - 48R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -\frac{5}{4} \\ 0 & 1 & -\frac{67}{24} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Problem 1 contd.

$$R_2 \leftarrow \frac{1}{24} R_2$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 1 & -\frac{67}{24} \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

$$R_1 \leftarrow R_1 - 6R_2, R_3 \leftarrow R_3 - 24R_2, R_4 \leftarrow R_4 - 48R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -\frac{5}{4} \\ 0 & 1 & -\frac{67}{24} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (\text{a row - reduced matrix})$$

Problem 1 contd.

$$R_2 \leftarrow \frac{1}{24} R_2$$

$$\sim \begin{bmatrix} 1 & 6 & -18 \\ 0 & 1 & -\frac{67}{24} \\ 0 & 24 & -67 \\ 0 & 48 & -134 \end{bmatrix}$$

$$R_1 \leftarrow R_1 - 6R_2, R_3 \leftarrow R_3 - 24R_2, R_4 \leftarrow R_4 - 48R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -\frac{5}{4} \\ 0 & 1 & -\frac{67}{24} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (a \text{ row-reduced matrix})$$

Thus

$$x_1 + 0x_2 - \frac{5}{4}x_3 = 0$$

$$0x_1 + x_2 - \frac{67}{24}x_3 = 0$$

Problem 1 contd.

Let $x_3 = a$.

Problem 1 contd.

Let $x_3 = a. \implies x_1 = \frac{5}{4}a, x_2 = \frac{67}{24}a$

Problem 1 contd.

Let $x_3 = a. \implies x_1 = \frac{5}{4}a, x_2 = \frac{67}{24}a$

Solution set, $S = \{(\frac{5}{4}a, \frac{67}{24}a, a) : a \in \mathbb{R}\}$

Problem 1 contd.

Let $x_3 = a. \implies x_1 = \frac{5}{4}a, x_2 = \frac{67}{24}a$

Solution set, $S = \{(\frac{5}{4}a, \frac{67}{24}a, a) : a \in \mathbb{R}\}$

Note that

- (i) x_3 is a called free variable and**
- (ii) x_1, x_2 are called pivot variables.**

Theorem 4

Every $m \times n$ matrix over the field F is row-equivalent to a row-reduced matrix.

Theorem 4

Every $m \times n$ matrix over the field F is row-equivalent to a row-reduced matrix.

Proof : (Assignment)

Problem 2

Find all solutions of the systems of linear equations $AX = 2X$ and $AX = 3X$ where

$$A = \begin{bmatrix} 6 & -4 & 0 \\ 4 & -2 & 0 \\ -1 & 0 & 3 \end{bmatrix}$$

Problem 2

Find all solutions of the systems of linear equations $AX = 2X$ and $AX = 3X$ where

$$A = \begin{bmatrix} 6 & -4 & 0 \\ 4 & -2 & 0 \\ -1 & 0 & 3 \end{bmatrix}$$

Solution : (i) The system $AX = 2X$ is

$$\begin{bmatrix} 6 & -4 & 0 \\ 4 & -2 & 0 \\ -1 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 2 \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

Problem 2

Find all solutions of the systems of linear equations $AX = 2X$ and $AX = 3X$ where

$$A = \begin{bmatrix} 6 & -4 & 0 \\ 4 & -2 & 0 \\ -1 & 0 & 3 \end{bmatrix}$$

Solution : (i) The system $AX = 2X$ is

$$\begin{bmatrix} 6 & -4 & 0 \\ 4 & -2 & 0 \\ -1 & 0 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = 2 \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\Rightarrow \left. \begin{array}{l} 6x - 4y = 2x \\ 4x - 2y = 2y \\ -x + 3z = 2z \end{array} \right\}$$

Problem 2 contd.

$$\Rightarrow \left. \begin{array}{l} 4x - 4y = 0 \\ 4x - 4y = 0 \\ -x + z = 0 \end{array} \right\}$$

Problem 2 contd.

$$\Rightarrow \left. \begin{array}{l} 4x - 4y = 0 \\ 4x - 4y = 0 \\ -x + z = 0 \end{array} \right\}$$

The coefficient matrix is

$$\begin{bmatrix} 4 & -4 & 0 \\ 4 & -4 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

Problem 2 contd.

$$\Rightarrow \left. \begin{array}{l} 4x - 4y = 0 \\ 4x - 4y = 0 \\ -x + z = 0 \end{array} \right\}$$

The coefficient matrix is

$$\begin{bmatrix} 4 & -4 & 0 \\ 4 & -4 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

Let us find a row-reduced matrix which is row-equivalent to the above matrix.

Problem 2 contd.

$$\Rightarrow \left. \begin{array}{l} 4x - 4y = 0 \\ 4x - 4y = 0 \\ -x + z = 0 \end{array} \right\}$$

The coefficient matrix is

$$\begin{bmatrix} 4 & -4 & 0 \\ 4 & -4 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

Let us find a row-reduced matrix which is row-equivalent to the above matrix. $R_3 \leftarrow (-1)R_3$, $R_3 \longleftrightarrow R_1$

Problem 2 contd.

$$\Rightarrow \left. \begin{array}{l} 4x - 4y = 0 \\ 4x - 4y = 0 \\ -x + z = 0 \end{array} \right\}$$

The coefficient matrix is

$$\begin{bmatrix} 4 & -4 & 0 \\ 4 & -4 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

Let us find a row-reduced matrix which is row-equivalent to the above matrix. $R_3 \leftarrow (-1)R_3$, $R_3 \longleftrightarrow R_1$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 4 & -4 & 0 \\ 4 & -4 & 0 \end{bmatrix}$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & 4 \\ 0 & -4 & 4 \end{bmatrix}$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & 4 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_2 \longleftarrow -\frac{1}{4}R_2$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & 4 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_2 \longleftarrow -\frac{1}{4}R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & -4 & 4 \end{bmatrix}$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & 4 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_2 \longleftarrow -\frac{1}{4}R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_3 \longleftarrow R_3 + 4R_2$$

Problem 2 contd.

$$R_2 \longleftarrow R_2 - 4R_1 \text{ and } R_3 \longleftarrow R_3 - 4R_1$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & 4 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_2 \longleftarrow -\frac{1}{4}R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & -4 & 4 \end{bmatrix}$$

$$R_3 \longleftarrow R_3 + 4R_2$$

$$\sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 2X\} = \{(a, a, a) : a \in \mathbb{R}\}$$

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 2X\} = \{(a, a, a) : a \in \mathbb{R}\}$$

Note : For all $X \in S$, $AX = 2X$. What is its geometrical interpretation?

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 2X\} = \{(a, a, a) : a \in \mathbb{R}\}$$

Note : For all $X \in S$, $AX = 2X$. **What is its geometrical interpretation?** (Hint : eigen vector)

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 2X\} = \{(a, a, a) : a \in \mathbb{R}\}$$

Note : For all $X \in S$, $AX = 2X$. **What is its geometrical interpretation?** (Hint : eigen vector)

(ii) Find all solutions of $AX = 3X$

Problem 2 contd.

The equivalent system is

$$\left. \begin{array}{l} x - z = 0 \\ y - z = 0 \end{array} \right\}$$

Let $z = a$ (Note that z is a free variable). Thus $x = a = y$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 2X\} = \{(a, a, a) : a \in \mathbb{R}\}$$

Note : For all $X \in S$, $AX = 2X$. **What is its geometrical interpretation?** (Hint : eigen vector)

(ii) Find all solutions of $AX = 3X$

The solution set is

$$S = \{X \in \mathbb{R}^3 : AX = 3X\} = \{(0, 0, a) : a \in \mathbb{R}\}$$

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.
3. The k_i denotes the column which contains leading one(if exists) of R_i (row i).

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.
3. The k_i denotes the column which contains leading one(if exists) of R_i (row i).
 $k_1 = 2,$

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.
3. The k_i denotes the column which contains leading one(if exists) of R_i (row i).
 $k_1 = 2, k_2 = 4,$

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.
3. The k_i denotes the column which contains leading one(if exists) of R_i (row i).
 $k_1 = 2$, $k_2 = 4$, and $k_3 = 5$.

Note

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

1. A is a row-reduced matrix.
2. All non-zero rows are above zero rows.
3. The k_i denotes the column which contains leading one(if exists) of R_i (row i).
 $k_1 = 2$, $k_2 = 4$, and $k_3 = 5$.

Note that $k_1 < k_2 < k_3$

Row-reduced echelon matrix

$$A = \begin{bmatrix} 0 & 1 & -3 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

blue zeros forms a staircase (echelon) from right to left

Row-reduced echelon matrix

An $m \times n$ matrix R is called a row-reduced echelon matrix if:

Row-reduced echelon matrix

An $m \times n$ matrix R is called a row-reduced echelon matrix if:

(a) R is row-reduced ;

Row-reduced echelon matrix

An $m \times n$ matrix R is called a row-reduced echelon matrix if:

- (a) R is row-reduced ;
- (b) every row of R which has all its entries 0 occurs below every row which has a non-zero entry;

Row-reduced echelon matrix

An $m \times n$ matrix R is called a row-reduced echelon matrix if:

- (a) R is row-reduced ;
- (b) every row of R which has all its entries 0 occurs below every row which has a non-zero entry;
- (c) if rows $1, 2, \dots, r$ are the non-zero rows of R , and if the leading non-zero entry of row i occurs in column k_i , $i = 1, 2, \dots, r$, then $k_1 < k_2 < \dots < k_r$.

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

$k_1 = 3, k_2 = 1, k_3 = 2$ which violates the condition (c)

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

$k_1 = 3, k_2 = 1, k_3 = 2$ which violates the condition (c)

Could you find a row-reduced echelon matrix C which is row-equivalent to B ?

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

$k_1 = 3, k_2 = 1, k_3 = 2$ which violates the condition (c)

Could you find a row-reduced echelon matrix C which is row-equivalent to B ? ($R_1 \longleftrightarrow R_2, R_2 \longleftrightarrow R_3$)

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

$k_1 = 3, k_2 = 1, k_3 = 2$ which violates the condition (c)

Could you find a row-reduced echelon matrix C which is row-equivalent to B ? ($R_1 \longleftrightarrow R_2, R_2 \longleftrightarrow R_3$)

$$C = \begin{bmatrix} 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 1 & -\frac{11}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

B is a row-reduced matrix, but not a row-reduced echelon matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & -\frac{11}{3} \\ 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Why?

$k_1 = 3, k_2 = 1, k_3 = 2$ which violates the condition (c)

Could you find a row-reduced echelon matrix C which is row-equivalent to B ? ($R_1 \longleftrightarrow R_2, R_2 \longleftrightarrow R_3$)

$$C = \begin{bmatrix} 1 & 0 & 0 & \frac{17}{3} \\ 0 & 1 & 0 & -\frac{5}{3} \\ 0 & 0 & 1 & -\frac{11}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad (\text{Is it unique?})$$

Problem

1. Prove or disprove that there is only one 3×3 row-reduced echelon matrix with 3 non-zero rows.

Problem

1. Prove or disprove that there is only one 3×3 row-reduced echelon matrix with 3 non-zero rows.
2. Find all 2×2 row-reduced echelon matrices.

Theorem 5

Every $m \times n$ matrix A is row-equivalent to a row-reduced echelon matrix.

Theorem 5

Every $m \times n$ matrix A is row-equivalent to a row-reduced echelon matrix.

Assignment

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

$$k_1 = 2, k_2 = 4$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

$$k_1 = 2, k_2 = 4 \implies \text{Pivot variables} = \{x_{k_1}, x_{k_2}\} = \{x_2, x_4\}$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

$$k_1 = 2, k_2 = 4 \implies \text{Pivot variables} = \{x_{k_1}, x_{k_2}\} = \{x_2, x_4\}$$

$$\text{No. of free variables} = n - r = 5 - 2 = 3,$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

$$k_1 = 2, k_2 = 4 \implies \text{Pivot variables} = \{x_{k_1}, x_{k_2}\} = \{x_2, x_4\}$$

$$\text{No. of free variables} = n - r = 5 - 2 = 3,$$

$$\text{Free variables} = \{u_1, u_2, u_3\} = \{x_1, x_3, x_5\}$$

Note 1:

Consider a row-reduced echelon matrix R and the system

$$RX = 0$$

$$R = \begin{bmatrix} 0 & 1 & -3 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\}$$

No. of non-zero rows of R , $r = 2$, No. of variables, $n = 5$

$$k_1 = 2, k_2 = 4 \implies \text{Pivot variables} = \{x_{k_1}, x_{k_2}\} = \{x_2, x_4\}$$

$$\text{No. of free variables} = n - r = 5 - 2 = 3,$$

$$\text{Free variables} = \{u_1, u_2, u_3\} = \{x_1, x_3, x_5\}$$

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \implies \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j}u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j}u_j = 0 \end{array} \right\} \text{ (general expression)}$$

Note 1 contd.

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0 \end{array} \right\} \text{general expression)}$$

Note 1 contd.

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0 \end{array} \right\} \text{general expression)}$$

Set the free variables as :

$$u_1 = x_1 = a, \quad u_2 = x_3 = b, \quad u_3 = x_5 = c$$

Note 1 contd.

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j}u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j}u_j = 0 \end{array} \right\} \text{general expression)}$$

Set the free variables as :

$$u_1 = x_1 = a, \quad u_2 = x_3 = b, \quad u_3 = x_5 = c$$

$$\Rightarrow x_2 = 3b - \frac{1}{2}c, \quad x_4 = -2c$$

Note 1 contd.

$$\left. \begin{array}{l} x_2 - 3x_3 + \frac{1}{2}x_5 = 0 \\ x_4 + 2x_5 = 0 \end{array} \right\} \Rightarrow \left. \begin{array}{l} x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j = 0 \\ x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0 \end{array} \right\} \text{general expression)}$$

Set the free variables as :

$$u_1 = x_1 = a, \quad u_2 = x_3 = b, \quad u_3 = x_5 = c$$

$$\Rightarrow x_2 = 3b - \frac{1}{2}c, \quad x_4 = -2c$$

$$\text{Solution set } S = \left\{ \left(a, 3b - \frac{1}{2}c, b, -2c, c \right) : a, b, c \in \mathbb{R} \right\}$$

Note 1 contd. (a visit to future !)

Solution set $S = \{(a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R}\}$

Note 1 contd. (a visit to future !)

Solution set $S = \{(a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R}\}$

$$S = \left\{ a(1, 0, 0, 0, 0) + b(0, 3, 1, 0, 0) + c(0, -\frac{1}{2}, 0, -2, 1) : a, b, c \in \mathbb{R} \right\}$$

Note 1 contd. (a visit to future !)

Solution set $S = \{(a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R}\}$

$$S = \left\{ a(1, 0, 0, 0, 0) + b(0, 3, 1, 0, 0) + c(0, -\frac{1}{2}, 0, -2, 1) : a, b, c \in \mathbb{R} \right\}$$

$$= \text{Span of } \left\{ (1, 0, 0, 0, 0), (0, 3, 1, 0, 0), (0, -\frac{1}{2}, 0, -2, 1) \right\}$$

Note 1 contd. (a visit to future !)

Solution set $S = \{(a, 3b - \frac{1}{2}c, b, -2c, c) : a, b, c \in \mathbb{R}\}$

$$S = \left\{ a(1, 0, 0, 0, 0) + b(0, 3, 1, 0, 0) + c(0, -\frac{1}{2}, 0, -2, 1) : a, b, c \in \mathbb{R} \right\}$$

$$= \text{Span of } \left\{ (1, 0, 0, 0, 0), (0, 3, 1, 0, 0), (0, -\frac{1}{2}, 0, -2, 1) \right\}$$

Dimension of $S = \dim S = 3 = n - r$ (Information for future)

Note 2

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows.

Note 2

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows. Let rows $1, 2, \dots, r$ be the non-zero rows of R , and suppose that the leading non-zero entry of row i occurs in column k_i .

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows. Let rows $1, 2, \dots, r$ be the non-zero rows of R , and suppose that the leading non-zero entry of row i occurs in column k_i . The system $RX = 0$ has r (non-trivial) equations.

Note 2

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows. Let rows $1, 2, \dots, r$ be the non-zero rows of R , and suppose that the leading non-zero entry of row i occurs in column k_i . The system $RX = 0$ has r (non-trivial) equations. Also the unknown x_{k_i} will occur (with non-zero coefficient) only in the i^{th} equation.

Note 2

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows. Let rows $1, 2, \dots, r$ be the non-zero rows of R , and suppose that the leading non-zero entry of row i occurs in column k_i . The system $RX = 0$ has r (non-trivial) equations. Also the unknown x_{k_i} will occur (with non-zero coefficient) only in the i^{th} equation. Let u_1, u_2, \dots, u_{n-r} denote the $(n - r)$ unknowns which are different from $x_{k_1}, x_{k_2}, \dots, x_{k_r}$.

Note 2

Consider an $m \times n$ row-reduced echelon matrix R with r non-zero rows. Let rows $1, 2, \dots, r$ be the non-zero rows of R , and suppose that the leading non-zero entry of row i occurs in column k_i . The system $RX = 0$ has r (non-trivial) equations. Also the unknown x_{k_i} will occur (with non-zero coefficient) only in the i^{th} equation. Let u_1, u_2, \dots, u_{n-r} denote the $(n - r)$ unknowns which are different from $x_{k_1}, x_{k_2}, \dots, x_{k_r}$. Then r non-trivial equations of $RX = 0$ are of the form

Note 2 contd.

$$x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j = 0$$

$$x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0$$

.....

$$x_{k_r} + \sum_{j=1}^{n-r} C_{rj} u_j = 0$$

Note 2 contd.

$$\begin{aligned}x_{k_1} + \sum_{j=1}^{n-r} C_{1j} u_j &= 0 \\x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j &= 0 \\&\dots\dots\dots \\x_{k_r} + \sum_{j=1}^{n-r} C_{rj} u_j &= 0\end{aligned}$$

All the solutions of the system of equations $RX = 0$ are obtained by assigning any value whatsoever to u_1, u_2, \dots, u_{n-r} , and then computing the corresponding values of $x_{k_1}, x_{k_2}, \dots, x_{k_r}$.

Remarks (Note 2 contd.)

- (i) **If $n - r > 0$, then the system $RX = 0$ has at least one free variable and thus it has a non-trivial solution.**

Remarks (Note 2 contd.)

- (i) **If $n - r > 0$, then the system $RX = 0$ has at least one free variable and thus it has a non-trivial solution.**
- (ii) **If $n - r = 0$, then the system $RX = 0$ has no free variable and thus it has only trivial solution**

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Proof: Let R be a row-reduced echelon matrix which is row-equivalent to A .

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Proof: Let R be a row-reduced echelon matrix which is row-equivalent to A . Then the systems $AX = 0$ and $RX = 0$ have same solutions by Theorem 3.

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Proof: Let R be a row-reduced echelon matrix which is row-equivalent to A . Then the systems $AX = 0$ and $RX = 0$ have same solutions by Theorem 3. If r is the number of non-zero rows of R , then $r \leq m$.

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Proof: Let R be a row-reduced echelon matrix which is row-equivalent to A . Then the systems $AX = 0$ and $RX = 0$ have same solutions by Theorem 3. If r is the number of non-zero rows of R , then $r \leq m$. Since $m < n$, we have $r < n$. Thus the system $RX = 0$ has $n - r (\geq 1)$ free variables and it admits a non-trivial solution.

Theorem 6

If A is an $m \times n$ matrix and $m < n$, then the homogeneous system of linear equations $AX = 0$ has a non-trivial solution.

Proof: Let R be a row-reduced echelon matrix which is row-equivalent to A . Then the systems $AX = 0$ and $RX = 0$ have same solutions by Theorem 3. If r is the number of non-zero rows of R , then $r \leq m$. Since $m < n$, we have $r < n$. Thus the system $RX = 0$ has $n - r (\geq 1)$ free variables and it admits a non-trivial solution. Hence $AX = 0$ has a non-trivial solution.

If R is an $n \times n$ (square) row-reduced echelon matrix with n non-zero rows,

If R is an $n \times n$ (square) row-reduced echelon matrix with n non-zero rows, then $R = I$ (the identity matrix)

If R is an $n \times n$ (square) row-reduced echelon matrix with n non-zero rows, then $R = I$ (the identity matrix)

Hint : (i) Every row has a leading one and (ii)

$$k_1 = 1 < k_2 = 2 < \dots < k_n = n$$

Theorem 7

If A is an $n \times n$ (square) matrix,

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I .

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I . By Theorem 3, $AX = 0$ and $IX = 0$ have the same solution set. Thus the solution set of $AX = 0$ is

$$S = \{X : AX = 0\}$$

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I . By Theorem 3, $AX = 0$ and $IX = 0$ have the same solution set. Thus the solution set of $AX = 0$ is

$$S = \{X : AX = 0\} = \{X : IX = 0\}$$

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I . By Theorem 3, $AX = 0$ and $IX = 0$ have the same solution set. Thus the solution set of $AX = 0$ is

$$S = \{X : AX = 0\} = \{X : IX = 0\} = \{X : X = 0\}$$

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I . By Theorem 3, $AX = 0$ and $IX = 0$ have the same solution set. Thus the solution set of $AX = 0$ is

$$S = \{X : AX = 0\} = \{X : IX = 0\} = \{X : X = 0\} = \{0\}$$

Theorem 7

If A is an $n \times n$ (square) matrix, then A is row equivalent to the $n \times n$ identity matrix if and only if the system of equations $AX = 0$ has only the trivial solution.

Proof

Case 1. Suppose that A is row-equivalent to the $n \times n$ identity matrix I . By Theorem 3, $AX = 0$ and $IX = 0$ have the same solution set. Thus the solution set of $AX = 0$ is

$$S = \{X : AX = 0\} = \{X : IX = 0\} = \{X : X = 0\} = \{0\}$$

Hence the system $AX = 0$ has only the trivial solution.

Proof of Theorem 7 contd.

Case 2. Suppose that the system $AX = 0$ has only the trivial solution.

Proof of Theorem 7 contd.

Case 2. Suppose that the system $AX = 0$ has only the trivial solution. Prove that A is row-equivalent to the $n \times n$ identity matrix.

Proof of Theorem 7 contd.

Case 2. Suppose that the system $AX = 0$ has only the trivial solution. Prove that A is row-equivalent to the $n \times n$ identity matrix. Let R be an $n \times n$ row-reduced echelon matrix which is row-equivalent to A . By Theorem 3, the systems $AX = 0$ and $RX = 0$ have exactly the same solutions.

Proof of Theorem 7 contd.

Case 2. Suppose that the system $AX = 0$ has only the trivial solution. Prove that A is row-equivalent to the $n \times n$ identity matrix. Let R be an $n \times n$ row-reduced echelon matrix which is row-equivalent to A . By Theorem 3, the systems $AX = 0$ and $RX = 0$ have exactly the same solutions. Since **$AX = 0$ has only the trivial solution**, $RX = 0$ has only the trivial solution.

Proof of Theorem 7 contd.

Case 2. **Suppose that the system $AX = 0$ has only the trivial solution.** Prove that A is row-equivalent to the $n \times n$ identity matrix. Let R be an $n \times n$ row-reduced echelon matrix which is row-equivalent to A . By Theorem 3, the systems $AX = 0$ and $RX = 0$ have exactly the same solutions. Since **$AX = 0$ has only the trivial solution**, $RX = 0$ has only the trivial solution. Hence the system $RX = 0$ has no free variables.

Proof of Theorem 7 contd.

Case 2. **Suppose that the system $AX = 0$ has only the trivial solution.** Prove that A is row-equivalent to the $n \times n$ identity matrix. Let R be an $n \times n$ row-reduced echelon matrix which is row-equivalent to A . By Theorem 3, the systems $AX = 0$ and $RX = 0$ have exactly the same solutions. Since **$AX = 0$ has only the trivial solution**, $RX = 0$ has only the trivial solution. Hence the system $RX = 0$ has no free variables. Thus the number of free variables (of the system $RX = 0$), $n - r = 0$ where r is the number of non-zero rows of R .

Proof of Theorem 7 contd.

Case 2. **Suppose that the system $AX = 0$ has only the trivial solution.** Prove that A is row-equivalent to the $n \times n$ identity matrix. Let R be an $n \times n$ row-reduced echelon matrix which is row-equivalent to A . By Theorem 3, the systems $AX = 0$ and $RX = 0$ have exactly the same solutions. Since **$AX = 0$ has only the trivial solution**, $RX = 0$ has only the trivial solution. Hence the system $RX = 0$ has no free variables. Thus the number of free variables (of the system $RX = 0$), $n - r = 0$ where r is the number of non-zero rows of R . So R is an $n \times n$ row-reduced echelon matrix with $n(= r)$ non-zero rows and thus $k_1 = 1 < k_2 = 2 < \dots < k_n = n$. This proves that $R = I$, an identity matrix.
Hence A is row-equivalent to $R = I$.

Section 1.5 Matrix multiplication

Solve all exercise problems in section 1.4 (pages 15-16)