

Systems of Linear Equations

Nachiketa Mishra
IIITDM Kancheepuram, Chennai


Linear Equations

Equation of a line :

$$y = mx + c$$

$$ax + by = c$$

Example of a line




Equation of a plane :

$$z = ax + by + c$$

$$Ax + By + Cz = D$$

Example of a plane



Systems of linear equations

$$A_{11}x_1 + A_{12}x_2 + \dots + A_{1n}x_n = y_1$$

$$A_{21}x_1 + A_{22}x_2 + \dots + A_{2n}x_n = y_2$$

⋮ ⋮ ⋮

$$A_{m1}x_1 + A_{m2}x_2 + \dots + A_{mn}x_n = y_m$$

- m equations
- n variables (x_1, x_2, \dots, x_n)
- $A_{ij}, y_i \in F$ (F is a field)

Matrix form

$$\begin{bmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ \dots & \dots & \dots & \dots \\ A_{m1} & A_{m2} & \dots & A_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_n \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ \dots \\ y_m \end{bmatrix}$$

$$AX = Y$$

$$A = [A_{ij}]_{m \times n}, 1 \leq i \leq m, 1 \leq j \leq n$$

$$X = [x_j]_{n \times 1} \quad Y = [y_i]_{m \times 1}$$

The solution set of the linear system $AX = Y$ is

$$S = \{X \in R^n : AX = Y\}$$

Problem 1

Solve the following system of linear equations

$$4x - y = 5$$

$$2x + y = 7$$

Problem 1

Solve the following system of linear equations

$$4x - y = 5$$

$$2x + y = 7$$

Matrix form

$$\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

Problem 1

Solve the following system of linear equations

$$4x - y = 5$$


$$2x + y = 7$$

Matrix form

$$\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

Solve graphically (sketch it!)

Problem 1 (solution)



Problem 1

Solve the following system of linear equations

$$4x - y = 5$$

$$2x + y = 7$$

Matrix form

$$\begin{bmatrix} 4 & -1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

Solve graphically (sketch it!)

Solution $S = \{(2, 3)\}$

**Our objective is to design an efficient machinery
to solve $AX = B$**

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (Eq1)$$

$$2x + y = 7 \quad (Eq2)$$

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (Eq1)$$

$$2x + y = 7 \quad (Eq2)$$

Let us employ a high school technique

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (\text{Eq1})$$

$$2x + y = 7 \quad (\text{Eq2})$$

Let us employ a high school technique

Multiply (Eq2) by 2

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (\text{Eq1})$$

$$2x + y = 7 \quad (\text{Eq2})$$

Let us employ a high school technique

Multiply (Eq2) by 2

$$4x - y = 5 \quad (\text{Eq1})$$

$$4x + 2y = 14 \quad (\text{Eq2})$$

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (\text{Eq1})$$

$$2x + y = 7 \quad (\text{Eq2})$$

Let us employ a high school technique

Multiply (Eq2) by 2

$$4x - y = 5 \quad (\text{Eq1})$$

$$4x + 2y = 14 \quad (\text{Eq2})$$

$$(\text{Eq2}) - (\text{Eq1}) \implies$$

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (\text{Eq1})$$

$$2x + y = 7 \quad (\text{Eq2})$$

Let us employ a high school technique

Multiply (Eq2) by 2

$$4x - y = 5 \quad (\text{Eq1})$$

$$4x + 2y = 14 \quad (\text{Eq2})$$

(Eq2)-(Eq1) \Rightarrow

$$4x - y = 5 \quad (\text{Eq1})$$

$$0x + 3y = 9 \quad (\text{Eq2})$$

Solution of Problem 1 through elementary row operations

$$4x - y = 5 \quad (\text{Eq1})$$

$$2x + y = 7 \quad (\text{Eq2})$$

Let us employ a high school technique

Multiply (Eq2) by 2

$$4x - y = 5 \quad (\text{Eq1})$$

$$4x + 2y = 14 \quad (\text{Eq2})$$

$$(\text{Eq2}) - (\text{Eq1}) \implies$$

$$4x - y = 5 \quad (\text{Eq1})$$

$$0x + 3y = 9 \quad (\text{Eq2})$$

We have three equivalent systems say red, blue and green

contd.

Let us express **red** system in augmented matrix form

contd.

Let us express **red** system in augmented matrix form

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right]$$

contd.

Let us express **red** system in augmented matrix form

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right]$$

Multiply second row by 2 $(R_2 \leftarrow 2 \times R_2)$

contd.

Let us express **red** system in augmented matrix form

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right]$$

Multiply second row by 2 $(R_2 \leftarrow 2 \times R_2)$

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right]$$

contd.

Let us express **red** system in augmented matrix form

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right]$$

Multiply second row by 2 ($R_2 \leftarrow 2 \times R_2$)

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right]$$

Subtract first row from second row ($R_2 \leftarrow R_2 - R_1$)

contd.

Let us express **red** system in augmented matrix form

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right]$$

Multiply second row by 2 ($R_2 \leftarrow 2 \times R_2$)

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right]$$

Subtract first row from second row ($R_2 \leftarrow R_2 - R_1$)

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

Contd.

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

Contd.

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

$$R_2 \leftarrow \frac{1}{3}R_2 \quad \Rightarrow$$

Contd.

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

$$R_2 \leftarrow \frac{1}{3}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 1 & 3 \end{array} \right]$$

Contd.

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

$$R_2 \leftarrow \frac{1}{3}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

Contd.

$$\left[\begin{array}{cc|c} 4 & -1 & 5 \\ 2 & 1 & 7 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 4 & 2 & 14 \end{array} \right] \sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 3 & 9 \end{array} \right]$$

$$R_2 \leftarrow \frac{1}{3}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 4 & -1 & 5 \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

Contd.

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

Contd.

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow R_1 + \frac{1}{4}R_2 \quad \Rightarrow$$

Contd.

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow R_1 + \frac{1}{4}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 3 \end{array} \right]$$

Contd.

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow R_1 + \frac{1}{4}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 3 \end{array} \right]$$

Let us write it in the matrix form

Contd.

$$R_1 \leftarrow \frac{1}{4}R_1 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & -\frac{1}{4} & \frac{5}{4} \\ 0 & 1 & 3 \end{array} \right]$$

$$R_1 \leftarrow R_1 + \frac{1}{4}R_2 \quad \Rightarrow$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 3 \end{array} \right]$$

Let us write it in the matrix form

$$x + 0y = 2$$

$$0x + y = 3$$

Salient points

- Multiplying an equation by a non-zero scalar preserves the solution space ($R_i \leftarrow cR_i, \quad c \neq 0$)

Salient points

- Multiplying an equation by a non-zero scalar preserves the solution space ($R_i \leftarrow cR_i, \quad c \neq 0$)
- Replacing i^{th} equation by sum of i^{th} equation and constant multiple of j^{th} equation preserves the solution space.
 $(R_i \leftarrow R_i + cR_j)$

Salient points

- Multiplying an equation by a non-zero scalar preserves the solution space ($R_i \leftarrow cR_i, \quad c \neq 0$)
- Replacing i^{th} equation by sum of i^{th} equation and constant multiple of j^{th} equation preserves the solution space.
 $(R_i \leftarrow R_i + cR_j)$
- Interchanging two equations preserves the solution space.
 $(R_i \longleftrightarrow R_j)$

Problem 2

Solve the following system of linear equations.

$$3x - 2y = -6, \quad x + 2y = -10$$

Solve graphically !

Problem 2

Solve the following system of linear equations.

$$3x - 2y = -6, \quad x + 2y = -10$$

Solve graphically !

Augmented matrix is

Problem 2

Solve the following system of linear equations.

$$3x - 2y = -6, \quad x + 2y = -10$$

Solve graphically !

Augmented matrix is

$$\left[\begin{array}{cc|c} 3 & -2 & -6 \\ 1 & 2 & -10 \end{array} \right]$$

Problem 2

Solve the following system of linear equations.

$$3x - 2y = -6, \quad x + 2y = -10$$

Solve graphically !

Augmented matrix is

$$\left[\begin{array}{cc|c} 3 & -2 & -6 \\ 1 & 2 & -10 \end{array} \right]$$

Interchange first and second rows ($R_1 \longleftrightarrow R_2$)

Problem 2

Solve the following system of linear equations.

$$3x - 2y = -6, \quad x + 2y = -10$$

Solve graphically !

Augmented matrix is

$$\left[\begin{array}{cc|c} 3 & -2 & -6 \\ 1 & 2 & -10 \end{array} \right]$$

Interchange first and second rows ($R_1 \longleftrightarrow R_2$)

$$\sim \left[\begin{array}{cc|c} 1 & 2 & -10 \\ 3 & -2 & -6 \end{array} \right]$$

Problem 2 contd.

$$R_2 \leftarrow R_2 - 3R_1 \implies$$

Problem 2 contd.

$$R_2 \leftarrow R_2 - 3R_1 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & -10 \\ 0 & -8 & 24 \end{array} \right]$$

Problem 2 contd.

$$R_2 \leftarrow R_2 - 3R_1 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & -10 \\ 0 & -8 & 24 \end{array} \right]$$

$$R_2 \leftarrow -\frac{1}{8}R_2$$

Problem 2 contd.

$$R_2 \leftarrow R_2 - 3R_1 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & -10 \\ 0 & -8 & 24 \end{array} \right]$$

$$R_2 \leftarrow -\frac{1}{8}R_2$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & -10 \\ 0 & 1 & -3 \end{array} \right]$$

Problem 2 contd.

$$R_1 \leftarrow R_1 - 2R_2 \implies$$

Problem 2 contd.

$$R_1 \leftarrow R_1 - 2R_2 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & -4 \\ 0 & 1 & -3 \end{array} \right]$$

Problem 2 contd.

$$R_1 \leftarrow R_1 - 2R_2 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & -4 \\ 0 & 1 & -3 \end{array} \right]$$

$$x = -4, \quad y = -3$$

Problem 2 contd.

$$R_1 \leftarrow R_1 - 2R_2 \implies$$

$$\sim \left[\begin{array}{cc|c} 1 & 0 & -4 \\ 0 & 1 & -3 \end{array} \right]$$

$$x = -4, \quad y = -3$$

Solution $S = \{(-4, -3)\}$

Problem 3

Solve the system of linear equations

$$x + y = 2, \quad 2x + 2y = 5$$

Solve graphically!

Problem 3

Solve the system of linear equations

$$x + y = 2, \quad 2x + 2y = 5$$

Solve graphically!

Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 5 \end{array} \right]$$

Problem 3

Solve the system of linear equations

$$x + y = 2, \quad 2x + 2y = 5$$

Solve graphically!

Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 5 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

Problem 3

Solve the system of linear equations

$$x + y = 2, \quad 2x + 2y = 5$$

Solve graphically!

Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 5 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

$$\sim \left[\begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]$$

Problem 3

Solve the system of linear equations

$$x + y = 2, \quad 2x + 2y = 5$$

Solve graphically!

Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 5 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

$$\sim \left[\begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]$$

Second row is $0x + 0y = 1$. No solution i.e., $S = \{ \}$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Solve graphically!

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Solve graphically! Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 2 & 5 \\ 2 & 4 & 10 \end{array} \right]$$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Solve graphically! Augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 2 & 5 \\ 2 & 4 & 10 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Solve graphically! **Augmented matrix is**

$$\left[\begin{array}{cc|c} 1 & 2 & 5 \\ 2 & 4 & 10 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & 5 \\ 0 & 0 & 0 \end{array} \right]$$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$

Solve graphically! **Augmented matrix is**

$$\left[\begin{array}{cc|c} 1 & 2 & 5 \\ 2 & 4 & 10 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$

$$\sim \left[\begin{array}{cc|c} 1 & 2 & 5 \\ 0 & 0 & 0 \end{array} \right] \quad \Rightarrow \quad x + 2y = 5$$

Problem 4

Solve the system

$$x + 2y = 5, \quad 2x + 4y = 10$$


Solve graphically! **Augmented matrix is**

$$\left[\begin{array}{cc|c} 1 & 2 & 5 \\ 2 & 4 & 10 \end{array} \right] \quad (R_2 \leftarrow R_2 - 2R_1)$$


$$\sim \left[\begin{array}{cc|c} 1 & 2 & 5 \\ 0 & 0 & 0 \end{array} \right] \quad \Rightarrow \quad x + 2y = 5$$

The solution set: $S = \{(5 - 2c, c) : c \in \mathbb{R}\}$


two dimensional problem and possible solutions



(a)




(b)





(c)



3-dimensional problem with a unique solution



3-dimensional problem with no solutions



3-dimensional problem with infinite number of solutions



Linear combination of equations

$$A_{11}x_1 + A_{12}x_2 + A_{13}x_3 = y_1 \quad (1)$$

$$A_{21}x_1 + A_{22}x_2 + A_{23}x_3 = y_2 \quad (2)$$

Linear combination of equations

$$A_{11}x_1 + A_{12}x_2 + A_{13}x_3 = y_1 \quad (1)$$

$$A_{21}x_1 + A_{22}x_2 + A_{23}x_3 = y_2 \quad (2)$$

Consider $c_1(1) + c_2(2)$ (a linear combination) \Rightarrow

Linear combination of equations

$$A_{11}x_1 + A_{12}x_2 + A_{13}x_3 = y_1 \quad (1)$$

$$A_{21}x_1 + A_{22}x_2 + A_{23}x_3 = y_2 \quad (2)$$

Consider $c_1(1) + c_2(2)$ (a linear combination) \Rightarrow

$$\begin{aligned} & c_1(A_{11}x_1 + A_{12}x_2 + A_{13}x_3) + c_2(A_{21}x_1 + A_{22}x_2 + A_{23}x_3) \\ &= c_1y_1 + c_2y_2 \quad (3) \end{aligned}$$

Suppose that $x_1 = a, x_2 = b, x_3 = c$ is solution of (1) and (2)

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

$$= c_1 y_1$$

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

$$= c_1 y_1 + c_2 y_2$$

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

$$= c_1 y_1 + c_2 y_2$$

So $x_1 = a, x_2 = b, x_3 = c$ **is solution of (3)**

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

$$= c_1 y_1 + c_2 y_2$$

So $x_1 = a, x_2 = b, x_3 = c$ **is solution of (3)**

Converse need not be true (Try !)

Suppose that $x_1 = a, x_2 = b, x_3 = c$ **is solution of (1) and (2)**

Show that above solution is also a solution of (3).

Consider L.H.S. of (3),

$$c_1 (A_{11}a + A_{12}b + A_{13}c) + c_2 (A_{21}a + A_{22}b + A_{23}c)$$

$$= c_1 y_1 + c_2 y_2$$

So $x_1 = a, x_2 = b, x_3 = c$ **is solution of (3)**

Converse need not be true (Try !)

Note: If X^* is a solution of k linear equations, then X^* is also a solution of a linear combination of those k equations.

Equivalent systems

We say two systems are **equivalent**

Equivalent systems

We say two systems are **equivalent** if each equation in each system is a linear combination of equations in the other system.

Equivalent systems

We say two systems are **equivalent** if each equation in each system is a linear combination of equations in the other system.

Why do we focus on equivalent systems?

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively.

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. Prove that $S_A = S_B$.

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. Prove that $S_A = S_B$.

Let $X \in S_A$.

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. Prove that $S_A = S_B$.

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) ,

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) .

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) . $\implies X$ satisfies every equation in (B) .

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) . $\implies X$ satisfies every equation in (B) . $\implies X \in S_B$

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) . $\implies X$ satisfies every equation in (B) . $\implies X \in S_B$

Hence $S_A \subseteq S_B$

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) . $\implies X$ satisfies every equation in (B) . $\implies X \in S_B$

Hence $S_A \subseteq S_B$

Similarly, $S_B \subseteq S_A$

Theorem 1: Equivalent systems of linear equations have exactly same solutions.

Proof:

Let (A) and (B) be two equivalent systems with solution sets S_A and S_B respectively. **Prove that $S_A = S_B$.**

Let $X \in S_A$. $\implies X$ satisfies every equation in (A) , and every equation in (B) is a linear combination of equations in (A) . $\implies X$ satisfies every equation in (B) . $\implies X \in S_B$

Hence $S_A \subseteq S_B$

Similarly, $S_B \subseteq S_A$

$\implies S_A = S_B$

Problem

Show that the following systems of linear equations are equivalent.

$$\begin{array}{l} x - y = 0 \\ 2x + y = 0 \end{array} \quad \left. \begin{array}{l} \\ \end{array} \right\} \quad \text{---} \quad (I)$$

$$\begin{array}{l} 3x + y = 0 \\ x + y = 0 \end{array} \quad \left. \begin{array}{l} \\ \end{array} \right\} \quad \text{---} \quad (II)$$

Solution

$$3x + y = \frac{1}{3}(x - y) + \frac{4}{3}(2x + y)$$

Solution

$$\begin{aligned}3x + y &= \frac{1}{3}(x - y) + \frac{4}{3}(2x + y) \\x + y &= -\frac{1}{3}(x - y) + \frac{2}{3}(2x + y)\end{aligned}$$

Solution

$$\begin{aligned}3x + y &= \frac{1}{3}(x - y) + \frac{4}{3}(2x + y) \\x + y &= -\frac{1}{3}(x - y) + \frac{2}{3}(2x + y) \\x - y &= (3x + y) - 2(x + y)\end{aligned}$$

Solution

$$\begin{aligned}3x + y &= \frac{1}{3}(x - y) + \frac{4}{3}(2x + y) \\x + y &= -\frac{1}{3}(x - y) + \frac{2}{3}(2x + y) \\x - y &= (3x + y) - 2(x + y) \\2x + y &= \frac{1}{2}(3x + y) + \frac{1}{2}(x + y)\end{aligned}$$

Solution

$$\begin{aligned}3x + y &= \frac{1}{3}(x - y) + \frac{4}{3}(2x + y) \\x + y &= -\frac{1}{3}(x - y) + \frac{2}{3}(2x + y) \\x - y &= (3x + y) - 2(x + y) \\2x + y &= \frac{1}{2}(3x + y) + \frac{1}{2}(x + y)\end{aligned}$$

Note : $AX = 0$ is called a homogeneous system.