

The Ratio Test

Theorem (The Ratio Test)

Let $\sum a_n$ be a series with positive terms and suppose that

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \rho.$$

Then

- (a) *the series converges if $\rho < 1$,*
- (b) *the series diverges if $\rho > 1$ or is infinite,*
- (c) *the test is inconclusive if $\rho = 1$.*

Proof

(a) $\rho < 1$.

Let r be a number between ρ and 1: $\rho < r < 1$.

Proof

(a) $\rho < 1$.

Let r be a number between ρ and 1: $\rho < r < 1$.

Then the number $\epsilon = r - \rho$ is positive.

Since

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \rho,$$

there is an integer N such that

$$n \geq N \Rightarrow \left| \frac{a_{n+1}}{a_n} - \rho \right| < \epsilon$$

This implies that, for $n \geq N$,

$$-\epsilon < \frac{a_{n+1}}{a_n} - \rho < \epsilon \quad \text{or} \quad \rho - \epsilon < \frac{a_{n+1}}{a_n} < \rho + \epsilon = r.$$

That is,

$$\begin{aligned}a_{N+1} &< ra_N, \\a_{N+2} &< ra_{N+1} < r^2 a_N, \\a_{N+3} &< ra_{N+2} < r^3 a_N, \\&\vdots \\a_{N+m} &< ra_{N+m-1} < r^m a_N.\end{aligned}$$

Consider the series $\sum c_n$, where $c_n = a_n$ for $n = 1, 2, \dots, N$ and $c_{N+1} = ra_N$, $c_{N+2} = r^2 a_N, \dots, c_{N+m} = r^m a_N, \dots$

Now $a_n \leq c_n$ for all n and

$$\begin{aligned}\sum_{n=1}^{\infty} c_n &= a_1 + a_2 + \dots + a_{N-1} + a_N + ra_N + r^2 a_N + \dots \\&= a_1 + a_2 + \dots + a_{N-1} + a_N(1 + r + r^2 + \dots)\end{aligned}$$

The geometric series $1 + r + r^2 + \dots$ converges as $|r| < 1$.

That is,

$$\begin{aligned}a_{N+1} &< ra_N, \\a_{N+2} &< ra_{N+1} < r^2 a_N, \\a_{N+3} &< ra_{N+2} < r^3 a_N, \\&\vdots \\a_{N+m} &< ra_{N+m-1} < r^m a_N.\end{aligned}$$

Consider the series $\sum c_n$, where $c_n = a_n$ for $n = 1, 2, \dots, N$ and $c_{N+1} = ra_N$, $c_{N+2} = r^2 a_N, \dots, c_{N+m} = r^m a_N, \dots$

Now $a_n \leq c_n$ for all n and

$$\begin{aligned}\sum_{n=1}^{\infty} c_n &= a_1 + a_2 + \dots + a_{N-1} + a_N + ra_N + r^2 a_N + \dots \\&= a_1 + a_2 + \dots + a_{N-1} + a_N(1 + r + r^2 + \dots)\end{aligned}$$

The geometric series $1 + r + r^2 + \dots$ converges as $|r| < 1$. So $\sum c_n$ converges.

That is,

$$\begin{aligned}a_{N+1} &< ra_N, \\a_{N+2} &< ra_{N+1} < r^2 a_N, \\a_{N+3} &< ra_{N+2} < r^3 a_N, \\&\vdots \\a_{N+m} &< ra_{N+m-1} < r^m a_N.\end{aligned}$$

Consider the series $\sum c_n$, where $c_n = a_n$ for $n = 1, 2, \dots, N$ and $c_{N+1} = ra_N$, $c_{N+2} = r^2 a_N, \dots, c_{N+m} = r^m a_N, \dots$

Now $a_n \leq c_n$ for all n and

$$\begin{aligned}\sum_{n=1}^{\infty} c_n &= a_1 + a_2 + \dots + a_{N-1} + a_N + ra_N + r^2 a_N + \dots \\&= a_1 + a_2 + \dots + a_{N-1} + a_N(1 + r + r^2 + \dots)\end{aligned}$$

The geometric series $1 + r + r^2 + \dots$ converges as $|r| < 1$. So $\sum c_n$ converges. Since $a_n \leq c_n$, $\sum a_n$ also converges.

(b) $1 < \rho \leq \infty$.

From some index M on,

$$\frac{a_{n+1}}{a_n} > 1 \quad \text{and} \quad a_M < a_{M+1} < a_{M+2} < \dots$$

So, the terms of the series do not approach zero as n becomes infinite. Hence the series diverges by the n th Term Test.

(c) $\rho = 1$.

Consider the following two series:

$$\sum_{n=1}^{\infty} \frac{1}{n} \quad \text{and} \quad \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

$$\text{For } \sum_{n=1}^{\infty} \frac{1}{n}: \frac{a_{n+1}}{a_n} = \frac{1/(n+1)}{1/n} = \frac{n}{n+1} \rightarrow 1.$$

$$\text{For } \sum_{n=1}^{\infty} \frac{1}{n^2}: \frac{a_{n+1}}{a_n} = \frac{1/(n+1)^2}{1/n^2} = \left(\frac{n}{n+1} \right)^2 \rightarrow 1.$$

In both cases, $\rho = 1$. But the first series diverges and the second converges.

Example

Investigate the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n + 5}{3^n}.$$

Example

Investigate the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n + 5}{3^n}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{(2^{n+1} + 5)/3^{n+1}}{(2^n + 5)/3^n} = \frac{1}{3} \cdot \frac{2^{n+1} + 5}{2^n + 5}$$

Example

Investigate the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n + 5}{3^n}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{(2^{n+1} + 5)/3^{n+1}}{(2^n + 5)/3^n} = \frac{1}{3} \cdot \frac{2^{n+1} + 5}{2^n + 5} \rightarrow \frac{2}{3}.$$

Example

Investigate the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n + 5}{3^n}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{(2^{n+1} + 5)/3^{n+1}}{(2^n + 5)/3^n} = \frac{1}{3} \cdot \frac{2^{n+1} + 5}{2^n + 5} \rightarrow \frac{2}{3}.$$

The series converges because here $\rho = 2/3 < 1$.

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n}{n!}.$$

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{2^n}{n!}.$$

Solution: For the series $\sum_{n=1}^{\infty} \frac{a^n}{n!}$,

$$\frac{a_{n+1}}{a_n} = \frac{a^{n+1}/(n+1)!}{a^n/n!} = \frac{a}{n+1} \rightarrow 0.$$

The series converges because here $\rho = 0 < 1$.

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

But

$$\frac{a_{n+1}}{a_n} = \frac{2(n+1)}{2n+1} > 1 \text{ for all } n.$$

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

But

$$\frac{a_{n+1}}{a_n} = \frac{2(n+1)}{2n+1} > 1 \text{ for all } n.$$

Thus $a_{n+1} > a_n$ for all n .

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

But

$$\frac{a_{n+1}}{a_n} = \frac{2(n+1)}{2n+1} > 1 \text{ for all } n.$$

Thus $a_{n+1} > a_n$ for all n .

So, $a_1 < a_2 < a_3 < \dots$

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

But

$$\frac{a_{n+1}}{a_n} = \frac{2(n+1)}{2n+1} > 1 \text{ for all } n.$$

Thus $a_{n+1} > a_n$ for all n .

So, $a_1 < a_2 < a_3 < \dots$

Also $a_1 = 2$. Thus a_n does not converge to 0.

Example

Test the convergence of the series

$$\sum_{n=1}^{\infty} \frac{4^n n! n!}{(2n)!}.$$

Solution: Here

$$\frac{a_{n+1}}{a_n} = \frac{4^{n+1}(n+1)!(n+1)!}{(2n+2)!} \cdot \frac{(2n)!}{4^n n! n!} = \frac{4(n+1)(n+1)}{(2n+2)(2n+1)} = \frac{2(n+1)}{2n+1} \rightarrow 1.$$

Because the limit $\rho = 1$, we cannot decide whether the series converges from the ratio test.

But

$$\frac{a_{n+1}}{a_n} = \frac{2(n+1)}{2n+1} > 1 \text{ for all } n.$$

Thus $a_{n+1} > a_n$ for all n .

So, $a_1 < a_2 < a_3 < \dots$

Also $a_1 = 2$. Thus a_n does not converge to 0.

Hence the series diverges.

The Root Test

Theorem (The Root Test)

Let $\sum a_n$ be a series with $a_n \geq 0$ for $n \geq N$ (N an integer) and suppose that

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \rho.$$

The Root Test

Theorem (The Root Test)

Let $\sum a_n$ be a series with $a_n \geq 0$ for $n \geq N$ (N an integer) and suppose that

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \rho.$$

Then

(a) the series converges if $\rho < 1$,

The Root Test

Theorem (The Root Test)

Let $\sum a_n$ be a series with $a_n \geq 0$ for $n \geq N$ (N an integer) and suppose that

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \rho.$$

Then

- (a) the series converges if $\rho < 1$,
- (b) the series diverges if $\rho > 1$ or ρ is infinite,

The Root Test

Theorem (The Root Test)

Let $\sum a_n$ be a series with $a_n \geq 0$ for $n \geq N$ (N an integer) and suppose that

$$\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \rho.$$

Then

- (a) the series converges if $\rho < 1$,
- (b) the series diverges if $\rho > 1$ or ρ is infinite,
- (c) the test is inconclusive if $\rho = 1$.

Examples

Which of the following series converges and which diverges?

(a) $\sum_{n=1}^{\infty} \frac{n^2}{2^n}$

Examples

Which of the following series converges and which diverges?

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{n^2}$$

Examples

Which of the following series converges and which diverges?

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{n^2} \quad (c) \sum_{n=1}^{\infty} \left(\frac{1}{1+n} \right)^n$$

Examples

Which of the following series converges and which diverges?

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{n^2} \quad (c) \sum_{n=1}^{\infty} \left(\frac{1}{1+n} \right)^n$$

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \text{ converges because } \sqrt[n]{\frac{n^2}{2^n}} = \frac{(\sqrt[n]{n})^2}{2} \rightarrow \frac{1}{2} < 1.$$

Examples

Which of the following series converges and which diverges?

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{n^2} \quad (c) \sum_{n=1}^{\infty} \left(\frac{1}{1+n} \right)^n$$

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \text{ converges because } \sqrt[n]{\frac{n^2}{2^n}} = \frac{(\sqrt[n]{n})^2}{2} \rightarrow \frac{1}{2} < 1.$$

$$(b) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \text{ diverges because } \sqrt[n]{\frac{2^n}{n^2}} = \frac{2}{(\sqrt[n]{n})^2} \rightarrow \frac{2}{1} > 1.$$

Examples

Which of the following series converges and which diverges?

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \quad (b) \sum_{n=1}^{\infty} \frac{2^n}{n^2} \quad (c) \sum_{n=1}^{\infty} \left(\frac{1}{1+n} \right)^n$$

$$(a) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \text{ converges because } \sqrt[n]{\frac{n^2}{2^n}} = \frac{(\sqrt[n]{n})^2}{2} \rightarrow \frac{1}{2} < 1.$$

$$(b) \sum_{n=1}^{\infty} \frac{n^2}{2^n} \text{ diverges because } \sqrt[n]{\frac{2^n}{n^2}} = \frac{2}{(\sqrt[n]{n})^2} \rightarrow \frac{2}{1} > 1.$$

$$(c) \sum_{n=1}^{\infty} \left(\frac{1}{1+n} \right)^n \text{ converges because } \sqrt[n]{\left(\frac{1}{1+n} \right)^n} = \frac{1}{1+n} \rightarrow 0 < 1.$$

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Solution: Here

$$\sqrt[n]{a_n} = \begin{cases} \sqrt[n]{n}/2, & n \text{ odd} \\ 1/2, & n \text{ even.} \end{cases}$$

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Solution: Here

$$\sqrt[n]{a_n} = \begin{cases} \sqrt[n]{n}/2, & n \text{ odd} \\ 1/2, & n \text{ even.} \end{cases}$$

Therefore,

$$\frac{1}{2} \leq \sqrt[n]{a_n} \leq \frac{\sqrt[n]{n}}{2}.$$

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Solution: Here

$$\sqrt[n]{a_n} = \begin{cases} \sqrt[n]{n}/2, & n \text{ odd} \\ 1/2, & n \text{ even.} \end{cases}$$

Therefore,

$$\frac{1}{2} \leq \sqrt[n]{a_n} \leq \frac{\sqrt[n]{n}}{2}.$$

We know that $\sqrt[n]{n} \rightarrow 1$.

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Solution: Here

$$\sqrt[n]{a_n} = \begin{cases} \sqrt[n]{n}/2, & n \text{ odd} \\ 1/2, & n \text{ even.} \end{cases}$$

Therefore,

$$\frac{1}{2} \leq \sqrt[n]{a_n} \leq \frac{\sqrt[n]{n}}{2}.$$

We know that $\sqrt[n]{n} \rightarrow 1$. So, $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 1/2$ by the Sandwich Theorem.

Example

Let $a_n = \begin{cases} n/2^n, & n \text{ odd} \\ 1/2^n, & n \text{ even.} \end{cases}$ Does $\sum a_n$ converge?

Solution: Here

$$\sqrt[n]{a_n} = \begin{cases} \sqrt[n]{n}/2, & n \text{ odd} \\ 1/2, & n \text{ even.} \end{cases}$$

Therefore,

$$\frac{1}{2} \leq \sqrt[n]{a_n} \leq \frac{\sqrt[n]{n}}{2}.$$

We know that $\sqrt[n]{n} \rightarrow 1$. So, $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = 1/2$ by the Sandwich Theorem.

Thus here the limit is $\rho < 1$. Hence the series converges by the Root Test.

Alternating Series

Definition

A series in which the terms are alternatively positive and negative is called an **alternating series**.

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

$$1 - 2 + 3 - 4 + 5 - 6 + \dots + (-1)^{n+1} n + \dots$$

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

$$1 - 2 + 3 - 4 + 5 - 6 + \dots + (-1)^{n+1} n + \dots$$

The first series, called the **alternating harmonic series**, converges.

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

$$1 - 2 + 3 - 4 + 5 - 6 + \dots + (-1)^{n+1} n + \dots$$

The first series, called the **alternating harmonic series**, converges.

The second series, a geometric series with common ratio $r = -1/2$, converges.

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

$$1 - 2 + 3 - 4 + 5 - 6 + \dots + (-1)^{n+1} n + \dots$$

The first series, called the **alternating harmonic series**, converges.

The second series, a geometric series with common ratio $r = -1/2$, converges.

The third series diverges because

Examples

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^{n+1}}{n} + \dots$$

$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

$$1 - 2 + 3 - 4 + 5 - 6 + \dots + (-1)^{n+1} n + \dots$$

The first series, called the **alternating harmonic series**, converges.

The second series, a geometric series with common ratio $r = -1/2$, converges.

The third series diverges because the n th term does not approach zero.

Homework

Let $\{a_n\}$ be a sequence such that the subsequences $\{a_{2m}\}$ and $\{a_{2m+1}\}$ both converge to the same limit l .

Homework

Let $\{a_n\}$ be a sequence such that the subsequences $\{a_{2m}\}$ and $\{a_{2m+1}\}$ both converge to the same limit l . Then show that $a_n \rightarrow l$.

The Alternating Series Test

Theorem (The Alternating Series Test (Leibniz's Theorem))

The series

$$\sum_{n=1}^{\infty} (-1)^{n+1} u_n = u_1 - u_2 + u_3 - u_4 + \dots$$

converges if all three of the following conditions are satisfied:

The Alternating Series Test

Theorem (The Alternating Series Test (Leibniz's Theorem))

The series

$$\sum_{n=1}^{\infty} (-1)^{n+1} u_n = u_1 - u_2 + u_3 - u_4 + \dots$$

converges if all three of the following conditions are satisfied:

1. *The u_n 's are all positive.*

The Alternating Series Test

Theorem (The Alternating Series Test (Leibniz's Theorem))

The series

$$\sum_{n=1}^{\infty} (-1)^{n+1} u_n = u_1 - u_2 + u_3 - u_4 + \dots$$

converges if all three of the following conditions are satisfied:

- 1. The u_n 's are all positive.*
- 2. $u_n \geq u_{n+1}$ for all $n \geq N$, for some integer N .*

The Alternating Series Test

Theorem (The Alternating Series Test (Leibniz's Theorem))

The series

$$\sum_{n=1}^{\infty} (-1)^{n+1} u_n = u_1 - u_2 + u_3 - u_4 + \dots$$

converges if all three of the following conditions are satisfied:

1. *The u_n 's are all positive.*
2. *$u_n \geq u_{n+1}$ for all $n \geq N$, for some integer N .*
3. *$u_n \rightarrow 0$.*

Proof:

Let us assume that $N = 1$.

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$s_{2m} = (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m})$$

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned} s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m} \end{aligned}$$

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

The second equality shows that $s_{2m} \leq u_1$.

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

The second equality shows that $s_{2m} \leq u_1$.

So, $\{s_{2m}\}$ is monotonically increasing and bounded above.

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

The second equality shows that $s_{2m} \leq u_1$.

So, $\{s_{2m}\}$ is monotonically increasing and bounded above. So, it converges, say

$$\lim_{n \rightarrow \infty} s_{2m} = l.$$

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

The second equality shows that $s_{2m} \leq u_1$.

So, $\{s_{2m}\}$ is monotonically increasing and bounded above. So, it converges, say

$$\lim_{n \rightarrow \infty} s_{2m} = l.$$

Also $u_{2m+1} \rightarrow 0$.

Proof:

Let us assume that $N = 1$.

If n is an even integer, say $n = 2m$, then the sum of the first n terms is

$$\begin{aligned}s_{2m} &= (u_1 - u_2) + (u_3 - u_4) + \dots + (u_{2m-1} - u_{2m}) \\ &= u_1 - (u_2 - u_3) - (u_4 - u_5) - \dots - (u_{2m-2} - u_{2m-1}) - u_{2m}\end{aligned}$$

The first equality shows that the s_{2m} is the sum of m non-negative terms. Hence $s_{2m+2} \geq s_{2m}$.

The second equality shows that $s_{2m} \leq u_1$.

So, $\{s_{2m}\}$ is monotonically increasing and bounded above. So, it converges, say

$$\lim_{n \rightarrow \infty} s_{2m} = l.$$

Also $u_{2m+1} \rightarrow 0$. Hence

$$s_{2m+1} = s_{2m} + u_{2m+1} \rightarrow l + 0 = l.$$

Thus we have that $s_{2m} \rightarrow l$ and $s_{2m+1} \rightarrow l$.

Thus we have that $s_{2m} \rightarrow l$ and $s_{2m+1} \rightarrow l$.

Hence $s_n \rightarrow l$.

Thus we have that $s_{2m} \rightarrow l$ and $s_{2m+1} \rightarrow l$.

Hence $s_n \rightarrow l$. This means that the alternating series converges.

Example

The alternating harmonic series

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

satisfies the three requirements of the alternating series theorem with $N = 1$.

Example

The alternating harmonic series

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

satisfies the three requirements of the alternating series theorem with $N = 1$. So, it converges.