

Analysis of Previous Satellite-Based Problem Statements and AI Solutions

1. Introduction

Satellite imagery has become one of the most valuable sources of geospatial information for solving real-world problems. Every day, satellites launched by organizations such as the Indian Space Research Organisation (ISRO), National Remote Sensing Centre (NRSC), NASA, and the European Space Agency (ESA) capture high-resolution images of the Earth's surface. These images provide crucial information about forests, agricultural fields, rivers, roads, cities, coastlines, and natural disasters.

Traditionally, experts analyze these satellite images manually, which is both time-consuming and labor-intensive. Since India covers more than 3.28 million square kilometres, continuously monitoring every region through manual inspection is practically impossible. Artificial Intelligence (AI) and Deep Learning provide an efficient solution by automatically analyzing satellite images, detecting changes, identifying objects, and generating useful reports for decision-makers.

Satellite Image AI combines Computer Vision, Deep Learning, Remote Sensing, and Geographic Information Systems (GIS) to automate tasks such as flood detection, urban expansion monitoring, crop health assessment, forest fire detection, and disaster management. Because of these applications, satellite image analysis has become a recurring theme in the Smart India Hackathon (SIH) and other government innovation programs.

This report analyzes several satellite-related problem statements and proposes AI-based solutions that improve accuracy, reduce manual effort, and support faster decision-making.

2. Why Satellite Image AI is Important

Satellite imagery offers several advantages over traditional field surveys and manual inspections.

Advantages

- Covers very large geographical areas
- Provides regular monitoring through periodic satellite observations
- Reduces the need for expensive field inspections
- Enables early detection of environmental changes
- Supports disaster management and emergency response

- Helps government agencies make data-driven decisions

Major Applications

- Human activity monitoring
- Flood detection
- Forest fire monitoring
- Crop health assessment
- Railway infrastructure monitoring
- Urban planning
- Environmental conservation
- Disaster management
- Air quality monitoring

These applications explain why government organizations continue to introduce satellite-related challenges in Smart India Hackathon.

Problem Statement 1: Human Activity Change Detection Using Satellite Images

Organization: Indian Space Research Organisation (ISRO)

Objective: Develop an AI system capable of automatically detecting and classifying human-induced land changes by comparing satellite images captured at different time periods.

Background

Human activities such as illegal construction, road expansion, deforestation, urban development, and land encroachment continuously modify the Earth's surface. Monitoring these changes manually requires experts to compare thousands of satellite images, making the process slow, expensive, and prone to human error.

Government agencies require continuous monitoring to protect forests, prevent illegal activities, update land records, and support urban planning. Since satellites regularly capture images of the same location, AI can automate this comparison process and significantly reduce manual effort.

Why this Problem Exists

Government departments need to continuously monitor land resources to:

- Detect unauthorized construction
- Monitor urban expansion
- Identify deforestation
- Protect environmentally sensitive areas
- Update land-use records

Manual comparison of satellite images is difficult because India covers a vast geographical area, making continuous monitoring almost impossible.

Current Challenges

- Large volume of satellite imagery
- Manual image comparison
- Seasonal vegetation changes causing false detections
- Cloud cover and lighting variations
- Difficulty identifying small changes over large regions

Existing Limitations

Traditional methods rely heavily on GIS experts manually comparing satellite images. These methods are time-consuming, expensive, and often fail to detect important changes quickly. Many illegal activities remain unnoticed until significant environmental damage has already occurred.

Proposed AI Solution

To overcome these limitations, we propose an AI-powered Human Activity Change Detection System that automatically compares satellite images captured at different time intervals. The system first identifies regions where significant changes have occurred and then analyzes whether those changes are caused by human activities or natural seasonal variations. This additional verification step helps reduce false alerts caused by vegetation growth or weather changes.

Once the changes are verified, the AI model classifies them into categories such as new construction, road expansion, deforestation, or land encroachment. The final results are displayed on an interactive GIS dashboard where government officials can review the affected areas and generate automated reports for further action.

Workflow



Expected Output

- Construction detection
- Road expansion monitoring
- Deforestation detection
- Urban growth monitoring
- Land encroachment identification

Suitable AI Models

AI Model	Purpose
ChangeFormer	Detect changes between satellite images
Siamese Neural Network	Compare images captured at different times
Vision Transformer (ViT)	Learn complex spatial features
U-Net	Pixel-level segmentation
SegFormer	Semantic segmentation

Suitable Datasets

- Sentinel-2
- Landsat-8/9
- LEVIR-CD (Building Change Detection)
- WHU Building Dataset
- Hansen Global Forest Change Dataset (for forest monitoring)

Problem Statement 2: SAR Image Colorization

Organization: Indian Space Research Organisation (ISRO)

Objective: Develop an AI-based system that converts grayscale Synthetic Aperture Radar (SAR) images into visually meaningful colour images while preserving important structural information.

Background

Unlike optical satellites, Synthetic Aperture Radar (SAR) satellites use microwave signals to capture images. This allows them to operate during nighttime and cloudy weather, making them extremely useful during disasters such as floods and cyclones. However, SAR images are grayscale and often difficult for non-experts to interpret.

Colourized SAR images can improve understanding and help government agencies quickly identify important geographical features.

Why this Problem Exists

Although SAR imagery is highly reliable, its grayscale appearance makes interpretation difficult. Disaster management teams and government officials often require visually understandable images to quickly identify water bodies, buildings, roads, vegetation, and damaged infrastructure.

Current Challenges

- Speckle noise
- Lack of colour information
- Difficult visual interpretation

- Structural information must be preserved
- Limited paired SAR-optical datasets

Existing Limitations


Traditional colourization methods mainly focus on improving image appearance but sometimes generate unrealistic colours or alter important structures, making the images unsuitable for practical applications.

Proposed AI Solution

The proposed system uses Generative Artificial Intelligence to automatically transform grayscale SAR images into realistic colourized images. Before colourization, the system removes unwanted image noise to improve image quality. It then learns the relationship between SAR images and corresponding optical satellite images, enabling it to generate visually meaningful colour representations while preserving important structures such as roads, rivers, buildings, and coastlines.

The final colourized images can be displayed alongside the original SAR images, allowing disaster management teams and researchers to better understand affected regions and make faster decisions during emergency situations.

Workflow



Expected Output

- Colourized SAR imagery

- Better disaster interpretation
- Improved land-use visualization
- Enhanced infrastructure identification

Suitable AI Models

AI Model	Purpose
Pix2Pix GAN	Image-to-image translation
CycleGAN	Colour generation
U-Net	Image reconstruction
Vision Transformer	Feature extraction
Stable Diffusion (Future Work)	High-quality image enhancement

Suitable Datasets

- SEN1-2 Dataset
- Sentinel-1
- Sentinel-2

Problem Statement 3: Satellite-Based Air Quality Downscaling

Organization: Indian Space Research Organisation (ISRO)

Objective: Develop an AI-powered system that generates high-resolution air quality maps by combining satellite imagery with environmental and meteorological information.

Background

Satellites measure atmospheric pollutants such as PM2.5 and NO₂ over large geographical regions. However, these measurements are generally available at coarse resolutions, making them less useful for city-level pollution monitoring. Government agencies require more

detailed pollution maps to identify local pollution hotspots and support environmental planning.

Why this Problem Exists

Air pollution depends on multiple factors such as traffic density, industrial emissions, weather conditions, and land use. Low-resolution satellite data cannot accurately capture these local variations, limiting its usefulness for public health and urban planning.

Current Challenges

- Low spatial resolution
- Missing observations due to cloud cover
- Complex environmental relationships
- Limited ground monitoring stations

Existing Limitations


Traditional interpolation methods estimate pollution between monitoring stations but often fail to represent local pollution hotspots accurately. They also struggle in areas where very few monitoring stations are available.


Proposed AI Solution

The proposed AI system combines satellite observations with weather information, land-use data, and ground monitoring station readings to generate high-resolution pollution maps. Instead of simply enlarging satellite images, the AI learns how environmental conditions influence pollution levels and produces more accurate city-scale estimates.

To improve reliability, the predicted pollution values are compared with actual readings collected from monitoring stations. Any differences are used to refine the final output, helping government agencies generate more accurate pollution maps for environmental monitoring and public health planning.

Workflow





Expected Output

- High-resolution PM2.5 maps
- NO₂ concentration maps
- Pollution hotspot identification
- Environmental monitoring dashboard

Suitable AI Models

AI Model	Purpose
CNN	Feature extraction
Swin Transformer	Spatial feature learning
Graph Neural Network (GNN) (Advanced)	Spatial relationship modeling
ESRGAN (Optional)	Image enhancement
U-Net	Spatial refinement

Suitable Datasets

- Sentinel-5P
- MODIS
- ERA5 Meteorological Dataset
- CPCB Air Quality Data
- OpenAQ

References

- Indian Space Research Organisation (ISRO)
- National Remote Sensing Centre (NRSC)
- European Space Agency (ESA)
- Sentinel-1 & Sentinel-2 Missions
- Landsat 8/9 (NASA & USGS)
- Hansen Global Forest Change Dataset
- LEVIR-CD Dataset
- WHU Building Dataset
- SEN1-2 Dataset
- CPCB Air Quality Monitoring Data

Advanced AI-Based Satellite Image Solutions for Smart India Hackathon (SIH)

Problem Statement 4: Flood Mapping and Damage Assessment using Satellite Images

Organization: Inspired by Disaster Management Applications (ISRO / NRSC)

Objective: Develop an AI-powered system that automatically detects flooded regions and estimates infrastructure damage using satellite imagery.

Background

Floods are among the most destructive natural disasters in India, affecting millions of people every year. States such as Assam, Bihar, Kerala, and Uttarakhand experience severe flooding due to heavy rainfall and overflowing rivers. Government agencies use satellite

images to understand the extent of flooding because they provide a large-scale view of affected regions.

However, manually analyzing these images takes several hours, delaying rescue operations and relief planning.

Why this Problem Exists

Floodwater spreads rapidly and continuously changes its boundaries. Authorities need immediate information about submerged villages, damaged roads, affected farmlands, and safe evacuation routes. Traditional field surveys are slow and cannot cover large areas within a short time.

Current Challenges

- Rapidly changing flood boundaries
- Cloud cover affecting optical imagery
- Large geographical coverage
- Delayed manual assessment
- Difficulty estimating infrastructure damage

Existing Limitations

Current flood assessment methods rely heavily on GIS experts manually interpreting satellite images. This process is time-consuming and often delays rescue operations and resource allocation.

Proposed AI Solution

The proposed system automatically compares satellite images captured before and after a flood event to identify flooded regions and estimate the extent of damage. By using deep learning models, the system can distinguish water bodies from roads, buildings, agricultural fields, and other land features without requiring manual inspection.

The detected flood boundaries are displayed on an interactive GIS dashboard along with information about damaged infrastructure and high-risk areas. This enables disaster management authorities to prioritize rescue operations, allocate emergency resources efficiently, and generate automated flood assessment reports.

Workflow



Expected Output

- Flood extent map
- Damaged road detection
- Damaged building estimation
- Agricultural loss assessment
- Rescue priority zones

Suitable AI Models

AI Model	Purpose
U-Net	Flood segmentation
DeepLabV3+	Water body extraction
SegFormer	Semantic segmentation
Vision Transformer	Feature extraction
ChangeFormer	Flood change detection

Suitable Datasets

- Sentinel-1 SAR
- Sentinel-2
- SpaceNet
- xView2 Disaster Dataset
- Copernicus Emergency Management Service

Problem Statement 5: Urban Encroachment Detection using Satellite Images

Organization: Inspired by Smart City and Urban Planning Applications

Objective: Develop an AI-powered system that automatically identifies illegal constructions and urban encroachments using satellite imagery.

Background

Rapid urbanization has increased unauthorized construction and encroachment on government land, rivers, forests, and protected areas. Monitoring these activities manually is difficult because cities expand continuously and require frequent inspections.

Satellite imagery provides a cost-effective way to monitor urban development over large areas and detect unauthorized land use.

Why this Problem Exists

Government authorities need accurate information about illegal settlements, construction activities, and land encroachments to enforce regulations and improve urban planning. Manual inspections require significant manpower and cannot keep pace with rapid urban expansion.

Current Challenges

- Rapid city expansion
- Frequent construction activities
- Large urban coverage

- Difficulty distinguishing legal and illegal structures
- Regular monitoring requirements

Existing Limitations


Traditional urban monitoring depends on periodic surveys and manual interpretation of satellite images. This often delays the identification of unauthorized developments and increases enforcement costs.

Proposed AI Solution

The proposed AI system continuously analyzes satellite images of urban regions to detect newly constructed buildings, unauthorized settlements, and encroachments on protected land. By comparing historical and recent satellite imagery, the AI identifies changes in built-up areas and classifies whether they represent planned development or suspicious land use.

The detected locations are displayed on a GIS dashboard where authorities can review changes, compare historical images, and generate reports for further investigation. Continuous monitoring enables faster identification of urban encroachments while reducing manual inspection efforts.

Workflow



Expected Output

- Illegal construction detection
- Urban expansion map

- Encroachment identification
- Infrastructure development monitoring

Suitable AI Models

AI Model	Purpose
ChangeFormer	Building change detection
Mask R-CNN	Building extraction
U-Net	Urban segmentation
Vision Transformer	Urban scene understanding
YOLOv11	Building detection

Suitable Datasets

- SpaceNet
- Sentinel-2
- LEVIR-CD
- WHU Building Dataset
- OpenStreetMap (OSM)

Problem Statement 6: Railway Track Hazard Detection using Satellite Images

Organization: Inspired by Railway Infrastructure Monitoring

Objective: Develop an AI-powered monitoring system that detects hazards affecting railway corridors using satellite imagery.

Background

India has one of the largest railway networks in the world. Heavy rainfall, landslides, flooding, soil erosion, and fallen trees frequently disrupt railway operations. Since railway

tracks pass through forests, mountains, and remote regions, continuous manual inspection is difficult.

Why this Problem Exists

Unexpected hazards can block railway tracks and increase the risk of accidents. Early detection allows authorities to perform preventive maintenance before train operations are affected.

Current Challenges

- Large railway network
- Difficult terrain
- Weather-related hazards
- High inspection costs
- Limited monitoring frequency

Existing Limitations


Current inspection methods depend on patrol teams, drones, and scheduled maintenance. These methods cannot continuously monitor thousands of kilometres of railway tracks.

Proposed AI Solution

The proposed system continuously analyzes satellite images of railway corridors to identify hazards such as landslides, flooding, erosion, and fallen trees. Deep learning models automatically detect high-risk locations and estimate the severity of each hazard.

The identified hazards are displayed on a GIS dashboard, allowing railway authorities to prioritize inspections and maintenance activities. Automated monitoring improves railway safety while reducing inspection time and operational costs.

Workflow



Risk Classification



GIS Dashboard



Maintenance Report

Expected Output

- Landslide detection
- Flooded railway sections
- Fallen tree identification
- High-risk railway corridors

Suitable AI Models

AI Model	Purpose
YOLOv11	Object detection
SegFormer	Landslide segmentation
Mask R-CNN	Obstacle extraction
Vision Transformer	Scene understanding
U-Net	Flood segmentation

Suitable Datasets

- Sentinel-2
- SpaceNet
- Landsat-8/9
- OpenRailwayMap
- Copernicus DEM

Problem Statement 7: Crop Health Monitoring using Satellite Images

Organization: Inspired by Precision Agriculture Applications

Objective: Develop an AI system that monitors crop health and identifies crop stress using satellite imagery.

Background

Agriculture is one of India's most important economic sectors. Crop diseases, water stress, nutrient deficiencies, and pest attacks significantly reduce agricultural productivity. Traditional crop monitoring requires frequent field visits, which are difficult over large agricultural regions.

Why this Problem Exists

Farmers and agricultural departments require continuous crop monitoring to detect problems early and improve crop yield. Satellite imagery enables regular monitoring of large agricultural areas without physical inspections.

Current Challenges

- Large agricultural coverage
- Seasonal weather changes
- Early disease detection
- Water stress identification
- Crop diversity

Existing Limitations


Traditional crop monitoring depends mainly on manual observations and localized surveys, making large-scale monitoring difficult.

Proposed AI Solution

The proposed AI platform analyzes multispectral satellite imagery to evaluate vegetation health and identify early signs of crop stress, disease, and water deficiency. By calculating vegetation indices and applying deep learning models, the system classifies crop conditions and predicts potential yield losses.

The generated crop health maps help farmers and agricultural departments identify problem areas quickly, enabling timely intervention and better resource management.

Workflow



Expected Output

- Healthy crop map
- Crop stress detection
- Disease hotspot identification
- Yield estimation

Suitable AI Models

AI Model	Purpose
EfficientNet	Crop classification
ResNet	Disease detection
U-Net	Crop segmentation
Vision Transformer	Feature extraction
LSTM	Yield prediction

Suitable Datasets

- Sentinel-2
- BigEarthNet
- EuroSAT
- Radiant Earth MLHub

Problem Statement 8: Forest Fire Detection and Burned Area Assessment

Organization: Inspired by Forest and Environmental Monitoring Applications

Objective: Develop an AI-powered system that automatically detects forest fires and estimates burned areas using satellite imagery.

Background

Forest fires destroy biodiversity, wildlife habitats, and natural resources while increasing carbon emissions. Detecting fires at an early stage is essential for minimizing environmental damage and supporting emergency response teams.

Why this Problem Exists

Traditional fire monitoring methods depend on watch towers, field observations, and delayed satellite interpretation. Fires often spread rapidly before authorities receive accurate information.

Current Challenges

- Large forest coverage
- Dense smoke
- Rapid fire spread
- Difficult terrain
- Delayed reporting

Existing Limitations


Existing monitoring systems often detect fires after significant damage has already occurred and cannot provide continuous large-scale monitoring.

Proposed AI Solution

The proposed AI system continuously analyzes satellite imagery to identify active fire regions, smoke plumes, and burned vegetation. Deep learning models automatically estimate the extent of fire damage and classify the severity of affected regions.

The results are displayed on an interactive GIS dashboard that helps forest departments monitor fire progression, prioritize emergency response, and assess post-fire damage efficiently.

Workflow



Expected Output

- Fire location map
- Burned area estimation
- Fire severity classification
- Emergency response planning

Suitable AI Models

AI Model	Purpose
----------	---------

YOLOv11	Fire detection
U-Net	Burned area segmentation
DeepLabV3+	Smoke segmentation
EfficientNet	Fire severity classification
Vision Transformer	Large-scale forest analysis

Suitable Datasets

- MODIS Fire Dataset
- VIIRS Active Fire Dataset
- Sentinel-2
- Landsat-9

Conclusion

The problem statements discussed in this section demonstrate how Artificial Intelligence, Computer Vision, and Remote Sensing can be applied to solve real-world challenges in disaster management, agriculture, transportation, urban planning, and environmental conservation. By combining satellite imagery with modern deep learning techniques, these systems can automate monitoring tasks, improve decision-making, and provide scalable solutions for government organizations.