

Review

Caveolin and cavin family members: Dual roles in cancer

Reshu Gupta^{*}, Chirine Toufaily, Borhane Annabi

Laboratoire d'Oncologie Moléculaire, Centre de Recherche BioMed, Département de Chimie, Université du Québec à Montréal, Québec H3C 3P8, Canada

ARTICLE INFO

Article history:

Received 30 January 2014

Accepted 4 September 2014

Available online 21 September 2014

Keywords:

Caveolin

Cavin

Cancer

ABSTRACT

Caveolae are specialized plasma membrane subdomains with distinct lipid and protein compositions, which play an essential role in cell physiology through regulation of trafficking and signaling functions. The structure and functions of caveolae have been shown to require the proteins caveolins. Recently, members of the cavin protein family were found to be required, in concert with caveolins, for the formation and function of caveolae. Caveolins have a paradoxical role in the development of cancer formation. They have been involved in both tumor suppression and oncogenesis, depending on tumor type and progress stage. High expression of caveolins and cavins leads to inhibition of cancer-related pathways, such as growth factor signaling pathways. However, certain cancer cells that express caveolins and cavins have been shown to be more aggressive and metastatic because of their increased potential for anchorage-independent growth. Here, we will survey the functional roles of caveolins and of different cavin family members in cancer regulation.

© 2014 Elsevier B.V. and Société française de biochimie et biologie Moléculaire (SFBBM). All rights reserved.

1. Introduction

The plasma membrane is a dynamic multidomain system heavily decorated with small «flask-shaped» invaginations of 60–80 nm diameter, known as caveolae, which were first observed in 1953 [1]. As further demonstrated by electron microscopy in unfixed fast-frozen material, caveolae have been shown to also exhibit an open neck structure [2,3]. These structures are enriched with various signaling molecules including some cell surface receptors which are attached through a lipid anchor. Caveolae occur at different densities in different cell types, being most prominent in fibroblasts, vascular endothelial cells, adipocytes and epithelial cells. The principal membrane components of caveolae are a family of integral membrane proteins known as caveolins, which are intimately involved in caveolar function. Caveolae and caveolins are involved in a variety of cellular processes including endocytosis, lipid homeostasis, signal transduction and tumorigenesis. Approximately 144

individual caveolin molecules have been estimated as being incorporated in a single caveolar structure [4]. Recently, a new family of membrane proteins was discovered, named cavins, and were shown to stabilize the oligodimerization of caveolins as well as caveolae formation (Fig. 1) [5–7]. Both caveolins and cavins play a major role in the formation and function of caveolae, and are involved in several physiological and pathological processes.

The roles of caveolin and of cavin in cancer regulation are now well recognized and known to be dependent on cancer cell types and conditions. Although, under some conditions, Caveolin-1 (Cav-1), a member of the caveolin family protein, may suppress tumorigenesis, it is also known for its association with and contribution to malignant progression through various mechanisms. Among these, Cav-1 expression was found to correlate with resistance to ionizing radiation [8,9]. Cav-1, via its scaffolding domain, interacts with a variety of specific proteins that are localized in lipid rafts and caveolar membranes, such as receptor tyrosine kinases, platelet-derived growth factor receptor (PDGFR), epidermal growth factor receptor (EGFR), endothelial nitric oxide synthase (eNOS), proteins involved in calcium transport, H-Ras, integrins, nerve growth factor, serine/threonine kinases, phospholipases, G protein-coupled receptors and Src family kinases [10–20]. Such interactions subsequently generate compartmentalized signaling platforms which facilitate the signaling cascades that contribute to cancer regulation. A high level of intracellular Cav-1 expression is associated with metastatic progression of many types of cancers, such as

Abbreviations: Cav, caveolin; CRC, colorectal cancer; CSD, caveolin scaffolding domain; PDGFR, platelet-derived growth factor receptor; EGFR, epidermal growth factor receptor; eNOS, endothelial nitric oxide synthase; ETS, erythroblastosis virus transforming protein; PEA3, polyoma-virus enhancer activator; MMP, matrix metalloproteinases; RCC, renal cell carcinoma; ROS, reactive oxygen species; STB, β -subunit of Shiga toxin.

^{*} Corresponding author. Current address: A-254, Sector-2, Rohini, New Delhi 110085, India. Tel.: +91 8800681780.

E-mail address: sgreshu12@outlook.com (R. Gupta).




Fig. 1. Caveolin oligomer formation is regulated upon interaction with cavin members. Caveolin-1 and caveolin-2 can exist as monomers, homodimers, or heterodimers. The presence of Cavin-1 and of Cavin-2 allows Caveolin-1/Caveolin-2 to interact within the cell membrane to form heterooligomers in caveolae. In the absence of Cavin-1 or Cavin-2, caveolin oligomers embedded in the cell membrane cannot contribute to caveolae formation.

prostate [21,22], lung [23], renal [24] and esophageal cancers [25]. The expression levels of caveolins and cavins vary in different types of tumor which affects their roles in carcinogenesis (Table 1). In this review, we will discuss the distinctive properties of caveolins and cavins in various cancer types, especially their structures, types, and involvement in cancer regulation. Finally, we will discuss the possibility of biologically targeting these proteins for the development of new and more efficient therapies for cancer cure. Unquestionably, the abilities of caveolins and cavins to modulate signaling has important implications in the process of tumor formation.

2. The caveolin family

The caveolin family is composed of three isoforms of caveolin proteins in mammalian cells: Caveolin-1 (Cav-1), Caveolin-2 (Cav-

2) and Caveolin-3 (Cav-3) [26,27]. Most caveolin proteins are expressed in different types of tissues. However, Cav-3 is predominantly expressed in cytoskeletal muscle tissue. All three proteins have an N-terminal Caveolin Scaffolding Domain (CSD), a transmembrane domain and a C-terminal domain. Both the C-terminal and CSD are sufficient for caveolin attachment to the membrane, even without the involvement of the membrane spanning domain [28].

3. Caveolin-1

Caveolin-1 (Cav-1), discovered by immunocytochemistry as a major component of caveolae, is a 178-amino acid protein that is localized in plasma membrane caveolae, Golgi apparatus and trans-Golgi-derived transport vesicles [28–30]. Cav-1 is expressed either as a soluble cytoplasmic form or as a secreted form, depending on the

Table 1

The dual promoting/suppressive roles of cavin and caveolin members in different types of cancer is a consequence of their expression levels.

| Type of cancer | Cav-1 | Cav-2 | Cav-3 | Cavin-1 | Cavin-2 | Cavin-3 | Mode of action | References |
|--------------------------|-------|-------|-------|---------|---------|---------|---------------------------|---------------|
| Thyroid | High | High | | | | | Tumor Suppressor/Promoter | [147] |
| Breast | Low | Low | | | | | Tumor Promoter | [231] |
| | High | High | | High | | | Tumor Promoter | [163,232,203] |
| | Low | Low | | Low | Low | Low | Tumor Suppressor | [164,209] |
| Prostate | High | High | | | | | Tumor Promoter | [219,235] |
| | | | | Low | | | Tumor Suppressor | [236] |
| Hepatocellular carcinoma | High | | | High | | | Tumor Promoter | [234] |
| Pancreatic | High | | | | | | Tumor Promoter | [198] |
| Urinary bladder | High | High | | | | | Tumor Promoter | [160] |
| Oesophagous | High | High | | | | | Tumor Promoter | [159] |
| Colon | High | | | | | | Tumor Promoter | [237] |
| Mammary | | | High | | | | Tumor Promoter | [238] |
| Colorectal | | | | | | Low | Tumor Suppressor | [249] |
| Ovarian | | | | | | Low | Tumor Suppressor | [239] |
| Lung | | Low | | Low | | Low | Tumor suppressor | [204,233] |
| Gastric | | | | | | Low | Tumor Suppressor | [207] |

cell type [31]. The first 31 amino acids are believed to be important in the selective targeting of Cav-1 to different subcellular compartments [32]. Cav-1 is expressed at different levels in multiple tissues, with the highest levels found in adipocytes, endothelial cells, fibroblasts, smooth muscle cells, and a variety of epithelial cells [33]. Cav-1 is preferentially targeted to the cytosol of skeletal muscle cells and keratinocytes, to secretory vesicles of endocrine and exocrine cells, to mitochondria of airway epithelial cells and to lipid droplets [34]. Moreover, Cav-1 can also localize away from caveolar regions in the cytoplasm, focal adhesion complexes, the extracellular matrix, and the nucleus, due to its function in a variety of cell signaling activities that may regulate tumor cell behavior [35–37].

Two isoforms of Cav-1 are expressed: Cav-1 α , a 24 kDa protein which contains residues 1–178, and Cav-1 β , a Cav-1 α N-truncated version of a 21 kDa protein which contains residues 32–178. Generation of these isoforms results from either alternative initiation of full-length mRNA or from transcription of a shorter splice variant [38–40]. Both isoforms show distinct but overlapping sub-localization revealing that the N-terminal domain of the protein is not essential for localization in caveolae [40]. *In vitro*, both isoforms were shown to undergo serine phosphorylation. However, only the β -isoform is serine-phosphorylated *in vivo*, suggesting different functions associated with each isoform [41]. Many studies also showed that tyrosine phosphorylation of Cav-1 was dependent on the attachment of the v-Src gene product on the membrane and is associated with different Src tyrosine kinase families. Cav-1 can be phosphorylated by Src at its Tyr14 residue [34,42]. Therefore, only Cav-1 α can be tyrosine-phosphorylated upon Src transformation [43]. More recently, other tyrosine kinases such as c-Abl and Fyn have been shown to phosphorylate Cav-1 at its Tyr14 residue [44,45]. Phosphorylation of Cav-1 at Ser80 converts it into a soluble, secreted protein [46]. It has also been observed that palmitoylation of Cav-1 at Cys133, Cys143 and Cys156 is required for its oligomerization [47,48]. The oligomerization of Cav-1, either with other Cav-1 molecules or with other caveolin proteins, is thought to be

involved in forming the filamentous structure of the caveolar coat. From the hydrophilicity plots of the primary sequence and from mutational analysis, Cav-1 is predicted to have a membrane-spanning hairpin-like structure, with both amino and carboxyl termini oriented towards the cytoplasm [47]. This membrane-spanning model is supported by findings that antibodies directed against either the Cav-1 amino or carboxyl termini require cell permeabilization prior to immunodetection. Furthermore, a sequence of 10 amino acids located within the amino terminal region of Cav-1 is required for the localization of the protein to the rear of migrating cells as well as for caveolar formation [49–54].

Because of the presence of a high number of aromatic residues, the carboxyl terminal portion of Cav-1 CSD (residues 82–101) can interact with numerous signaling molecules localized within caveolae, including Src family tyrosine kinases, growth factor receptors, eNOS, integrins, and G-protein coupled receptors (GPCR) [34,55–58]. In fact, the signaling of proteins such as c-Neu [59], eNOS [60], EGFR [56], estrogen receptor α [61], Id-1 (inhibitor of differentiation/DNA synthesis) [62], TGF- β 1 [63], IGFR1 [64], HGF [65] and others, have been shown to be tightly linked to Cav-1 and to its CSD (Fig. 2). However, recent findings challenge the roles attributed to the caveolin binding motifs in driving direct protein recruitment to the CSD. Therefore, previous work implicating caveolin as a scaffold for direct protein recruitment may need to be re-evaluated to uncover the actual mechanisms by which caveolins modulate specific signaling pathways [66].

4. Caveolin-1 in cancer

Cancer is a multistep process, in which activation of oncogenes or suppression of tumor suppressor genes are phenomena playing critical roles [67]. Cancer is largely characterized by the uncontrolled growth of abnormal cells escaping the immune system, resistance to apoptosis, and the ability to trigger angiogenesis and metastasis. The molecular changes responsible for malignant cell




Fig. 2. Compartmentalization of caveolin and cavin proteins in caveolae. Compartmentalization of caveolin and cavin proteins serves as a platform in caveolae to organize various signalling molecules to effectively transduce extracellular stimuli into intracellular signals through the anchoring of a variety of cancer-related signalling interactors to the caveolin scaffolding domain (CSD). However a recent study does not entirely support this hypothesis as explained in the text [66]. ENOS, endothelial nitric oxide synthase; GPCRs, G-protein coupled receptors; TGF, transforming growth factor; VEGFR, vascular endothelial growth factor receptor; uPAR, urokinase plasminogen activator receptor.

function and tumorigenesis are caused by both genetic mutations and epigenetic mechanisms related to changes in gene expression that are primarily mediated by DNA methylation and histone modifications. Disruption of the epigenome is also a fundamental mechanism in cancer where regulation of tumor development is controlled by a balance between oncogenes and tumor suppressors. Several studies have revealed the importance of Cav-1 during the process of tumor progression and its expression was detected in different types of tumors [68]. In several reports, Cav-1 expression was found to be associated with tumor grade, size and stage, which showed its correlation with the evolution of cancers (Table 2). However, it is evident that Cav-1 also acts as a tumor suppressor [69]. The dual role of Cav-1 was shown to depend on tumor types. In the following sections, we will be discussing the role of caveolin as a tumor promoter in tissue invasion and metastasis, and as a tumor suppressor during apoptosis and inflammation.

5. Caveolin-1 as a tumor suppressor

Over the past 10–15 years, a large amount of data has demonstrated that Cav-1 functions as a tumor suppressor. For instance, expression of Cav-1 in transformed NIH-3T3 cells with different oncogenes such as *rAb1* and *H-Ras* (G12V), resulted in the abrogation of anchorage-independent cell growth [70]. Cav-1 regulates multiple cancer-associated processes including cellular transformation, tumor growth, cell death and survival, multidrug resistance, angiogenesis, cell migration, invasion, apoptosis and metastasis [71]. Cav-1 mRNA and protein levels were down-regulated in oncogene-transformed fibroblasts and re-expression of Cav-1 was sufficient to reverse the transformed phenotype and to prevent anoikis [70,72]. Recently, PI3K/AKT/mTOR was reported to be a suitable target for cancer therapy. Activation of this pathway

has, in fact, been associated with breast cancer progression. Interestingly, Cav-1 deficiency has been shown to promote PI3K/AKT, EGFR-MAPK and MAPK signaling in breast cancer [73], pancreatic cancer [74] and in NIH 3T3 cells [75]. Additionally, the presence of single nucleotide polymorphisms in an allele of Cav-1 T29107A was found to be protective against the development of nasopharyngeal carcinoma, while Cav-1 G14713A was shown to enhance tumor progression [76,77]. These results indicate that the loss of Cav-1 is capable of promoting cell transformation in some cell lines [13,27,78]. Furthermore, reduced expression of Cav-1 is seen in several tumors, including lung [21,79,80], mammary [81,82], colon [52,83], ovarian carcinoma [84], breast [85], sarcomas [86] as well as osteosarcomas [87], while re-expression of Cav-1 again reversed the transformed phenotype. Instead of promoting tumor incidence, Cav-1 deficiency promoted susceptibility to tumor formation in knockout mice, and it may be more suitable to consider Cav-1 as a «tumor susceptibility» factor rather than a tumor suppressor gene. A correlation between Cav-1 and cellular differentiation has also been shown where exposure of pre-adipocytes to differentiation-inducing agents dramatically induced Cav-1 expression at both the transcriptional and translational levels [88].

Human genes encoding Cav-1 and Cav-2 were localized to the q31.1–q31.2 region of chromosome 7 at the D7S522 locus as confirmed by FISH analysis. Interestingly, in a variety of human cancer cell types this locus is frequently deleted [60]. Recombinant expression of the Cav-1 P132L mutant or sporadic mutation in Cav-1 at codon 132 (P132L) induced cellular transformation in NIH 3T3 fibroblasts and breast cancer cell lines, respectively, by acting as a negative dominant of wild-type Cav-1 and by exhibiting a defect in intracellular oligomerization and conformation that, in turn, affects its interactions with cellular partners [89–91]. The Cav-1 promoter region was also found to be hypermethylated in prostate cancer

Table 2
Caveolin-1 expression levels in different types of cancer and different cancer stages.

| Type of cancer | Number of samples | Grade | High expression of Cav-1 (n = 3) | p value | Size (cm) | Expression of Cav-1 | Stage | Expression of Cav-1 | p value | References |
|---------------------------------|-------------------|----------------------|---|---------|------------------------------|--------------------------------------|----------------------------|---|---------|------------|
| Lobular breast | 136 | I II III | 14 (61) 21 (55) 4 (20) | 0.0001 | | | | | | [240] |
| Basal-like breast cancer | 449 | I II III | 6 (84) 14 (140) 39 (225) | 0.026 | | | | | | [163] |
| Salivary gland | 75 | I + II III + IV | 33 (51) 8 (24) | 0.011 | | | | | | [241] |
| Renal cell carcinoma | 114 | I II III | 9 (79) 9 (28) 3 (7) | 0.0008 | <5 5–7 >7 | 4 (31) 2 (29) 15 (53) | | | | [242] |
| Bladder carcinoma | 94 | I II III | 0 (6) 0 (25) 35 (63) | <0.001 | | | Ta T1 T2 T3 T4 | 0 (20) 3 (14) 16 (29) 13 (23) 1 (3) | <0.001 | [160] |
| NSCLC | 115 | I II III IV | 21 (36) 16 (27) 19 (41) 4 (11) | 0.07 | ≤3 ≥3 | 13 (32) 47 (83) | | | | [243] |
| Gastrointestinal stromal tumors | 108 | I II III IV | 1 (7) 1 (16) 0 (27) 4 (58) | 0.334 | <2 >2–≤5 >5–≤10 >10 | 1 (11) 1 (30) 2 (45) 2 (22) | 0.086 | | | [255] |
| Tongue carcinoma | 61 | | | | | | T1 T2 + T3 + T4 | 10 (24) 2 (37) | <0.05 | [245] |
| Upper urinary tract carcinoma | 98 | I II + III | 1 (40) 9 (58) | 0.036 | | | Ta + T1 T2+T4 | 1 (64) 9 (34) | <0.001 | [244] |
| Gastric | 405 | I II + III + IV | 3 (144) 19 (261) | 0.027 | | | | | | [244] |
| Pancreatic cancer | 38 | | | | | | 1 + 2 3 + 4 | 3 (19) 15 (19) | <0.0001 | [246] |

[92] and DNA methyltransferase inhibitor was found to induce Cav-1 expression via C-pG Island (CGI) shore demethylation [93] and to suppress breast cancer. Similarly, polymorphisms in Cav-1 were shown to be involved in colorectal cancer (CRC) and may be used as novel and useful genomic markers for early detection of CRC [94]. In one hospital-based case–control study, the association of Cav-1 polymorphisms with oral cancer risk in a central Taiwanese population was investigated. Cav-1 was concluded to be involved in oral cancer where the A allele of the Cav-1 G14713A was risky, the A allele of the Cav-1 T29107A protective, and AA/TT on these two polymorphisms found to be the most risky combined genotype for the development of oral cancer [76]. Recently, it has been shown that mutations (missense and frameshifts) in the gene encoding Cav-1 altered its tumor suppressive functions in breast cancer and found to promote mammary tumorigenesis in a Kashmiri population [95]. Therefore, both expression levels and functional mutations in Cav-1 contribute to the promotion of cancer.

6. Caveolin-1 functions as a tumor promoter

Despite the evidence favoring the tumor suppressive role of Cav-1, several studies also support an opposite role for Cav-1 as a protein that actually promotes tumor growth. The expression of Cav-1 was shown to be increased upon tumor formation in mouse models and human patients [21,96], and thus its presence promoted tumor growth and metastasis of cancer cells via an autocrine/paracrine mechanism [97–99]. Post-translational modifications are also believed to play a role in the function of Cav-1 as a tumor promoter gene such as the phosphorylation of Cav-1 on Tyr14 by Src [100]. These results were confirmed by studies showing that Cav-1 downregulation converted androgen-insensitive metastatic mouse prostate cancer cells into an androgen-sensitive phenotype [101], and could independently promote prostate cancer cell survival and metastatic activities [98].

Similarly, clinicopathological analyses of esophageal [102], breast [103], renal [104], brain [105], lung [23,106], ependymoma [107], and prostate [108] cancers revealed a role for Cav-1 as a tumor promoter. In prostate cancer patients, the presence of Cav-1 was found to be associated with angiogenesis, cancer recurrence, elevated metastasis and poor patient prognosis [68,109–111]. Likewise, upregulation of Cav-1 in MDR colon cancer cells led to a reduced rate of proliferation and thus contributed to their selective growth advantage under chemotherapy [112]. Similarly, Cav-1 expression in MCF-7 human breast cancer cells prevented anoikis [113,114]. Cav-1 also promoted chemoresistance in gastric cancer [115] and in lymph node metastasis of non-small cell lung cancer [116]. Overexpression of Cav-1 has also been associated with metastasis and poor patient prognosis in breast carcinomas [117] and in oesophageal squamous cell carcinoma [94]. Interestingly, Cav-1 upregulates glucose uptake and ATP production through HMGA1-mediated GLUT3 transcription, suggesting that Cav-1 may increase tumor cell growth by enhancing aerobic glycolysis in colorectal cancers [118,119]. In summary, there are equally convincing data which show a dual role for Cav-1 as both tumor promoter and tumor suppressor. Perhaps one way to resolve this discrepancy is by defining Cav-1 as a « conditional tumor suppressor » protein. Along these lines, a recent study demonstrated that Cav-1 expression was increased by an erythroblastosis virus transforming protein (ETS) such as the polyoma-virus enhancer activator (PEA3) in mice epithelial lung cells [120]. It was suggested that PEA3, a pro-metastatic ETS protein involved in breast cancer, was able to increase Cav-1 expression in metastatic lymph node sites [80]. In contrast, the ETS repressor factor Net suppressed Cav-1 transcription in primary lung tumors and PEA3 was found to activate Cav-1 transcription in metastatic lymph

nodes in non-small cell lung carcinoma [120]. Similarly, expression of Mgat5/galectin-3 promoted the tumor-enhancing role of Cav-1, and these conditions were associated with poor prognosis in several tumor types [121]. Although we do not know to what extent the absence of Mgat5/galectin-3 regulates other functions of Cav-1, damage to Mgat5/galectin-3 is frequently associated with diminished growth and metastasis of tumor cells, and thus represents an interesting candidate protein for tumor inhibition.

7. Caveolin-1 in invasion and metastasis

As previously mentioned, Cav-1 was first identified as the major tyrosine-phosphorylated protein in Rous sarcoma virus-transformed chicken embryo fibroblasts [30]. It is known that Rho signaling, via Rho kinase (ROCK), is closely associated with stem cells and tumor cell migration and metastasis [122]. Tyrosine-phosphorylated Cav-1 functions as an effector of Rho/ROCK signaling in the regulation of focal adhesion lifetime [100] and, thereby, in tumor cell migration and invasion. This defines a feedback loop between Rho/ROCK, Src, and phosphorylated Cav-1 in tumor cell protrusions, identifying a novel function for Cav-1 in tumor metastasis that may contribute to the poor prognosis associated with some Cav-1-expressing tumors [123]. Moreover, the interaction between Cav-1 and Rho-GTPases (most likely RhoC, but not RhoA) promoted metastasis by stimulating $\alpha 5$ -integrin expression and by regulating the Src-dependent activation of the p130 (Cas)/Rac1, FAK/Pyk2 and Ras/Erk1/2 signaling cascades [124]. Furthermore, the association of Cav-1 with invasion, survival, and poor prognosis in hepatocellular cancer was associated with an increased expression of matrix metalloproteinases (MMP) [125].

Export of Cav-1 to secretory vesicles has also been found to be functionally associated with melanoma progression [126]. Consistently, fibroblast expression of Cav-1, through p190RhoGAP, favors directional migration and invasiveness of carcinoma cells *in vitro*. *In vivo*, stromal Cav-1 remodels peritumoral and intratumoral microenvironments to facilitate tumor invasion, correlating with increased metastatic potency. Thus, biomechanical remodeling of the microenvironment by stromal Cav-1 favors tumor invasion and metastasis [127]. In contrast, Cav-1 can inhibit pancreatic carcinoma cell invasion via the Erk-MMP signaling pathway, suggesting that the endogenous expression or re-expression of Cav-1 might help reduce pancreatic carcinoma cells invasive potential [128]. Cav-1 interaction with integrins has also been shown to regulate focal adhesion turnover [19,129,130], and Cav-1 inhibited metastatic potential in melanomas through suppression of the integrin/Src/FAK signaling pathway [131].

8. Caveolin-1 in apoptosis

Cav-1 is believed to regulate apoptosis in different cell lines. Given its dual role as a tumor suppressor or as a tumor promoter gene, it can also regulate either anti- or pro-apoptotic functions. Cav-1 was further demonstrated to promote cell-cycle arrest through a p53/p21-dependent pathway where it was demonstrated that its overexpression induced the cells to exit the S phase and to increase the G0/G1 cell population in mouse embryonic fibroblasts derived from Cav-1 transgenic mice [132].

In cancer therapies, most of the agents used promote apoptosis and result in decreased tumor masses. A defective apoptotic machinery would therefore lead to the failure of treatment, which is commonly observed in most types of cancers these days. Several studies demonstrated an enhanced sensitivity to chemically-induced apoptosis when Cav-1 is expressed. For instance, Cav-1 enhanced resveratrol-mediated transport and cytotoxicity in a hepatocellular carcinoma model [133] and paclitaxel-mediated

apoptosis in MCF-7 breast cancer cells which is regulated via Tyr14 phosphorylation of Cav-1 by inactivating Bcl-2 and increasing mitochondrial permeability [134]. Interaction of multidrug resistance protein with Cav-1 was shown to be increased in bleomycin-induced cell cycle arrest and subsequent cellular senescence [135]. Exposure to sub-toxic concentrations of cisplatin also induced hydrogen peroxide generation, and the subsequent increase of reactive oxygen species (ROS) further up-regulated Cav-1 levels and anoikis resistance which is crucial to the survival of metastatic cancer cells [136,137].

Aside from its pro-apoptotic functions, there is also strong evidence which indicates Cav-1 plays a role as an anti-apoptotic protein. For example, in the case of colon cancer, it was shown that Ku70, a protein identified as part of the Ku70/Ku80 complex that sequesters the pro-apoptotic gene Bax and mediates DNA repair, bound to Cav-1 and led to the inhibition of apoptosis mediated by chemotherapy drugs [138]. Similarly, an inhibition of apoptosis induced by the tumor necrosis factor-related apoptosis-inducing ligand TRAIL was also due to Cav-1 expression in hepatocarcinoma cells [139]. Increased expression of p-Cav-1 was shown to promote cell survival after oxidative stress and to play a role as an anti-apoptotic protein [137]. Cav-1 was shown to play an active role in mediating the transformation and survival of mouse hepatoma cells, and as such might represent a potential target for gene and anti-tumor drug treatment [140]. Furthermore, Cav-1 was demonstrated to inhibit cell detachment-induced p53 activation and anoikis by upregulating insulin-like growth factor-1 receptor's signaling and transcription in breast cancer cells [141,142], to regulate hyperoxia/ROS-induced apoptosis through interactions with Fas and BID [143], and to alter TrkA modification including Tyr490 phosphorylation [144]. These observations suggest that Cav-1 could suppress the TrkA-mediated phenotypic effects by altering TrkA modification via functional interaction. Similarly, RNA interference-directed Cav-1 knockdown sensitized renal carcinoma cells to doxorubicin-induced apoptosis and reduced lung metastasis [145]. In conclusion, the functional role of Cav-1 in apoptosis appears to be linked to tumor type as well as to the nature and the potency of the apoptosis inducers which are used to trigger death signalling.

9. Caveolin-2

Caveolin-2 (Cav-2) was discovered following the micro-sequencing of a 20 kDa protein that co-purified with adipocyte-derived caveolar membranes [146]. Three isoforms, alpha, beta and gamma, exist for Cav-2 which is located on human chromosome 7q31.1, a locus often deleted in breast and prostate cancers [147]. Cav-2 can form stable hetero-oligomeric complexes of 14–16 molecules and can coalesce into large macromolecular complexes that define caveolar architecture at the plasma membrane [148,149]. Similar to Cav-1, Cav-2 is expressed in white adipose tissue, fibroblasts, and epithelial cells, and can be found expressed alone in lungs [149,150]. Caveolar formation can still occur even when Cav-2 is not expressed suggesting a role for Cav-2 as a partner binding to Cav-1 [33]. Otherwise trapped in the Golgi apparatus [151], Cav-2 transport to the plasma membrane is dependent upon Cav-1 expression [152]. Cav-2 differs from Cav-1 in functional domains including a G-protein binding domain and a CSD located at its N-terminal region [153,154]. Additionally, the Cav-1 N-terminal amino acids Lys47–Lys57 are highly conserved in Cav-2, and disruption of this region impedes cell polarity and directional migration [17]. Furthermore, the CSD of Cav-2 is responsible for its localization in the Golgi apparatus of polarized cells derived from colon cancer [155]. Cav-2 also possesses the capacity to modulate several signaling pathways as it has been shown that insulin triggered Tyr19 phosphorylation of Cav-2 which increased the

interaction between phospho-Tyr19 and phospho-ERK, indicating that this phosphorylation is required for actin cytoskeleton-dependent ERK nuclear import [156]. Cav-2 expression is also involved in pulmonary defects, lipid metabolism and cancer formation [150]. It has been shown that Cav-2 expression is more frequent in anaplastic carcinoma, papillary thyroid carcinoma and diffuse sclerosing variant of papillary carcinoma. This evidence confirms a role for Cav-2 in the modulation of cancer progression [157,158].

10. Caveolin-2 in cancer

Cav-2 expression was shown not to be altered in oncogenic transformation. Along this evidence, dual roles for Cav-2 have been documented in various cancer cells, as Cav-2 was neither expressed in HepG2 hepatocellular carcinoma cells, nor in SH-SY5Y neuroblastoma cells [158]. However, an increase in Cav-2 expression was documented in esophageal and urothelial carcinomas [159,160], as well as in C6 glioma, HeLa epithelial cervical cancer, A549 lung adenocarcinoma, and MCF7 breast cancer cells [161]. Cav-2 transfection of HepG2 hepatocellular carcinoma cells and Cav-2 knockdown in C6 glioma cells caused reductions in cell proliferation and growth, while Cav-2 re-expression in SH-SY5Y neuroblastoma, and depletion in HeLa epithelial cervical cancer and A549 lung adenocarcinoma cells promoted cancer cell proliferation. Collectively, these data suggest that Cav-2 acts as a modulator of cancer progression. Interestingly, Cav-2 was shown to negatively regulate lung endothelial cell proliferation and cell cycle progression [162], and to be associated with breast cancer basal-like and triple-negative immunophenotypes [163,164]. Cav-2 expression was up-regulated during the late stages of C6 glioma cell differentiation while Cav-3 was gradually down-regulated during this differentiation process [165]. In the case of renal cell carcinoma (RCC), Cav-2 was shown to be a direct target of the specific miRNA miE-218 which is considered to act as a tumor suppressor in RCC. Silencing of Cav-2 mRNA and protein produced a high degree of inhibition of cellular proliferation and migration in RCC [166]. These data suggest that, like Cav-1, Cav-2 could be considered as a tumor suppressor or tumor promoter gene.

11. Caveolin-3

Caveolin-3 (Cav-3), the third member of the caveolin family, is more restricted in distribution than are Cav-1 and Cav-2, and has been particularly studied in vascular smooth muscle, cardiac and skeletal muscles [167], in glial cells [165], and in myoepithelial cells within the mammary gland [164]. Cav-3 was identified through database searches and through classical cDNA library screening in an attempt to find Cav-1 homologs. Human Cav-3 proteins contain 151 amino acids and contain specific, separate domains: A N-terminal domain (aa 1–53) which contains a signature sequence (aa 41–48) that is present in all caveolins, a scaffolding segment (aa 54–73) responsible for homo-oligomerization, a transmembrane domain (aa 74–106) and a C-terminal (aa 107–151) domain. The Cav-3 oligomerization process is believed to be initiated in the endoplasmic reticulum and presages the organization of detergent-resistant caveolar complexes of approximately 25–50 nm in diameter. These structures then fuse with the plasma membrane forming the final caveolae. The transmembrane domain forms a hairpin loop in the sarcolemma, allowing both the N- and C-terminal ends to face the cytoplasm. Cav-3 has a small ubiquitin-like modifier (SUMO) consensus motif (ΨKXD/E, where Ψ is a hydrophobic residue) near the scaffolding domain, allowing SUMOylation in a manner that is enhanced by the SUMO E3 ligase PIASy (protein inhibitor of activated STAT-y). Thus, SUMOylation is a

covalent modification of caveolins that influences the regulation of signaling partners [168]. Interestingly, Cav-3 also forms large oligomeric complexes of approximately 350–400 kDa *in vivo*. Cav-3-generated caveolae have been shown to compartmentalize, and to modulate a number of signaling proteins including eNOS, β -adrenergic receptors, protein kinase C isoforms, G proteins, Src-family kinases, multiple components of the dystrophin-glycoprotein complex [169,170], and natriuretic peptide signaling [171]. Additionally, it has been shown that point mutations in Cav-3 (P140L) can attenuate p38, AKT [172] and endoplasmic reticulum stress signaling [173].

In Cav-1-deficient knockout mice, caveolae were only detectable in skeletal and heart muscle cells [174,175], whereas no caveolar formation is observed in skeletal and heart muscle from Cav-3 knockout mice [176]. Tissues from Cav-1/Cav-3 double knockout mice are completely devoid of caveolae [177]. Together with a large body of data in the literature, these results validate the concept that either Cav-1 or Cav-3 is required to form caveolae. Cav-2, therefore, appears less important in this respect although the presence of this isoform does modulate morphological traits and efficacy in caveolar formation. The role of Cav-3 in cancer is still not well established and needs further investigation. In thyroid cancer, Cav-3 was highly expressed in the stroma of anaplastic carcinoma (AC) compared to papillary thyroid carcinoma (PTC) where Cav-1 and Cav-2 were highly expressed [158].

12. The cavins

Evidence for the presence of another protein family responsible for caveolae formation support the role of cavin proteins [178]. These cavins are composed of four types which include Cavin-1 (Polymerase 1 and Transcript Release Factor, PTRF), Cavin-2 (Serum-Deprivation Response Protein, SDPR), Cavin-3 (SDR related gene product that binds to c-kinase, SRBC), and Cavin-4 (Muscle-Restricted Coiled-Coil Protein, MURC), all of which share a role with Cavin-1 in the regulation of caveolae formation.

13. Cavin-1/PTRF

Cavin-1 (PTRF, Cav-p60) was identified by a yeast two hybrid screen as the first caveolin regulatory protein [179]. Originally believed to be a transcript release factor, Cavin-1 is a soluble protein with a putative leucine zipper, nuclear localization sequences and PEST (Proline, Glutamic acid, Serine and Threonine) domains. Studies have shown its colocalization and codistribution with Cav-1 in adipose tissue and in lipid rafts. In particular, Cavin-1 was recently shown to be an abundant caveolar coat protein required for caveolae stabilization and probably for interaction with the cytoskeleton [5,180,181]. Cavin-1 was shown to colocalize with Cav-1 at the plasma membrane but not in the Golgi apparatus [5]. It is recruited by Cav-1 and Cav-3 to plasma membrane domains where it binds to phosphatidylserine, cholesterol and oligomerized caveolins [182]. Binding of Cavin-1 to these domains stabilizes the membrane curvature as evidenced by the flask shape of caveolae. The interaction of Cavin-1 with Cav-1 is not direct, but is possibly mediated through cytoskeletal interactions between microtubules and actin filaments [5]. Live cell imaging demonstrated that the lateral mobility and lysosomal degradation of Cav-1 is enhanced by the loss of Cavin-1 [183]. Therefore, Cavin-1 is a potential caveolar coat protein that regulates caveolar structure and caveolin function.

14. Cavin-2/SDPR

Cavin-2 (SDPR) was purified as a phosphatidylserine-binding protein from human platelets [182] with greater expression in a

serum-deprived microenvironment, giving rise to the name of serum deprivation protein [184]. Cavin-2 was also initially shown to potentially bind PKC α and to localize within caveolae [185]. It is now well recognized that biogenesis of caveolae requires both caveolin and Cavin-1 proteins although the identity, function and molecular interaction of other caveolar components remain unresolved. It was demonstrated that Cavin-2 recruited Cavin-1 to the plasma membrane, and was required for stable expression levels of both Cav-1 and Cavin-1 proteins [186]. This interaction leads to the formation of caveolin-containing complexes and to the stabilization of caveolar structures. Interestingly, overexpression of Cavin-2 results in the formation of elongated tubular caveolae, which implies that, unlike Cav-1 or Cavin-1, Cavin-2 does not increase caveolar number but rather induces changes in caveolar morphology [187]. The domains of Cavin-2 responsible for the formation of caveolae curvature comprise a potential helical coiled-coil forming region followed by another conserved domain containing multiple basic amino acids [186]. The membrane tubulations closely resemble those induced by the β -subunit of Shiga toxin (STB). It has been demonstrated that STB induces the formation of membrane tubes even in ATP-depleted cells [188], but whether such tubes arise because of the interaction of the toxin with membrane lipids, or whether additional membrane or cytosolic proteins are required remains unclear. Cavin-2-induced tubes originate from caveolae, and incorporate STB. Loss of Cavin-1, Cavin-2 or Cav-1 reduces the propensity of STB to induce membrane tubulation. Thus, formation of caveolae is not solely dependent on oligomerization of caveolin proteins, but also requires the involvement of other factors [189]. As there is no obvious sequence similarity between Cavin-2 and known curvature-inducing proteins, it is hypothesized that Cavin-2 may induce curvature by a novel mechanism [190]. Further insights into this will necessitate structural information.

15. Cavin-3/SRBC

Cavin-3 (SDR related gene product that binds to c-kinase, SRBC) localizes within caveolae [148] and was initially identified as a phosphatidylserine-binding protein and as a substrate of PKC [191]. Similarly to Cavin-2, its expression was also shown to be induced upon serum deprivation [191,192]. Binding of Cavin-3 to PKC δ depends on the presence of phosphatidylserine, which stabilizes its interaction with PKC δ through bridging. Cavin-3 also binds to PKC α to the same extent as to PKC δ , but it barely binds to PKC ζ . Immunofluorescence-based evidence shows that all of the cavin proteins are required for the formation of caveolae, and that an individual caveola can contain more than one member of the cavin family [148]. The ability of these proteins to concentrate in caveolae is believed to be directly or indirectly dependent on the expression of Cav-1. Therefore, all cavins appear to have the properties of a caveolin adaptor molecule that links signal transduction to the regulation of caveolar function. Furthermore, Cavin-3 leads to the formation of caveolar vesicles as it remains associated with caveolae when budding occurs, and its absence impairs intracellular vesicular trafficking [193].

16. Cavin-4/MURC

It has been demonstrated that Cavin-4 (Muscle-Restricted Coiled-Coil Protein, MURC), the fourth member of the cavin family, is an evolutionarily-conserved, muscle-specific component of the cavin complex [194–196] which is associated with the sarcolemmal caveolae complexes. Cavin-4 is also present at low levels in other cell types, such as embryonic fibroblasts [197], and is able to interact with Cavin-2 [194]. Cavin-4 influences skeletal muscle differentiation by activating ERK and RhoA through the Rho-ROCK

pathway, which also modulates cardiac function. Overexpression of Cavin-4 in skeletal and cardiac muscle promotes myogenesis and causes cardiac dysfunction and conduction disturbance. These facts raise the possibility that mutations in Cavin-4 may directly cause muscle disease [197]. Therefore, Cavin-4 is considered to be a new potential candidate for muscle-related caveolinopathies.

17. Cavins in cancer

The role of cavins in caveolae formation is now well recognized. Recently, it has been proposed that Cavin-1 stabilizes Cav-1 expression and promotes its role as a tumor promoter in pancreatic cancer cells [197]. Gene silencing of Cavin-1 decreased cellular invasiveness and metastasis suggesting Cavin-1 as a potential therapeutic target [197]. In contrast, PC3 pancreatic cancer cells that expressed Cavin-1 showed a decrease in invasiveness and migration which correlated with reduced production of the matrix metalloproteinase MMP-9 [198], whose lowered expression associated with reduced angiogenesis and inflammation [199,200]. Cavin-1 and, to a lesser extent, cavins-2, -3, and -4, also decreased MMP-9. Genetic ablation of Cavin-1 also resulted in the loss of caveolae although Cav-1 expression remained albeit at a reduced level [186,201]. Taken together, these observations suggest that Cavin-1 expression alters prostate and pancreatic cancer aggressiveness via two different roles either as tumor suppressor or as tumor promoter respectively. Furthermore, it has been demonstrated that changes occur in cell membranes involving the loss of caveolae and of Cavin-1 expression concurrent with the development of prostate cancer. These changes were also accompanied by an up-regulation of Cav-2 [202]. Cavin-1 and Cavin-2 were shown to be expressed in MDR cell lines, and Cavin-1 was demonstrated to be involved in drug resistance by rendering lipid rafts more rigid [203]. In another study, Cavin-1 was confirmed as a non-small cell lung cancer biomarker by label-free proteomics and was shown to have a relationship with the EGFR pathway [204]. Similarly, Cav-1-induced tubule formation in breast cancer cells was suppressed by the presence of Cavin-1 [205]. Cavin-1 also acts as a novel regulator of oxidative stress-induced premature senescence by linking free radicals to activation of the p53/p21Waf1/Cip1-pathway [206].

In contrast to Cavin-1 and Cavin-2, Cavin-3's function and role in caveolae formation are not well established. The human Cavin-3 gene maps to 11q15.5–15.4 where it is located in a tumor suppressor region and is inactivated in breast and lung cancers [207]. The loss of Cavin-3 in lung cancer appears to be a consequence of DNA methylation and subsequent gene silencing [208], although other studies have suggested that alternate mechanisms could be involved. Targeting of Cavin-3 to caveolae depends on both its LZ domain and the expression of Cav-1. A recent study showed that Cavin-1 and Cavin-3 were down-regulated in breast cancer cell lines and breast tumor tissue [209]. However, given that cavins are required in conjunction with caveolins for caveolae formation, their role in tumor regulation will still require more investigation.

18. Mechanisms involved in the dual role of caveolins and cavins in cancer

Three distinct mechanisms, i.e. tyrosine phosphorylation, serine phosphorylation and P132L dominant negative point mutation, have been shown capable of counteracting the growth inhibitory function of Cav-1. These findings may, in part, explain the dual role of Cav-1; they depend on the tumor type/stage. Cav-1 undergoes tyrosine phosphorylation at its NH₂-terminus and then recruits several SH2-domain containing proteins, such as Grb-7. This, in turn, can increase either the anchorage-independent growth of cells or decrease anoikis [210]. Cav-2 tyrosine phosphorylation at

Tyr19 and Tyr27 enables the recruitment of other SH2-domain containing proteins, such as c-Src, Nck, and Ras-GAP. Likewise, Cav-1 serine phosphorylation at Ser80 can serve to convert Cav-1 into a soluble, secreted protein [211]. The secreted form of Cav-1 apparently acts as a growth factor and can enable cells to defend against apoptotic stimuli. It has also been shown that tumor cells which secreted Cav-1 triggered proangiogenic activities in prostate cancer [212]. These are the possible mechanisms by which Cav-1 functions as a growth stimulator in prostate carcinomas and perhaps also in other cancers.

The caveolin scaffolding domain (CSD; residues 82–101) is also believed to inhibit the growth stimulatory activity of several signaling molecules, including receptor tyrosine kinases, Src-family kinases, and members of the Ras-p42/44 MAP kinase cascade. In this regard, the CSD may function as an endogenous kinase inhibitor by recognizing a conserved aromatic amino acid motif that is present within the kinase domain of most known kinases. However, it can also inhibit other signaling molecules, such as eNOS, by recognizing the same aromatic motif within the NOS catalytic domain [57,213]. However, only a limited number of studies have attempted to directly test binding of the CSD peptide to signaling proteins, and in these cases binding was not investigated in the context of an interaction with alleged caveolin binding motifs. Thus, the mechanisms involved in the inhibition of signaling by CSD and, more generally, in the effects of caveolin and/or caveolar loss of function on specific signaling pathways will necessitate reinvestigation, and potentially argues against a role for caveolin binding motifs in driving direct protein recruitment to caveolae via CSD [66].

Caveolin-related cancer research is still finding novel roles for the caveolins. In a recent study, Cav-1 was shown to be important for the endocytosis of E-cadherin and its downregulation resulted in decreased E-cadherin expression, increased β -catenin transcriptional activation, and increased invasiveness in neoplastic cells [214]. Furthermore, loss of Cav-1 function in the tumor microenvironment contributes to the metastatic behavior of prostate tumor cells by a mechanism that involves up-regulation of TGF- β 1 and SNCG through Akt activation [215]. Similarly, in follicular thyroid carcinoma, reduced expression of Cav-1 down-regulated PTEN and tumor suppressive functions. Given that PTEN is required for activation of the phospho-Akt apoptotic pathway, down-regulation of its activity consequently leads to increased cell survival and proliferation [147]. In contrast, knockdown of Cav-1 in hepatocellular carcinoma cells strongly reduced TGF- β -mediated AKT phosphorylation, thus sensitizing primary murine hepatocytes for pro-apoptotic TGF- β signaling [216]. Gene expression microarray analyses in glioma cells demonstrated significant enrichment in genes corresponding to downregulation of MAPK, PI3K/AKT and mTOR signaling [217], as well as activation of apoptotic pathways in Cav-1-overexpressing U87-MG cells, as well as upregulation of MAPK/AP1 signaling in Cav-1 knocked-down cutaneous squamous cell carcinoma [218], supporting Cav-1's tumor suppressive role.

Several research groups have shown localization of MT1-MMP and MMP-2 to caveolae and depicted caveolae as being required for proper MT1-MMP localization and function in tumor development [36,219]. Furthermore, recombinant expression of Cav-1 in mammary epithelial cells was reported to suppress their metastatic capacity, to inhibit their invasiveness, and to prevent their ability to secrete MMP-2 and MMP-9. Given the heterogeneity of Cav-1 expression in various tumors, therapies targeting Cav-1 will need to consider the specific roles of this protein within a given specific type of tumor. In one exciting gene therapy strategy, the Cav-1 promoter was used to specifically target prostate cancer cells *in vitro* and *in vivo* with minimal toxicity [220]. Another type of anti-tumor therapy strategy was used to show that Cav-1 was

highly expressed in the vascular endothelium, suggesting that targeting a tumor's blood supply may provide an interesting opportunity for Cav-1-based directed therapies [221].

Cavin-1/PTRF is purported to be an essential component of caveolae, as it has been shown that recruitment of Cavin-1 to membrane domains, enriched in phosphatidylserine, cholesterol and oligomerized caveolins, was responsible for the stabilization of membrane curvature enabling the classic flask shape of caveolae. Loss of Cavin-1 releases caveolar components, including caveolins into the plasma membrane and the level of non-caveolar caveolins is thought to be tightly regulated via endolysosomal degradation [5]. In a recent report, it has been shown that Cavin-1 and Cav-1 can bind IGF-IR and regulate IGF-IR internalization and plasma membrane replacement through mechanisms frequently dysregulated in cancer cells, although the exact roles of Cav-1 and IGF-IR's in human cancer continue to be debated. With the discovery of IGF-IR interaction with Cavin-1 in caveolae, new insight has emerged in understanding the functions of these domains in IGF-I action [222]. Similarly, in metastatic PC3 prostate cancer cells that express abundant Cav-1 but no Cavin-1, heterologous Cavin-1 expression restored caveolae formation and Cav-1 distribution [5]. Recently, it has been shown that Cavin-1-expressing PC3 cells exhibited decreased migration, and that this effect was mediated by reduced MMP-9 production. Cavin-1, and to a lesser extent, Cavin-2, -3, and -4, all decreased MMP-9, suggesting that MMP-9 production may not solely depend on caveolae formation. Taken together, these observations suggest that Cavin-1 expression can inhibit prostate cancer aggressiveness by reducing MMP-9-mediated functions [201]. Interestingly, it was found that Cavin-1 was not expressed in prostate cancer epithelium [202,223,224]. In agreement with a protective role for Cavin-1 in prostate cancer, its down regulation in DU145 cells enhanced cell 3D migration [225]. Fhl1 induced the expression of Cavin-1 in Src-transformed cells, independent of MAPK activity, while Fhl1 and Cavin-1 expression were found to be suppressed in breast, kidney, and prostate tumors [226].

Several lines of evidence suggest that Cavin-3 may function as a tumor suppressor, but the molecular basis of its action remains unknown [227]. Cavin-3 was originally identified in screens looking for PKC δ -binding proteins, and was found to be phosphorylated *in vivo* by PKC δ , a potential tumor suppressor involved in the regulation of cell proliferation, differentiation, and apoptosis [191]. The mRNA for *Cavin-3* is induced in response to serum deprivation and is downregulated during G₀/G₁ transition, suggesting that it

may be involved in cell-cycle control [191]. Through a yeast 2-hybrid screen, Cavin-3 was also identified as a BRCA1-interacting protein, raising the possibility that it may participate in DNA-damage response, and that its inactivation may compromise BRCA1-mediated tumor suppression functions [228]. Cavin-3 was also shown to increase p53 protein stability, and its proapoptotic effect stems partially from the p53-enhancing activity suggesting that dysregulation of Cavin-3 may attenuate p53's response to stresses and thus contribute to malignant tumor progression [229].

Cavin-3 is a proapoptotic tumor suppressor which is commonly altered in colorectal cancer by promoter hypermethylation, and whose gene transcription is directly activated by NF- κ B in response to TNF α . This suggests that Cavin-3 inactivation may contribute to tumor progression by reducing cellular sensitivity to TNF α and to other stresses such as those due to a chronic inflammatory microenvironment [227]. It was further inferred that Cavin-3 induced G₁ arrest of the cell cycle partially through CDKN1A induction and that this increased the apoptotic response of tumor cells to various stresses including etoposide, 5-FU, γ -irradiation, H₂O₂, and serum deprivation. Consistent with these effects, Cavin-3 significantly decreased the colony-forming ability of tumor cells and delayed the formation and growth of xenograft tumors. It was also found that *Cavin-3* mRNA was strongly elevated in response to genotoxic or nongenotoxic stimuli, raising the possibility that it was controlled by stress signaling and involved in damage response. Moreover, Cavin-3 suppressed both basal and IGF-induced phosphorylation of AKT, without having any detectable effect on ERK and JNK [227]. It was further demonstrated that the p53-enhancing function of Cavin-3 might be associated with its regulatory role on AKT [230]. Therefore, these findings suggest that Cavin-3-mediated tumor suppression might be, in part, attributed to its ability to inhibit the PI3K-AKT signaling pathway which plays a crucial role in the development and progression of a variety of human tumors. Despite several lines of evidence for Cavin-3's tumor suppressive role, the molecular mechanisms underlying these actions in tumorigenesis are not fully understood (Table 3).

19. Concluding remarks

Caveolae formation and function is a crucial phenomenon in several normal or pathological events. Caveolae formation is dependent on the expression and interaction of two protein families, namely caveolins and cavins and current research has clearly

Table 3

The promoting and suppressive roles of cavins 1–3 in different types of cancer.

| Tumor | Protein involved | Mechanisms involved | Role in cancer | References |
|-----------------------------------|------------------------|--|------------------|------------|
| Prostate cancer | Stromal Cav-1 | TGF- β , Akt, γ -synuclein (SNCG) | Tumor Suppressor | [215] |
| Melanoma | Cav-1 | Shh heterotypic | Tumor Suppressor | [254] |
| Thyroid cancer | Cav-1 | PTEN, Akt | Tumor Suppressor | [147] |
| Mammary epithelial cells | Cav-1 | p38 MAPK/JNK MAPK activation via TLR4 | Tumor Suppressor | [247] |
| Glioblastoma | Cav-1 | Inactivation of MAPK & PI3K/mTOR | Tumor Suppressor | [217] |
| Cutaneous squamous cell carcinoma | Cav-1 | MAPK/AP1 | Tumor Suppressor | [218] |
| Mammary cancer cells | Cav-1 | Integrin/EGF | Tumor Promoter | [248] |
| Hepatocellular cancer | Cav-1 | TGF- β | Tumor Promoter | [216] |
| Prostate cancer | Cav-1 | IGF-IR/IR | Tumor Promoter | [220] |
| Colon cancer | Cav-1 | AMPK-TP53/p53 | Tumor Promoter | [36] |
| Breast cancer | Cav-2 | MIR-199a-3p | Tumor Suppressor | [250] |
| Osteosarcoma | Cav-2 | NFBD1/Erk | Tumor Suppressor | [251] |
| Renal cell carcinoma | Cav-2 | miR-218 | Tumor Promoter | [166] |
| Mammary tumor | Cav-3 | Elf5, Stat5a, c-Myc | Tumor Promoter | [238] |
| Glioma | Cav-1 & Cav-2 | VEGF | Tumor Promoter | [252] |
| Colon carcinoma | Cav-1 & Cav-2 | PPAR- γ | Tumor Promoter | [253] |
| Prostate cancer | Cavin-1 | MMP-9 | Tumor Suppressor | [223] |
| Colorectal cancer | Cavin-3 | NF- κ B, TNF- α | Tumor Suppressor | [249] |
| Gastric cancer | Stromal Cav-1, Cavin-3 | p53, p21, PUMA, OXA | Tumor Suppressor | [255,207] |

established a role for caveolins and cavinins as “tumor and metastasis-modifying genes”. Understanding the cross-interactions of these proteins, their structure and their function will in turn give insight into their involvement in diseases such as cancer where the formation or loss of caveolae modulators affects oncogenic functions. Cav-1 and Cav-2 were attributed dual roles in cancer, either as tumor suppressors or tumor promoters depending on the type and stage of cancer. Cav-3, expressed strictly in muscle, has also been documented as an essential protein for caveolae formation but its exact role in cancer remains to be defined. In some types of cancer such as breast cancer, overexpression of Cav-1 inhibits cellular metastasis, cellular adhesion, proliferation and migration whereas in other cancers, Cav-1 was reported to induce these phenomena. Cavin proteins were discovered after caveolin proteins and were also found to be essential components in caveolae formation. Four family members were identified and are thought to interact with caveolin in order to stabilize caveolae. More studies are needed to understand the roles and implications of cavinins in cancer although several studies have revealed that expression of Cavin-1 and of Cavin-2 are downregulated in breast cancer cells and tissues. On the other hand, overexpression of cavinins in prostate cancer has been shown to be associated with reduced expression of MMPs, which highlights its potential role in tumor suppression. Current research has clearly established roles for caveolins and cavinins as «tumor- and metastasis-modifying genes» and, due to the heterogeneity of caveolin and cavin expression in different tumors, these proteins could therefore be targeted in cancer therapies. Future studies will undoubtedly reveal novel relationships between these proteins and their effects on a variety of signaling pathways. This will offer exciting opportunities to develop anti-cancer therapies that target them, both in primary tumors and in metastatic disease.

Authors' contributions

RG, CT and BA contributed to manuscript conception and writing. RG contributed to literature search and manuscript writing. CT and BA contributed to manuscript writing and critically revised the paper. All authors read and approved the final manuscript.

Conflict of interest

All authors declare no conflict of interest on the topics covered by this review.

Acknowledgments

B.A. holds an institutional Chair in Cancer Prevention and Treatment at UQAM. This study was funded by a grant from the Natural Sciences and Engineering Research Council of Canada (NSERC) (288249) to B.A.

References

- [1] G.E. Palade, Fine structure of blood capillaries, *J. Appl. Phys.* 24 (1953) 1424–1436.
- [2] T. Richter, M. Floetenmeyer, C. Ferguson, J. Galea, J. Goh, M.R. Lindsay, G.P. Morgan, B.J. Marsh, R.G. Parton, High-resolution 3D quantitative analysis of caveolar ultrastructure and caveola-cytoskeleton interactions, *Traffic* 9 (2008) 893–909.
- [3] W. Schlormann, F. Steiniger, W. Richter, R. Kaufmann, G. Hause, C. Lemke, M. Westermann, The shape of caveolae is omega-like after glutaraldehyde fixation and cup-like after cryofixation, *Histochem. Cell. Biol.* 133 (2010) 223–228.
- [4] L. Pelkmans, M. Zerial, Kinase-regulated quantal assemblies and kiss-and-run recycling of caveolae, *Nature* 436 (2005) 128–133.
- [5] M.M. Hill, M. Bastiani, R. Luetterforst, M. Kirkham, A. Kirkham, S.J. Nixon, P. Walser, D. Abankwa, V.M. Oorschot, S. Martin, J.F. Hancock, R.G. Parton, PTRF-Cavin, a conserved cytoplasmic protein required for caveola formation and function, *Cell* 132 (2008) 113–124.
- [6] R. Chadda, S. Mayor, PTRF triggers a cave in, *Cell* 132 (2008) 23–24.
- [7] R. Nabi, Cavin fever: regulating caveolae, *Nat. Cell. Biol.* 11 (2009) 789–791.
- [8] N. McLaughlin, B. Annabi, M. Bouzeghrane, A. Temme, J.P. Bahary, R. Moudmjan, R. Béliveau, The Survivin-mediated radioresistant phenotype of glioblastomas is regulated by RhoA and inhibited by the green tea polyphenol (-)-epigallocatechin-3-gallate, *Brain Res.* 1071 (1) (2006) 1–9.
- [9] B. Annabi, Y.T. Lee, C. Martel, A. Pilorget, J.P. Bahary, R. Béliveau, Radiation induced-tubulogenesis in endothelial cells is antagonized by the anti-angiogenic properties of green tea polyphenol (-)-epigallocatechin-3-gallate, *Cancer Biol. Ther.* 2 (6) (2003) 642–649.
- [10] C.S. Huang, J. Zhou, A.K. Feng, C.C. Lynch, J. Klumperman, S.J. DeArmond, W.C. Mobley, Nerve growth factor signaling in caveolae-like domains at the plasma membrane, *J. Biol. Chem.* 274 (1999) 36707–36714.
- [11] M. Isshiki, R.G. Anderson, Calcium signal transduction from caveolae, *Cell. Calcium* 26 (1999) 201–208.
- [12] S. Roy, R. Luetterforst, A. Harding, A. Apolloni, M. Etheridge, E. Stang, B. Rolls, J.F. Hancock, R.G. Parton, Dominant-negative caveolin inhibits H-Ras function by disrupting cholesterol-rich plasma membrane domains, *Nat. Cell. Biol.* 1 (1999) 98–105.
- [13] E.J. Smart, G.A. Graf, M.A. McNiven, W.C. Sessa, J.A. Engelman, P.E. Scherer, T. Okamoto, M.P. Lisanti, Caveolins, liquid-ordered domains, and signal transduction, *Mol. Cell. Biol.* 19 (1999) 7289–7304.
- [14] M. Bucci, J.P. Gratton, R.D. Rudic, L. Acevedo, F. Rovietto, G. Crino, W.C. Sessa, In vivo delivery of the caveolin-1 scaffolding domain inhibits nitric oxide synthesis, and reduces inflammation, *Nat. Med.* 6 (2000) 1362–1367.
- [15] C. Dessy, R.A. Kelly, J.L. Balligand, O. Feron, Dynamin mediates caveolar sequestration of muscarinic cholinergic receptors, and alteration in NO signaling, *EMBO J.* 19 (2000) 4272–4280.
- [16] B. Ranzani, A. Schlegel, M.P. Lisanti, Caveolin proteins in signaling, oncogenic transformation, and muscular dystrophy, *J. Cell. Sci.* 113 (2000) 2103–2109.
- [17] M. Stahlhut, K. Sandvig, B. Van Deurs, Caveolae: uniform structures with multiple functions in signaling, cell growth, and cancer, *Exp. Cell. Res.* 261 (2000) 111–118.
- [18] K.K. Wary, F. Mainiero, S.J. Isakoff, E.E. Marcantonio, F.G. Giancotti, The adaptor protein Shc couples a class of integrins to the control of cell cycle progression, *Cell* 87 (1996) 733–743.
- [19] Y. Wei, X. Yan, Q. Liu, J.A. Wilkins, H.A. Chapman, A role for caveolin and the urokinase receptor in integrin-mediated adhesion and signaling, *J. Cell. Biol.* 144 (1999) 1285–1294.
- [20] M. Shatz, M. Liscovitch, Caveolin-1: a tumor-promoting role in human cancer, *Int. J. Radiat. Biol.* 84 (2008) 177–189.
- [21] G. Yang, L.D. Truong, T.L. Timme, C. Ren, T.M. Wheeler, S.H. Park, Y. Nasu, C.H. Bangma, M.W. Kattan, P.T. Scardino, T.C. Thompson, Elevated expression of caveolin is associated with prostate and breast cancer, *Clin. Cancer Res.* 4 (1998) 1873–1880.
- [22] G. Yang, L.D. Truong, T.M. Wheeler, T.C. Thompson, Caveolin-1 expression in clinically confined human prostate cancer: a novel prognostic marker, *Cancer Res.* 59 (1999) 5719–5723.
- [23] C.C. Ho, P.H. Huang, H.Y. Huang, Y.H. Chen, P.C. Yang, S.M. Hsu, Up-regulated caveolin-1 accentuates the metastasis capability of lung adenocarcinoma by inducing filopodia formation, *Am. J. Pathol.* 161 (2002) 1647–1656.
- [24] H.J. Joo, D.K. Oh, Y.S. Kim, K.B. Lee, S.J. Kim, Increased expression of caveolin-1 and microvessel density correlates with metastasis and poor prognosis in clear cell renal cell carcinoma, *BJU Int.* 93 (2004) 291–296.
- [25] K. Kato, Y. Hida, M. Miyamoto, H. Hashida, T. Shinohara, T. Itoh, S. Okushiba, S. Kondo, H. Katoh, Overexpression of caveolin-1 in esophageal squamous cell carcinoma correlates with lymph node metastasis and pathologic stage, *Cancer* 94 (2002) 929–933.
- [26] T.M. Williams, M.P. Lisanti, The caveolin proteins, *Genome Biol.* 4 (2004) 214–221.
- [27] T. Okamoto, A. Schlegel, P.E. Scherer, M.P. Lisanti, Caveolins, a family of scaffolding proteins for organizing 'preassembled signaling complexes' at the plasma membrane, *J. Biol. Chem.* 273 (1998) 5419–5422.
- [28] A. Schlegel, M.P. Lisanti, A molecular dissection of caveolin-1 membrane attachment and oligomerization. Two separate regions of the caveolin-1 carboxy-terminal domain mediate membrane binding and oligomer/oligomer interactions in vivo, *J. Biol. Chem.* 275 (2000) 21605–21617.
- [29] T.V. Kurzchalia, P. Dupree, R.G. Parton, R. Kellner, H. Virta, M. Lehnert, K. Simons, VIP 21, A 21-kDa membrane protein is an integral component of trans-Golgi-network-derived transport vesicles, *J. Cell. Biol.* 118 (1992) 1003–1014.
- [30] J.R. Glenney, D. Soppet Jr., Sequence and expression of caveolin, a protein component of caveolae plasma membrane domains phosphorylated on tyrosine in Rous sarcoma virus-transformed fibroblasts, *Proc. Natl. Acad. Sci. U S A* 89 (1992) 10517–10521.
- [31] P. Liu, M. Rudick, R.G. Anderson, Multiple functions of caveolin-1, *J. Biol. Chem.* 277 (2002) 41295–41298.
- [32] W.P. Li, P. Liu, B.K. Pilcher, R.G. Anderson, Cell-specific targeting of caveolin-1 to caveolae, secretory vesicles, cytoplasm or mitochondria, *J. Cell. Sci.* 114 (2001) 1397–1408.

- [33] B. Razani, S.E. Woodman, M.P. Lisanti, Caveolae: from cell biology to animal physiology, *Pharmacol. Rev.* 54 (2002) 431–467.
- [34] S. Li, J. Couet, M.P. Lisanti, Src tyrosine kinases, Alpha subunits, and H-Ras share a common membrane anchored scaffolding protein, caveolin. Caveolin binding negatively regulates the auto-activation of Src tyrosine kinases, *J. Biol. Chem.* 271 (1996) 29182–29190.
- [35] N. Nohata, T. Hanazawa, N. Kikkawa, M. Mutallip, L. Fujimura, H. Yoshino, K. Kawakami, T. Chiyomaru, H. Enokida, M. Nakagawa, Y. Okamoto, N. Seki, Caveolin-1 mediates tumor cell migration and invasion and its regulation by miR-133a in head and neck squamous cell carcinoma, *Int. J. Oncol.* 38 (2011) 209–217.
- [36] B. Annabi, M. Lachambre, N. Bousquet-Gagnon, M. Pagé, D. Gingras, R. Béliveau, Localization of membrane-type 1 matrix metalloproteinase in caveolae membrane domains, *Biochem. J.* 353 (Pt 3) (2001) 547–553.
- [37] B. Annabi, S. Thibault, R. Moudmjan, R. Béliveau, Hyaluronan cell surface binding is induced by type I collagen and regulated by caveolae in glioma cells, *J. Biol. Chem.* 279 (21) (2004) 21888–21896.
- [38] H. Kogo, T. Aiba, T. Fujimoto, Cell type-specific occurrence of caveolin-1 α and -1 β in the lung caused by expression of distinct mRNAs, *J. Biol. Chem.* 279 (2004) 25574–25581.
- [39] H. Kogo, T. Fujimoto, Caveolin-1 isoforms are encoded by distinct mRNAs. Identification of mouse caveolin-1 mRNA variants caused by alternative transcription initiation and splicing, *FEBS Lett.* 465 (2000) 119–123.
- [40] P.E. Scherer, Z. Tang, M. Chun, M. Sargiacomo, H.F. Lodish, M.P. Lisanti, Caveolin isoforms differ in their N-terminal protein sequence and subcellular distribution. Identification and epitope mapping of an isoform-specific monoclonal antibody probe, *J. Biol. Chem.* 270 (1995) 16395–16401.
- [41] G. Sowa, M. Pyper, D. Fulton, W.C. Sessa, The phosphorylation of caveolin-2 on serines 23 and 36 modulates caveolin-1-dependent caveolae formation, *Proc. Natl. Acad. Sci. U S A* 100 (11) (2003) 6511–6516.
- [42] J.R. Glenney, L. Zokas, Novel tyrosine kinase substrates from Rous sarcoma virus-transformed cells are present in the membrane skeleton, *J. Cell. Biol.* 108 (1989) 2401–2408.
- [43] S. Li, R. Seitz, M.P. Lisanti, Phosphorylation of caveolin by Src tyrosine kinases: the α -isoform of caveolin is selectively phosphorylated by v-Src in vivo, *J. Biol. Chem.* 271 (1996) 3863–3868.
- [44] A.R. Sanguinetti, C.C. Mastick, c-Abl is required for oxidative stress-induced phosphorylation of caveolin-1 on tyrosine 14, *Cell. Signal.* 15 (2003) 289–298.
- [45] A.R. Sanguinetti, H. Cao, C. Corley Mastick, Fyn is required for oxidative- and hyperosmotic-stress-induced tyrosine phosphorylation of caveolin-1, *Biochem. J.* 376 (2003) 159–168.
- [46] A. Schlegel, P. Arvan, M.P. Lisanti, Caveolin-1 binding to endoplasmic reticulum membranes and entry into the regulated secretory pathway are regulated by serine phosphorylation. Protein sorting at the level of the endoplasmic reticulum, *J. Biol. Chem.* 276 (2001) 4398–4408.
- [47] S. Monier, D.J. Dietzen, W.R. Hastings, D.M. Lublin, T.V. Kurzchalia, Oligomerization of VIP21-caveolin in vitro is stabilized by long chain fatty acylation or cholesterol, *FEBS Lett.* 388 (1996) 143–149.
- [48] D.J. Dietzen, W.R. Hastings, D.M. Lublin, Caveolin is palmitoylated on multiple cysteine residues. Palmitoylation is not necessary for localization of caveolin to caveolae, *J. Biol. Chem.* 270 (1995) 6838–6842.
- [49] D. Volonte, F. Galbiati, R.G. Pestell, M.P. Lisanti, Cellular stress induces the tyrosine phosphorylation of caveolin-1 (Tyr14) via activation of p38 mitogen-activated protein kinase and c-Src kinase. Evidence for caveolae, the actin cytoskeleton, and focal adhesions as mechanical sensors of osmotic stress, *J. Biol. Chem.* 276 (2001) 8094–8103.
- [50] B. Van Deurs, K. Roepstorff, A.M. Hommelgaard, K. Sandvig, Caveolae: anchored, multifunctional platforms in the lipid ocean, *Trends Cell. Biol.* 13 (2003) 92–100.
- [51] F. Sotgia, J.K. Lee, K. Das, M. Bedford, T.C. Petrucci, P. Macioce, M. Sargiacomo, F.D. Bricarelli, C. Minetti, M. Sudol, M.P. Lisanti, Caveolin-3 directly interacts with the carboxy-terminal tail of beta-dystroglycan. Identification of a central WW-like domain within caveolin family members, *J. Biol. Chem.* 275 (2000) 38048–38058.
- [52] F. Bender, M. Montoya, V. Monardes, L. Leyton, A.F. Quest, Caveolae and caveolae like membrane domains in cellular signaling and disease: identification of downstream targets for the tumor suppressor protein caveolin-1, *Biol. Res.* 35 (2002) 151–167.
- [53] T. Machleidt, W.P. Li, P. Liu, R.G. Anderson, Multiple domains in caveolin-1 control its intracellular traffic, *J. Cell. Biol.* 148 (2000) 17–28.
- [54] X.H. Sun, D.C. Flynn, V. Castranova, L.L. Millecchia, A.R. Beardsley, J. Liu, Identification of a novel domain at the N-terminus of caveolin-1 that controls rear polarization of the protein and caveolae formation, *J. Biol. Chem.* 282 (2007) 7232–7241.
- [55] H.L. Chen, L.F. Fan, J. Gao, J.P. Ouyang, Y.X. Zhang, Differential expression and function of the caveolin-1 gene in non-small cell lung carcinoma, *Oncol. Rep.* 25 (2011) 359–366.
- [56] J. Couet, S. Li, T. Okamoto, T. Ikezu, M.P. Lisanti, Identification of peptide and protein ligands for the caveolin-scaffolding domain. Implications for the interaction of caveolin with caveolae-associated proteins, *J. Biol. Chem.* 272 (1997) 6525–6533.
- [57] G. Garcia-Cardena, P. Martasek, B.S. Siler-Masters, P.M. Skidd, J.C. Couet, S. Li, M.P. Lisanti, W.C. Sessa, Dissecting the interaction between nitric oxide synthase (NOS) and caveolin. Functional significance of the nos caveolin binding domain in vivo, *J. Biol. Chem.* 272 (1997) 25437–25440.
- [58] R.S. Ostrom, P.A. Insel, The evolving role of lipid rafts and caveolae in G protein-coupled receptor signaling: implications for molecular pharmacology, *Br. J. Pharmacol.* 143 (2004) 235–245.
- [59] J.A. Engelman, R.J. Lee, A. Karnezis, D.J. Bearss, M. Webster, P. Siegel, W.J. Muller, J.J. Windle, R.G. Pestell, M.P. Lisanti, Reciprocal regulation of neu tyrosine kinase activity and caveolin-1 protein expression in vitro and in vivo: Implications for human breast cancer, *J. Biol. Chem.* 273 (1998) 20448–20455.
- [60] L. Druhan, K. Karupiah, C.A. Chen, T. Smith, J.L. Zweier, W.C. Sessa, A.J. Cardounel, Suppression of eNOS-derived superoxide by caveolin-1: a biotin-dependent mechanism, *Am. J. Physiol. Heart Circ. Physiol.* 301 (2011) H903–H911.
- [61] X. Wang, S. Feng, H. Zhang, Y. Wang, Y. Cui, Z. Wang, J. Liu, W. Zou, RNA interference-mediated caveolin-1 down-regulation decrease estrogen receptor α (ER α) signaling in human mammary epithelial cells, *Mol. Biol. Rep.* 38 (2011) 761–768.
- [62] X. Zhang, M.T. Ling, Q. Wang, C.K. Lau, S.C. Leung, T.K. Lee, A.L. Cheung, Y.C. Wong, X. Wang, Identification of a novel inhibitor of differentiation-1 (ID-1) binding partner, caveolin-1, and its role in epithelial-mesenchymal transition and resistance to apoptosis in prostate cancer cells, *J. Biol. Chem.* 282 (2007) 33284–33294.
- [63] S.K. Miyasato, J. Loeffler, R. Shohet, J. Zhang, M. Lindsey, C.J. Le Saux, Caveolin-1 modulates TGF- β 1 signaling in cardiac remodeling, *Matrix Biol.* 30 (2011) 318–329.
- [64] A.S. Martins, J.L. Ordóñez, A.T. Amaral, F. Prins, G. Floris, M. Debiec-Rychter, P.C. Hogendoorn, E. de Alava, IGF1R signaling in Ewing sarcoma is shaped by clathrin/caveolin-dependent endocytosis, *PLoS One* 6 (2011) 19846–19857.
- [65] L. Zhang, Y. Wang, L. Wang, T. Cao, S. Hyuga, T. Sato, Y. Wu, S. Yamagata, T. Yamagata, Ganglioside GD1a negatively regulates hepatocyte growth factor expression through caveolin-1 at the transcriptional level in murine osteosarcoma cells, *Biochim. Biophys. Acta* 1810 (2011) 759–768.
- [66] B.M.1 Collins, M.J. Davis, J.F. Hancock, R.G. Parton, Structure-based reassessment of the caveolin signaling model: do caveolae regulate signaling through caveolin-protein interactions? *Dev. Cell.* 23 (2012) 11–20.
- [67] D. Hanahan, R.A. Weinberg, Hallmarks of cancer: the next generation, *Cell* 144 (5) (2011) 646–674.
- [68] T.M. Williams, G.S. Hassan, J. Li, A.W. Cohen, F. Medina, P.G. Frank, R.G. Pestell, D. Di Vizio, M. Loda, M.P. Lisanti, Caveolin-1 promotes tumor progression in an autochthonous mouse model of prostate cancer: genetic ablation of Cav-1 delays advanced prostate tumor development in trans mice, *J. Biol. Chem.* 280 (2005) 25134–25145.
- [69] K. Quann, D.M. Gonzales, I. Mercier, C. Wang, F. Sotgia, R.G. Pestell, M.P. Lisanti, J.F. Jasmin, Caveolin-1 is a negative regulator of tumor growth in glioblastoma and modulates chemosensitivity to temozolomide, *Cell. Cycle* 12 (10) (2013) 1510–1520.
- [70] J.A. Engelman, C.C. Wykoff, S. Yashuhara, K.S. Song, T. Okamoto, M.P. Lisanti, Recombinant expression of caveolin-1 in oncogenically transformed cells abrogates anchorage-independent growth, *J. Biol. Chem.* 272 (1997) 16374–16381.
- [71] M. Sáinz-Jaspeado, J. Martín-Liberal, L. Lagares-Tena, S. Mateo-Lozano, X. Garcia del Muro, O.M. Tirado, Caveolin-1 in sarcomas: friend or foe? *Oncotarget* 2 (4) (2011) 305–312.
- [72] A.J. Koleske, D. Baltimore, M.P. Lisanti, Reduction of caveolin and caveolae in oncogenically transformed cells, *Proc. Natl. Acad. Sci. U S A* 92 (1995) 1381–1385.
- [73] S. Feng, Y. Wang, X. Wang, Z. Wang, Y. Cui, J. Liu, C. Zhao, M. Jin, W. Zou, Caveolin-1 gene silencing promotes the activation of PI3K/AKT dependent on Eralpha36 and the transformation of MCF10ACE, *Sci. China Life Sci.* 53 (2010) 598–605.
- [74] F. Han, D. Gu, Q. Chen, H. Zhu, Caveolin-1 acts as a tumor suppressor by down-regulating epidermal growth factor receptor-mitogen-activated protein kinase signaling pathway in pancreatic carcinoma cell lines, *Pancreas* 38 (2009) 766–774.
- [75] F. Galbiati, D. Volonte, J.A. Engelman, G. Watanabe, R. Burk, R.G. Pestell, M.P. Lisanti, Targeted down-regulation of caveolin-1 is sufficient to drive cell transformation and hyperactivate the p42/44 MAP kinase cascade, *EMBO J.* 17 (1998) 6633–6648.
- [76] D.T. Bau, M.H. Tsai, Y.A. Tsou, C.H. Wang, C.W. Tsai, S.S. Sun, C.H. Hua, S.K. Shyue, R.Y. Tsai, The association of caveolin-1 genotypes with oral cancer susceptibility in Taiwan, *Ann. Surg. Oncol.* 18 (2011) 1431–1438.
- [77] C.M. Hsu, M.D. Yang, C.W. Tsai, C.Y. Ho, W.S. Chang, S.C. Chang, L.B. Jeng, Y. Tsai, F.J. Tsai, D.T. Bau, The contribution of caveolin-1 genotype and phenotype to hepatocellular carcinoma, *Anticancer Res.* 33 (2013) 671–677.
- [78] F. Capozza, T.M. Williams, W. Schubert, S. McClain, B. Bouzahzah, F. Sotgia, M.P. Lisanti, Absence of caveolin-1 sensitizes mouse skin to carcinogen-induced epidermal hyperplasia and tumor formation, *Am. J. Pathol.* 162 (2003) 2029–2039.
- [79] H. Wikman, E. Kettunen, J.K. Seppanen, A. Karjalainen, J. Hollmen, S. Anttila, S. Knuutila, Identification of differentially expressed genes in pulmonary adenocarcinoma by using cDNA array, *Oncogene* 21 (2002) 5804–5813.

- [80] C. Racine, M. Belanger, H. Hirabayashi, M. Boucher, J. Chakir, J. Couet, Reduction of caveolin 1 gene expression in lung carcinoma cell lines, *Biochem Biophys. Res. Commun.* 255 (1999) 580–586.
- [81] S.W. Lee, C.L. Reimer, P. Oh, D.B. Campbell, J.E. Schnitzer, Tumor cell growth inhibition by caveolin re-expression in human breast cancer cells, *Oncogene* 16 (1998) 1391–1397.
- [82] D.A. Zuccari, R. Castro, A.F. Gavioli, U.M. Mancini, C.S. Frade, C. Leonel, Immunohistochemical and molecular analysis of caveolin-1 expression in canine mammary tumors, *Genet. Mol. Res.* 11 (2012) 153–165.
- [83] F.C. Bender, M.A. Reymond, C. Bron, A.F. Quest, Caveolin-1 levels are down-regulated in human colon tumors, and ectopic expression of caveolin-1 in colon carcinoma cell lines reduces cell tumorigenicity, *Cancer Res.* 60 (2000) 5870–5878.
- [84] K. Wiechen, L. Diatchenko, A. Agoulnik, K.M. Scharff, H. Schober, K. Arlt, B. Zhumabayeva, P.D. Siebert, M. Dietel, R. Schafer, C. Sers, Caveolin-1 is downregulated in human ovarian carcinoma and acts as a candidate tumor suppressor gene, *Am. J. Pathol.* 159 (2001) 1635–1643.
- [85] A.K. Witkiewicz, A. Dasgupta, F. Sotgia, I. Mercier, R.G. Pestell, M. Sabel, C.G. Kleer, J.R. Brody, M.P. Lisanti, An absence of stromal caveolin-1 expression predicts early tumor recurrence and poor clinical outcome in human breast cancers, *Am. J. Pathol.* 174 (2009) 2023–2034.
- [86] K. Wiechen, C. Sers, A. Agoulnik, K. Arlt, M. Dietel, P.M. Schlag, U. Schneider, Down-regulation of caveolin-1, a candidate tumor suppressor gene, in sarcomas, *Am. J. Pathol.* 158 (2001) 833–839.
- [87] L. Cantiani, M.C. Manara, C. Zucchini, P. De Sanctis, M. Zuntini, L. Valvassori, M. Serra, M. Olivero, M.F. Di Renzo, M.P. Colombo, P. Picci, K. Scotlandi, Caveolin-1 reduces osteosarcoma metastases by inhibiting c-Src activity and met signaling, *Cancer Res.* 67 (2007) 7675–7685.
- [88] D.A. Rodriguez, J.C. Tapia, J.G. Fernandez, V.A. Torres, N. Muñoz, D. Galleguillos, L. Leyton, A.F. Quest, Caveolin-1-mediated suppression of cyclooxygenase-2 via a beta-catenin-Tcf/Lef-dependent transcriptional mechanism reduced prostaglandin E2 production and survivin expression, *Mol. Biol. Cell.* 20 (2009) 2297–2310.
- [89] C.A. Hanson, K.R. Drake, M.A. Baird, B. Han, L.J. Kraft, M.W. Davidson, A.K. Kenworthy, Overexpression of caveolin-1 is sufficient to phenocopy the behavior of a disease-associated mutant, *Traffic* 14 (2013) 663–677.
- [90] P.E. Scherer, M.P. Lisanti, G. Baldini, M. Sargiacomo, C.C. Mastick, H.F. Lodish, Induction of caveolin during adipogenesis and association of GLUT4 with caveolin-rich vesicles, *J. Cell. Biol.* 127 (1994) 1233–1243.
- [91] K. Hayashi, S. Matsuda, K. Machida, T. Yamamoto, Y. Fukuda, Y. Nimura, T. Hayakawa, M. Hamauchi, Invasion activating caveolin-1 mutation in human scirrhous breast cancers, *Cancer Res.* 61 (2001) 2361–2364.
- [92] J. Cui, L.R. Rohr, G. Swanson, V.O. Speights, T. Maxwell, A.R. Brothman, Hypermethylation of the caveolin-1 gene promoter in prostate cancer, *Prostate* 46 (2001) 249–256.
- [93] X. Rao, J. Evans, H. Chae, J. Pirose, S. Kim, P. Yan, R.L. Huang, H.C. Lai, H. Lin, Y. Liu, J.K. Rhee, Y.W. Huang, F. Gu, J.W. Gray, T.M. Huang, K.P. Nephew, CpG island shore methylation regulates caveolin-1 expression in breast cancer, *Oncogene* (2012), <http://dx.doi.org/10.1038/ncr.2012.474>.
- [94] M.D. Yang, R.Y. Tsai, C.S. Liu, C.H. Chang, H.C. Wang, Y.A. Tsou, C.H. Wang, C.C. Lin, S.K. Shyue, D.T. Bau, Association of Caveolin-1 polymorphisms with colorectal cancer susceptibility in Taiwan, *World J. Gastrointest. Oncol.* 2 (2010) 326–331.
- [95] N. Syeed, S.A. Husain, S. Abdullah, A.S. Sameer, N.A. Chowdri, M.S. Nanda, M.A. Siddiqi, Mutational profile of the CAV-1 gene in breast cancer cases in the ethnic Kashmiri population, *Asian Pac. J. Cancer Prev.* 11 (2010) 1099–1105.
- [96] K.H. Nam, B.L. Lee, J.H. Park, J. Kim, N. Han, H.E. Lee, M.A. Kim, H.S. Lee, W.H. Kim, Caveolin 1 expression correlates with poor prognosis and focal adhesion kinase expression in gastric cancer, *Pathobiology* 80 (2013) 87–94.
- [97] R. Bartz, J. Zhou, J.T. Hsieh, Y. Ying, W. Li, P. Liu, Caveolin-1 secreting LNCaP cells induce tumor growth of caveolin-1 negative LNCaP cells in vivo, *Int. J. Cancer* 122 (2007) 520–525.
- [98] L. Li, G. Yang, S. Ebara, T. Satoh, Y. Nasu, T.L. Timme, C. Ren, J. Wang, S.A. Tahir, T.C. Thompson, Caveolin-1 mediates testosterone-stimulated survival/clonal growth and promotes metastatic activities in prostate cancer cells, *Cancer Res.* 61 (2001) 4386–4392.
- [99] S.A. Tahir, G. Yang, S. Ebara, T.L. Timme, T. Satoh, L. Li, A. Goltsov, M. Ittmann, J.D. Morrisett, T.C. Thompson, Secreted caveolin-1 stimulates cell survival/clonal growth and contributes to metastasis in androgen-insensitive prostate cancer, *Cancer Res.* 61 (2001) 3882–3885.
- [100] M.A. Del Pozo, N. Balasubramanian, N.B. Alderson, W.B. Kiosses, A. Grande-García, R.G. Anderson, M.A. Schwartz, Phospho-caveolin-1 mediates integrin-regulated membrane domain internalization, *Nat. Cell. Biol.* 7 (2005) 901–908.
- [101] Y. Nasu, T.L. Timme, G. Yang, C.H. Bangma, L. Li, C. Ren, S.H. Park, M. DeLeon, J. Wang, T.C. Thompson, Suppression of caveolin expression induces androgen sensitivity in metastatic androgen-insensitive mouse prostate cancer cells, *Nat. Med.* 4 (1998) 1062–1064.
- [102] T. Ando, H. Ishiguro, M. Kimura, A. Mitsui, Y. Mori, N. Sugito, K. Tomoda, R. Mori, K. Harada, T. Katada, R. Ogawa, Y. Fujii, Y. Kuwabara, The overexpression of caveolin-1 and caveolin-2 correlates with a poor prognosis and tumor progression in esophageal squamous cell carcinoma, *Oncol. Rep.* 18 (2007) 601–609.
- [103] W.T. Chiu, H.T. Lee, F.J. Huang, K.D. Aldape, J. Yao, P.S. Steeg, C.Y. Chou, Z. Lu, K. Xie, S. Huang, Caveolin-1 upregulation mediates suppression of primary breast tumor growth and brain metastases by Stat3 inhibition, *Cancer Res.* 71 (2011) 4932–4943.
- [104] S. Waalkes, H. Eggers, H. Blasig, F. Atschekzei, M.W. Kramer, J. Hennenlotter, W. Tränkle, A. Stenzl, J. Serth, A.J. Schrader, M.A. Kuczyk, A.S. Merseburger, Caveolin-1 mRNA is overexpressed in malignant renal tissue and might serve as a novel diagnostic marker for renal cancer, *Biomark. Med.* 5 (2011) 219–225.
- [105] V. Barresi, S. Cerasoli, G. Paioli, E. Vitarelli, G. Giuffrè, G. Guiducci, et al., Caveolin-1 in meningiomas: expression and clinico-pathological correlations, *Acta Neuropathol.* 112 (2006) 617–626.
- [106] F. Pancotti, L. Roncuzzi, M. Maggolini, A. Gasperi-Campani, Caveolin-1 silencing arrests the proliferation of metastatic lung cancer cells through the inhibition of STAT3 signaling, *Cell. Signal* 24 (2012) 1390–1397.
- [107] R. Senetta, C. Miracco, S. Lanzafame, L. Chiusa, R. Caltabiano, A. Galia, G. Stella, P. Cassoni, Epidermal growth factor receptor and caveolin-1 coexpression identifies adult supratentorial ependymomas with rapid unfavorable outcomes, *Neuro Oncol.* 13 (2011) 176–183.
- [108] K.G. Bryant, J. Camacho, J.F. Jasmin, C. Wang, S. Addya, M.C. Casimiro, P. Fortina, S. Balasubramanian, K.E. Knudsen, R. Schwarting, M.P. Lisanti, I. Mercier, Caveolin-1 overexpression enhances androgen-dependent growth and proliferation in the mouse prostate, *Int. J. Biochem. Cell. Biol.* 43 (2011) 1318–1329.
- [109] J.A. Karam, Y. Lotan, C.G. Roehrborn, R. Ashfaq, P.I. Karakiewicz, S.F. Shariat, Caveolin-1 overexpression is associated with aggressive prostate cancer recurrence, *Prostate* 67 (2007) 614–622.
- [110] G. Yang, J. Addai, T.M. Wheeler, A. Frolov, B.J. Miles, D. Kadmon, T.C. Thompson, Correlative evidence that prostate cancer cell-derived caveolin-1 mediates angiogenesis, *Hum. Pathol.* 38 (2007) 1688–1695.
- [111] I. Steiner, K. Jung, K. Miller, C. Stephan, A. Erbersdobler, Expression of endothelial factors in prostate cancer: a possible role of caveolin-1 for tumour progression, *Oncol. Rep.* 27 (2012) 389–395.
- [112] Y. Lavie, G. Ficuci, M. Liscovitch, Up-regulation of caveolae and caveolar constituents in multidrug-resistant cancer, *J. Biol. Chem.* 273 (1998) 32380–32383.
- [113] G. Ficuci, D. Ravid, R. Reich, M. Liscovitch, Caveolin-1 inhibits anchorage-independent growth, anoikis and invasiveness in MCF-7 human breast cancer cells, *Oncogene* 21 (2002) 2365–2375.
- [114] D. Ravid, S. Maor, H. Werner, M. Liscovitch, Caveolin-1 inhibits anoikis and promotes survival signaling in cancer cells, *Adv. Enzyme Regul.* 46 (2006) 163–175.
- [115] G. Yuan, I. Regel, F. Lian, T. Friedrich, I. Hitkova, R.D. Hofheinz, P. Ströbel, R. Langer, G. Keller, C. Röcken, W. Zimmermann, R.M. Schmid, M.P. Ebert, E. Burgermeister, WNT6 is a novel target gene of caveolin-1 promoting chemoresistance to epirubicin in human gastric cancer cells, *Oncogene* 32 (2013) 375–387.
- [116] M. Li, H. Chen, L. Diao, Y. Zhang, C. Xia, F. Yang, Caveolin-1 and VEGF-C promote lymph node metastasis in the absence of intratumoral lymphangiogenesis in non-small cell lung cancer, *Tumori* 96 (2010) 734–743.
- [117] S. Garcia, J.P. Dales, E. Charafe-Jauffret, S. Carpentier-Meunier, L. Andrac-Meyer, J. Jacquemier, C. Andonian, M.N. Lavaut, C. Allasia, P. Bonnier, C. Charpin, Poor prognosis in breast carcinomas correlates with increased expression of targetable CD146 and c-Met and with proteomic basal-like phenotype, *Hum. Pathol.* 38 (2007) 830–841.
- [118] T.K. Ha, N.G. Her, M.G. Lee, B.K. Ryu, J.H. Lee, J. Han, S.I. Jeong, M.J. Kang, N.H. Kim, H.J. Kim, S.G. Chi, Caveolin-1 increases aerobic glycolysis in colorectal cancers by stimulating HMG1-mediated GLUT3 transcription, *Cancer Res.* 72 (2012) 4097–4109.
- [119] T.K. Ha, S.G. Chi, CAV1/caveolin 1 enhances aerobic glycolysis in colon cancer cells via activation of SLC2A3/GLUT3 transcription, *Autophagy* 8 (2012) 1684–1685.
- [120] K.A. Sloan, H.A. Marquez, J. Li, Y. Cao, A. Hinds, C.J. O'Hara, S. Kathuria, M.I. Ramirez, C.M. Williams, H. Kathuria, Increased PEA3/E1AF and decreased Net/Elk-3, both ETS proteins, characterize human NSCLC progression and regulate caveolin-1 transcription in Calu-1 and NCI-H23 NSCLC cell lines, *Carcinogenesis* 30 (2009) 1433–1442.
- [121] J.G. Goetz, B. Joshi, P. Lajoie, Concerted regulation of focal adhesion dynamics by galectin-3 and tyrosine phosphorylated caveolin-1, *J. Cell. Biol.* 180 (2008) 1261–1275.
- [122] B. Annabi, M. Bouzeghrane, R. Moumdjian, A. Moghrabi, R. Béliveau, Probing the infiltrating character of brain tumors: inhibition of RhoA/ROK-mediated CD44 cell surface shedding from glioma cells by the green tea catechin EGCG, *J. Neurochem.* 94 (4) (2005) 906–916.
- [123] B. Joshi, S.S. Strugnell, J.G. Goetz, M.E. Cox, O.L. Griffith, S.K. Chan, S.J. Jones, S.P. Leung, H. Masoudi, S. Leung, S.M. Wiseman, I.R. Nabi, Phosphorylated Caveolin-1 regulates Rho/ROK-dependent focal adhesion dynamics and tumor cell migration and invasion, *Cancer Res.* 68 (2008) 8210–8220.
- [124] E. Arpaia, H. Blaser, M. Quintela-Fandino, G. Duncan, H.S. Leong, A. Ablack, S.C. Nambiar, E.F. Lind, J.J. Silvester, C.K. Fleming, A. Rufini, M.W. Tusche, A. Brüstle, P.S. Ohashi, J.D. Lewis, T.W. Mak, The interaction between caveolin-1 and Rho-GTPases promotes metastasis by controlling the expression of alpha5-integrin and the activation of Src, Ras and Erk, *Oncogene* 31 (2012) 884–896.

- [125] Y. Tang, X. Zeng, F. He, Y. Liao, N. Qian, M. Toi, Caveolin-1 is related to invasion, survival, and poor prognosis in hepatocellular cancer, *Med. Oncol.* 29 (2012) 977–984.
- [126] F. Felicetti, I. Parolini, L. Bottero, K. Fecchi, M.C. Errico, C. Raggi, M. Biffoni, F. Spadaro, M.P. Lisanti, M. Sargiacomo, A. Carè, Caveolin-1 tumor-promoting role in human melanoma, *Int. J. Cancer* 125 (2009) 1514–1522.
- [127] J.G. Goetz, S. Minguet, I. Navarro-Lérida, J.J. Lazzano, R. Samaniego, E. Calvo, M. Tello, T. Osteso-Ibáñez, T. Pellinen, A. Echarri, A. Cerezo, A.J. Klein-Szanto, R. Garcia, P.J. Keely, P. Sánchez-Mateos, E. Cukierman, M.A. Del Pozo, Biomechanical remodeling of the microenvironment by stromal caveolin-1 favors tumor invasion and metastasis, *Cell* 146 (2011) 148–163.
- [128] K.K. Wary, A. Mariotti, C. Zurzolo, F.G. Giancotti, A requirement for caveolin-1 and associated kinase Fyn in integrin signaling and anchorage-dependent cell growth, *Cell* 94 (1998) 625–634.
- [129] K. Gaus, S. Le Lay, N. Balasubramanian, M.A. Schwartz, Integrin-mediated adhesion regulates membrane order, *J. Cell. Biol.* 174 (2006) 725–734.
- [130] M. Nethe, P.L. Hordijk, A model for phospho-caveolin-1-driven turnover of focal adhesions, *Cell. Adh. Migr.* 5 (2011) 59–64.
- [131] C. Trimmer, D. Whitaker-Menezes, G. Bonuccelli, J.N. Millman, K.M. Daumer, A.E. Aplin, R.G. Pestell, F. Sotgia, M.P. Lisanti, F. Capozza, CAV1 inhibits metastatic potential in melanomas through suppression of the integrin/Src/FAK signaling pathway, *Cancer Res.* 70 (2010) 7489–7499.
- [132] F. Galbiati, D. Volonté, J. Liu, F. Capozza, P.G. Frank, L. Zhu, R.G. Pestell, M.P. Lisanti, Caveolin-1 expression negatively regulates cell cycle progression by inducing G(0)/G(1) arrest via a p53/p21(WAF1/Cip1)-dependent mechanism, *Mol. Biol. Cell.* 12 (2001) 2229–2244.
- [133] A.N. Shajahan, A. Wang, M. Decker, R.D. Minshall, M.C. Liu, R. Clarke, Caveolin-1 tyrosine phosphorylation enhances paclitaxel-mediated cytotoxicity, *J. Biol. Chem.* 282 (8) (2007) 5934–5943.
- [134] A. Linge, P. Meleady, M. Henry, M. Clynes, M. Kasper, K. Barth, Bleomycin treatment of A549 human lung cancer cells results in association of MGr1-Ag and caveolin-1 in lipid rafts, *Int. J. Biochem. Cell. Biol.* 43 (2011) 98–105.
- [135] T. Songserm, V. Pongrakhananon, P. Chanvorachote, Sub-toxic cisplatin mediates anoikis resistance through hydrogen peroxide-induced caveolin-1 up-regulation in non-small cell lung cancer cells, *Anticancer Res.* 32 (2012) 1659–1669.
- [136] P. Rungtapanap, U. Nimmannit, H. Halim, Y. Rojanasakul, P. Chanvorachote, Hydrogen peroxide inhibits non-small cell lung cancer cell anoikis through the inhibition of caveolin-1 degradation, *Am. J. Physiol. Cell. Physiol.* 300 (2011) C235–C245.
- [137] C.J. Percy, B.K. Pat, H. Healy, D.W. Johnson, G.C. Gobe, Phosphorylation of caveolin-1 is anti-apoptotic and promotes cell attachment during oxidative stress of kidney cells, *Pathology* 40 (2008) 694–701.
- [138] H. Zou, D. Volonté, F. Galbiati, Interaction of caveolin-1 with Ku70 inhibits Bax-mediated apoptosis, *PLoS One* 7 (2012) e39379.
- [139] X. Zhao, Y. Liu, Q. Ma, X. Wang, H. Jin, M. Mehrpour, Q. Chen, Caveolin-1 negatively regulates TRAIL-induced apoptosis in human hepatocarcinoma cells, *Biochem. Biophys. Res. Commun.* 378 (2009) 21–26.
- [140] S. Wang, L. Jia, H. Zhou, X. Wang, J. Zhang, Caveolin-1 promotes the transformation and anti-apoptotic ability of mouse hepatoma cells, *IUBMB Life* 60 (2008) 693–699.
- [141] D. Ravid, S. Maor, H. Werner, M. Liscovitch, Caveolin-1 inhibits cell detachment-induced p53 activation and anoikis by upregulation of insulin-like growth factor-I receptors and signaling, *Oncogene* 24 (2005) 1338–1347.
- [142] C. Glait, L. Tencer, D. Ravid, R. Sarfstein, M. Liscovitch, H. Werner, Caveolin-1 up-regulates IGF-I receptor gene transcription in breast cancer cells via Sp1- and p53-dependent pathways, *Exp. Cell. Res.* 312 (2006) 3899–3908.
- [143] M. Zhang, S.J. Lee, C. An, J.F. Xu, B. Joshi, I.R. Nabi, A.M. Choi, Y. Jin, Caveolin-1 mediates Fas-BID signaling in hyperoxia-induced apoptosis, *Free Radic. Biol. Med.* 50 (2011) 1252–1262.
- [144] E.J. Jung, C.W. Kim, Caveolin-1 inhibits TrkA-induced cell death by influencing on TrkA modification associated with tyrosine-490 phosphorylation, *Biochem. Biophys. Res. Commun.* 402 (2010) 736–741.
- [145] J. Park, E. Bae, C. Lee, S.S. Yoon, Y.S. Chae, K.S. Ahn, N.H. Won, RNA interference-directed caveolin-1 knockdown sensitizes SN12CPM6 cells to doxorubicin-induced apoptosis and reduces lung metastasis, *Tumor Biol.* 31 (2010) 643–650.
- [146] P.E. Scherer, T. Okamoto, M. Chun, I. Nishimoto, H.F. Lodish, M.P. Lisanti, Identification, sequence and expression of caveolin-2 defines a caveolin gene family, *Proc. Natl. Acad. Sci. U S A* 93 (1996) 131–135.
- [147] M.A. Aldred, M.E. Ginn-Pease, C.D. Morrison, A.P. Popkie, O. Gimm, C. Hoang-Vu, U. Krause, H. Dralle, S.M. Jhiang, C. Plass, C. Eng, Caveolin-1 and caveolin-2, together with three bone morphogenetic protein-related genes, may encode novel tumor suppressors down-regulated in sporadic follicular thyroid carcinogenesis, *Cancer Res.* 63 (2003) 2864–2871.
- [148] K.A. McMahon, H. Zajicek, W.P. Li, M.J. Peyton, J.D. Minna, V.J. Hernandez, L.K. Phelps, R.G. Anderson, SRBC/cavin-3 is a caveolin adapter protein that regulates caveolae function, *EMBO J.* 28 (2009) 1001–1015.
- [149] P.E. Scherer, R.Y. Lewis, D. Volonté, J.A. Engelman, F. Galbiati, J. Couet, D.S. Kohtz, E. van Donselaar, P. Peters, M.P. Lisanti, Cell-type and tissue-specific expression of caveolin-2. Caveolins 1 and 2 co-localize and form a stable hetero-oligomeric complex in vivo, *J. Biol. Chem.* 272 (1997) 29337–29346.
- [150] L.L. Zhu, Y. Cui, Y.S. Chang, F.D. Fang, A second protein marker of caveolae: caveolin-2, *Chin. Med. Sci. J.* 25 (2010) 119–124.
- [151] M. Hahn-Obercyger, L. Graeve, Z. Madar, A high-cholesterol diet increases the association between caveolae and insulin receptors in rat liver, *J. Lipid Res.* 50 (2009) 98.
- [152] I. Parolini, M. Sargiacomo, F. Galbiati, G. Rizzo, F. Grignani, J.A. Engelman, T. Okamoto, T. Ikezu, P.E. Scherer, R. Moraet, et al., Expression of caveolin-1 is required for the transport of caveolin-2 to the plasma membrane. Retention of caveolin-2 at the level of the golgi complex, *J. Biol. Chem.* 274 (1999) 25718–25725.
- [153] M.P. Lisanti, P.E. Scherer, Z. Tang, M. Sargiacomo, Caveolae, caveolin and caveolin-rich membrane domains: a signalling hypothesis, *Trends Cell. Biol.* 4 (1994) 231–235.
- [154] L. Breuza, S. Corby, J.P. Arsanto, M.H. Delgrossi, P. Scheiffele, A. Le Bivic, The scaffolding domain of caveolin 2 is responsible for its golgi localization in caco-2 cells, *J. Cell. Sci.* 115 (2002) 4457–4467.
- [155] X.H. Sun, Z.Y. Liu, H. Chen, A.R. Beardsley, Q. Qi, J.A. Liu, conserved sequence in caveolin-1 is both necessary and sufficient for caveolin polarity and cell directional migration, *FEBS Lett.* 583 (2009) 3681–3689.
- [156] H. Kwon, K. Jeong, Y. Pak, Identification of pY19-caveolin-2 as a positive regulator of insulin-stimulated actin cytoskeleton-dependent mitogenesis, *J. Cell. Mol. Med.* 13 (2009) 1549–1564.
- [157] S. Lee, H. Kwon, K. Jeong, Y. Pak, Regulation of cancer cell proliferation by caveolin-2 down-regulation and re-expression, *Int. J. Oncol.* 38 (2011) 1395–1402.
- [158] D. Kim, H. Kim, J.S. Koo, Expression of caveolin-1, caveolin-2 and caveolin-3 in thyroid cancer and stroma, *Pathobiology* 79 (2012) 1–10.
- [159] Y.C. Hu, K.Y. Lam, S. Law, J. Wong, G. Srivastava, Profiling of differentially expressed cancer-related genes in esophageal squamous cell carcinoma (ESCC) using human cancer cDNA arrays: overexpression of oncogene MET correlates with tumor differentiation in ESCC, *Clin. Cancer Res.* 7 (11) (2001) 3519–3525.
- [160] A. Fong, E. Garcia, L. Gwynn, M.P. Lisanti, M.J. Fazzari, M. Li, Expression of caveolin-1 and caveolin-2 in urothelial carcinoma of the urinary bladder correlates with tumor grade and squamous differentiation, *Am. J. Clin. Pathol.* 120 (1) (2003) 93–100.
- [161] R. Hnasko, M.P. Lisanti, The biology of caveolae: lessons from caveolin knockout mice and implications for human disease, *Mol. Interv.* 3 (8) (2003) 445–464.
- [162] L. Xie, P.G. Frank, M.P. Lisanti, G. Sowa, Endothelial cells isolated from caveolin-2 knockout mice display higher proliferation rate and cell cycle progression relative to their wild-type counterparts, *Am. J. Physiol. Cell. Physiol.* 298 (2010) C693–C701.
- [163] S.E. Elsheikh, A.R. Green, E.A. Rakha, R.M. Samaka, A.A. Ammar, D. Powe, J.S. Reis-Filho, I.O. Ellis, Caveolin 1 and caveolin 2 are associated with breast cancer basal-like and triple-negative immunophenotype, *Br. J. Cancer* 99 (2008) 327–334.
- [164] Y. Sagara, K. Mimori, K. Yoshinaga, F. Tanaka, K. Nishida, S. Ohno, H. Inoue, M. Mori, Clinical significance of Caveolin-1, Caveolin-2 and HER2/neu mRNA expression in human breast cancer, *Br. J. Cancer* 91 (2004) 959–965.
- [165] W.I. Silva, H.M. Maldonado, G. Velazquez, M. Rubio-Davila, J.D. Miranda, E. Aquino, N. Mayol, A. Cruz-Torres, J. Jardon, I.K. Salgado-Villanueva, Caveolin isoform expression during differentiation of C6 glioma cells, *Int. J. Dev. Neurosci.* 23 (2005) 599–612.
- [166] T. Yamasaki, N. Seki, H. Yoshino, T. Itesako, H. Hidaka, Y. Yamada, S. Tatarano, T. Yonezawa, T. Kinoshita, M. Nakagawa, H. Enokida, MicroRNA-218 inhibits cell migration and invasion in renal cell carcinoma through targeting Caveolin-2 involved in focal adhesion pathway, *J. Urol.* 190 (2013) 1059–1068.
- [167] Z. Tang, P.E. Scherer, T. Okamoto, K. Song, C. Chu, D.S. Kohtz, I. Nishimoto, H.F. Lodish, M.P. Lisanti, Molecular cloning of caveolin-3, a novel member of the caveolin gene family expressed predominantly in muscle, *J. Biol. Chem.* 271 (1996) 2255–2261.
- [168] S.R. Fuhs, P.A. Insel, Caveolin-3 undergoes SUMOylation by the SUMO E3 ligase PIASy: sumoylation affects G-protein-coupled receptor desensitization, *J. Biol. Chem.* 286 (2011) 14830–14841.
- [169] K.S. Song, P.E. Scherer, Z. Tang, T. Okamoto, S. Li, M. Chafel, C. Chu, D.S. Kohtz, M.P. Lisanti, Expression of caveolin-3 in skeletal, cardiac, and smooth muscle cells. Caveolin-3 is a component of the sarcolemma and co-fractionates with dystrophin and dystrophin-associated glycoproteins, *J. Biol. Chem.* 271 (1996) 15160–15165.
- [170] V.O. Rybin, X. Xu, M.P. Lisanti, S.F. Steinberg, Differential targeting of beta-adrenergic receptor subtypes and adenylyl cyclase to cardiomyocyte caveolae. A mechanism to functionally regulate the cAMP signaling pathway, *J. Biol. Chem.* 275 (2000) 41447–41457.
- [171] Y.T. Horikawa, M. Panneerselvam, Y. Kawaraguchi, Y.M. Tsutsumi, S.S. Ali, R.C. Balijepalli, F. Murray, B.P. Head, I.R. Niesman, T. Rieg, V. Vallon, P.A. Insel, H.H. Patel, D.M. Roth, Cardiac-specific over-expression of caveolin-3 attenuates cardiac hypertrophy and increases natriuretic peptide expression and signaling, *J. Am. Coll. Cardiol.* 57 (2011) 2273–2283.
- [172] E. Stoppini, S. Rossi, E. Meacci, F. Penna, P. Costelli, A. Bellucci, F. Faggi, D. Maiolo, E. Monti, A. Fanzani, Point mutated caveolin-3 form (P104L) impairs myoblast differentiation via Akt and p38 signalling reduction, leading to an immature cell signature, *Biochim. Biophys. Acta* 1812 (2011) 468–479.

- [173] A. Kuga, Y. Ohsawa, T. Okada, F. Kanda, M. Kanagawa, T. Toda, Y. Sunada, Endoplasmic reticulum stress response in P104L mutant caveolin-3 transgenic mice, *Hum. Mol. Genet.* 20 (2011) 2975–2983.
- [174] B. Razani, J.A. Engelman, X.B. Wang, W. Schubert, X.L. Zhang, C.B. Marks, F. Macaluso, R.G. Russell, M. Li, R.G. Pestell, D.D. Vizio, H. Hou, H.B. Kneitz, G. Lagaud, G.J. Christ, W. Edelmann, M.P. Lisanti, Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities, *J. Biol. Chem.* 276 (2001) 38121–38138.
- [175] M. Drab, P. Verkade, M. Elger, M. Kasper, M. Lohn, B. Lauterbach, J. Menne, C. Lindschau, F. Mende, F.C. Luft, A. Schedl, H. Haller, T.V. Kurzchalia, Loss of caveolae, vascular dysfunction, and pulmonary defects in caveolin-1 gene-disrupted mice, *Science* 293 (2001) 2449–2452.
- [176] F. Galbiati, J.A. Engelman, D. Volonte, X.L. Zhang, C. Minetti, M. Li, H. Hou, B. Kneitz, W. Edelmann, M.P. Lisanti, Caveolin-3 null mice show a loss of caveolae, changes in the microdomain distribution of the dystrophin-glycoprotein complex, and t-tubule abnormalities, *J. Biol. Chem.* 276 (2001) 21425–21433.
- [177] D.S. Park, S.E. Woodman, W. Schubert, A.W. Cohen, P.G. Frank, M. Chandra, J. Shirani, B. Razani, B. Tang, L.A. Jelicics, S.M. Factor, L.M. Weiss, H.B. Tanowitz, M.P. Lisanti, Caveolin-1/3 double-knockout mice are viable, but lack both muscle and non-muscle caveolae, and develop a severe cardiomyopathic phenotype, *Am. J. Pathol.* 160 (2002) 2207–2217.
- [178] B. Nolwenn, D. Isabelle, L.L. Oazig, Cavin proteins: new players in the caveolae field, *Biochimie* 93 (2011) 71–77.
- [179] P. Jansa, S.W. Mason, U. Hoffmann-Rohrer, I. Grummt, Cloning and functional characterization of PTRF, a novel protein which induces dissociation of paused ternary transcription complexes, *EMBO J.* 17 (1998) 2855–2864.
- [180] L. Liu, P.F. Pilch, A critical role of cavin (polymerase I and transcript release factor) in caveolae formation and organization, *J. Biol. Chem.* 283 (2007) 4314–4322.
- [181] K.L. Inder, Y.Z. Zheng, M.J. Davis, H. Moon, D. Loo, H. Nguyen, J.A. Clements, R.G. Parton, L.J. Foster, M.M. Hill, Expression of PTRF in PC-3 Cells modulates cholesterol dynamics and the actin cytoskeleton impacting secretion pathways, *Mol. Cell. Proteomics* 11 (2012), <http://dx.doi.org/10.1074/mcp.M111.012245>.
- [182] R. Burgener, M. Wolf, T. Ganz, M. Baggiolini, Purification and characterization of a major phosphatidylserine-binding phosphoprotein from human platelets, *Biochem. J.* (1990) 729–734.
- [183] A. Hayer, M. Stoeber, C. Bissig, A. Helenius, Biogenesis of caveolae: stepwise assembly of large caveolin and cavin complexes, *Traffic* 11 (2010) 361–382.
- [184] S. Gustincich, P. Vatta, S. Goruppi, M. Wolf, S. Saccone, G. Della Valle, M. Baggiolini, C. Schneider, The human serum deprivation response gene (SDPR) maps to 2q32-q33 and codes for a phosphatidylserine-binding protein, *Genomics* (1999) 120–129.
- [185] C. Mineo, Y.S. Ying, C. Chapline, S. Jaken, R.G. Anderson, Targeting of protein kinase C α to caveolae, *J. Cell. Biol.* 141 (1998) 601–610.
- [186] C.G. Hansen, N.A. Bright, G. Howard, B.J. Nichols, SDPR induces membrane curvature and functions in the formation of caveolae, *Nat. Cell. Biol.* 11 (2009) 807–814.
- [187] C.G. Hansen, B.J. Nichols, Exploring the caves: cavins, caveolins and caveolae, *Trends Cell. Biol.* 20 (2010) 177–186.
- [188] W. Romer, L. Berland, V. Chambon, K. Gaus, B. Windschiegel, D. Tenza, M.R. Aly, V. Fraissier, J.C. Florent, D. Perrais, C. Lamaze, G. Raposo, C. Steinem, P. Sens, P. Bassereau, L. Johannes, Shiga toxin induces tubular membrane invaginations for its uptake into cells, *Nature* 450 (2007) 670–675.
- [189] M. Bauer, L. Pelkmans, A new paradigm for membrane-organizing and –shaping scaffolds, *FEBS Lett.* 580 (2006) 5559–5564.
- [190] J. Zimmerberg, M.M. Kozlov, How proteins produce cellular membrane curvature, *Nat. Rev. Mol. Cell. Biol.* 7 (2006) 9–19.
- [191] Y. Izumi, S. Hirai, Y. Tamai, A. Fujise-Matsuoka, Y. Nishimura, S. Ohno, A protein kinase C δ -binding protein SRBC whose expression is induced by serum starvation, *J. Biol. Chem.* 272 (1997) 7381–7389.
- [192] X.L. Xu, L.C. Wu, F. Du, A. Davis, M. Peyton, Y. Tomizawa, A. Maitra, G. Tomlinson, A.F. Gazdar, B.E. Weissman, A.M. Bowcock, R. Baer, J.D. Minna, Inactivation of human SRBC, located within the 11p15.5–p15.4 tumor suppressor region, in breast and lung cancers, *Cancer Res.* 61 (2001) 7943–7949.
- [193] C. Minetti, M. Bado, P. Broda, F. Sotgia, C. Bruno, F. Galbiati, D. Volonte, G. Lucania, A. Pavan, E. Bonilla, M.P. Lisanti, G. Cordone, Impairment of caveolae formation and T-system disorganization in human muscular dystrophy with caveolin-3 deficiency, *Am. J. Pathol.* 160 (2002) 265–270.
- [194] T. Ogata, T. Ueyama, K. Isodono, M. Tagawa, N. Takehara, T. Kawashima, K. Harada, T. Takahashi, T. Shioi, H. Matsubara, H. Oh, MURC, a muscle-restricted coiled-coil protein that modulates the Rho/ROCK pathway, induces cardiac dysfunction and conduction disturbance, *Mol. Cell. Biol.* 28 (2008) 3424–3436.
- [195] M. Tagawa, T. Ueyama, T. Ogata, N. Takehara, N. Nakajima, K. Isodono, S. Asada, T. Takahashi, H. Matsubara, H. Oh, MURC, a muscle-restricted coiled-coil protein, is involved in the regulation of skeletal myogenesis, *Am. J. Physiol. Cell. Physiol.* 295 (2008) 490–498.
- [196] B. Thisse, C. Thisse, Fast Release Clones: a High Throughput Expression Analysis, 2004 (ZFIN Direct Data Submission).
- [197] M. Bastiani, L. Liu, M.M. Hill, M.P. Jedrychowski, S.J. Nixon, H.P. Lo, D. Abankwa, R. Luetterforst, M. Fernandez-Rojo, M.R. Breen, S.P. Gygi, J. Vinten, P.J. Walser, K.N. North, J.F. Hancock, P.F. Pilch, R.G. Parton, MURC/Cavin-4 and cavin family members form tissue-specific caveolar complexes, *J. Cell. Biol.* 185 (2009) 1259–1273.
- [198] L. Liu, H.X. Xu, W.Q. Wang, C.T. Wu, T. Chen, Y. Qin, C. Liu, J. Xu, J. Long, B. Zhang, Y.F. Xu, Q.X. Ni, M. Li, X.J. Yu, Cavin-1 is essential for the tumor-promoting effect of caveolin-1 and enhances its prognostic potency in pancreatic cancer, *Oncogene* (2013 Jun 17), <http://dx.doi.org/10.1038/nc.2013.223>.
- [199] E. Tahanian, L.A. Sanchez, T.C. Shiao, R. Roy, B. Annabi, Flavonoids targeting of I κ B phosphorylation abrogates carcinogen-induced MMP-9 and COX-2 expression in human brain endothelial cells, *Drug. Des. Dev. Ther.* 5 (2011) 299–309.
- [200] E. Tahanian, S. Peiro, B. Annabi, Low intracellular ATP levels exacerbate carcinogen-induced inflammatory stress response and inhibit in vitro tubulogenesis in human brain endothelial cells, *J. Inflamm. Res.* 4 (2011) 1–10.
- [201] C.S. Aung, M.M. Hill, M. Bastiani, R.G. Parton, M.O. Parat, PTRF-cavin-1 expression decreases the migration of PC3 prostate cancer cells: role of matrix metalloproteinase 9, *Eur. J. Cell. Biol.* 90 (2011) 136–142.
- [202] M.L. Gould, G. Williams, H.D. Nicholson, Changes in caveolae, caveolin, and polymerase I and transcript release factor (PTRF) expression in prostate cancer progression, *Prostate* 70 (2010) 1609–1621.
- [203] J.S. Yi, D.G. Mun, H. Lee, J.S. Park, J.W. Lee, J.S. Lee, S.J. Kim, B.R. Cho, S.W. Lee, Y.G. Ko, PTRF/Cavin-1 is essential for multidrug resistance in cancer cells, *J. Proteome Res.* 12 (2013) 605–614.
- [204] A. Gámez-Pozo, I. Sánchez-Navarro, E. Calvo, M.T. Agulló-Ortuño, R. López-Vacas, E. Díaz, E. Camafrita, M. Nistal, R. Madero, E. Espinosa, J.A. López, J.A. Fresno Vara, PTRF/cavin-1 and MIF proteins are identified as non-small cell lung cancer biomarkers by label-free proteomics, *PLoS One* 7 (2012) e33752–e33761.
- [205] D. Volonte, F. Galbiati, Polymerase I and transcript release factor (PTRF)/cavin-1, a novel regulator of stress-induced premature senescence, *J. Biol. Chem.* 286 (2011) 28657–28661.
- [206] P. Verma, A.G. Ostermeyer-Fay, D.A. Brown, Caveolin-1 induces formation of membrane tubules that sense actomyosin tension and are inhibited by polymerase I and transcript release factor/cavin-1, *Mol. Biol. Cell.* 21 (2010) 2226–2240.
- [207] J.H. Lee, D.S. Byun, M.G. Lee, B.K. Ryu, M.J. Kang, K.S. Chae, K.Y. Lee, H.J. Kim, H. Park, S.G. Chi, Frequent epigenetic inactivation of hSRBC in gastric cancer and its implication in attenuated p53 response to stresses, *Int. J. Cancer* 122 (2008) 1573–1584.
- [208] S. Zochbauer-Muller, K.M. Fong, J. Geradts, X. Xu, S. Seidl, A. End-Pfutzenreuter, G. Lang, G. Heller, C.C. Zielinski, A.F. Gazdar, J.D. Minna, Expression of the candidate tumor suppressor gene hSRBC is frequently lost in primary lung cancers with and without DNA methylation, *Oncogene* 24 (2005) 6249–6255.
- [209] L. Bai, X. Deng, Q. Li, M. Wang, W. An, A. Deli, Z. Gao, Y. Xie, Y. Dai, Y.S. Cong, Down-regulation of the cavin family proteins in breast cancer, *J. Cell. Biochem* 113 (2012) 322–328.
- [210] H. Lee, D. Volonte, F. Galbiati, P. Iyengar, D.M. Lublin, D.B. Bregman, M.T. Wilson, R. Campos-Gonzalez, B. Bouzahzah, R.G. Pestell, P.E. Scherer, M.P. Lisanti, Constitutive and growth factor-regulated phosphorylation of caveolin-1 occurs at the same site (Tyr-14) in vivo: identification of a c-Src/Cav-1/Grb7 signaling cassette, *Mol. Endocrinol.* 14 (2000) 1750–1775.
- [211] H. Lee, D.S. Park, X.B. Wang, P.E. Scherer, P.E. Schwartz, M.P. Lisanti, Src-induced phosphorylation of caveolin-2 on tyrosine 19. Phospho-caveolin-2 [Tyr(P)19] is localized near focal adhesions, remains associated with lipid rafts/caveolae, but no longer forms a high molecular mass hetero-oligomer with caveolin-1, *J. Biol. Chem.* 277 (2002) 34556–34567.
- [212] S.A. Tahir, G. Yang, A.A. Goltsov, M. Watanabe, K. Tabata, J. Addai, M.A. Fattah el, D. Kadmon, T.C. Thompson, Tumor cell-secreted caveolin-1 has proangiogenic activities in prostate cancer, *Cancer Res.* 68 (2008) 731–739.
- [213] H. Ju, R. Zou, V.J. Venema, R.C. Venema, Direct interaction of endothelial nitric-oxide synthase and caveolin-1 inhibits synthase activity, *J. Biol. Chem.* 272 (1997) 18522–18525.
- [214] Z. Lu, S. Ghosh, Z. Wang, T. Hunter, Downregulation of caveolin-1 function by EGF leads to the loss of E-cadherin, increased transcriptional activity of β -catenin, and enhanced tumor cell invasion, *Cancer Cell.* 4 (2003) 499–515.
- [215] G. Ayala, M. Morello, A. Frolov, S. You, R. Li, F. Rosati, G. Bartolucci, G. Danza, R.M. Adam, T.C. Thompson, M.P. Lisanti, M.R. Freeman, D. Di Vizio, Loss of caveolin-1 in prostate cancer stroma correlates with reduced relapse-free survival and its functionally relevant to tumor progression, *J. Pathol.* 231 (2013) 77–87.
- [216] C. Meyer, Y. Liu, A. Kaul, I. Peipe, S. Dooley, Caveolin-1 abrogates TGF β -mediated hepatocyte apoptosis, *Cell. Death Dis.* 4 (2013) e466, <http://dx.doi.org/10.1038/cddis.2012.204>.
- [217] K. Quann, D.M. Gonzales, I. Mercier, C. Wang, F. Sotgia, R.G. Pestell, M.P. Lisanti, J.F. Jasmin, Caveolin-1 is a negative regulator of tumor growth in glioblastoma and modulates, *Cell. Cycle* 12 (2013) 1510–1520.
- [218] C. Trimmer, G. Bonuccelli, S. Katiyar, F. Sotgia, R.G. Pestell, M.P. Lisanti, F. Capozza, Cav1 suppresses tumor growth and metastasis in a murine model of cutaneous SCC through modulation of MAPK/AP-1 activation, *Am. J. Pathol.* 182 (2013) 992–1004.
- [219] A. Puyraimond, R. Fridman, M. Lemesle, B. Arbeille, S. Menashi, MMP-2 colocalizes with caveolae on the surface of endothelial cells, *Exp. Cell. Res.* 262 (2001) 28–36.

- [220] C. Pramudji, S. Shimura, S. Ebara, G. Yang, J. Wang, C. Ren, Y. Yuan, S.A. Tahir, T.L. Timme, T.C. Thompson, In situ prostate cancer gene therapy using a novel adenoviral vector regulated by the caveolin-1 promoter, *Clin. Cancer Res.* 7 (2001) 4272–4279.
- [221] J.P. Gratton, M.I. Lin, J. Yu, E.D. Weiss, Z.L. Jiang, T.A. Fairchild, Y. Iwakiri, R. Groszmann, K.P. Claffey, Y.C. Cheng, W.C. Sessa, Selective inhibition of tumor microvascular permeability by cavtratin blocks tumor progression in mice, *Cancer Cell* 4 (2003) 31–39.
- [222] M. Hamoudane, S. Maffioli, R. Cordera, D. Maggi, B. Salani, Caveolin-1 and polymerase I and transcript release factor: new players in insulin-like growth factor-I receptor signaling, *J. Endocrinol. Invest* 36 (2013) 204–208.
- [223] H. Moon, C.S. Lee, K.L. Inder, S. Sharma, E. Choi, D.M. Black, K.A. Lê Cao, C. Winterford, J.I. Coward, M.T. Ling, D.J. Craik, R.G. Parton, P.J. Russell, M.M. Hill, PTRF/cavin-1 neutralizes non-caveolar caveolin-1 microdomains in prostate cancer, *Oncogene* 33 (27) (2014) 3561–3570.
- [224] K.L. Inder, Y.Z. Zheng, M.J. Davis, H. Moon, D. Loo, H. Nguyen, J.A. Clements, R.G. Parton, L.J. Foster, M.M. Hill, Expression of PTRF in PC-3 cells modulates cholesterol dynamics and the actin cytoskeleton impacting secretion pathways, *Mol. Cell. Proteomics* 11 (2) (2012 Feb). M111.012245.
- [225] M.M. Hill, N.H. Daud, C.S. Aung, D. Loo, S. Martin, S. Murphy, D.M. Black, R. Barry, F. Simpson, L. Liu, P.F. Pilch, J.F. Hancock, M.O. Parat, R.G. Parton, Co-regulation of cell polarization and migration by caveolar proteins PTRF/Cavin-1 and caveolin-1, *PLoS One* 7 (2012) e43041.
- [226] X. Li, Z. Jia, Y. Shen, H. Ichikawa, J. Jarvik, R.G. Nagele, G.S. Goldberg, Coordinate suppression of Sdpr and Fhl1 expression in tumors of the breast, kidney, and prostate, *Cancer Sci.* 99 (2008) 1326–1333.
- [227] Jin-Hee Lee, Min-Ju Kang, Hye-Yeon Han, Min-Goo Lee, Seong-In Jeong, Byung-Kyu Ryu, Tae-Kyu Ha, Nam-Goo Her, Jikhyon Han, Sun Jin Park, Kil Yeon Lee, Hyo-Jong Kim, Sung-Gil Chi, Epigenetic alteration of *PRKCDDBP* in colorectal cancers and its implication in tumor cell resistance to TNF α -induced apoptosis, *Clin. Cancer Res.* 17 (December, 2011) 7551.
- [228] X.L. Xu, L.C. Wu, F. Du, A. Davis, M. Peyton, Y. Tomizawa, Inactivation of human SRBC, located within the 11p15.5-p15.4 tumor suppressor region, in breast and lung cancers, *Cancer Res.* 61 (2001) 7943–7949.
- [229] J.H. Lee, D.S. Byun, M.G. Lee, B.K. Ryu, M.J. Kang, K.S. Chae, Frequent epigenetic inactivation of hSRBC in gastric cancer and its implication in attenuated p53 response to stresses, *Int. J. Cancer* 122 (2008) 1573–1584.
- [230] Y. Ogawara, S. Kishishita, T. Obata, Y. Isazawa, T. Suzuki, K. Tanaka, Akt enhances Mdm2-mediated ubiquitination and degradation of p53, *J. Biol. Chem.* 277 (2002) 21843–21850.
- [231] S. Paskas, J. Janković, I. Marečko, T.I. Dencić, S. Statić, D. Cvejić, S. Savin, Caveolin-1 expression in papillary thyroid carcinoma: correlation with clinicopathological parameters and BRAF mutation status, *Otolaryngol. Head. Neck Surg.* 150 (2014) 201–209.
- [232] V.D. Eynden, Overexpression of caveolin-1 and -2 in cell lines and in human samples of inflammatory breast cancer, *Breast Cancer Res. Treat.* 95 (2005) 219–228.
- [233] F. Bianchi, J. Hu, G. Pelosi, R. Cirincione, M. Ferguson, C. Ratcliffe, P. Paolo, D. Fiore, K. Gatter, F. Pezzella, U. Pastorino, Lung cancers detected by screening with spiral computed tomography have a malignant phenotype when analyzed by cDNA microarray, *Clin. Cancer Res.* 10 (2004) 6023–6028.
- [234] H.N. Choi, K.R. Kim, H.S. Park, K.Y. Jang, M.J. Kang, D.G. Lee, Y.K. Kim, B.H. Cho, E.J. Cha, W.S. Moon, Expression of caveolin in hepatocellular carcinoma: association with unpaired artery formation and radiologic findings, *Korean J. Hepatol.* 13 (2007) 396–408.
- [235] G. Yang, A. Goltsov, C. Ren, S. Kurosaka, K. Edamura, R. Logothetis, F. DeMayo, P. Troncoso, J. Blando, J. DiGiovanni, T. Thompson, Caveolin-1 upregulation contributes to c-Myc-induced high-grade prostatic intraepithelial neoplasia and prostate cancer, *Mol. Cancer Res.* 10 (2012) 218–229.
- [236] Z.D. Nassar, H. Moon, T. Duong, L. Neo, M.M. Hill, M. Francois, R.G. Parton, M.O. Parat, PTRF/Cavin-1 decreases prostate cancer angiogenesis and lymphangiogenesis, *Oncotarget* 4 (2013) 1844–1855.
- [237] S.W. Fine, M.P. Lisanti, F. Galbiati, M. Li, Elevated expression of caveolin-1 in adenocarcinoma of the colon, *Am. J. Clin. Pathol.* 115 (2001) 719–724.
- [238] F. Sotgia, M.C. Casimiro, G. Bonuccelli, M. Liu, D. Whitaker-Menezes, O. Er, K.M. Daumer, I. Mercier, A.K. Witkiewicz, C. Minetti, F. Capozza, M. Gormley, A.A. Quong, H. Rui, P.G. Frank, J.N. Milliman, E.S. Knudsen, J. Zhou, C. Wang, R.G. Pestell, M.P. Lisanti, Loss of caveolin-3 induces a lactogenic microenvironment that is protective against mammary tumor formation, *Am. J. Pathol.* 174 (2009) 613–629.
- [239] S.Y. Tong, K.D. Ki, J.M. Lee, M.J. Kang, T.K. Ha, S.I. Chung, S.G. Chi, S.K. Lee, Frequent inactivation of hSRBC in ovarian cancers by promoter CpG island hypermethylation, *Acta Obstet. Gynecol. Scand.* 89 (2010) 629–635.
- [240] G. Perrone, V. Altomare, M. Zagami, S. Morini, T. Petitti, C. Battista, A. Onetti Muda, C. Rabitti, Caveolin-1 expression in human breast lobular cancer progression, *Mod. Pathol.* 22 (2009) 71–78.
- [241] L. Shi, Chen Li Wang, L. Zhang, Z. Chen, Expression of caveolin-1 in mucoidermoid carcinoma of the salivary glands: correlation with vascular endothelial growth factor, microvessel density, and clinical outcome, *Cancer* 109 (2007) 1523–1531.
- [242] L. Campbell, M. Gumbleton, D.F. Griffiths, Caveolin-1 overexpression predicts poor disease-free survival of patients with clinically confined renal cell carcinoma, *Br. J. Cancer* 89 (2003) 1909–1913.
- [243] P. Zhan, X.K. Shen, Q. Qian, Q. Wang, J.P. Zhu, Y. Zhang, H.Y. Xie, C.H. Xu, K.K. Hao, W. Hu, N. Xia, G.J. Lu, L.K. Yu, Expression of caveolin-1 is correlated with disease stage and survival in lung adenocarcinomas, *Oncol. Rep.* 27 (2012) 1072–1078.
- [244] D.S. Cho, H. Yim, K.S. Cho, S.J. Hong, N.H. Cho, S.I. Kim, H.S. Ahn, S.J. Kim, Impact of caveolin-1 expression on the prognosis of transitional cell carcinoma of the upper urinary tract, *J. Korean Med. Sci.* 23 (2008) 296–301.
- [245] J. Xue, H. Chen, L. Diao, X. Chen, D. Xia, Expression of Caveolin-1 in tongue squamous cell carcinoma by quantum dots, *Eur. J. Histochem.* 54 (2010) e20.
- [246] C.P. Tanase, S. Dima, M. Mihai, E. Raducan, M.I. Nicolescu, L. Albuiescu, B. Voiculescu, T. Dumitrascu, L.M. Cruceru, M. Leabu, I. Popescu, M.E. Hinescu, Caveolin-1 overexpression correlates with tumor progression markers in pancreatic ductal adenocarcinoma, *J. Mol. Histol.* 40 (2009) 23–29.
- [247] X.X. Wang, Z. Wu, H.F. Huang, C. Han, W. Zou, J. Liu, Caveolin-1, through its ability to negatively regulate TLR4, is crucial determinant of MAPK activation in LPS-challenged mammary epithelial cells, *Asian Pac. J. Cancer Prev.* 14 (2013) 2295–2299.
- [248] C. Boscher, I.R. Nabi, Galectin-3 and phosphor-caveolin-1 dependent outside-integrin signaling mediated EGF mitogenic response in mammary cancer cells, *Mol. Biol. Cell.* 24 (2013) 2134–2145.
- [249] J.H. Lee, M.J. Kang, H.Y. Han, M.G. Lee, S.I. Jeong, B.K. Ryu, T.K. Ha, N.G. Her, J. Han, S.J. Park, K.Y. Lee, H.J. Kim, S.G. Chi, Epigenetic alteration of *PRKCDDBP* in colorectal cancers and its implication in tumor cell resistance to TNF-induced apoptosis, *Clin. Cancer Res.* 17 (2011) 7551–7562.
- [250] T. Shatseva, D.Y. Lee, Z. Deng, B.B. Yang, MicroRNA miR-199a-3p regulates cell proliferation and survival by targeting caveolin-2, *J. Cell. Sci.* 124 (16) (2011) 2826–2836.
- [251] K.A. Wilson, S.A. Colavito, V. Schulz, P.H. Wakefield, W. Sessa, D. Tuck, D.F. Stern, NFB1/MDC1 regulates Cav1 and Cav2 independently of DNA damage and p53, *Mol. Cancer Res.* 9 (2011) 766–781.
- [252] L.N. Zhao, Z.H. Yang, Y.H. Liu, H.Q. Ying, H. Zhang, Y.X. Xue, Vascular endothelial growth factor increases permeability of the blood-tumor barrier via caveolae-mediated transcellular pathway, *J. Mol. Neurosci.* 44 (2011) 122–129.
- [253] E. Burgermeister, L. Tencer, M. Liscovitch, Peroxisome proliferator-activated receptor-gamma upregulates caveolin-1 and caveolin-2 expression in human carcinoma cells, *Oncogene* 22 (2003) 3888–3900.
- [254] F. Capozza, C. Trimmer, R. Castello-Cros, S. Katiyar, D. Whitaker-Menezes, A. Follenzi, M. Crosariol, G. Llaverias, F. Sotgia, R.G. Pestell, M.P. Lisanti, Genetic ablation of Cav1 differentially affects melanoma tumor growth and metastasis in mice: role of Cav1 in Shh heterotypic signaling and trans-endothelial migration, *Cancer Res.* 72 (2012) 2262–2274.
- [255] E.J. Kim, J.H. Sohn, M.K. Kim, S.W. Chae, H.S. Lee, E.Y. Cho, W.H. Kim, The loss of expression of caveolin-1 in gastrointestinal stromal tumors, *Korean J. Pathol.* 39 (2005) 338–344.