

Assignment 2
IB3702 Mathematics for Machine learning

Harman Singh

November 7, 2025

Problem 1

exercise a

$$Av = \Lambda v$$

First I am going to multiply A to the first column of Q

So $v_1 = [-a, 0, b]^T$ with eigenvalue 25.

This gets me the equation:

$$A \begin{pmatrix} -a \\ 0 \\ b \end{pmatrix} = 25 \begin{pmatrix} -a \\ 0 \\ b \end{pmatrix}$$

Now I calculate $Av_1 =$

$$Av_1 = \begin{pmatrix} -2 & 0 & -36 \\ 0 & -3 & 0 \\ -36 & 0 & -23 \end{pmatrix} \begin{pmatrix} -a \\ 0 \\ b \end{pmatrix}$$

Now we multiply horizontally on the left, with vertically on the right. So for example the element on the top left will be $(-2)(-a)$ and then $0(0)$ and then $-36(b)$ and so on.

$$\begin{pmatrix} (-2)(-a) & +0(0) & +(-36)(b) \\ 0(-a) & +(-3)(0) & +0(b) \\ (-36)(-a) & +0(0) & +(-23)(b) \end{pmatrix} = \begin{pmatrix} 2a - 36b \\ 0 \\ 36a - 23b \end{pmatrix}$$

Now I equate this to $25v_1$:

$$\begin{pmatrix} 2a - 36b \\ 0 \\ 36a - 23b \end{pmatrix} = 25 \begin{pmatrix} -a \\ 0 \\ b \end{pmatrix} = \begin{pmatrix} -25a \\ 0 \\ 25b \end{pmatrix}$$

And now I can just solve the equations for a and b :

$$-2a - 36b = -25a \implies 27a - 36b = 0 \implies 27a = 36b \implies b = \frac{27}{36}a = \frac{3}{4}a$$

$$36a - 23b = 25b \implies 36a - 48b = 0 \implies 36a = 48b \implies b = \frac{36}{48}a = \frac{3}{4}a$$

Since Q orthogonally diagonalizes A , the following must be true

$$(-a)^2 + (0)^2 + (b)^2 = 1 \implies a^2 + b^2 = 1$$

I can enter the value of b I found above into this equation so we get:

$$a^2 + \left(\frac{3}{4}a\right)^2 = 1 \implies a^2 + \frac{9}{16}a^2 = 1 \implies \frac{25}{16}a^2 = 1 \implies a^2 = \frac{16}{25} \implies a = \frac{4}{5}$$

Then enter this value of a into the equation for b :

$$b = \frac{3}{4}a = \frac{3}{4} \cdot \frac{4}{5} = \frac{12}{20} = \frac{3}{5}$$

$$\boxed{a = \frac{4}{5}, b = \frac{3}{5}}$$

exercise b

...

exercise c

...

exercise d

...

exercise e

...

Problem 2

exercise a

...

exercise b

...

exercise c

...

exercise d

...

exercise e

...

Problem 3

exercise a

...

exercise b

...

Problem 4

I construct the following matrix:

$$A = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 5 \end{pmatrix}$$

exercise a

I chose an upper triangular matrix, because in this type the determinant is easy to calculate, it's just the multiplication of the diagonal elements.

Here: $\det(A) = 1 \cdot 1 \cdot 5 = 5$

And I also made sure to use simple numbers so the calculations would be easier.

exercise b

(10) True, because the determinant is 5 (calculated in exercise a), $\neq 0$

(1) True, because a square matrix is invertible if the determinant is nonzero (our determinant is 5)

(7)

To calculate the RREF of A I perform the following operations:

$$\begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 5 \end{pmatrix} \xrightarrow{R_3 = \frac{1}{5} \cdot R_3} \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 3 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_2 = R_2 - 3R_3} \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \xrightarrow{R_1 = R_1 - 2R_2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
$$= \mathbb{I}_3$$

(3) True, because the null space only contains the zero vector when the RREF is the identity matrix, which we have shown above.

(4) True. From (3) we know that the null space only contains the zero vector, so the only solution to $A\vec{x} = \vec{0}$ is the trivial solution $\vec{x} = \vec{0}$

(6) The rank of a matrix is the number of pivot columns in the RREF and ours has 3 pivots, so $\text{rank}(A) = 3 = n$

(2) To prove this, we need to first find the existence of a solution, and then prove the uniqueness of it.

Existence of solution

Since A is invertible (see (1)), we can multiply both sides of the equation $A\vec{x} = \vec{b}$ with A^{-1} :

$$A^{-1}A\vec{x} = A^{-1}\vec{b} \implies \vec{x} = A^{-1}\vec{b}$$

So there exists a solution for each $\vec{b} \in \mathbb{R}^n$

Uniqueness of solution

If I assume there are two solutions $\vec{x}_1$ and $\vec{x}_2$ so that:

$$A\vec{x}_1 = \vec{b} \quad \text{and} \quad A\vec{x}_2 = \vec{b}$$

Then that would mean that:

$$A(\vec{x}_1 - \vec{x}_2) = \vec{0}$$

From (4) we know that the only solution to this equation is the trivial solution $\vec{x}_1 - \vec{x}_2 = \vec{0}$, so $\vec{x}_1 = \vec{x}_2$

(5)

- (5.1) True, the columns are linearly independent when the RREF is the identity matrix (see above)
- (5.2) True, because the columns of A span $\mathbb{R}^n$ when A is invertible (see (1))

(8) The transpose of A is:

$$A^T = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 3 & 5 \end{pmatrix}$$

The determinant of A^T is also 5 ($1 \times 1 \times 5 = 5$), and a square matrix is invertible when the determinant is nonzero, so A^T is invertible.

(9) The rows of A are the columns of A^T , and since I showed in (5) that the columns of a matrix form a basis for $\mathbb{R}^n$ when the matrix is invertible, the rows of A also form a basis for $\mathbb{R}^n$

Problem 5

...

...