



PS Space Charge Simulations

Updates

haroon.rafique@cern.ch

August 22, 2019

Table of Contents

MD4224 Injection Bump

PS Transfer

MD4224 Injection Bump

PS Transfer

Methods Explored

At each of the four sections (40, 42, 43, 44) in the PS we use:

- ▶ Two 10 cm HKICKER with MULTIPOLE inbetween for the sextupole component.
- ▶ A single zero-length MULTIPOLE with dipolar and sextupolar components.
- ▶ A single 20 cm (zero gradient) QUADRUPOLE with dipolar and sextupolar components added as errors.
- ▶ A single 20 cm (zero angle) SBEND with dipolar and sextupolar components added as errors.

Issues

- ▶ Make sure to update PTC and PyORBIT every turn.
- ▶ Include calls to correct PTC files (energize lattice, update twiss, read final settings etc).
- ▶ Unknown environment bug on LXPlus and HPC-Batch: causes simulation to run without updating PTC every turn - for unknown reason. Now seems to work - not satisfied with not knowing more - will keep an eye out for future weirdness.
- ▶ Dipole component needed to be normalised when used in QUADRUPOLE and SBEND methods.

Horizontal Closed Orbit Maximum and Minimum

Extrema of the closed orbit as a function of turn.




Figure: PyORBIT Horizontal Orbit Extrema: Only the negative extrema for the split HKICKER method is different.

Injection Bump

- ▶ All single element methods agree, only the split HKICKER method is different.
- ▶ Tunes however are not in agreement.

Note: all simulations run at the most extreme vertical working point of $(Q_x, Q_y) = (6.21, 6.10)$.

Split HKICKER + MULTIPOLE




Figure: Horizontal Orbit Extrema for split HKICKER + MULTIPOLE: PyORBIT and MADX agree.

Split HKICKER + MULTIPOLE





Figure: Tunes for split HKICKER + MULTIPOLE: magnitude and Q_y sign incorrect.




Figure: Horizontal Orbit Extrema for MULTIPOLE: MADX and PTC show no change, however effect is visible in PyORBIT.

MULTIPOLE





Figure: Tunes for MULTIPOLe: magnitude correct, sign opposite from expected, no change in MADX or PTC.

QUADRUPOLE




Figure: Horizontal Orbit Extrema for QUADRUPOLE: No bump in MADX, PTC acting strangely, PyORBIT as expected

QUADRUPOLE





Figure: Tunes for QUADRUPOLE: MADX and PTC disagree, PyORBIT magnitude and sign incorrect.




Figure: Horizontal Orbit Extrema for SBEND: unexplained behaviour in MADX and PTC, PyORBIT as expected.





Figure: Tunes for SBEND: MADX and PTC agree but magnitude is large, PyORBIT magnitude incorrect but sign ok.

Test Conversion from MADX Table to PyORBIT Table

- ▶ As MULTIPOLE method seems almost correct (tunes are incorrect sign but good magnitude) check for sign error in tables.
- ▶ As Split HKICKER + MULTIPOLE tunes both increase (expect Q_x to increase but Q_y to decrease) check for sign error in tables.
- ▶ As Split HKICKER + MULTIPOLE tunes are \approx half magnitude expected, check if factorial should not be included.
- ▶ Abandon SBEND and QUADRUPOLE methods for now due to unexpected behaviours and disagreements between PTC and MADX.

MULTIPOLE Sign Check

Multiply all sextupole components by -1. Bump unchanged.





Figure: Tunes for MULTIPOLE Sign Check: Magnitudes changed unexpectedly - too large.

Split HKICKER + MULTIPOLE Sign Check

Multiply all sextupolar components (i.e. only the MULTIPOLE) by -1. Change in bump negligible.





Figure: Tunes for Split HKICKER + MULTIPOLE Sign Check: both tunes flipped.



Split HKICKER + MULTIPOLE Factorial Check

Do not include the factorial in sextupolar components (i.e. only the MULTIPOLE).
Bump unchanged.





Figure: Tunes for Split HKICKER + MULTIPOLE Factorial Check: Some improvement in magnitude.

Injection Bump Conclusions

- ▶ Split HKICKER + MULTIPOLE: Bump OK, tune swing small, Q_y incorrect sign.
- ▶ MULTIPOLE: Bump OK, tune swing correct magnitude, tunes incorrect sign.
- ▶ QUADRUPOLE: Bump OK, tune swing small, tunes incorrect sign.
- ▶ SBEND: Bump OK, tune swing small, tunes correct sign.
- ▶ Tests show that multipole method is perhaps the closest to the desired tune swing
 - only when flipping sign on sextupolar components.

MD4224 Injection Bump

PS Transfer

Goals

- ▶ To investigate the space charge contribution to beam envelope oscillations observed in the PS immediately after injection.
- ▶ Reproduce measurements performed using BGI and SEM Grids in PTC-PyORBIT in order to understand the contribution of space charge.

Current Plan

- ▶ Start with matched distribution and track without space charge.
- ▶ Add dispersion mismatch as constant factor (will obtain parameterised values from Selim/Matthew soon).
- ▶ Start with PTC TWISS to match distribution, later use end of transfer line values transported to BGI/SEM position (to be provided by Selim/Matthew).

Note: all simulations run at the nominal working point of $(Q_x, Q_y) = (6.21, 6.24)$.

Matched Distribution

No Space Charge

Mismatch of factor 0% and +10%.

- ▶ Had to correct units of dE (factor 1E3).
- ▶ Matching not correct.
- ▶ Abandon and use tomo distribution.

Matched Distribution Longitudinal Motion




Figure: Longitudinal motion for matched distribution: oscillations larger than expected - indicates mismatch.

Matched Distribution Horizontal Dispersion





Figure: Horizontal dispersion for matched distribution: 10% mismatch identical to 0% mismatch - indicates existing mismatch already larger than 10%.

Matched Distribution Emittances





Figure: Emittances for matched distribution: 10% mismatch identical to 0% mismatch - indicates existing mismatch already larger than 10%.

No Space Charge

Mismatch of factor 0%, +10%, +20%, and +30%.

Tomo Distribution Longitudinal Motion (No Space Charge)




Figure: Longitudinal motion for `tomo` distribution: oscillation magnitude as expected.

Tomo Distribution Horizontal Dispersion (No Space Charge)




Figure: Horizontal dispersion for tomo distribution: oscillation magnitude is a function of mismatch as expected.

Tomo Distribution Emittances (No Space Charge)




Figure: Emittances for tomo distribution: horizontal emittance growth is a function of dispersion mismatch.

With Space Charge

Mismatch of factor +10%, +20%, and +30%.

Tomo Distribution Longitudinal Motion (With Space Charge)




Figure: Longitudinal motion for tomo distribution with space charge: oscillation has opposite sign to no space charge case - SC drives bunch lengthening at the start?

Tomo Distribution Horizontal Dispersion (With Space Charge)




Figure: Horizontal dispersion for tomo distribution with space charge: dispersion oscillations have longer period, highest mismatch in anti-phase rather than larger magnitude.

Tomo Distribution Emittances (With Space Charge)




Figure: Emittances for tomo distribution with space charge: clear emittance exchange.



www.cern.ch