

Package ‘StatEngine’

July 6, 2020

Type Package
Title An R Package for the UofSC Course STAT 509 ``Statistics for Engineers"
Version 1.0.0
Author Dewei Wang
Maintainer Dewei Wang <deweiwang@stat.sc.edu>
Description This R package provides tools for methods covered in the course STAT 509 ``Statistics for Engineers" at the University of South Carolina
Depends R (>= 3.1.0),
car (>= 3.0.0)
License MIT
Encoding UTF-8
LazyData true

R topics documented:

Continuous Distributions	1
Counting Techniques	4
Descriptive Statistics	6
Discrete Distributions	7
One-Sample Confidence Intervals	10
One-Sample Tests	12
Index	15

Continuous Distributions	
<i>Continuous distributions</i>	

Description

Plot pdf/cdf, calculate mean/variance/standard deviation, and compute probabilities for various continuous distributions

Usage

```

# continuous Uniform Distribution
uniform.summary(a,b,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
uniform.prob(a,b,lb,ub)
uniform.quantile(a,b,q)

# Normal distribution
normal.summary(mu,sigma,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
normal.prob(mu,sigma,lb,ub)
normal.quantile(mu,sigma,q)

# Exponential distribution
exponential.summary(lambda,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
exponential.prob(lambda,lb,ub)
exponential.quantile(lambda,q)

# Gamma distribution
gamma.summary(r,lambda,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
gamma.prob(r,lambda,lb,ub)
gamma.quantile(r,lambda,p)

# Weibull distribution
weibull.summary(beta,delta,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
weibull.prob(beta,delta,lb,ub)
weibull.quantile(beta,delta,p)

# Lognormal distribution
lognormal.summary(theta,omega,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
lognormal.prob(theta,omega,lb,ub)
lognormal.quantile(theta,omega,p)

#Beta distribution
beta.summary(alpha,beta,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
beta.prob(alpha,beta,lb,ub)
beta.quantile(alpha,beta,p)

```

Arguments

plotpdf	TRUE or FALSE, if TRUE, it plots the pmf
plotcdf	TRUE or FALSE, if TRUE, it plots the cdf
lb,ub	lower bound (lb) and upper bound (ub) in a probability statement; lb could be -Inf; ub could be Inf; ub cannot be less than lb
a,b	lower bound and upper bound of the support of a continuous uniform distribution
mu,sigma	mean and standard deviation of a normal distribution
lambda	parameter of an exponential distribution
r,lambda	shape and scale parameters of a gamma distribution
beta,delta	shape and scale parameters of a Weibull distribution
theta,omega	shape and scale parameters of a lognormal distribution
alpha,beta	parameters of a beta distribution

Details

Plot the probability density function (pdf) and the cumulative distribution function (cdf) and calculate probability, mean, variance and standard deviation, and compute probabilities of various discrete distributions (continuous uniform distribution, normal distribution, exponential distribution, gamma distribution, Weibull distribution, lognormal distribution, beta distribution).

Value

<code>uniform.summary</code>	a list of the mean, variance, and standard deviation of a continuous uniform distribution, plot of the pdf/cdf or not
<code>uniform.prob</code>	probability of X between lb and ub based on a continuous uniform distribution
<code>uniform.quantile</code>	quantile of a continuous uniform distribution
<code>normal.summary</code>	a list of the mean, variance, and standard deviation of a normal distribution, plot of the pdf/cdf or not
<code>normal.prob</code>	probability of X between lb and ub based on a normal distribution
<code>normal.quantile</code>	quantile of a normal distribution
<code>exponential.summary</code>	a list of the mean, variance, and standard deviation of an exponential distribution, plot of the pdf/cdf or not
<code>exponential.prob</code>	probability of X between lb and ub based on an exponential distribution
<code>exponential.quantile</code>	quantile of an exponential distribution
<code>gamma.summary</code>	a list of the mean, variance, and standard deviation of a gamma distribution, plot of the pdf/cdf or not
<code>gamma.prob</code>	probability of X between lb and ub based on a gamma distribution
<code>gamma.quantile</code>	quantile of a gamma distribution
<code>weibull.summary</code>	a list of the mean, variance, and standard deviation of a Weibull distribution, plot of the pdf/cdf or not
<code>weibull.prob</code>	probability of X between lb and ub based on a Weibull distribution
<code>weibull.quantile</code>	quantile of a Weibull distribution
<code>lognormal.summary</code>	a list of the mean, variance, and standard deviation of a lognormal distribution, plot of the pdf/cdf or not
<code>lognormal.prob</code>	probability of X between lb and ub based on a lognormal distribution
<code>lognormal.quantile</code>	quantile of a lognormal distribution
<code>beta.summary</code>	a list of the mean, variance, and standard deviation of a beta distribution, plot of the pdf/cdf or not
<code>beta.prob</code>	probability of X between lb and ub based on a beta distribution
<code>beta.quantile</code>	quantile of a beta distribution

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Chapter 4 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
# Continuous uniform distribution
a=4.9;b=5.1;uniform.summary(a,b)
uniform.prob(a,b,4.95,5) # P(4.95<X<5)
uniform.quantile(a,b,0.9) # x such that P(X>x)=0.1

#Normal distribution
normal.summary(10,2) #mu=10,sigma=2,variance=4
normal.prob(10,2,9,11)
normal.quantile(10,2,0.98)

#Exponential distribution
exponential.summary(25) #lambda=25
exponential.prob(25,0.1,Inf) #P(X>0.1)
exponential.quantile(25,0.1) # x such that P(X>x)=0.9

#Gamma distribution
gamma.summary(10,.5) #r=10,lambda=0.5
gamma.prob(10,0.5,25,Inf) #P(X>25)
gamma.quantile(10,0.5,0.95) # x such that P(X<x)=0.95

#Weibull distribution
weibull.summary(2,5000) #beta=2,delta=5000
weibull.prob(2,5000,6000,Inf) #P(X>6000)
weibull.quantile(2,5000,1-0.05) # x such that P(X>x)=0.05

#Lognormal distribution
lognormal.summary(10,1.5) #theta=10,omega=1.5
lognormal.prob(10,1.5,10000,Inf) #P(X>10000)
lognormal.quantile(10,1.5,1-0.99) # x such that P(X>x)=0.99

#Beta distribution
beta.summary(2.5,1) #alpha=2.5, beta=1
beta.prob(2.5,1,0.7,Inf) # P(X>0.7)
beta.quantile(2.5,1,0.99) # x such that P(X<x)=0.99
```

Description

Use permutations and combinations to count outcomes

Usage

`nPr(n,r)`: permute r items from a set of n distinct items

`nCr(n,r)`: select r items from a set of n items where order does not matter

`SimPerm(n_vec)`: permute n items when they are not totally distinct

Arguments

<code>n</code>	total n itmes
<code>r</code>	select r
<code>n_vec</code>	a vector of (n_1, n_2, \dots, n_r)

Details

Permutation, Combination, and Permutation of similar items

Value

`nPr(n,r)` provides the number of different ways to permute r items from a set of n distinct items.

`nCr(n,r)` provides the number of different ways to select r items from a set of n items where order does not matter.

`SimPerm(n_vec)` provides the number of different ways to permute n items when they are not totally distinct.

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Section 2.2 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
nPr(8,4)
nCr(3,2)
nCr(47,4)
SimPerm(c(2,3,2))
```

Descriptive Statistics*Descriptive Statistics*

Description

Calculate and plot descriptive statistics

Usage

```
data.summary(x)
```

Arguments

x	the sample
plot	True or False; plot or not the stem and leaf diagram, scatterplot, histogram, boxplot, QQ-plot

Details

Calculate various numerical descriptive statistics and plot the stem and leaf diagram, scatterplot, histogram, boxplot, QQ-plot

Value

data.summary	a table of numerical descriptive statistics (std stands for standard deviation), a stem and leaf diagram, and a figure consisting of scatterplot, histogram, boxplot, QQ-plot of the sample
--------------	---

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Chapter 6 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
x=scan("https://raw.githubusercontent.com/Harrindy/StatEngine/master/Data/CompressiveStrength.csv")
data.summary(x)
```

Discrete Distributions*Discrete distributions*

Description

Plot pmf/cdf, calculate mean/variance/standard deviation, and compute probabilities for various discrete distributions

Usage

```
# For use-defined discrete distribution
discrete.plotpdf(x,fx)
discrete.plotcdf(x,fx)
discrete.summary(x,fx,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
discrete.prob(x,fx,lb)
discrete.prob(x,fx,lb,ub,inclusive=c("none","left","right","both"))

# Discrete Uniform Distribution
duniform.summary(range,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
duniform.prob(range,lb)
duniform.prob(range,lb,ub,inclusive=c("none","left","right","both"))

# Binomial distribution
binomial.summary(n,p,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
binomial.prob(n,p,lb)
binomial.prob(n,p,lb,ub,inclusive=c("none","left","right","both"))

# Geometric distribution
geometric.summary(p,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
geometric.prob(p,lb)
geometric.prob(p,lb,ub,inclusive=c("none","left","right","both"))

# Negative Binomial distribution
negbinom.summary(r,p,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
negbinom.prob(r,p,lb)
negbinom.prob(r,p,lb,ub,inclusive=c("none","left","right","both"))

# Hypergeometric distribution
hypergeo.summary(N,K,n,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
hypergeo.prob(N,K,n,lb)
hypergeo.prob(N,K,n,lb,ub,inclusive=c("none","left","right","both"))

# Poisson distribution
poisson.summary(lambda,L,plotpdf=c("TRUE","FALSE"), plotcdf=c("TRUE","FALSE"))
poisson.prob(lambda,L,lb)
poisson.prob(lambda,L,lb,ub,inclusive=c("none","left","right","both"))
```

Arguments

x possible values of a user defined discrete random variable

<code>fx</code>	probabilities of $X=x$, the order of entries in <code>fx</code> must matches the order of entries in <code>x</code> .
<code>plotpdf</code>	TRUE or FALSE, if TRUE, it plots the pmf
<code>plotcdf</code>	TRUE or FALSE, if TRUE, it polts the cdf
<code>lb,ub</code>	lower bound (lb) and upper bound (ub) in a probability statement; lb could be $-\text{Inf}$; ub could be Inf ; ub cannot be less than lb
<code>inclusive</code>	"none": $\text{lb} < X < \text{ub}$; "left": $\text{lb} \leq X < \text{ub}$; "right": $\text{lb} < X \leq \text{ub}$; "both": $\text{lb} \leq X \leq \text{ub}$
<code>Range</code>	contains all possible values of a discrete uniform random variable
<code>p</code>	parameter p of a Binomial distribution/Geometric distribution/Negative Binomial distribution
<code>n</code>	parameter n of a Binomial distribution
<code>r</code>	parameter r of a negative Binomial distribution
<code>N,K,n</code>	parameters N , K , and n of a hypergeometric distribution
<code>lambda,L</code>	parameters λ and L of a Poisson distribution. Default: $L=1$

Details

Plot the probability mass function (pmf) and the cumulative distribution function (cdf) and calculate probability, mean, variable and standard deviation, and compute probabilities of various discrete distributions (a self-defined discrete distribution, discrete uniform distribution, binomial distribution, geometric distribution, negative binomial distribution, hypergeometric distribution, and Poisson distribution).

Value

<code>discrete.plotpdf</code>	a figure of the pmf of the user-defined discrete distribution
<code>discrete.plotcdf</code>	a figure of the cdf of the user-defined discrete distribution
<code>discrete.summary</code>	a list of the mean, variance, and standard deviation of the user-defined discrete distribution, plot of the pmf/cdf or not
<code>discrete.prob</code>	probability of X between lb and ub based on the user-defined discrete distribution
<code>duniform.summary</code>	a list of the mean, variance, and standard deviation of a discrete uniform distribution, plot of the pmf/cdf or not
<code>duniform.prob</code>	probability of X between lb and ub based on a uniform distribution
<code>binomial.summary</code>	a list of the mean, variance, and standard deviation of a binomial distribution, plot of the pmf/cdf or not
<code>binomial.prob</code>	probability of X between lb and ub based on a binomial distribution
<code>geometric.summary</code>	a list of the mean, variance, and standard deviation of a geometric distribution, plot of the pmf/cdf or not
<code>geometric.prob</code>	probability of X between lb and ub based on a geometric distribution
<code>negbinom.summary</code>	a list of the mean, variance, and standard deviation of a negative binomial distribution, plot of the pmf/cdf or not

<code>negbinom.prob</code>	probability of X between lb and ub based on a negative binomial distribution
<code>hypergeo.summary</code>	a list of the mean, variance, and standard deviation of a hypergeometric distribution, plot of the pmf/cdf or not
<code>hypergeo.prob</code>	probability of X between lb and ub based on a hypergeometric distribution
<code>poisson.summary</code>	a list of the mean, variance, and standard deviation of a Poisson distribution, plot of the pmf/cdf or not
<code>poisson.prob</code>	probability of X between lb and ub based on a Poisson distribution

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Chapter 3 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
x=c(0,1,2,3,4);fx=c(0.6561,0.2916,0.0486,0.0036,0.0001);
```

```
discrete.summary(x,fx)
discrete.summary(x,fx,plotpdf=FALSE,plotcdf=TRUE)
discrete.summary(x,fx,plotpdf=TRUE,plotcdf=FALSE)
```

```
discrete.prob(x,fx,2) #P(X=2)
```

```
discrete.prob(x,fx,2,4,inclusive="none") #P(2<X<4)
discrete.prob(x,fx,2,4,inclusive="left") #P(2<=X<4)
discrete.prob(x,fx,2,4,inclusive="right") #P(2<X<=4)
discrete.prob(x,fx,2,4,inclusive="both") #P(2<=X<=4)
```

```
discrete.prob(x,fx,2,Inf,inclusive="none") #P(2<X)
discrete.prob(x,fx,2,Inf,inclusive="left") #P(2<=X)
discrete.prob(x,fx,2,Inf,inclusive="right") #P(2<X)
discrete.prob(x,fx,2,Inf,inclusive="both") #P(2<=X)
```

```
discrete.prob(x,fx,-Inf,4,inclusive="none") #P(X<4)
discrete.prob(x,fx,-Inf,4,inclusive="left") #P(X<4)
discrete.prob(x,fx,-Inf,4,inclusive="right") #P(X<=4)
discrete.prob(x,fx,-Inf,4,inclusive="both") #P(X<=4)
```

```
#X~Discrete Uniform(1,2,4,5,8,10)
range=c(1,2,4,5,8,10)
duniform.summary(range)
duniform.prob(range,2) #P(X=2)
duniform.prob(range,2,4,inclusive="left") #P(2<=X<4)
```

```
#X~Binomial(n=5,p=0.3)
binomial.summary(5,0.3)
```

```

binomial.prob(5,0.3,2) #P(X=2)
binomial.prob(5,0.3,2,4,inclusive="left") #P(2<=X<4)

#X~Geometric(p=0.3)
geometric.summary(0.3)
geometric.prob(0.3,2) #P(X=2)
geometric.prob(0.3,2,4,inclusive="left") #P(2<=X<4)

#X~Negative Binomial(r=3,p=0.3)
negbinom.summary(3,0.3)
negbinom.prob(3,0.3,5) #P(X=5)
negbinom.prob(3,0.3,5,7,inclusive="left") #P(5<=X<7)

#X~Hypergeometric(N=300,K=100,n=4)
hypergeo.summary(300,100,4)
hypergeo.prob(300,100,4,2) #P(X=2)
hypergeo.prob(300,100,4,2,4,inclusive="left") #P(2<=X<4)

#X~Poisson(lambda=2.3,L=5)
poisson.summary(2.3,5)
poisson.prob(2.3,5,10) #P(X=10)
poisson.prob(2.3,5,1,Inf,inclusive="left") #P(1<=X)

```

One-Sample Confidence Intervals

One-Sample Tests

Description

Conduct hypothesis testing on population mean, population variance, and population proportion.
 Compute power of a test, and calculate the required sample size for a desired power.

Usage

```

#Test on population mean of a normal distribution when the population variance is known:
Ztest(mu0,H1,alpha,sigma,sample) # if sample available
Ztest(mu0,h1,alpha,sigma,n,barx) # if statistics are provided
# Power calculation
Ztest.power(H1,alpha,sigma,n,delta)
# Sample size calculation
sample.size.Ztest(H1,sigma,alpha,beta,delta)

#Test on population mean of a normal distribution when the population variance is unknown:
Ttest(mu0=?,H1=?,alpha=?,sample=?) #If data are available
Ttest(mu0=?,H1=?,alpha=?,n=?,barx=?,s=?) #If statistics are provided s=sd(data)
# Power calculation
Ttest.power(H1=?,est.sigma=sn,alpha=?,n=?,delta=?)
# Sample size calculation
sample.size.Ttest(H1=?,est.sigma=sn, beta=?,delta=?,alpha=?)

#One-Sample Proportion Z-tests on a population proportion:
Proptest(p0=?,H1=?,alpha=?,n=?,X=?)
# Power calculation

```

```

Proptest.power(H1=?,alpha=?,p0=?,p1=?,n=?)
# Sample size calculation
sample.size.Proptest(H1=?,beta=?,alpha=?,p0=?,p1=?)

```

Arguments

<code>mu0</code>	the hypothesized value of the population mean in Z-tests and T-tests
<code>H1</code>	type of alternative hypothesis: "two", "left", or "right"
<code>alpha</code>	the significance level
<code>sample</code>	a vector of the observed sample
<code>sigma</code>	the known population standard deviation
<code>n</code>	the sample size
<code>barx</code>	the observed sample mean
<code>delta</code>	the delta value in power/sample size calculation of Z-test/T-test
<code>beta</code>	the desired power for sample size calculation
<code>s</code>	the observed sample standard deviation
<code>est.sigma</code>	an estimate of the population standard deviation in power/sample size calculation of T-test
<code>sigma0</code>	the hypothesized value of the population standard deviation in Chi-square-tests
<code>lambda</code>	the lambda value in power/sample size calculation of Chi-square-tests
<code>p0</code>	the hypothesized value of the population proportion
<code>X</code>	number of observations belongs to a class of interest
<code>p1</code>	the p1 value in power/sample size calculation of One-sample Proportion Z-tests

Details

Conduct hypothesis testing on population mean, population variance, and population proportion. Compute power of a test, and calculate the required sample size for a desired power.

Value

<code>test</code>	As long as the function has "test", it produces the test results of using three approaches.
<code>power</code>	As long as the function has "test", it computes the power of the test.
<code>sample.size</code>	As long as the function has "sample.size", the outcome is the minimum sample size required to reach the given power.

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Chapter 9 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
#Ztest
#must include the = sign
Ztest(mu0=3,H1="two",alpha=0.05,sigma=0.9,n=15,barx=2.78)
Ztest.power(H1="two",alpha=0.05,sigma=0.9,n=15,delta=3.25-3)
sample.size.Ztest(H1="two",sigma=0.9,alpha=0.05,beta=1-0.9,delta=3.75-3)

#Ttest
#must include the = sign
x=c(131.15, 130.69, 130.91, 129.54, 129.64, 128.77, 130.72, 128.33,
128.24, 129.65, 130.14, 129.29, 128.71, 129.00, 129.39, 130.42,
129.53, 130.12, 129.78, 130.92)
data.summary(x)
Ttest(mu0=130,H1="two",alpha=0.05,sample=x)
Ttest.power(H1="two",est.sigma=sd(x),alpha=0.05, n=length(x),delta=0.5)
sample.size.Ttest(H1="two",est.sigma =sd(x),beta=0.25,delta=0.1,alpha=0.05)

#Chi-square test
#must include the = sign
Chi2test(sigma0=sqrt(0.01),H1="right",alpha=0.05,n=20,s=sqrt(0.0153))
Chi2test.power(H1="right",alpha=0.05,n=20,lambda=1.25)
sample.size.Chi2test(H1="right",beta=0.2,lambda=1.25,alpha=0.05)

Chi2test(sigma0=0.01,H1="right",alpha=0.01,n=15,s=0.008)
Chi2test.power(H1="right",alpha=0.01,n=15,lambda=1.5)
sample.size.Chi2test(H1="right",beta=0.2,lambda=1.25,alpha=0.01)

#One-sample proportion Z-test
#must include the = sign
Proptest(p0=0.05, H1="left",alpha=0.05,n=300,X=7)
Proptest.power(H1="left",alpha=0.05,p0=0.05,p1=0.03,n=300)
sample.size.Proptest(H1="left",beta=0.1,alpha=0.05,p0=0.05,p1=0.03)
```

One-Sample Tests

One-Sample Confidence Intervals

Description

Compute confidence intervals on the population mean, population variance, and population proportion. In addition, it computes prediction interval for a single future observation from a normal distribution.

Usage

```
#CI for pupulation mean of a normal distribution when the population variance is known:
Zinterval(level,sigma,sample) # if sample available
Zinterval(level,sigma,n,barx) # if stats are provided
# Choice of sample size for estimating the population mean when error is specified
sample.size.Zinterval(level,sigma,E)

#CI for pupulation mean when the population variance is unknown and the distribution is normal
#or when the sample size is smaller than 25:
Tinterval(level,sample) # if sample available
```

```

Tinterval(level,n,barx,s) # if stats are provided

#Large-sample CI for pupulation mean:
AZinterval(level,sample) # if sample available
AZinterval(level,n,barx,s) # if stats are provided

#CI for pupulation variance (or standard deviation) of a normal distribution:
Chi2interval(level,sample) # if sample available
Chi2interval(level,n,s) # if stats are provided

#Large-sample CI for a pupulation proportion:
Propinterval(level,n,X)
# Choice of sample size for estimating a population proportion when error is specified
sample.size.Propinterval(level,ini.p,E) # using an intial guess
sample.size.Propinterval(level,ini.p=0.5,E) # using the conservative apporach

# Prediction interval of a single future observation form a normal distribution:
Predinterval(level,sample) # if sample available
Predinterval(level,n,barx,s) # if stats are provided

```

Arguments

level	the confidence level
sample	a vector of the observed sample
sigma	the known population standard deviation
s	the observed sample standard deviation
barx	the observed sample mean
n	the sample size
X	number of observations belongs to a class of interest
E	specified error in sample size calculation
ini.p	A initial estimate of the populatin proportion. Default is 0.5 which corresponds to the conservative approach
df	the degrees of freedom of a t or chi.square distribution
q	a quantile value

Details

Compute CIs for the population mean, population variance, and population proportaion and PI for a single future observation from a normal distribution.

Value

interval	As long as the function has "interval", the outcome contains a two-sided CI (or PI) and the two one-sided confidence bounds.
sample.size	As long as the function has "sample.size", the outcome is the minimum sample size required to control the error to be no larger than E.
t.quantile	quantile of a t distribution
Chi2.quantile	quantile of a chi.square distribution

Note

deweiwang@stat.sc.edu

Author(s)

Dewei Wang

References

Chapter 8 of the textbook "Applied Statistics and Probability for Engineers" 7th edition

Examples

```
#Zinterval
#must include the = sign
x=c(64.1, 64.7, 64.5, 64.6, 64.5, 64.3, 64.6, 64.8, 64.2, 64.3)
Zinterval(level=0.95,sigma=1,sample=x)
Zinterval(level=0.99,sigma=1,sample=x)
sample.size.Zinterval(E=0.5,sigma=1,level=0.95)
# Using stats, must include the = sign
Zinterval(level=0.95,sigma=2,n=9,barx=98)

#Tinterval
#must include the = sign
Tinterval(level=0.95,n=10,barx=1000,s=20)
Tinterval(level=0.95,n=25,barx=1000,s=20)
Tinterval(level=0.99,n=10,barx=1000,s=20)
Tinterval(level=0.99,n=25,barx=1000,s=20)

#Large-sample Zinterval
#must include the = sign
x=scan("https://raw.githubusercontent.com/Harrindy/StatEngine/master/Data/Mercury.csv")
AZinterval(level=0.95,sample=x)

#Chi.square interval for variance/standard deviation
#must include the = sign
Chi2interval(level=0.95,n=20,s=0.01532)

#CIs for a population proportion
#must include the = sign
Propinterval(level=0.95,n=85,X=10)
sample.size.Propinterval(level=0.95,ini.p=0.12,E=0.05)
sample.size.Propinterval(level=0.95,ini.p=0.5,E=0.05)

#Prediction interval for normal distribution
#must include the = sign
x=c(19.8, 10.1, 14.9, 7.5, 15.4, 15.4, 15.4, 18.5, 7.9, 12.7, 11.9, 11.4, 11.4,
14.1, 17.6, 16.7, 15.8, 19.5, 8.8, 13.6, 11.9, 11.4)
Tinterval(level=0.95,sample=x)
Predinterval(level=0.95,sample=x)
```

Index

AZinterval (One-Sample Tests), 12

beta.prob (Continuous Distributions), 1

beta.quantile (Continuous Distributions), 1

beta.summary (Continuous Distributions), 1

binomial.prob (Discrete Distributions), 7

binomial.summary (Discrete Distributions), 7

Chi2.quantile (One-Sample Tests), 12

Chi2interval (One-Sample Tests), 12

Chi2test (One-Sample Confidence Intervals), 10

Continuous Distributions, 1

Counting Techniques, 4

DataSummary (Descriptive Statistics), 6

Descriptive Statistics, 6

Discrete Distributions, 7

discrete.plotcdf (Discrete Distributions), 7

discrete.plotpdf (Discrete Distributions), 7

discrete.summary (Discrete Distributions), 7

duniform.prob (Discrete Distributions), 7

duniform.summary (Discrete Distributions), 7

exponential.prob (Continuous Distributions), 1

exponential.quantile (Continuous Distributions), 1

exponential.summary (Continuous Distributions), 1

gamma.prob (Continuous Distributions), 1

gamma.quantile (Continuous Distributions), 1

gamma.summary (Continuous Distributions), 1

geometric.prob (Discrete Distributions), 7

geometric.summary (Discrete Distributions), 7

hypergeo.prob (Discrete Distributions), 7

hypergeo.summary (Discrete Distributions), 7

lognormal.prob (Continuous Distributions), 1

lognormal.quantile (Continuous Distributions), 1

lognormal.summary (Continuous Distributions), 1

nCr (Counting Techniques), 4

negbinom.prob (Discrete Distributions), 7

negbinom.summary (Discrete Distributions), 7

normal.prob (Continuous Distributions), 1

normal.quantile (Continuous Distributions), 1

normal.summary (Continuous Distributions), 1

nPr (Counting Techniques), 4

One-Sample Confidence Intervals, 10

One-Sample Tests, 12

poisson.prob (Discrete Distributions), 7

poisson.summary (Discrete Distributions), 7

Predinterval (One-Sample Tests), 12

Propinterval (One-Sample Tests), 12

Proptest (One-Sample Confidence Intervals), 10

sample.size.Chi2test (One-Sample Confidence Intervals), 10

sample.size.Propinterval (One-Sample Tests), 12

sample.size.Proptest (One-Sample
Confidence Intervals), [10](#)
sample.size.Ttest (One-Sample
Confidence Intervals), [10](#)
sample.size.Zinterval (One-Sample
Tests), [12](#)
sample.size.Ztest (One-Sample
Confidence Intervals), [10](#)
SimPerm (Counting Techniques), [4](#)

t.quantile (One-Sample Tests), [12](#)
Tinterval (One-Sample Tests), [12](#)
Ttest (One-Sample Confidence
Intervals), [10](#)

uniform.prob (Continuous
Distributions), [1](#)
uniform.quantile (Continuous
Distributions), [1](#)
uniform.summary (Continuous
Distributions), [1](#)

weibull.prob (Continuous
Distributions), [1](#)
weibull.quantile (Continuous
Distributions), [1](#)
weibull.summary (Continuous
Distributions), [1](#)

Zinterval (One-Sample Tests), [12](#)
Ztest (One-Sample Confidence
Intervals), [10](#)