



WIKIPEDIA
The Free Encyclopedia

[Main page](#)
[Contents](#)
[Featured content](#)
[Current events](#)
[Random article](#)
[Donate to Wikipedia](#)

Interaction
[Help](#)
[About Wikipedia](#)
[Community portal](#)
[Recent changes](#)
[Contact Wikipedia](#)

[Toolbox](#)
[Print/export](#)

Languages
[Français](#)
[Norsk bokmål](#)
[Polski](#)

[Create account](#)  [Log in](#)

Article

[Talk](#)

Read

[Edit](#)

[View history](#)

Search



Rosenbrock function

From Wikipedia, the free encyclopedia


In [mathematical optimization](#), the **Rosenbrock function** is a non-[convex function](#) used as a performance test problem for optimization [algorithms](#) introduced by [Howard H. Rosenbrock](#) in 1960.^[1] It is also known as **Rosenbrock's valley** or **Rosenbrock's banana function**.

The global minimum is inside a long, narrow, [parabolic](#) shaped flat valley. To find the valley is trivial. To converge to the global [minimum](#), however, is difficult.

It is defined by

$$f(x, y) = (1 - x)^2 + 100(y - x^2)^2.$$

It has a global minimum at $(x, y) = (1, 1)$ where $f(x, y) = 0$. A different coefficient of the second term is sometimes given, but this does not affect the position of the global minimum.



Contents [\[hide\]](#)

[1 Multidimensional generalisations](#)

- 1 Multidimensional generalisations
- 2 Stationary points
- 3 An example of optimization
- 4 See also
- 5 Notes
- 6 References
- 7 External links

Multidimensional generalisations [\[edit\]](#)

Two variants are commonly encountered. One is the sum of $N/2$ uncoupled 2D Rosenbrock problems,

$$f(\mathbf{x}) = f(x_1, x_2, \dots, x_N) = \sum_{i=1}^{N/2} [100(x_{2i-1}^2 - x_{2i})^2 + (x_{2i-1} - 1)^2] \quad [2]$$

This variant is only defined for even N and has predictably simple solutions.

A more involved variant is

$$f(\mathbf{x}) = \sum_{i=1}^{N-1} [(1 - x_i)^2 + 100(x_{i+1} - x_i^2)^2] \quad \forall x \in \mathbb{R}^N. \quad [3]$$

This variant has been shown to have exactly one minimum for $N = 3$ (at $(1, 1, 1)$) and exactly two minima for $4 \leq N \leq 7$ -- the global minimum of all ones and a local minimum near $(x_1, x_2, \dots, x_N) = (-1, 1, \dots, 1)$. This result is obtained by setting the gradient of the function equal to zero, noticing that the resulting equation is a rational function of x . For small N the polynomials can be determined exactly and [Sturm's theorem](#) can be used to determine the number of real roots, while the roots can be [bounded](#) in the region of $|x_i| < 2.4$.^[4] For larger N this method breaks down due to the size of the coefficients involved.

Stationary points [\[edit\]](#)

Many of the stationary points of the function exhibit a regular pattern when plotted.^[4] This structure can be exploited to locate them.

An example of optimization [\[edit\]](#)



The Rosenbrock function can be efficiently optimized by adapting appropriate coordinate system without using any [gradient information](#) and without building local approximation models (in contrast to many derivate-free optimizers). The following figure illustrates an example of 2-dimensional Rosenbrock function optimization by [Adaptive coordinate descent](#) from starting point $x_0 = (-3, -4)$. The solution with the function value 10^{-10} can be found after 325 function evaluations.


See also [\[edit\]](#)



- [Test functions for optimization](#)


Notes [\[edit\]](#)

- [^] Rosenbrock, H.H. (1960). "An automatic method for finding the greatest or least value of a function". *The Computer Journal* **3**: 175–184. doi:10.1093/comjnl/3.3.175 [↗](#). ISSN 0010-4620 [↗](#).
- [^] L C W Dixon, D J Mills. Effect of Rounding errors on the Variable Metric Method. *Journal of Optimization Theory and Applications* **80**,






1994. [1] 




3. ^ "Generalized Rosenbrock's function" .
Retrieved 2008-09-16.
4. ^ ^{*a*} ^{*b*} Schalk Kok, Carl Sandrock. Locating and Characterizing the Stationary Points of the Extended Rosenbrock Function. *Evolutionary Computation* **17**, 2009. [2] 



References [\[edit\]](#)

- Rosenbrock, H. H. (1960), "An automatic method for finding the greatest or least value of a function", *The Computer Journal* **3**: 175–184, doi:10.1093/comjnl/3.3.175 , ISSN 0010-4620 , MR0136042 

External links [\[edit\]](#)

- Rosenbrock function plot in 3D 
- Minimizing the Rosenbrock Function  by Michael Croucher, The Wolfram Demonstrations Project.
- Weisstein, Eric W., "Rosenbrock Function ", *MathWorld*.

Categories: [Mathematical optimization](#)

This page was last modified on 22 May 2013 at 14:39.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#).

Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.

[Privacy policy](#) [About Wikipedia](#) [Disclaimers](#) [Contact Wikipedia](#) [Mobile view](#)

