

Clinical Knowledge Reference - Symptoms, Conditions and Medication Timing

Synthetic enterprise knowledge document for MAOS development. It demonstrates structured clinical knowledge retrieval, exact term matching, hybrid search metadata, and evidence-backed responses. **It is not medical advice.**

Cardiometabolic Conditions

Condition	Typical symptoms / signals	Medication & timing
Hypertension	Persistent elevated blood pressure; common symptoms can be absent or include headache or dizziness.	Amlodipine — commonly taken once daily at a consistent time, according to the clinician's prescription.
Type 2 Diabetes	Common symptoms may include increased thirst, frequent urination, fatigue, or blurred vision.	Metformin — commonly taken with meals; exact schedule follows the clinician-approved regimen.
Hyperlipidemia	Often asymptomatic; discovered through lipid testing.	Atorvastatin — commonly taken once daily at a consistent time; timing follows the prescribed schedule.

Respiratory Conditions

Condition	Typical symptoms / signals	Medication & timing
Asthma	Episodes of wheezing, chest tightness, cough, and shortness of breath.	Albuterol — rescue use follows the clinician's action plan; controller medicines have separate prescribed schedules.
Community-acquired respiratory infection	May include fever, cough, fatigue, and shortness of breath.	Amoxicillin — when prescribed, doses are taken at the exact interval supplied by the clinician.
Allergic rhinitis	Sneezing, nasal congestion, itchy eyes, and runny nose.	Cetirizine — commonly taken once daily; timing follows the prescribed instructions and may be adjusted for drowsiness.

Gastrointestinal Conditions

Condition	Typical symptoms / signals	Medication & timing
Gastroesophageal reflux disease	Heartburn, regurgitation, or upper abdominal discomfort.	Omeprazole — commonly taken before a meal when prescribed; exact timing depends on the prescription.
Gastritis	Upper abdominal discomfort, nausea, or early fullness.	Omeprazole — commonly taken before a meal when prescribed; the treatment plan determines the schedule.
Irritable bowel syndrome	Abdominal pain associated with bowel habit changes.	Dicyclomine — timing follows the prescribed regimen and is individualized to the patient's treatment plan.

MAOS synthetic demonstration data - not for clinical use

Metadata for Retrieval and Authorization

Each document/chunk in the knowledge layer should carry metadata used for routing, filtering, and access control.

Field	Example	Purpose
document_type	clinical_reference	Controls document class
condition_code	HTN-001	Exact-code / keyword matching
specialty	cardiology	Departmental filtering
access_level	doctor	RBAC filtering
source_system	clinical_knowledge_base	Citation/source tracking
version	3.2	Document version control
effective_from	2026-01-01	Policy/knowledge validity
review_status	approved	Only approved content is answerable

Example retrieval behavior

User: "HTN-001 medication timing"
Retriever: combine semantic similarity with BM25 keyword matching for the exact code.
Filter: specialty=cardiology AND access_level appropriate for the authenticated role.
Answer: return only approved evidence, cite the source document/version, and avoid making an autonomous diagnosis.