

Problem 1

MATH 421 HW4, Harry Luo

Prove or disprove the following statements:

1. The set $\{x \in \mathbb{R} : x \geq 2\}$ is open.
2. The set $\{x \in \mathbb{R} : x \neq 2\}$ is open.

solution:

1. Let $\varepsilon > 0$. Consider $2 \in [2, \infty)$, and interval $(2 - \varepsilon, 2 + \varepsilon)$:

Since $2 - \frac{\varepsilon}{2} \in (2 - \varepsilon, 2 + \varepsilon)$, but $2 - \frac{\varepsilon}{2} \notin [2, \infty)$,

it follows that for any $\varepsilon > 0$, the interval $(2 - \varepsilon, 2 + \varepsilon)$ is not a subset of $\{x \in \mathbb{R} : x \geq 2\}$, so the set $\{x \in \mathbb{R} : x \geq 2\}$ is not open.

-
2. Let $\varepsilon > 0, x \in \{x \in \mathbb{R} : x \neq 2\}$. Let $\varepsilon = \left| \frac{x-2}{2} \right|$.

Then for any $y \in (x - \varepsilon, x + \varepsilon)$, we have

$$\begin{aligned} y &< x + \varepsilon, \quad y > x - \varepsilon \\ \Rightarrow |y - x| &< \varepsilon = \left| \frac{x - 2}{2} \right| \end{aligned} \tag{1}$$

Thus by triangle inequality,

$$\begin{aligned} |y - 2| &= |y - x + x - 2| \\ &\geq |x - 2| - |y - x| \\ &\geq |x - 2| - \left| \frac{x - 2}{2} \right| \\ &= \frac{|x - 2|}{2} \\ &= \varepsilon > 0 \end{aligned} \tag{2}$$

Therefore $y \neq 2 \Rightarrow y \in \{x \in \mathbb{R} : x \neq 2\}$. So the set is open. ■

Problem 2:

Let $A, B \subseteq \mathbb{R}$ be subsets. Prove the following statements:

1. (De Morgan's Laws) $(A \cap B)^c = A^c \cup B^c$ and $(A \cup B)^c = A^c \cap B^c$
 2. If A and B are closed then $A \cap B$ and $A \cup B$ are closed.
-

solution:

1. • Let $x \in (A \cap B)^c$, then $x \notin (A \cap B) \Rightarrow (x \notin A) \text{ or } (x \notin B)$

This is equivalent to $x \in A^c$ or $x \in B^c \Rightarrow x \in (A^c \cup B^c)$.

So for any $x \in (A \cap B)^c$, $x \in (A^c \cup B^c)$, thus the two sets are equal. ■

- Let $x \in (A \cup B)^c$, then $x \notin (A \cup B) \Rightarrow x \notin A$ and $x \notin B$.

So $x \in A^c$ and $x \in B^c \Rightarrow x \in (A^c \cap B^c)$. So for any $x \in (A \cup B)^c$, $x \in (A^c \cap B^c)$, thus the two sets are equal. ■

2. • If A is closed and B is closed, then A^c and B^c are open. Since unions of open sets are open, then $A^c \cup B^c$ is open.

By De Morgan's Laws, $A^c \cup B^c = (A \cap B)^c$ is closed.

Thus $A \cap B$ is open. ■

- If A is closed and B is closed, then A^c and B^c are open. Since intersections of open sets are open, then $A^c \cap B^c$ is open.

By De Morgan's Laws, $A^c \cap B^c = (A \cup B)^c$ is open.

Thus $A \cup B$ is closed. ■

Problem 3:

Let $\varepsilon > 0$. For each of the following functions $\mathbb{R} \rightarrow \mathbb{R}$ and numbers $l \in \mathbb{R}$, find a δ s.t. $0 < |x - 1| < \delta$ implies $|f(x) - l| < \varepsilon$.

1. $f(x) = x^4$ and $l = 1$
 2. $g(x) = \begin{cases} 1/x & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$, and $l = 1$
 3. $h(x) = f(x) + g(x)$ and $l = 2$. hint: in the proof of the corresponding limit laws, we saw how to pick this δ based on our answers for (a) and (b).
-

solution:

1. For any arbitrary ε , there exists a $\delta = \min\{1, \varepsilon/15\}$, s.t. $0 < |x - 1| < \delta$, so

$$0 < |x - 1| < 1 \Rightarrow \begin{cases} |x + 1| < 3 \\ |x^2 + 1| < 5 \end{cases} \quad (3)$$

and

$$\begin{aligned} |f(x) - l| &= |x^4 - 1| = |x - 1||x + 1||x^2 + 1| \\ &< \delta * 3 * 5 \\ &= 15\delta = \varepsilon \end{aligned} \quad (4)$$

■

2. For any arbitrary ε there exists $\delta = \min\{\frac{1}{2}, \frac{\varepsilon}{2}\}$, s.t. $0 < |x - 1| < \delta$, so

$$1 - \delta < x < \delta + 1 \Rightarrow \frac{1}{2} < \frac{1}{x} < 2. \quad (5)$$

and

$$|g(x) - 1| = \left| \frac{1}{x} - 1 \right| = \frac{|x - 1|}{x} < 2|x - 1| = 2\delta = \varepsilon. \quad (6)$$

■

3. $|h(x) - 2| = |f(x) - 1 + g(x) - 1| < |f(x) - 1| + |g(x) - 1| \quad (7)$

From the previous two parts, we know that we can choose $\delta_1 = \min\{1, \frac{\varepsilon_1}{15}\}$ and $\delta_2 = \min\{\frac{1}{2}, \frac{\varepsilon_2}{2}\}$. To ensure Equation 7 is smaller than ε , we choose

$$\delta = \min\left\{\frac{1}{2}, 1, \frac{\varepsilon}{2}, \frac{\varepsilon}{15}\right\} = \min\left\{\frac{1}{2}, \frac{\varepsilon}{15}\right\} \quad (8)$$

Therefore,

$$|h(x) - 2| < \varepsilon. \quad (9)$$

■

Problem 4:

let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be functions s.t. $\lim_{x \rightarrow a} f(x) = l$ and $\lim_{x \rightarrow a} g(x) = m$ for some numbers $a, l, m \in \mathbb{R}$. Prove that if $\forall x \in \mathbb{R} f(x) \leq g(x)$, then $l \leq m$.

solution:

Problem 5:

Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be functions and $a \in \mathbb{R}$. Prove or disprove the following statements:

- (a) If $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ both do not exist, then $\lim_{x \rightarrow a} (f + g)(x)$ does not exist.
- (b) If $\lim_{x \rightarrow a} f(x)$ exists and $\lim_{x \rightarrow a} (f + g)(x)$ does not exist, then $\lim_{x \rightarrow a} g(x)$ does not exist.
- (c) If $\lim_{x \rightarrow a} f(x)$ exists and $\lim_{x \rightarrow a} g(x)$ does not exist, then $\lim_{x \rightarrow a} (f + g)(x)$ does not exist.

(hint: Each statement is either an application of the limit law for addition, or it is false.

Remember, if the statement is false, then we need to come up with a counterexample.)

solution: