



Calculus of Variations

History: In 1696 Johann Bernoulli of the University of Basel posed the following problem: given two points A and B in a vertical plane, find the path $A \rightarrow B$ which the moveable particle M will traverse in the shortest time, assuming that its acceleration is due to gravity only and there is no friction. This is famous BRACHISTOCHROME shortest Time from the Greek.



the path $A \rightarrow B$ which the moveable particle M will traverse in the shortest time, assuming that its acceleration is due to gravity only and there is no friction. This is famous BRACHISTOCHROME shortest Time from the Greek.



$$dt = \frac{ds}{v} \text{ instantaneous velocity}$$

$$ds^2 = dx^2 + dy^2 \Rightarrow ds = \sqrt{1 + (y')^2} dx$$

$$\frac{mv^2}{2} = mgy \Rightarrow v = \sqrt{2gy}$$


$$t[y(x)] = \frac{1}{\sqrt{2g}} \int_{x_A}^{x_B} \frac{\sqrt{1 + (y')^2}}{\sqrt{y}} dx$$

$t[y(x)]$ is called a functional

Minimization (maximization) of a functional is the subject of calculus of variations.

⇒ Euler equation for the functionals of the type:

$$v[y(x)] = \int_{x_0}^{x_1} F(x, y(x), y'(x)) dx$$



$$y(x, \varepsilon) = y(x) + \varepsilon \delta y(x).$$

$$\varepsilon \rightarrow 0$$

$\delta y(x)$ is called variation

$$v[y(x, \varepsilon)] = \varphi(\varepsilon) \text{ so}$$

that in the sense of parameter ε extremum of the functional corresponds to the extremum of the function

$$\varphi(\varepsilon), \text{ namely: } \varphi'(0) = 0$$

$$\varphi(\varepsilon) = \int_{x_0}^{x_1} F(x, y(x, \varepsilon), y'(x, \varepsilon)) dx$$

$$\varphi'(\varepsilon) = \int_{x_0}^{x_1} \left[\frac{\partial F}{\partial y} \frac{\partial}{\partial \varepsilon} y(x, \varepsilon) + \frac{\partial F}{\partial y'} \frac{\partial}{\partial \varepsilon} y'(x, \varepsilon) \right] dx$$

$$F_y = \frac{\partial}{\partial y} F(x, y(x, \varepsilon), y'(x, \varepsilon))$$

$$F_{y'} = \frac{\partial}{\partial y'} F(x, y(x, \varepsilon), y'(x, \varepsilon))$$

$$\frac{\partial}{\partial \varepsilon} y(x, \varepsilon) = \delta y \quad \frac{\partial}{\partial \varepsilon} y'(x, \varepsilon) = \delta y'$$

$$\varphi'(\varepsilon) = \int_{x_0}^{x_1} [F_y \delta y + F_{y'} \delta y'] dx$$

$$\delta y' = \delta \left(\frac{dy}{dx} \right) = \frac{d}{dx} \delta y$$

Integrate this term by parts

$$\varphi'(\varepsilon) = \int_{x_0}^{x_1} \left[F_y \delta y + \overbrace{F_{y'} \frac{d}{dx} \delta y} \right] dx$$

$$\psi'(\epsilon) = (F_{y'} \delta y)_{x=x_0}^{x=x_1} + \int_{x_0}^{x_1} \left[F_y - \frac{d}{dx} (F_{y'}) \right] \delta y \, dx$$

Since $\delta y(x_0) = \delta y(x_1) = 0 \Rightarrow$ fixed end points

$$\psi'(0) = \int_{x_0}^{x_1} \left(F_y - \frac{d}{dx} F_{y'} \right) \delta y(x) \, dx$$

Fundamental Theorem: If for any continuous function $\gamma(x)$ defined at the interval $x \in [x_0, x_1]$

$$\int_{x_0}^{x_1} P(x) \gamma(x) \, dx = 0$$

where function $P(x)$ is also continuous at that interval, then

$$P(x) \equiv 0$$


Applying the (FT) to above variation and from condition $\psi'(0) = 0$ we find:

$$F_y - \frac{d}{dx} F_{y'} = 0$$

Euler (1744)

Some famous problems


(#1) The minimum surface of revolution: find curve $y(x)$



which upon revolution covers the smallest area?

(#2) Find the plane curve of given length which encloses the greatest possible area?

(#3) freely suspended chain:



Exp. 1 $v[y(x)] = \int_0^{\pi/2} [(y')^2 - y^2] dx \quad y(0)=0 \quad y(\pi/2)=1$

$$F_y = -2y \quad F_{y'} = 2y'$$

$$F_y - \frac{d}{dx} F_{y'} = 0 \implies y'' + y = 0$$

$$y(x) = C_1 \cos x + C_2 \sin x$$

$$y(0) = C_1 = 0$$

$$y(\pi/2) = C_2 = 1$$

$$y(x) = \sin x$$

Exp. 2 $v[y(x)] = \int_0^1 [(y')^2 + 12xy] dx \quad y(0)=0 \quad y(1)=1$

$$F_y = 12x \quad F_{y'} = 2y'$$

$$F_y - \frac{d}{dx} F_{y'} = 0 \implies y'' - 6x = 0$$

$$y(x) = x^3 + C_1 x + C_2$$

$$y(0) = C_2 = 0$$

$$y(1) = 1 + C_1 = 1 \implies C_1 = 0$$

$$y(x) = x^3$$

If F depends only on y and y' Euler equation has first integral. Indeed:

$$F_y - \frac{d}{dx} F_{y'} = F_y - F_{xy'} - y' F_{y'y} - y'' F_{y'y'} = 0$$

Assume $F = F(y, y')$ so $\Rightarrow F_{xy'} = 0$

$$y' (F_y - y' F_{y'y} - y'' F_{y'y'}) = \frac{d}{dx} (F - y' F_{y'}) = 0$$

$$\boxed{F - y' F_{y'} = C}$$

In physical terms this integral corresponds to energy conservation.

Solution to Special problem #3

$$E = \rho g \int_A^B y \, ds = \rho g \int_A^B y \sqrt{1 + (y')^2} \, dx$$

$$L = \int_A^B ds = \int_A^B \sqrt{1 + (y')^2} \, dx$$

$$v[y(x)] = \int_{x_A}^{x_B} \left[\rho g y \sqrt{1 + (y')^2} - \lambda \sqrt{1 + (y')^2} \right] dx$$

Lagrange multiplier

It is convenient to choose $\lambda = \rho g y_0$ where y_0 is some constant, so:

$$v[y(x)] = \rho g \int_{x_A}^{x_B} (y - y_0) \sqrt{1 + (y')^2} \, dx$$

Change now variables $y - y_0 = z \Leftrightarrow y' = z'$

$$v[z(x)] = \rho g \int_A^B z \sqrt{1 + (z')^2} \, dx$$

This functional is independent of x so that has first

$$\underbrace{z \sqrt{1+(z')^2} - z' \frac{z \cdot z'}{\sqrt{1+(z')^2}}} = C$$

$$\frac{z}{\sqrt{1+(z')^2}} = C \Rightarrow z^2 = C^2 [1+(z')^2] \Rightarrow$$

$$\frac{dz}{dx} = \frac{\sqrt{z^2 - C^2}}{C} \Rightarrow \frac{dz}{\sqrt{z^2 - C^2}} = \frac{dx}{C}$$

Change variables now $z = c \cosh t$ $dz = \frac{c \sinh t}{dt} dt$

$$\frac{c}{c} \cdot \frac{\sinh t}{c \sqrt{\cosh^2 t - 1}} dt = \frac{dx}{C} \Rightarrow t = \frac{x - x_0}{C}$$

So that $z(x) = C \cosh \left(\frac{x - x_0}{C} \right)$ and finally,

$$\boxed{y(x) = y_0 + C \cosh \left(\frac{x - x_0}{C} \right)}$$

The three constants y_0, x_0 and C are chosen so that the curve passes through (x_A, y_A) and (x_B, y_B) and so that the length is L . This presents numerical problems, but is always possible in principle.

Generalized functionals

$$J[y(x)] = \int_{x_0}^{x_1} F(x, y(x), y'(x), \dots, y^{(n)}(x)) dx$$

$$\begin{cases} y(x_0) = y_0 & y'(x_0) = y_0', \dots, y_0^{(n-1)}(x_0) = y_0^{(n-1)} \\ y(x_1) = y_1 & y'(x_1) = y_1', \dots, y_1^{(n-1)}(x_1) = y_1^{(n-1)} \end{cases}$$

Introduce a variation $y(\epsilon, x) = y(x) + \epsilon \delta y(x) :$

$$\delta v = \frac{d}{d\epsilon} \left[\int_{x_0}^{x_1} F(x, y(\epsilon, x), y'(\epsilon, x), \dots, y^{(n)}(\epsilon, x)) dx \right]_{\epsilon=0} =$$

$$= \int_{x_0}^{x_1} \left[F_y \delta y + F_{y'} \delta y' + F_{y''} \delta y'' + \dots + F_{y^{(n)}} \delta y^{(n)} \right] dx$$

$$a) \int_{x_0}^{x_1} F_{y'} \delta y' dx = \underbrace{F_{y'} \delta y(x) \Big|_{x_0}^{x_1}}_{\equiv 0} - \int_{x_0}^{x_1} \delta y(x) \frac{d}{dx} (F_{y'}) dx$$

$$b) \int_{x_0}^{x_1} F_{y''} \delta y'' dx = F_{y''} \delta y'(x) \Big|_{x_0}^{x_1} - \int_{x_0}^{x_1} \delta y'(x) \frac{d}{dx} F_{y''} dx =$$

$$= \underbrace{F_{y''} \delta y'(x) \Big|_{x_0}^{x_1}}_{\equiv 0} - \underbrace{\frac{d}{dx} F_{y''} \delta y(x) \Big|_{x_0}^{x_1}}_{\equiv 0} + \int_{x_0}^{x_1} \frac{d^2}{dx^2} (F_{y''}) \delta y(x) dx$$

$$\therefore n) \int_{x_0}^{x_1} F_{y^{(n)}} \delta y^{(n)} dx = (-1)^{n-1} \int_{x_0}^{x_1} \frac{d^{n-1}}{dx^{n-1}} F_{y^{(n)}} \delta y(x) dx$$

$$\delta v = \int_{x_0}^{x_1} \delta y(x) \left[F_y - \frac{d}{dx} F_{y'} + \frac{d^2}{dx^2} F_{y''} + \dots + (-1)^{n-1} \frac{d^{n-1}}{dx^{n-1}} F_{y^{(n)}} \right] dx$$

$$\boxed{F_y - \frac{d}{dx} F_{y'} + \frac{d^2}{dx^2} F_{y''} + \dots + (-1)^{n-1} \frac{d^{n-1}}{dx^{n-1}} F_{y^{(n)}} = 0}$$

Euler - Poisson equation.

Exp. 1

$$v[y(x)] = \int_0^{\pi/2} [(y'')^2 - y^2 + x^2] dx \quad \begin{array}{l} y(0)=1 \quad y'(0)=0 \\ y(\pi/2)=0 \quad y'(\pi/2)=1 \end{array}$$

$$\{ F_y = -2y; F_{y'} = 0; F_{y''} = 2y'' \} : y^{(4)} - y = 0$$

$$K^4 - 1 = 0 \Rightarrow K^2 = \pm 1 \Rightarrow K_1 = \pm 1 \quad K_{2,3} = \pm i$$

$$y(x) = C_1 e^x + C_2 e^{-x} + C_3 \cos x + C_4 \sin x$$

By using boundary conditions :

$$C_1 = 0 \quad C_2 = 0 \quad C_3 = 1 \quad C_4 = 0$$

$$\boxed{y(x) = \cos x}$$

Exp. 2

$$v[y(x)] = \int_{-l}^l \left[\frac{m}{2} (y'')^2 + P y \right] dx \quad \begin{array}{l} y(-l) = 0 \quad y'(-l) = 0 \\ y(l) = 0 \quad y'(l) = 0 \end{array}$$

$$F_y = P \quad F_{y'} = 0 \quad F_{y''} = m y''$$

$$P + \frac{d^2}{dx^2} (m y'') = 0 \Rightarrow y^{(4)} = - \frac{P}{m}$$

$$y(x) = - \frac{P x^4}{24m} + C_1 x^3 + C_2 x^2 + C_3 x + C_4$$

By using boundary conditions

$$y(x) = - \frac{P}{24m} (x^4 - 2l^2 x^2 + l^4) \Rightarrow \boxed{y(x) = - \frac{P}{24m} (x^2 - l^2)^2}$$

If functionals depends on many functions then one has to do variational analysis for each

$$v[y_1, y_2, \dots, y_m] = \int_{x_0}^{x_1} F(x, y_1, y_1', \dots, y_1^{(n_1)}, y_2, y_2', \dots, y_2^{(n_2)}, \dots, y_m, y_m', \dots, y_m^{(n_m)}) dx$$

$$\boxed{F_{y_i} - \frac{d}{dx} F_{y_i'} + \frac{d^2}{dx^2} F_{y_i''} + \dots + (-1)^{n_i} \frac{d^{n_i}}{dx^{n_i}} F_{y_i^{(n_i)}} = 0}$$

$$i = 1, 2, \dots, m$$

System of coupled differential equations of order $[2n_i]_{\max}$