

Contents

I Vector algebra	1
I.1 Coordinate Transformation	1
I.1.1 cylindrical	1
I.1.2 spherical	1
I.2 Dot product	2
I.3 cross product	2
I.4 Projection	2
II Vector calculus	2
II.1 Arc length	2
II.2 Line integration	3
II.3 Surface integration	3
II.4 Jacobian	3
II.5 Gradient	4
II.6 Divergence	5
II.7 Green's theorem	5
II.8 Surface Integral	5
II.9 Stokes' theorem	6
III Complex analysis	6
III.1 Complex numbers and basic operations	6
III.1.1 Definitions	6
III.1.2 operations	6
III.2 Differentiation	6
III.2.1 open sets in \mathbb{C}	6
III.2.2 Holomorphic function	6
III.2.3 Differentiation operations	7
III.2.4 Cauchy-Riemann equations	7

I Vector algebra

I.1 Coordinate Transformation

I.1.1 cylindrical

$$x = \rho \cos \varphi$$

$$y = \rho \sin \varphi$$

$$z = z$$

reverse

$$\rho = \sqrt{x^2 + y^2}$$

$$\cos \varphi = \frac{x}{\rho}$$

$$\sin \varphi = \frac{y}{\rho}$$

I.1.2 spherical

$$x = \rho \sin \varphi \cos \theta$$

$$y = \rho \sin \varphi \sin \theta$$

$$z = \rho \cos \varphi$$

reverse

$$\rho = \sqrt{x^2 + y^2 + z^2}$$

$$\cos \varphi = \frac{z}{\rho}$$

$$\cos \theta = \frac{x}{r}$$

$$\sin \theta = \frac{y}{r}$$

I.2 Dot product

- commutative
- positive definite
- distributive
- cauchy-schwarz inequality

I.3 cross product

- anticommutative $\vec{u} \times \vec{v} = -(\vec{v} \times \vec{u})$
- distributive $\vec{u} \times (\vec{v} + \vec{w}) = \vec{u} \times \vec{v} + \vec{u} \times \vec{w}$
- scalar multiplication
- triple scalar product $\vec{u} \cdot (\vec{v} \times \vec{w}) = (\vec{u} \times \vec{v}) \cdot \vec{w}$
- triple vector product $\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{b} \cdot \vec{a})\vec{c} - (\vec{c} \cdot \vec{a})\vec{b}$

I.4 Projection

The projection of \vec{a} onto \vec{b} is given by

$$\frac{\vec{a} \cdot \vec{b}}{\|\vec{b}\|^2} \vec{b} = (a \cdot \hat{b}) \hat{b}$$

II Vector calculus

II.1 Arc length

- Def: Given a curve $\vec{r}(u) = (x(u), y(u), z(u))$ for $a \leq t \leq b$ the length of the curve S, as a function of time is given by

$$S(t) = \int_a^t \|\dot{r}(u)\| du$$

where $\|\dot{r}(u)\| = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2}$

- Curvature:

$$K(t) = \frac{\|\dot{T}(t)\|}{\|\dot{r}(t)\|} = \frac{\|(\dot{r}(t) \times \ddot{r}(t))\|}{(\|\dot{r}(t)\|)^3}, \text{ where } T(t) = \frac{\dot{r}(t)}{\|\dot{r}(t)\|}$$

II.2 Line integration

- for curve $\vec{r}(t) = (x(t), y(t))$

$$\int_C f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

- center of mass $(\bar{x}, \bar{y}, \bar{z})$, where

$$\begin{cases} \bar{x} = \left(\frac{1}{M}\right) \int_C \rho(x, y, z) x ds \\ \bar{y} = \left(\frac{1}{M}\right) \int_C y \rho(x, y, z) ds \\ \bar{z} = \left(\frac{1}{M}\right) \int_C z \rho(x, y, z) ds \end{cases}$$

- Work done by force F along curve, $\vec{r}(t)$, which can be generalized into the formula for line integration,

$$W = \int_C F \cdot d\vec{r} = \int_C \vec{F} \cdot \vec{T} ds = \boxed{\int_a^b F[x(t), y(t)] \cdot (\dot{r}(t)) dt}$$

- When vector field $\vec{F} = \vec{F}(x, y, z) = (P, Q, R)$,

$$\int_C \vec{F} \cdot d\vec{r} = \int_C P dx + Q dy + R dz$$

II.3 Surface integration

- Parametric representation of surface:

$$\begin{cases} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{cases}$$

- Use normal vector at a point (u_0, v_0) of surface to represent tangent plane.

$$\vec{r}_v = \frac{\partial \vec{r}}{\partial v}(u_0, v_0), \vec{r}_u = \frac{\partial \vec{r}}{\partial u}(u_0, v_0)$$

$$\vec{N} = \vec{r}_u \times \vec{r}_v$$

- Surface area of a surface S with $(u, v) \in D$

$$A(S) = \iint_D \|\vec{r}_u \times \vec{r}_v\| du dv$$

II.4 Jacobian

- Def: Given a transformation $(u, v) \in D \longrightarrow [x(u, v), y(u, v)] \in S$, the Jacobian is given by

$$J(u, v) = \frac{\partial(x, y)}{\partial(u, v)} \equiv \det \begin{pmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{pmatrix}$$

- Jacobian in coordinate transformation

Upon evaluating an integral, we can change the coordinates of the integral from $\{x, y\} \rightarrow \{u, v\}$ by parametrize the variables:

$$x = x(u, v) \quad y = y(u, v)$$

Then the integral becomes

$$\iint_S f(x, y) \, dA = \iint_D f(x(u, v), y(u, v)) |J(u, v)| \, du \, dv$$

II.5 Gradient

- Nabla operation:

$$\nabla = \frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z}$$

- Gradient in cartesian Scalar field $f = f(x, y, z)$

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

- Gradient in polar coordinates $f = f(r, \theta)$

$$\nabla f = \vec{e}_r \frac{\partial f}{\partial r} + \vec{e}_\theta \frac{1}{r} \frac{\partial f}{\partial \theta}$$

$$\text{where } \vec{e}_r = \frac{x}{\|x\|} = (\cos \theta, \sin \theta) \vec{e}_\theta = (-\sin \theta, \cos \theta)$$

$$\nabla = \vec{e}_r \partial_r + \vec{e}_\theta \frac{1}{r} \partial_\theta$$

- Gradient in spherical

$$\nabla f = \hat{\rho} \partial_\rho + \hat{\varphi} \frac{1}{\rho} \partial_\varphi + \hat{\theta} \frac{1}{\rho \sin \varphi} \partial_\theta$$

- Gradient of scalar field in spherical coordinates

Let $g(\rho, \phi, \theta) = f(x, y, z)$

$$\begin{cases} x = \rho \sin \phi \cos \theta \\ y = \rho \sin \phi \sin \theta \\ z = \rho \cos \phi \end{cases} \quad \begin{bmatrix} \partial_\rho g \\ \partial_\phi g \\ \partial_\theta g \end{bmatrix} = \begin{bmatrix} \partial_\rho x & \partial_\rho y & \partial_\rho z \\ \partial_\phi x & \partial_\phi y & \partial_\phi z \\ \partial_\theta x & \partial_\theta y & \partial_\theta z \end{bmatrix} \begin{bmatrix} \partial_x f \\ \partial_y f \\ \partial_z f \end{bmatrix}$$

$$\begin{aligned} \hat{\rho} &= (\partial_\rho x, \partial_\rho y, \partial_\rho z) = \frac{(x, y, z)}{\rho} \\ \hat{\phi} &= \frac{1}{\rho} (\partial_\phi x, \partial_\phi y, \partial_\phi z) \\ \hat{\theta} &= \frac{1}{\rho \sin \phi} (\partial_\theta x, \partial_\theta y, \partial_\theta z) \end{aligned} \quad \Rightarrow \quad \begin{bmatrix} \partial_x f \\ \partial_y f \\ \partial_z f \end{bmatrix} = \begin{bmatrix} \hat{\rho}_1 & \hat{\phi}_1 & \hat{\theta}_1 \\ \hat{\rho}_2 & \hat{\phi}_2 & \hat{\theta}_2 \\ \hat{\rho}_3 & \hat{\phi}_3 & \hat{\theta}_3 \end{bmatrix} \begin{bmatrix} \partial_\rho g \\ \frac{1}{\rho} \partial_\phi g \\ \frac{1}{\rho \sin \phi} \partial_\theta g \end{bmatrix}$$

II.6 Divergence

- div of vec field:

3D:

$$\nabla \cdot \vec{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

- Div in polar 2D

$$\vec{U} = U_r \hat{r} + U_\theta \hat{\theta}, \text{ where } U_r = U \cdot \hat{r}, U_\theta = U \cdot \hat{\theta}$$

$$\nabla \cdot U = \left(\frac{1}{r} \right) \frac{\partial(r U_r)}{\partial r} + \frac{\partial U_\theta}{\partial \theta}$$

- Div in spherical coord

$$\vec{U} = U_\rho \hat{\rho} + U_\theta \hat{\theta} + U_\phi \hat{\phi},$$

$$\nabla \cdot \vec{U} = \frac{1}{\rho^2} \frac{\partial(\rho^2 U_\rho)}{\partial \rho} + \frac{1}{\rho} \sin \phi \frac{\partial(U_\theta)}{\partial \theta} + \frac{1}{\rho \sin \phi} \frac{\partial(U_\phi \sin \phi)}{\partial \phi}$$

II.7 Green's theorem

$$\int_C P dx + Q dy = \iint_D \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dA = \iint_C \vec{F} \cdot d\vec{r}$$

II.8 Surface Integral

- for a surface,

$$\vec{r}(u, v) = (x(u, v), y(u, v), z(u, v))$$

$$\Rightarrow \iint_S \vec{F} \cdot d\vec{S} = \iint_S \vec{F} \cdot \vec{n} dS = \iint_D \vec{F}(\vec{r}(u, v)) \cdot (\vec{r}_u \times \vec{r}_v) dA$$

- if the surface is a graph of a function $z = g(x, y)$, $(x, y) \in D$, $\vec{F} = (P, Q, R)$, then

$$\int_S \vec{F} \cdot d\vec{s} = \iint_D (P, Q, R) \cdot (-\partial_x g, -\partial_y g, 1) dA$$

II.9 Stokes' theorem

Let $F : R^3 \rightarrow R^3$ be a vector field on R^3 , then

$$\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl}(\vec{F}) \cdot d\vec{s},$$

where $\text{curl}(\vec{F}) = \nabla \times \vec{F}$

III Complex analysis

III.1 Complex numbers and basic operations

III.1.1 Definitions

- Def: $i^2 = -1$
- Complex number: $z = x + iy$
- Conjugate: $z = x - iy$
- Real part: $\Re(z) = x$, Imaginary part: $\Im(z) = y$
- Modulus/ Norm/ Magnitude: $|z| = \sqrt{x^2 + y^2}$
- Polar form: $z = |z| (\cos \theta + i \sin \theta) = re^{i\theta}$
- Argument(angle) : $\arg(z) = \theta$ such that $z = |z| (\cos \theta + i \sin \theta)$. Angle between vector (x, y) with real axis

III.1.2 operations

- addition: $z_1 + z_2 = (x_1 + x_2) + i(y_1 + y_2)$
- multiplication: $z_1 z_2 = (x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1)$
(normal multiplication with $i^2 = -1$)
- Division:

$$\frac{z_1}{z_2} = \frac{z_1 z_2^*}{z_2 z_2^*} = \frac{x_1 x_2 + y_1 y_2}{x_2^2 + y_2^2} + i \frac{x_2 y_1 - x_1 y_2}{x_2^2 + y_2^2}$$

- Commutativity: $z_1 z_2 = z_2 z_1$ $z_1 + z_2 = z_2 + z_1$
- associativity: $(z_1 z_2) z_3 = z_1 (z_2 z_3)$ $(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3)$
- distributivity: $z_1 (z_2 + z_3) = z_1 z_2 + z_1 z_3$
- Trig inequality: $|z_1 + z_2| \leq |z_1| + |z_2|$

III.2 Differentiation

III.2.1 open sets in \mathbb{C}

- Def: Let $z_0 \in \mathbb{C}$, $r > 0$. Disk $B_{r(z_0)} = \{z \in \mathbb{C} \mid |z - z_0| < r\}$ It is very important to note that it's not "less or equal"

Given a set $\Omega \in \mathbb{C}$, A point $z_0 \in \Omega$ is called an interior point of Ω if there exists $r > 0$ s.t. $B_{r(z_0)} \subset \Omega$.

A set Ω is **open** if every point of Ω is an interior point of Ω . In other words, there are no points on the boundary of Ω that are included in Ω .

III.2.2 Holomorphic function

Let Ω be an open set in \mathbb{C} , A function $f : \Omega \rightarrow \mathbb{C}$ is called **holomorphic** at $z_0 \in \Omega$ if the limit

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} (h \in \mathbb{C}, h \neq 0)$$

exists.

- The said function $f(z)$ is holomorphic on Ω if it is holomorphic on every point of Ω .
- In the special case that f is holomorphic on \mathbb{C} , f is an **entire** function.
- Holomorphic in 1st order guarantees holomorphic and analytic in any order and thus continuous.

III.2.3 Differentiation operations

If f and g are holomorphic on Ω , then

- $f + g$ is holomorphic on Ω ,

$$(f + g)' = f' + g'$$

- fg is analytic on Ω ,

$$(fg)' = f'g + fg'$$

- $\frac{f}{g}$ is analytic and, if $g(z) \neq 0$,

$$\frac{f}{g} = \frac{f'g - fg'}{g^2}$$

III.2.4 Cauchy-Riemann equations

for complex function $f : \Omega \rightarrow \mathbb{C}$, $f(z) = u(x, y) + iv(x, y)$ that is holomorphic at $z_0 = x_0 + iy_0$, then the partial derivatives of u and v exist and satisfy the Cauchy-Riemann equations:

$$\partial_x u = \partial_y v, \quad \partial_y u = -\partial_x v$$

Conversely, if u and v are continuously differentiable on an open set Ω and satisfy the Cauchy-Riemann equations, then $f(z) = u(x, y) + iv(x, y)$ is holomorphic on Ω .

In the language of logic, let C be “satisfying cauchy-riemann equations”, and H be “function is holomorphic”, then $H \rightarrow C$. If D is “ u and v have continuous partial derivatives with respect to x and y ”, then $(C \& D) \leftrightarrow H$