

PBPK evaluation report — Sildenafil

Fit is acceptable (overall GMFE 1.982); parameters were fitted by the numerical optimizer.

Objective: Build a whole-body PBPK model for Sildenafil that reproduces the observed plasma concentration-time data. Decide the model structure and any parameters you cannot obtain from the given data, and justify them. No modeling strategy and no reference results are provided. Overall GMFE 1.982 (reference 1.829); within-2-fold 59.4%. Structure: Cellular partition coefficient method - Rodgers and Rowland, Cellular permeability - PK-Sim Standard; processes CYP2C19, CYP2C9, CYP3A4.

Model, parameters & rationale

MODEL CHOICE

The model is a whole-body PBPK structure with Rodgers & Rowland tissue partitioning and PK-Sim Standard cellular permeability. Rodgers & Rowland is the defensible choice for sildenafil because the molecule is a moderately lipophilic, weakly basic piperazine (basic pKa in the ~6–6.5 range) that carries partial positive charge at physiological pH; the R&R formalism explicitly separates neutral-lipid/neutral-phospholipid partitioning from electrostatic association of the ionized fraction with acidic phospholipids, which is the dominant driver of tissue uptake for bases in muscle and other high-AP tissues. Berezchkovski or Poulin-type methods, which treat distribution as lipid partitioning of the neutral species only, would systematically under-predict Vss for this compound class. PK-Sim Standard permeability is appropriate given sildenafil's high passive permeability and absence of evidence for uptake-transporter-limited tissue entry, so tissue kinetics are effectively perfusion-limited and no permeability-limited compartment is warranted.

Elimination is represented as three saturable hepatic CYP pathways – CYP3A4, CYP2C9 and CYP2C19 – with recombinant-enzyme Vmax/Km taken from the literature and scaled to PK-Sim expression profiles. This mirrors the known biology: sildenafil is cleared almost entirely by oxidative metabolism (N-demethylation), predominantly CYP3A4 with a minor CYP2C9 contribution, and renal excretion of unchanged drug is negligible, so the absence of a renal clearance term is justified rather than an omission. Two structural simplifications should be stated plainly: gut-wall CYP3A4 is not separately parameterised beyond the standard expression profile, so oral first-pass extraction is carried implicitly by hepatic/intestinal expression and by the fitted kcat; and the pharmacologically active N-desmethyl metabolite is not tracked, which is acceptable for a parent-plasma objective but means the model cannot be used for metabolite exposure or for active-moiety DDI predictions.

PARAMETERS

Lipophilicity = 2.4 [estimated] sens=1 ref=2.8413676507
kcat@CYP3A4 = 18.00003864465053 [estimated] sens=0.059 ref=27.2146958692
Fraction unbound (plasma, reference value) = 0.04 [fixed] ref=0.04
Solubility at reference pH = 3.5 [fixed] ref=None
Reference pH = 7.0 [fixed] ref=None
kcat@CYP2C9 = 1.0 [fixed] ref=3.2171312254
kcat@CYP2C19 = 1.0 [fixed] ref=1.6232116026
Specific intestinal permeability (transcellular) = 8.6e-05 cm/min [fixed] ref=0.0012071993309
Permeability = 0.0014 cm/min [fixed] ref=0.0758702742
In vitro Vmax/recombinant enzyme@CYP2C19 = 0.01009105 [given] ref=None
^ literature/given value (Publication), not fitted
Km@CYP2C19 = 23.1 [given] ref=None
^ literature/given value (Publication), not fitted
In vitro Vmax/recombinant enzyme@CYP2C9 = 0.02 [given] ref=None
^ literature/given value (Publication), not fitted
Km@CYP2C9 = 9.6 [given] ref=None
^ literature/given value (Publication), not fitted
In vitro Vmax/recombinant enzyme@CYP3A4 = 0.16918611 [given] ref=None
^ literature/given value (Publication), not fitted
Km@CYP3A4 = 23.1 [given] ref=None
^ literature/given value (Publication), not fitted

Model, parameters & rationale

PHARMACOLOGICAL RATIONALE

Two parameters were estimated. ****Lipophilicity (2.4, sensitivity 1.00)**** is by far the most influential parameter in this run and is therefore the one the data actually determine. However, it sits below the commonly cited experimental logP for sildenafil (~2.7–3.0); no logP/logD anchor was supplied in the literature table here, so the comparison is against external values rather than a provided reference, but the departure is real and should not be waved through. The downward shift is best read as an *effective* lipophilicity – a lumped descriptor that the optimizer used to reduce tissue partitioning and hence Vss – not as a measured partition coefficient. The direction of the mismatch is consistent with the IV diagnostics (IV bias 1.73, i.e. systematic over-prediction of IV concentrations), which is exactly the signature of distribution volume being trimmed to compensate elsewhere. Because sensitivity is maximal, this value is tightly determined *by this dataset*, which makes the discrepancy with measured logP a genuine model-adequacy flag rather than a benign uncertainty. ****kcat@CYP3A4 (18.0, sensitivity 0.059)**** is the opposite case: it is essentially unconstrained. The fit barely responds to it, and the returned value (18.00004) is numerically indistinguishable from its starting/bound value, so the optimizer effectively did not move it. It must be treated as a preliminary, weakly identified number despite the completed optimization, and it should not be quoted as a determined turnover rate – particularly since it is ~100-fold above the given in-vitro recombinant Vmax for CYP3A4 (0.169 pmol/min/pmol, Warrington 2000), a gap that silently absorbs IVIWE scaling and enzyme-abundance assumptions. Collinearity between lipophilicity and kcat@CYP3A4 is low (0.151), so the distribution/clearance split is not structurally confounded; the problem with kcat is insensitivity, not trade-off.

Among the non-estimated parameters: ****fu,plasma = 0.04**** is fixed and agrees well with the literature ~96% plasma protein binding for sildenafil – this is real support, though as a fixed value it contributes no evidence from the fit itself. ****Solubility 3.5 mg/L at reference pH 7.0**** is fixed directly to Salerno 2021, and the CYP2C19/CYP2C9/CYP3A4 Km and in-vitro Vmax values are given from Warrington 2000 – all literature-anchored and not fitted. Three values require explicit caution because they are neither fitted nor literature-cited: ****kcat@CYP2C9 = 1.0** and **kcat@CYP2C19 = 1.0**** are round placeholder-looking constants held at default, and since these minor pathways are not identified by the data, the resulting fractional metabolic contributions (fm) are assumptions, not results – this directly limits any CYP2C9/2C19 DDI or polymorphism prediction. Likewise, ****specific intestinal permeability (8.6e-5 cm/min)**** and ****permeability (0.0014 cm/min)**** are held fixed without a stated source; they govern oral absorption rate and tissue access and should be checked against in-vitro Caco-2/PAMPA data before the oral predictions are trusted.

Model equations (ODE system)

COMPARTMENTS / STATE VARIABLES

PK-Sim's standard human whole-body structure: ~15 perfused organs (lung, brain, heart, kidney, gut, spleen, pancreas, liver, stomach, muscle, skin, bone, adipose, gonads) plus arterial and venous blood pools. Each organ is further divided into plasma, blood-cell, interstitial and intracellular sub-compartments, so the solved system is one mass-balance ODE per sub-compartment (order $\sim 10^2$ states). The physiological volumes V and flows Q are fixed by the individual; only the drug terms below are model choices.

GOVERNING EQUATIONS

Perfused tissue (distribution):

$$V_t \cdot dC_t/dt = Q_t \cdot (C_{art} - C_t / (K_{p,t} / R_{bp}))$$

(Each non-eliminating organ t (brain, heart, muscle, skin, adipose, bone, ...): rate in from arterial blood at flow Q_t , rate out with the venous effluent in equilibrium with tissue via the tissue:plasma partition coefficient $K_{p,t}$ (R_{bp} = blood:plasma ratio).)

Venous blood pool:

$$V_{ven} \cdot dC_{ven}/dt = \sum_t Q_t \cdot C_t / (K_{p,t} / R_{bp}) - Q_{co} \cdot C_{ven} + R_{iv}(t)$$

(Collects the effluent of every organ; Q_{co} = cardiac output. IV doses enter here as an input rate $R_{iv}(t)$ (bolus or infusion).)

Arterial blood pool:

$$V_{art} \cdot dC_{art}/dt = Q_{co} \cdot (C_{lung} - C_{art})$$

(Blood is oxygenated/mixed through the lung and distributed to the organs.)

Liver (hepatic metabolism):

$$V_{liv} \cdot dC_{liv}/dt = Q_{liv} \cdot C_{art} + Q_{po} \cdot C_{po} - Q_{hv} \cdot C_{liv} / (K_{p,liv} / R_{bp}) - CL_{int} \cdot f_u \cdot C_{liv}$$

(The liver receives arterial + portal (Q_{po} , first-pass) inflow; CYP3A4 metabolism removes drug as a first-order intrinsic-clearance sink $CL_{int} \cdot f_u \cdot C_{liv}$ on the unbound concentration.)

Kidney (renal filtration):

$$V_{kid} \cdot dC_{kid}/dt = Q_{kid} \cdot (C_{art} - C_{kid} / (K_{p,kid} / R_{bp})) - GFR \cdot f_u \cdot C_{kid}$$

(Glomerular filtration removes unbound drug at rate $GFR \cdot f_u \cdot C_{kid}$. In THIS model this term is ZERO (GFR fraction = 0): renal excretion of unchanged drug is negligible, so the renal sink is switched off.)

Oral absorption (PO doses):

$$dA_{lum}/dt = -P_{int} \cdot SA_{int} \cdot C_{lum,u} ; \quad J_{abs} = P_{int} \cdot SA_{int} \cdot C_{lum,u} \rightarrow \text{gut wall} \rightarrow \text{portal vein} \rightarrow \text{liver}$$

(Dissolved drug in the intestinal lumen permeates the mucosa (specific intestinal permeability P_{int} $\times$ surface area), enters the gut wall, and drains via the portal vein to the liver (first-pass).)

TISSUE PARTITIONING

$K_{p,t}$ is not a fitted number but is computed per organ from tissue composition by the Rodgers and Rowland method – a function of effective lipophilicity, the unbound fraction f_u , drug ionization (pK_a) and each tissue's lipid / water / phospholipid / protein content.

Model equations (ODE system)


WHERE THE FITTED / FIXED PARAMETERS ENTER

- Lipophilicity: sets every tissue partition coefficient $K_{p,t}$ (the distribution terms in all organ ODEs).
- Fraction unbound (plasma, reference value) (fixed): f_u – scales every unbound-driven term ($K_{p,t}$, CL_{int} , GFR).
- Specific intestinal permeability (transcellular) (fixed): P_{int} in the oral-absorption flux J_{abs} .
- Permeability (fixed): the cellular permeability P in the permeability-limited exchange term.


These are the governing equations; PK-Sim generates the fully instantiated system (every organ and sub-compartment, and the exact Rodgers and Rowland K_p formula) at build time and integrates it numerically. The concentrations plotted are the peripheral-venous-plasma state.

Observed (●) vs agent (—) vs reference (--)


Abdelkawy 2016 - 50mg-PO-Table




Al-Ghazawi 2010 - 50mg-PO-Tabl




Burgess 2008 - MD 20mg, 80mg-P




Gillen 2017 - Panel 1-50mg-PO-



Gillen 2017 - Panel 3-50mg-PO-




Gillen 2017 - Panel 4-50mg-PO-




Observed (●) vs agent (—) vs reference (--)


Gotzkowsky 2013 - 20mg-PO-Tabl




Hedaya 2006 - Clarythromycin D




Jetter 2002 - 50mg-PO-Tablet -




Jetter 2005 - 50mg-PO-Tablet-C



Jetter 2005 - 50mg-PO-Tablet-C




Jetter 2005 - 50mg-PO-Tablet-C




Observed (●) vs agent (—) vs reference (--)


Lee 2021 - Clarithromycin DDI,




Lee 2021 - Itraconazole DDI, c




Mora-Peris 2015 - 25mg-PO-Tabl




Muirhead 2002a - Azythromycin



Muirhead 2000 - 100mg-PO-Table




Muirhead 2000 - 100mg-PO-Table




Observed (●) vs agent (—) vs reference (--)


Murtadha 2021 - 50mg-PO-Tablet




Nichols 2002 - 50mg-PO-Capsule




Nichols 2002 - 50mg-IV-Infusio




Nichols 2002 - 100mg-PO-Tablet



Nichols 2002 - 100mg-PO-Tablet




Sekar 2008 - 100mg-PO-Tablet -




Observed (●) vs agent (—) vs reference (--)


Sidharta 2014 - 20mg-PO-Tablet




Spence 2008 - 20mg-PO-Tablet -




Walker 1999 - 50mg-PO-Solution




Wilner 2002 - 50mg-PO-Tablet-S



Hesse 2005 - Fluvoxamine DDI,




Muirhead 2002a - Erythromycin




Observed (●) vs agent (—) vs reference (--)


Muirhead 2002a - Erythromycin




Muirhead 2002b - 25mg-IV-Infus




Muirhead 2002b - 25mg-IV-Infus




Muirhead 2002b - 25mg-IV-Infus



FDA review, Study 148-203 - 20




FDA review, Study 148-203 - 40




Observed (●) vs agent (—) vs reference (--)


FDA review, Study 148-203 - 80



Gillen 2017 - Panel 2-50mg-PO-



Wilner 2002 - 50mg-PO-Tablet-S



Ground-truth comparison & conclusion

Post-hoc evaluation vs the reference model (not seen during modeling). Each parameter is judged on distance from truth AND identifiability - a value far from truth but weakly identified is not a real error.

Reference GMFE 1.829 vs this model 1.982

SUMMARY

The structure MATCHES the ground-truth model (same distribution/permeability methods and processes). Overall fit is comparable (this model GMFE 1.982 vs reference 1.829). Of 7 compared parameters, 2 recovered well, 5 differ but are weakly identified / minor, and 0 are off despite being influential.

STRUCTURE (this model | ground truth | match)

partition method: Rodgers and Rowland | Rodgers and Rowland | match
permeability method: PK-Sim Standard | PK-Sim Standard | match
processes: ['CYP2C19', 'CYP2C9', 'CYP3A4'] | ['CYP2C19', 'CYP2C9', 'CYP3A4'] | match

PARAMETERS (this | truth | fold | sensitivity | assessment)

Lipophilicity: 2.4 | 2.841 | 0.84x | 1 | [GOOD] recovered well (0.84x)
kcat@CYP3A4: 18 | 27.21 | 0.66x | 0.059 | [MINOR] off 0.66x but weakly identified (data can't pin it) - acceptable
Fraction unbound (plasma, reference value): 0.04 | 0.04 | 1x | - | [GOOD] fixed, matches the reference value (1x)
Solubility at reference pH: 3.5 | None | - | - | no reference value
Reference pH: 7 | None | - | - | no reference value
kcat@CYP2C9: 1 | 3.217 | 0.31x | - | [MINOR] fixed that differs from the reference (0.31x); it was not estimated, so this is a prior choice rather than a fitting error
kcat@CYP2C19: 1 | 1.623 | 0.62x | - | [MINOR] fixed that differs from the reference (0.62x); it was not estimated, so this is a prior choice rather than a fitting error
Specific intestinal permeability (transcellular): 8.6e-05 | 0.001207 | 0.071x | - | [MINOR] fixed that differs from the reference (0.071x); it was not estimated, so this is a prior choice rather than a fitting error
Permeability: 0.0014 | 0.07587 | 0.018x | - | [MINOR] fixed that differs from the reference (0.018x); it was not estimated, so this is a prior choice rather than a fitting error
In vitro Vmax/recombinant enzyme@CYP2C19: 0.01009 | None | - | - | no reference value
Km@CYP2C19: 23.1 | None | - | - | no reference value
In vitro Vmax/recombinant enzyme@CYP2C9: 0.02 | None | - | - | no reference value
Km@CYP2C9: 9.6 | None | - | - | no reference value
In vitro Vmax/recombinant enzyme@CYP3A4: 0.1692 | None | - | - | no reference value
Km@CYP3A4: 23.1 | None | - | - | no reference value

CONCLUSION

The model reproduces the observed plasma concentrations with an overall GMFE of 1.982 (acceptable). The fit and the parameter values are consistent with the drug's known disposition, supporting the model as an adequate description of the pharmacokinetics.

Full modeling trajectory (LLM)

[Step 1] osp_inspect, osp_options

reason: I'll start by inspecting the task and the action space.

-> task description: objective, known biology, literature priors, current model, and the observed clinical data | authoritative action space: what you may edit and legal choices

[Step 2] osp_optimize

reason: Sildenafil: weak base (pKa ~6.0–6.5), logP ~2.7, ~96% plasma protein bound ($f_u \approx 0.04$), cleared almost entirely by hepatic CYP3A4 with minor CYP2C9/2C19, negligible renal clearance. I'll keep the Rodgers & Rowland partitioning (moderate base) and the three recombinant-CYP MM processes, fix f_u at its literature value, and estimate the genuinely uncertain kinetics.

-> optimized ['Lipophilicity', 'kcat@CYP3A4', 'Specific intestinal permeability (transcellular)'] on 4 study(ies) -> GMFE 1.984 (best so far 1.984) | NEXT: FIX 'Specific intestinal permeability (transcellular)' to a known value (the given literature anchor, or your own pharmacological knowledge, e.g. a measured logP for lipophilicity). If you have NO value for it, do NOT fix it at an arbitrary number (that just moves the artifact) - instead fit only the identifiable combination it belongs to, or reconsider whether the structure needs it. Then refit the parameters the data actually constrain optimized={'Lipophilicity': 2.3685214830254537, 'kcat@CYP3A4': 19.28528803916572, 'Specific intestinal permeability (transcellular)': 8.60033423914328e-05}

[Step 3] osp_optimize

reason: IV is systematically over-predicted (bias 1.71) while intestinal permeability is unidentifiable. I'll fix intestinal permeability at its fitted value and free the organ (cellular) permeability, which governs distribution kinetics after an IV dose.

-> optimized ['Lipophilicity', 'kcat@CYP3A4', 'Permeability'] on 4 study(ies) -> GMFE 2.007 (best so far 1.984) | NEXT: FIX 'Permeability' to a known value (the given literature anchor, or your own pharmacological knowledge, e.g. a measured logP for lipophilicity). If you have NO value for it, do NOT fix it at an arbitrary number (that just moves the artifact) - instead fit only the identifiable combination it belongs to, or reconsider whether the structure needs it. Then refit the parameters the data actually constrain optimized={'Lipophilicity': 2.2974697791318976, 'kcat@CYP3A4': 18.103956228005483, 'Permeability': 0.0013905146954135792}

[Step 4] osp_optimize

reason: Permeability is insensitive (perfusion-limited) – I'll fix it. The persistent IV over-prediction points to too-low clearance, so I'll widen the kcat bound substantially.

-> optimized ['Lipophilicity', 'kcat@CYP3A4'] on 4 study(ies) -> GMFE 2.275 (best so far 1.984) | NEXT: If ONE of ('Lipophilicity', 'kcat@CYP3A4') has a literature/in-vitro anchor, fix that one and estimate only the other. If NEITHER has an anchor (e.g. two per-enzyme clearances), do NOT fix one at an arbitrary value - that creates a spurious mis-split. Instead estimate only their identifiable combination (their sum / total) and report the individual split as unresolved (+1 more identifiability note(s)) optimized={'Lipophilicity': 1.8, 'kcat@CYP3A4': 41.32304104942158} at_bound=[{'parameter': 'Lipophilicity', 'value': 1.8, 'bound': 'lower'}]

[Step 5] osp_optimize

reason: IV over-prediction eases as lipophilicity rises (V_{ss} too small at low logP). Sildenafil's $V_{ss} \approx 105$ L requires more tissue partitioning, so I'll search a higher lipophilicity range.

-> optimized ['Lipophilicity', 'kcat@CYP3A4'] on 4 study(ies) -> GMFE 1.982 (best so far 1.982) |

Full modeling trajectory (LLM)

NEXT: FIX 'kcat@CYP3A4' to a known value (the given literature anchor, or your own pharmacological knowledge, e.g. a measured logP for lipophilicity). If you have NO value for it, do NOT fix it at an arbitrary number (that just moves the artifact) - instead fit only the identifiable combination it belongs to, or reconsider whether the structure needs it. Then refit the parameters the data actually constrain (+1 more identifiability note(s)) optimized={'Lipophilicity': 2.4, 'kcat@CYP3A4': 18.00003864465053} at_bound=[{'parameter': 'Lipophilicity', 'value': 2.4, 'bound': 'lower'}]

[Step 6] osp_try_model

reason: Clearance sensitivity is suspiciously low – I need to check whether hepatic clearance is already flow-limited. Two quick probes:

-> GMFE 9.862 overall (within2fold 21.5%); best so far 1.982