Program 1

Write a Python program to print "Hello Python".

In [1]:
```
1 print("Hello Python")
```

```
Hello Python
```

Program 2

Write a Python program to do arithmetical operations addition and division.

In [2]:
```
1 # Addition
2 num1 = float(input("Enter the first number for addition: "))
3 num2 = float(input("Enter the second number for addition: "))
4 sum_result = num1 + num2
5 print(f"sum: {num1} + {num2} = {sum_result}")
```

```
Enter the first number for addition: 5
Enter the second number for addition: 6
sum: 5.0 + 6.0 = 11.0
```

In [3]:
```
1 # Division
2 num3 = float(input("Enter the dividend for division: "))
3 num4 = float(input("Enter the divisor for division: "))
4 if num4 == 0:
5     print("Error: Division by zero is not allowed.")
6 else:
7     div_result = num3 / num4
8     print(f"Division: {num3} / {num4} = {div_result}")
```

```
Enter the dividend for division: 25
Enter the divisor for division: 5
Division: 25.0 / 5.0 = 5.0
```

Program 3

Write a Python program to find the area of a triangle.

In [4]:
```
1 # Input the base and height from the user
2 base = float(input("Enter the length of the base of the triangle: "))
3 height = float(input("Enter the height of the triangle: "))
4 # Calculate the area of the triangle
5 area = 0.5 * base * height
6 # Display the result
7 print(f"The area of the triangle is: {area}")
```

```
Enter the length of the base of the triangle: 10
Enter the height of the triangle: 15
The area of the triangle is: 75.0
```

Program 4

Write a Python program to swap two variables.

In [5]:

```
# Input two variables
a = input("Enter the value of the first variable (a): ")
b = input("Enter the value of the second variable (b): ")
# Display the original values
print(f"Original values: a = {a}, b = {b}")
# Swap the values using a temporary variable
temp = a
a = b
b = temp
# Display the swapped values
print(f"Swapped values: a = {a}, b = {b}")
```

```
Enter the value of the first variable (a): 5
Enter the value of the second variable (b): 9
Original values: a = 5, b = 9
Swapped values: a = 9, b = 5
```

Program 5

Write a Python program to generate a random number.

In [6]:

```
import random
print(f"Random number: {random.randint(1, 100)}")
```

```
Random number: 89
```

Program 6

Write a Python program to convert kilometers to miles.

In [7]:

```
kilometers = float(input("Enter distance in kilometers: "))

# Conversion factor: 1 kilometer = 0.621371 miles
 conversion_factor = 0.621371

miles = kilometers * conversion_factor


print(f"{kilometers} kilometers is equal to {miles} miles")
```

```
Enter distance in kilometers: 100
100.0 kilometers is equal to 62.137100000000004 miles
```

Program 7

Write a Python program to convert Celsius to Fahrenheit.

In [8]:
```
1 celsius = float(input("Enter temperature in Celsius: "))
2
3 # Conversion formula: Fahrenheit = (Celsius * 9/5) + 32 fahrenheit =
4 (celsius * 9/5) + 32
5
6 print(f"{celsius} degrees Celsius is equal to {fahrenheit} degrees Fah
```

```
Enter temperature in Celsius: 37
37.0 degrees Celsius is equal to 98.6 degrees Fahrenheit
```

Program 8

Write a Python program to display calendar.

In [9]:
```
1 import calendar
2
3 year = int(input("Enter year: ")) month = int(input("Enter month: "))
4
5 cal = calendar.month(year, month)
6
7 print(cal)
```

```
Enter year: 2023
Enter month: 11
   November 2023
Mo Tu We Th Fr Sa Su
            1  2  3  4  5
 6  7  8  9 10 11 12
13 14 15 16 17 18 19
20 21 22 23 24 25 26
27 28 29 30
```

Program 9

Write a Python program to solve quadratic equation.

The standard form of a quadratic equation is:

$$ax2 + bx + c = 0$$

where

a, b and c are real numbers and

$a \neq 0$

The solutions of this quadratic equation is given by:

$$(-b \pm (b^2 - 4ac)1/2)/(2a)$$

In [10]:

```
import math

# Input coefficients a = float(input("Enter coefficient a: ")) b =
float(input("Enter coefficient b: ")) c = float(input("Enter
coefficient c: "))

# Calculate the discriminant discriminant = b**2 - 4*a*c

# Check if the discriminant is positive, negative, or zero if
discriminant > 0:

# Two real and distinct roots root1 = (-b +
math.sqrt(discriminant)) / (2*a) root2 = (-b -
math.sqrt(discriminant)) / (2*a) print(f"Root 1: {root1}")
print(f"Root 2: {root2}")


elif discriminant == 0:
    # One real root (repeated)
    root = -b / (2*a)
    print(f"Root: {root}")
else:
    # Complex roots real_part = -b / (2*a) imaginary_part
    = math.sqrt(abs(discriminant)) / (2*a) print(f"Root 1:
    {real_part} + {imaginary_part}i") print(f"Root 2:
    {real_part} - {imaginary_part}i")
```

```
Enter coefficient a: 1
Enter coefficient b: 4
Enter coefficient c: 8
Root 1: -2.0 + 2.0i
Root 2: -2.0 - 2.0i
```

Program 10

Write a Python program to swap two variables without temp variable.

In [11]:

```
a = 5 b = 10

# Swapping without a temporary variable
a, b = b, a


print("After swapping:")
print("a =", a)
print("b =", b)
```

```
After swapping:
a = 10
b = 5
```

Program 11

Write a Python Program to Check if a Number is Positive, Negative or Zero.

In [12]:

```
num = float(input("Enter a number: "))
if num > 0:
    print("Positive number")
elif num == 0:
    print("Zero")
else:
    print("Negative number")
```

```
Enter a number: 6.4
Positive number
```

Program 12

Write a Python Program to Check if a Number is Odd or Even.

In [13]:

```
num = int(input("Enter a number: "))

i f    n u m % 2    = =    0 :
    print("This is a even number")
else:
    print("This is a odd number")
```

```
Enter a number: 3
This is a odd number
```

Program 13

Write a Python Program to Check Leap Year.

In [14]:

```
year = int(input("Enter a year: "))

# divided by 100 means century year (ending with 00) # century year
divided by 400 is leap year if (year % 400 == 0) and (year % 100 == 0):

    print("{0} is a leap year".format(year))

# not divided by 100 means not a century year # year divided by 4
is a leap year elif (year % 4 ==0) and (year % 100 != 0):

    print("{0} is a leap year".format(year))

# if not divided by both 400 (century year) and 4 (not century year)
# year is not leap year
else:
    print("{0} is not a leap year".format(year))
```

```
Enter a year: 2024
2024 is a leap year
```

Program 14

Write a Python Program to Check Prime Number.

Prime Numbers:

A prime number is a whole number that cannot be evenly divided by any other number except for 1 and itself. For example, 2, 3, 5, 7, 11, and 13 are prime numbers because they cannot be divided by any other positive integer except for 1 and their own value.

In [15]:

```
num = int(input("Enter a number: "))

# define a flag variable
flag = False

if num == 1:
    print(f"{num}, is not a prime number")
elif num > 1:
    # check for factors
    for i in range(2, num):
        if (num % i) == 0:
            flag = True     # if factor is found, set flag to True
            # break out of loop
            break

    # check if flag is True
if flag:
    print(f"{num}, is not a prime number")
else:
    print(f"{num}, is a prime number")
```

```
Enter a number: 27
27, is not a prime number
```

Program 15

Write a Python Program to Print all Prime Numbers in an Interval of 1-10.

In [20]:

```
# Python program to display all the prime numbers within an interval

lower = 1 upper = 10

print("Prime numbers between", lower, "and", upper, "are:")

for num in range(lower, upper + 1):

   # all prime numbers are greater than 1
   if num > 1:
        for i in range(2, num):
            if (num % i) == 0:
                break
        else:
            print(num)
```

```
Prime numbers between 1 and 10 are:
2 3 5 7
```

Program 16

Write a Python Program to Find the Factorial of a Number.

In [21]:

```
num = int(input("Enter a number: "))
factorial = 1
if num <0:
    print("Factirial does not exist for negative numbers")
elif num == 0:
    print("Factorial of 0 is 1")
else:
    for i in range(1, num+1):
        factorial = factorial*i
    print(f'The factorial of {num} is {factorial}')
```

```
Enter a number: 4
The factorial of 4 is 24
```

Program 17

Write a Python Program to Display the multiplication Table.

In [22]:
```
num = int(input("Display multiplication table of: "))

for i in range(1, 11):
    print(f"{num} X {i} = {num*i}")
```

```
Display multiplication table of: 19
19 X 1 = 19 19 X 2 = 38 19 X 3 = 57
19 X 4 = 76 19 X 5 = 95 19 X 6 =
114 19 X 7 = 133 19 X 8 = 152 19 X
9 = 171 19 X 10 = 190
```

Program 18

Write a Python Program to Print the Fibonacci sequence.

Fibonacci sequence:

The Fibonacci sequence is a series of numbers where each number is the sum of the two preceding ones, typically starting with 0 and 1. So, the sequence begins with 0 and 1, and the next number is obtained by adding the previous two numbers. This pattern continues indefinitely, generating a sequence that looks like this:

0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, and so on.

Mathematically, the Fibonacci sequence can be defined using the following recurrence relation:

$F(0)=0\ F(1)=1\ F(n)=F(n-1)+F(n-2) for n>1$

In [23]:
```
nterms = int(input("How many terms? "))

# first two terms n1, n2 = 0, 1 count = 0

# check if the number of terms is valid if nterms
<= 0:



    print("Please enter a positive integer")
# if there is only one term, return n1
elif nterms == 1:
    print("Fibonacci sequence upto",nterms,":")
    print(n1)
# generate fibonacci sequence
else:
    print("Fibonacci sequence:")
    while count < nterms:
        print(n1)
        nth = n1 + n2
        # update values
         n1 = n2
         n2 = nth
         count += 1
```

```
How many terms? 10 Fibonacci
sequence: 0 1 1 2 3 5 8 13 21 34
```

Program 19

Write a Python Program to Check Armstrong Number?

Armstrong Number:

It is a number that is equal to the sum of its own digits, each raised to a power equal to the number of digits in the number.

For example, let's consider the number 153:

- It has three digits (1, 5, and 3).
- If we calculate 13 + 53+33, we get 1+125+27, which is equal to 153.

So, 153 is an Armstrong number because it equals the sum of its digits raised to the power of the number of digits in the number.
Another example is 9474:

- It has four digits (9, 4, 7, and 4).
    - If we calculate $94+44+7^4+4^4$, we get 6561 + 256 + 2401 + 256, which is also equal to 9474.

Therefore, 9474 is an Armstrong number as well.

In [25]:

```
num = int(input("Enter a number: "))

# Calculate the number of digits in num num_str = str(num)
num_digits = len(num_str)

# Initialize variables sum_of_powers = 0 temp_num = num

# Calculate the sum of digits raised to the power of num_digits

while temp_num > 0:



    digit = temp_num % 10
    sum_of_powers += digit ** num_digits
    temp_num //= 10

# Check if it's an Armstrong number
if sum_of_powers == num:
    print(f"{num} is an Armstrong number.")
else:
    print(f"{num} is not an Armstrong number.")

```

```
Enter a number: 9474
9474 is an Armstrong number.
```

In [26]:

```
# Input the interval from the user lower = int(input("Enter
the lower limit of the interval: ")) upper = int(input("Enter
the upper limit of the interval: "))


for num in range(lower, upper + 1): # Iterate through the numbers i
    order = len(str(num))  # Find the number of digits in 'num'
    temp_num = num
    sum = 0

    while temp_num > 0:
        digit = temp_num % 10
        sum += digit ** order
        temp_num //= 10

    # Check if 'num' is an Armstrong number
    if num == sum: print(num)
```

```
Enter the lower limit of the interval: 10
Enter the upper limit of the interval: 1000
153 370 371 407
```

Program 21

Write a Python Program to Find the Sum of Natural Numbers.

Natural numbers are a set of positive integers that are used to count and order objects. They are the numbers that typically start from 1 and continue indefinitely, including all the whole numbers greater than 0. In mathematical notation, the set of natural numbers is often denoted as "N" and can be expressed as:

$$N = 1,2,3,4,5,6,7,8,...$$

In [27]:

```
limit = int(input("Enter the limit: "))

# Initialize the sum sum = 0

# Use a for loop to calculate the sum of natural numbers
for i in range(1, limit + 1):

    sum += i

# Print the sum print("The sum of natural numbers up to",
limit, "is:", sum)
```

```
Enter the limit: 10
The sum of natural numbers up to 10 is: 55
```