

UNIT – III, Module – 1

Lecture 26: Graphs & Trees

[ Special graphs; Bipartite graphs; Adjacency matrix, power; Adjacency list; Incidence, degree matrices; Graph isomorphism; Graph invariants ]

Dr. Suddhasil De

---

---

---

---

---

---

---

Notation table

Symbol / Notation	Meaning
$K_n$	Complete undirected graph of $n$ vertices.
$C_n$	Cycle graph of $n$ vertices.
$W_n$	Wheel graph of $n + 1$ vertices.
$Q_n$	$n$ -dimensional hypercube graph (also called $n$ -cube graph).
$K_{m,n}$	Complete bipartite graph with $m$ -vertex and $n$ -vertex partitions.
$Adj_G$	Adjacency matrix of simple graph $G$ .
$Adj_G^r$	$r$ -th power adjacency matrix of simple graph $G$ .
$I_G$	Incidence matrix of simple graph $G$ .
$\tilde{I}_G$	Oriented incidence matrix of simple graph $G$ .
$D_G$	Degree matrix of simple graph $G$ .
$L_G$	Laplacian matrix of simple graph $G$ .

Graph Theory

Dept. of CSE, NITP  
2

Dr. Suddhasil De

---

---

---

---

---


---

---

Special graphs

- Special graphs:
  - Property::  $K_n$  ( $n > 0$ ): **complete** undirected **graph**  $G = (V, E)$  with  $|V| = n$  vertices, where every distinct vertex pair adjacent to each other (i.e., simple undirected graph  $G$ , with exactly one edge between every pair of distinct vertices).

Figures of  $K_n$ , for  $1 \leq n \leq 6$



- Property::  $K_n \rightarrow$  applicable to unlabeled graph.

[Ref: Kenneth H. Rosen, Discrete Mathematics and Its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory

Dept. of CSE, NITP  
3

Dr. Suddhasil De

---

---

---

---

---

---

---

Special graphs

- Special graphs:
  - Property: Degree of every vertex in  $K_n$  ( $n > 0$ ):  $(n - 1)$ .
  - Property: (Theorem): Number of edges in  $K_n$  ( $n > 0$ ):  $\binom{n}{2} = \frac{1}{2} \cdot n \cdot (n - 1)$ .

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

---

---

---

---

---


---

---

Special graphs

- Special graphs:
  - Property:  $C_n$  ( $n \geq 3$ ): cycle graph  $G = (V, E)$ , being simple undirected graph with  $|V| = n$  vertices, vertex set  $V = \{v_1, v_2, v_3, \dots, v_{n-1}, v_n\}$ , and edge set  $E = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}, \{v_n, v_1\}\}$ .

Figures of  $C_n$ , for  $3 \leq n \leq 6$



[Ref: Kenneth H. Rosen, Discrete Mathematics and its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

---

---

---

---

---


---

---

Special graphs

- Special graphs:
  - Property:  $W_n$  ( $n \geq 3$ ): wheel graph  $G = (V, E)$ , being simple undirected graph with  $|V| = n + 1$ , obtained by adding new vertex to  $C_n$  and connecting this new vertex to each of  $n$  vertices in  $C_n$  by new edges, i.e.,  $W_n = C_n + K_1$ .

Figures of  $W_n$ , for  $3 \leq n \leq 6$



[Ref: Kenneth H. Rosen, Discrete Mathematics and its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

---

---

---

---

---

---


---

2

### Special graphs

- Special graphs:
  - Property:  $Q_n$  ( $n > 0$ ):  $n$ -dimensional hypercube graph (also called,  $n$ -cube graph)  $G = (V, E)$ , being simple undirected graph with vertices representing  $2^n$  bit strings of length  $n$  and adjacent vertices to differ by exactly one bit position's value in bit strings represented by them.

Figures of  $Q_n$ , for  $1 \leq n \leq 3$



[Ref: Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory


Dept. of CSE, NITP

Dr. Suddhasil De

7

### Special graphs

- Bipartite graph: possibility of bi-partitioning of vertex set of graph.
  - Property: **Bipartite graph**  $G = (V, E)$ : Vertex set  $V$  to be bi-partitioned into two disjoint subsets  $V_1$  and  $V_2$ , such that every edge in  $G$  to join vertex  $u \in V_1$  with vertex  $v \in V_2$ .
  - Property: Bipartite graph: also called bigraph, bicolored graph, pair graph.
  - Property: Partite sets:  $V_1$  and  $V_2$  in bipartite graph  $G = (V, E)$ .



[Ref: Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

8

### Special graphs

- Bipartite graph:
  - Property: Bipartite graph to have — (i) no self-loop, (ii) parallel edges between vertices to be represented by single edge.

Discrete Mathematics


Dept. of CSE, NITP

Dr. Suddhasil De

9

Special graphs

- Bipartite graph:
  - Property:  $K_{m,n}$  ( $m, n > 0$ ): **complete bipartite graph**  $G = (V, E)$ , with bi-partitioned  $V = V_1 \cup V_2$ ,  $|V_1| = m$ ,  $|V_2| = n$ , and  $V_1 \cap V_2 = \emptyset$ , such that edge present between every vertex of  $V_1$  to every vertex of  $V_2$ .



[Ref: Kenneth H. Rosen, Discrete Mathematics and its Applications, Eighth edition, McGraw-Hill Education, 2019.]  
Discrete Mathematics Dept. of CSE, NITP Dr. Suddhasil De  
10

---

---

---

---

---

---

---

Special graphs

- Bipartite graph:
  - Property:  $K_{1,n}$  ( $n > 0$ ): also called star.
  - Property:  $K_{1,3}$  : also called claw, cherry etc.
  - Property: (**Theorem**): Number of edges in  $K_{m,n}$  ( $m, n > 0$ ) =  $(m \cdot n)$ .

Discrete Mathematics Dept. of CSE, NITP Dr. Suddhasil De  
11

---

---

---

---

---

---

---

Graph representations

- Graph representation: convenient and simplified representation of graphs for carrying out graph algorithms; common approaches —
  - adjacency matrix;
  - adjacency list;
  - incidence matrix;
  - degree matrix.

Discrete Mathematics Dept. of CSE, NITP Dr. Suddhasil De  
12

---

---

---

---

---

---

---

### Graph representations

- Graph representation:
  - Property: Adjacency matrix of simple graph  $G = (V, E)$ :  $n \times n$  zero-one matrix  $Adj_G = (a_{ij})_{1 \leq i, j \leq n}$ , where  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ , such that
 
$$a_{ij} = \begin{cases} 1, & \text{if } \{v_i, v_j\} \in E \\ 0, & \text{otherwise.} \end{cases}$$
  - Property: Adjacency matrix of graph based on ordering chosen for vertices in its vertex set.
  - Property: (Theorem):  $n!$  different adjacency matrices for graph with  $n$  vertices.

[Ref: Kenneth H. Rosen, Discrete Mathematics and Its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

13

### Graph representations

- Graph representation:
  - Property: For simple graph  $G = (V, E)$ ,  $|V| = n$ , symmetric adjacency matrix  $Adj_G = (a_{ij})_{1 \leq i, j \leq n} = (a_{ji})_{1 \leq i, j \leq n}$ ,  $a_{ii} = 0$  (no self-loop).
  - Property: Adjacency matrix of multigraph  $G = (V, E)$ ,  $|V| = n$ , [also applicable to pseudograph]:
 
$$Adj_G = (a'_{ij})_{1 \leq i, j \leq n}$$
, where
 
$$a'_{ij} = \begin{cases} \text{multiplicity}(v_i, v_j), & \text{if } \{v_i, v_j\} \in E \\ 0, & \text{if } \{v_i, v_j\} \notin E. \end{cases}$$

[Ref: Kenneth H. Rosen, Discrete Mathematics and Its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

14

### Graph representations

- Graph representation:
  - Property: Adjacency matrix: also called **connection matrix**.
  - Property: For disconnected undirected graph  $G' = (V', E')$ , possibility of two components  $g'_1 = (v'_1, e'_1)$ ,  $g'_2 = (v'_2, e'_2)$ , where  $g'_1 \cup g'_2 = G'$ ,  $v'_1 \cap v'_2 = \emptyset$ , if and only if possibility to partition  $Adj_{G'}$  as:
 
$$Adj_{G'} = \begin{bmatrix} Adj_{g'_1} & | & [0] \\ \hline [0] & | & Adj_{g'_2} \end{bmatrix}$$
, where  $[0]$  = zero matrix.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

15

### Graph representations

- Graph representation:
  - Property: Power of adjacency matrix of simple graph  $G = (V, E)$ :  
 $Adj_G^2 = Adj_G \cdot Adj_G = ({}^2a_{ij})_{1 \leq i, j \leq n}$ , where  
 if  $i \neq j$ ,  ${}^2a_{ij}$  = number of 1's in product of row  $i$  and column  $j$   
 = number of 1's in product of row  $i$  and row  $j$  (due to symmetry)  
 = number of positions in both row  $i$  and row  $j$  to have 1's  
 = number of vertices adjacent to both  $i$ -th vertex  $v_i$  and  $j$ -th vertex  $v_j$   
 = number of different paths between  $v_i$  and  $v_j$ , where  $len(path P)=2$ ,  
 if  $i = j$ ,  ${}^2a_{ij}$  = number of 1's in row  $i$  (or column  $i$ )  
 = degree of vertex  $v_i$ ,  $deg(v_i)$  (provided no self-loops in  $G$ ).


Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

16

### Graph representations

- Graph representation:
  - Property: Power of adjacency matrix of simple graph  $G = (V, E)$ :  
 example.  

 $Adj_G =$ 

	$v_1$	$v_2$	$v_3$	$v_4$	$v_5$	$v_6$
$v_1$	0	1	0	0	1	1
$v_2$	1	0	0	1	1	0
$v_3$	0	0	0	1	0	0
$v_4$	0	1	1	0	1	1
$v_5$	1	1	0	1	0	0
$v_6$	1	0	0	1	0	0

 $Adj_G^2 =$ 

3	1	0	3	1	0
1	3	1	1	2	2
0	1	1	0	1	1
3	1	0	4	1	0
1	2	1	1	3	2
0	2	1	0	2	2

 Two paths between  $v_5$  and  $v_2$  where  $len(path)=2$   
 $deg(v_4) = 4$

[Ref: Narsingh Deo, Graph Theory with Applications to Engineering and Computer Science, Prentice-Hall, 1974.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

17

### Graph representations

- Graph representation:
  - Property: Power of adjacency matrix of simple graph  $G = (V, E)$ :  
 $Adj_G^3 = Adj_G^2 \cdot Adj_G = ({}^3a_{ij})_{1 \leq i, j \leq n}$ , where  
 if  $i \neq j$ ,  ${}^3a_{ij}$  = product of row  $i$  of  $Adj_G^2$  and column/row  $j$  of  $Adj_G$   
 $= \sum_{k=1}^n ({}^2a_{ik} \cdot a_{kj}) = \sum_{k=1}^n (\text{number of all different paths of length 3 from } v_i \text{ to } v_j \text{ via } v_k)$   
 = number of different paths between  $v_i$  and  $v_j$ , where  $len(path P)=3$ ,  
 if  $i = j$ ,  ${}^3a_{ij}$  = twice of number of different circuits of length 3 (i.e., triangles) passing through  $v_i$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

18

### Graph representations

- Graph representation:
  - Property:: (**Theorem**): For any undirected graph  $G = (V, E)$  [also applicable to directed graphs], with adjacency matrix  $Adj_G$  for  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ ,  $1 \leq i, j \leq n$ , number of different paths of length  $r$  ( $r \in \mathbb{Z}^+$ ) from  $v_i$  to  $v_j$  (i.e., count of different edge sequences of  $r$  edges between them) to be obtained from  $(i, j)$ -th entry, i.e.,  ${}^r a_{ij} = \sum_{k=1}^n ({}^{r-1} a_{ik} \cdot a_{kj})$  of  $Adj_G^r$ , where  $Adj_G^r = Adj_G^{r-1} \cdot Adj_G$ ,  $Adj_G^r = ({}^r a_{ij})_{1 \leq i, j \leq n}$ ,  $Adj_G^{r-1} = ({}^{r-1} a_{ij})_{1 \leq i, j \leq n}$ ,  $Adj_G = (a_{ij})_{1 \leq i, j \leq n}$ .
  - Proof**: Proof by mathematical induction.[Ros19:723-724;Deo74:161]

Discrete Mathematics

Dept. of CSE, NITP  
19

Dr. Suddhasil De

### Graph representations

- Graph representation:
  - Property:: (**Corollary**): For any connected graph  $G = (V, E)$ , with adjacency matrix  $Adj_G$  for  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ ,  $1 \leq i, j \leq n$ , distance between vertices  $v_i$  and  $v_j$ ,  $d(v_i, v_j) = \ell$  (for  $i \neq j$ ), if and only if  $\ell$  being smallest integer for which  ${}^\ell a_{ij} \neq 0$  in  $Adj_G^\ell$ .
  - Property:: (**Corollary**): If any undirected graph  $G = (V, E)$ , with adjacency matrix  $Adj_G$  for  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ , and  $\mathcal{A} = Adj_G + Adj_G^2 + Adj_G^3 + \dots + Adj_G^{n-1}$ , where  $\mathcal{A} = (\alpha_{ij})_{1 \leq i, j \leq n}$ ,  $\alpha_{ij} \in \mathbb{Z}$ , then  $G$  to become disconnected if and only if  $\exists \alpha_{ij} (\alpha_{ij} = 0)$ .

Discrete Mathematics

Dept. of CSE, NITP  
20

Dr. Suddhasil De

### Graph representations

- Graph representation:
  - Benefit of adjacency matrix: efficient for simple dense graph (i.e., graph with many edges, such as graph with more than half of all possible edges), due to very few numerical comparisons to decide presence of edges.
  - Shortcomings of adjacency matrix: special sparse matrix related algorithms required for adjacency matrix of simple sparse graph.


Discrete Mathematics

Dept. of CSE, NITP  
21

Dr. Suddhasil De

### Graph representations

- Graph representation:
  - Property: Adjacency list of simple graph  $G = (V, E)$ : tabular view of vertices and their corresponding list (like, array) of adjacent vertices.



- Benefit of adjacency list: efficient for simple sparse graph.

[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics


Dept. of CSE, NITP

Dr. Suddhasil De

22

### Graph representations

- Graph representation:
  - Property: **Incidence matrix** of simple graph  $G = (V, E)$  [also applicable to multigraph, pseudograph]:  $n \times m$  zero-one matrix  $I_G = (b_{ij})_{1 \leq i \leq n, 1 \leq j \leq m}$ , where  $V = \{v_1, v_2, \dots, v_n\}$ ,  $E = \{e_1, e_2, \dots, e_m\}$ ,  $|V| = n$ ,  $|E| = m$ , such that  $b_{ij} = \begin{cases} 1, & \text{if } e_j \text{ incident with } v_i \\ 0, & \text{otherwise.} \end{cases}$
  - Property: For incidence matrix of multigraph, pseudograph,  $E$  to contain parallel edges, self-loops also.



[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics


Dept. of CSE, NITP

Dr. Suddhasil De

23

### Graph representations

- Graph representation:
  - Property: Incidence matrix of pseudograph: example.



[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

24



Graph representations

- Graph representation:
- Property:: **Oriented incidence matrix** of simple graph  $G = (V, E): n \times m$  matrix  $\tilde{I}_G = (\mathbf{b}'_j)_{1 \leq j \leq m}$ , where  $V = \{v_1, v_2, \dots, v_n\}$ ,  $E = \{e_1, e_2, \dots, e_m\}$ ,  $|V| = n$ ,  $|E| = m$ , and  $\mathbf{b}'_j$  to be obtained from  $j$ -th column of  $I_G$  by arbitrarily replacing anyone of two 1's in that column by  $-1$ .

---

---

---

---

---

---

---

---

Graph representations

- Graph representation:
- Property:: **Degree matrix** of simple graph  $G = (V, E): n \times n$  diagonal matrix  $D_G = (\deg_{ij})_{1 \leq i, j \leq n}$ , where  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ , and  $\deg_{ij} = \begin{cases} \deg(v_i), & \text{if } i = j \\ 0, & \text{otherwise.} \end{cases}$
- Property:: **Laplacian matrix** of simple graph  $G = (V, E): n \times n$  matrix  $L_G = D_G - Adj_G = (L_{ij})_{1 \leq i, j \leq n}$ , where  $V = \{v_1, v_2, \dots, v_n\}$ ,  $|V| = n$ , and  $L_{ij} = \begin{cases} \deg(v_i), & \text{if } i = j \\ -1, & \text{if } i \neq j, \text{ and } v_i \text{ adjacent to } v_j \\ 0, & \text{otherwise.} \end{cases}$

---

---

---

---

---

---

---

---

Graph representations

- Graph representation:
- Property:: (**Theorem**): For simple connected undirected graph  $G = (V, E)$  with  $|V| = n$  vertices,  $\tilde{I}_G \cdot (\tilde{I}_G^T) = D_G - Adj_G = L_G$ .  
[Above equation indicating relation between adjacency and incidence matrices.]

---

---

---

---

---


---

---

---

## Graph

- Special graphs:
  - Property: **Graph isomorphism**:  $G = (V, E)$  and  $G' = (V', E')$  to become equivalent (and called **isomorphic** to each other) on establishing one-to-one correspondence between their vertices and between their edges preserving incidence relations (for simple graphs  $G, G'$ ).



[Ref: Narsingh Deo, Graph Theory with Applications to Engineering and Computer Science, Prentice-Hall, 1974.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

28

## Graph

- Special graphs:
  - Property: Isomorphic graphs to be same, perhaps drawn differently, with different labels (i.e., names) of their vertices and edges.



- Condition for isomorphism between  $G = (V, E)$  and  $G' = (V', E')$ :  
 $f: V \rightarrow V'$ , with  $\forall e \in E ((e = \{u, v\}) \rightarrow (\exists e' \in E' (e' = \{f(u), f(v)\})))$ ,  
 where,  $u, v \in V, f(u), f(v) \in V'$ .

[Ref: Narsingh Deo, Graph Theory with Applications to Engineering and Computer Science, Prentice-Hall, 1974.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

29

## Graph

- Special graphs:
  - Property: **Graph invariants** of isomorphic simple graphs —
    - same number of vertices, due to one-to-one correspondence between vertex sets of isomorphic simple graphs;
    - same number of edges, again due to one-to-one correspondence between edges implied from one-to-one correspondence between vertices of isomorphic simple graphs;
    - same degrees of vertices in isomorphic simple graphs;
    - existence of circuit of same length (say,  $k > 2$ ) in isomorphic simple graphs;

(contd. to next slide)

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

30

Graph

- Special graphs:
  - Property:: **Graph invariants** of isomorphic simple graphs (contd.) —
    - v. existence of path traversing through all vertices of isomorphic simple graphs, so that corresponding vertices in two graphs to have same degree.
  - Property:: Determining graph isomorphism based on —
    - (a)  $Adj_G, Adj_{G'}$ ,
    - (b)  $Adj_G^T, Adj_{G'}^T$ , to justify satisfying these graph invariants.

---

---

---

---

---

---

---

Summary

- Focus: Special graphs, graph representation, isomorphism.
- Special graphs  $K_n, C_n, W_n, Q_n$ , with examples.
- Bipartite graphs and  $K_{m,n}$ , with examples.
- Graph representations.
- Adjacency matrix of simple graphs, multigraphs, disconnected graphs, with examples.
- Power of adjacency matrix of simple graph, and its properties and related theorems, with examples.
- Adjacency list of simple graph, with examples.

---

---

---

---

---

---

---

Summary

- Incidence matrix of simple graphs, multigraphs, with examples.
- Oriented incidence matrix of simple undirected graphs.
- Degree matrix of simple graphs.
- Graph isomorphism.
- Isomorphic graphs, with examples.
- Graph invariants of isomorphic simple graphs.

---

---

---

---

---

---

---

References

1. [Ros19] Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.

2. [Lip07] Seymour Lipschutz and Marc Lars Lipson, *Schaum's Outline of Theory and Problems of Discrete Mathematics*, Third edition, McGraw-Hill Education, 2007.

3. [Wes01] Douglas Brent West, *Introduction to Graph Theory*, Second edition, Prentice-Hall, 2001.

4. [Deo74] Narsingh Deo, *Graph Theory with Applications to Engineering and Computer Science*, Prentice-Hall, 1974.

5. [Har69] Frank Harary, *Graph Theory*, Addison-Wesley, 1969.

---

---

---

---

---

---

---