

UNIT – II, Module – 1

Lecture 09: Discrete Structures

[ Cartesian product of finite sets; Union and intersection; Principle of inclusion-exclusion; Set operations; Set identities ]

Dr. Suddhasil De

---

---

---

---

---

---

---

Recap:: Discrete structures

- Discrete structures examples:
  - (i) sets (collections of objects) and multisets;
  - (ii) combinations (built from sets; unordered collections of objects to be used in counting);
  - (iii) relations (sets of ordered pairs representing relationships between objects);
  - (iv) graphs (sets of vertices and edges connecting vertices);
  - (v) sequences (ordered lists of elements, as well as special type of functions expressing relationships among elements);
  - (vi) **matrices**.

Discrete Mathematics

Dept. of CSE, NITP  
2

Dr. Suddhasil De

---

---

---

---

---

---

---

Notation table

Symbol / Notation	Meaning
$\cdot$	Arithmetic and algebraic multiplication in number theory.
$\times$	Multiply operation for Cartesian product in set theory.
$\cup, \cup_{i=1}^n$	Union operation of two, more (arbitrary no.) sets in set theory resp.
$\cap, \cap_{i=1}^n$	Intersection operation of two, more (arbitrary) sets in set theory resp.
$\vee, \vee_{i=1}^n$	Disjunction of two, more (arbitrary no.) assertions resp.
$\wedge, \wedge_{i=1}^n$	Conjunction of two, more (arbitrary no.) assertions resp.
$\setminus$	Set subtraction of two sets in set theory.
$\Delta$	Symmetric difference of two sets in set theory.
$\bar{A}$	Complement of set <b>A</b> in set theory.
$\mathcal{U}$	Universal set (or universe) in set theory.
$\emptyset$	Empty set in set theory.

Discrete Mathematics

Dept. of CSE, NITP  
3

Dr. Suddhasil De

---

---

---

---

---

---

---

## Set

- Set operations:
  - Product property:: **Cartesian product** of finite sets  $A_1, A_2, \dots, A_n$ :  
 $A_1 \times A_2 \times \dots \times A_n = \prod_{i=1}^n A_i = \{(a_1, a_2, \dots, a_n) \mid a_i \in A_i, i = 1, 2, \dots, n\}$ .  
 where,  $(a_1, a_2, \dots, a_n)$  = ordered  $n$ -tuple.
    - Property:: Ordered  $n$ -tuple: ordered collection of  $n$  elements.
    - Product equality property:: **Equality** of two ordered  $n$ -tuples  $(a_1, a_2, \dots, a_n)$ ,  $(b_1, b_2, \dots, b_n)$ :  $(a_1, a_2, \dots, a_n) = (b_1, b_2, \dots, b_n)$ , if and only if  $a_i = b_i$  for all  $i = 1, 2, \dots, n$ .
  - Property:: Cartesian product of finite sets  $A$  and  $B$ : set of all ordered pairs  $(a, b)$ , i.e.,  $A \times B = \{(a, b) \mid (a \in A) \wedge (b \in B)\}$ .
  - Property:: Ordered pair: ordered 2-tuple.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

4

## Set

- Set operations:
  - Property:: **Cartesian product** of finite set  $A^m$ :  
 $A^m = \{(a_1, a_2, \dots, a_m) \mid a_i \in A, i = 1, 2, \dots, m\}$ .
    - Property::  $|A_1 \times A_2 \times \dots \times A_n| = |A_1| \cdot |A_2| \cdot \dots \cdot |A_n| = \prod_{i=1}^n |A_i|$ .
    - Property::  $A \times B = B \times A$ , if  $A = \emptyset$  or  $B = \emptyset$  (so,  $A \times B = \emptyset$ ) or  $A = B$ .
  - Relation property:: **Relation** from set  $A$  to set  $B$ : subset of Cartesian product  $A \times B$ , more specifically called binary relation, denoted by  $\mathcal{R}$ .
  - Property:: Elements of relation  $\mathcal{R}$  from set  $A$  to set  $B$ : ordered pairs, where first element to belong to  $A$  and second to  $B$ .

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

5

## Set

- Set operations:
  - Union property:: **Union** of finite family of sets:  $A_1 \cup A_2 \cup \dots \cup A_n = \bigcup_{i=1}^n A_i = \{x \mid \exists i (1 \leq i \leq n) (x \in A_i)\}$ .
  - Generalization property:: For indexed set  $S$ , such that  $i \in S$ , union of finite family of sets:  $\bigcup_{i \in S} A_i = \{x \mid \exists i (i \in S) (x \in A_i)\}$ .
  - Property:: Union of two sets  $A$  and  $B$ :  $A \cup B = \{x \mid (x \in A) \vee (x \in B)\}$ .
  - Property:: Union of infinite family of sets:  $A_1 \cup A_2 \cup \dots \cup A_n \cup \dots = \bigcup_{i=1}^{\infty} A_i = \{x \mid \exists i (i \in \mathbb{N}) (x \in A_i)\}$ .

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

6

## Set

- Set operations:
  - Intersection property:: **Intersection** of finite family of sets:  
 $A_1 \cap A_2 \cap \dots \cap A_n = \bigcap_{i=1}^n A_i = \{x \mid \bigwedge_{i=1}^n (x \in A_i)\} = \{x \mid \forall i (x \in A_i), i = 1, 2, \dots, n\}.$
  - Generalization property:: For indexed set  $S$ , such that  $i \in S$ , intersection of finite family of sets:  $\bigcap_{i \in S} A_i = \{x \mid \forall i \in S (x \in A_i)\}.$
  - Intersection property:: Intersection of two sets  $A$  and  $B$ :  $A \cap B = \{x \mid (x \in A) \wedge (x \in B)\}.$
  - Property:: Intersection of infinite family of sets:  
 $A_1 \cap A_2 \cap \dots \cap A_n \cap \dots = \bigcap_{i=1}^{\infty} A_i = \{x \mid \bigwedge_{i=1}^{\infty} (x \in A_i)\}.$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

7

## Set

- Set operations:
  - Property:: **Principle of inclusion-exclusion**: Generalization of following two-set relation —  $|A \cup B| = |A| + |B| - |A \cap B|.$
  - Disjoint property:: If  $A \cap B = \emptyset$ , then  $A$  and  $B$  to become **disjoint**.
  - Set subtraction property:: **Set subtraction** of two finite sets  $A$  and  $B$  = Difference between  $A$  and  $B$  = complement of  $B$  with respect to  $A$ :  
 $A \setminus B = A - B = \{x \mid (x \in A) \wedge (x \notin B)\}.$
  - Successor property:: Successor of  $A$ :  $\text{succ}(A) = A \cup \{A\}.$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

8

## Set

- Set operations:
  - Symmetric difference property:: **Symmetric difference** of  $A$  and  $B$  (also disjunctive union):  $A \Delta B = A \ominus B = A \oplus B = \{x \mid (x \in A) \oplus (x \in B)\}.$
  - Property::  $A \Delta B = (A \setminus B) \cup (B \setminus A)$   
 $= (A \cup B) \setminus (A \cap B).$  Exclusive disjunction
  - Complement property:: **Complement** of  $A$  (denoted by  $\bar{A}$ , also by  $A'$ , or  $A^c$ ):  $\bar{A} = \{x \in U \mid x \notin A\} = U \setminus A.$
  - Property:: Complement of  $A$  with respect to superset  $B$  (such that  $A \subseteq B$ ):  $\{x \in B \mid x \notin A\}.$
  - Property::  $A \setminus B = A \cap \bar{B}.$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

9

## Set

- Set operations: examples.
  - Example-1:: for sets  $A = \{1, 2\}$ ,  $B = \{a, b, c\}$ , their Cartesian products  $A \times B$  and  $B \times A$ :
 
$$A \times B = \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c)\}.$$

$$B \times A = \{(a, 1), (a, 2), (b, 1), (b, 2), (c, 1), (c, 2)\}.$$
 Also,  $A^2 = A \times A = \{(1, 1), (1, 2), (2, 1), (2, 2)\}$ ,  
 $A^3 = \{(1, 1, 1), (1, 1, 2), (1, 2, 1), (1, 2, 2), (2, 1, 1), (2, 1, 2), (2, 2, 1), (2, 2, 2)\}$ ,  
 $B^2 = B \times B = \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c), (c, a), (c, b), (c, c)\}.$ 
 Further,  $|A \times B| = |B \times A| = |A| \cdot |B| = 6$ ;  
 $|A^2| = |A| \cdot |A| = 4$ ;  $|A^3| = |A| \cdot |A| \cdot |A| = 8$ ;  $|B^2| = |B| \cdot |B| = 9$ .

Discrete Mathematics

Dept. of CSE, NITP  
10

Dr. Suddhasil De

## Set

- Set operations: examples.
  - Example-2:: for sets  $A = \{1, 3, 5\}$ , and  $B = \{1, 2, 3\}$ ,  
 $A \cup B = B \cup A = \{1, 2, 3, 5\}$ ;  $A \cap B = B \cap A = \{1, 3\}$ ;  
 $A \setminus B = \{5\}$ ;  $B \setminus A = \{2\}$ ;  $A \Delta B = B \Delta A = \{2, 5\}.$
  - Example-3:: for  $A = \{1, 3, 5, 7, 9\}$ , and  $B = \{2, 4, 6, 8, 10\}$ ,  $A \cap B = \emptyset$ ,  
 so,  $A$  and  $B$  to be disjoint.  
 $A \cup B = B \cup A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ ;  
 $A \setminus B = \{1, 3, 5, 7, 9\}$ ;  $B \setminus A = \{2, 4, 6, 8, 10\}$ ;  
 $A \Delta B = B \Delta A = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}.$

Discrete Mathematics

Dept. of CSE, NITP  
11

Dr. Suddhasil De

## Set

- Set operations: examples.
  - Example-4:: for set  $A = \{a, e, i, o, u\}$ , and universal set  $\mathcal{U} = \{\text{set of letters of English alphabet}\}$ , so  
 $\bar{A} = A^c = \{b, c, d, f, g, h, j, k, l, m, n, p, q, r, s, t, v, w, x, y, z\}.$
  - Example-5:: for set  $A_i = \{i, i + 1, i + 2, i + 3, \dots\}$ , where  $i = 1, 2, \dots, n$ ,  
 $\bigcup_{i=1}^n A_i = \bigcup_{i=1}^n \{i, i + 1, i + 2, i + 3, \dots\} = \{1, 2, 3, \dots\}$ , and  
 $\bigcap_{i=1}^n A_i = \bigcap_{i=1}^n \{i, i + 1, i + 2, i + 3, \dots\} = \{n, n + 1, n + 2, \dots\} = A_n.$
  - Example-6:: for set  $B_i = \{1, 2, 3, \dots, i\}$ , where  $i = 1, 2, 3, \dots, \infty$ ,  
 $\bigcup_{i=1}^{\infty} B_i = \bigcup_{i=1}^{\infty} \{1, 2, 3, \dots, i\} = \{1, 2, 3, \dots\} = \mathbb{Z}^+$ , and  
 $\bigcap_{i=1}^{\infty} B_i = \bigcap_{i=1}^{\infty} \{1, 2, 3, \dots, i\} = \{1\} = B_1.$

Discrete Mathematics

Dept. of CSE, NITP  
12

Dr. Suddhasil De

Set

- Computer representation of set: techniques for storing elements of set, so as to carry out set operations efficiently.
- Bit string representation: storing elements of given set A, based on arbitrary ordering of elements of  $\mathcal{U}$ , assuming  $\mathcal{U}$  to be finite (and of reasonable size not larger than memory capacity of computer), i.e.,  $b_1b_2\cdots b_n$ , where  $|A| = n$  and  $b_i = \begin{cases} 1, & \text{if } a_i \in A \\ 0, & \text{otherwise} \end{cases}$  for  $i = 1, 2, \dots, n$ .  
 $n$  bits for A
- Benefit: easy to find complements of sets and unions, intersections and differences of sets.

---

---

---

---

---

---

---

---

Set

- Set operation identity: logically equal resultant sets.
- Set identities  $\rightarrow$  **Laws of algebra of set theory.**
- Proof of each law based on 3 proof methods —

Methods of Proving Set Identities	
Description	Method
Subset method	Show that each side of the identity is a subset of the other side.
Membership table	For each possible combination of the atomic sets, show that an element in exactly these atomic sets must either belong to both sides or belong to neither side (for any arbitrary element $x \in \mathcal{U}$ )
Apply existing identities	Start with one side, transform it into the other side using a sequence of steps by applying an established identity.

[Ref: Kenneth H. Rosen, Discrete Mathematics and Its Applications, Eighth edition, McGraw-Hill Education, 2019.]

---

---

---

---

---

---

---

---

Laws of algebra of set theory

Idempotent laws:	(III.1a) $A \cup A = A$	(III.1b) $A \cap A = A$
Identity laws:	(III.2a) $A \cup \emptyset = A$	(III.2b) $A \cap \mathcal{U} = A$
Domination laws:	(III.3a) $A \cup \mathcal{U} = \mathcal{U}$	(III.3b) $A \cap \emptyset = \emptyset$
Associative laws:	(III.4a) $(A \cup B) \cup C = A \cup (B \cup C)$	(III.4b) $(A \cap B) \cap C = A \cap (B \cap C)$
Commutative laws:	(III.5a) $A \cup B = B \cup A$	(III.5b) $A \cap B = B \cap A$
Distributive laws:	(III.6a) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$	(III.6b) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
Involution law:	(III.7) $\overline{\overline{A}} = A, (A^c)^c = A$	(also complementation law)
De Morgan's laws:	(III.8a) $\overline{(A \cup B)} = \overline{A} \cap \overline{B}, (A \cup B)^c = (A^c) \cap (B^c)$	(III.8b) $\overline{(A \cap B)} = \overline{A} \cup \overline{B}, (A \cap B)^c = (A^c) \cup (B^c)$

---

---

---

---

---

---

---

---

Laws of algebra of set theory

Complement laws:	(III.9a) $A \cup \bar{A} = U$ , $A \cup A^c = U$	(III.9c) $A \cap \bar{A} = \emptyset$ , $A \cap A^c = \emptyset$
	(III.9b) $\bar{U} = \emptyset$ , $U^c = \emptyset$	(III.9d) $\bar{\emptyset} = U$ , $\emptyset^c = U$
Absorption laws:	(III.10a) $A \cup (A \cap B) = A$	(III.10b) $A \cap (A \cup B) = A$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

16

---

---

---

---

---

---

---

---

Set

- Set operation identity: examples.
- Example-1:: Proof for law-(III.8b):  $\overline{(A \cap B)} = \bar{A} \cup \bar{B}$ .

Proof: using **subset method** [one set shown to be subset of other].

(1) Proving  $\overline{(A \cap B)} \subseteq \bar{A} \cup \bar{B}$ , i.e., if  $x \in \overline{(A \cap B)}$ , then  $x \in (\bar{A} \cup \bar{B})$ .  
Let  $x \in \overline{(A \cap B)}$ . By definition of complement,  $x \notin (A \cap B)$ , i.e.,  $\neg((x \in A) \wedge (x \in B))$  to be TRUE as per definition of intersection.  
By De Morgan's law for propositions in (I.10b),  $\neg(x \in A) \vee \neg(x \in B)$ .  
Using definition of negation of propositions,  $(x \notin A)$  or  $(x \notin B)$ , i.e.,  $(x \in \bar{A})$  or  $(x \in \bar{B})$  as per definition of complement of set.  
So, by definition of union,  $x \in (\bar{A} \cup \bar{B})$ . (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

17

---

---

---

---

---

---

---

---

Set

- Set operation identity: examples.
- Example-1 proof (contd.):

(2) Proving  $\bar{A} \cup \bar{B} \subseteq \overline{(A \cap B)}$ , i.e., if  $x \in (\bar{A} \cup \bar{B})$ , then  $x \in \overline{(A \cap B)}$ .  
Let  $x \in (\bar{A} \cup \bar{B})$ . By definition of union,  $(x \in \bar{A})$  or  $(x \in \bar{B})$ .  
As per definition of complement of set,  $(x \notin A)$  or  $(x \notin B)$ . So, proposition  $\neg(x \in A) \vee \neg(x \in B)$  to become TRUE.  
By De Morgan's law for propositions in (I.10b),  $\neg((x \in A) \wedge (x \in B))$  to be TRUE. Then, as per definition of intersection,  $\neg(x \in (A \cap B))$ .  
So, by definition of complement,  $x \in \overline{(A \cap B)}$ .  
Combining (1) and (2),  $\overline{(A \cap B)} = \bar{A} \cup \bar{B}$ . ■

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

18

---

---

---

---

---

---

---

---

## Set

- Set operation identity: examples.
- Example-2:: Same as Example-1, i.e., law-(III.8b):  $\overline{(A \cap B)} = \bar{A} \cup \bar{B}$ .

Proof: using compact form of subset method.

$$\begin{aligned}
 \overline{(A \cap B)} &= \{x \mid x \notin (A \cap B)\} && \text{by set-builder of set complement} \\
 &= \{x \mid \neg(x \in (A \cap B))\} && \text{by definition of set "not belong to"} \\
 &= \{x \mid \neg((x \in A) \wedge (x \in B))\} && \text{by definition of set intersection} \\
 &= \{x \mid \neg(x \in A) \vee \neg(x \in B)\} && \text{by De Morgan's law in (I.10b)} \\
 &= \{x \mid (x \notin A) \vee (x \notin B)\} && \text{by definition of set "not belong to"} \\
 &= \{x \mid (x \in \bar{A}) \vee (x \in \bar{B})\} && \text{by definition of set complement} \\
 &= \{x \mid (x \in (\bar{A} \cup \bar{B}))\} = \bar{A} \cup \bar{B} && \text{by definition of set union.} \blacksquare
 \end{aligned}$$

Discrete Mathematics

Dept. of CSE, NITP  
19

Dr. Suddhasil De

## Set

- Set operation identity: examples.
- Example-3:: Proof for law-(III.6b):  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

Proof: using compact subset method.

$$\begin{aligned}
 A \cap (B \cup C) &= \{x \mid x \in (A \cap (B \cup C))\} && \text{by set-builder of set intersection} \\
 &= \{x \mid (x \in A) \wedge (x \in (B \cup C))\} && \text{by definition of set intersection} \\
 &= \{x \mid (x \in A) \wedge ((x \in B) \vee (x \in C))\} && \text{by definition of set union} \\
 &= \{x \mid ((x \in A) \wedge (x \in B)) \vee ((x \in A) \wedge (x \in C))\} && \text{by (I.6b)} \\
 &= \{x \mid (x \in (A \cap B)) \vee (x \in (A \cap C))\} && \text{by definition of intersection} \\
 &= \{x \mid x \in ((A \cap B) \cup (A \cap C))\} && \text{by definition of set union} \\
 &= (A \cap B) \cup (A \cap C) && \text{by set-builder of set union.} \blacksquare
 \end{aligned}$$

Discrete Mathematics

Dept. of CSE, NITP  
20

Dr. Suddhasil De

## Set

- Set operation identity: examples.
- Example-4:: Proof for law-(III.6b):  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

Proof: using membership table method.

Adjacent table:	A	B	C	BUC	A∩(BUC)	A∩B	A∩C	(A∩B)∪(A∩C)
membership	T	T	T	T	T	T	T	T
table for given	T	T	F	T	T	T	F	T
combinations of	T	F	T	T	T	F	T	T
sets.	F	T	T	T	F	F	F	F
Same 5 <sup>th</sup> and 8 <sup>th</sup>	F	T	F	T	F	F	F	F
columns.	F	F	T	T	F	F	F	F
	F	F	F	F	F	F	F	F

Discrete Mathematics

Dept. of CSE, NITP  
21

Dr. Suddhasil De

Set

- Set operation identity: examples.
  - Example-5:: Proof for:  $\overline{(A \cup (B \cap C))} = (\overline{C} \cup \overline{B}) \cap \overline{A}$ .
- Proof:** apply **existing set identities** to prove new identities.
- |  |                                   |
|--|-----------------------------------|
| $\overline{(A \cup (B \cap C))} = \overline{A} \cap \overline{(B \cap C)}$ | by De Morgan's law in (III.8a)    |
| $= \overline{A} \cap (\overline{B} \cup \overline{C})$                     | by De Morgan's law in (III.8b)    |
| $= (\overline{B} \cup \overline{C}) \cap \overline{A}$                     | by commutative law in (III.5b)    |
| $= (\overline{C} \cup \overline{B}) \cap \overline{A}$                     | by commutative law in (III.5a). ■ |

---

---

---

---

---

---

---

Summary

- Focus: Set operations.
- Cartesian product of finite sets, with examples.
- Union and intersection of finite sets, and related theorems with examples.
- Principle of inclusion-exclusion in set theory.
- Set operations, with examples.
- Set identities, i.e., laws of algebra of set theory.
- Set operation identity examples.

---

---

---

---

---

---

---

References

1. [Ros21] Kenneth H. Rosen, Kamala Krithivasan, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2021.
2. [Ross12] Kenneth A. Ross, Charles R. B. Wright, *Discrete Mathematics*, Fifth edition, Pearson Education, 2012.
3. [Mot15] Joe L. Mott, Abraham Kandel, Theodore P. Baker, *Discrete Mathematics for Computer Scientists and Mathematicians*, Second edition, Pearson Education, 2015.
4. [Lip07] Seymour Lipschutz, Marc L. Lipson, *Schaum's Outline of Theory and Problems of Discrete Mathematics*, Third edition, McGraw-Hill Education, 2007.

---

---

---

---

---

---

---



Further Reading

- Cartesian product of sets:: [Ros21]:128-130.
- Union of sets:: [Ros21]:133-134,139-141.
- Intersection of sets:: [Ros21]:134,139-141.
- Principle of inclusion-exclusion in set theory:: [Ros21]:134.
- Set operations:: [Ros21]:135-136.
- Set identities:: [Ros21]:136-139.

Discrete Mathematics

Dept. of CSE, NITP  
25

Dr. Suddhasil De

---

---

---

---

---

---

---

Lecture Exercises: Problem 1 [Ref: Gate 2015, Q.2, p.1 (Set-3)]

Suppose  $U$  is the power set of the set  $S = \{1,2,3,4,5,6\}$ . For any  $T \in U$ , let  $|T|$  denote the number of elements in  $T$  and  $T'$  denote the complement of  $T$ . For any  $T, R \in U$ , let  $T \setminus R$  be the set of all elements in  $T$  which are not in  $R$ . Which one of the following is TRUE?

- (a)  $\forall X \in U (|X| = |X'|)$ .
- (b)  $\exists X \in U \exists Y \in U (|X| = 5, |Y| = 5 \text{ and } X \cap Y = \emptyset)$ .
- (c)  $\forall X \in U \forall Y \in U (|X| = 2, |Y| = 3 \text{ and } X \setminus Y = \emptyset)$ .
- (d)  $\forall X \in U \forall Y \in U (X \setminus Y = Y' \setminus X')$ .

Discrete Mathematics

Dept. of CSE, NITP  
26

Dr. Suddhasil De

---

---

---

---

---

---

---

Lecture Exercises: Problem 1 Ans

(D) Let arbitrary variables  $X, Y \in U$ .

Then, L.H.S.:  $X \setminus Y = X \cap Y'$

by set subtraction property

Also, R.H.S.:  $Y' \setminus X' = Y' \cap (X')'$

by set subtraction property

$= Y' \cap X$

by involution law (III.7)

$= X \cap Y'$

by commutative law (III.5b)

So,  $(X \setminus Y = Y' \setminus X')$ . ①

After applying universal generalization (II.8) on ① for both variables successively,  $\forall X \in U \forall Y \in U (X \setminus Y = Y' \setminus X') \equiv \forall Y \in U \forall X \in U (X \setminus Y = Y' \setminus X')$  (as per nested universal quantification ordering property).

- What about options (A), (B), (C)?

Discrete Mathematics

Dept. of CSE, NITP  
27

Dr. Suddhasil De

---

---

---

---

---

---

---