

UNIT – IV, Module – 1

Lecture 35: Graph Planarity

[Planar, nonplanar graphs; Region, degree of region; Elementary subdivision; Homeomorphism; Kuratowski's two graphs; Euler's Formula]


Dr. Suddhasil De

Notation table

Symbol / Notation	Meaning
R	Region in planar representation of graph.
$\deg(R)$	Degree of region R in planar representation of graph.
K_5	Kuratowski's first graph.
$K_{3,3}$	Kuratowski's second graph.

Graph planarity

- Motivation for graph planarity: to find answers to some real-life problems. E.g., buildings-and-utilities problem.
 - Property:: Possibility to join these buildings and utilities without connections crossing each other?
- Rephrased problem: Capturing given scenario as graph, resulting into complete bipartite graph $K_{3,3}$, possibility to draw $K_{3,3}$ in a plane without crossing any of its edges?




[Ref: Kenneth H. Rosen, Discrete Mathematics and its Applications, Eighth edition, McGraw-Hill Education, 2019.]

Graph planarity

- Planar graph: graph or multigraph, where its **drawing** (i.e., **geometric representation**) in single plane **possible** without crossing any of its edges.

- Property:: Possibility of **planar graph** to be **embedded** (i.e., drawn) in plane surface.

- Property:: Crossing of edges: intersection of lines or arcs representing edges at some point other than their common vertices.



[Ref: Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory


Dept. of CSE, NITP

Dr. Suddhasil De

4

Graph planarity

- Planar graph:
 - Property:: **Planar representation** of graph: embedding of graph in single plane without crossing any of its edges.



- Note:** planar graph usually drawn with crossings; still planar, due to possibility to draw it in different way without crossings.

[Ref: Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory


Dept. of CSE, NITP

Dr. Suddhasil De

5

Graph planarity

- Planar graph:
 - Property:: **Regions** in planar representation of graph: subareas of plane generated while drawing planar representation of graph (i.e., embedding graph in plane); also called **faces**.
 - Property:: Both bounded and unbounded regions possible. In fig. $R_6 \rightarrow$ unbounded.
 - Property:: **Unbounded region**: also called **exterior face**.
 - Property:: **Nonplanar graph**: graph not possible to be drawn on plane without crossover between its edges.



[Ref: Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory


Dept. of CSE, NITP

Dr. Suddhasil De

6

Graph planarity

- Planar graph:
 - Property: **Degree of region R** in planar representation of graph: denoted by $\deg(R)$, where $\deg(R)$ = number of edges on boundary of R , such that an edge to contribute two to $\deg(R)$ when that edge to be traced out twice while tracing boundary of R .



[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

7

Graph planarity

- Planar graph:
 - Property: **Elementary subdivision** of planar graph $G = (V, E)$: graph $G' = (V', E')$ obtained after performing following operations in sequence — (i) removing existing edge $\{u, v\}$ such that $E \setminus \{u, v\}$, i.e., $G - \{u, v\}$; (ii) adding new vertex w , such that $V' = V \cup \{w\}$; and (iii) adding new edges such that $E' = E \setminus \{u, v\} \cup \{\{u, w\}, \{w, v\}\}$.
 - Property: **Homeomorphism**: Homeomorphic graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ of planar graph $G = (V, E)$, if G_1 and G_2 obtained from G by a sequence of elementary subdivisions.

Graph Theory


Dept. of CSE, NITP

Dr. Suddhasil De

8

Graph planarity

- Planar graph:
 - Example-1: Homeomorphic graphs G , G_1 and G_2 .



$G \xrightarrow{\text{elementary subdivision}} G$: empty sequence of elementary subdivisions.
 $G \xrightarrow{\text{elementary subdivisions}} G_1$: (1) removing $\{a, c\}$, adding f , adding $\{a, f\}$, $\{f, c\}$; (2) removing $\{b, c\}$, adding g , adding $\{b, g\}$, $\{g, c\}$; (3) removing $\{b, g\}$, adding h , adding $\{b, h\}$, $\{h, g\}$.

(contd. to next slide)

[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Graph Theory

Dept. of CSE, NITP

Dr. Suddhasil De

9

Graph planarity

- Planar graph:
- Example-1 contd.:
 $G \xrightarrow{\text{elementary subdivisions}} G_2$: (1) removing $\{b, c\}$, adding g , adding $\{b, g\}$, $\{g, c\}$; (2) removing $\{b, g\}$, adding i , adding $\{b, i\}$, $\{i, g\}$; (3) removing $\{b, e\}$, adding j , adding $\{b, j\}$, $\{j, e\}$; (4) removing $\{b, j\}$, adding k , adding $\{b, k\}$, $\{k, j\}$.

Graph Theory

Dept. of CSE, NITP
10

Dr. Suddhasil De

Graph planarity

- Planar graph:
- Benefits: (i) planar graphs for modeling of electronic circuits to print on single board with no wires crossing;
 (ii) planar subgraphs (after vertex partition) for multi-layered circuit print, with insulated wires at crossings, and then objective to draw with fewest possible crossings;
 (iii) planar graphs in design of road networks to connect a group of cities by highways without using underpasses or overpasses.


Graph Theory

Dept. of CSE, NITP
11

Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:
 - Property: (Theorem: Planarity of K_5): K_5 nonplanar.
 [K_5 : called Kuratowski's first graph]
- Proof:** [Based on Jordan curve theorem: "every simple curve divides plane into two regions;" Special case: every simple polygon divides plane into two regions — interior (consisting of points inside curve), and exterior (consisting of points outside curve)]
- Idea:** drawing $K_5 = (V, E)$, where $V = \{v_1, v_2, v_3, v_4, v_5\}$, in sequence, starting from simple pentagon, and successively adding edge(s) (incident on existing vertex) at each stage.



(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
12

Dr. Suddhasil De

Graph planarity


- Nonplanarity of graph:

Proof of (Theorem: Planarity of K_5) contd.:

In figure-(1), simple pentagon to connect all vertices of K_5 by one edge, forming one circuit.

As per **Jordan curve theorem**, number of regions = 2, interior region $R^{(in)}$ and exterior region $R^{(ex)}$.

Adding edge $\{v_1, v_3\}$ to figure-(1) resulting in figure-(2), whose drawing possible through inside or outside pentagon, without intersecting edges of figure-(1). Let $\{v_1, v_3\}$ be drawn through inside.



(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
13

Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:


Proof of (Theorem: Planarity of K_5) contd-2.:

In case when $\{v_1, v_3\}$ be drawn through outside, then also similar arguments possible to prove theorem.

$\{v_1, v_3\}$ splitting $R^{(in)}$ into $R_1^{(in)}$ and $R_2^{(in)}$ regions.

In figure-(2), adding edges $\{v_2, v_4\}$ and $\{v_2, v_5\}$ resulting in figure-(3), whose drawing through inside of pentagon of figure-(2) not possible without intersecting edges.

So, these two edges drawn through outside of pentagon of figure-(2), splitting $R^{(ex)}$ into $R_1^{(ex)}, R_3^{(in)}, R_4^{(in)}$ (contd. to next slide)



Graph Theory

Dept. of CSE, NITP
14

Dr. Suddhasil De

Graph planarity


- Nonplanarity of graph:

Proof of (Theorem: Planarity of K_5) contd-3.:

In figure-(3), adding edge $\{v_3, v_5\}$ resulting in figure-(4), whose drawing through outside of pentagon of figure-(3) not possible without intersecting edge $\{v_2, v_4\}$.

So, this edge drawn through inside of pentagon of figure-(3), splitting $R_2^{(in)}$ into $R_{21}^{(in)}, R_{22}^{(in)}$ regions.

At this point, adding edge $\{v_1, v_4\}$ in figure-(4) resulting in figure-(5), whose drawing through inside of pentagon of figure-(4) not possible without intersecting edge $\{v_1, v_4\}$.



(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
15

Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:


Proof of (Theorem: Planarity of K_5) contd-4.:

But drawing of $\{v_1, v_4\}$ through outside of pentagon of figure-(4) also not possible without intersecting edge $\{v_2, v_5\}$.

Thus, at least one edge of K_5 found, with no possibility to be placed inside or outside pentagon without crossover.

Hence, K_5 not possible to be embedded in plane. ■

- Property: K_5 (or, Kuratowski's first graph) = nonplanar graph with smallest number of vertices (only 5 vertices).



Graph Theory

Dept. of CSE, NITP
16


Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:

- Property: (Theorem: Planarity of $K_{3,3}$): $K_{3,3}$ nonplanar. [$K_{3,3}$ also called Kuratowski's second graph]

Proof: In any planar representation of $K_{3,3}$, vertices v_1 and v_2 must be connected to both v_4 and v_5 through four edges $\{v_1, v_4\}$, $\{v_1, v_5\}$, $\{v_2, v_4\}$, $\{v_2, v_5\}$, shown in figure-(1). These four edges form 'closed curve' square, and as per Jordan curve theorem, number of regions = 2, exterior region R_1 and interior region R_2 . (contd. to next slide)



Graph Theory

Dept. of CSE, NITP
17

Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:


Proof of (Theorem: Planarity of $K_{3,3}$) contd.:

Adding vertex v_3 to figure-(1) resulting in figure-(2), whose drawing possible through inside or outside of square in either R_1 or R_2 .

Let v_3 be drawn through R_2 .

In case if v_3 drawn through R_1 , then also similar arguments possible to prove theorem.

In figure-(2), v_3 must be connected to both v_4 and v_5 through edges $\{v_3, v_4\}$, $\{v_3, v_5\}$, splitting R_2 into R_{21} and R_{22} regions. (contd. to next slide)



Graph Theory

Dept. of CSE, NITP
18

Dr. Suddhasil De

Graph planarity

- Nonplanarity of graph:
Proof of (Theorem: Planarity of $K_{3,3}$) contd-2.:
At this point, adding vertex v_6 in R_{21} region of figure-(2) not possible without crossing, due to drawing of edge $\{v_2, v_6\}$ from v_6 not possible without intersecting any of existing edges $\{v_1, v_4\}, \{v_1, v_5\}, \{v_3, v_4\}, \{v_3, v_5\}$.
But, adding v_6 in R_{22} region of figure-(2) also not possible without crossing, due to drawing of edge $\{v_1, v_6\}$ without intersecting any of existing edges $\{v_2, v_4\}, \{v_2, v_5\}, \{v_3, v_4\}, \{v_3, v_5\}$ not possible.

(contd. to next slide)

Graph planarity

- Nonplanarity of graph:
Proof of (Theorem: Planarity of $K_{3,3}$) contd-3.:
Further, drawing of v_6 in R_1 region of figure-(2) also not possible without crossing, due to drawing of edge $\{v_3, v_6\}$ not possible without intersecting any of existing edges $\{v_1, v_4\}, \{v_1, v_5\}, \{v_2, v_4\}, \{v_2, v_5\}$.
Thus, at least one vertex of $K_{3,3}$ found, with no possibility to be placed inside or outside square of figure-(2) without crossover.
Hence, $K_{3,3}$ not possible to be embedded in plane. ■
• **Note::** Answer to buildings-and-utilities problem: not possible to connect three houses and three utilities in plane without a crossing.

Graph planarity

- Nonplanarity of graph:
 - Property:: $K_{3,3}$ (or, Kuratowski's second graph) = nonplanar graph with smallest number of edges (only 9 edges).

Graph planarity

- Euler's formula:
 - Property: (**Theorem: Euler's Formula**): If $G = (V, E)$ be connected planar simple graph with $|E| = e$ edges and $|V| = v$ vertices, and if r be number of regions in planar representation of G , then $r = e - v + 2$.

Proof: [By principle of mathematical induction]

Idea: constructing sequence of planar subgraphs inductively as $G_1, G_2, \dots, G_e = G$, successively adding one edge (incident on existing vertex) at each stage; such construction possible, G being connected. For any arbitrary integer n ($1 \leq n \leq e$), r_n, e_n, v_n to represent count of regions, edges, vertices of planar representation of G_n respectively.

(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
22

Dr. Suddhasil De

Graph planarity

- Euler's formula:


Proof of (**Euler's Formula**) contd:

(*Basis step*) Arbitrarily picking one edge of G from E to obtain G_1 .

Then, for planar G_1 , $r_1 = e_1 - v_1 + 2$, because $e_1 = 1$ (single edge), $v_1 = 2$ (two vertices $u_{1,1}$ and $u_{1,2}$), and $r_1 = 1$ (single region $R_{1,1}$), as evident in figure-(1).

(*Inductive step*) Inductive hypothesis: premise that Euler's formula to become true for planar subgraph G_k ($1 < k < n$), i.e., $r_k = e_k - v_k + 2$.

Let arbitrary edge $\{u_{k+1,i}, u_{k+1,j}\}$ of G , picked from E , be added to G_k to construct G_{k+1} . Two possible cases to handle for G_{k+1} . (contd. to next slide)



Graph Theory

Dept. of CSE, NITP
23

Dr. Suddhasil De

Graph planarity


- Euler's formula:

Proof of (**Euler's Formula**) contd-2:

(A) Both vertex $u_{k+1,i}$ and vertex $u_{k+1,j}$ already included in G_k in figure-(2).

Then, $u_{k+1,i}$ and $u_{k+1,j}$ must be on boundary of common region $R_{k+1,l}$ of G_k , else impossible to add edge $\{u_{k+1,i}, u_{k+1,j}\}$ to G_k without crossing other edges of G_k (thereby satisfying G_{k+1} planar), as shown in figure-(2).

Then, addition of $\{u_{k+1,i}, u_{k+1,j}\}$ splitting $R_{k+1,l}$ into two regions.



(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
24

Dr. Suddhasil De

Graph planarity

- Euler's formula:

Proof of (Euler's Formula) contd-3:

Consequently, two regions $R_{k+1,l}^{(1)}$ and $R_{k+1,l}^{(2)}$ generated, shown in figure-(3).

Then, in \mathcal{G}_{k+1} , $r_{k+1} = r_k + 1$, $e_{k+1} = e_k + 1$, and

$$v_{k+1} = v_k.$$


So, $r_{k+1} = r_k + 1 = e_k - v_k + 2 + 1 = (e_k + 1) - v_{k+1} + 2 = e_{k+1} - v_{k+1} + 2$. Because, each side of formula ' $r_k = e_k - v_k + 2$ ' in \mathcal{G}_k to increase by exactly one, thereby formula ' $r_{k+1} = e_{k+1} - v_{k+1} + 2$ ' still true for \mathcal{G}_{k+1} .

(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
25

Dr. Suddhasil De



Graph planarity

- Euler's formula:

Proof of (Euler's Formula) contd-4:

(B) For edge $\{u_{k+1,i}, u_{k+1,j}\}$, any one of two vertices of that edge not present in \mathcal{G}_k .

Let vertex $u_{k+1,i}$ already included in \mathcal{G}_k , but vertex $u_{k+1,j}$ not in \mathcal{G}_k in figure-(4).


Adding $\{u_{k+1,i}, u_{k+1,j}\}$ to \mathcal{G}_k not producing any new region in \mathcal{G}_{k+1} , because $u_{k+1,j}$ must be within region $R'_{k+1,l}$ of \mathcal{G}_k with $u_{k+1,i}$ on its boundary, else impossible to add that edge to \mathcal{G}_k without crossing existing edges (satisfying \mathcal{G}_{k+1} planar) in figure-(4).

(contd. to next slide)

Graph Theory

Dept. of CSE, NITP
26

Dr. Suddhasil De



Graph planarity

- Euler's formula:

Proof of (Euler's Formula) contd-5:

Then, in \mathcal{G}_{k+1} , $r_{k+1} = r_k$, $e_{k+1} = e_k + 1$, and $v_{k+1} = v_k + 1$.

So, $r_{k+1} = r_k = e_k - v_k + 2 = (e_k + 1) - (v_k + 1) + 2 = e_{k+1} - v_{k+1} + 2$.

Because, each side of formula ' $r_k = e_k - v_k + 2$ ' in \mathcal{G}_k to remain same, and increase in e_{k+1} and v_{k+1} to cancel each other, thereby formula ' $r_{k+1} = e_{k+1} - v_{k+1} + 2$ ' still true for \mathcal{G}_{k+1} .

Hence, Euler's formula to be true for planar subgraph \mathcal{G}_n ($1 \leq n \leq e$).

As $\mathcal{G}_e = \mathcal{G}$, by Mathematical Induction, $r = e - v + 2$. ■

Graph Theory

Dept. of CSE, NITP
27

Dr. Suddhasil De

Summary

- Focus: Planarity of graph.
- Motivation for graph planarity.
- Planar graph, with properties, examples and benefits.
- Planar representation of graph.
- Regions in planar graph, with properties and examples.
- Nonplanar graph.
- Graph nonplanarity theorems on Kuratowski’s first graph and second graph, with proofs.
- Euler’s Formula on graph planarity, with proof.
- Degree of region in planar graph.

References

1. [Ros19] Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.
2. [Lip07] Seymour Lipschutz and Marc Lars Lipson, *Schaum’s Outline of Theory and Problems of Discrete Mathematics*, Third edition, McGraw-Hill Education, 2007.
3. [Wes01] Douglas Brent West, *Introduction to Graph Theory*, Second edition, Prentice-Hall, 2001.
4. [Deo74] Narsingh Deo, *Graph Theory with Applications to Engineering and Computer Science*, Prentice-Hall, 1974.
5. [Har69] Frank Harary, *Graph Theory*, Addison-Wesley, 1969.
