

CS34110

Discrete Mathematics and Graph Theory

UNIT – II, Module – 2

Lecture 16: Counting

[Binomial coefficient, binomial expression;
Binomial theorem; Binomial identities;
Pascal's identity theorem, Pascal's triangle]

Dr. Suddhasil De

Binomial coefficients

- Binomial coefficient: coefficient of any term after expansion of powers of binomial expressions (or simply, binomials).
- Property:: **Binomial**: polynomial expressed as sum of two monomials.
- Property:: **Monomial**: polynomial expressed by only one term with nonzero coefficient.
- Property:: **Coefficient**: generally, multiplicative factor involved in any of all terms of polynomial, or series, or expression.
- Property:: Types of coefficient (in general case): (i) number (dimensionless), also called **numerical factor**; (ii) constant (with units of measurement), also called **constant multiplier**.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

2

Binomial coefficients

- Binomial expression's expansion:
 - Property:: Binomial expression: of general form $(a + b)^n$.
 - Property:: Expansion of binomial: using combinatorial reasoning.
- Example-1:: $(a + b)^3 = (a + b) \cdot (a + b) \cdot (a + b)$ (1)
 $= (a \cdot a + a \cdot b + b \cdot a + b \cdot b) \cdot (a + b)$
 $= a \cdot a \cdot a + a \cdot a \cdot b + a \cdot b \cdot a + a \cdot b \cdot b + b \cdot a \cdot a + b \cdot a \cdot b + b \cdot b \cdot a + b \cdot b \cdot b$
To obtain any (blue) term, say $a \cdot a \cdot a$, a MUST be chosen from every multiplier $(a + b)$, i.e., each choice of which possible independent of previous choice and each choice possible in only 1 way, so as per **product rule**, no. of ways $= 1 \cdot 1 \cdot 1 = 1 =$ **coefficient value**. (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

3

Binomial coefficients

- Binomial expression's expansion:
 - Example-1 (contd.): $(a+b)^3 = (a+b) \cdot (a+b) \cdot (a+b)$ (1)
 $= a \cdot a \cdot a + a \cdot a \cdot b + a \cdot b \cdot a + a \cdot b \cdot b + b \cdot a \cdot a + b \cdot a \cdot b + b \cdot b \cdot a + b \cdot b \cdot b$.
 For $a^2 \cdot b$, three (red) terms in expansion, where each term to start either by a or by b only; their ways also independent. Reason:
 If a chosen 1st, then after choosing a , for next choice, another a or b to be independently chosen from remaining 2 multipliers in 2 ways, and then last choice also independent. So, by **product rule**, no. of addends = $1 \cdot 2 \cdot 1 = 2$.
 If b chosen 1st, by **product rule**, no. of addends = $1 \cdot 1 \cdot 1 = 1$. (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
4

Dr. Suddhasil De

Binomial coefficients

- Binomial expression's expansion:
 - Example-1 (contd-2.): $(a+b)^3 = (a+b) \cdot (a+b) \cdot (a+b)$ (1)
 $= a \cdot a \cdot a + a \cdot a \cdot b + a \cdot b \cdot a + a \cdot b \cdot b + b \cdot a \cdot a + b \cdot a \cdot b + b \cdot b \cdot a + b \cdot b \cdot b$.
 Then for $a^2 \cdot b$, as per **sum rule**, no. of addends = no. of ways = $2+1 = 3 = \text{coefficient value}$.
 Alternative explanation:: number of ways to select 1st a from three $(a+b)$ multipliers, leaving rest multipliers for choice of 2nd a and one b .
 Same explanation for $b^2 \cdot a$.

Discrete Mathematics

Dept. of CSE, NITP
5

Dr. Suddhasil De

Binomial coefficients

- Binomial expression's expansion:
 - Example-2: $(a+b)^4 = (a+b) \cdot (a+b) \cdot (a+b) \cdot (a+b)$ (2)
 $= (a \cdot a \cdot a \cdot a + b \cdot b \cdot b \cdot b) + (a \cdot a \cdot a \cdot b + a \cdot a \cdot b \cdot a + a \cdot b \cdot a \cdot a + b \cdot a \cdot a \cdot a) + (b \cdot b \cdot b \cdot a + b \cdot b \cdot a \cdot b + b \cdot a \cdot b \cdot b + a \cdot b \cdot b \cdot b) + (a_1 \cdot a_2 \cdot b_3 \cdot b_4 + a_1 \cdot a_2 \cdot b_4 \cdot b_3 + a_1 \cdot a_3 \cdot b_2 \cdot b_4 + a_1 \cdot a_3 \cdot b_4 \cdot b_2 + a_1 \cdot a_4 \cdot b_2 \cdot b_3 + a_1 \cdot a_4 \cdot b_3 \cdot b_2 + \dots)$.
 For brown terms of form $a^2 \cdot b^2$, to select 1st a from four $(a+b)$ multipliers in 4 ways; for each of these 4 ways, to select 2nd a from remaining $(a+b)$ multipliers in 3 ways independently.
 So, by **product rule**, no. of ways = $4 \cdot 3$, where 2 ways be equivalent.
 So, by **division rule**, distinct no. of summands = $\frac{4}{2} \cdot \frac{3}{1} = 6 = \text{coefficient}$.

Discrete Mathematics

Dept. of CSE, NITP
6

Dr. Suddhasil De

Binomial theorem

- Binomial theorem: expanding nonnegative power of generalized binomial expression as sum of terms involving binomial coefficients.

• Property:: (Binomial theorem): For variables (& constants) x and y ,

$$(x+y)^n = \binom{n}{0} \cdot x^n + \binom{n}{1} \cdot x^{n-1} \cdot y + \dots + \binom{n}{n-1} \cdot x \cdot y^{n-1} + \binom{n}{n} \cdot y^n$$

$$= \sum_{r=0}^n \binom{n}{r} \cdot x^{n-r} \cdot y^r = \sum_{r=0}^n \binom{n}{r} \cdot x^r \cdot y^{n-r}$$

where, $n \in \mathbb{N}$, and "n-choose-r" denoted by $\binom{n}{r}$ = number of ways of selecting r elements from set of n elements, $r \in \mathbb{N}$, $r \leq n$.

Discrete Mathematics

Dept. of CSE, NITP
7

Dr. Suddhasil De

Binomial theorem

- Binomial theorem:

Proof of (Binomial theorem): [Using counting principles].

Expanded product to contain summands of form: $x^{n-r} \cdot y^r$, for $r = 0, 1, 2, 3, \dots, n$. Each summand = one/more addends summed together. Number of addends resulting in summand of form $x^{n-r} \cdot y^r$ (for some r) = **distinct** ways to select x for $(n-r)$ times from n no. of $(x+y)$ multipliers, leaving rest r no. of $(x+y)$ for choice of r times of y .

\therefore Coefficient of $x^{n-r} \cdot y^r$ (as per **product rule & division rule**): $\frac{n!}{r! \cdot (n-r)!}$.
 $\dots, \frac{(n-(r-1))!}{1} = \frac{n \cdot (n-1) \cdot \dots \cdot (n-(r-1))}{r \cdot (r-1) \cdot \dots \cdot 1} = \frac{n!}{r! \cdot (n-r)!}$ denoted by $\binom{n}{r}$.

Also, same coefficient $\binom{n}{r}$ for $x^r \cdot y^{n-r}$. ■

Discrete Mathematics

Dept. of CSE, NITP
8

Dr. Suddhasil De

Binomial theorem

- Binomial theorem examples:

• Example-1:: Expansion of $(x+y)^4$. From binomial theorem,

$$(x+y)^4 = \sum_{j=0}^4 \binom{4}{j} \cdot x^{4-j} \cdot y^j = \sum_{j=0}^4 \binom{4}{j} \cdot x^j \cdot y^{4-j}$$

$$= \binom{4}{0} \cdot x^4 + \binom{4}{1} \cdot x^3 \cdot y + \binom{4}{2} \cdot x^2 \cdot y^2 + \binom{4}{3} \cdot x \cdot y^3 + \binom{4}{4} \cdot y^4$$

$$= 1 \cdot x^4 + 4 \cdot x^3 \cdot y + 6 \cdot x^2 \cdot y^2 + 4 \cdot x \cdot y^3 + 1 \cdot y^4.$$

• Example-2:: Coefficient of $x^{12} \cdot y^{13}$ in $(x+y)^{25}$. From binomial theorem, given term's coefficient (from 1st expression of summands)

$$= \binom{25}{13} = \frac{25!}{13! \cdot 12!} = 5200300.$$

From 2nd expression of summands in binomial theorem, given term's coefficient = $\binom{25}{12} = \frac{25!}{12! \cdot 13!} = 5200300.$

Discrete Mathematics

Dept. of CSE, NITP
9

Dr. Suddhasil De

Binomial theorem

- Binomial theorem examples:
 - Example-3: Coefficient of $x^{12} \cdot y^{13}$ in expansion of $(2 \cdot x - 3 \cdot y)^{25}$.
 Let $u = 2 \cdot x$, $v = -3 \cdot y$; so given binomial to become $(u + v)^{25}$, and required term to become $2^{12} \cdot u^{12} \cdot (-1)^{13} \cdot 3^{13} \cdot v^{13}$.
 From 1st expression of summands in binomial theorem, given term's coefficient = $\binom{25}{13} \cdot 2^{12} \cdot (-1)^{13} \cdot 3^{13} = -\frac{25!}{13! \cdot 12!} \cdot 2^{12} \cdot 3^{13}$.
 From 2nd expression of summands in binomial theorem, given term's coefficient = $\binom{25}{12} \cdot 2^{12} \cdot (-1)^{13} \cdot 3^{13} = -\frac{25!}{13! \cdot 12!} \cdot 2^{12} \cdot 3^{13}$.

Discrete Mathematics

Dept. of CSE, NITP
10

Dr. Suddhasil De

Binomial identities

- Binomial identity: formula expressing products of binomial factors as sum over terms, each including a binomial coefficient.
 - Property: (Corollary): $\sum_{r=0}^n \binom{n}{r} = 2^n$, where $n \in \mathbb{N}$.
 Proof of above corollary to obtain straight from binomial theorem on assigning $x = y = 1$, and simplifying.
 - Property: (Corollary): $\sum_{r=0}^n (-1)^r \cdot \binom{n}{r} = 0$, where $n \in \mathbb{Z}^+$.
 - Property: (Corollary): $\binom{n}{0} + \binom{n}{2} + \binom{n}{4} + \dots + \binom{n}{1} + \binom{n}{3} + \binom{n}{5} + \dots$, where $n \in \mathbb{Z}^+$.
 - Property: (Corollary): $\sum_{r=0}^n \left(2^r \cdot \binom{n}{r} \right) = 3^n$, where $n \in \mathbb{N}$.

Discrete Mathematics

Dept. of CSE, NITP
11

Dr. Suddhasil De

Binomial identities

- Binomial identity:
 - Property: (Pascal's identity theorem): $\binom{n+1}{r} = \binom{n}{r-1} + \binom{n}{r}$, where $n, r \in \mathbb{Z}^+$, $r \leq n$.
Proof: Using counting principles. Let T be any set, where $|T| = n + 1$.
 Let any arbitrary element $a \in T$, and let $S = T \setminus \{a\}$, where $|S| = n$.
 Then, number of subsets of T containing r elements = $\binom{n+1}{r}$, $r \leq n$.
 Such subset formation of T containing r elements possible in two procedures — either (i) each subset to contain a , together with $r - 1$ elements taken from n elements of S , or (ii) each subset to contain r elements taken from n elements of S , and not containing a . (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
12

Dr. Suddhasil De

Binomial identities

- Binomial identity:
 - Proof of (Pascal's identity theorem) contd.
In first case, number of ways to form subsets of $r - 1$ elements from $S = \binom{n}{r-1}$. \therefore Number of subsets of T (of r elements) with $a = \binom{n}{r-1}$.
In second case, number of ways to form subsets of r elements from $S = \binom{n}{r}$. \therefore Number of subsets of T (of r elements) without $a = \binom{n}{r}$.
Then, according to **sum rule**,
$$\binom{n+1}{r} = \binom{n}{r-1} + \binom{n}{r}.$$
 ■

Binomial identities

- Binomial identity:
 - (Pascal's identity theorem) **Alternate proof**: based on algebraic manipulations. To derive L.H.S., starting with R.H.S. of given equation. Use of expression of binomial coefficients.
$$\binom{n}{r-1} + \binom{n}{r} = \frac{n!}{(r-1)! \cdot (n-r+1)!} + \frac{n!}{r! \cdot (n-r)!} = \frac{n! \cdot r}{r! \cdot (n-r+1)!} + \frac{n! \cdot (n-r+1)}{r! \cdot (n-r+1)!} = \frac{n! \cdot (r+n-r+1)}{r! \cdot (n-r+1)!} = \frac{(n+1)!}{r! \cdot (n+1-r)!} = \binom{n+1}{r}.$$
 ■


Binomial identities

- Binomial identity:
 - Property:: (Theorem): $\binom{n}{r} = \binom{n}{n-r}$, where $n \in \mathbb{Z}^+$, $r \in \mathbb{N}$, $r \leq n$.
Proof: Based on algebraic manipulations. To derive L.H.S., starting with R.H.S. of given equation. Use of expression of binomial coefficients.
$$\binom{n}{n-r} = \frac{n!}{(n-r)! \cdot (n-(n-r))!} = \frac{n!}{(n-r)! \cdot r!} = \binom{n}{r}.$$
 ■

Binomial identities

- Binomial identity examples:

- Example-1:: Pascal's triangle.



[Ref: Kenneth H. Rosen, Kamala Krithivasan, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2021.]

Discrete Mathematics

Dept. of CSE, NITP
16

Dr. Suddhasil De

Extended binomial coefficients

- Extended binomial coefficient: for $u \in \mathbb{R}$, $r \in \mathbb{N}$, **extended**

$$\text{binomial coefficient, } \binom{u}{r} = \begin{cases} \frac{u \cdot (u-1) \cdot \dots \cdot (u-(r-1))}{r!}, & r > 0 \\ 1, & r = 0 \end{cases}$$

$$= \begin{cases} \frac{u!}{r! \cdot (u-r)!}, & r > 0 \\ 1, & r = 0. \end{cases}$$

- Property:: **Negative extended binomial coefficient**: for $u \in \mathbb{R}^+$, $r \in \mathbb{N}$,

$$\binom{-u}{r} = \begin{cases} \frac{(-1)^r \cdot (u+r-1)!}{r! \cdot (u-1)!}, & r > 0 \\ 1, & r = 0 \end{cases}, \text{ where } \frac{(-1)^r \cdot (u+r-1)!}{r! \cdot (u-1)!} = (-1)^r \cdot \binom{u+r-1}{r}.$$

Discrete Mathematics

Dept. of CSE, NITP
17

Dr. Suddhasil De

Extended binomial theorem

- Extended binomial theorem: extension of binomial theorem to \mathbb{R} from \mathbb{Z}^+ .
- Property:: (**Extended binomial theorem**): For variable (x) & constant (u)
 $x \in \mathbb{R}$, where $|x| < 1$, and for $u \in \mathbb{R}$, $(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} \cdot x^k$.

Discrete Mathematics

Dept. of CSE, NITP
18

Dr. Suddhasil De

Summary

- Focus: Binomial coefficients and identities.
- Binomial coefficients, and binomial expression expansion.
- Binomial theorem, with proof and examples.
- Binomial identities, with proofs.
- Pascal's identity theorem, with proof.
- Extended binomial coefficients.
- Extended binomial theorem.

References

1. [Ros21] Kenneth H. Rosen, Kamala Krithivasan, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2021.
2. [Ross12] Kenneth A. Ross, Charles R. B. Wright, *Discrete Mathematics*, Fifth edition, Pearson Education, 2012.
3. [Mot15] Joe L. Mott, Abraham Kandel, Theodore P. Baker, *Discrete Mathematics for Computer Scientists and Mathematicians*, Second edition, Pearson Education, 2015.
4. [Lip17] Seymour Lipschutz, Marc L. Lipson, Varsha H. Patil, *Discrete Mathematics (Schaum's Outlines)*, Revised Third edition, McGraw-Hill Education, 2017.

Further Reading

- Binomial coefficients:: [Ros21]:437.
- Binomial expression:: [Ros21]:437.
- Binomial theorem:: [Ros21]:437-438.
- Binomial identities:: [Ros21]:439-440,442-443.
- Pascal's identity theorem:: [Ros21]:440.
- Pascal's triangle:: [Ros21]:440-441.
