

UNIT – II, Module – 3

Lecture 19: Relations

[Binary relation; Homogeneous, heterogeneous relations; Reflexivity, symmetry, antisymmetry, transitivity; Binary relation operations]

Dr. Suddhasil De

Notation table

Symbol / Notation	Meaning
\mathcal{R}	Relation.
$\bar{\mathcal{R}}$	No relation.
\mathcal{R}^{-1}	Inverse of relation \mathcal{R} .
$\bar{\mathcal{R}}$	Complementary of relation \mathcal{R} .
Δ	Diagonal relation.
\mathcal{R}^p -relation	Relation fulfilling property \mathcal{R}^p .

Relations

- Binary relation (also commonly referred as *relation*): For sets **A** and **B**, binary relation from **A** to **B** = **subset** of $A \times B$.
- Property:: Binary relation from **A** to **B** = set \mathcal{R} of ordered pairs (i.e., ordered 2-tuples) as elements, i.e., $\mathcal{R} = \{(a, b) \mid (a \in A) \wedge (b \in B)\}$.
- Property:: $a\mathcal{R}b \equiv (a, b) \in \mathcal{R} \equiv$ "*a is related to b by \mathcal{R}* " \equiv "*a is \mathcal{R} -related to b*"; Domain of $\mathcal{R} = A$; Range of $\mathcal{R} = B$.
No relation: $a\bar{\mathcal{R}}b \equiv \neg(a\mathcal{R}b) \equiv \neg((a, b) \in \mathcal{R}) \equiv (a, b) \notin \mathcal{R}$.
- Applicability:: to model wide variety of concepts in mathematics —
 - (i) "greater than", "equal to", "divides" relations in arithmetic,
 - (ii) "is adjacent to" relation in graph theory etc.

Relations


- Binary relation:
 - Property:: Relation on set A = *Homogeneous* relation on A = Binary relation from A to A = subset of $A \times A$.
 - Property:: *Universal relation* = $A \times A$. *Empty relation* = \emptyset .
 - Property:: (*Theorem*): " 2^{n^2} distinct homogeneous relations possible on a set A of n elements."
- Proof: As $|A| = n$, so $|A \times A| = n^2$. Again, relation on $A \subseteq A \times A$.
 \therefore Number of relations on A = numbers of subsets of $A \times A = 2^{n^2}$. ■
- Property:: *Heterogeneous* relation: binary relation from A to B , with possibly A and B containing distinct elements (i.e., $A \cap B \neq \emptyset$).

Relations

- Binary relation:
 - Property:: *Inverse* binary relation of 'relation \mathcal{R} from A to B ': denoted as \mathcal{R}^{-1} ; $\mathcal{R}^{-1} = \{(b, a) \mid (a, b) \in \mathcal{R}\} = \{(b, a) \mid (a \in A) \wedge (b \in B)\}$.
 - Property:: *Complementary* binary relation of 'relation \mathcal{R} from A to B ': denoted as $\bar{\mathcal{R}}$; $\bar{\mathcal{R}} = \{(a, b) \mid (a, b) \notin \mathcal{R}\} = \{(a, b) \mid (a \in \bar{A}) \vee (b \in \bar{B})\}$.
 - Property:: *Diagonal* relation on set A : denoted as Δ ; $\Delta = \{(a, a) \mid a \in A\}$; subset of $A \times A$.

Relations

- Binary relation examples:
 - Example-1:: For sets $A = \{0, 1, 2\}$, $B = \{a, b\}$, their Cartesian products $A \times B$ to form relation \mathcal{R} from A to B : $A \times B = \mathcal{R} = \{(0, a), (0, b), (1, a), (2, b)\}$.
So, $0\mathcal{R}a$, but $1\not\mathcal{R}b$.
 - Example-2:: For set $A = \{1, 2, 3, 4\}$, to find ordered pairs in relation $\mathcal{R} = \{(a, b) \mid a, b \in A, a \text{ divides } b\}$.
So, $\mathcal{R} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}$.
Above elements of satisfying " a divides b ", and $a, b \in A$.



[Ref: Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.]

Relations

- Binary relation examples:
 - Example-3:: For following relations (on infinite set), where $a, b \in \mathbb{Z}$:
 $\mathcal{R}_1 = \{(a, b) \mid a \leq b\}$. $\mathcal{R}_2 = \{(a, b) \mid a > b\}$. $\mathcal{R}_3 = \{(a, b) \mid a = b\}$.
 $\mathcal{R}_4 = \{(a, b) \mid a = b \text{ or } a = -b\}$. $\mathcal{R}_5 = \{(a, b) \mid a = b + 1\}$.
 $\mathcal{R}_6 = \{(a, b) \mid a + b \leq 3\}$,
to find relations containing each of pairs:
(1,1), (1,2), (2,1), (1,-1), and (2,2).
(1,1): member of $\mathcal{R}_1, \mathcal{R}_3, \mathcal{R}_4, \mathcal{R}_6$.
(1,2): member of $\mathcal{R}_1, \mathcal{R}_6$. (2,1): member of $\mathcal{R}_2, \mathcal{R}_5, \mathcal{R}_6$.
(1,-1): member of $\mathcal{R}_2, \mathcal{R}_4, \mathcal{R}_6$. (2,2): member of $\mathcal{R}_1, \mathcal{R}_3, \mathcal{R}_4$.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

7

Relations

- Binary relation *types*: based on property \mathcal{P} being satisfied or not.
- Property:: \mathcal{P} -relation: any binary relation fulfilling property \mathcal{P} .
- Property:: Typical types of \mathcal{P} : reflexivity, irreflexivity, symmetry, asymmetry, antisymmetry, transitivity etc.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

8

Relations

- Binary relation *types*:
 - Reflexivity property:: **Reflexive** binary relation \mathcal{R} on finite set A : $a\mathcal{R}a$ for every element $a \in A$, i.e., $\mathcal{R} = \{(a, a) \mid \forall a \in A\}$, i.e.,
 $\forall a \in A ((a, a) \in \mathcal{R}) \equiv \forall a ((a \in A) \rightarrow ((a, a) \in \mathcal{R}))$.
In words, reflexive binary relation \mathcal{R} on finite set A to relate every element of A to itself by \mathcal{R} .
 - Irreflexivity property:: **Irreflexive** binary relation \mathcal{R} on finite set A : $a\mathcal{R}a$ for every element $a \in A$, i.e., $\forall a \in A ((a, a) \notin \mathcal{R})$.
In words, irreflexive binary relation \mathcal{R} on finite set A not to relate any element of A to itself by \mathcal{R} .

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

9

Relations

- Binary relation *types*:
- Property:: Relation \mathcal{R} on finite set A **neither reflexive nor irreflexive**:
 $(\exists a \in A ((a, a) \notin \mathcal{R})) \wedge (\exists b \in A ((b, b) \in \mathcal{R}))$.
- Property:: Binary relation \mathcal{R} on finite set A irreflexive, if and only if binary relation $((A \times A) \setminus \mathcal{R})$ reflexive.

Relations

- Binary relation *types*:
- Symmetry property:: **Symmetric** binary relation \mathcal{R} on finite set A : **$b\mathcal{R}a$ whenever $a\mathcal{R}b$** for every element $a, b \in A$, i.e.,
$$\forall a \in A \forall b \in A ((a, b) \in \mathcal{R}) \rightarrow ((b, a) \in \mathcal{R})$$

In words, symmetric binary relation \mathcal{R} on finite set A to relate element b of A to element a of A by \mathcal{R} , whenever a related to b by \mathcal{R} .
- Property:: \mathcal{R} on set A **not symmetric**: $\exists a \in A \exists b \in A (((a, b) \in \mathcal{R}) \wedge ((b, a) \notin \mathcal{R}))$.

Relations

- Binary relation *types*:
- Asymmetry property:: **Asymmetric** binary relation \mathcal{R} on finite set A : **$b\mathcal{R}a$ whenever $a\mathcal{R}b$** for every element $a, b \in A$, i.e.,
$$\forall a \in A \forall b \in A (((a, b) \in \mathcal{R}) \rightarrow ((b, a) \notin \mathcal{R}))$$
$$\equiv \forall a \in A \forall b \in A \neg (((a, b) \in \mathcal{R}) \wedge ((b, a) \in \mathcal{R}))$$

In words, asymmetric binary relation \mathcal{R} on finite set A to not relate every element b of A to every other element a of A by \mathcal{R} , whenever a related to b by \mathcal{R} .

Relations

- Binary relation *types*:
 - Property: \mathcal{R} on set A **neither symmetric nor asymmetric**:
$$\left(\exists a \in A \exists b \in A \left((a, b) \in \mathcal{R} \wedge (b, a) \notin \mathcal{R} \right) \right) \wedge$$
$$\left(\exists c \in A \exists d \in A \left((c, d) \in \mathcal{R} \wedge (d, c) \in \mathcal{R} \right) \right).$$

Relations

- Binary relation *types*:
 - Antisymmetry property: **Antisymmetric** binary relation \mathcal{R} on finite set A : $a = b$ whenever $a \mathcal{R} b$ and $b \mathcal{R} a$ for every element $a, b \in A$, i.e.,
$$\forall a \in A \forall b \in A \left(((a, b) \in \mathcal{R}) \wedge ((b, a) \in \mathcal{R}) \rightarrow (a = b) \right).$$

In words, antisymmetric binary relation \mathcal{R} on finite set A to not relate any pairs of distinct elements of A to each other by \mathcal{R} .
 - Property: \mathcal{R} on set A **not antisymmetric**:
$$\exists a \in A \exists b \in A \left((a \neq b) \wedge ((a, b) \in \mathcal{R}) \wedge ((b, a) \in \mathcal{R}) \right).$$

Relations

- Binary relation *types*:
 - Property: Possibility of relation \mathcal{R} to **hold** both antisymmetric and symmetric properties. E.g., (i) $\mathcal{R} = \emptyset$, $A = \{a\}$ (vacuously satisfied); (ii) $\mathcal{R} = \{(1,1), (2,2)\}$, $A = \mathbb{Z}$.
 - Property: Possibility of relation \mathcal{R} to **not satisfy** combined symmetric, asymmetric and antisymmetric properties (i.e., **neither symmetric nor asymmetric nor antisymmetric**).
E.g., $\mathcal{R} = \{(a, b), (b, a), (a, c)\}$, $A = \{a, b, c\}$ (where, $a \neq b \neq c$); reasons: $(c, a) \notin \mathcal{R}$, $(b, a) \in \mathcal{R}$, and $a \neq b$.

Relations

- Binary relation *types*:
- Transitivity property: **Transitive** binary relation \mathcal{R} on finite set A : $a\mathcal{R}c$ whenever $a\mathcal{R}b$ and $b\mathcal{R}c$ for every element $a, b, c \in A$, i.e.,

$$\forall a \in A \forall b \in A \forall c \in A \left(((a, b) \in \mathcal{R}) \wedge ((b, c) \in \mathcal{R}) \rightarrow ((a, c) \in \mathcal{R}) \right).$$

In words, transitive binary relation \mathcal{R} on finite set A to relate element a of A to element c of A by \mathcal{R} , whenever a related to element b of A by \mathcal{R} and b related to c by \mathcal{R} .
- Property: \mathcal{R} on set A **not transitive**:

$$\exists a \in A \exists b \in A \exists c \in A \left(((a, b) \in \mathcal{R}) \wedge ((b, c) \in \mathcal{R}) \wedge ((a, c) \notin \mathcal{R}) \right).$$

Discrete Mathematics

Dept. of CSE, NITP
16

Dr. Suddhasil De

Relations

- Binary relation *types*:
 - Property: (**Theorem**): " $2^{n \cdot (n-1)}$ reflexive relations possible on a set of n elements."
- Proof:** [Using basic counting principle] Constituents of job: 2 tasks.
 Let A = arbitrary set of n elements, \mathcal{R} = some arbitrary relation.
 Then, \mathcal{R} on A = subset of $A \times A$, i.e., \mathcal{R} to specify whether each of n^2 ordered pairs in $A \times A$ to belong to \mathcal{R} .
 Task-1: \mathcal{R} reflexive. Then, \mathcal{R} to contain all n ordered pairs (a, a) for $a \in A$. So, 1 subset of $A \times A$ holding all (a, a) ordered pairs of reflexive relation formed based on ordered pairing of same element. (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
17

Dr. Suddhasil De

Relations

- Binary relation *types*:
- Proof** contd.:
 For any $a, b \in A$, considering ordered pairs of form (a, b) , where $a \neq b$, as per product rule, number of such ordered pairs = $n \cdot (n-1)$.
 However, each of $n \cdot (n-1)$ ordered pairs may or may not be in \mathcal{R} .
 Task-2: selecting $n \cdot (n-1)$ ordered pairs.
 Constituent: $n \cdot (n-1)$ sub-tasks.
 Sub-task T_i ($i = 1, 2, \dots, n \cdot (n-1)$): choice of adding or not i -th ordered pair of distinct elements in \mathcal{R} .
 Single procedure to perform sub-task T_i . (contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
18

Dr. Suddhasil De

Relations

- Binary relation *types*:

Proof contd-2.:

Sub-task T_i performing regardless of how T_1, T_2, \dots, T_{i-1} performed.

Then, according to product rule, number of subsets of $A \times A$ for all (a, b) ordered pairs in reflexive relation formed based on ordered

pairing of distinct elements $= \underbrace{2 \cdot 2 \cdot 2 \cdot \dots \cdot 2}_{n \cdot (n-1) \text{ factors}} = 2^{n \cdot (n-1)}$.

So, task-2 providing $2^{n \cdot (n-1)}$ relations \mathcal{R} .

Task-2 performing regardless of how task-1 performed.

So, again applying product rule, number of $\mathcal{R} = 1 \cdot 2^{n \cdot (n-1)}$. ■

Discrete Mathematics

Dept. of CSE, NITP
19

Dr. Suddhasil De

Relations

- Binary relation *types* examples:

- Example-1:: Given relations on $\{1,2,3,4\}$, to find reflexive relations:

$\mathcal{R}_1 = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,1), (4,4)\}$.

$\mathcal{R}_2 = \{(1,1), (1,2), (2,1)\}$.

$\mathcal{R}_3 = \{(1,1), (1,2), (1,4), (2,1), (2,2), (3,3), (4,1), (4,4)\}$.

$\mathcal{R}_4 = \{(2,1), (3,1), (3,2), (4,1), (4,2), (4,3)\}$.

$\mathcal{R}_5 = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,3), (2,4), (3,3), (3,4), (4,4)\}$.

$\mathcal{R}_3 \rightarrow$ reflexive: as all $a\mathcal{R}_3a$ present, i.e. $\forall a((a,a) \in \mathcal{R}_3)$; here,
 $(1,1), (2,2), (3,3), (4,4) \in \mathcal{R}_3$.

$\mathcal{R}_5 \rightarrow$ reflexive: as $(1,1), (2,2), (3,3), (4,4) \in \mathcal{R}_5$.

(contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
20

Dr. Suddhasil De

Relations

- Binary relation *types* examples:

- Example-1 contd.

For other given relations: $(3,3) \notin \mathcal{R}_1$; $(2,2), (3,3), (4,4) \notin \mathcal{R}_2$;

$(1,1), (2,2), (3,3), (4,4) \notin \mathcal{R}_4$.

So, $\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_4 \rightarrow$ not reflexive.

- Example-2:: To find reflexive nature of "divides" ($|$) relation on \mathbb{Z}^+ .

Considering \mathcal{R} as ' $|$ ', $\mathcal{R} = \{(a,b) \mid a, b \in \mathbb{Z}^+, a|b\}$, where $a|b =$
 a divides b .

As " a divides a ", so $(a,a) \in \mathcal{R}$, i.e., $a\mathcal{R}a$.

So, ' $|$ ' \rightarrow reflexive.

Discrete Mathematics

Dept. of CSE, NITP
21

Dr. Suddhasil De

Relations

- Binary relation *types* examples:
 - Example-3:: Given relations on {1,2,3,4}, to find symmetric relations:
 $\mathcal{R}_1 = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,1), (4,4)\}$.
 $\mathcal{R}_2 = \{(1,1), (1,2), (2,1)\}$.
 $\mathcal{R}_3 = \{(1,1), (1,2), (1,4), (2,1), (2,2), (3,3), (4,1), (4,4)\}$.
 $\mathcal{R}_4 = \{(2,1), (3,1), (3,2), (4,1), (4,2), (4,3)\}$.
 $\mathcal{R}_5 = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,3), (2,4), (3,3), (3,4), (4,4)\}$.
 $\mathcal{R}_2 \rightarrow$ symmetric: as for each such case whenever $a\mathcal{R}_2b$ present, then $b\mathcal{R}_2a$ also present; here, $(1,2), (2,1) \in \mathcal{R}_2$, and no other case in \mathcal{R}_2 .

(contd. to next slide)

Relations

- Binary relation *types* examples:
 - Example-3 contd.
 $\mathcal{R}_3 \rightarrow$ symmetric: as, $(1,2), (2,1), (1,4), (4,1) \in \mathcal{R}_3$.
For other given relations: $(4,1) \in \mathcal{R}_1$, but $(1,4) \notin \mathcal{R}_1$;
 $(2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \in \mathcal{R}_4$, but
 $(1,2), (1,3), (2,3), (1,4), (2,4), (3,4) \notin \mathcal{R}_4$;
 $(1,2), (1,3), (1,4), (2,3), (2,4), (3,4) \in \mathcal{R}_5$, but
 $(2,1), (3,1), (4,1), (3,2), (4,2), (4,3) \notin \mathcal{R}_5$.
So, $\mathcal{R}_1, \mathcal{R}_4, \mathcal{R}_5 \rightarrow$ not symmetric.

Relations

- Binary relation *types* examples:
 - Example-4:: To find symmetric nature of "equality" ('=') relation on \mathbb{Z} .
Considering \mathcal{R} as '=', $\mathcal{R} = \{(a,b) \mid a,b \in \mathbb{Z}, a \text{ equals } b\}$.
As " a equals b " \rightarrow " b equals a ", so $\left(((a,b) \in \mathcal{R}) \rightarrow ((b,a) \in \mathcal{R}) \right)$,
i.e., $b\mathcal{R}a$ whenever $a\mathcal{R}b$.
So, '=' \rightarrow symmetric.

Relations

- Binary relation *types* examples:
 - Example-5:: Given relations on {1,2,3,4}, to find antisymmetric relations:
 $\mathcal{R}_1 = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,1), (4,4)\}$.
 $\mathcal{R}_2 = \{(1,1), (1,2), (2,1)\}$.
 $\mathcal{R}_3 = \{(1,1), (1,2), (1,4), (2,1), (2,2), (3,3), (4,1), (4,4)\}$.
 $\mathcal{R}_4 = \{(2,1), (3,1), (3,2), (4,1), (4,2), (4,3)\}$.
 $\mathcal{R}_5 = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,3), (2,4), (3,3), (3,4), (4,4)\}$.
For determining relations satisfying antisymmetric property, to start with relations not holding symmetric property.

Relations

- Binary relation *types* examples:
 - Example-5 contd.
 $\mathcal{R}_4 \rightarrow$ antisymmetric: as no such case possible, whenever $a\mathcal{R}_4b$ and $b\mathcal{R}_4a$ present, then $a = b$; here,
 $(2,1), (3,1), (3,2), (4,1), (4,2), (4,3) \in \mathcal{R}_4$, but
 $(1,2), (1,3), (2,3), (1,4), (2,4), (3,4) \notin \mathcal{R}_4$, and so no equality scenario.
 $\mathcal{R}_5 \rightarrow$ antisymmetric: as, similar to \mathcal{R}_4 , no such case possible; here,
 $(1,2), (1,3), (1,4), (2,3), (2,4), (3,4) \in \mathcal{R}_5$, but
 $(2,1), (3,1), (4,1), (3,2), (4,2), (4,3) \notin \mathcal{R}_5$, and so no equality comparison scenario.

Relations

- Binary relation *types* examples:
 - Example-5 contd-2.
For other given relations: $(1,2) \in \mathcal{R}_1$ and $(2,1) \in \mathcal{R}_1$, but $1 \neq 2$;
Same reasoning of \mathcal{R}_1 also for \mathcal{R}_2 ; and \mathcal{R}_3 .
So, $\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3 \rightarrow$ not antisymmetric.

Relations

- Binary relation *types* examples:
 - Example-6:: To find antisymmetric nature of "less than or equal to" (\leq) relation on \mathbb{Z} .
 Considering \mathcal{R} as ' \leq ', $\mathcal{R} = \{(a, b) \mid a, b \in \mathbb{Z}, a \text{ less or equal to } b\}$.
 As " a less or equal to b " and " b less or equal to a " \rightarrow " a equals b ", so

$$\left(((a, b) \in \mathcal{R}) \wedge ((b, a) \in \mathcal{R}) \right) \rightarrow (a = b),$$
 i.e., when $a\mathcal{R}b$ and $b\mathcal{R}a$, then $a = b$.
 So, ' \leq ' \rightarrow antisymmetric.

Discrete Mathematics

Dept. of CSE, NITP
28

Dr. Suddhasil De

Relations

- Binary relation *types* examples:
 - Example-7:: Given relations on $\{1,2,3,4\}$, to find transitive relations:
 $\mathcal{R}_1 = \{(1,1), (1,2), (2,1), (2,2), (3,4), (4,1), (4,4)\}$.
 $\mathcal{R}_2 = \{(1,1), (1,2), (2,1)\}$.
 $\mathcal{R}_3 = \{(1,1), (1,2), (1,4), (2,1), (2,2), (3,3), (4,1), (4,4)\}$.
 $\mathcal{R}_4 = \{(2,1), (3,1), (3,2), (4,1), (4,2), (4,3)\}$.
 $\mathcal{R}_5 = \{(1,1), (1,2), (1,3), (1,4), (2,2), (2,3), (2,4), (3,3), (3,4), (4,4)\}$.
 Here, $\mathcal{R}_4 \rightarrow$ transitive: as for each such case, whenever $a\mathcal{R}_4b$ and $b\mathcal{R}_4c$ present, then $a\mathcal{R}_4c$ present; here, $(3,2), (2,1), (3,1) \in \mathcal{R}_4$,
 $(4,2), (2,1), (4,1) \in \mathcal{R}_4$, $(4,3), (3,2), (4,2) \in \mathcal{R}_4$, $(4,3), (3,1), (4,1) \in \mathcal{R}_4$.
(contd. to next slide)

Discrete Mathematics

Dept. of CSE, NITP
29

Dr. Suddhasil De

Relations

- Binary relation *types* examples:
 - Example-7 contd.
 $\mathcal{R}_5 \rightarrow$ transitive: as $(1,2), (2,3), (1,3) \in \mathcal{R}_5$, $(1,2), (2,4), (1,4) \in \mathcal{R}_5$,
 $(1,3), (3,4), (1,4) \in \mathcal{R}_5$, $(2,3), (3,4), (2,4) \in \mathcal{R}_5$, to satisfy all possible
 transitive conditions in \mathcal{R}_5 .
 For other given relations: $(3,4) \in \mathcal{R}_1$ and $(4,1) \in \mathcal{R}_1$, but $(3,1) \notin \mathcal{R}_1$,
 $(4,1) \in \mathcal{R}_1$ and $(1,2) \in \mathcal{R}_1$, but $(4,2) \notin \mathcal{R}_1$;
 $(2,1) \in \mathcal{R}_2$ and $(1,2) \in \mathcal{R}_2$, but $(2,2) \notin \mathcal{R}_2$;
 $(4,1) \in \mathcal{R}_3$ and $(1,2) \in \mathcal{R}_3$, but $(4,2) \notin \mathcal{R}_3$.
 So, $\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3 \rightarrow$ not transitive.

Discrete Mathematics

Dept. of CSE, NITP
30

Dr. Suddhasil De

Relations

- Binary relation *types* examples:
 - Example-8:: To find transitive nature of "divides" ($|$) relation on \mathbb{Z}^+ .
Considering \mathcal{R} as ' $|$ ', $\mathcal{R} = \{(a, b) \mid a, b \in \mathbb{Z}^+, a|b\}$, where $a|b = a$ divides b .
For any $a, b, c \in \mathbb{Z}^+$, let " a divides b ". Then, $b = k \cdot a$, for some $k \in \mathbb{Z}^+$.
Again let " b divides c ". Then, $c = l \cdot b$, for some $l \in \mathbb{Z}^+$.
 $\therefore c = l \cdot k \cdot a = (l \cdot k) \cdot a$ where $(l \cdot k) \in \mathbb{Z}^+$, i.e., " a divides c ". So,
 $\left(((a, b) \in \mathcal{R}) \wedge ((b, c) \in \mathcal{R}) \right) \rightarrow ((a, c) \in \mathcal{R})$,
i.e., when $a\mathcal{R}b$ and $b\mathcal{R}c$, then $a\mathcal{R}c$.
So, ' $|$ ' \rightarrow transitive.

Discrete Mathematics

Dept. of CSE, NITP
31

Dr. Suddhasil De

Relations

- Binary relation *operations*: similar to operations on sets.
 - Union property:: **Union** of two binary relations \mathcal{R}_1 and \mathcal{R}_2 on finite sets **A** and **B**: $\mathcal{R}_1 \cup \mathcal{R}_2 = \{(a, b) \mid ((a, b) \in \mathcal{R}_1) \vee ((a, b) \in \mathcal{R}_2)\}$.
 - Intersection property:: **Intersection** of two binary relations \mathcal{R}_1 and \mathcal{R}_2 on finite sets **A** and **B**: $\mathcal{R}_1 \cap \mathcal{R}_2 = \{(a, b) \mid ((a, b) \in \mathcal{R}_1) \wedge ((a, b) \in \mathcal{R}_2)\}$.
 - Subtraction property:: **Subtraction** of two binary relations \mathcal{R}_1 and \mathcal{R}_2 on finite sets **A** and **B**: $\mathcal{R}_1 \setminus \mathcal{R}_2 = \{(a, b) \mid ((a, b) \in \mathcal{R}_1) \wedge ((a, b) \notin \mathcal{R}_2)\}$.

Discrete Mathematics

Dept. of CSE, NITP
32

Dr. Suddhasil De

Relations

- Binary relation *operations*:
 - Property:: **Symmetric difference** of two binary relations \mathcal{R}_1 and \mathcal{R}_2 on finite sets **A** and **B**: $\mathcal{R}_1 \oplus \mathcal{R}_2 = \mathcal{R}_1 \ominus \mathcal{R}_2 = \mathcal{R}_1 \Delta \mathcal{R}_2 = \{(a, b) \mid ((a, b) \in \mathcal{R}_1) \oplus ((a, b) \in \mathcal{R}_2)\}$.
 - Composition property:: **Composition** (or **composite**) of binary relations \mathcal{R} (from finite sets **A** to **B**) and \mathcal{S} (from finite sets **B** to **C**): $\mathcal{S} \circ \mathcal{R} = \{(a, c) \mid a \in \mathbf{A}, c \in \mathbf{C}, \exists b \in \mathbf{B} ((a, b) \in \mathcal{R}) \wedge ((b, c) \in \mathcal{S})\}$.
 - Power property:: **Powers** of binary relation \mathcal{R} on finite set **A**: $\mathcal{R}^{n+1} = \mathcal{R}^n \circ \mathcal{R}$, provided $\mathcal{R}^1 = \mathcal{R}$, where, $n \in \mathbb{N} \setminus \{0\}$.
 - Property:: $\mathcal{R}^2 = \mathcal{R} \circ \mathcal{R}$.

Discrete Mathematics

Dept. of CSE, NITP
33

Dr. Suddhasil De

Relations

- Binary relation *operations*:
 - Property:: (**Theorem**): Binary relation R on finite set A to become transitive if and only if $R^n \subseteq R$ for $n \in \mathbb{N} \setminus \{0\}$.

Discrete Mathematics

Dept. of CSE, NITP
34

Dr. Suddhasil De

Summary

- Focus: Relations.
- Binary relation, and properties.
- Homogeneous and heterogeneous relations, and properties.
- Binary relation types, and \mathcal{P} -relation.
- Reflexive and irreflexive binary relations, with examples.
- Symmetric and asymmetric binary relations, with examples.
- Antisymmetric binary relation, with examples.
- Transitive binary relation, with examples.
- Union of two binary relations.
- Intersection of two binary relations.

Discrete Mathematics

Dept. of CSE, NITP
35

Dr. Suddhasil De

Summary

- Subtraction of two binary relations.
- Symmetric difference of two binary relations.
- Composition of two binary relations.
- Powers of binary relation.

Discrete Mathematics

Dept. of CSE, NITP
36

Dr. Suddhasil De

References

1. [Ros19] Kenneth H. Rosen, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2019.

2. [Mot08] Joe L. Mott, Abraham Kandel, Theodore P. Baker, *Discrete Mathematics for Computer Scientists and Mathematicians*, PHI, Second edition, 2008.

3. [Lip07] Seymour Lipschutz and Marc Lars Lipson, *Schaum's Outline of Theory and Problems of Discrete Mathematics*, Third edition, McGraw-Hill Education, 2007.

Further Reading

- Binary relations:: [Ros19]:599-601.
- Homogeneous relations:: [Ros19]:601-602.
- Binary relation types:: [Ros19]:602-605.
- Reflexive binary relations:: [Ros19]:602-603,605.
- Symmetric binary relations:: [Ros19]:603-604.
- Antisymmetric binary relations:: [Ros19]:603-604.
- Transitive binary relations:: [Ros19]:604-605.
- Union, intersection of binary relations:: [Ros19]:606.
- Composite of binary relations:: [Ros19]:606-608.
