

CS34110

Discrete Mathematics and Graph Theory

UNIT – II, Module – 1

Lecture 10: Discrete Structures

[Multiset; Multiset operations; Function; Injection, surjection, bijection; Increasing, decreasing functions; Identity, inverse functions]

Dr. Suddhasil De

Recap:: Discrete structures

- Discrete structures examples:
 - (i) **sets** (collections of objects) and **multisets**;
 - (ii) combinations (built from sets; unordered collections of objects to be used in counting);
 - (iii) relations (sets of ordered pairs representing relationships between objects);
 - (iv) graphs (sets of vertices and edges connecting vertices);
 - (v) sequences (ordered lists of elements, as well as special type of functions expressing relationships among elements)
 - (vi) **matrices**.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

2

Multiset

- Multiset** (short for **Multiple-membership set**; also called bag, or mset): unordered collection of multiple instances of distinct members, i.e., possibility of any element to occur as member more than once.
- Property:: Multiset $A = \{m_1 \cdot a_1, m_2 \cdot a_2, \dots, m_r \cdot a_r\}$, where element a_1 occurring m_1 times, element a_2 occurring m_2 times, and so on. Alternate representation:: $A = \{a_1^{m(a_1)}, a_2^{m(a_2)}, \dots, a_r^{m(a_r)}\}$, where m = multiplicity function.
- Property:: **Multiplicities** of members of multiset: m_i ($i = 1, 2, \dots, r$), of elements a_i , each m_i being nonnegative integer.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

3

Multiset

- Multiset:
 - Property:: **Cardinality of multiset \mathbf{M}** : $|\mathbf{M}| = \sum_{i=1}^r m_i = \sum_{x \in \mathcal{U}} m(x)$
 - Inclusion property:: Multiset \mathbf{M} to be **subset** of multiset \mathbf{N} :
 $\mathbf{M} \subseteq \mathbf{N} (\equiv \mathbf{N} \supseteq \mathbf{M})$, if and only if $\forall x \left((x \in \mathcal{U}) \wedge (m_{\mathbf{M}}(x) \leq m_{\mathbf{N}}(x)) \right)$.
 - Union property:: **Union** of multisets \mathbf{M} and \mathbf{N} :
 $\mathbf{M} \cup \mathbf{N} = \{x \in \mathcal{U} \mid m_{\mathbf{M} \cup \mathbf{N}}(x) = \max(m_{\mathbf{M}}(x), m_{\mathbf{N}}(x))\}$.
 - Intersection property:: **Intersection** of multisets \mathbf{M} and \mathbf{N} :
 $\mathbf{M} \cap \mathbf{N} = \{x \in \mathcal{U} \mid m_{\mathbf{M} \cap \mathbf{N}}(x) = \min(m_{\mathbf{M}}(x), m_{\mathbf{N}}(x))\}$.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

4

Multiset

- Multiset:
 - Subtraction property:: **Difference** between multisets \mathbf{M} and \mathbf{N} :
 $\mathbf{M} \setminus \mathbf{N} = \mathbf{M} - \mathbf{N} = \{x \in \mathcal{U} \mid m_{\mathbf{M} \setminus \mathbf{N}}(x) = \max(m_{\mathbf{M}}(x) - m_{\mathbf{N}}(x), 0)\}$.
 - Addition property:: **Sum** of multisets \mathbf{M} and \mathbf{N} :
 $\mathbf{M} + \mathbf{N} = \{x \in \mathcal{U} \mid m_{\mathbf{M} + \mathbf{N}}(x) = (m_{\mathbf{M}}(x) + m_{\mathbf{N}}(x))\}$.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

5

Multiset

- Multiset examples:
 - Example-1:: For multisets $\mathbf{P} = \{4 \cdot a, 1 \cdot b, 3 \cdot c\}$, $\mathbf{Q} = \{3 \cdot a, 4 \cdot b, 2 \cdot d\}$:
 $\mathbf{P} \cup \mathbf{Q} = \{\max(m_{\mathbf{P}}(a), m_{\mathbf{Q}}(a)) \cdot a, \max(m_{\mathbf{P}}(b), m_{\mathbf{Q}}(b)) \cdot b, \max(m_{\mathbf{P}}(c), m_{\mathbf{Q}}(c)) \cdot c, \max(m_{\mathbf{P}}(d), m_{\mathbf{Q}}(d)) \cdot d\}$
 $= \{\max(4, 3) \cdot a, \max(1, 4) \cdot b, \max(3, 0) \cdot c, \max(0, 2) \cdot d\}$
 $= \{4 \cdot a, 4 \cdot b, 3 \cdot c, 2 \cdot d\} = \{a^{(4)}, b^{(4)}, c^{(3)}, d^{(2)}\}$.
 - $\mathbf{P} \cap \mathbf{Q} = \{\min(m_{\mathbf{P}}(a), m_{\mathbf{Q}}(a)) \cdot a, \min(m_{\mathbf{P}}(b), m_{\mathbf{Q}}(b)) \cdot b, \min(m_{\mathbf{P}}(c), m_{\mathbf{Q}}(c)) \cdot c, \min(m_{\mathbf{P}}(d), m_{\mathbf{Q}}(d)) \cdot d\}$
 $= \{\min(4, 3) \cdot a, \min(1, 4) \cdot b, \min(3, 0) \cdot c, \min(0, 2) \cdot d\} = \{3 \cdot a, 1 \cdot b\}$.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

6

Multiset

- Multiset examples:
 - Example-1 (contd.)::
$$\mathbf{P} \setminus \mathbf{Q} = \left\{ \max \left((m_{\mathbf{P}}(a) - m_{\mathbf{Q}}(a)), 0 \right) \cdot a, \max \left((m_{\mathbf{P}}(b) - m_{\mathbf{Q}}(b)), 0 \right) \cdot b, \max \left((m_{\mathbf{P}}(c) - m_{\mathbf{Q}}(c)), 0 \right) \cdot c, \max \left((m_{\mathbf{P}}(d) - m_{\mathbf{Q}}(d)), 0 \right) \cdot d \right\}$$
$$= \{ \max(1,0) \cdot a, \max(-3,0) \cdot b, \max(3,0) \cdot c, \max(-2,0) \cdot d \}$$
$$= \{ 1 \cdot a, 3 \cdot c \}.$$
$$\mathbf{P} + \mathbf{Q} = \left\{ (m_{\mathbf{P}}(a) + m_{\mathbf{Q}}(a)) \cdot a, (m_{\mathbf{P}}(b) + m_{\mathbf{Q}}(b)) \cdot b, (m_{\mathbf{P}}(c) + m_{\mathbf{Q}}(c)) \cdot c, (m_{\mathbf{P}}(d) + m_{\mathbf{Q}}(d)) \cdot d \right\} = \{ 7 \cdot a, 5 \cdot b, 3 \cdot c, 2 \cdot d \}.$$

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

7

Recap:: Discrete structures

- Discrete structures examples:
 - (i) **sets** (collections of objects) and **multisets**;
 - (ii) combinations (built from sets; unordered collections of objects to be used in counting);
 - (iii) relations (sets of ordered pairs representing relationships between objects);
 - (iv) graphs (sets of vertices and edges connecting vertices);
 - (v) sequences (ordered lists of elements, as well as special type of **functions** expressing relationships among elements)
 - (vi) **matrices**.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

8

Function

- **Function** f from nonempty set \mathbf{A} to nonempty set \mathbf{B} : $f: \mathbf{A} \rightarrow \mathbf{B}$ = " f maps \mathbf{A} to \mathbf{B} " = assignment " $f(a) = b$ " of exactly one element $b \in \mathbf{B}$ corresponding to each element $a \in \mathbf{A}$.
 - [Alternate] $f: \mathbf{A} \rightarrow \mathbf{B}$ = relation from \mathbf{A} to \mathbf{B} containing one, and only one, ordered pair (a, b) , for every element $a \in \mathbf{A}$, with a as first element and $b \in \mathbf{B}$ as last element, leading to assignment $f(a) = b$.
 - Property:: \mathbf{A} : **domain** of f .
 - Property:: \mathbf{B} : **codomain** of f .
 - Property:: b : **image** of a .
 - Property:: a : **preimage** of b .

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

9

Function

- Function:
 - Property: **Range** of f — set of all images of elements of A .
 - Property: Range: also called image. Range of $f \subseteq$ codomain B .
 - Property: **one-to-one** (or **injective**) function f —
 $\forall a \forall b ((f(a) = f(b)) \rightarrow (a = b)) \equiv \forall a \forall b ((a \neq b) \rightarrow (f(a) \neq f(b)))$,
where domain of discourse = domain of f .
 - Property: **onto** (or **surjective**) function f — $\forall y \exists x (f(x) = y)$, where
domain for x = domain of f , domain for y = codomain of f .
 - Property: **one-to-one correspondence** (or **bijjective**) function f — if
both one-to-one and onto.

Discrete Mathematics

Dept. of CSE, NITP
10

Dr. Suddhasil De

Function

- Function:
 - Property: **Real-valued function** — function $f: A \rightarrow \mathbb{R}$ to be real-valued,
if codomain of f = set of real numbers = \mathbb{R} .
 - Property: **Integer-valued function** — function $f: A \rightarrow \mathbb{Z}$ to be integer-
valued, if codomain of f = set of integers = \mathbb{Z} .
 - Property: **Sum** of real-valued functions $f_1: A \rightarrow \mathbb{R}, f_2: A \rightarrow \mathbb{R}$ — real-
valued sum function $f_1 + f_2: A \rightarrow \mathbb{R}$, where $(f_1 + f_2)(x) = f_1(x) + f_2(x)$,
for all $x \in A$.

Discrete Mathematics

Dept. of CSE, NITP
11

Dr. Suddhasil De

Function

- Function:
 - Property: **Product** of real-valued functions $f_1: A \rightarrow \mathbb{R}, f_2: A \rightarrow \mathbb{R}$ — real-
valued product function $f_1 \cdot f_2: A \rightarrow \mathbb{R}$, where $(f_1 \cdot f_2)(x) = f_1(x) \cdot$
 $f_2(x)$, for all $x \in A$.
 - Property: **Sum** of integer-valued functions $g_1: A \rightarrow \mathbb{Z}, g_2: A \rightarrow \mathbb{Z}$ —
integer-valued sum function $g_1 + g_2: A \rightarrow \mathbb{Z}$, where $(g_1 + g_2)(x) =$
 $g_1(x) + g_2(x)$, for all $x \in A$.
 - Property: **Product** of integer-valued functions $g_1: A \rightarrow \mathbb{Z}, g_2: A \rightarrow \mathbb{Z}$ —
integer-valued product function $g_1 \cdot g_2: A \rightarrow \mathbb{Z}$, where $(g_1 \cdot g_2)(x) =$
 $g_1(x) \cdot g_2(x)$, for all $x \in A$.

Discrete Mathematics

Dept. of CSE, NITP
12

Dr. Suddhasil De

Function

- Function:
 - Property: **Increasing** function $f: A \rightarrow B$, where $A, B \subseteq \mathbb{R}$ —
 $\forall a \forall b ((a < b) \rightarrow (f(a) \leq f(b)))$, $a, b \in A$,
where domain of discourse = domain of $f = A$.
 - Property: **Strictly increasing** function $f: A \rightarrow B$, where $A, B \subseteq \mathbb{R}$ —
 $\forall a \forall b ((a < b) \rightarrow (f(a) < f(b)))$, $a, b \in A$,
where domain of discourse = domain of $f = A$.
 - Property: **Decreasing** function $f: A \rightarrow B$, where $A, B \subseteq \mathbb{R}$ —
 $\forall a \forall b ((a < b) \rightarrow (f(a) \geq f(b)))$, $a, b \in A$,
where domain of discourse = domain of $f = A$.

Discrete Mathematics

Dept. of CSE, NITP
13

Dr. Suddhasil De

Function

- Function:
 - Property: **Strictly decreasing** function $f: A \rightarrow B$, where $A, B \subseteq \mathbb{R}$ —
 $\forall a \forall b ((a < b) \rightarrow (f(a) > f(b)))$, $a, b \in A$,
where domain of discourse = domain of $f = A$.
 - Property: One-to-one (or injective) function: either (i) strictly increasing function, or (ii) strictly decreasing function.
 - Property: Not one-to-one (or NOT injective) function:
either (i) increasing function, but not strictly increasing function,
or (ii) decreasing function, but not strictly decreasing function.
 - Property: Onto (or surjective) function: equal range and codomain.


Discrete Mathematics

Dept. of CSE, NITP
14

Dr. Suddhasil De

Function

- Function:



[Ref: Kenneth H. Rosen, Kamala Krithivasan, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2021.]

Discrete Mathematics

Dept. of CSE, NITP
15

Dr. Suddhasil De

Function

- Function examples:
 - Example-1:: For function g assigning grades $\{A+, A, B, C, D, F\}$ to set of students $\{\text{stud}_1, \text{stud}_2, \text{stud}_3, \dots, \text{stud}_n\}$ of discrete mathematics course, with no D grades being assigned to any of n students,
domain of g = set $\{\text{stud}_1, \text{stud}_2, \text{stud}_3, \dots, \text{stud}_n\}$,
codomain of g = set $\{A+, A, B, C, D, F\}$,
range of g = set $\{A+, A, B, C, F\}$.

Function

- Function examples:
 - Example-2:: For function f extracting last two bits of bit string of length 2 or greater course,
domain of f = set of all bit strings of length 2 or greater,
codomain of f = set $\{00, 01, 10, 11\}$,
range of f = set $\{00, 01, 10, 11\}$.
 - Example-3:: For $f: \mathbb{Z} \rightarrow \mathbb{Z}$ assigning squares of integer, i.e. $f(x) = x^2$,
domain of f = set of all integers \mathbb{Z} ,
codomain of f = set of all integers \mathbb{Z} ,
range of f = set of all perfect square integers = $\{0, 1, 4, 9, \dots\}$.

Function

- Function examples:
 - Example-4:: For functions $f_1: \mathbb{R} \rightarrow \mathbb{R}$, $f_2: \mathbb{R} \rightarrow \mathbb{R}$, such that $f_1(x) = x^2$, $f_2(x) = x - x^2$, their **sum** and **product** to be —
 $(f_1 + f_2)(x) = f_1(x) + f_2(x) = x^2 + x - x^2 = x$, for all $x \in \mathbb{R}$; and
 $(f_1 \cdot f_2)(x) = f_1(x) \cdot f_2(x) = x^2 \cdot (x - x^2) = x^3 - x^4$, for all $x \in \mathbb{R}$.
 - Example-5:: For function $f: \mathbb{Z}^+ \rightarrow \mathbb{Z}$, $f(x) = x^2$, to show its **one-to-one** nature, then in turn, to show $f(x)$ as **strictly increasing** —
For any $x, y \in \mathbb{Z}^+$ and assuming $x < y$, to obtain $x^2 < x \cdot y$, and $x \cdot y < y^2$. Combining, $f(x) = x^2 < x \cdot y < y^2 = f(y)$, i.e., $f(x) \neq f(y)$ if $x \neq y$. Thus, $\forall x \forall y ((x \neq y) \rightarrow (f(x) \neq f(y)))$.

Function

- Function examples:
 - Example-6:: For function $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$ —
Considering $x = 1$ and $y = -1$, where $x \neq y$, $f(x) = 1 = f(y)$.
So, $\exists x \exists y \neg((x \neq y) \rightarrow (f(x) \neq f(y)))$, i.e., $\neg(\forall x \forall y((x \neq y) \rightarrow (f(x) \neq f(y))))$.
So, $f: \mathbb{Z} \rightarrow \mathbb{Z}$ **not one-to-one**.
 - Example-7:: For function $g: \mathbb{R} \rightarrow \mathbb{R}$, $g(x) = x^2$ **not one-to-one**, for same reason as above.
 - Example-8:: For functions $g_1: \mathbb{R}^+ \rightarrow \mathbb{R}$, $g_1(x) = x^2$, and $g_2: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, $g_2(x) = x^2$ both **one-to-one**, for same reason as given in Example-5.

Discrete Mathematics

Dept. of CSE, NITP
19

Dr. Suddhasil De

Function

- Function examples:
 - Example-9:: For function $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x + 1$, to show **one-to-one** (alternately) leading to show unique codomain-domain mapping —
For any $x, y \in \mathbb{R}$, and assuming $f(x) = f(y)$, resulting $x + 1 = y + 1$, i.e., $x = y$. So, $\forall x \forall y((f(x) = f(y)) \rightarrow (x = y))$.
So, $f: \mathbb{R} \rightarrow \mathbb{R}$ **one-to-one**.
 - Example-10:: For function $g: \mathbb{Z} \rightarrow \mathbb{Z}$, $g(x) = x + 1$, to show **onto**, thus to show entire codomain-domain mapping (i.e. codomain=range) —
For arbitrary $y \in \mathbb{Z}$, and assuming $g(x) = y$, resulting in $x + 1 = y$, i.e., $x = y - 1$. So, $x \in \mathbb{Z}$. So, $\forall y \exists x(f(x) = y)$. So, $g: \mathbb{Z} \rightarrow \mathbb{Z}$ **onto**.

Discrete Mathematics

Dept. of CSE, NITP
20

Dr. Suddhasil De

Function

- Function examples:
 - Example-11:: for function $g: \mathbb{Z} \rightarrow \mathbb{Z}$, $g(x) = x^2$ —
Considering $y = -1$, leading to $x^2 = -1$, resulting in $x \notin \mathbb{Z}$.
So, $\exists y \neg(\exists x(g(x) = y))$, i.e. $\neg(\forall y \exists x(g(x) = y))$.
So, $g: \mathbb{Z} \rightarrow \mathbb{Z}$ **not onto**.


Discrete Mathematics

Dept. of CSE, NITP
21

Dr. Suddhasil De

Function

- Function:
 - Property: **Identity function** on A , $\iota_A: A \rightarrow A$ — assignment " $\iota_A(a) = a$ ", for all $a \in A$, i.e., identity function on set assigning each element to itself.
 - Property: $\iota_A: A \rightarrow A$ of nature bijective (one-to-one correspondence), i.e. both one-to-one (injective) and onto (surjective).
 - Property: **Inverse function** of bijective function $f: A \rightarrow B$ — assignment " $f^{-1}(b) = a$ " only when $f(a) = b$, where $a \in A$, $b \in B$.



[Ref: Kenneth H. Rosen, Kamala Krithivasan, Discrete Mathematics and Its Applications, Eighth edition, McGraw-Hill Education, 2021.]

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

22

Function

- Function examples:
 - Example-12: For function $f: \mathbb{Z} \rightarrow \mathbb{Z}$, $f(x) = x + 1$, shown one-to-one and onto (in Example-9, 10 of previous lecture), to obtain **inverse** — For any $y \in \mathbb{Z}$, and considering $y = f(x) = x + 1$, resulting in $x = y - 1$, i.e. $f^{-1}(y) = x = y - 1$. So, $x \in \mathbb{Z}$. So, $f: \mathbb{Z} \rightarrow \mathbb{Z}$ **invertible**, where $f^{-1}(y) = y - 1$.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

23

Summary

- Focus: Multiset, function as discrete structures.
- Multiset, and its representation.
- Multiset inclusion, union and intersection, with examples.
- Cardinality of multiset.
- Multiset operations, with examples.
- Function, its representation and related terminologies.
- Injective, surjective, bijective functions, with examples.
- Real-valued, integer-valued functions.
- Increasing, decreasing functions.
- Identity function, inverse function.

Discrete Mathematics

Dept. of CSE, NITP

Dr. Suddhasil De

24

References

1. [Ros21] Kenneth H. Rosen, Kamala Krithivasan, *Discrete Mathematics and its Applications*, Eighth edition, McGraw-Hill Education, 2021.

2. [Ross12] Kenneth A. Ross, Charles R. B. Wright, *Discrete Mathematics*, Fifth edition, Pearson Education, 2012.

3. [Mot15] Joe L. Mott, Abraham Kandel, Theodore P. Baker, *Discrete Mathematics for Computer Scientists and Mathematicians*, Second edition, Pearson Education, 2015.

4. [Lip07] Seymour Lipschutz, Marc L. Lipson, *Schaum's Outline of Theory and Problems of Discrete Mathematics*, Third edition, McGraw-Hill Education, 2007.

Further Reading

- Multiset:: [Ros21]:143-144.
- Function:: [Ros21]:147-149.
- Injective, surjective, bijective functions:: [Ros21]:150-153.
- Real-valued, integer-valued functions:: [Ros21]:149-150.
- Increasing, decreasing functions:: [Ros21]:151.

Lecture Exercises: Problem 1 [Ref: Gate 2024, Set-1, Q.32, p.20]

Let **A** and **B** be non-empty finite sets such that there exist one-to-one and onto functions — (i) from **A** to **B**, and (ii) from $(A \times A)$ to $(A \cup B)$. The number of possible values of $|A|$ is _____.

Lecture Exercises: Problem 1 Ans

- Given: $|A| \neq 0$, $|B| \neq 0$, bijection $f: A \rightarrow B$, bijection $g: (A \times A) \rightarrow (A \cup B)$.
- For arbitrary $n \in \mathbb{Z}^+$, let " $|A| = n$ " be TRUE. Then, " $|B| = n$ " becomes TRUE, due to existence of bijection $f: A \rightarrow B$.
- Also, " $|A \times A| = |A| \cdot |A| = n^2$ " becomes TRUE. Then, " $|A \cup B| = n^{2n}$ " becomes TRUE, due to existence of bijection $g: (A \times A) \rightarrow (A \cup B)$.
- Following principle of inclusion-exclusion, $|A \cup B| = |A| + |B| - |A \cap B|$.
So, $|A \cup B| \leq |A| + |B| \equiv n^2 \leq 2 \cdot n \equiv n^2 - 2 \cdot n \leq 0 \equiv n \cdot (n - 2) \leq 0$.
- For the last simplified inequality to hold, value of n to be within closed interval $[0, 2]$. But, $n \neq 0$, as per premise.
- So, $n \in [1, 2]$. Only **two** possible values of $|A|$.

Discrete Mathematics

Dept. of CSE, NITP
28

Dr. Suddhasil De
