

Don

MATHEMATICS

BASED ON LATEST PATTERN



FREE
QBOOK

- ⌘ KEY POINTS & IMPORTANT FORMULAE
- ⌘ ALL TEXTUAL QUESTIONS SOLVED
- ⌘ ENORMOUS CREATIVE QUESTIONS
- ⌘ SOLUTIONS TO MCQs
- ⌘ GOVT. MODEL QUESTION PAPER



Don Publications (P) Ltd.

▼ WARNING: All rights reserved. No part of this publication may be reproduced or stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the Publisher, the copy-right holder.

PREFACE

Don Mathematics for 10th standard is released with much pride. This guide is prepared based on the Tamilnadu Government's latest new syllabus.

In this book, 'KEY POINTS' and 'IMPORTANT FORMULAE' are given for headings. All the textual questions are solved. Also enormous additional creative questions are solved. Latest Govt. model question paper is given at the end of this book.

All the Textual MCQs and additional creative MCQs are given with full solution. We firmly believe that this book will be of immense help to the students to score centum in their exam.

Wishing you all the best!

S.A. Suresh Kumar MCA, MBA
for **Don** Publications (P) Ltd

Authored by

Mr. SANKARAN BALASUBRAMANIAN M.Sc., M.Phil., BGL
Chennai.

Mrs. S. P. RAJINI M.Sc., B.Ed., (Maths) M.Com., M.Phil.,
Virugambakkam, Chennai.

Reviewed by

Mr. A. ANTONY XAVIER RAJ M.Sc., M.A., M.Ed., M.Phil., PGDCA.,
Maths Assistant (Rtd.), St. Xavier's Hr.Sec. School, Tirunelveli.

Compiled by

Dr. S.A. RAJKUMAR MBBS., MS (EM)
Tirunelveli.

Published by:



Don Publications (P) Ltd.

3/100/1, Uthamapandiankulam, Tuticorin Main Road, **TIRUNELVELI** - 627351.

☎ : (0462) 2520977, 2520978, Cell: 73730 88821, 94433 39941. E-Mail: donbooks02@gmail.com

New No. 17, Old No. 21 - A, 2nd Floor, 11th Avenue, Ashok Nagar, **CHEENNAI** - 600 083.

☎ : (044) 25892474, 48565226, Cell: 73730 88893, 77083 30666.

E-Mail: donbooks.chennai01@gmail.com

Contents

1	MATHEMATICAL SYMBOLS	6
1	RELATIONS AND FUNCTIONS	1
2	NUMBERS AND SEQUENCES	42
3	ALGEBRA	113
4	GEOMETRY	223
5	COORDINATE GEOMETRY	294
6	TRIGONOMETRY	348
7	MENSURATION	407
8	STATISTICS AND PROBABILITY	449
	GOVT.MODEL QUESTION PAPER 2020	508

TO ORDER BOOKS

Dear Teachers and Principals

Thank you for your patronage to **Don Book**. You are instrumental in our phenomenal success.
Now you can order your required copies by sending the same in your letter head.

For order please

Mobile No. : **7373088821, 0462 - 2520977, 2520978**

Whatsapp No. : **9443339941**

Website : www.donbookkart.com

E-mail ID : order@donbooks.co.in, enquire@donbooks.co.in

BANK DETAILS (FOR DIRECT DEPOSIT)

STATE BANK OF INDIA

A/C No. : **35655497778**

IFSC Code : SBIN0000932

(Sripuram Branch, Tirunelveli)

INDIAN BANK

A/C No. : **6204213630**

IFSC Code : IDIB000K184

(KTC Nagar Branch, Tirunelveli)

KARUR VYSYA BANK

A/C No. : **1174135000000133**

IFSC Code : KVBL0001174

(Palayamkottai Branch)

KARUR VYSYA BANK

A/C No. : **1651135000001796**

IFSC Code : KVBL0001651

(Saidapet Branch, Chennai)

SYNDICATE BANK

A/C No. : **60501010001381**

IFSC Code : SYNB0006050

(Ashok Nagar Branch, Chennai)

CITY UNION BANK

A/C No. : **230109000157474**

IFSC Code : CIUB000230

(Palayamkottai Branch, Tirunelveli)

D.D./Cheque

Should be drawn in favour of '**Don Publication (P) Ltd**' payable at Tirunelveli.

(Please send the chellan along with your order)

You can purchase our books through online by visiting the website www.donbookkart.com
Here you can purchase book in bulk quantities.

For students:

You can buy our books through our website www.donbookkart.com or sending money order
(by clearly writing your address)

• Address for correspondence: •



Don Publications (P) Ltd.

3/100/1, Uthamapandiankulam, Tuticorin Main Road, **TIRUNELVELI** - 627351.

☎ : (0462) 2520977, 2520978, Cell: 73730 88821, 94433 39941. E-Mail: donbooks02@gmail.com

New No. 17, Old No. 21-A, 2nd Floor, 11th Avenue, Ashok Nagar, **CHEENNAI** - 600 083.

☎ : (044) 24892474, 48565226. Cell: 73730 88893, 77083 30666, E-Mail: donbooks.chennai01@gmail.com

OUR AREA REPRESENTATIVES CONTACT NUMBERS

You can buy books at the following wholesale / retail stores:

Don

Chennai	94432 21264, 94890 51855	Krishnagiri	94426 08863
Chennai South	73587 26333, 94890 51855	Dharmapuri	94426 08863
Chennai Central	90033 59974, 94890 51855	Tuticorin	94890 51851
Tiruvallur	73587 26333	Trichy, Ariyalur, Perambalur	73587 61666, 99430 22335
Vellore	94899 06831	Vilupuram	94890 51854
Kanchipuram	94426 08864, 94890 51855	Thiruvarur, Thanjavur, Nagapattinam	94421 19859
Tirunelveli	94433 39971, 73730 88835	Thiruvannamalai :	94426 08864
Nagercoil	94433 39971, 97905 77993, 77082 23111	Virudhunagar, Ramnad, Sivagangai	75980 38097
Takkalai	73587 64222, 77082 23111	Madurai, Theni, Pudukkottai	75980 38096
Kuzhithurai	94443 39971, 73730 88835	Coimbatore	73958 32888
Erode	94426 08863	Tirupur, Nilgiris	73958 32888
Dindugul, Karur	94890 51852, 94426 08863	Puducherry, Cuddalore	94426 01362
Salem, Namakkal	94426 08863		

OUR BOOKS ARE AVAILABLE AT

CHENNAI		MARTHANDAM	
1. M.K.StoresChennai	044 - 25386955	33. Olympic Traders	04651 - 324533
2. Bismi Book Centre	044 - 25363375/25383264		
3. Kings Book House	044 - 25367660		
4. Limra Book Centre	044 - 42032081	34. Vignesh Book Centre	0427 - 4020407 / 4020409
5. F.A.Stores	044 - 25392240	35. AK Chandraiah Chettiar & Sons	0427 - 2410096
6. Prince Book House	044 - 42053926	36. Kalaivani Book Centre	0427 - 2411083
7. A.R.Book Mart	99415 16187		
8. M.R.Book Store	044 - 25364596 / 42080600		
9. Arraba Book Traders	044 - 25387868		
TAMBARAM (W)		DHARMAPURI	
10. Sri Ram & Co Book Shop	044 - 22266431	37. Vimal Book Centre	94434 13414
11. Kalaimagal Book Centre	99524 99827		
CHIDAMBARAM		TANJORE	
12. Vetri Book Centre	04144 - 222648 / 254359	38. Sri Murugan Publications	04362 - 272922
COVAI		39. Srinivasa Book Centre	04362 - 238985
13. Majestic Book House	0422 - 2382333 / 2384333		
14. Cheran Book House	0422 - 4355453/2396623		
15. Radhamani Stores	0422 - 2392122		
16. CBSC Book Shop	0422 - 2393093		
17. Kannan Puthaga Nilayam	0422 - 2398890, 4369890		
18. Murugan Book Depot	0422 - 2396531		
CUDDALORE		TENKASI	
19. Indian Book Centre	04142 - 231780	41. Maheshwari Book Centre	04633 - 224406
20. Bell Book House	94432 65414		
PANRUTTI		TIRUNELVELI	
21. Sri Vinayaga Stores	81241 01837	42. Eagle Book CentrePalai, Tvl	0462 - 2578899
DINDIGUL		43. Padma Book Stall	0462 - 2337650
22. Ayyanar Book Centre	0451 - 2426561	44. Shyamala Book Centre	0462 - 2338800
ERODE		45. Chitradevi Book Centre	0462 - 2334344
23. Selvam Book Centre	0424 - 2257027	46. Velavan Book Centre	0462 - 2574457
24. Dhana Book Company	0424 - 2221230	47. Ponnaiya Pillai Book Shop	0462 - 2572564
25. Kalaivani Book Centre	0424 - 3205711		
26. Senthil Book Palace	0424 - 2214886		
KALLAKURICHI		TIRUVANNAMALAI	
27. Sri Kiruba Stationary	04151 - 222114	48. APM Book Centre	04175 - 224725
KOVILPATTI			
28. Chitradevi Book Centre	04632 - 220733	TIRUPPUR	
MADURAI		49. Senthil Stores	0421 - 2247274
29. Mano Book Centre	0452 - 2621577	50. Chola Book House	0421 - 4323668
30. Jeyam Book Centre	0452 - 2623636	51. Maheshwari Book Stall	94420 04254, 0421 - 2236414
31. Sunmathi Traders	0452 - 2338966		
NAGERCOIL		TIRUCHIRAPPALLI	
32. M. ARUMUGAMPILLAI SONS BOOK CENTRE	96004 23549, 0452 - 230728	52. Sri Murugan Book Centre	0431 - 2703076
		53. Rasi Publications	0431 - 2703692
		54. Tamilnadu Book House	96592 99661
		55. Sri Raghavendra Stores	0431 - 2707506
TUTICORIN		TUTICORIN	
		56. Sri Durga Stores	0461 - 2326694
		57. Eagle Book Centre	0461 - 2392333
VILUPPURAM		VILUPPURAM	
		58. Sabarinathan Brothers	04146 - 222581
VIRUDHUNAGAR		59. Sethus Book Centre	04562 - 248400 / 94864 61400

MATHEMATICAL SYMBOLS


Don

SYMBOL	MEANING	SYMBOL	MEANING
+	Addition	$\sqrt{}$	Square root
-	Subtraction	$\sqrt[3]{}$	Cubic root
\times	Multiplication	\parallel	Parallel to
\div	Division	\perp	Perpendicular to
\pm	Plus or Minus	\angle	Angle
=	Equal to	\triangle	Triangle
\neq	Not equal to	\circ	Circle
\approx	Equivalent to	\square	Square
\simeq	Approximately equal to	\square	Rectangle
\equiv (or) \cong	Congruent	\therefore	Therefore
\equiv	Identically equal to	\because	Since (or) because
<	Less than	π	Pi
\leq	Less than or equal to	Σ	Summation
>	Greater than	A' (or) A^c	Complement of A
\geq	Greater than or equal to	\emptyset (or) {}	Empty set or null set or void set
$ \quad $	Absolute value	$n(A)$	Number of elements in the set A
\propto	Proportional to	$P(A)$	Power set of A
∞	Infinity	$\ \quad \ ^{ly}$	Similarly
\cup	Union	Δ	Symmetric difference
\cap	Intersection	\mathbb{N}	Set of Natural numbers
\mathbb{U}	Universal set	\mathbb{W}	Set of Whole numbers
\in	Belongs to	\mathbb{Z}	Set of all Integers
\notin	Does not belong to	\mathbb{R}	Set of Real numbers
\subset	Proper subset of	$ $ (or) :	Such that
\subseteq	Subset of or is contained in	x''	x seconds
$\not\subset$	Not a proper subset of	x'	x minutes
$\not\subseteq$	Not a subset of or is not contained in	x°	x degrees
\implies	Implies	\overline{AB}	Segment AB
\Leftrightarrow	Implies and is implied by	\overrightarrow{AB}	Ray AB
:	Ratio	\overleftrightarrow{AB}	Line AB
.	Decimal	x^n	x to the power n
%	Percent [out of 100]		

UNIT 1

RELATIONS AND FUNCTIONS

MIND MAP



CARTESIAN PRODUCT

Key Points

- ❖ The set of all ordered pairs (a, b) such that $a \in A$ and $b \in B$ is called the Cartesian product of non-empty sets A and B and is denoted by $A \times B$.
- ❖ For three non-empty sets A , B and C , $A \times B \times C$ is the set of all ordered triplets having first element from A , second element from B and third element from C .
- ❖ If A and B are two finite sets, then the number of elements in $A \times B$, i.e., $n(A \times B) = n(A) \times n(B)$.
- ❖ If either A or B is an infinite set, then $A \times B$ is an infinite set.
- ❖ If A , B and C are finite sets, then $n(A \times B \times C) = n(A) \times n(B) \times n(C)$.
- ❖ Two ordered pairs (a, b) and (c, d) are equal if and only if $a = c$, $b = d$.
- ❖ $A \times B \neq B \times A$ unless $A = B$.

Don

- ⇒ If one of A and B or both A and B are null sets then $A \times B = \emptyset$.
- ⇒ $A \times B$ and $B \times A$ are equivalent sets.
- ⇒ For any three sets A, B and C
 - (i) $A \times (B \cup C) = (A \times B) \cup (A \times C)$
 - (ii) $A \times (B \cap C) = (A \times B) \cap (A \times C)$
- ⇒ $A \times (B - C) = (A \times B) - (A \times C)$
- ⇒ If $A \subseteq B$, then $(A \times B) \cap (B \times A) = A^2$

Worked Examples

1.1 If $A = \{1, 3, 5\}$ and $B = \{2, 3\}$ then

- (i) find $A \times B$ and $B \times A$.
- (ii) Is $A \times B \neq B \times A$? If not why?
- (iii) Show that $n(A \times B) = n(B \times A) = n(A) \times n(B)$

Sol: Given that $A = \{1, 3, 5\}$ and $B = \{2, 3\}$

$$\begin{aligned} A \times B &= \{1, 3, 5\} \times \{2, 3\} \\ &= \{(1, 2), (1, 3), (3, 2), (3, 3), \\ &\quad (5, 2), (5, 3)\} \dots (1) \\ B \times A &= \{2, 3\} \times \{1, 3, 5\} \\ &= \{(2, 1), (2, 3), (2, 5), (3, 1), \\ &\quad (3, 3), (3, 5)\} \dots (2) \end{aligned}$$

From (1) and (2) we conclude that $A \times B \neq B \times A$ as $(1, 2) \neq (2, 1)$ and $(1, 3) \neq (3, 1)$, etc.

$$n(A) = 3; n(B) = 2$$

From (1) and (2) we observe that,

$$n(A \times B) = n(B \times A) = 6$$

we see that, $n(A) \times n(B) = 3 \times 2 = 6$ and

$$n(B) \times n(A) = 2 \times 3 = 6$$

Hence,

$$n(A \times B) = n(B \times A) = n(A) \times n(B) = 6$$

Thus, $n(A \times B) = n(B \times A) = n(A) \times n(B)$.

1.2 If $A \times B = \{(3, 2), (3, 4), (5, 2), (5, 4)\}$ then find A and B.

Sol: $A \times B = \{(3, 2), (3, 4), (5, 2), (5, 4)\}$

We have $A = \{\text{set of all first co-ordinates of elements of } A \times B\}$.

Therefore, $A = \{3, 5\}$

$B = \{\text{set of all second co-ordinates of elements of } A \times B\}$.

Therefore, $B = \{2, 4\}$

Thus $A = \{3, 5\}$ and $B = \{2, 4\}$.

1.3 Let $A = \{x \in \mathbb{N} \mid 1 < x < 4\}$,
 $B = \{x \in \mathbb{W} \mid 0 \leq x < 2\}$ and $C = \{x \in \mathbb{N} \mid x < 3\}$.
 Then verify that

$$(i) A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$(ii) A \times (B \cap C) = (A \times B) \cap (A \times C)$$

Sol: $A = \{x \in \mathbb{N} \mid 1 < x < 4\} = \{2, 3\}$

$$B = \{x \in \mathbb{W} \mid 0 \leq x < 2\} = \{0, 1\}$$

$$C = \{x \in \mathbb{N} \mid x < 3\} = \{1, 2\}$$

$$(i) A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$B \cup C = \{0, 1\} \cup \{1, 2\}$$

$$= \{0, 1, 2\}$$

$$A \times (B \cup C) = \{2, 3\} \times \{0, 1, 2\}$$

$$= \{(2, 0), (2, 1), (2, 2), (3, 0),$$

$$(3, 1), (3, 2)\} \dots (1)$$

$$A \times B = \{2, 3\} \times \{0, 1\}$$

$$= \{(2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$A \times C = \{2, 3\} \times \{1, 2\}$$

$$= \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cup (A \times C) = \{(2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$\cup \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cup (A \times C) = \{(2, 0), (2, 1), (2, 2), (3, 0),$$

$$(3, 1), (3, 2)\} \dots (2)$$

From (1) and (2),

$A \times (B \cup C) = (A \times B) \cup (A \times C)$ is verified.

$$(ii) A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$B \cap C = \{0, 1\} \cap \{1, 2\} = \{1\}$$

$$A \times (B \cap C) = \{2, 3\} \times \{1\} = \{(2, 1), (3, 1)\} \dots (1)$$

$$A \times B = \{2, 3\} \times \{0, 1\} = \{(2, 0),$$

$$(2, 1), (3, 0), (3, 1)\}$$

$$A \times C = \{2 \times 3\} \times \{1, 2\}$$

$$= \{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cap (A \times C) = \{(2, 0), (2, 1), (3, 0), (3, 1)\} \cap$$

$$\{(2, 1), (2, 2), (3, 1), (3, 2)\}$$

$$(A \times B) \cap (A \times C) = \{(2, 1), (3, 1)\} \dots (2)$$

From (1) and (2),

$A \times (B \cap C) = (A \times B) \cap (A \times C)$ is verified.

Unit - 1 | RELATIONS AND FUNCTIONS

Don

 Progress Check

1. For any two non-empty sets A and B, $A \times B$ is called as _____

Ans : Cartesian Product

2. If $n(A \times B) = 20$ and $n(A) = 5$ then $n(B)$ is _____

Ans : $n(B) = 4$

3. If $A = \{-1, 1\}$ and $B = \{-1, 1\}$ then geometrically describe the set of points of $A \times B$

Ans : $A \times B = \{(-1, -1), (-1, 1), (1, -1), (1, 1)\}$

4. If A, B are the line segments given by the intervals $(-4, 3)$ and $(-2, 3)$ respectively, represent the cartesian product of A and B.

Ans : $A \times B = \{(-4, -2), (-4, 3), (3, -2), (3, 3)\}$

 Thinking Corner

1. When will $A \times B$ be equal to $B \times A$?

Ans : $A \times B = B \times A$, When $A = B$.

2. Let $A = \{1, 2, 3\}$ and $B = \{x \mid x \text{ is a prime number less than } 10\}$. Find $A \times B$ and $B \times A$.

Sol : $A = \{1, 2, 3\}; B = \{2, 3, 5, 7\}$

$A \times B = \{(1, 2), (1, 3), (1, 5), (1, 7), (2, 2), (2, 3), (2, 5), (2, 7), (3, 2), (3, 3), (3, 5), (3, 7)\}$

$B \times A = \{(2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3), (5, 1), (5, 2), (5, 3), (7, 1), (7, 2), (7, 3)\}$

3. If $B \times A = \{(-2, 3), (-2, 4), (0, 3), (0, 4), (3, 3), (3, 4)\}$ find A and B.

Sol : From $B \times A$, All the first entries belong to the set B and all the second entries belong to A.

$$\therefore A = \{3, 4\}$$

$$B = \{-2, 0, 3\}$$

4. If $A = \{5, 6\}$, $B = \{4, 5, 6\}$, $C = \{5, 6, 7\}$, Show that $A \times A = (B \times B) \cap (C \times C)$.

Sol : $A = \{5, 6\}, B = \{4, 5, 6\}, C = \{5, 6, 7\}$

$$\text{LHS: } A \times A = \{5, 6\} \times \{5, 6\}$$

$$= \{(5, 5), (5, 6), (6, 5), (6, 6)\}$$

$$B \times B = \{4, 5, 6\} \times \{4, 5, 6\}$$

$$= \{(4, 4), (4, 5), (4, 6), (5, 4), (5, 5), (5, 6), (6, 4), (6, 5), (6, 6)\}$$

RHS:

$$C \times C = \{5, 6, 7\} \times \{5, 6, 7\}$$

$$= \{(5, 5), (5, 6), (5, 7), (6, 5), (6, 6), (6, 7), (7, 5), (7, 6), (7, 7)\}$$

$$(B \times B) \cap (C \times C) = \{(5, 5), (5, 6), (6, 5), (6, 6)\}$$

$$\therefore \text{LHS} = \text{RHS}$$

$$A \times A = (B \times B) \cap (C \times C)$$

Hence proved.

Exercise 1.1

1. Find $A \times B$, $A \times A$ and $B \times A$ if

$$(i) A = \{2, -2, 3\} \text{ and } B = \{1, -4\}$$

$$(ii) A = B = \{p, q\}$$

$$(iii) A = \{m, n\}; B = \emptyset$$

Sol :

$$(i) A = \{2, -2, 3\} \text{ and } B = \{1, -4\}$$

$$\begin{aligned} A \times B &= \{2, -2, 3\} \times \{1, -4\} \\ &= \{(2, 1), (2, -4), (-2, 1), \\ &\quad (-2, -4), (3, 1), (3, -4)\} \end{aligned}$$

$$A \times A = \{2, -2, 3\} \times \{2, -2, 3\}$$

$$\begin{aligned} &= \{(2, 2), (2, -2), (2, 3), (-2, 2), (-2, -2), \\ &\quad (-2, 3), (3, 2), (3, -2), (3, 3)\} \end{aligned}$$

$$B \times A = \{1, -4\} \times \{2, -2, 3\}$$

$$= \{(1, 2), (1, -2), (1, 3), \\ &\quad (-4, 2), (-4, -2), (-4, 3)\}$$

$$(ii) A = B = \{p, q\}$$

$$\begin{aligned} A \times B &= \{p, q\} \times \{p, q\} \\ &= \{(p, p), (p, q), (q, p), (q, q)\} \end{aligned}$$

$$A \times A = \{p, q\} \times \{p, q\}$$

$$= \{(p, p), (p, q), (q, p), (q, q)\}$$

$$B \times A = \{p, q\} \times \{p, q\}$$

$$= \{(p, p), (p, q), (q, p), (q, q)\}$$

$$\therefore A \times B = A \times A = B \times A$$

Since $A = B$

$$(iii) A = \{m, n\}, B = \emptyset$$

If either A or B are null sets, then $A \times B$ will also be an empty set.

$$\text{i.e., } A = \emptyset \text{ (or) } B = \emptyset$$

$$\text{then } A \times B = \emptyset, B \times A = \emptyset$$

$$\text{and } A \times A = \{m, n\} \times \{m, n\}$$

$$= \{(m, m), (m, n), (n, m), (n, n)\}$$

Don

5. Given $A = \{1, 2, 3\}$, $B = \{2, 3, 5\}$, $C = \{3, 4\}$

and $D = \{1, 3, 5\}$, check if

$$(A \cap C) \times (B \cap D) = (A \times B) \cap (C \times D) \text{ is true?}$$

Sol: $A = \{1, 2, 3\}$, $B = \{2, 3, 5\}$, $C = \{3, 4\}$, $D = \{1, 3, 5\}$

$$A \cap C = \{3\}$$

$$B \cap D = \{3, 5\}$$

$$(A \cap C) \times (B \cap D) = \{3\} \times \{3, 5\}$$

$$= \{(3, 3), (3, 5)\} \quad \dots (1)$$

$$\begin{aligned} A \times B &= \{(1, 2), (1, 3), (1, 5), (2, 2), \\ &\quad (2, 3), (2, 5), (3, 2), \\ &\quad (3, 3), (3, 5)\} \end{aligned}$$

$$\begin{aligned} C \times D &= \{(3, 1), (3, 3), (3, 5), (4, 1), \\ &\quad (4, 3), (4, 5)\} \end{aligned}$$

$$(A \times B) \cap (C \times D) = \{(3, 3), (3, 5)\} \quad \dots (2)$$

From (1) and (2), it is clear that

$$\therefore (A \cap C) \times (B \cap D) = (A \times B) \cap (C \times D)$$

Hence it is true.

6. Let $A = \{x \in \mathbb{W} \mid x < 2\}$, $B = \{x \in \mathbb{N} \mid 1 < x \leq 4\}$ and $C = \{3, 5\}$. Verify that

$$(i) A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$(ii) A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$(iii) (A \cup B) \times C = (A \times C) \cup (B \times C)$$

Sol: Given $A = \{x \in \mathbb{W} \mid x < 2\} \Rightarrow A = \{0, 1\}$

$$B = \{x \in \mathbb{N} \mid 1 < x \leq 4\} \Rightarrow B = \{2, 3, 4\}$$

$$C = \{3, 5\}$$

$$(i) A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$B \cup C = \{2, 3, 4, 5\}$$

$$A \times (B \cup C) = \{0, 1\} \times \{2, 3, 4, 5\}$$

$$= \{(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\} \dots (1)$$

$$A \times B = \{0, 1\} \times \{2, 3, 4\}$$

$$= \{(0, 2), (0, 3), (0, 4), (1, 2), (1, 3), (1, 4)\}$$

$$A \times C = \{0, 1\} \times \{3, 5\}$$

$$= \{(0, 3), (0, 5), (1, 3), (1, 5)\}$$

$$(A \times B) \cup (A \times C) = \{(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 3), (1, 4), (1, 5)\} \dots (2)$$

From (1) \times (2), it is clear that

$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$

Hence verified.

$$(ii) A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$A = \{0, 1\}, B = \{2, 3, 4\}, C = \{3, 5\}$$

$$B \cap C = \{3\}$$

$$A \times (B \cap C) = \{0, 1\} \times \{3\}$$

$$= \{(0, 3), (1, 3)\} \dots (1)$$

$$A \times B = \{(0, 2), (0, 3), (0, 4), (1, 2), (1, 3), (1, 4)\}$$

$$A \times C = \{(0, 3), (0, 5), (1, 3), (1, 5)\}$$

$$(A \times B) \cap (A \times C) = \{(0, 3), (1, 3)\} \dots (2)$$

From (1) and (2), it is clear that

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

Hence verified.

$$(iii) (A \cup B) \times C = (A \times C) \cup (B \times C)$$

$$A = \{0, 1\}, \quad B = \{2, 3, 4\}, \quad C = \{3, 5\}$$

$$(A \cup B) = \{0, 1, 2, 3, 4\}$$

$$(A \cup B) \times C = \{0, 1, 2, 3, 4\} \times \{3, 5\}$$

$$= \{(0, 3), (0, 5), (1, 3), (1, 5), (2, 3), (2, 5), (3, 3), (3, 5), (4, 3), (4, 5)\} \dots (1)$$

$$A \times C = \{0, 1\} \times \{3, 5\}$$

$$= \{(0, 3), (0, 5), (1, 3), (1, 5)\}$$

$$B \times C = \{2, 3, 4\} \times \{3, 5\}$$

$$= \{(2, 3), (2, 5), (3, 3), (3, 5), (4, 3), (4, 5)\}$$

$$(A \times C) \cup (B \times C) = \{(0, 3), (0, 5), (1, 3), (1, 5), (2, 3), (2, 5), (3, 3), (3, 5), (4, 3), (4, 5)\} \dots (2)$$

From (1) and (2), it is clear that

$$(A \cup B) \times C = (A \times C) \cup (B \times C)$$

Hence verified.

7. Let A is the set of all natural numbers less than 8, B is the set of all prime numbers less than 8, C is the set of even prime number. Verify that

$$(i) (A \cap B) \times C = (A \times C) \cap (B \times C)$$

$$(ii) A \times (B - C) = (A \times B) - (A \times C)$$

Sol: Given: 'A' is the set of all natural numbers less than 8.

$$A = \{1, 2, 3, 4, 5, 6, 7\}$$

'B' is the set of all prime numbers less than 8

$$B = \{2, 3, 5, 7\}$$

'C' is the set of all even prime number

$$C = \{2\}$$

(i) Verify

$$(A \cap B) \times C = (A \times C) \cap (B \times C)$$

$$A \cap B = \{2, 3, 5, 7\}$$

$$(A \cap B) \times C = \{2, 3, 5, 7\} \times \{2\}$$

$$= \{(2, 2), (3, 2), (5, 2), (7, 2)\} \dots (1)$$

$$A \times C = \{1, 2, 3, 4, 5, 6, 7\} \times \{2\}$$

$$= \{(1, 2), (2, 2), (3, 2), (4, 2), (5, 2), (6, 2), (7, 2)\}$$

$$B \times C = \{2, 3, 5, 7\} \times \{2\}$$

$$= \{(2, 2), (3, 2), (5, 2), (7, 2)\}$$

$$(A \times C) \cap (B \times C) = \{(2, 2), (3, 2), (5, 2), (7, 2)\} \dots (2)$$

From (1) and (2) it is clear that

$$(A \cap B) \times C = (A \times C) \cap (B \times C)$$

Hence verified.

(ii) Verify

$$A \times (B - C) = (A \times B) - (A \times C)$$

$$B - C = \{2, 3, 5, 7\} - \{2\} = \{3, 5, 7\}$$

Unit - 1 | RELATIONS AND FUNCTIONS

$$\begin{aligned}
 A \times (B - C) &= \{1, 2, 3, 4, 5, 6, 7\} \times \{3, 5, 7\} \\
 &= \{(1, 3), (1, 5), (1, 7), (2, 3), (2, 5), \\
 &\quad (2, 7), (3, 3), (3, 5), (3, 7), (4, 3), \\
 &\quad (4, 5), (4, 7), (5, 3), (5, 5), (5, 7), \\
 &\quad (6, 3), (6, 5), (6, 7), (7, 3), (7, 5), \\
 &\quad (7, 7)\} \quad \dots (1)
 \end{aligned}$$

$$\begin{aligned} A \times B &= \{1, 2, 3, 4, 5, 6, 7\} \times \{2, 3, 5, 7\} \\ &= \{(1, 2), (1, 3), (1, 5), (1, 7), (2, 2), \\ &\quad (2, 3), (2, 5), (2, 7), (3, 2), (3, 3), \\ &\quad (3, 5), (3, 7), (4, 2), (4, 3), (4, 5), \\ &\quad (4, 7), (5, 2), (5, 3), (5, 5), (5, 7), \\ &\quad (6, 2), (6, 3), (6, 5), (6, 7), (7, 2), \\ &\quad (7, 3), (7, 5), (7, 7)\} \end{aligned}$$

$$\begin{aligned} A \times C &= \{1, 2, 3, 4, 5, 6, 7\} \times \{2\} \\ &= \{(1,2), (2,2), (3,2), (4,2), (5,2), (6,2), \\ &\quad (7,2)\} \end{aligned}$$

$$(A \times B) - (A \times C) = \{(1, 3), (1, 5), (1, 7), (2, 3), (2, 5), \\ (2, 7), (3, 3), (3, 5), (3, 7), (4, 3), \\ (4, 5), (4, 7), (5, 3), (5, 5), (5, 7), \\ (6, 3), (6, 5), (6, 7), (7, 3), (7, 5), \\ (7, 7)\} \quad \dots (2)$$

From (1) and (2), it is clear that

$$A \times (B - C) = (A \times B) - (A \times C)$$

Hence verified.

RELATIONS

Key Points

Worked Examples

- 1.4** Let $A = \{3, 4, 7, 8\}$ and $B = \{1, 7, 10\}$. Which of the following sets are relations from A to B?

 - (i) $R_1 = \{(3, 7), (4, 7), (7, 10), (8, 1)\}$
 - (ii) $R_2 = \{(3, 1), (4, 12)\}$
 - (iii) $R_3 = \{(3, 7), (4, 10), (7, 7), (7, 8), (8, 11), (8, 7), (8, 10)\}$


Sol : $A \times B = \{(3, 1), (3, 7), (3, 10), (4, 1),$
 $(4, 7), (4, 10), (7, 1), (7, 7),$
 $(7, 10), (8, 1), (8, 7), (8, 10)\}$

- (i)** We note that, $\mathbb{R}_1 \subseteq A \times B$. Thus, \mathbb{R}_1 is a relation from A to B.

(ii) Here, $(4, 12) \in \mathbb{R}_2$, but $(4, 12) \notin A \times B$. So, \mathbb{R}_2 is not a relation from A to B.

(iii) Here, $(7, 8) \in \mathbb{R}_3$, but $(7, 8) \notin A \times B$. So, \mathbb{R}_3 is not a relation from A to B.

- 1.5** The arrow diagram shows a relationship between the sets P and Q. Write the relation in (i) Set builder form (ii) Roster form (iii) What is the domain and range of \mathbb{R} ?



Sol:

- (i) Set builder form of
 $\mathbb{R} = \{(x, y) \mid y = x - 2, x \in P, y \in Q\}$

(ii) Roster form $\mathbb{R} = \{(5, 3), (6, 4), (7, 5)\}$

(iii) Domain of $\mathbb{R} = \{5, 6, 7\}$;
 Range of $\mathbb{R} = \{3, 4, 5\}$

Don

 Progress Check

1. Which of the following are relations from A to B?

- (i) $\{(1, b), (1, c), (3, a), (4, b)\}$
- (ii) $\{(1, a), (b, 4), (c, 3)\}$
- (iii) $\{(1, a), (a, 1), (2, b), (b, 2)\}$

Ans : Let $A = \{1, 2, 3, 4\}$ and $B = \{a, b, c\}$

Relations from A to B

- (i) $\{(1, b), (1, c), (3, a), (4, b)\}$

2. Which of the following are relations from B to A?

- (i) $\{(c, a), (c, b), (c, 1)\}$
- (ii) $\{(c, 1), (c, 2), (c, 3), (c, 4)\}$
- (iii) $\{(a, 4), (b, 3), (c, 2)\}$

Ans : Let $A = \{1, 2, 3, 4\}$ and $B = \{a, b, c\}$

Relations from B to A

- (iii) $\{(a, 4), (b, 3), (c, 2)\}$

Exercise 1.2

1. Let $A = \{1, 2, 3, 7\}$ and $B = \{3, 0, -1, 7\}$, which of the following are relation from A to B?

- (i) $R_1 = \{(2, 1), (7, 1)\}$
- (ii) $R_2 = \{(-1, 1)\}$
- (iii) $R_3 = \{(2, -1), (7, 7), (1, 3)\}$
- (iv) $R_4 = \{(7, -1), (0, 3), (3, 3), (0, 7)\}$

Sol : $A = \{1, 2, 3, 7\}$, $B = \{3, 0, -1, 7\}$

$A \times B = \{(1, 3), (1, 0), (1, -1), (1, 7), (2, 3), (2, 0), (2, -1), (2, 7), (3, 3), (3, 0), (3, -1), (3, 7), (7, 3), (7, 0), (7, -1), (7, 7)\}$

- (i) $R_1 = \{(2, 1), (7, 1)\}$

Since $(2, 1)$ and $(7, 1)$ are not the elements of $A \times B$, R_1 is not a relation from A to B. Moreover $1 \notin B$.

- (ii) $R_2 = \{(-1, 1)\}$, $(-1, 1) \notin A \times B$, $\therefore R_2$ is not a relation from A to B.

But $(-1, 1) \in (B \times A)$ as $-1 \in B$ and $1 \in A$.

- (iii) $R_3 = \{(2, -1), (7, 7), (1, 3)\}$

It is clear that $R_3 \subseteq A \times B$

$\therefore R_3$ is a relation from A to B.

- (iv) $R_4 = \{(7, -1), (0, 3), (3, 3), (0, 7)\}$

In this $(0, 3)$ and $(0, 7) \in R_4$

But $(0, 3)$ and $(0, 7)$ are not the elements of $A \times B$. Hence R_4 is not a relation from A to B.

2. Let $A = \{1, 2, 3, 4, \dots, 45\}$ and R be the relation defined as "is square of" on A. Write R as a subset of $A \times A$. Also, find the domain and range of R .

Sol : $A = \{1, 2, 3, 4, \dots, 45\}$

Relation is "is square of" and $A \rightarrow A$ on A

$A \times A = \{(1, 1), (1, 2), (1, 3), (1, 4), \dots, (45, 45)\}$

The square of '1' is $1 \in A$ and $(1, 1) \in A \times A$

The square of 2 is $4 \in A$ and $(4, 2) \in A \times A$

The square of 3 is $9 \in A$ and $(9, 3) \in A \times A$

The square of 4 is $16 \in A$ and $(16, 4) \in A \times A$

The square of 5 is $25 \in A$ and $(25, 5) \in A \times A$

The square of 6 is $36 \in A$ and $(36, 6) \in A \times A$

The square of 7 is $49 \notin A$.

$R = \{(1, 1), (4, 2), (9, 3), (16, 4), (25, 5), (36, 6)\}$

Domain of R = $\{1, 4, 9, 16, 25, 36\}$

Range of R = $\{1, 2, 3, 4, 5, 6\}$

3. A Relation R is given by the set $\{(x, y) / y = x + 3, x \in \{0, 1, 2, 3, 4, 5\}\}$. Determine its domain and range.

Sol :

Given Set = $\{(x, y) / y = x + 3, x \in \{0, 1, 2, 3, 4, 5\}\}$

When $x = 0$, $y = 0 + 3 = 3$

When $x = 1$, $y = 1 + 3 = 4$

When $x = 2$, $y = 2 + 3 = 5$

When $x = 3$, $y = 3 + 3 = 6$

When $x = 4$, $y = 4 + 3 = 7$

When $x = 5$, $y = 5 + 3 = 8$

\therefore Relation $R = \{(0, 3), (1, 4), (2, 5), (3, 6), (4, 7), (5, 8)\}$

Domain = $\{0, 1, 2, 3, 4, 5\}$

Range = $\{3, 4, 5, 6, 7, 8\}$

4. Represent each of the given relations by

(a) an arrow diagram, (b) a graph and (c) a set in roster, wherever possible.

- (i) $\{(x, y) | x = 2y, x \in \{2, 3, 4, 5\}, y \in \{1, 2, 3, 4\}\}$

- (ii) $\{(x, y) | y = x + 3, x, y \text{ are natural numbers} < 10\}$

Sol :

- (i) Given Set-Builder form

$\{(x, y) / x = 2y, x \in \{2, 3, 4, 5\},$

$y \in \{1, 2, 3, 4\}\}$

$x = 2y$


When $y = 1$, $x = 2$ ($1 = 2 \in x$)

Unit - 1 | RELATIONS AND FUNCTIONS


Don

When $y = 2$, $x = 2$ ($2 = 4 \in x$)When $y = 3$, $x = 2$ ($3 = 6 \notin x$)When $y = 4$, $x = 2$ ($4 = 8 \notin x$) \therefore Relation $R = \{(2, 1), (4, 2)\}$

(a) Arrow diagram



(b) Graph


(c) Roster form $R = \{(2, 1), (4, 2)\}$

(ii) Given set


$$\{(x, y) / y = x + 3, x, y \text{ are natural numbers} < 10\}$$

When $x = 1$, $y = 1 + 3 = 4$ When $x = 2$, $y = 2 + 3 = 5$ When $x = 3$, $y = 3 + 3 = 6$ When $x = 4$, $y = 4 + 3 = 7$ When $x = 5$, $y = 5 + 3 = 8$ When $x = 6$, $y = 6 + 3 = 9$ When $x = 7$, $y = 7 + 3 = 10$ is not possibleSince x and y are less than 10. \therefore Relation $R = \{(1, 4), (2, 5), (3, 6), (4, 7), (5, 8), (6, 9)\}$

(a) Arrow diagram



(b) Graph



(c) Roster form


$$R = \{(1, 4), (2, 5), (3, 6), (4, 7), (5, 8), (6, 9)\}$$

5. A company has four categories of employees given by Assistants (A), Clerks (C), Managers (M) and an Executive Officer (E). The company provide ₹ 10,000, ₹ 25,000, ₹ 50,000 and ₹ 1,00,000 as salaries to the people who work in the categories A, C, M and E respectively. If A_1, A_2, A_3, A_4 and A_5 were Assistants; C_1, C_2, C_3, C_4 were Clerks; M_1, M_2, M_3 were managers and E_1, E_2 were Executive officers and if the relation R is defined by xRy , where x is the salary given to person y , express the relation R through an ordered pair and an arrow diagram.

Sol:

Ordered Pair : The Domain of the relation is about the salaries given to person.

Relation is $R = \{(10000, A_1), (10000, A_2), (10000, A_3), (10000, A_4), (10000, A_5), (25000, C_1), (25000, C_2), (25000, C_3), (25000, C_4), (50000, M_1), (50000, M_2), (50000, M_3), (100000, E_1), (100000, E_2)\}$

Relation R defined by $x R y$ ' x ' is the salary given to person ' y '.**Arrow diagram**

Don

FUNCTIONS


Key Points

- Let A and B be two non-empty sets, then a relation from A to B i.e., a subset of $A \times B$ is called a function from A to B, if
- for each $a \in A$ there exists $b \in B$ such that $(a, b) \in f$
 - $(a, b) \in f$ and $(a, c) \in f \Rightarrow b = c$
- A function 'f' from a set 'A' to set 'B' associates each element of set A to a unique element of set B.
- Let $f : A \rightarrow B$. Then, the set A is domain of 'f' and B is co-domain of 'f'. The set of all images of elements of A is known as the range of 'f' or image set of A under f and is denoted by $f(A)$.
- i.e., $f(A) = \{f(x) : x \in A\} = \text{Range of } f$
 $\therefore f(A) \subseteq B$
- Not every curve in the cartesian plane is the graph of a function.
- Vertical line test: A set of points in the cartesian plane is the graph of a function if and only if no vertical straight line intersects the curve more than once.

Worked Examples

- 1.6 Let $X = \{1, 2, 3, 4\}$ and $Y = \{2, 4, 6, 8, 10\}$ and $R = \{(1, 2), (2, 4), (3, 6), (4, 8)\}$. Show that R is a function and find its domain, co-domain and range?

Sol: Pictorial representation of R is given in the figure. From the diagram, we see that for each $x \in X$, there exists only one $y \in Y$. Thus all elements in X have only image in Y. Therefore R is a function.



$$\begin{aligned}\text{Domain } X &= \{1, 2, 3, 4\}; \\ \text{Co-domain } Y &= \{2, 4, 6, 8, 10\}; \\ \text{Range of } f &= \{2, 4, 6, 8\}.\end{aligned}$$

- 1.7 A relation 'f' is defined by $f(x) = x^2 - 2$ where, $x \in \{-2, -1, 0, 3\}$

- (i) List the elements of f (ii) Is f a function?

Sol: $f(x) = x^2 - 2$ where $x \in \{-2, -1, 0, 3\}$

(i) $f(-2) = (-2)^2 - 2 = 2$
 $f(-1) = (-1)^2 - 2 = -1$
 $f(0) = (0)^2 - 2 = -2$
 $f(3) = (3)^2 - 2 = 7$

Therefore, $f = \{(-2, 2), (-1, -1), (0, -2), (3, 7)\}$

- (ii) We note that each element in the domain of f has a unique image. Therefore f is a function.


- 1.8 If $X = \{-5, 1, 3, 4\}$ and $Y = \{a, b, c\}$, then which of the following relations are functions from X to Y?

- $R_1 = \{(-5, a), (1, a), (3, b)\}$
- $R_2 = \{(-5, b), (1, b), (3, a), (4, c)\}$
- $R_3 = \{(-5, a), (1, a), (3, b), (4, c), (1, b)\}$

Sol:

- (i) $R_1 = \{(-5, a), (1, a), (3, b)\}$


We may represent the relation R_1 in an arrow diagram.



R_1 is not a function as $4 \in X$ does not have an image in Y.


Unit - 1 | RELATIONS AND FUNCTIONS**Don**

- (ii) $R_2 = \{(-5, b), (1, b), (3, a), (4, c)\}$
Arrow diagram of R_2 is shown in Figure.



R_2 is a function as each element of X has an unique image in Y.

- (iii) $R_3 = \{(-5, a), (1, a), (3, b), (4, c), (1, b)\}$
Representing R_3 in an arrow diagram.



R_3 is not a function as $1 \in X$ has two images $a \in Y$ and $b \in Y$.

Note that the image of an element should always be unique.

- 1.9 Given $f(x) = 2x - x^2$, find

- (i) $f(1)$ (ii) $f(x+1)$ (iii) $f(x) + f(1)$

Sol :

- (i) Replacing x with 1, we get

$$f(1) = 2(1) - (1)^2 = 2 - 1 = 1$$

- (ii) Replacing x with $x+1$, we get

$$\begin{aligned} f(x+1) &= 2(x+1) - (x+1)^2 \\ &= 2x + 2 - (x^2 + 2x + 1) \\ &= -x^2 + 1 \end{aligned}$$

- (iii) $f(x) + f(1) = (2x - x^2) + 1 = -x^2 + 2x + 1$
[Note that $f(x) + f(1) \neq f(x+1)$.]

In general $f(a+b)$ is not equal to $f(a) + f(b)$]

Progress Check

1. Relations are subsets of ____; Functions are subsets of ____.

Ans : Cartesian product; Relations.

2. True or False: All the elements of a relation should have images.

Ans : False

3. True or False: All the elements of a function should have images.

Ans : True

4. True or False: If $R : A \rightarrow B$ is a relation then the domain of $R = A$

Ans : True

5. If $f: \mathbb{N} \rightarrow \mathbb{N}$ is defined as $f(x) = x^2$ the pre-image(s) of 1 and 2 are ____ and ____.

Ans : 1 and None

6. The difference between relation and function is ____.

Ans : Every function is a Relation, but the relation need not be a function.

7. Let A and B be two non-empty finite sets. Then which one among the following two collection is large?

- (i) The number of relations between A and B.
(ii) The number of functions between A and B.

Ans :

(i) The number of relations between A and B is large.

(ii) Number of relation is always greater than number of functions.

Thinking Corner

1. Is the relation representing the association between planets and their respective moons a function?

Ans : Yes.


Exercise 1.3

1. Let $f = \{(x, y) | x, y \in \mathbb{N} \text{ and } y = 2x\}$ be a relation on N. Find the domain, co-domain and range. Is this relation a function?

Sol :

$$f = \{(x, y) | x, y \in \mathbb{N} \text{ and } y = 2x\}$$

Given that $y = 2x$



'x' is always a natural number. Domain is the set of all first entries.

So, Domain-Set of natural numbers = N and y is always an even number as $y = 2x$

Don

Range = Set of even natural numbers
 Co-domain = Set of natural numbers = N
 Here, the first elements (x) are having unique images. So, this relation is a function.

- 2. Let $X = \{3, 4, 6, 8\}$. Determine whether the relation $R = \{(x, f(x)) \mid x \in X, f(x) = x^2 + 1\}$ is a function from X to N ?**

Sol : Given $X = \{3, 4, 6, 8\}$

$$\text{Relation } R = \{(x, f(x)) \mid x \in X, f(x) = x^2 + 1\}$$

$$\text{When } x = 3, f(3) = (3)^2 + 1 = 9 + 1 = 10 \in N$$

$$\text{When } x = 4, f(4) = (4)^2 + 1 = 16 + 1 = 17 \in N$$

$$\text{When } x = 6, f(6) = (6)^2 + 1 = 36 + 1 = 37 \in N$$

$$\text{When } x = 8, f(8) = (8)^2 + 1 = 64 + 1 = 65 \in N$$

$$R = \{(3, 10), (4, 17), (6, 37), (8, 65)\}$$

Since, all the elements of X are having natural numbers as images, it is a function from X to N.

- 3. Given the function $f : x \rightarrow x^2 - 5x + 6$, evaluate**

$$(i) f(-1)$$

$$(ii) f(2a)$$

$$(iii) f(2)$$

$$(iv) f(x-1)$$

Sol : $f(x) = x^2 - 5x + 6$

$$(i) f(-1) = (-1)^2 - 5(-1) + 6 = 1 + 5 + 6 = 12$$

$$(ii) f(2a) = (2a)^2 - 5(2a) + 6 = 4a^2 - 10a + 6$$


$$(iii) f(2) = (2)^2 - 5(2) + 6 = 4 - 10 + 6 = 0$$

$$(iv) f(x-1) = (x-1)^2 - 5(x-1) + 6$$

$$= x^2 - 2x + 1 - 5x + 5 + 6$$

$$= x^2 - 7x + 12$$

- 4. A graph representing the function $f(x)$ is given below figure. From figure it is clear that $f(9) = 2$.**



- (i) Find the following values of the function**

$$(a) f(0)$$

$$(b) f(7)$$

$$(c) f(2)$$

$$(d) f(10)$$

- (ii) For what value of x is $f(x) = 1$?**

- (iii) Describe the following (a) Domain
 (b) Range.**

- (iv) What is the image of 6 under f?**

Sol :

$$(i) (a) f(0) = 9$$

$$(b) f(7) = 6$$

$$(c) f(2) = 6$$

$$(d) f(10) = 0$$

- (ii) For what value of x is $f(x) = 1$?**

From the graph, it is known that when $x = 9.5$, $f(9.5) = 1$

- (iii) (a) Domain = $\{x \mid 0 \leq x \leq 10, x \in \mathbb{R}\}$**

- (b) Range = $\{x \mid 0 \leq x \leq 9, x \in \mathbb{R}\}$**

- (iv) Image of '6' under f is '5'.**

- 5. Let $f(x) = 2x + 5$. If $x \neq 0$ then find $\frac{f(x+2) - f(2)}{x}$.**

Sol :

$$f(x) = 2x + 5, x \neq 0$$

$$\frac{f(x+2) - f(2)}{x} = \frac{[2(x+2)+5] - [2(2)+5]}{x}$$

$$= \frac{2x+4+5-9}{x} = \frac{2x+9-9}{x}$$

$$= \frac{2x}{x} = 2$$

- 6. A function f is defined by $f(x) = 2x - 3$**

$$(i) \text{ find } \frac{f(0) + f(1)}{2}$$

$$(ii) \text{ find } x \text{ such that } f(x) = 0$$

$$(iii) \text{ find } x \text{ such that } f(x) = x.$$

$$(iv) \text{ find } x \text{ such that } f(x) = f(1-x).$$

Sol : $f(x) = 2x - 3$

$$(i) \frac{f(0) + f(1)}{2} = \frac{[2(0)-3] + [2(1)-3]}{2}$$

$$= \frac{0-3+2-3}{2} = \frac{2-6}{2}$$

$$= \frac{-4}{2} = -2$$

$$(ii) \text{ Given } f(x) = 0$$

$$\therefore 2x - 3 = 0$$

$$2x = 3 \Rightarrow x = 3/2$$

$$(iii) \text{ Given } f(x) = x$$

$$2x - 3 = x$$

$$2x - x = 3 \Rightarrow x = 3$$

$$(iv) \text{ Given } f(x) = f(1-x)$$

$$2x - 3 = 2(1-x) - 3$$

$$2x - 3 = 2 - 2x - 3$$

$$2x + 2x = 2 - 3 + 3$$



$$4x = 2$$

$$x = 2/4 = 1/2$$

Unit - 1 | RELATIONS AND FUNCTIONS

Don

7. An open box is to be made from a square piece of material, 24 cm on a side, by cutting equal squares from the corners and turning up the sides as shown figure. Express the volume V of the box as a function of x.



Sol: From the diagram, the solid is a cuboid.
 Volume of cuboid = length × breadth × height
 where $l = 24 - 2x$, $b = 24 - 2x$, $h = x$
 \therefore Volume $V(x) = (24 - 2x)(24 - 2x)x$
 $V(x) = x(24 - 2x)^2, x > 0$
 $= 4x^3 - 96x^2 + 576x, x > 0$

So, the domain is $0 < x < 12$

8. A function f is defined by $f(x) = 3 - 2x$. Find x such that $f(x^2) = (f(x))^2$.

Sol: $f(x) = 3 - 2x$
 Given $f(x^2) = [f(x)]^2$
 $3 - 2x^2 = (3 - 2x)^2$
 $3 - 2x^2 = 9 - 12x + 4x^2$
 $6x^2 - 12x + 6 = 0$
 $6(x^2 - 2x + 1) = 0$
 $(x - 1)^2 = 0 \Rightarrow x = 1.$

9. A plane is flying at a speed of 500 km per hour. Express the distance d travelled by the plane as function of time t in hours.

Sol: Let the distance be 'd'
 Speed = 500 km/hr,
 Time = 't' hours
 \therefore Distance = Speed × time
 $d(t) = 500t$

10. The data in the adjacent table depicts the length of a woman's forehand and her corresponding height. Based on this data, a student finds a relationship between the height (y) and the forehand length (x) as $y = ax + b$, where a, b are constants.

Length x of forehand (in cm)	Height y (in inch)
35	56
45	65
50	69.5
55	74

- (i) Check if this relation is a function.
- (ii) Find a and b.
- (iii) Find the height of a woman whose forehand length is 40 cm.
- (iv) Find the length of forehand of a woman if her height is 53.3 inches.

Sol: $y = ax + b$; x = forehand length; y = height

x	y
35	56
45	65
50	69.5
55	74

For all the x-values, there is an image which is 'y'. Moreover, the difference between two consecutive 'y' values is constant.

\therefore In $y = ax + b$,

- (i) the Relation

$R = \{(35, 56), (45, 65), (50, 69.5), (55, 74)\}$ is a function.

- (ii) In $y = ax + b$

when $x = 35, y = 56$
 $\Rightarrow 56 = 35a + b$ (1)

when $x = 45, y = 65$
 $\Rightarrow 65 = 45a + b$ (2)

Solving (1) and (2), we get $a = 0.90$ and $b = 24.5$

- (iii) Given, forehand length is 40 cm

i.e., when $x = 40, y = ax + b$

So, $y = (0.90)(40) + 24.5 = 60.5$

\therefore Height of woman is 60.5 inches

- (iv) Given height is 53.3 inches

ie. when $y = 53.3, x = ?$

$$\Rightarrow 53.3 = 0.9x + 24.5$$

$$53.3 - 24.5 = 0.9x$$

$$\Rightarrow x = \frac{28.8}{0.9} = 32$$

length of forehand is 32 cm.

Don**REPRESENTATION OF FUNCTIONS****Worked Examples**

- 1. 10** Using vertical line test, determine which of the following curves (figure a, b, c, d) represent a function?




Figure (a)




Figure (b)




Figure (c)




Figure (d)

Sol :

The curves in figure (a) and (c) do not represent a function as the vertical lines meet the curves in two points P and Q.

The curves in figure (b) and (d) represent a function as the vertical lines meet the curve in at most one point.

- 1. 11** Let $A = \{1, 2, 3, 4\}$ and $B = \{2, 5, 8, 11, 14\}$ be two sets. Let $f : A \rightarrow B$ be a function given by $f(x) = 3x - 1$. Represent this function as


- (i) by arrow diagram
- (ii) in a table form
- (iii) as a set of ordered pairs
- (iv) in a graphical form

Sol :

$$\begin{aligned} A &= \{1, 2, 3, 4\}; B = \{2, 5, 8, 11, 14\}; \\ f(x) &= 3x - 1 \\ f(1) &= 3(1) - 1 = 3 - 1 = 2 \\ f(2) &= 3(2) - 1 = 6 - 1 = 5 \\ f(3) &= 3(3) - 1 = 9 - 1 = 8 \\ f(4) &= 3(4) - 1 = 12 - 1 = 11 \end{aligned}$$

(i) Arrow diagram

Let us represent the function $f: A \rightarrow B$ by an arrow diagram

**(ii) Table form**

The given function f can be represented in a tabular form as given below

x	1	2	3	4
$f(x)$	2	5	8	11

(iii) Set of ordered pairs


The function f can be represented as a set of ordered pairs as

$$f = \{(1, 2), (2, 5), (3, 8), (4, 11)\}$$


(iv) Graphical form

In the adjacent xy -plane the points


$(1, 2), (2, 5), (3, 8), (4, 11)$ are plotted.




- 1.12.** Using horizontal line test fig (a), fig (b), fig (c), determine which of the following functions are one-one.



(a)



(b)



(c)


Sol : The curves in fig (a) and fig (c), represent a one-one function as the horizontal lines meet the curves in only one point P.

The curve in fig (b) does not represent a one-one function, since, the horizontal line meet the curve in two points P and Q.

- 1.13.** Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and $f = \{(1, 4), (2, 5), (3, 6)\}$ be a function from A to B. Show that f is one-one but not onto function.

Sol : $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$;
 $f = \{(1, 4), (2, 5), (3, 6)\}$

Then f is a function from A to B and for different elements in A, there are different images in B. Hence f is one-one function. Note that the element 7 in the co-domain does not have any pre-image in the domain. Hence f is not onto.



Therefore f is one-one but not an onto function.

Don

- 1.14.** If $A = \{-2, -1, 0, 1, 2\}$ and $f: A \rightarrow B$ is an onto function defined by $f(x) = x^2 + x + 1$ then find B.

Sol : Given $A = \{-2, -1, 0, 1, 2\}$ and

$$\begin{aligned}f(x) &= x^2 + x + 1 \\f(-2) &= (-2)^2 + (-2) + 1 = 3 \\f(-1) &= (-1)^2 + (-1) + 1 = 1 \\f(0) &= 0^2 + 0 + 1 = 1 \\f(1) &= 1^2 + 1 + 1 = 3 \\f(2) &= 2^2 + 2 + 1 = 7\end{aligned}$$

Since, f is an onto function, range of $f = B$

Co-domain.

Therefore, $B = \{1, 3, 7\}$.

- 1.15.** Let f be a function $f: \mathbb{N} \rightarrow \mathbb{N}$ defined by $f(x) = 3x + 2, x \in \mathbb{N}$

- (i) Find the images of 1, 2, 3
- (ii) Find the pre-images of 29, 53
- (iii) Identify the type of function

Sol : The function $f: \mathbb{N} \rightarrow \mathbb{N}$ defined by $f(x) = 3x + 2$

- (i) If $x = 1, f(1) = 3(1) + 2 = 5$
If $x = 2, f(2) = 3(2) + 2 = 8$
If $x = 3, f(3) = 3(3) + 2 = 11$
The images of 1, 2, 3 are 5, 8, 11 respectively.

- (ii) If x is the pre-image of 29, then $f(x) = 29$.
Hence $3x + 2 = 29 \Rightarrow 3x = 27 \Rightarrow x = 9$.
Similarly, if x is the pre-image of 53, then $f(x) = 53$. Hence $3x + 2 = 53$
 $3x = 51 \Rightarrow x = 17$.
Thus the pre-images of 29 and 53 are 9 and 17 respectively.

- (iii) Since different elements of \mathbb{N} have different images in the co-domain, the function f is one-one function.
The co-domain of f is \mathbb{N} .
But the range of f = {5, 8, 11, 14, 17, ...} is a subset of \mathbb{N} .
Therefore f is not an onto function. That is, f is an into function.
Thus f is one-one and into function.

- 1.16.** Forensic scientists can determine the height (in cms) of a person based on the length of their thigh bone. They usually do so using the function $h(b) = 2.47b + 54.10$ where b is the length of the thigh bone.

- (i) Check if the function h is one-one
- (ii) Also find the height of a person if the length of his thigh bone is 50 cms.
- (iii) Find the length of the thigh bone if the height of a person is 147.96 cms.

Sol :

- (i) To check if h is one-one, we assume that $h(b_1) = h(b_2)$.

Then we get,

$$\begin{aligned}2.47b_1 + 54.10 &= 2.47b_2 + 54.10 \\2.47b_1 &= 2.47b_2 \Rightarrow b_1 = b_2\end{aligned}$$

Thus, $h(b_1) = h(b_2) \Rightarrow b_1 = b_2$.
So, the function h is one-one.

- (ii) If the length of the thigh bone b = 50, then the height is
 $h(50) = (2.47 \times 50) + 54.10 = 177.6$ cms.
- (iii) If the height of a person is 147.96 cms, then $h(b) = 147.96$ and so the length of the thigh bone is given by $2.47b + 54.10 = 147.96$.

$$b = \frac{93.86}{2.47} = 38$$

Therefore, the length of the thigh bone is 38 cms.

- 1.17.** Let f be a function from \mathbb{R} to \mathbb{R} defined by $f(x) = 3x - 5$. Find the values of a and b given that (a, 4) and (1, b) belong to f.

Sol : $f(x) = 3x - 5$ can be written as

$$f = \{(x, 3x - 5) | x \in \mathbb{R}\}$$

(a, 4) means the image of a is 4.

That is, $f(a) = 4$

$$\begin{aligned}3a - 5 &= 4 \\ \Rightarrow a &= 3\end{aligned}$$

(1, b) means the image of 1 is b.

That is, $f(1) = b$

$$3(1) - 5 = b \Rightarrow b = -2$$

- 1.18.** The distance S (in kms) travelled by a particle in time 't' hours is given by $S(t) = \frac{t^2 + t}{2}$. Find the distance travelled by the particle after

- (i) three and half hours
- (ii) eight hours and fifteen minutes.

Sol : The distance travelled by the particle is given by

$$S(t) = \frac{t^2 + t}{2}$$

- (i) $t = 3.5$ hours. Therefore,

Unit - 1 | RELATIONS AND FUNCTIONS

$$\begin{aligned} S(3.5) &= \frac{(3.5)^2 + 3.5}{2} \\ &= \frac{15.75}{2} = 7.875 \end{aligned}$$

The distance travelled in 3.5 hours is 7.875 kms.

(ii) $t = 8.25$ hours. Therefore,

$$\begin{aligned} S(8.25) &= \frac{(8.25)^2 + 8.25}{2} \\ &= \frac{76.3125}{3} \\ &= 38.15625 \end{aligned}$$

The distance travelled in 8.25 hours is 38.16 kms, approximately.

- 1.19.** If the function $f: R \rightarrow R$ defined by

$$f(x) = \begin{cases} 2x+7, & x < -2 \\ x^2-2, & -2 \leq x < 3 \\ 3x-2, & x \geq 3 \end{cases}$$


(i) $f(4)$

(ii) $f(-2)$

(iii) $f(4) + 2f(1)$

(iv) $\frac{f(1)-3f(4)}{f(-3)}$

Sol: The function f is defined by three values in intervals I, II, III as shown below



For a given value of $x = a$, find out the interval at which the point a is located, there after find $f(a)$ using the particular value defined in that interval.

(i) First, we see that $x = 4$ lie in the third interval. Therefore,

$$f(x) = 3x - 2; f(4) = 3(4) - 2 = 10$$

(ii) $x = -2$ lies in the second interval.

Therefore,

$$f(x) = x^2 - 2; f(-2) = (-2)^2 - 2 = 2$$

(iii) From (i), $f(4) = 10$.

To find $f(1)$, first we see that $x = 1$ lies in the second interval.

Therefore,

$$f(x) = x^2 - 2 \Rightarrow f(1) = 1^2 - 2 = -1$$

$$\text{Therefore, } f(4) + 2f(1) = 10 + 2(-1) = 8$$

(iv) We know that $f(1) = -1$ and $f(4) = 10$.

For finding $f(-3)$, we see that $x = -3$, lies in the first interval.

Therefore, $f(x) = 2x + 7$; thus,
 $f(-3) = 2(-3) + 7 = 1$
Hence, $\frac{f(1)-3f(4)}{f(-3)} = \frac{-1-3(10)}{1} = -31$

Progress Check

State True (or) False

1. All one - one functions are onto functions.

Ans : False

2. There will be no one - one function from A to B when $n(A) = 4, n(B) = 3$.

Ans : True

3. All onto functions are one - one functions.

Ans : False

4. There will be no onto function from A to B when $n(A) = 4, n(B) = 5$.

Ans : True

5. If f is a bijection from A to B, then $n(A) = n(B)$.

Ans : True

6. If $n(A) = n(B)$, then f is a bijection from A to B.

Ans : False

7. All constant functions are bijections.

Ans : False

Thinking Corner

1. Can there be a one to many function?


Ans : There cannot be a one to many function as the elements in Co-domain should have only one pre-image in the domain.

2. Is an identity function one-one function?


Ans : Yes. It is one-to-one function.

Exercise 1.4

1. Determine whether the graph given below represent functions. Give reason for your answers concerning each graph.



Don

**Sol :**

- (i) It is not a function. Since, a vertical line intersects the curve in two points.
- (ii) It is a function. Any vertical line drawn, will intersect the curve at only one point.
- (iii) It is not a function. Vertical line intersecting the curve at two points.
- (iv) It is a function. Vertical line intersects the curve at only one point.

2. Let $f : A \rightarrow B$ be a function defined by

$$f(x) = \frac{x}{2} - 1, \text{ where } A = \{2, 4, 6, 10, 12\}, \\ B = \{0, 1, 2, 4, 5, 9\}. \text{ Represent } f \text{ by}$$

- (i) set of ordered pairs; (ii) a table;
- (iii) an arrow diagram; (iv) a graph


Sol : $f : A \rightarrow B, f(x) = \frac{x}{2} - 1$

$A = \{2, 4, 6, 10, 12\}, B = \{0, 1, 2, 4, 5, 9\}$


- (i) Set of ordered pairs
 $= \{(2, 0), (4, 1), (6, 2), (10, 4), (12, 5)\}$
- (ii) Table

x	2	4	6	10	12
$f(x)$	0	1	2	4	5

(iii) Arrow diagram




(iv) Graph

**3. Represent the function $f = \{(1, 2), (2, 2), (3, 2), (4, 3), (5, 4)\}$ through**

- (i) an arrow diagram
- (ii) a table form (iii) a graph

Sol : Given function $f = \{(1, 2), (2, 2), (3, 2), (4, 3), (5, 4)\}$


(i) Arrow diagram



(ii) Table form

x	1	2	3	4	5
$f(x)$	2	2	2	3	4

(iii) Graph

**4. Show that the function $f : \mathbb{N} \rightarrow \mathbb{N}$ be defined by**

$$f(x) = 2x - 1 \text{ is one - one but not onto.}$$

Sol : Given function $f : \mathbb{N} \rightarrow \mathbb{N}$

$$f(x) = 2x - 1$$

This function maps every element from the domain to element that is twice minus one the original. $2x - 1$ is always an odd number when $x \in \mathbb{N}$.

Clearly, each element from the domain is mapped to different element in the co-domain. So, the function is one-to-one. On the other hand, there are no elements in the domain that would map to even numbers. So, the function is not onto.

Don

10. A function $f : [-5, 9] \rightarrow \mathbb{R}$ is defined as follows:

$$f(x) = \begin{cases} 6x + 1 & \text{if } -5 \leq x < 2 \\ 5x^2 - 1 & \text{if } 2 \leq x < 6 \\ 3x - 4 & \text{if } 6 \leq x \leq 9 \end{cases}$$

Find (i) $f(-3) + f(2)$ (ii) $f(7) - f(1)$

$$\text{(iii) } 2f(4) + f(8) \quad \text{(iv) } \frac{2f(-2) - f(6)}{f(4) + f(-2)}$$

Sol :

$$f : [-5, 9] \rightarrow \mathbb{R}$$

$$f(x) = \begin{cases} 6x + 1 & \text{if } -5 \leq x < 2 \\ 5x^2 - 1 & \text{if } 2 \leq x < 6 \\ 3x - 4 & \text{if } 6 \leq x \leq 9 \end{cases}$$

$$\text{(i) } f(-3) + f(2) = [6(-3) + 1] + [5(2)^2 - 1] \\ = (-18 + 1) + (20 - 1) \\ = -17 + 19 = 2 \quad [\because -5 \leq -3 < 2 \\ 2 \leq 2 < 6]$$

$$\text{(ii) } f(7) - f(1) = [3(7) - 4] - [6(1) + 1] \\ = (21 - 4) - (6 + 1) \\ = 17 - 7 = 10 \quad [\because 6 \leq 7 \leq 9 \\ -5 \leq 1 < 2]$$

$$\text{(iii) } 2f(4) + f(8) = 2[5(4)^2 - 1] + [3(8) - 4] \\ = 2[80 - 1] + [24 - 4] \\ = 158 + 20 = 178 \quad [\because 2 \leq 4 < 6]$$

$$\text{(iv) } \frac{2f(-2) - f(6)}{f(4) + f(-2)} \quad 6 \leq 8 \leq 9$$

$$f(-2) = 6(-2) + 1 = -12 + 1 = -11 \quad [\because -5 \leq -2 < 2]$$

$$f(6) = 3(6) - 4 = 18 - 4 = 14 \quad [\because 6 \leq 6 < 9]$$

$$f(4) = 5(4)^2 - 1 = 80 - 1 = 79 \quad [\because 2 \leq 4 < 6]$$

$$f(-2) = -11$$

$$\therefore \frac{2f(-2) - f(6)}{f(4) + f(-2)} = \frac{2(-11) - 14}{79 - 11} \\ = \frac{-22 - 14}{68} = -\frac{36}{68} = -\frac{9}{17}$$

11. The distance S an object travels under the influence of gravity in time t seconds is given by

$$S(t) = \frac{1}{2}gt^2 + at + b, \text{ where, (}g\text{ is the acceleration}$$

due to gravity), a, b are constants. Check if the function $S(t)$ is one-one.

Sol : Distance travelled by an object is given to

$$\text{be } S(t) = \frac{1}{2}gt^2 + at + b$$

' g ' - acceleration due to gravity is a constant.

' g ' is the acceleration due to gravity 'a' and 'b' are constants.

' t ' is a variable. (t - time)

At different values of ' t ', $S(t)$ is having different values. (images in codomain) clearly $f : t \rightarrow S(t)$ is one-to-one function.

Given $S(t) = \frac{1}{2}gt^2 + at + b$ (a, b are constants)


t = time in seconds

Let us take $t = 1, 2, 3, \dots$ Seconds

$$\text{When } t = 1 \Rightarrow S(1) = \frac{g}{2} + a + b$$

$$\text{When } t = 2 \Rightarrow S(2) = 2g + 2a + b$$

$$\text{When } t = 3 \Rightarrow S(3) = \frac{9}{2}g + 3a + b \text{ and so on.}$$



All the elements of t , having different images in $S(t)$. Hence it is an One-to-one function.

12. The function 't' which maps temperature in Celsius (C) into temperature in Fahrenheit (F) is defined by $t(C) = F$ where $F = \frac{9C}{5} + 32$. Find,

$$\text{(i) } t(0) \quad \text{(ii) } t(28)$$

$$\text{(iii) } t(-10)$$

$$\text{(iv) the value of C when } t(C) = 212$$

$$\text{(v) the temperature when the Celsius value is equal to the Fahrenheit value.}$$

Sol :

$$\text{Given } t(C) = F \text{ where } F = \frac{9C}{5} + 32.$$

C - Celsius, F - Fahrenheit

$$\therefore t(C) = \frac{9C}{5} + 32$$

$$\text{(i) } t(0) = \frac{9(0)}{5} + 32 = 0 + 32 = 32^\circ F$$

$$\text{(ii) } t(28) = \frac{9(28)}{5} + 32 = \frac{252}{5} + 32 \\ = 50.4 + 32 = 82.4^\circ F$$

$$\text{(iii) } t(-10) = \frac{9(-10)}{5} + 32 = -18 + 32 = 14^\circ F$$

Unit - 1 | RELATIONS AND FUNCTIONS**Don**

- (iv) Given $t(C) = 212$

$$\therefore \frac{9C}{5} + 32 = 212 \Rightarrow \frac{9C}{5} = 212 - 32 \\ C = 180 \times \frac{5}{9} = 100^{\circ}\text{C}$$

- (v) The temperature when the Celsius value is equal to the Fahrenheit value.

$$\therefore F = C$$

$$\begin{aligned} \frac{9C}{5} + 32 &= C \\ \frac{9C}{5} - C &= -32 \Rightarrow \frac{9C - 5C}{5} = -32 \\ 4C &= -32 \times 5 \Rightarrow C = -\frac{160}{4} \\ {}^{\circ}\text{C} &= -40 \end{aligned}$$

COMPOSITION OF FUNCTIONS**Key Points**

- ↗ Let $f : X \rightarrow R$ and $g : X \rightarrow R$ be any two real functions, where $X \subset R$, then their sum $f + g$ i.e., $(f + g) : X \rightarrow R$ is the function defined by $(f + g)(x) = f(x) + g(x)$, for all $x \in X$.
- ↗ Their difference i.e., $(f - g) : X \rightarrow R$ is the function defined by $(f - g)(x) = f(x) - g(x)$, for all $x \in X$.
- ↗ The multiplication (αf) is a function from $X \rightarrow R$ defined by a scalar by $(\alpha f)(x) = \alpha f(x)$, $x \in X$.
- ↗ The product $fg : X \rightarrow R$ is defined by $(fg)(x) = f(x).g(x)$, for all $x \in X$.
- ↗ The quotient $\frac{f}{g}$ is a function defined by $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$, provided $g(x) \neq 0$, $x \in X$.
- ↗ The composition of two functions f and g is denoted by $f \circ g$ and $(f \circ g)(x) = f[g(x)]$.
- ↗ Composition of functions is not always commutative i.e., $f \circ g \neq g \circ f$.
- ↗ Composition of functions is always associative.

Worked Examples

- 1. 20.** Find $f \circ g$ and $g \circ f$ when $f(x) = 2x + 1$ and $g(x) = x^2 - 2$.

Sol : $f(x) = 2x + 1, g(x) = x^2 - 2$

$$(f \circ g)(x) = f(g(x)) = f(x^2 - 2) \\ = 2(x^2 - 2) + 1 = 2x^2 - 3$$

$$(g \circ f)(x) = g(f(x)) = g(2x + 1) \\ = (2x + 1)^2 - 2 = 4x^2 + 4x - 1$$

$$\text{Thus } f \circ g = 2x^2 - 3, g \circ f = 4x^2 + 4x - 1.$$

From the above, we see that $f \circ g \neq g \circ f$.

- 1. 21.** Represent the function $f(x) = \sqrt{2x^2 - 5x + 3}$ as a composition of two functions.

Sol :

We set $f_2(x) = 2x^2 - 5x + 3$ and $f_1(x) = \sqrt{x}$

Then, $f(x) = \sqrt{2x^2 - 5x + 3}$

$$\begin{aligned} &= \sqrt{f_2(x)} \\ &= f_1[f_2(x)] = f_1 f_2(x) \end{aligned}$$

- 1. 22.** If $f(x) = 3x - 2$, $g(x) = 2x + k$ and $f \circ g = g \circ f$, then find the value of k .

Sol : $f(x) = 3x - 2, g(x) = 2x + k$

$$(f \circ g)(x) = f(g(x)) = f(2x + k) \\ = 3(2x + k) - 2 = 6x + 3k - 2$$

Thus, $(f \circ g)(x) = 6x + 3k - 2$.

$$(g \circ f)(x) = g(f(x)) = 2(3x - 2) + k$$

Thus, $(g \circ f)(x) = 6x - 4 + k$.

Given that $f \circ g = g \circ f$

$$\text{Therefore, } 6x + 3k - 2 = 6x - 4 + k$$

$$6x - 6x + 3k - k = -4 + 2 \Rightarrow k = -1$$

- 1. 23.** Find k if $f \circ f(k) = 5$ where $f(k) = 2k - 1$.

Sol :

$$f \circ f(k) = f(f(k)) = f(2k - 1)$$

Don

$$= 2(2k - 1) - 1 = 4k - 3$$

Thus, $f \circ f(k) = 4k - 3$

But, it is given that $f \circ f(k) = 5$

Therefore $4k - 3 = 5 \Rightarrow k = 2$.

- 1.24.** If $f(x) = 2x + 3$, $g(x) = 1 - 2x$ and $h(x) = 3x$.
Prove that $f \circ (g \circ h) = (f \circ g) \circ h$.

Sol :

$$f(x) = 2x + 3, g(x) = 1 - 2x, h(x) = 3x$$

$$\text{Now, } (f \circ g)(x) = f(g(x)) = f(1 - 2x)$$

$$= 2(1 - 2x) + 3 = 5 - 4x$$

$$\text{Since, } (f \circ g) \circ h(x) = (f \circ g)(h(x))$$

$$= (f \circ g)(3x)$$

$$= 5 - 4(3x) = 5 - 12x \quad \dots (1)$$

$$(g \circ h)(x) = g(h(x)) = g(3x)$$

$$= 1 - 2(3x) = 1 - 6x$$

$$\text{Since, } f \circ (g \circ h)(x) = f(1 - 6x) = 2(1 - 6x) + 3 = 5 - 12x \quad \dots (2)$$

From (1) and (2), we get

$$(f \circ g) \circ h = f \circ (g \circ h)$$

- 1.25.** Find x if $gff(x) = fgg(x)$, given $f(x) = 3x + 1$ and $g(x) = x + 3$.

Sol : $gff(x) = g[f\{f(x)\}]$

(This means "g of f of f of x")

$$= g[f(3x + 1)] = g[3(3x + 1) + 1] = g(9x + 4) = [(9x + 4) + 3] = 9x + 7$$

$$fgg(x) = f[g\{g(x)\}]$$

(This means "f of g of g of x")

$$= f[g(x + 3)] = f[(x + 3) + 3]$$

$$f(x + 6) = [3(x + 6) + 1] = 3x + 19$$

These two quantities being equal,

we get $9x + 7 = 3x + 19$. Solving this equation we obtain $x = 2$.

Progress Check

State your answer for the following questions by selecting the correct option.

- 1. Composition of functions is commutative**
- (a) Always true (b) Never true
 - (c) Sometimes true

Ans : (c) Sometimes true

- 2. Composition of functions is associative**
- (a) Always true (b) Never true
 - (c) Sometimes true

Ans : (a) Always true

- 3. Is a constant function a linear function?**

Ans : Yes

- 4. Is quadratic function a one-one function?**

Ans : No

- 5. Is cubic function a one-one function?**

Ans : Yes

- 6. Is the reciprocal function a bijection?**

Ans : Yes

- 7. Is $f : A \rightarrow B$ is a constant function, then the range of f will have _____ elements.**

Ans : One element

Thinking Corner

- 1. If $f(x) = x^m$ and $g(x) = x^n$ does $f \circ g = g \circ f$?**

Ans : $f(x) = x^m, g(x) = x^n$
 $f \circ g = f[g(x)] = f(x^n) = (x^n)^m = x^{nm}$
 $g \circ f = g[f(x)] = g[x^m] = (x^m)^n = x^{mn}$
 $\therefore f \circ g = g \circ f$

Exercise 1.5

- 1. Using the functions f and g given below, find $f \circ g$ and $g \circ f$. Check whether $f \circ g = g \circ f$.**

(i) $f(x) = x - 6, g(x) = x^2$

(ii) $f(x) = \frac{2}{x}, g(x) = 2x^2 - 1$

(iii) $f(x) = \frac{x+6}{3}, g(x) = 3 - x$

(iv) $f(x) = 3 + x, g(x) = x - 4$

(v) $f(x) = 4x^2 - 1, g(x) = 1 + x$

Sol :

(i) $f(x) = x - 6, g(x) = x^2$
 $f \circ g = f[g(x)] = f[x^2] = x^2 - 6$
 $g \circ f = g[f(x)] = g[x - 6] = (x - 6)^2 = x^2 - 12x + 36$
 $\therefore f \circ g \neq g \circ f$

(ii) $f(x) = \frac{2}{x}, g(x) = 2x^2 - 1$

$$f \circ g = f[g(x)] = f[2x^2 - 1] = \frac{2}{2x^2 - 1}$$

$$g \circ f = g[f(x)] = g\left[\frac{2}{x}\right] = 2\left(\frac{2}{x}\right)^2 - 1$$

$$= 2\left(\frac{4}{x^2}\right) - 1 = \frac{8 - x^2}{x^2}$$

$$\therefore f \circ g \neq g \circ f$$

Unit - 1 | RELATIONS AND FUNCTIONS

Don

(iii) $f(x) = \frac{x+6}{3}$, $g(x) = 3-x$

$$f \circ g = f[g(x)] = f(3-x) = \frac{3-x+6}{3} = \frac{9-x}{3}$$

$$\begin{aligned} g \circ f &= g[f(x)] = g\left(\frac{x+6}{3}\right) = 3 - \left(\frac{x+6}{3}\right) \\ &= \frac{9-x-6}{3} = \frac{3-x}{3} \therefore f \circ g \neq g \circ f \end{aligned}$$

(iv) $f(x) = 3+x$, $g(x) = x-4$

$$f \circ g = f[g(x)] = f[x-4] = 3+x-4 = x-1$$

$$g \circ f = g[f(x)] = g[3+x] = 3+x-4 = x-1$$

$$\therefore f \circ g = g \circ f$$

(v) $f(x) = 4x^2 - 1$, $g(x) = 1+x$

$$f \circ g = f[g(x)] = f(1+x) = 4(1+x)^2 - 1$$

$$= 4(1+2x+x^2) - 1$$

$$= 4+8x+4x^2 - 1$$

$$= 4x^2 + 8x + 3$$

$$g \circ f = g[f(x)] = 1+4x^2 - 1$$

$$= 4x^2$$

$$\therefore f \circ g \neq g \circ f$$

2. Find the value of k , such that $f \circ g = g \circ f$.

(i) $f(x) = 3x+2$, $g(x) = 6x-k$

(ii) $f(x) = 2x-k$, $g(x) = 4x+5$

Sol :

(i) $f(x) = 3x+2$, $g(x) = 6x-k$

$$f \circ g = f[g(x)] = f(6x-k)$$

$$= 3(6x-k) + 2$$

$$= 18x - 3k + 2$$

$$g \circ f = g[f(x)] = 6(3x+2) - k$$

$$= 18x + 12 - k$$

Given $f \circ g = g \circ f$

$$\therefore 18x - 3k + 2 = 18x + 12 - k$$

$$3k - k = -12 + 2$$

$$2k = -10$$

$$k = -5$$

(ii) $f(x) = 2x-k$, $g(x) = 4x+5$

$$f \circ g = f[g(x)] = f[4x+5]$$

$$= 2(4x+5) - k$$

$$= 8x + 10 - k$$

$$g \circ f = g[f(x)] = g(2x-k) = 4(2x-k) + 5$$

$$= 8x - 4k + 5$$

Given $f \circ g = g \circ f$

$$\therefore 8x + 10 - k = 8x - 4k + 5$$

$$4k - k = 5 - 10$$

$$3k = -5$$

$$k = -5/3$$

3. If $f(x) = 2x-1$, $g(x) = \frac{x+1}{2}$, show that

$$f \circ g = g \circ f = x.$$

Sol : $f(x) = 2x-1$, $g(x) = \frac{x+1}{2}$

$$f \circ g = f[g(x)] = f\left(\frac{x+1}{2}\right) = 2\left(\frac{x+1}{2}\right) - 1 = x + 1 - 1 = x$$

$$g \circ f = g[f(x)] = g(2x-1) = \frac{2x-1+1}{2}$$

$$= \frac{2x}{2} = x$$

$\therefore f \circ g = g \circ f = x$ Hence proved.

4. (i) If $f(x) = x^2 - 1$, $g(x) = x - 2$ find a , if $g \circ f(a) = 1$.

(ii) Find k , if $f(k) = 2k - 1$ and $f \circ f(k) = 5$

Sol :

(i) $f(x) = x^2 - 1$, $g(x) = x - 2$

$$f(a) = a^2 - 1$$

Given, $(g \circ f)(a) = 1$

$$g[f(a)] = 1$$

$$g[a^2 - 1] = 1$$

$$a^2 - 1 - 2 = 1$$

$$a^2 - 3 = 1$$

$$a^2 = 1 + 3 = 4$$

$$a = \pm 2$$

(ii) $f(k) = 2k - 1$

Given $(f \circ f)(k) = 5$

$$f[f(k)] = 1$$

$$f[2k - 1] = 1$$

$$\therefore 2(2k - 1) - 1 = 5$$

$$4k - 2 - 1 = 5$$

$$4k - 3 = 5$$

$$4k = 5 + 3 = 8$$

$$k = 8/4 = 2$$

5. Let $A, B, C \subseteq \mathbb{N}$ and a function $f : A \rightarrow B$ be defined by $f(x) = 2x+1$ and $g : B \rightarrow C$ be defined by $g(x) = x^2$. Find the range of $f \circ g$ and $g \circ f$.

Sol : $A, B, C \subseteq \mathbb{N}$

$$f : A \rightarrow B \text{ defined by } f(x) = 2x+1$$

$$g : B \rightarrow C \text{ defined by } g(x) = x^2$$

$$f \circ g = f[g(x)] = f(x^2) = 2x^2 + 1$$

$$g \circ f = g[f(x)] = g(2x+1) = (2x+1)^2$$

$$= 4x^2 + 4x + 1$$

' x ' can take any real value and can produce any real value. Thus, the domain and range of $f \circ g$ and $g \circ f$ is \mathbb{R} (Set of real numbers).

Don

- 6. If $f(x) = x^2 - 1$. Find (a) $f \circ f$ (b) $f \circ f \circ f$**

Sol: $f(x) = x^2 - 1$

(a) $f \circ f = f[f(x)] = f(x^2 - 1)$
 $= (x^2 - 1)^2 - 1 = x^4 - 2x^2 + 1 - 1 = x^4 - 2x^2$

(b) $f \circ f \circ f = f[f[f(x)]] = f[f(x^2 - 1)] = f[x^4 - 2x^2]$
 $= (x^4 - 2x^2)^2 - 1 = x^8 - 4x^6 + 4x^4 - 1$

- 7. If $f : R \rightarrow R$ and $g : R \rightarrow R$ are defined by $f(x) = x^5$ and $g(x) = x^4$ then check if f, g are one-one and $f \circ g$ is one-one.**

Sol:

$$f(x) = x^5, g(x) = x^4$$

$$f(x) = x^5$$

For any value of 'x', $f(x)$ gives us a different value (image) in co domain.

$\therefore f(x)$ is one-one function

$$g(1) = 1; g(-1) = 1$$

Hence $g(x)$ is not one-one function

$$f \circ g = f[g(x)] = f(x^4) = (x^4)^5 = x^{20}$$

$f \circ g$ is also One - One function as x is mapped with different value of $f \circ g$.

- 8. Consider the function $f(x), g(x), h(x)$ as given below. Show that $(f \circ g) \circ h = f \circ (g \circ h)$ in each case.**

(i) $f(x) = x - 1, g(x) = 3x + 1$ and $h(x) = x^2$

(ii) $f(x) = x^2, g(x) = 2x$ and $h(x) = x + 4$

(iii) $f(x) = x - 4, g(x) = x^2$ and $h(x) = 3x - 5$

Sol:

(i) $f(x) = x - 1, g(x) = 3x + 1, h(x) = x^2$

$$f \circ g = f[g(x)] = f[3x + 1]$$

$$= (3x + 1) - 1 = 3x$$

$$(f \circ g) \circ h = (f \circ g)[h(x)] = (f \circ g)[x^2] = 3x^2$$

$$g \circ h = g[h(x)] = g[x^2] = 3x^2 + 1$$

$$f \circ (g \circ h) = f[g(h(x))] = f[g(x^2)] = f[3x^2 + 1]$$

$$= (3x^2 + 1) - 1 = 3x^2$$

$$\therefore (f \circ g) \circ h = f \circ (g \circ h)$$

Hence proved.

(ii) $f(x) = x^2, g(x) = 2x, h(x) = x + 4$

$$f \circ g = f[g(x)] = f[2x]$$

$$= f(2x) = (2x)^2 = 4x^2$$

$$(f \circ g) \circ h = (f \circ g)[h(x)] = (f \circ g)[x + 4]$$

$$= 4(x + 4)^2$$

$$= 4(x^2 + 8x + 16) = 4x^2 + 32x + 64$$

$$g \circ h = g[h(x)] = g[x + 4] = 2(x + 4)$$

$$\begin{aligned} f \circ (g \circ h) &= f[g(h(x))] = f[g(x + 4)] \\ &= f[2(x + 4)] = [2(x + 4)]^2 = 4(x + 4)^2 \\ &= 4(x^2 + 8x + 16) = 4x^2 + 32x + 64 \\ \therefore (f \circ g) \circ h &= f \circ (g \circ h) \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad f(x) &= x - 4, g(x) = x^2, h(x) = 3x - 5 \\ f \circ g &= f[g(x)] = f(x^2) = x^2 - 4 \end{aligned}$$

$$\begin{aligned} (f \circ g) \circ h &= (f \circ g)[h(x)] = (f \circ g)[3x - 5] \\ &= (3x - 5)^2 - 4 = 9x^2 - 30x + 25 - 4 \\ &= 9x^2 - 30x + 21 \end{aligned}$$

$$\begin{aligned} g \circ h &= g[h(x)] \\ &= g(3x - 5) = (3x - 5)^2 = 9x^2 - 30x + 25 \\ f \circ (g \circ h) &= f[g(h(x))] = f[g(3x - 5)] = f[(3x - 5)^2] \\ &= 9x^2 - 30x + 25 - 4 = 9x^2 - 30x + 21 \\ \therefore (f \circ g) \circ h &= f \circ (g \circ h) \end{aligned}$$

- 9. Let $f = \{(-1, 3), (0, -1), (2, -9)\}$ be a linear function from Z into Z . Find $f(x)$.**

Sol: $f = \{(-1, 3), (0, -1), (2, -9)\}, f: Z \rightarrow Z$

Since 'f' is a linear function

$$f(x) = ax + b$$

$$\text{in } (0, -1), \text{ when } x = 0, f(0) = -1$$

$$\therefore a(0) + b = -1 \Rightarrow b = -1$$

$$\text{in } (-1, 3), \text{ when } x = -1, f(-1) = 3$$

$$\therefore a(-1) + b = 3 \Rightarrow -a - 1 = 3$$

$$-a = 4 \Rightarrow a = -4$$

$$\therefore f(x) = -4x - 1$$

- 10. In electrical circuit theory, a circuit $C(t)$ is called a linear circuit if it satisfies the superposition principle given by $C(at_1 + bt_2) = aC(t_1) + bC(t_2)$, where a, b are constants. Show that the circuit $C(t) = 3t$ is linear.**

Sol: Given superposition principle is

$$C(at_1 + bt_2) = aC(t_1) + bC(t_2) \quad a, b \text{ are constants}$$

If all the independent sources except for $C(t_1)$ have known fixed values, then

$$C(t) = aC(t_1) + d$$

$$\text{where } d = bC(t_2)$$

$\therefore C(t)$ is linear.

Aliter:

$C(t)$ is linear and $t = t_1 + t_2$

$$\text{Let } C(t_1) = t \text{ and } C(t_2) = 2t$$

By Given data,

$$C(at_1 + bt_2) = aC(t_1) + bC(t_2) \quad \dots (1)$$

$$\text{Now } C(t) = C(t_1 + t_2) \quad [\because t = t_1 + t_2]$$

$$= C(t_1) + C(t_2) \text{ from (1)}$$


$$= t + 2t = 3t$$

Hence the function $C(t)$ is linear.

Don

9. Let $A = \{1, 2, 3, 4\}$ and $B = \{4, 8, 9, 10\}$. A function $f : A \rightarrow B$ given by $f = \{(1, 4), (2, 8), (3, 9), (4, 10)\}$ is a
- Many-one function
 - Identity function
 - One-to-one function
 - Into function
- [Ans: (3)]

Sol: $A = \{1, 2, 3, 4\}$, $B = \{4, 8, 9, 10\}$



and $f = \{(1, 4), (2, 8), (3, 9), (4, 10)\}$
One-to-one function

10. If $f(x) = 2x^2$ and $g(x) = \frac{1}{3x}$, Then $f \circ g$ is

- $\frac{3}{2x^2}$
- $\frac{2}{3x^2}$
- $\frac{2}{9x^2}$
- $\frac{1}{6x^2}$

[Ans: (3)]

Sol: $f(x) = 2x^2$, $g(x) = 1/3x$
 $f \circ g = f[g(x)] = f\left[\frac{1}{3x}\right]$
 $= 2\left(\frac{1}{3x}\right)^2 = 2\left(\frac{1}{9x^2}\right)$
 $= \left(\frac{2}{9x^2}\right)$

11. If $f : A \rightarrow B$ is a bijective function and if $n(B) = 7$, then $n(A)$ is equal to

- 7
- 49
- 1
- 14

[Ans: (1)]

Sol:

$f : A \rightarrow B$ is a bijective function and $n(B) = 7$.
 f is a bijective function i.e., one-to-one onto function.
 \therefore A and B should have equal number of elements.

$$n(A) = 7$$

12. Let f and g be two functions given by
 $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 7)\}$
 $g = \{(0, 2), (1, 0), (2, 4), (-4, 2), (7, 0)\}$ then the range of $f \circ g$ is

- $\{0, 2, 3, 4, 5\}$
- $\{-4, 1, 0, 2, 7\}$
- $\{1, 2, 3, 4, 5\}$
- $\{0, 1, 2\}$

Sol: $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 7)\}$

$$g = \{(0, 2), (1, 0), (2, 4), (-4, 2), (7, 0)\}$$

$$\text{Now, } f \circ g(0) = f[2] = 0$$

$$f \circ g(1) = f[0] = 1$$

$$f \circ g(2) = f[4] = 2$$

$$f \circ g(-4) = f[2] = 0$$

$$f \circ g(7) = f[0] = 1$$

$$\text{Range of } f \circ g = \{0, 1, 2\}$$

13. Let $f(x) = \sqrt{1+x^2}$ then

- $f(xy) = f(x).f(y)$
- $f(xy) \geq f(x).f(y)$
- $f(xy) \leq f(x).f(y)$
- None of these

[Ans: (3)]

Sol: $f(x) = \sqrt{1+x^2}$

$$\text{then } f(y) = \sqrt{1+y^2}$$

$$\text{and } f(xy) = \sqrt{1+x^2y^2}$$

Hence, it is clear that $f(xy) \leq f(x)f(y)$

14. If $g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$ is a function given by $g(x) = \alpha x + \beta$ then the values of α and β are

- (-1, 2)
- (2, -1)
- (-1, -2)
- (1, 2)

[Ans: (2)]

Sol:

$$g = \{(1, 1), (2, 3), (3, 5), (4, 7)\}$$

$$g(x) = \alpha x + \beta$$

When $\alpha = 2$ and $\beta = -1$, $g(x) = 2x - 1$

Which is satisfying 'g'.

15. $f(x) = (x+1)^3 - (x-1)^3$ represents

- a linear function
- a cubic function
- a reciprocal function
- a quadratic function

[Ans: (4)]

Sol:

$$f(x) = (x+1)^3 - (x-1)^3$$

$$= (x^3 + 3x^2 + 3x + 1) - (x^3 - 3x^2 + 3x - 1)$$

$$= 6x^2 + 2$$

Don

$$\begin{aligned} B \times D &= \{1, 2, 3, 4\} \times \{5, 6, 7, 8\} \\ &= \{(1, 5), (1, 6), (1, 7), (1, 8), \\ &\quad (2, 5), (2, 6), (2, 7), (2, 8), \\ &\quad (3, 5), (3, 6), (3, 7), (3, 8), \\ &\quad (4, 5), (4, 6), (4, 7), (4, 8)\} \\ \therefore (A \times C) &\text{ is a subset of } (B \times D) \end{aligned}$$

8. If $f(x) = \frac{x-1}{x+1}$, $x \neq 1$ show that $f(f(x)) = -\frac{1}{x}$, provided $x \neq 0$.

Sol: $f(x) = \frac{x-1}{x+1}, x \neq -1$

$$\begin{aligned} f[f(x)] &= f\left[\frac{x-1}{x+1}\right] \\ &= \frac{\frac{x-1}{x+1}-1}{\frac{x-1}{x+1}+1} = \frac{x-1-(x+1)}{x-1+(x+1)} \\ &= \frac{x-1-x-1}{x-1+x+1} = -\frac{2}{2x} = -\frac{1}{x} \end{aligned}$$

Hence proved.

9. The function f and g are defined by $f(x) = 6x + 8$;
 $g(x) = \frac{x-2}{3}$

- (i) Calculate the value of $gg\left(\frac{1}{2}\right)$
(ii) Write an expression for $gf(x)$ in its simplest form.

Sol: $f(x) = 6x + 8$, $g(x) = \frac{x-2}{3}$

$$\begin{aligned} \text{(i)} \quad g\left(\frac{1}{2}\right) &= \frac{\frac{1}{2}-2}{3} = \frac{1-4}{2 \times 3} = -\frac{3}{6} = -\frac{1}{2} \\ \therefore gg\left(\frac{1}{2}\right) &= g\left[g\left(\frac{1}{2}\right)\right] \\ &= g\left(-\frac{1}{2}\right) = -\frac{-\frac{1}{2}-2}{3} = -\frac{1-4}{2 \times 3} = -\frac{5}{6} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad g f(x) &= g[f(x)] = g[6x + 8] \\ &= \frac{6x+8-2}{3} = \frac{6x+6}{3} \\ &= \frac{6(x+1)}{3} = 2(x+1) \end{aligned}$$

10. Write the domain of the following real functions


(i) $f(x) = \frac{2x+1}{x-9}$ (ii) $p(x) = \frac{-5}{4x^2+1}$

(iii) $g(x) = \sqrt{x-2}$ (iv) $h(x) = x + 6$

Sol:

(i) $f(x) = \frac{2x+1}{x-9}$

If $x - 9 = 0$, then
 $x = 9$



∴ The domain is all values of x that make the expression defined.

i.e., $(-\infty, 9) \cup (9, \infty)$

i.e., $(x / x \neq 9) \Rightarrow R - \{9\}$

(ii) $p(x) = \frac{-5}{4x^2+1}$

Here, the expression is defined for all real values of 'x'.

i.e., $x \in R$

(iii) $g(x) = \sqrt{x-2}$

$g(x)$ is defined real only when $x \geq 2$

(iv) $h(x) = x + 6$

$h(x)$ is defined for all real values of 'x'.

i.e., $x \in R$.



CREATIVE QUESTIONS

I. Multiple Choice Questions

Cartesian Product

1. If $A = \{1, 2\}$, $B = \{0, 1\}$, then $A \times B$ is
 (1) $\{(1, 0), (1, 1), (2, 0), (2, 1)\}$
 (2) $\{(1, 0), (2, 1)\}$
 (3) $\{(1, 1), (1, 2), (0, 1), (0, 2)\}$
 (4) None of these [Ans: (1)]

Sol :

$$\begin{aligned} A &= \{1, 2\}, B = \{0, 1\} \\ A \times B &= \{(1, 0), (1, 1), (2, 0), (2, 1)\} \end{aligned}$$

2. If the set A has 'p' elements, B has 'q' elements, then the number of elements in $A \times B$ is
 (1) $p+q$ (2) $p+q+1$
 (3) pq (4) p^2 [Ans: (3)]

Sol :

$$\begin{aligned} \text{Number of elements in } A &= p \\ \text{Number of elements in } B &= q \\ \text{Number of elements in } A \times B &= pq \end{aligned}$$

3. If A, B, C are any three sets, then $A \times (B \cup C)$ is equal to
 (1) $(A \times B) \cup (A \times C)$ (2) $(A \cup B) \cup (A \cup C)$
 (3) Both (a) and (b) (4) None of these [Ans: (1)]

4. Let $A = \{a, b, c, d\}$, $B = \{b, c, d, e\}$, then $n\{(A \times B) \cap (B \times A)\} =$

- (1) 3
 (2) 6
 (3) 9
 (4) None of these [Ans: (3)]

Sol :

$$A = \{a, b, c, d\}, B = \{b, c, d, e\}$$

$$A \times B = \{(a, b), (a, c), (a, d), (a, e), (b, b), (b, c), (b, d), (b, e), (c, b), (c, c), (c, d), (c, e), (d, b), (d, c), (d, d), (d, e)\}$$

$$B \times A = \{(b, a), (b, b), (b, c), (b, d), (c, a), (c, b), (c, c), (c, d), (d, a), (d, b), (d, c), (d, d), (e, a), (e, b), (e, c), (e, d)\}$$

$$(A \times B) \cap (B \times A) = \{(b, b), (b, c), (b, d), (c, b), (c, c), (c, d), (d, b), (d, c), (d, d)\}$$

$$n\{(A \times B) \cap (B \times A)\} = 9$$

Relations

6. If A is the set of even numbers less than 8 and B is the set of prime numbers less than 7, then the number of relations from A to B is
 (1) 2^9 (2) 9^2
 (3) 3^2 (4) 2^{9-1} [Ans: (1)]

Sol :

$$\begin{aligned} A &= \{2, 4, 6\} \Rightarrow n(A) = 3 \\ B &= \{2, 3, 5\} \Rightarrow n(B) = 3 \\ \therefore n(A \times B) &= 3 \times 3 = 9 \\ \text{Number of relations from } A \text{ to } B &= 2^9 \end{aligned}$$

7. Let N be the set of all natural numbers and let R be a relation on N defined as $R = \{(x, y) / x \in N, y \in N \text{ and } x + 3y = 15\}$. Then R as set of ordered pairs is

- (1) $\{(3, 4), (5, 3), (9, 2), (13, 2)\}$
 (2) $\{(3, 5), (2, 7), (9, 2), (12, 1)\}$
 (3) $\{(3, 4), (6, 3), (9, 2), (12, 1)\}$
 (4) $\{(4, 5), (7, 3), (4, 5), (4, 2)\}$ [Ans: (3)]

Sol :

$$\begin{aligned} R &= \{(x, y), x \in N, y \in N \text{ and } x + 3y = 15\} \\ x + 3y &= 15 \\ \text{When } x = 3, y &= 4, \text{ then } 3 + 3(4) = 15 \\ \text{When } x = 6, y &= 3, \text{ then } 6 + 3(3) = 15 \\ \text{When } x = 9, y &= 2, \text{ then } 9 + 3(2) = 15 \\ \text{When } x = 12, y &= 1, \text{ then } 12 + 3(1) = 15 \\ \therefore R &= \{(3, 4), (6, 3), (9, 2), (12, 1)\} \end{aligned}$$

8. A relation R is defined from $\{2, 3, 4, 5\}$ to $\{3, 6, 7, 10\}$ by : $x R y \Leftrightarrow x$ is relatively prime to y. Then, domain of R is

- (1) $\{2, 3, 5\}$ (2) $\{3, 5\}$
 (3) $\{2, 3, 4\}$ (4) $\{2, 3, 4, 5\}$ [Ans: (4)]

Sol :

$$\begin{aligned} \text{Since } x &\text{ is relatively prime to } y, \\ \text{Domain} &= \{2, 3, 4, 5\} \end{aligned}$$

9. Let R be a relation from set A to a set B, then

- (1) $R = A \cup B$ (2) $A \cap B$
 (3) $R \subseteq A \times B$ (4) $R \subseteq B \times A$ [Ans: (3)]

Sol :

Relation is a subset of Cartesian Product.

Unit - 1 | RELATIONS AND FUNCTIONS

(1) $x^2 - 2$
(3) $f\left(-\frac{a}{a+1}\right)$

(2) $x^2 - 1$
(4) $f(a)$

[Ans: (1)]

Sol :

Given $f\left(x + \frac{1}{x}\right) = x^2 + \frac{1}{x^2}$

Let us find $\left(x + \frac{1}{x}\right)^2 = x^2 + \frac{1}{x^2} + 2x \cdot \frac{1}{x}$
 $= x^2 + \frac{1}{x^2} + 2$

Subtracting 2 from this, we get $x^2 + \frac{1}{x^2}$

$$\therefore f(x) = x^2 - 2$$

17. If $f(x) = x - 2$, $g(x) = \sqrt{x^2 + 1}$, then
 $(g \circ f)(x) = ?$

- (1) $\sqrt{x^2 + 1} - 2$ (2) $\sqrt{x^2 + 4x + 5}$
(3) $x^2 - 1$ (4) $x^2 - 4x + 5$

[Ans: (2)]

Sol :

$$\begin{aligned}(g \circ f)(x) &= g[f(x)] \\&= g[x - 2] \\&= \sqrt{(x - 2)^2 + 1} \\&= \sqrt{x^2 - 4x + 4 + 1} \\&= \sqrt{x^2 - 4x + 5}\end{aligned}$$

18. Given $f(2) = 3$, $g(3) = 2$ and $g(2) = 5$, then
 $(f \circ g)(3) = ?$

- (1) 2 (2) 3
(3) 4 (4) 5

[Ans: (2)]

Sol :

$$\begin{aligned}(f \circ g)(3) &= f[g(3)] \\&= f[2] = 3\end{aligned}$$

19. Given $f = \{(-2, 1), (0, 3), (4, 5)\}$, $g = \{(1, 1), (3, 3), (4, 5)\}$ then, Domain and range of $g \circ f$.

- (1) D = {3, 0}, R = {-2, 1}
(2) D = {3, -2}, R = {1, 5}
(3) D = {-2, 0}, R = {1, 3}
(4) D = {-2, 1}, R = {0, 3}

[Ans: (3)]

Sol :

$$\begin{aligned}(g \circ f)(-2) &= g[f(-2)] = g(1) = 1 \\(g \circ f)(0) &= g[f(0)] = g(3) = 3 \\(g \circ f)(4) &= g[f(4)] \\&= g(5) = \text{undefined}\end{aligned}$$

$$\therefore g \circ f = \{(-2, 1), (0, 3)\}$$

Hence the Domain = {-2, 0},

Range = {1, 3}

Don**II. Very Short Answer Questions**

1. If $\left(\frac{x}{3} + 1, y - \frac{2}{3}\right) = \left(\frac{5}{3}, \frac{1}{3}\right)$, find the values of 'x' and 'y'

Sol : $\frac{x}{3} + 1 = \frac{5}{3}, \quad y - \frac{2}{3} = \frac{1}{3}$

$$\frac{x}{3} = \frac{5}{3} - 1, \quad y = \frac{1}{3} + \frac{2}{3}$$

$$\frac{x}{3} = \frac{5-3}{3}, \quad y = \frac{3}{3}$$

$$\frac{x}{3} = \frac{2}{3}, \quad y = 1$$

$$x = 2$$

2. If the Set 'A' has 3 elements and the Set B = {3, 4, 5}, find the number of elements in $(A \times B)$.

Sol :

'A' has 3 elements and 'B' has 3 elements, then ' $A \times B$ ' will have 9 elements.

3. If G = {7, 8} and H = {5, 4, 2}, find $G \times H$ and $H \times G$.

Sol :

$$\begin{aligned}G \times H &= \{7, 8\} \times \{5, 4, 2\} \\&= \{(7, 5), (7, 4), (7, 2), (8, 5), (8, 4), (8, 2)\}\end{aligned}$$

$$\begin{aligned}H \times G &= \{5, 4, 2\} \times \{7, 8\} \\&= \{(5, 7), (5, 8), (4, 7), (4, 8), (2, 7), (2, 8)\}\end{aligned}$$

4. If A = {-1, 1}, find $A \times A \times A$.

Sol :

$$\begin{aligned}A \times A &= \{-1, 1\} \times \{-1, 1\} \\&= \{(-1, -1), (-1, 1), (1, -1), (1, 1)\}\end{aligned}$$

$$\begin{aligned}A \times A \times A &= \{(-1, -1), (-1, 1), (1, -1), (1, 1)\} \\&\quad \times \{-1, 1\} \\&= \{(-1, -1, -1), (-1, -1, 1), (-1, 1, -1), \\&\quad (-1, 1, 1), (1, -1, -1), (1, -1, 1), \\&\quad (1, 1, -1), (1, 1, 1)\}\end{aligned}$$

5. If $A \times B = \{(a, x), (a, y), (b, x), (b, y)\}$, find 'A' and 'B'.

Sol :

$$\begin{aligned}A &= \text{Set of all first entries} = \{a, b\} \\B &= \text{Set of all second entries} = \{x, y\}\end{aligned}$$

Ques:

6. Let $A = \{3, 5\}$ and $B = \{7, 11\}$. Let $R = \{(a, b); a \in A, b \in B, a - b \text{ is odd}\}$. Show that R is an empty relation from A into B .

Sol :

Since $a \in A$ and $b \in B$,

$$\begin{aligned} a - b &= (3 - 7), (3 - 11), (5 - 7), (5 - 11) \\ &= -4, -8, -2, -6 \quad \text{None of them is an odd number.} \end{aligned}$$

Therefore, R is an empty relation.

7. What is the number of relations on A if R is a relation on a finite set A having ' n ' elements?

Sol :

Set A has ' n ' elements.

$$\therefore \text{Number of elements in } A \times A = n \times n = n^2$$

The number of possible subsets of $A \times A = 2^{n^2}$

Since, each subset of $A \times A$ is a relation on A , the total number of relations on A is 2^{n^2} .

8. Let set $A = \{\text{January, February, April, June, September, October, November, December}\}$ and set $B = \{28, 29, 30, 31\}$. Let R be a relation from A to B defined by $R = \{(a, b) \in A \times B : 'a' \text{ month has 'b' number of days}\}$. Write a subset of relation R connected with 'Teachers Day'.

Sol :

$$R = \{(a, b) \in A \times B, 'a' \text{ month has 'b' number of days}\}$$

$$\therefore R = \{(\text{January}, 31), (\text{February}, 28), (\text{February}, 29), (\text{April}, 30), (\text{June}, 30), (\text{September}, 30), (\text{October}, 31), (\text{November}, 30), (\text{December}, 31)\}$$

\therefore The subset of relation R connected with Teachers Day is $\{(\text{September}, 30)\}$

9. A function ' f ' is defined by $f(x) = 2x - 5$. Write down the values of $f(0)$, $f(7)$ and $f(-3)$.

Sol :

$$f(x) = 2x - 5$$

$$f(0) = 2(0) - 5 = -5$$

$$f(7) = 2(7) - 5 = 14 - 5 = 9$$

$$f(-3) = 2(-3) - 5 = -6 - 5 = -11$$

10. If $A = \{1, 2, 3\}$ and $B = \{a, b, c\}$ then do

- $\{(1, a), (2, b), (2, c), (3, c)\}$ and
- $\{(2, b), (3, b)\}$ represent a function $A \rightarrow B$?

Sol :

- The two ordered pairs $(2, b)$ and $(2, c)$ have the same first co-ordinate. Therefore, $(1, a), (2, b), (2, c), (3, c)$ does not represent a function $A \rightarrow B$.

- (ii) Since one element 1 of A is not associated with some element of B . i.e., 1 is not the first co-ordinate of any ordered pair. So $\{(2, b), (3, b)\}$ does not represent a function from $A \rightarrow B$.

11. Let $A = \{1, 2, 3\}$ and $B = \{4, 5\}$. Let $f = \{(1, 4), (1, 5), (2, 4), (3, 5)\}$. Is ' f ' a function from A into B ?

Sol :

The first elements of two ordered pairs $(1, 4)$ and $(1, 5)$ in ' f ' are same.

Therefore $f = \{(1, 4), (1, 5), (2, 4), (3, 5)\}$ is not a function from A to B .

12. Find the domain of $f(x) = \frac{x}{x^2 - 5x + 6}$.

Sol :

$$f(x) = \frac{x}{x^2 - 5x + 6} = \frac{x}{(x-3)(x-2)}$$

When $x = 2$ or 3 , $f(x)$ becomes undefined.

\therefore Domain of $f = R - \{2, 3\}$

13. Let $f : R \rightarrow R$ defined by $f(x) = \begin{cases} 3x - 2, & x < 0 \\ 1, & x = 0 \\ 4x + 1, & x > 0 \end{cases}$.
Find $f(-3)$, $f(0)$ and $f(5)$.

Sol :

$$f(-3) = 3(-3) - 2 = -11 \text{ as } -3 < 0$$

$$f(0) = 1 \text{ as } x = 0$$

$$f(5) = 4(5) + 1 = 20 + 1 = 21 \text{ as } x > 0$$

14. Let a function $f : R \rightarrow A$ be defined as

$$f(x) = \begin{cases} 1, & \text{if } x \text{ is rational} \\ -1, & \text{if } x \text{ is irrational, where } x \in R \end{cases}$$


Find $f\left(\frac{1}{3}\right)$ and $f(\sqrt{5})$.

Sol :

As $\frac{1}{3}$ is a rational number, $f\left(\frac{1}{3}\right) = 1$ and

as $\sqrt{5}$ is an irrational number, $f(\sqrt{5}) = -1$.

15. Consider the following arrow diagram.



Unit - 1 | RELATIONS AND FUNCTIONS

i) Is it a function? If so, write the equation in form.

ii) List the elements of this function.

Sol :

Yes, it is a function and $f(x) = x^3$

Elements of $f = \{(-2, -8), (-1, -1), (0, 0), (1, 1), (2, 8)\}$

16. If $f(x) = \frac{x-1}{x+1}$, then show that $f\left(\frac{1}{x}\right) = -f(x)$

Sol :

$$\begin{aligned} f\left(\frac{1}{x}\right) &= \frac{\frac{1}{x}-1}{\frac{1}{x}+1} = \frac{\frac{1-x}{x}}{\frac{1+x}{x}} = \frac{1-x}{1+x} \\ &= -\frac{(x-1)}{x+1} = -f(x) \end{aligned}$$

Hence proved.

17. Given $f(x) = 3x + 2$, $g(x) = x + 5$ find $f[g(x)]$ and $g[f(x)]$.

Sol :

$$\begin{aligned} f[g(x)] &= f[x+5] = 3(x+5) + 2 \\ &= 3x + 15 + 2 = 3x + 17 \\ g[f(x)] &= g[3x+2] = 3x + 2 + 5 = 3x + 7 \end{aligned}$$

18. If $f(x) = 2x - 1$, then find $f \circ f$.

Sol :

$$\begin{aligned} f(x) &= 2x - 1 \\ f \circ f &= f[f(x)] \\ &= f(2x - 1) = 2(2x - 1) - 1 \\ &= 4x - 2 - 1 = 4x - 3 \end{aligned}$$

19. Given $f(x) = x^2 + 6$ and $g(x) = 2x + 1$, find $(g \circ f)(2)$.

Sol :

$$\begin{aligned} g \circ f &= g[f(x)] \\ &= g[x^2 + 6] \\ &= 2(x^2 + 6) + 1 \\ &= 2x^2 + 12 + 1 = 2x^2 + 13 \end{aligned}$$

$$\begin{aligned} \text{Now, } (g \circ f)(2) &= 2(2)^2 + 13 \\ &= 2(4) + 13 = 8 + 13 = 21 \end{aligned}$$

III. Short Answer Questions:

1. If $A = \{1, 2, 3\}$, $B = \{3, 4\}$ and $C = \{4, 5, 6\}$, then find $(A \times B) \cup (A \times C)$.

Sol :

$$\begin{aligned} A \times B &= \{1, 2, 3\} \times \{3, 4\} \\ &= \{(1, 3), (1, 4), (2, 3), (2, 4), (3, 3), (3, 4)\} \\ A \times C &= \{1, 2, 3\} \times \{4, 5, 6\} \end{aligned}$$

$$\begin{aligned} &= \{(1, 4), (1, 5), (1, 6), (2, 4), (2, 5), \\ &\quad (2, 6), (3, 4), (3, 5), (3, 6)\} \end{aligned}$$

$$\begin{aligned} \therefore (A \times B) \cup (A \times C) &= \{(1, 3), (1, 4), (1, 5), (1, 6), \\ &\quad (2, 3), (2, 4), (2, 5), (2, 6), \\ &\quad (3, 3), (3, 4), (3, 5), (3, 6)\} \end{aligned}$$

2. If $A = \{1, 2, 4\}$, $B = \{2, 4, 5\}$, $C = \{2, 5\}$, find $(A - B) \times (B - C)$

Sol :

$$A - B = \{1, 2, 4\} - \{2, 4, 5\}$$

$$= \{1\}$$

$$B - C = \{2, 4, 5\} - \{2, 5\}$$

$$= \{4\}$$

$$\therefore (A - B) \times (B - C) = \{1\} \times \{4\}$$

$$= \{(1, 4)\}$$

3. If $R_1 = \{(x, y) / y = 2x + 7$, where $x \in R$ and $-5 \leq y \leq 5\}$ is a relation, then find the domain and range of R_1 .

Sol :

$$R_1 = \{(x, y) / y = 2x + 7, x \in R, -5 \leq y \leq 5\}$$

$$x = \{..., -6, -5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, ...\}$$

$$y = \{-5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5\}$$

$$\text{When } x = -6, y = -5$$

$$x = -5, y = -3$$

$$x = -4, y = -1$$

$$x = -3, y = 1$$

$$x = -2, y = 3$$

$$x = -1, y = 5$$

$$\therefore R = \{(-6, -5), (-5, -3), (-4, -1), (-3, 1), (-2, 3), (-1, 5)\}$$

$$\text{Domain} = \{-6, -5, -4, -3, -2, -1\}$$

$$\text{Range} = \{-5, -3, -1, 1, 3, 5\}$$

4. Let R be a relation in N defined by $R = \{(1+x, 1+x^2) / x \leq 4, x \in N\}$. Find

(i) R

(ii) domain of R

(iii) range of R

Sol :

$$R = \{(1+x, 1+x^2) / x \leq 4, x \in N\}$$

as $x \leq 4$, $x \in N$, x takes the values 1, 2, 3, 4

$$\therefore R = \{(2, 2), (3, 5), (4, 10), (5, 17)\}$$


$$\text{Domain of } R = \{2, 3, 4, 5\}$$

$$\text{Range of } R = \{2, 5, 10, 17\}$$

Ques

5. The figure shows a relationship between the Sets P and Q. Write this relation in

- (i) Set builder form
- (ii) Roster form
- (iii) What is its domain and Range?

**Sol:**

- (i) Set builder form $R = \{(x, y) / y = x - 2 \text{ for } x = 5, 6, 7\}$
- (ii) Roster form $R = \{(5, 3), (6, 4), (7, 5)\}$
- (iii) Domain of R = {5, 6, 7}
Range of R = {3, 4, 5}

6. Find the domain and range for the functions

$$(a) f(x) = x^2 + 2, \quad (b) f(t) = \frac{1}{t+2}$$

Sol:

(a) $f(x) = x^2 + 2$ is defined for all real values of 'x'. Hence the domain of $f(x)$ is "all real values of x". Since ' x^2 ' is always positive, $x^2 + 2$ is never less than 2.
 \therefore The range of $f(x)$ is "all real numbers $f(x) \geq 2$ ".

$$(b) f(t) = \frac{1}{t+2}$$

Domain: The function is not defined for $t = -2$. Hence, the domain of $f(t)$ is "all real numbers except -2"

Range: "all real numbers except zero".

7. Let $f = \{(1, 1), (2, 3), (0, -1), (-1, -3)\}$ be a function from Z to Z defined by $f(x) = ax + b$, for some integers a, b. Determine a, b.

Sol:

$$f(x) = ax + b$$

$$\text{for } x = 1, f(x) = 1$$

$$\therefore a(1) + b = 1 \Rightarrow a + b = 1 \quad \dots (1)$$

$$\text{for } x = 2, f(x) = 3$$

$$\therefore a(2) + b = 3 \Rightarrow 2a + b = 3 \quad \dots (2)$$

Solving (1) and (2)

We get, a = 2, b = -1

$$\therefore f(x) = 2x - 1$$

Also for $x = 0, f(0) = -1$ and
when $x = -1, f(-1) = -3$.

8. Find the domain and Range of the real function

$$'f' \text{ defined by } f(x) = \frac{4-x}{x-4}.$$

Sol:

When $x = 4, f(x)$ is not defined as $f(4) = \frac{4-4}{4-4} = \frac{0}{0}$. So, the domain is the set of all real numbers except $x = 4$ i.e., $R - \{4\}$.

$$\text{and } f(x) = \frac{4-x}{x-4} = -\frac{(x-4)}{x-4} = -1, x \neq 4$$

$$\therefore \text{Range of } f(x) = \frac{4-x}{x-4} \text{ is } -1.$$

$$\text{Domain} = R - \{4\}, \text{Range} = \{-1\}$$

9. Express the following function as set of ordered pairs and determine the range: $f : X \rightarrow R, f(x) = x^3 + 1$, where $X = \{-1, 0, 3, 9, 7\}$.

Sol:

$$f : X \rightarrow R, f(x) = x^3 + 1 \text{ where } X = \{-1, 0, 3, 9, 7\}$$

$$f(-1) = (-1)^3 + 1 = -1 + 1 = 0$$

$$f(0) = (0)^3 + 1 = 0 + 1 = 1$$

$$f(3) = (3)^3 + 1 = 27 + 1 = 28$$

$$f(9) = (9)^3 + 1 = 729 + 1 = 730$$

$$f(7) = (7)^3 + 1 = 343 + 1 = 344$$

$\therefore f = \{(-1, 0), (0, 1), (3, 28), (9, 730), (7, 344)\}$ and
Range of $f = \{0, 1, 28, 730, 344\}$

10. If $A = \{-1, 0, 2, 5, 1\}$ and $f : A \rightarrow Z$ is defined by $f(x) = x^2 - x - 2$. Find the range of 'f'.

Sol:

$$\text{Given } A = \{-1, 0, 2, 5, 1\}$$

$$f(x) = x^2 - x - 2$$

$$f(-1) = (-1)^2 - (-1) - 2 = 0$$

$$f(0) = (0)^2 - 0 - 2 = -2$$

$$f(2) = (2)^2 - 2 - 2 = 0$$

$$f(5) = (5)^2 - 5 - 2 = 18$$

$$f(1) = (1)^2 - 1 - 2 = -2$$

\therefore Range of $f = \{-2, 0, 18\}$

11. If $f(x) = x^2$, find $\frac{f(1.1) - f(1)}{(1.1 - 1)}$

Sol:

$$f(x) = x^2$$

$$f(1.1) = (1.1)^2 = 1.21$$

$$f(1) = (1)^2 = 1$$

$$\therefore \frac{f(1.1) - f(1)}{(1.1 - 1)} = \frac{1.21 - 1}{1.1 - 1}$$

$$= \frac{0.21}{0.1} \times \frac{100}{100} = \frac{21}{10} = 2.1$$

Unit - 1 | RELATIONS AND FUNCTIONS

12. If $f(x) = x^2 - 2$ and $g(x) = 2x - 3$, then Evaluate :

- (i) $f(-3)$ (ii) $g(5)$ (iii) $f(a + 1)$

Sol :

- (i) $f(-3)$

$$\begin{aligned} f(x) &= x^2 - 2 \\ f(-3) &= (-3)^2 - 2 = 9 - 2 = 7 \end{aligned}$$

- (ii) $g(5)$

$$\begin{aligned} g(x) &= 2x - 3 \\ g(5) &= 2(5) - 3 = 10 - 3 = 7 \end{aligned}$$

- (iii) $f(a + 1)$

$$\begin{aligned} f(x) &= x^2 - 2 \\ f(a + 1) &= (a + 1)^2 - 2 \\ &= a^2 + 2a + 1 - 2 = a^2 + 2a - 1 \end{aligned}$$

13. Given $f(x) = 4x + 2$, Evaluate $\frac{f(a+h)-f(a)}{h}$.

Sol :

$$\begin{aligned} f(x) &= 4x + 2 \\ f(a+h) &= 4(a+h) + 2 \\ &= 4a + 4h + 2 \\ f(a) &= 4a + 2 \end{aligned}$$

$$\begin{aligned} \therefore \frac{f(a+h)-f(a)}{h} &= \frac{4a + 4h + 2 - 4a - 2}{h} \\ &= \frac{4h}{h} = 4 \end{aligned}$$

14. Given $f(x) = 3x + 7$, $g(x) = x + 1$, $h(x) = 2x - 3$. Find $f(2) + h(1)$ and $f(0) + g(0) - h(0)$.

Sol :

$$\begin{aligned} f(2) + h(1) &= 3(2) + 7 + 2(1) - 3 \\ &= 6 + 7 + 2 - 3 = 12 \end{aligned}$$

$$\begin{aligned} f(0) + g(0) - h(0) &= 3(0) + 7 + 0 + 1 - (2(0) - 3) \\ &= 0 + 7 + 0 + 1 - 0 + 3 = 11 \end{aligned}$$

15. If $f(x) = 2x^4 + x^2 + 1$ and $g(x) = \sqrt{x}$. Find $f \circ g$.

Sol :

$$\begin{aligned} f \circ g &= f[g(x)] = f[\sqrt{x}] \\ &= 2(\sqrt{x})^4 + (\sqrt{x})^2 + 1 = 2x^2 + x + 1 \end{aligned}$$

16. If $f(x) = -2x + 9$ and $g(x) = -4x^2 + 5x - 3$, find $f \circ g$ and $g \circ f$.

Sol :

$$\begin{aligned} f \circ g &= f[g(x)] \\ &= f[-4x^2 + 5x - 3] \\ &= -2(-4x^2 + 5x - 3) + 9 \\ &= 8x^2 - 10x + 15 \end{aligned}$$

$$\begin{aligned} g \circ f &= g[f(x)] \\ &= g[-2x + 9] \\ &= -4(-2x + 9)^2 + 5(-2x + 9) - 3 \end{aligned}$$

$$\begin{aligned} &= -4(4x^2 - 36x + 81) - 10x + 45 - 3 \\ &= -16x^2 + 144x - 324 - 10x + 42 \\ &= -16x^2 + 134x - 282 \end{aligned}$$

17. If $g(x) = \sqrt[3]{x-4}$ and $h(x) = x^3 + 4$, find $(h \circ g)(-15)$.

Sol :

$$\begin{aligned} h \circ g &= h[g(x)] \\ &= h[\sqrt[3]{x-4}] \\ &= (\sqrt[3]{x-4})^3 + 4 \\ &= (x-4)^{\frac{3}{3}} + 4 \\ &= x-4+4 = x \\ \therefore (h \circ g)(-15) &= -15 \end{aligned}$$

IV. Long Answer Questions

1. If $A = \{x / x^2 - 4x + 3 = 0\}$, $B = \{x / x^2 - x - 6 = 0\}$, $C = \{x / x^3 - 4x = 0\}$, find (i) $A \times B$ (ii) $A \times C$ (iii) $(A - B) \times C$

Sol :

$$\begin{aligned} A &= \{x / x^2 - 4x + 3 = 0\} \\ &= \{x / (x-3)(x-1) = 0\} \\ &= \{x / x = 1, 3\} \\ B &= \{x / x^2 - x - 6 = 0\} \\ &= \{x / (x+2)(x-3) = 0\} \\ &= \{x / x = -2, 3\} \\ C &= \{x / x^3 - 4x = 0\} \\ &= \{x / x(x^2 - 4) = 0\} \\ &= \{x / x = 0, -2, 2\} \\ A &= \{1, 3\}, B = \{-2, 3\}, C = \{-2, 0, 2\} \end{aligned}$$

$$\begin{aligned} \text{(i)} \quad A \times B &= \{1, 3\} \times \{-2, 3\} \\ &= \{(1, -2), (1, 3), (3, -2), (3, 3)\} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad A \times C &= \{1, 3\} \times \{-2, 0, 2\} \\ &= \{(1, -2), (1, 0), (1, 2), (3, -2), \\ &\quad (3, 0), (3, 2)\} \end{aligned}$$

$$\begin{aligned} \text{(iii)} \quad A - B &= \{1, 3\} - \{-2, 3\} = \{1\} \\ (A - B) \times C &= \{1\} \times \{-2, 0, 2\} \\ &= \{(1, -2), (1, 0), (1, 2)\}. \end{aligned}$$

2. If A and B be two non-empty sets, then show that $A \times B = B \times A$ if $A = B$.

Sol :

$$\begin{aligned} \text{Let } A \times B &= B \times A \text{ and } x \in A \\ x \in A &\Rightarrow (x, b) \in A \times B \forall b \in B \\ &\Rightarrow (x, b) \in (B \times A) \end{aligned}$$

Don

$$\begin{aligned}\because A \times B &= B \times A \\ \Rightarrow x &\in B \\ \Rightarrow A &\subseteq B\end{aligned}$$

By the definition of Cartesian Product and let y be an arbitrary element of B
 $y \in B \Rightarrow (a, y) \in A \times B \forall a \in A$

$$\begin{aligned}\Rightarrow (a, y) &\in (B \times A) \\ \therefore A \times B &= B \times A\end{aligned}$$

By the definition of Cartesian Product

$$\begin{aligned}\Rightarrow y &\in A \\ B &\subseteq A\end{aligned}$$

From these two results, $A = B$

Conversely, let $A = B$, then

$$\begin{aligned}A \times B &= A \times A \text{ and } B \times A = A \times A \\ \therefore A \times B &= B \times A\end{aligned}$$

Hence $A \times B = B \times A$ if $A = B$.

3. In the given ordered pairs $(4, 6), (8, 4), (3, 3), (9, 11), (6, 3), (3, 0), (2, 3)$. Find the following relations. Also, find the domain and range.

- | | |
|-----------------------|-------------------|
| (i) is two less than | (ii) is less than |
| (iii) is greater than | (iv) is equal to |

Sol :

(i) R_1 is the set of all ordered pairs whose first component is two less than the second component.

$$\therefore R_1 = \{(4, 6), (9, 11)\}$$

$$\begin{aligned}\text{Domain of } R_1 &= \text{Set of all first components} \\ &= \{4, 9\}\end{aligned}$$

$$\begin{aligned}\text{Range of } R_1 &= \text{Set of all second components} \\ &= \{6, 11\}\end{aligned}$$

(ii) R_2 is the set of all ordered pairs whose first component is less than the second component.

$$\therefore R_2 = \{(4, 6), (9, 11), (2, 3)\}$$

$$\text{Domain of } R_2 = \{4, 9, 2\}$$

$$\text{Range of } R_2 = \{6, 11, 3\}$$

(iii) R_3 is the set of all ordered pairs whose first component is greater than the second component.

$$\therefore R_3 = \{(8, 4), (6, 3), (3, 0)\}$$

$$\text{Domain of } R_3 = \{8, 6, 3\}$$

$$\text{Range of } R_3 = \{4, 3, 0\}$$

(iv) R_4 is the set of all ordered pairs whose first component is equal to the second component.

$$\therefore R_4 = \{(3, 3)\}$$

$$\text{Domain of } R_4 = \{3\} \text{ and Range of } R_4 = \{3\}$$


10th Std | MATHEMATICS

4. Let $A = \{2, 3, 4, 5\}$ and $B = \{8, 9, 10, 11\}$. Let R be the relation “is factor of” from A and B .
- Write R in the roster form. Also find Domain and Range of R .
 - Draw an arrow diagram to represent the relation.

Sol :

(i) Given, that the Relation R consists of elements (a, b) where ‘ a ’ is a factor of ‘ b ’. Therefore, Relation ‘ R ’ in the roster form is
 $R = \{(2, 8), (2, 10), (3, 9), (4, 8), (5, 10)\}$
 Domain of $R = \{2, 3, 4, 5\}$
 Range of $R = \{8, 10, 9\}$

(ii) Arrow diagram representing ‘ R ’



5. Given $f(x) = 3x + 7$, $g(x) = -x + 8$, $h(x) = x^2 + 3x - 1$. Prove that Composition of functions is associative.

Sol :

$$\begin{aligned}f \circ g &= f[g(x)] \\ &= f[-x + 8] \\ &= 3(-x + 8) + 7 \\ &= -3x + 24 + 7 = -3x + 31 \\ (f \circ g) \circ h &= (f \circ g)[h(x)] \\ &= (f \circ g)[x^2 + 3x - 1] \\ &= -3(x^2 + 3x - 1) + 31 \\ &= -3x^2 - 9x + 3 + 31 \\ &= -3x^2 - 9x + 34\end{aligned} \quad \dots (1)$$

$$\begin{aligned}g \circ h &= g[h(x)] \\ &= g[x^2 + 3x - 1] \\ &= -(x^2 + 3x - 1) + 8 \\ &= -x^2 - 3x + 1 + 8 = -x^2 - 3x + 9\end{aligned}$$

$$\begin{aligned}f \circ (g \circ h) &= f[g(h(x))] = f[g(x^2 + 3x - 1)] \\ &= f[-x^2 - 3x + 9] \\ &= 3(-x^2 - 3x + 9) + 7 \\ &= -3x^2 - 9x + 27 + 7\end{aligned}$$

Unit - 1 | RELATIONS AND FUNCTIONS**Don**

$$= -3x^2 - 9x + 34 \quad \dots (2)$$

From (1) and (2)

$$(f \circ g) \circ h = f \circ (g \circ h)$$

\therefore Composition of functions is Associative.

6. If 'f' and 'g' are two real valued functions defined as $f(x) = 2x + 1$, $g(x) = x^2 + 1$, then find

- (i) $f + g$ (ii) $f - g$ (iii) fg (iv) $\frac{f}{g}$

Sol:

$$\begin{aligned} \text{(i)} \quad (f+g)(x) &= f(x) + g(x) \\ &= 2x + 1 + x^2 + 1 \\ &= x^2 + 2x + 2 \end{aligned}$$

(ii) $(f-g)(x) = f(x) - g(x)$

$$= (2x+1) - (x^2+1)$$

$$= 2x + 1 - x^2 - 1 = 2x - x^2$$

$$= -x^2 + 2x$$

(iii) $(fg)(x) = [f(x)][g(x)]$

$$= (2x+1)(x^2+1)$$

$$= 2x^3 + x^2 + 2x + 1$$

(iv) $\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$, provided $g(x) \neq 0$

$$= \frac{2x+1}{x^2+1}$$

