

# Matgeo Presentation - Problem 12.797

ee25btech11056 - Suraj.N

October 11, 2025

## Problem Statement

Let  $\mathbf{A}$  be an  $n \times n$  real matrix. Consider the following statements:

- (I) If  $\mathbf{A}$  is symmetric, then there exists  $c \geq 0$  such that  $\mathbf{A} + c\mathbf{I}_n$  is symmetric and positive definite, where  $\mathbf{I}_n$  is the  $n \times n$  identity matrix.
- (II) If  $\mathbf{A}$  is symmetric and positive definite, then there exists a symmetric and positive definite  $\mathbf{B}$  such that  $\mathbf{A} = \mathbf{B}^2$ .

Which of the above statements is/are true?

- (a) Only (I)
- (b) Only (II)
- (c) Both (I) and (II)
- (d) Neither (I) nor (II)

<b>Name</b>	<b>Description</b>
<b>A</b>	Matrix

Table : Matrix

## Solution

Checking statement (I)

If  $\mathbf{A}$  is symmetric, its eigenvalues are real. Let the minimum eigenvalue of  $\mathbf{A}$  be  $\lambda_{\min}$ . Then choose  $c > -\lambda_{\min}$ .

The Eigen values of  $\mathbf{A}$  are given as :

$$|\mathbf{A} - \lambda_i \mathbf{I}| = 0 \quad (0.1)$$

The Eigen values of  $\mathbf{A} + c\mathbf{I}_n$  are given as :

$$|\mathbf{A} - (\lambda_k - c)\mathbf{I}| = 0 \quad (0.2)$$

$$\lambda_k = \lambda_i + c \quad (0.3)$$

$$\lambda_i + c > 0 \quad (0.4)$$

Since  $\lambda_i + c > 0$  for all  $i$ ,  $\mathbf{A} + c\mathbf{I}_n$  is positive definite and symmetric.

Hence, statement (I) is **true**.

## Solution

Checking statement (II)

If  $\mathbf{A}$  is symmetric and positive definite, then it can be diagonalized as:

$$\mathbf{A} = \mathbf{P}\mathbf{D}\mathbf{P}^{\top} \quad (0.5)$$

where  $\mathbf{P}$  is orthogonal and  $\mathbf{D}$  is a diagonal matrix with positive entries (since  $\mathbf{A}$  is positive definite). Define

$$\mathbf{B} = \mathbf{P}\mathbf{D}^{1/2}\mathbf{P}^{\top} \quad (0.6)$$

Then,

$$\mathbf{B}^2 = \mathbf{P}\mathbf{D}^{1/2}\mathbf{P}^{\top}\mathbf{P}\mathbf{D}^{1/2}\mathbf{P}^{\top} = \mathbf{P}\mathbf{D}\mathbf{P}^{\top} = \mathbf{A} \quad (0.7)$$

Hence,  $\mathbf{B}$  is symmetric and positive definite. Therefore, statement (II) is also **true**.

**Final Answer:** (c) Both (I) and (II)

# Solution

## Examples

### Example a for (I)

$$\mathbf{A} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (0.8)$$

To find eigenvalues, evaluate:

$$|\mathbf{A} - \lambda \mathbf{I}| = 0 \quad (0.9)$$

$$\begin{vmatrix} -\lambda & 1 \\ 1 & -\lambda \end{vmatrix} = 0 \quad (0.10)$$

$$\begin{vmatrix} -\lambda & 1 \\ 1 & -\lambda \end{vmatrix} \xleftrightarrow{R_2 \rightarrow R_2 + \frac{1}{\lambda} R_1} \begin{vmatrix} -\lambda & 1 \\ 0 & \frac{1-\lambda^2}{\lambda} \end{vmatrix} = 0 \quad (0.11)$$

$$\lambda^2 - 1 = 0 \quad (0.12)$$

$$\lambda = \pm 1 \quad (0.13)$$

$$\lambda_1 = 1, \lambda_2 = -1 \quad (0.14)$$

## Solution

The minimum eigenvalue  $\lambda_{\min} = -1$ . Choose  $c = 2$ . Then:

$$\lambda_1 + c = 3, \quad \lambda_2 + c = 1 \quad (0.15)$$

All eigenvalues are positive, so  $\mathbf{A} + 2\mathbf{I}$  is symmetric positive definite.

**Example b for (I)**

$$\mathbf{A} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix} \quad (0.16)$$

Compute eigenvalues from  $|\mathbf{A} - \lambda\mathbf{I}| = 0$ :

$$\begin{vmatrix} -2 - \lambda & 0 & 0 \\ 0 & 1 - \lambda & 0 \\ 0 & 0 & 3 - \lambda \end{vmatrix} = 0 \quad (0.17)$$

$$(-2 - \lambda)(1 - \lambda)(3 - \lambda) = 0 \quad (0.18)$$

Hence

$$\lambda_1 = -2, \quad \lambda_2 = 1, \quad \lambda_3 = 3 \quad (0.19)$$

## Solution

Choose  $c = 3$  so  $c > -\lambda_{\min} = 2$ . Then  $\lambda_i + c > 0$  for all  $i$ . Thus  $\mathbf{A} + 3\mathbf{I}$  is symmetric positive definite.

### Example a for (II)

$$\mathbf{A} = \begin{pmatrix} 5 & 2 \\ 2 & 5 \end{pmatrix} \quad (0.20)$$

Find eigenvalues from  $|\mathbf{A} - \lambda\mathbf{I}| = 0$ :

$$\begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix} = 0 \quad (0.21)$$

$$\begin{vmatrix} 5 - \lambda & 2 \\ 2 & 5 - \lambda \end{vmatrix} \xleftrightarrow{R_2 \rightarrow R_2 - \frac{2}{5-\lambda} R_1} \begin{vmatrix} 5 - \lambda & 2 \\ 0 & (5 - \lambda) - \frac{4}{5-\lambda} \end{vmatrix} = 0 \quad (0.22)$$

$$(5 - \lambda)^2 - 4 = 0 \quad (0.23)$$

$$\lambda^2 - 10\lambda + 21 = 0 \quad (0.24)$$

$$\lambda_1 = 7, \lambda_2 = 3 \quad (0.25)$$



## Solution

For eigenvectors:

$$(\mathbf{A} - 7\mathbf{I})\mathbf{v} = 0, \quad (\mathbf{A} - 3\mathbf{I})\mathbf{v} = 0 \quad (0.26)$$

They correspond to  $\mathbf{v}_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$  and  $\mathbf{v}_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ . Form  $\mathbf{P}$  and  $\mathbf{D}$ :

$$\mathbf{P} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} 7 & 0 \\ 0 & 3 \end{pmatrix} \quad (0.27)$$

Then

$$\mathbf{B} = \mathbf{P}\mathbf{D}^{1/2}\mathbf{P}^T, \quad \mathbf{B}^2 = \mathbf{A} \quad (0.28)$$

Hence verified for  $2 \times 2$ .

**Example b for (II)**

$$\mathbf{A} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix} \quad (0.29)$$

## Solution

Compute  $|\mathbf{A} - \lambda \mathbf{I}| = 0$ :

$$\begin{vmatrix} 2 - \lambda & 0 & 0 \\ 0 & 3 - \lambda & 0 \\ 0 & 0 & 4 - \lambda \end{vmatrix} = 0 \quad (0.30)$$

$$(2 - \lambda)(3 - \lambda)(4 - \lambda) = 0 \quad (0.31)$$

$$\lambda_1 = 2, \lambda_2 = 3, \lambda_3 = 4 \quad (0.32)$$

Diagonalization:

$$\mathbf{A} = \mathbf{P} \mathbf{D} \mathbf{P}^{\top}, \quad \mathbf{P} = \mathbf{I}, \quad \mathbf{D} = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix} \quad (0.33)$$

## Solution

Define

$$\mathbf{B} = \mathbf{P}\mathbf{D}^{1/2}\mathbf{P}^{\top} = \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & \sqrt{3} & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad (0.34)$$

Then

$$\mathbf{B}^2 = \mathbf{A} \quad (0.35)$$

Hence  $\mathbf{B}$  is symmetric positive definite.

**Conclusion:** In all four examples, both statements (I) and (II) hold true.