Assignment6

G HARSHA VARDHAN REDDY (CS21BTECH11017)

May 15, 2022

Outline

- Abstract
- Problem
- Theory
- Solution

Abstract

This document contains 6^th problem from exercise 2 of CBSE Class 12 (Probability).



Problem

Exercise 2 Problem 3 Let E and F be events with $P(E) = \frac{3}{5}$, $P(F) = \frac{3}{10}$ and $P(EF) = \frac{1}{5}$. Are E and F independent?



Theory

Independent events:

Two events are independent if the incidence of one event does not affect the probability of the other event.(or)

Two events A, B(say) are said to be independent if P(A|B) = P(A)

$$\implies P(A|B) = \frac{P(AB)}{P(B)} = P(A) \tag{1}$$

$$\implies P(A) \times P(B) = P(AB) \tag{2}$$



Solution

Let's denote the outcome of the experiment by a random variable $X \in \{0,1\}$, where X = 0 denotes occurrence of event E and X = 1denotes occurrence of event F.

$$\implies P(X=0) = \frac{3}{5},\tag{3}$$

$$P(X=1) = \frac{3}{10} \text{ and} \tag{4}$$

$$\Rightarrow P(X = 0) = \frac{3}{5},$$

$$P(X = 1) = \frac{3}{10} \text{ and}$$

$$P(X = 0, X = 1) = \frac{1}{5}$$
(5)



Let's check whether the above events are independent or not. From (3),(4)

$$P(X = 0) \times P(X = 1) = \frac{3}{5} \times \frac{3}{10}$$
 (6)

$$\implies P(X=0) \times P(X=1) = \frac{9}{50} \tag{7}$$

From (5) and (7) it's clear that

$$P(X = 0, X = 1) \neq P(X = 0) \times P(X = 1)$$
 (8)

Which says that the events E and F are not independent.

