

A THEORETICAL STUDY OF THE IMPLICATIONS OF RESOURCE COMPETITION FOR ADAPTIVE THERAPY OF CASTRATION-RESISTANT PROSTATE CANCER

THESIS DEFENCE

Harshavardhan BV
Supervisor: Prof. Sutirth Dey
July 2021

IISER Pune

INTRODUCTION

HOW IS CANCER USUALLY TREATED?




Figure 1: Competitive release under SOC¹

- Conventional therapy aims to reduce tumour burden (Frei & Canellos, 1980)

¹SOC: Standard-Of-Care

²MTD: Maximum Tolerated Dose

HOW IS CANCER USUALLY TREATED?




Figure 1: Competitive release under SOC¹

- Conventional therapy aims to reduce tumour burden (Frei & Canellos, 1980)
- SOC¹: Drugs applied at MTD²

¹SOC: Standard-Of-Care

²MTD: Maximum Tolerated Dose

HOW IS CANCER USUALLY TREATED?




Figure 1: Competitive release under SOC¹

- Conventional therapy aims to reduce tumour burden (Frei & Canellos, 1980)
- SOC¹: Drugs applied at MTD²
- Cells have heterogeneous sensitivity → without therapy - sensitive keep resistant in check

¹SOC: Standard-Of-Care

²MTD: Maximum Tolerated Dose

HOW IS CANCER USUALLY TREATED?




Figure 1: Competitive release under SOC¹

- Conventional therapy aims to reduce tumour burden (Frei & Canellos, 1980)
- SOC¹: Drugs applied at MTD²
- Cells have heterogeneous sensitivity → without therapy - sensitive keep resistant in check
- MTD² eliminates sensitive → resistant tumour doesn't respond to therapy (Scott & Marusyk, 2017)

¹SOC: Standard-Of-Care

²MTD: Maximum Tolerated Dose

HOW TO AVOID COMPETITIVE RELEASE?




Figure 2: Control under AT¹

- AT¹: apply drugs at lower, fluctuating doses (Gatenby et al., 2009)

¹AT: Adaptive Therapy

HOW TO AVOID COMPETITIVE RELEASE?




Figure 2: Control under AT¹

- AT¹: apply drugs at lower, fluctuating doses (Gatenby et al., 2009)
- During drug holiday → sensitive compete with resistant

¹AT: Adaptive Therapy

HOW TO AVOID COMPETITIVE RELEASE?




Figure 2: Control under AT¹

- AT¹: apply drugs at lower, fluctuating doses (Gatenby et al., 2009)
- During drug holiday → sensitive compete with resistant
- AT¹ outcome depends on competition

¹AT: Adaptive Therapy

- Castration-Resistant Prostate Cancer (CRPC)

- Castration-Resistant Prostate Cancer (CRPC)
- Difficult to cure with current treatments
- Shift in goal to extend survival

- Castration-Resistant Prostate Cancer (CRPC)
- Difficult to cure with current treatments
- Shift in goal to extend survival




Figure 3: Schematic representation of cell types

Cell type	Testosterone dependent	Testosterone producing	Mechanism of resistance
T^+	Yes	No	N/A
T^p	Yes	Yes	Cholesterol $\xrightarrow{CYP17\alpha}$ Testosterone
T^-	No	No	AR ¹ mutation

¹AR: Androgen Receptors

- Castration-Resistant Prostate Cancer (CRPC)
- Difficult to cure with current treatments
- Shift in goal to extend survival
- Prostate cells: AR¹ that trigger proliferation when activated by testosterone (Heinlein & Chang, 2004)




Figure 3: Schematic representation of cell types

Cell type	Testosterone dependent	Testosterone producing	Mechanism of resistance
T ⁺	Yes	No	N/A
T ^p	Yes	Yes	Cholesterol $\xrightarrow{CYP17\alpha}$ Testosterone
T ⁻	No	No	AR ¹ mutation

¹AR: Androgen Receptors

HOW DO WE MODEL THIS?

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{\sum_j y_j}{1 + K_{i,max} f_i(O_2) f_i(test)} \right) - \delta_i y_i \quad (1)$$

¹ $i \in \{T^+, T^p, T^-\}$, $res \in \{O_2, test\}$

² ll : lower limit, ul : upper limit

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{\sum_j y_j}{1 + K_{i,max} f_i(O_2) f_i(test)}\right) - \delta_i y_i \quad (1)$$

$$f_i(res) = \begin{cases} 1 & \text{if } ul_{res,i} \leq res \\ \frac{res - ll_{res,i}}{ul_{res,i} - ll_{res,i}} & \text{if } ll_{res,i} < res < ul_{res,i} \\ 0 & \text{if } res \leq ll_{res,i} \end{cases} \quad (2)$$




Figure 4: Function dependence of carrying capacity on resource

¹ $i \in \{T^+, T^p, T^-\}$, $res \in \{O_2, test\}$

² ll : lower limit, ul : upper limit

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{\sum_j y_j}{1 + K_{i,max} f_i(O_2) f_i(test)}\right) - \delta_i y_i \quad (1)$$

$$f_i(res) = \begin{cases} 1 & \text{if } ul_{res,i} \leq res \\ \frac{res - ll_{res,i}}{ul_{res,i} - ll_{res,i}} & \text{if } ll_{res,i} < res < ul_{res,i} \\ 0 & \text{if } res \leq ll_{res,i} \end{cases} \quad (2)$$

$$\frac{dO_2}{dt} = p_{O_2} - \sum_i \mu_{O_2,i} y_i - \lambda_{O_2} O_2 \quad (3)$$

$$\frac{d(test)}{dt} = p_{test} y_{TP} - \sum_i \mu_{test,i} y_i - \lambda_{test} test \quad (4)$$




Figure 4: Function dependence of carrying capacity on resource

¹ $i \in \{T^+, T^p, T^-\}$, $res \in \{O_2, test\}$

² ll : lower limit, ul : upper limit

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{\sum_j y_j}{1 + K_{i,max} f_i(O_2) f_i(test)}\right) - \delta_i y_i \quad (1)$$

$$f_i(res) = \begin{cases} 1 & \text{if } ul_{res,i} \leq res \\ \frac{res - ll_{res,i}}{ul_{res,i} - ll_{res,i}} & \text{if } ll_{res,i} < res < ul_{res,i} \\ 0 & \text{if } res \leq ll_{res,i} \end{cases} \quad (2)$$

$$\frac{dO_2}{dt} = p_{O_2} - \sum_i \mu_{O_2,i} y_i - \lambda_{O_2} O_2 \quad (3)$$

$$\frac{d(test)}{dt} = p_{test} y_{TP} - \sum_i \mu_{test,i} y_i - \lambda_{test} test \quad (4)$$




Figure 4: Function dependence of carrying capacity on resource

- Assumptions: No mutation, no spatial structure, well mixed
- Defined $\mathbb{R}_{\geq 0}$, $y_i < 1$ = extinction

¹ $i \in \{T^+, T^p, T^-\}$, $res \in \{O_2, test\}$

² ll : lower limit, ul : upper limit


- Some values directly from literature based on cell line data (Hail et al., 2010; Jain et al., 2011)
 - T^+ : LNCaP
 - T^p : 22Rv1
 - T^- : PC3

- Some values directly from literature based on cell line data (Hail et al., 2010; Jain et al., 2011)
 - T^+ : LNCaP
 - T^p : 22Rv1
 - T^- : PC3
- Some values derieved from literature via constraint equations (ATCC, 2021; Stewart et al., 2010; Titus et al., 2005)


- Some values directly from literature based on cell line data (Hail et al., 2010; Jain et al., 2011)
 - T^+ : LNCaP
 - T^p : 22Rv1
 - T^- : PC3
- Some values derived from literature via constraint equations (ATCC, 2021; Stewart et al., 2010; Titus et al., 2005)
- Study Parameters:
 - Resource limitations: varied using lower and upper limits
 - Initial seeding: different ratios of cell types and total population

WHAT HAPPENS IN THE ABSENCE OF THERAPY?

ALL CELL-TYPE COMPETITION OUTCOMES



(a) Equal seeding - 1:1:1




(b) High T^p seeding - 8:1:1


Figure 5: Final ratio of all cell types. (Stacked bar plot)

- All the cells have the same limitations

ALL CELL-TYPE COMPETITION OUTCOMES



(a) Equal seeding - 1:1:1




(b) High T^p seeding - 8:1:1


Figure 5: Final ratio of all cell types. (Stacked bar plot)

- All the cells have the same limitations
- Higher T^p seeding ratios \Rightarrow increased testosterone production

ALL CELL-TYPE COMPETITION OUTCOMES



(a) Equal seeding - 1:1:1




(b) High T^p seeding - 8:1:1


Figure 5: Final ratio of all cell types. (Stacked bar plot)

- All the cells have the same limitations
- Higher T^p seeding ratios \Rightarrow increased testosterone production
- Coexistence is important: tumour with T^p and T^+ would only respond to therapy

ALL CELL-TYPE COMPETITION OUTCOMES



(a) Equal seeding - 1:1:1




(b) High T^p seeding - 8:1:1

Figure 5: Final ratio of all cell types. (Stacked bar plot)


- All the cells have the same limitations
- Higher T^p seeding ratios \Rightarrow increased testosterone production
- Coexistence is important: tumour with T^p and T^+ would only respond to therapy

- Testosterone: drastic effect on coexistence as only $T^p - T^+$ affected

ALL CELL-TYPE COMPETITION OUTCOMES



(a) Equal seeding - 1:1:1



(b) High T^p seeding - 8:1:1

Figure 5: Final ratio of all cell types. (Stacked bar plot)

- All the cells have the same limitations
- Higher T^p seeding ratios \Rightarrow increased testosterone production
- Coexistence is important: tumour with T^p and T^+ would only respond to therapy

- Testosterone: drastic effect on coexistence as only $T^p - T^+$ affected
- Oxygen: minor effect, pushes to extinction if combined limitation on the edge

WHAT HAPPENS WHEN WE ADD THERAPY?

HOW IS IT IMPLEMENTED?

- Therapy: modelled as boolean

1 = MTD
0 = no dose (5)

¹MTD: Maximum tolerated dose, SOC: Standard-Of-Care, AT: Adaptive Therapy

²abi: abiraterone, dtx: docetaxel

HOW IS IT IMPLEMENTED?

- Therapy: modelled as boolean

$$\begin{aligned} 1 &= \text{MTD} \\ 0 &= \text{no dose} \end{aligned} \quad (5)$$

- Abiraterone: blocks $CYP17\alpha$
 T^p , T^+ affected

$$p_{test}(abi) = \begin{cases} p_{test,max} & \text{if } abi = 0 \\ p_{test,min} & \text{if } abi = 1 \end{cases} \quad (6)$$

¹MTD: Maximum tolerated dose, SOC: Standard-Of-Care, AT: Adaptive Therapy

²abi: abiraterone, dtx: docetaxel

HOW IS IT IMPLEMENTED?

- Therapy: modelled as boolean

$$\begin{aligned} 1 &= \text{MTD} \\ 0 &= \text{no dose} \end{aligned} \quad (5)$$

- Abiraterone: blocks $CYP17\alpha$
 T^p, T^+ affected

$$p_{test}(abi) = \begin{cases} p_{test,max} & \text{if } abi = 0 \\ p_{test,min} & \text{if } abi = 1 \end{cases} \quad (6)$$

- Docetaxel: disrupts microtubule
All 3 affected

$$r_i(dtx) = \begin{cases} r_{i,max} & \text{if } dtx = 0 \\ r_{i,min} & \text{if } dtx = 1 \end{cases} \quad (7)$$

¹MTD: Maximum tolerated dose, SOC: Standard-Of-Care, AT: Adaptive Therapy

²abi: abiraterone, dtx: docetaxel

HOW IS IT IMPLEMENTED?

- Therapy: modelled as boolean

$$\begin{aligned} 1 &= \text{MTD} \\ 0 &= \text{no dose} \end{aligned} \quad (5)$$

- SOC¹: dose given at MTD¹ from the beginning

$$\text{dose}(y, t) = 1 \quad \forall t, y \quad (8)$$

- Abiraterone: blocks $\text{CYP17}\alpha$
 T^p, T^+ affected

$$p_{\text{test}}(\text{abi}) = \begin{cases} p_{\text{test}, \max} & \text{if } \text{abi} = 0 \\ p_{\text{test}, \min} & \text{if } \text{abi} = 1 \end{cases} \quad (6)$$

- Docetaxel: disrupts microtubule
All 3 affected

$$r_i(\text{dtx}) = \begin{cases} r_{i, \max} & \text{if } \text{dtx} = 0 \\ r_{i, \min} & \text{if } \text{dtx} = 1 \end{cases} \quad (7)$$

¹MTD: Maximum tolerated dose, SOC: Standard-Of-Care, AT: Adaptive Therapy

²abi: abiraterone, dtx: docetaxel

HOW IS IT IMPLEMENTED?

- Therapy: modelled as boolean

$$\begin{aligned} 1 &= \text{MTD} \\ 0 &= \text{no dose} \end{aligned} \quad (5)$$

- Abiraterone: blocks CYP17 α
 T^p , T^+ affected

$$p_{\text{test}}(\text{abi}) = \begin{cases} p_{\text{test},\text{max}} & \text{if } \text{abi} = 0 \\ p_{\text{test},\text{min}} & \text{if } \text{abi} = 1 \end{cases} \quad (6)$$

- Docetaxel: disrupts microtubule
All 3 affected

$$r_i(\text{dtx}) = \begin{cases} r_{i,\text{max}} & \text{if } \text{dtx} = 0 \\ r_{i,\text{min}} & \text{if } \text{dtx} = 1 \end{cases} \quad (7)$$

- SOC¹: dose given at MTD¹ from the beginning

$$\text{dose}(y, t) = 1 \quad \forall t, y \quad (8)$$

- AT¹: binary mode considered


- Dose at MTD¹ when on
- Therapy turned on when population above On threshold
- Therapy turned off when population below Off threshold

$$\text{dose}(y, t) = \begin{cases} 0 & \text{if } \text{dose}(y, t - \Delta t) = 0 \text{ and } y < \text{On} \\ 1 & \text{if } \text{dose}(y, t - \Delta t) = 0 \text{ and } y \geq \text{On} \\ 1 & \text{if } \text{dose}(y, t - \Delta t) = 1 \text{ and } y > \text{Off} \\ 0 & \text{if } \text{dose}(y, t - \Delta t) = 1 \text{ and } y \leq \text{Off} \end{cases} \quad (9)$$


¹MTD: Maximum tolerated dose, SOC: Standard-Of-Care, AT: Adaptive Therapy

²abi: abiraterone, dtx: docetaxel

WHAT HAPPENS WITH STANDARD-OF-CARE?



(a) Equal seeding - 1:1:1



(b) High T^P seeding - 8:1:1

Figure 6: Final ratio of all cell types under standard-of-care. (Stacked bar plot)

- T^+ , T^P go extinct in all cases

- Testosterone levels insufficient for growth

THRESHOLDS FOR ADAPTIVE THERAPY




Figure 7: Standardisation of threshold for adaptive therapy

- Low threshold: T^- inhibits $T^p - T^+$ and causes extinction

THRESHOLDS FOR ADAPTIVE THERAPY




Figure 7: Standardisation of threshold for adaptive therapy

- Low threshold: T^- inhibits $T^p - T^+$ and causes extinction
- High threshold: $T^+ - T^p$ can compete with T^- (Hansen & Read, 2020)

THRESHOLDS FOR ADAPTIVE THERAPY




Figure 7: Standardisation of threshold for adaptive therapy

- Low threshold: T^- inhibits $T^p - T^+$ and causes extinction
- High threshold: $T^+ - T^p$ can compete with T^- (Hansen & Read, 2020)
- Too High: No therapy applied as On threshold never crossed

THRESHOLDS FOR ADAPTIVE THERAPY




Figure 7: Standardisation of threshold for adaptive therapy

- Low threshold: T^- inhibits $T^p - T^+$ and causes extinction
- High threshold: $T^+ - T^p$ can compete with T^- (Hansen & Read, 2020)
- Too High: No therapy applied as On threshold never crossed
- Chosen- On: 6000, Off: 4000

THRESHOLDS FOR ADAPTIVE THERAPY




Figure 7: Standardisation of threshold for adaptive therapy

- Low threshold: T^- inhibits $T^p - T^+$ and causes extinction
- High threshold: $T^+ - T^p$ can compete with T^- (Hansen & Read, 2020)
- Too High: No therapy applied as On threshold never crossed
- Chosen- On: 6000, Off: 4000
- $T^+ + T^p$ only for threshold
- With total: $T^+ - T^p$ go extinct before therapy turned off

IS ADAPTIVE THERAPY BETTER?




Figure 8: Time-series of all cell types with adaptive therapy. (On:6000, Off:4000)

- $T^+ - T^P$ extinct just by competition: no effect

IS ADAPTIVE THERAPY BETTER?






Figure 8: Time-series of all cell types with adaptive therapy. (On:6000, Off:4000)

- $T^+ - T^P$ extinct just by competition: no effect
- More $T^+ - T^P \rightarrow$ more responsive to abiraterone and better suppress T^-

IS ADAPTIVE THERAPY BETTER?



(a) Equal seeding - 1:1:1



(b) High T^p seeding - 8:1:1

Figure 8: Time-series of all cell types with adaptive therapy. (On:6000, Off:4000)

- $T^+ - T^p$ extinct just by competition: no effect
- More $T^+ - T^p \rightarrow$ more responsive to abiraterone and better suppress T^-

- Success of Adaptive therapy:

IS ADAPTIVE THERAPY BETTER?






Figure 8: Time-series of all cell types with adaptive therapy. (On:6000, Off:4000)

- $T^+ - T^P$ extinct just by competition: no effect
- More $T^+ - T^P \rightarrow$ more responsive to abiraterone and better suppress T^-
- Success of Adaptive therapy:
 - ✓ Preventing competitive release of resistant

IS ADAPTIVE THERAPY BETTER?



(a) Equal seeding - 1:1:1



(b) High T^P seeding - 8:1:1


Figure 8: Time-series of all cell types with adaptive therapy. (On:6000, Off:4000)

- $T^+ - T^P$ extinct just by competition: no effect
- More $T^+ - T^P \rightarrow$ more responsive to abiraterone and better suppress T^-


- Success of Adaptive therapy:
 - ✓ Preventing competitive release of resistant
 - × Reducing tumour burden: T^- replace dead cells

CAN ADAPTIVE THERAPY BE EVEN MADE
BETTER?

CAN DELAYING TREATMENT HELP?



(a) Equal seeding - 1:1:1




(b) High T^P seeding - 8:1:1


Figure 9: Time-series of all cell types with adaptive therapy delayed by 200 days. (On:6000, Off:4000)

- $T^+ - T^P \uparrow$ in periods of no therapy

CAN DELAYING TREATMENT HELP?



(a) Equal seeding - 1:1:1




(b) High T^P seeding - 8:1:1


Figure 9: Time-series of all cell types with adaptive therapy delayed by 200 days. (On:6000, Off:4000)

- $T^+ - T^P \uparrow$ in periods of no therapy
- Speculation: Can delaying bring better balance between $T^+ - T^P$ and T^- ?

CAN DELAYING TREATMENT HELP?



(a) Equal seeding - 1:1:1



(b) High T^p seeding - 8:1:1

Figure 9: Time-series of all cell types with adaptive therapy delayed by 200 days. (On:6000, Off:4000)

- $T^+ - T^p \uparrow$ in periods of no therapy
- Speculation: Can delaying bring better balance between $T^+ - T^p$ and T^- ?

- No advantage found as they have similar temporal dynamics

WHAT ABOUT USING MULTIPLE DRUGS?




Figure 10: Time-series of all cell types with combination adaptive therapy of abi and dtx¹. abi(On:6000, Off:4000; T^+ + T^P), dtx(On:6000, Off:4000; Total)

- Hormonal (abi¹) + cytotoxic (dtx¹) (West et al., 2019)

¹abi: abiraterone, dtx: docetaxel

WHAT ABOUT USING MULTIPLE DRUGS?




Figure 10: Time-series of all cell types with combination adaptive therapy of abi and dtx¹. abi(On:6000, Off:4000; $T^+ + T^P$), dtx(On:6000, Off:4000; Total)

- Hormonal (abi¹) + cytotoxic (dtx¹) (West et al., 2019)
- Method of action different \Rightarrow Minimal cross resistance

¹abi: abiraterone, dtx: docetaxel

WHAT ABOUT USING MULTIPLE DRUGS?




Figure 10: Time-series of all cell types with combination adaptive therapy of abi and dtx¹. abi(On:6000, Off:4000; T^+ + T^P), dtx(On:6000, Off:4000; Total)

- Hormonal (abi¹) + cytotoxic (dtx¹) (West et al., 2019)
- Method of action different \Rightarrow Minimal cross resistance

¹abi: abiraterone, dtx: docetaxel

- Test-of-concept: would require extensive standardization in future

WHAT ABOUT USING MULTIPLE DRUGS?




Figure 10: Time-series of all cell types with combination adaptive therapy of abi and dtx¹. abi(On:6000, Off:4000; $T^+ + T^P$), dtx(On:6000, Off:4000; Total)

- Hormonal (abi¹) + cytotoxic (dtx¹) (West et al., 2019)
- Method of action different \Rightarrow Minimal cross resistance

¹abi: abiraterone, dtx: docetaxel

- Test-of-concept: would require extensive standardization in future
- -ve effect on coexistence by $\downarrow T^+ - T^P$ outweigh +ve effect on coexistence by $\downarrow T^-$

CONCLUSION

- Resource levels can control strength of competition

- Resource levels can control strength of competition
- Balance of limitations promote coexistence

- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction

- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction
- With adaptive therapy: competitive release avoided
 - Effectiveness depends on T^+ and T^p population
 - Population controlled by resource limitations
 - Maximum limit on T^+ and T^p by thresholds of adaptive therapy

- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction
- With adaptive therapy: competitive release avoided
 - Effectiveness depends on T^+ and T^p population
 - Population controlled by resource limitations
 - Maximum limit on T^+ and T^p by thresholds of adaptive therapy
- Future directions:

- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction
- With adaptive therapy: competitive release avoided
 - Effectiveness depends on T^+ and T^p population
 - Population controlled by resource limitations
 - Maximum limit on T^+ and T^p by thresholds of adaptive therapy
- Future directions:
 - Make adaptive therapy effective at reducing tumour burden








- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction
- With adaptive therapy: competitive release avoided
 - Effectiveness depends on T^+ and T^p population
 - Population controlled by resource limitations
 - Maximum limit on T^+ and T^p by thresholds of adaptive therapy
- Future directions:
 - Make adaptive therapy effective at reducing tumour burden
 - Dynamic thresholds for turning on/off based on composition of the tumour

- Resource levels can control strength of competition
- Balance of limitations promote coexistence
- With standard-of-care: testosterone limitation is increased leading to extinction
- With adaptive therapy: competitive release avoided
 - Effectiveness depends on T^+ and T^P population
 - Population controlled by resource limitations
 - Maximum limit on T^+ and T^P by thresholds of adaptive therapy
- Future directions:
 - Make adaptive therapy effective at reducing tumour burden
 - Dynamic thresholds for turning on/off based on composition of the tumour
 - Different limitations for different cell types

I would like to thank the following people:

- Supervisor: Prof. Sutirth Dey
- Expert: Dr. M.S. Madhusudhan
- Mentor: Vibishan B
- PBL Members
- Friends and Family
- KVPY and IISER Pune

REFERENCES

-  ATCC. (2021). Atcc: The global bioresource center. <https://www.atcc.org/>
-  Frei, E., & Canellos, G. P. (1980). Dose: A critical factor in cancer chemotherapy. *The American Journal of Medicine*, 69(4), 585–594. [https://doi.org/https://doi.org/10.1016/0002-9343\(80\)90472-6](https://doi.org/https://doi.org/10.1016/0002-9343(80)90472-6)
-  Gatenby, R. A., Silva, A. S., Gillies, R. J., & Frieden, B. R. (2009). Adaptive therapy. *Cancer Research*, 69(11), 4894–4903. <https://doi.org/10.1158/0008-5472.CAN-08-3658>
-  Hail, N., Chen, P., & Bushman, L. R. (2010). Teriflunomide (leflunomide) promotes cytostatic, antioxidant, and apoptotic effects in transformed prostate epithelial cells: Evidence supporting a role for teriflunomide in prostate cancer chemoprevention. *Neoplasia*, 12(6), 464–475. <https://doi.org/https://doi.org/10.1593/neo.10168>
-  Hansen, E., & Read, A. F. (2020). Modifying adaptive therapy to enhance competitive suppression. *Cancers*, 12(12). <https://doi.org/10.3390/cancers12123556>
-  Heinlein, C. A., & Chang, C. (2004). Androgen Receptor in Prostate Cancer. *Endocrine Reviews*, 25(2), 276–308. <https://doi.org/10.1210/er.2002-0032>
-  Jain, H. V., Clinton, S. K., Bhinder, A., & Friedman, A. (2011). Mathematical modeling of prostate cancer progression in response to androgen ablation therapy. *Proceedings of the National Academy of Sciences*, 108(49), 19701–19706. <https://doi.org/10.1073/pnas.1115750108>
-  Scott, J., & Marusyk, A. (2017). Somatic clonal evolution: A selection-centric perspective [Evolutionary principles - heterogeneity in cancer?]. *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer*, 1867(2), 139–150. <https://doi.org/https://doi.org/10.1016/j.bbcan.2017.01.006>



Stewart, G. D., Ross, J. A., McLaren, D. B., Parker, C. C., Habib, F. K., & Riddick, A. C. (2010). The relevance of a hypoxic tumour microenvironment in prostate cancer. *BJU International*, 105(1), 8–13.
<https://doi.org/https://doi.org/10.1111/j.1464-410X.2009.08921.x>



Titus, M. A., Schell, M. J., Lih, F. B., Tomer, K. B., & Mohler, J. L. (2005). Testosterone and dihydrotestosterone tissue levels in recurrent prostate cancer. *Clinical Cancer Research*, 11(13), 4653–4657. <https://doi.org/10.1158/1078-0432.CCR-05-0525>



West, J. B., Dinh, M. N., Brown, J. S., Zhang, J., Anderson, A. R., & Gatenby, R. A. (2019). Multidrug cancer therapy in metastatic castrate-resistant prostate cancer: An evolution-based strategy. *Clinical Cancer Research*, 25(14), 4413–4421.
<https://doi.org/10.1158/1078-0432.CCR-19-0006>