

# Studying the effects of competition on adaptive therapy

## Mid Year Presentation

Harshavardhan BV (20161100)  
5th Year BS-MS

Under the guidance of Prof. Sutirth Dey,  
IISER Pune

January 2021

# Adaptive Therapy

- ▶ Conventional therapy @ MTD  $\rightarrow$  min tumour burden (Frei et al. 1980)
- ▶ Heterogenous sensitivity  $\rightarrow$  sensitive eliminated  $\rightarrow$  resistant population (Scott et al. 2017)
- ▶ AT = lower, fluctuating dose  $\rightarrow$  sensitive preserved
- ▶ AT dose =  $f(\text{tumour size})$  (Gatenby et al. 2009)

- ▶ System of study: Metastatic Castration-Resistant Prostate Cancer
- ▶ History of Adaptive therapy work (Cunningham et al. 2018)
- ▶ Therapy: ADT + Abiraterone

Cell type	Test. dependent	Test. Producing	Ab. sensitive	Mechanism
$T^+$	Yes	No	Yes	N/A
$T^p$	Yes	Yes	Yes	Cholestrol $\xrightarrow{CYP17\alpha}$ Testosterone
$T^-$	No	No	No	Androgen receptor mutations

# Competition between cells

- ▶ AT outcome  $\sim$  competition b/w sensitive and resistant
- ▶ sensitive keeps resistant in check
- ▶ sensitive killed by drug dose
- ▶ Competitive strategies through traits from cancer progression (Hanahan et al. 2011)
  - ▶ Higher proliferation rate
  - ▶ Better survival @ sub-optimal conditions
  - ▶ Lower death rate

# ODE Model

- ▶ Starting point: forming expectations, parameterization
- ▶ Logistic framework with dynamic carrying capacity  $\sim$  environmental conditions
- ▶ Environment = resource = {oxygen, testosterone}




Figure 1:  $f_i(res)$

$$\frac{dy_i}{dt} = r_i y_i \left(1 - \frac{\sum_j y_j}{1 + K_{i,max} f_i(O_2) f_i(test)}\right) - \delta_i y_i \quad (1)$$

$$\frac{dO_2}{dt} = p_{O_2} - \sum_i \mu_{O_2,i} y_i - \lambda_{O_2} O_2 \quad (2)$$

$$\frac{dtest}{dt} = p_{test} y_{TP} - \sum_i \mu_{test,i} y_i - \lambda_{test} test \quad (3)$$

$$f_i(res) = \begin{cases} 1 & \text{if } ul_{res,i} \leq res \\ \frac{res - ll_{res,i}}{ul_{res,i} - ll_{res,i}} & \text{if } ll_{res,i} < res < ul_{res,i} \\ 0 & \text{if } res \leq ll_{res,i} \end{cases} \quad (4)$$

$i \in \{T^+, T^p, T^-\}$  and  $res \in \{O_2, test\}$ .

# Parameters & Standardization

Some parameters directly from literature

- ▶  $\delta_i$ : Death rate (Jain et al. 2011)

$$T^+ \quad 2.5 \times 10^{-3} \text{ min}^{-1}$$

$$T^p \quad 2.5 \times 10^{-3} \text{ min}^{-1}$$

$$T^- \quad 1.6 \times 10^{-4} \text{ min}^{-1}$$

- ▶  $\mu_{O_2,i}$ : Oxygen uptake (Hail et al. 2010)

$$T^+ \quad 1.63 \times 10^{-6} \text{ min}^{-1} \text{ cell}^{-1}$$

$$T^p \quad 1.63 \times 10^{-6} \text{ min}^{-1} \text{ cell}^{-1}$$

$$T^- \quad 1.04 \times 10^{-6} \text{ min}^{-1} \text{ cell}^{-1}$$

- ▶  $\lambda_{res}$ : Decay rate (Jain et al. 2011)

$$O_2 \quad 0.100 \text{ min}^{-1}$$

$$test \quad 0.004 \text{ min}^{-1}$$

# Parameters & Standardization

Some parameters from assumptions & constraints

- $r_i$ : Growth rate (Eq 5)

$$T^+ \quad 2.84 \times 10^{-3} \text{ min}^{-1}$$

$$T^P \quad 2.79 \times 10^{-3} \text{ min}^{-1}$$

$$T^- \quad 6.23 \times 10^{-4} \text{ min}^{-1}$$

- $K_{i,max}$ : Maximum Carrying capacity (Eq 6)

$$T^+ \quad 8.35 \times 10^4$$

$$T^P \quad 9.62 \times 10^4$$

$$T^- \quad 1.34 \times 10^4$$

- $p_{res}$ : Production rate (Eq 7,8)

$$O_2 \quad 0.11 \text{ min}^{-1}$$

$$test \quad 5 \times 10^{-7} \text{ min}^{-1} \text{ cell}^{-1}$$

- $\mu_{test,i}$ : Testosterone uptake (Eq 8)

$$T^+ \quad 2.34 \times 10^{-8} \text{ min}^{-1} \text{ cell}^{-1}$$

$$T^P \quad 6.00 \times 10^{-8} \text{ min}^{-1} \text{ cell}^{-1}$$

$$T^- \quad 0 \text{ min}^{-1} \text{ cell}^{-1}$$

- $ll_{res,i}$ : Lower limit/threshold level  $\in [0, 1]$

- $ul_{res,i}$ : Upper limit/saturation level  $\in [0, 1]$

$$r_i = \frac{\ln(2)}{\tau_{d,i}} + \delta_i \quad (5)$$

$$K_{i,max} = \frac{r_i}{r_i - \delta_i} y_i^* \quad (6)$$

$$p_{O_2} = \lambda_{O_2} O_2^* + y_i^* \mu_i \quad (7)$$

$$p_{test} - \mu_{test,TP} = \frac{test^* \lambda_{test}}{y_{TP}^*} = 4 \times 10^{-4} \quad (8)$$

# Pairwise Competition: $T^P - T^-$

1.  $T^P$  test &  $O_2$  limited.  $T^-$  only  $O_2$  limited




Figure 2: Pairwise  $T^P - T^-$  timeseries, when  $T^P$  is testosterone limited and not testosterone limited (columns) and at different initial proportions of  $T^P$  (rows)



## Pairwise Competition: $T^P - T^-$

1.  $T^P$  test &  $O_2$  limited.  $T^-$  only  $O_2$  limited
2.  $T^P$  not severely testosterone limited
  - ▶  $T^P$  coexist or outcompete  $T^-$
  - ▶  $T^-$  outcompetes in other cases




Figure 2: Pairwise  $T^P - T^-$  timeseries, when  $T^P$  is testosterone limited and not testosterone limited (columns) and at different initial proportions of  $T^P$  (rows)

## Pairwise Competition: $T^P - T^-$

1.  $T^P$  test &  $O_2$  limited.  $T^-$  only  $O_2$  limited
2.  $T^P$  not severely testosterone limited
  - ▶  $T^P$  coexist or outcompete  $T^-$
  - ▶  $T^-$  outcompetes in other cases
3. Outcomes dependent on the initial proportion of  $T^P$




Figure 2: Pairwise  $T^P - T^-$  timeseries, when  $T^P$  is testosterone limited and not testosterone limited (columns) and at different initial proportions of  $T^P$  (rows)

# Pairwise Competition: $T^P - T^-$

1.  $T^P$  test &  $O_2$  limited.  $T^-$  only  $O_2$  limited
2.  $T^P$  not severely testosterone limited
  - ▶  $T^P$  coexist or outcompete  $T^-$
  - ▶  $T^-$  outcompetes in other cases
3. Outcomes dependent on the initial proportion of  $T^P$
4.  $T^-$  strongly oxygen limited
  - ▶  $T^P$  still testosterone limited
  - ▶  $T^-$  wins eventually
  - ▶ Oxygen levels rise faster than testosterone




Figure 2: Pairwise  $T^P - T^-$  timeseries, when  $T^P$  is testosterone limited and not testosterone limited (columns) and at different initial proportions of  $T^P$  (rows)




Figure 3: Pairwise  $T^P - T^-$  timeseries, when  $T^-$  is severely oxygen limited

## Pairwise Competition: $T^+ - T^P$

### 1. Both cell type limited by both resource




Figure 4: Pairwise  $T^+ - T^P$  timeseries, when both cell types are testosterone limited and when  $T^+$  is limited more than  $T^P$

## Pairwise Competition: $T^+ - T^P$

1. Both cell type limited by both resource
2. Both severely testosterone limited
  - ▶  $T^+$  consume & grows on limited testosterone
  - ▶ Density-dependent competition drive  $T^P$  extinct
  - ▶ No  $T^P$  = No testosterone  $\rightarrow T^+$  extinct




Figure 4: Pairwise  $T^+ - T^P$  timeseries, when both cell types are testosterone limited and when  $T^+$  is limited more than  $T^P$

## Pairwise Competition: $T^+ - T^P$

1. Both cell type limited by both resource
2. Both severely testosterone limited
  - ▶  $T^+$  consume & grows on limited testosterone
  - ▶ Density-dependent competition drive  $T^P$  extinct
  - ▶ No  $T^P$  = No testosterone  $\rightarrow T^+$  extinct
3.  $T^P$  is weakly limited by testosterone relative to  $T^+$ 
  - ▶ Both coexist
  - ▶  $T^P$  grow initially & not affected by  $T^+$




Figure 4: Pairwise  $T^+ - T^P$  timeseries, when both cell types are testosterone limited and when  $T^+$  is limited more than  $T^P$

## Pairwise Competition: $T^+ - T^P$

1. Both cell type limited by both resource
2. Both severely testosterone limited
  - ▶  $T^+$  consume & grows on limited testosterone
  - ▶ Density-dependent competition drive  $T^P$  extinct
  - ▶ No  $T^P$  = No testosterone  $\rightarrow T^+$  extinct
3.  $T^P$  is weakly limited by testosterone relative to  $T^+$ 
  - ▶ Both coexist
  - ▶  $T^P$  grow initially & not affected by  $T^+$
4.  $T^+$  severely oxygen limited
  - ▶  $T^P$  grow initially & secrete testosterone
  - ▶ Sustain small  $T^+$  if not extinct




Figure 4: Pairwise  $T^+ - T^P$  timeseries, when both cell types are testosterone limited and when  $T^+$  is limited more than  $T^P$




Figure 5: Pairwise  $T^+ - T^P$  timeseries, when both cell types are testosterone limited and not oxygen limited and when  $T^+$  is severely oxygen limited

# Future Plans

- ▶ Testosterone limitation relaxed
- ▶ Oxygen limit exploration with lower  $ul_{test,i}$
- ▶ Make oxygen more limiting than testosterone via production rates
- ▶ 3 cell-type competition
- ▶ Simulate AT regimens with therapy as  $p_{test} = f(dose)$
- ▶ Replicate in ABM & Compare



Thank You



Figure 6: Meme, Original Image by USDA

## References I

- [1] *ATCC: The Global Bioresource Center.*
- [2] Jessica J. Cunningham, Joel S. Brown, Robert A. Gatenby, and Kateřina Staňková. “Optimal control to develop therapeutic strategies for metastatic castrate resistant prostate cancer”. In: *Journal of Theoretical Biology* 459 (2018), pp. 67–78.
- [3] Emil Frei and George P. Canellos. “Dose: A critical factor in cancer chemotherapy”. In: *The American Journal of Medicine* 69.4 (1980), pp. 585–594.
- [4] Robert A. Gatenby, Ariosto S. Silva, Robert J. Gillies, and B. Roy Frieden. “Adaptive Therapy”. In: *Cancer Research* 69.11 (2009), pp. 4894–4903.
- [5] Numsen Hail, Ping Chen, and Lane R. Bushman. “Teriflunomide (Leflunomide) Promotes Cytostatic, Antioxidant, and Apoptotic Effects in Transformed Prostate Epithelial Cells: Evidence Supporting a Role for Teriflunomide in Prostate Cancer Chemoprevention”. In: *Neoplasia* 12.6 (2010), pp. 464–475.
- [6] Douglas Hanahan and Robert A Weinberg. “Hallmarks of cancer: the next generation”. In: *cell* 144.5 (2011), pp. 646–674.
- [7] Harsh Vardhan Jain, Steven K. Clinton, Arvinder Bhinder, and Avner Friedman. “Mathematical modeling of prostate cancer progression in response to androgen ablation therapy”. In: *Proceedings of the National Academy of Sciences* 108.49 (2011), pp. 19701–19706.

## References II

- [8] Jacob Scott and Andriy Marusyk. "Somatic clonal evolution: A selection-centric perspective". In: *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer* 1867.2 (2017). Evolutionary principles - heterogeneity in cancer?, pp. 139–150.
- [9] Thomas A. Stamey, Fuad S. Freiha, John E. McNeal, Elise A. Redwine, Alice S. Whittemore, and Hans-Peter Schmid. "Localized prostate cancer. Relationship of tumor volume to clinical significance for treatment of prostate cancer". In: *Cancer* 71.S3 (1993), pp. 933–938.
- [10] Grant D. Stewart, James A. Ross, Duncan B. McLaren, Christopher C. Parker, Fouad K. Habib, and Antony C.P. Riddick. "The relevance of a hypoxic tumour microenvironment in prostate cancer". In: *BJU International* 105.1 (2010), pp. 8–13.
- [11] Maximilian Strobl, Jeffrey West, Yannick Viossat, Mehdi Damaghi, Mark Robertson-Tessi, Joel Brown, Robert Gatenby, Philip Maini, and Alexander Anderson. "Turnover modulates the need for a cost of resistance in adaptive therapy". In: *bioRxiv* (2020).
- [12] Mark A. Titus, Michael J. Schell, Fred B. Lih, Kenneth B. Tomer, and James L. Mohler. "Testosterone and Dihydrotestosterone Tissue Levels in Recurrent Prostate Cancer". In: *Clinical Cancer Research* 11.13 (2005), pp. 4653–4657.